The thalamus drives light-evoked activity in the habenula of larval zebrafish

◆Ruey-Kuang Cheng[1], ◆Seetha Krishnan[2], Qian Lin[2], David G. C. Hildebrand[3, 4], Isaac H. Bianco[4], Caroline Kibat[5] and *Suresh Jesuthasan[1, 5, 6, 7]

[1]Lee Kong Chian School of Medicine, Nanyang Technological University, Singapore 636921.

[2]NUS Graduate School for Integrative Sciences and Engineering, 28 Medical Drive, National University of Singapore, Singapore 117456.

[3]Graduate Program in Neuroscience, Division of Medical Sciences, Graduate School of Arts and Sciences, Harvard University, Cambridge, Massachusetts 02138 United States of America.

[4]Department of Molecular and Cell Biology, Harvard University, Cambridge, Massachusetts 02138, United States of America.

[5]Neural Circuitry and Behavior Laboratory, Institute of Molecular and Cell Biology, Singapore 138673.

[6]Neuroscience and Behavioral Disorders Program, Duke-NUS Graduate Medical School, 8 College Road, Singapore 169857.

[7]Department of Physiology, National University of Singapore, Singapore 117597.

*Corresponding author: sureshj@ntu.edu.sg; +65 65869545

◆equal contribution

Abstract

The habenula integrates sensory stimuli and reward information to regulate the release of neuromodulators with broad effects on brain state and behavior. One stimulus that affects habenula activity is light, but how it does so is unknown. Here, we address this question using larval zebrafish. Calcium imaging shows that light evokes widespread activity in habenula neurons, coupled with a prominent early response in the dorsal left neuropil. Injection of a lipophilic dye into this region retrogradely labels a retino-recipient thalamic nucleus. Anterograde tracing of the thalamus demonstrates a projection to the habenula, while optogenetic and lesion experiments confirm functional connectivity. An analysis of the mouse mesoscale connectome indicates that a visual nucleus in the thalamus, the ventral lateral geniculate nucleus, projects to the habenula in this species also. Together, these data suggest the existence of a conserved thalamo-habenula projection that enables light to affect habenula activity in vertebrates.

Introduction

The habenula is an evolutionarily conserved structure (Stephenson-Jones et al., 2012) that influences multiple behaviors, ranging from fear (Agetsuma et al., 2010; Lee et al., 2010; Zhang et al., 2016), to learning (Matsumoto and Hikosaka, 2007; 2009; Amo et al., 2014), addiction (Fowler et al., 2011), sleep (Aizawa et al., 2013), aggression (Chou et al., 2016; Golden et al., 2016) and performance under stress (Thornton and Davies, 1991). It acts by regulating the release of broadly-acting neuromodulators such as serotonin, dopamine, epinephrine and histamine (Wang and Aghajanian, 1977; Morley et al., 1985; Jhou et al., 2009; Quina et al., 2014). To precisely control these neuromodulators, such that behavior is appropriate for a given context, the habenula integrates diverse information including internal state, sensory stimuli and reward value.

45  These different types of information reach the habenula via different pathways.

Internal states such as the circadian clock may be conveyed from the hypothalamus, for

example by hypocretin-secreting neurons (Appelbaum et al., 2009). Negative reward

causes the entopeduncular nucleus, or internal segment of the globus pallidus, to send

excitatory input to the habenula (Hong and Hikosaka, 2008). Sensory stimuli such as

odours evoke activity in the zebrafish habenula (Dreosti et al., 2014; Krishnan et al.,

2014) via a direct pathway from the olfactory bulb (Miyasaka et al., 2009). Another

stimulus that causes activity in the habenula is light, as has been demonstrated in rat

(Zhao and Rusak, 2005), pigeon (Semm and Demaine, 1984) and zebrafish (Dreosti et

al., 2014). How light affects the habenula is not known, implying the existence of an

input pathway that is not yet well defined.

56  The habenula is divided into two major regions based on pattern of connectivity. In

mammals, these are called the medial and lateral habenula, while in fish these are the

dorsal and ventral habenula. In larval zebrafish, short pulses of red light cause

asymmetric depolarization of the dorsal habenula, with more cells in the left side

showing response; the response in the ventral habenula is symmetric (Dreosti et al.,

2014). This response is dependent on the eyes (Dreosti et al., 2014), but no direct

pathway from the retina to the habenula has been documented (Burrill and Easter, 1994;

Robles et al., 2014). By retrograde tracing in adult zebrafish, Turner et al (Turner et al.,

2016) proposed that the habenula receives input from the nucleus rostrolateralis, a

thalamic nucleus with retinal input that is found in fish (Butler and Saidel, 2003; Saidel,

2013). However, no connectivity with the retina was shown, and because the thalamic

nucleus does not asymmetrically innervate the left habenula and potential artifacts in

labeling, the source of light-evoked activity in the habenula could not be determined

(Turner et al., 2016). Here, using a combination of imaging, tracing and manipulation, we

demonstrate that light-evoked activity in the habenula of zebrafish larva is mediated by a

thalamic nucleus with retinal input, i.e. the putative nucleus rostrolateralis. We then ask

whether a similar pathway could exist in mammals.

Results

The habenula displays a broad and dynamic response to irradiance change

The zebrafish habenula consists of neurons surrounding neuropils that are

innervated by afferent neurons (Hendricks and Jesuthasan, 2007; Miyasaka et al., 2009;

Amo et al., 2014; Turner et al., 2016). To gain insight into the neural circuits that enable

light to influence the habenula, we first characterized habenula activity evoked by pulses

of light. Two-photon imaging was performed on a transgenic zebrafish line expressing

the calcium indicator GCaMP3 throughout the habenula (Krishnan et al., 2014) (Figure

1A). Resonant-scanning, combined with piezo-driven focusing, was used to record the

activity of cells at multiple focal planes (Figure 1B, C). With a step size of 10 µm, so that

each cell would be sampled only once, most of habenula could be covered with 5 planes

at a rate of 1 Hz. Habenula activity was monitored as the larva was exposed to 20-

second pulses of blue light. We used relatively long pulses, rather than brief flashes, to

allow responses to transition as well as steady state to be identified. Pixel-wise analysis

in one 7-day old fish indicates that evoked activity – both transient and sustained -

occurred throughout the habenula in response to increase and decrease in irradiance

and to light and darkness (Figure 1D, E). The spatio-temporal pattern of activity was

reproducible across several cycles, as shown by the trajectory of the system through

state space (Figure 1F).

To assess if these responses were reproducible across multiple fish, we imaged

the habenula in 6 fish. Habenula neurons were segmented (Figure 2A-C; total of 4986

cells, with an average of 831 ± 53 cells (95% CI) per fish) and their activity was clustered

by *k*-means. Cluster centers were classified by response type. Transient and sustained

responses to increase and decrease in irradiance could be reliably evoked (Figure 2D-G).

The mean percentage of responding cells per fish (±95% CI) were ON: 30.98 ± 9.94%;

OFF: 19.03 ± 3.93%; Inhibitory: 7.93 ± 3.50%. Correlating the cells corresponding to the

different response types revealed that, in general, neurons that were excited by an

increase of irradiance did not fire to a decrease (Figure 2H, I). Some neurons that were

inhibited by light did, however, show a response at light offset (Figure 2F, blue trace; see

also Figure 1E, red trace). These observations confirm that the activity of zebrafish

habenula neurons is affected by change in irradiance, and that in addition to excitation

there is inhibition by light, as well as excitation to loss of light.

The s1011tGAL4 line drives GCaMP3 expression throughout the habenula, and

the response in all focal planes imaged suggests that all domains of the habenula

respond to change in light. To further test this, we imaged fish where specific domains

could be identified. Using the narp promoter (Agetsuma et al., 2010) (Figure 3A-C),

responses to light ON and OFF could be detected in the lateral subdomain of the dorsal

habenula (dHbL) (Figure 3 D, G, H; n = 5 fish). As we lacked a driver that is specific for

the medial subdomain of the dorsal habenula (dHbM), we assessed activity here by

analyzing narp-negative regions of the dorsal habenula in *narp:GAL4, UAS:DsRed,*

elavl3:GCaMP6f fish (Figure 3E, G, I; n = 5 fish). Additionally, we analyzed the

interpeduncular nucleus (IPN), which receives input from all regions of the dorsal

habenula (deCarvalho et al., 2014). Evoked activity was seen in all domains of the

interpeduncular nucleus (Figure 3K, L), consistent with all domains of the dorsal

habenula showing a response to change in irradiance. To assess the ventral habenula,

we imaged a line with a calcium indicator under the control of the *dao* promoter (Amo et

al., 2014). Again, evoked activity was seen (Figure 3F, G, J; n = 8 fish). These results

indicate that all domains of the habenula in larval zebrafish respond to change in

illumination.

To further characterize light evoked activity, we used high speed widefield

microscopy. With 200 Hz imaging, a strong and rapid increase in fluorescence was

detected in a discrete region in the dorsal left habenula (Figure 4A-F; n = 6 fish), as well

as bilaterally in the thalamic region, soon after the onset of visible light. Two-photon

microscopy at 13 Hz suggests that the rapid increase in the habenula occurs in the

dorsal left neuropil (Figure 4G-I; n = 4 fish). The increase suggests that there is

asymmetric activation of the dorsal habenula by light, potentially caused by a stronger

input to the dorsal left neuropil. The bilateral activation in the thalamic region argues

against the habenula asymmetry being an artifact of asymmetric illumination or reporter

expression.

Taken together, these results suggest that the habenula afferents providing

information about illumination should have the following properties. Firstly, they should

terminate broadly within the habenula. Secondly, there should be stronger activity at light

onset in terminals innervating the dorsal left neuropil. Thirdly, they should respond to

increase or decrease in irradiance. Fourthly, they should cause excitation and inhibition,

and may thus include excitatory and inhibitory neurons

The thalamus provides input to the habenula

We next searched for afferents that could provide such properties in larval

zebrafish. We focused initially on inputs that could account for the asymmetry in the

dorsal habenula. The entopeduncular nucleus (EN) is the major source of habenula

afferents in teleosts (Yañez and Anadon, 1994), including zebrafish (Amo et al., 2014;

Turner et al., 2016). This nucleus is labeled in the *Et(sqKR11)* line (Lee et al., 2010),

providing a simple way of visualizing EN afferents to the habenula. Some labeled fibers

were detected in the dorsal left neuropil, indicating that the EN does provide some

innervation to this region of the habenula (Figure 5A). However, labeling in this neuropil

was relatively sparse, compared with other neuropils. We therefore hypothesized that

there may be other inputs, from an anatomically distinct population, that targets this

neuropil. To test this, the lipophilic tracer DiD was injected into the dorsal left neuropil (n

= 6 fish). In all cases, neurons in the dorsal left habenula (which extend dendrites into

the neuropil), the parapineal, and a thalamic nucleus located ventrally to both habenula

(Figure 5B-D) were labeled.

DiD label was not detected in any other regions of the brain, and only rarely in the

entopeduncular nucleus, suggesting that the thalamus is the major source of input to the

dorsal left neuropil. This is consistent with the early activity seen in the thalamus (Figure

4A-F). The label in the thalamus cannot represent anterograde label from the habenula,

as tracing of projections from the habenula by expressing fluorescent proteins

specifically in the habenula does not result in a projection to the thalamus (Movie 1).

Moreover, the labeling of cell bodies in the thalamus (Figure 5D inset, E) indicates that

this is likely to be a retrograde label. The neuropil of this thalamic nucleus contains

terminals of retinal ganglion cells, as shown by DiI injection into the retina (Figure 5E).

Thus, the habenula neuropil with a strong response to light is innervated by thalamic

nuclei that receive retinal input. This is likely to be the nucleus rostrolateralis.

Expressing a fluorescent protein in thalamic neurons, using the *s1020tGAL4* driver

(Figure 5F, G), led to label of terminals throughout the habenula (Movie 2). Thus,

although the dorsal left neuropil receives input from the thalamus, the thalamus

projection is not restricted to this region of the habenula.

The habenula may receive glutamatergic and GABAergic input from the thalamus

As noted above, light caused both increase and decrease in activity of habenula

neurons, implying that there may be excitatory and inhibitory afferents. Using an

antibody to vGlut1/2, glutamatergic pre-synapses were detected in all neuropils of the

habenula (Figure 5H), indicating the existence of excitatory afferents. GAD65/67 labeled

puncta could also be detected in habenula neuropils (Figure 5I). In the lateral regions,

corresponding to the ventral habenula, labeled streaks were detected adjacent to cell

bodies. These labels were not located within habenula neurons, as they did not co-

localize with cytoplasmic label provided by GCaMP3, nor did they fill the cytoplasm,

implying that these puncta and streaks must reside in habenula afferents (i.e. axon

terminals) such as those labeled in Movie 2. Labeled cell bodies were seen below the

level of the habenula (see Movie 3). Consistent with this, GABAergic neurons could be

detected in the dorsal thalamus using the transgenic line *Tg(gad1b:RFP, vGlut2a:GAL4,*

UAS:eGFP) (Satou et al., 2013) (Figure 5J, K; Movie 4). No label was seen in the

entopeduncular nucleus, which has previously been shown to be glutamatergic (Amo et

al., 2014; Turner et al., 2016). These observations confirm that the thalamus contains

both glutamatergic and GABAergic neurons, as described previously (Mueller, 2012),

which may mediate light-evoked excitation and inhibition of habenula neurons.

Optogenetic stimulation of the thalamus drives habenula activity

The lipophilic and transgenic tracing experiments documented in Figure 5

demonstrate anatomical connectivity from the thalamus to the habenula. To test

functional connectivity, we used optogenetics. Channelrhodopsin-2 (ChR2) was

expressed in thalamic neurons with the s1020tGAL4 driver (Fig. 6A). Experiments were

carried out on fish lacking eyes, to prevent a response to visual stimulation. Short pulses

of blue light reproducibly caused an increase in fluorescence of GCaMP6f in habenula

neurons of fish with expression of ChR2 in the thalamus (Figure 6B, D, G). Some

response was seen in fish without ChR2 expression (Fig. 6F), suggesting that some

habenula response may be due to non-ocular sensors such as deep brain

photoreceptors (Matos-Cruz et al., 2011; Fernandes et al., 2013). The larger response in

fish with ChR2 expression, however, suggests that there is functional connectivity

between the thalamus and the habenula.

Irradiance change evokes activity in thalamus neurons

If the thalamus provides afferents mediating illumination-dependent activity in the

habenula, thalamic neurons should respond to increase and decrease of illumination. To

test this, calcium imaging was carried out in *s1020tGAL4, UAS:GCaMP6s* transgenic

fish. A response to increase and decrease in illumination was detected in cell bodies in

the anterior thalamus (Figure 7A-F) in all fish imaged (n = 5). Dendrites in the neuropil of

the thalamus also responded to change in illumination. Increase in irradiance caused

activity more dorsally, while decrease caused activity more ventrally (Figure 7G-J). Thus,

the thalamus has a response to both increase and decrease of illumination.

Habenula response to light has been shown to depend on the eyes (Dreosti et al.,

2014). Thus, if the thalamus mediates habenula response to light, light-evoked activity

here should depend on the eyes. Indeed, the robust responses to light were lost in fish

lacking eyes (Figure 7K-S), consistent with this hypothesis.

Functional asymmetry in light-evoked activity

As shown in Figure 4G-I, light evokes strong activity in the dorsal neuropil of the

left habenula. As the line used in this imaging experiment contains only labelled

habenula neurons, the response probabaly occur in dendrites. Given that there is no

obvious anatomical asymmetry in thalamic input to the habenula (Figure 5F, G; see also

Turner et al, 2016), we hypothesized that there may be functional asymmetry in activity

within thalamic inputs. To test this, we imaged the terminals of thalamic axons using

s1020tGAL4, UAS:GCaMP6s fish. Thalamic terminals in the dorsal left neropil showed a

greater activity compared to those in the right neuropil (Figure 8). Thus, preferential light-

evoked activity in thalamic afferents that project to the dorsal left neuropil may underlie

the asymmetric dorsal habenula response.

Thalamic lesion inhibits habenula response to illumination change

To further test if the thalamus contributes to light evoked activity in the habenula,

we lesioned the thalamic neuropil with a two-photon laser. This technique is expected to

injure fibers innervating the neuropil (Semmelhack et al., 2014). The laser was targeted

to the neuropil of the putative nucleus rostrolateralis, which was identified by first

imaging the response to light pulses (Figure 9A, B). Lesioning led visible damage in the

neuropil (Figure 9D), and to a reduction of evoked activity in the thalamus and habenula

(Figure 9C, E-G, I, L, M). There was some variability in the effect (Figure 9N), possibly

reflecting the limitations of this technique in enabling consistent ablation. Lesioning

other targets of retinal ganglion cell axons, which are located more posteriorly, did not

lead to a loss of light-evoked activity in the habenula (Figure 9G, I, J, K), indicating that

this technique did not cause indiscriminate damage. These observations support the

hypothesis that the putative nucleus rostrolateralis of the thalamus has a role in light-

evoked activity in the habenula of larval zebrafish.

A thalamo-habenula projection in the mouse

Finally, we asked whether a projection from the thalamus to the habenula,

239 especially from a visual nucleus, is restricted to zebrafish, or whether it could also exist

240 in a mammal. To do this, we examined the mouse mesoscale connectome (Oh et al.,

241 2014). A search of the connectivity database derived from AAV injections into p56 mice,

242 using thalamus as the injection site and epithalamus as the target site, yielded 18 hits

243 covering different thalamic nuclei. Three experiments (numbers 267538006, 525796603

244 and 147212977) had been targeted to the ventral lateral geniculate nucleus. In one case

245 (Figure 10), the anterograde tracer virus had been injected into a Slc32a1-IRES-Cre

246 mouse, which expresses Cre in GABAergic neurons. Viral tracing in the mouse thus

247 suggests that a visual nucleus in the thalamus projects to the habenula in a mammal.

248 ## Discussion

249 We have investigated how illumination conditions influence activity in the habenula

250 of larval zebrafish. The pineal, although light responsive, does not innervate the

251 zebrafish habenula (Yáñez et al., 2009) while the parapineal, which innervates the

252 habenula, has been shown to be dispensable (Dreosti et al., 2014). Calcium imaging

253 suggests that the afferent neurons mediating responses to light should terminate broadly

254 in the habenula, cause stronger activity in the dorsal left neuropil, depolarize to increase

255 or decrease in irradiance, and potentially include excitatory and inhibitory neurons.

256 Several observations suggest that afferent neurons with these properties reside in the

257 thalamus.

258 Lipophilic tracing of the habenula and transgenic labeling of thalamic neurons

259 demonstrate that the thalamus directly innervates the larval zebrafish habenula. High-

260 speed imaging with widefield microscopy suggests that there is correlated activity in the

261 thalamus and habenula, consistent with functional connectivity. Although widefield

262 imaging uses visible light to excite the reporter, the rate of imaging used here is faster

263 than the rise time of GCaMP6f (Chen et al., 2013), so initial images reflect activity prior

to onset of the stimulating light. Further evidence for functional connectivity is provided

by optogenetic stimulation of the thalamus, which causes a response in the habenula.

Moreover, lesion of the thalamus reduced light-evoked activity in the habenula. The

thalamus responds to both increase and decrease in illumination, and contains both

excitatory and inhibitory neurons; no other source of GABAergic inputs to the zebrafish

habenula has been described. Thus, by optical recording, anatomical tracing, activation

and lesion, our data suggests that the thalamus mediates the habenula responses to

irradiance change in larval zebrafish.

The region of the thalamus mediating activity in the habenula appears to be the

nucleus rostrolateralis, as proposed by Turner et al (2016). The neuropil here can be

functionally separated into two domains, based on the response to light – excitation to

light OFF in the ventral regions and excitation to light ON more dorsally. This neuropil

contains two previously defined targets of retinal ganglion cells, AF2 and AF4 (Burrill and

Easter, 1994). AF4 is innervated predominantly by M3 and M4 retinal ganglion cells

(Robles et al., 2014), which extend their dendritic tree into the proximal layer of the inner

plexiform layer and are considered ON neurons. AF2 is innervated by B1 retinal ganglion

cells that have dendrites in the distal layer (Robles et al., 2014), and these may account

for the OFF responses in the thalamus and habenula. This thalamic nucleus may also

receive input from non-retinal sources, but this remains to be investigated. The loss of a

thalamic response to light in fish lacking eyes, however, suggests that such contribution

may be minor.

Light is a potent regulator of brain function. It can affect mood (Vandewalle et al.,

2010), alertness (Badia et al., 1991), cognitive ability (LeGates et al., 2012) and

movement (Aschoff, 1960; Burgess and Granato, 2007). These phenomena are

sensitive to irradiance, not image formation, and are mediated by a number of sensors

including intrinsically-sensitive retinal ganglion cells whose targets include the thalamus

(Hattar et al., 2006). The ability of light to affect normal movement patterns (Burgess et

al., 2010) or to disrupt mood and cognition (LeGates et al., 2012) involves

neuromodulators such as serotonin, and changing irradiance affects activity in the dorsal

raphe (Fite et al., 2005; Cheng et al., 2016). Based on the data here, and the well-

established roles of the habenula in regulating neuromodulators, we suggest that some

of these effects of light may be mediated by the thalamic projection to the habenula.

A projection from the thalamus to the habenula may be evolutionarily conserved in

vertebrates. In humans and rabbits, a thalamo-habenula projection was proposed many

years ago based on degeneration experiments (Marburg, 1944; Cragg, 1961). Using

retrograde tracing with horseradish peroxidase, a projection from the dorsal thalamus to

the habenula was reported in a lizard (Díaz and Puelles, 1992). Hints of a projection can

also be seen in a tracing experiment performed in rats (Moore et al., 2000). The large-

scale mouse mesoscale connectome project (Oh et al., 2014), which uses viral-based

anterograde tracing, provides the most recent evidence for a thalamo-habenula

projection in a mammal. This technique suggests a projection from the ventral lateral

geniculate nucleus and other thalamic nuclei to the habenula. It will be interesting to

determine whether a similar anatomical connection exists in humans, as this may

contribute to the high functional connectivity between the thalamus and habenula

reported recently (Torrisi et al., 2016).

Materials and Methods

Fish lines

Experiments were performed in accordance with guidelines issued by the

Institutional Animal Care and Use Committee of the Biological Resource Centre at

Biopolis, Singapore. Zebrafish (*Danio rerio*) lines used for this study were:

Tg(UAS:GCaMP6s)sq205, SqKR11Et (Lee et al., 2010), *GAL4s1011t* (Scott and Baier,

2009), *GAL4s1020t* (Scott and Baier, 2009)*, Tg(UAS:GCaMP3)sq200*,

Tg(elavl3:GCaMP6f)a12200, *Tg(UAS:ChR2-eYFP)* (Arrenberg et al., 2009) and AB

wildtype.

Tg(elavl3:GCaMP6f)a12200 was generated by PCR amplification of the GCaMP6f

open reading frame (Addgene plasmid 40755 (Chen et al., 2013)) with forward primer

ataACTAGTgccaccATGGGTTCTCATCATCAT   and   reverse

ataCCGCGGcTCACTTCGCTGTCATCATTTGTAC   (restriction   site   and   coding

sequences are in upper case). This fragment was cloned into a plasmid with Tol2 arms

flanking an upstream attR1-R2 cassette and the insertion site using restriction enzymes

SpeI and SacII. Previously described *elavl3* (*HuC*) *cis*-regulatory elements (Higashijima

et al., 2003) were placed upstream via LR recombination (Invitrogen) with an attL flanked

elavl3 entry clone. The resulting plasmid was then co-injected into 1-cell stage embryos

at a concentration of 30 ng/µL with Tol2 transposase mRNA at a concentration of 30

ng/µL. A single founder was selected based on high and spatially broad expression.

Outcrossing this founder generated 50% GCaMP6f-positive embryos, which were

selected to establish the line.

Imaging

Zebrafish larvae (aged 5 - 10 dpf) were anaesthetized in mivacurium and

embedded in low-melting temperature agarose (1.2-2.0 % in E3) in a glass-bottom dish

(Mat Tek). They were imaged on a Nikon two-photon microscope (A1RMP), attached to

a fixed stage upright microscope, using a 25x water immersion objective (NA = 1.1). The

femtosecond laser (Coherent Vision II) was tuned to 920 nm for GCaMP imaging. Stacks

337 were collected in resonant-scanning mode with a 525/50 nm bandpass emission filter

338 and with 8x pixel averaging; single-plane images were collected in galvano-scanning

339 mode with 2x pixel averaging. The sample size was based on (Dreosti et al., 2014).

340 Light stimuli were generated by 5 mm blue LEDs (458 nm peak emission). They

341 were powered by a 5 V TTL signal from a control computer and synchronized with image

342 capture using a National Instruments DAQ board, controlled by the Nikon Elements

343 software. Light intensity at the sample was 0.13 mW/cm^2.

344 For widefield microscopy, excitation was provided by LEDs (Cairn OptoLED) at

345 470 nm. Images were captured on a Zeiss Axio Examiner with a 20x water immersion

346 objective, using a Flash4 camera (Hamamatsu) controlled by MetaMorph. After

347 background subtraction, change in fluorescence was measured using the deltaF-up

348 command in Fiji.

349 Data analysis

350 **Initial Data Preprocessing:** Raw images obtained were first registered to correct

351 for any vertical/horizontal movement artifacts using cross correlation. Then, a median

352 filter of size 3 was applied to remove noise. A darker region outside the region of

353 interest was chosen as the background and subtracted from the image to remove any

354 background noise. Non linear trends in the data were detrended using polynomials of

355 order 2-5. Data was then normalized into Z-scores by subtracting the overall mean and

356 dividing by the standard deviation. A rolling window average was then used to smooth

357 noisy traces where necessary. Where possible, cells were segmented or images were

358 directly analysed as pixels (see Below).

359 **Correlation to light evoked activity:** Temporal traces from pixels (Thalamus or

Thalamic afferents) or segmented cells (in the habenula) were classified as responding to light ON or OFF by calculating their correlation coefficient to a square wave that is 1 when the light is ON or when light is OFF and 0 during other time periods. High correlation to these traces indicated that the pixel or cell is responding to light ON or OFF respectively.

Pixel based analysis of the habenula and thalamus: To show the spatial and temporal distribution of light evoked activity in the habenula and thalamus, the Thunder platform (Freeman et al., 2014) was used for fast pixel based clustering and factorization.

PCA: Principal Component Analysis (PCA) was used to obtain a low dimensional representation of the population. In normal fish, in the habenula (Figure 1D) and thalamus (Figure 7M), the first two Principal Components (PC) easily picked up the evoked responses to light ON and OFF. The explained variance, though small, decayed rapidly after the first component (Figure 1D, variance explained PC1 12.92%, PC2 6.12% and PC3 4.57%; first 20 PCs 55.60% ; Figure 7M, variance explained PC1 11.87%, PC2 3.61% and PC3 2.67%; first 20 PCs 40.82%) and hence the first two PCs were chosen to plot the data in low dimension. They showed reproducible state changes in the population to changes in irradiance. In eye lesioned fish (Figure 7P) however, neither of the first twenty PCs (variance explained PC1 4.92%, PC2 4.24% and PC3 3.26%; first 20 PCs 39.53%) showed any discernable correlation to light evoked activity (The inset in Figure 7P). The first two are plotted in Figure 7P for comparison with controls. Correlation coefficients plotted in insets of Figure 7M and 7P were obtained by correlating the PCs with a trace that was 1 when light was ON and 0 otherwise.

K-means: K-means clustering was performed to identify pixels with similar responses profiles. Given the uncertainty of k-means to the optimal cluster number, an

iterative approach was used to separate pixels relating to evoked responses versus

pixels that don't (here referred to as noise clusters). The number of clusters were chosen

to reveal as many stimulus related clusters as possible, until there was little change in

the number and types of stimulus related clusters and increase in noise related clusters.

Noise clusters were then removed from the spatial and temporal plots for clarity. It is to

be noted that this analysis does have its drawbacks when used to analyse data with

variable variance and cannot not capture all possible types of responses. Multiple runs

were made to ensure that noisy pixels were not clustered into evoked clusters. Where

applicable cell segmentation and manual classification was used. For example, we were

not able to separate inhibitory clusters from off custers in Figure 3E-G. This was

revealed when the cells were segmented, and their temporal traces inspected. In all

cases, K-means cluster center showing evoked responses to light ON were colored in

shades of blue and those showing responses to light OFF were colored in shades of red.

Pixels belonging to the cluster were colored similarly and superimposed on an average

image of the plane analysed. In different datasets, on average, this analysis provided an

optimal k of 6-10; 3-4 clusters that didn't correspond to evoked activity were not included

while plotting.

For data following eye lesion (FIgure 7) and thalamic neuropil/AF9/AF7 lesion

(Figure 9), k-means was performed to differentiate responses between the controls and

the lesion. Hence, number of clusters were chosen such that cluster center adequately

differentiated responses before and after lesion. Cluster center not responding to light

were also plotted here. To verify the results of k-means, the presence of light response

was verified by examining the pixels (Figure 8 Q-S) or segmenting the cells (Figure 9 J-

N) in both control and lesioned animals.

The scripts used for analysis are provided at http://dx.doi.org/10.5061/dryad.q0171.

Cell segmentation: Each stack was scaled 2x in imageJ (RRID:SCR_003070),

then maximally projected to a single image, which was then subjected to a minimum filter

and unsharp mask to sharpen the boundary of cells. ROIs were identified using the "find

maxima…" command, as a way to localize regional darkest point as the center of each

ROI. The boundary of the ROI was outlined by "analyze particle…" that connects bright

pixels into mosaic-like tessellated plane, encircling each darkest point. Each ROI was

then numbered sequentially using the ImageJ ROI Manager and mapped back to the

original despeckled image stack. Manual segmentation was done here to delete

extraneous ROIs outside the habenula and to encircle cells that were not detected by

the algorithm (<10% of total ROIs). In the last step, "Set measurements…" and

"measure" in ImageJ provided the mean fluorescence value of all pixels within each ROI

across the entire image stack and the x-y coordinates of each ROI. Time-lapse series in

which z drifting occurred were excluded, as in this case ROIs could not be defined.

K-means on segmented cells: For Figures 2D-F, *k-means* was performed from

cells segmented by the semi automated algorithm described above. The purpose is to

determine heterogeneity of temporal responses to changes in irradiance, accurately

classify cells into ON, OFF and Inhibitory responses and perform correlation between

them. Analysis was done on traces from 4986 habenula cells from 6 fish. Traces were

detrended, smoothed and normalized to z-scores using baseline as the time before the

first blue light. Traces that did not reach a Z-score of 2 during the period of irradiance

change were classified as not having an evoked response and not included in the

clustering analysis. 2456 of 4986 cells were thus removed. *K-means* was first run with an

arbitrary *k* = 60. This generated a wide range of clusters capturing the temporal

heterogeneity of the responses. The clusters were then divided into ON, OFF, Inhibitory

and no evoked response. Neurons belonging to each cluster were correlated among

each other and to the cluster center. If they had an evoked response, they were

correlated with other clusters and assigned to one with the highest correlation. Otherwise

they were classified into the no evoked response category; 138 such cells were

reclassified. Traces of cells belonging to ON, OFF and Inhibitory clusters are plotted as a

heatmap in Figure 2G and their correlation coefficients in Figure 2H. Similarly, cells

segmented from multiple fish were classified to reveal responses in different habenula

regions as plotted in Figure 3H-J.

Boxplots: Boxplots in Figures 7Q, 8D-E and 9N were plotted to show the full

distribution of the data. The box in the boxplot ranges from the first quartile to the third

quartile, and the box shows the interquartile range (IQR). The line across the box is the

median of the data. The whiskers extend to 1.5*IQR on either side of the box. Anything

above this range are defined as outliers and plotted as black diamonds in the plots.

Neural tracing

DiD (Life Technologies) was dissolved in 50 µl ethanol to make a saturated

solution. This was heated to 55˚C for 5 minutes prior to injection into the fish that had

been fixed in 4% paraformaldehyde. Fish were mounted in 1.2% low melting

temperature agarose dissolved in PBS. The dye was pressure injected into the habenula

under a compound microscope (Leica DM LFS), using a 20X water immersion objective.

For labeling the retina, a saturated solution of DiI in chloroform was used. Injections

were carried out under a stereo microscope (Zeiss Stemi 2000). After injections, fish

were stored at 4˚C overnight to allow tracing, and then imaged with a 40x water

immersion objective on a Zeiss LSM 710 confocal microscope.

 Antibody label

Larvae were fixed in 4% para-formaldehyde/PBS overnight at 4˚C. They were then

rinsed in PBS. The brains were dissected out, and permeabalized using 1% BSA

(fraction V; Sigma), 0.1% DMSO and 0.1% Triton X-100. The antibodies used here, anti-

vGlut1/2 (Synaptic Systems 135503, RRID:AB_1279466; 1:100) and anti-GAD65/67

(Abcam ab11070, RRID:AB_297722; 1:500), have previously been used in zebrafish

(Wyart et al., 2009; Lee et al., 2010). The brains were incubated in the primary antibody

overnight, rinsed several times in PBS, then incubated in secondary antibody (Alexa 488

goat anti-rabbit; 1:1000). After washing, these were mounted in 1.2% agarose/PBS.

Imaging was carried out using a Zeiss LSM 510 laser scanning confocal microscope,

with a 40x water immersion objective.

Optogenetic stimulation

5 dpf *Tg(s1020GAL4, UAS:ChR2-eYFP, elavl3:GCaMP6f)* larvae were used. The

eyes were removed using fine tungsten needles in fish that were anesthetized with

MS222. This procedure was carried out in Ringers saline. Fish were then mounted in

1.2% agarose in Ringers saline, and imaged using two-photon microscopy as described

above, at 1 Hz. Optical stimulation was carried out using a 50 μm fiber optic probe (Doric

Lenses), placed approximately 20 μm from the thalamus. The 465 nm LED (Doric) was

driven with a current of 900 mA, 30 seconds after the start of imaging. 10 pulses were

provided, with a pulse duration of 25 milliseconds and a frequency between 1 and 8 Hz.

Each fish was exposed to at least 3 pulse trains. For Figure 6B-C, the average of the first

29 frames was used as a reference. The ratio of all frames relative to this reference was

obtained using FIJI (RRID:SCR_002285). The analysis to generate Figure 6G was blind

to the genotype.

 Enucleation

    5 day-old fish were anaesthetized in Ringer's saline containing MS222. The eyes

 were removed using electrolytically sharpened tungsten needles. Fish were allowed to

 recover for several hours in anesthetic-free saline. Activity recorded 2 - 4 hours after eye

 removal. To enable lateral imaging of the thalamus (Figure 7H-J), one eye was removed

 using this method.

 Laser ablation

    Tg(Elavl3:GCaMP6f) larvae were anaesthetized and then mounted in 2% low-

 melting temperature agarose. Lesions were created with the femto-second laser tuned to

 960 nm and fixed on a single point. Several pulses, each lasting 100 - 500 msec, were

 used. Lesioning was monitored by time-lapse imaging before and after each pulse, and

 was terminated when there was a localized increase in GCaMP6f fluorescence. The

 formation of a bubble indicates that lesioning involved plasma formation (Venugopalan

 et al., 2002). These can be imaged in the red channel of the two-photon microscope.

 Sample size was chosen based on (Aizenberg and Schuman, 2011). Animals with

 bleeding in the brain after lesioning, due to bursting of a blood vessel in the thalamis,

 were discarded.

 Analysis of the mouse mesoscale connectome

    The mouse connectivity atlas (http://connectivity.brain-map.org/) was searched in

 ""Target Search" mode, using "TH" as the source structure and "EPI" as the target

 structure. The minimum target volume listed was 0.005 mm^3. For each hit, serial

 transverse sections containing the habenula, which had been imaged by two-photon

 microscopy, were screened manually to verify that there were labelled fibers within the

503   habenula.

Acknowledgements

505   This work was funded by core funding from the Institute of Molecular and Cell

506   Biology and a grant from the Ministry of Education, Singapore. SK and QL were

507   supported by fellowships from the National University of Singapore. DGCH was

508   supported by NIH grant 5T32HL007901. We thank Claire Wyart for providing the

509   *UAS:ChR2-eYFP* transgenic line.

References

Agetsuma M, Aizawa H, Aoki T, Nakayama R, Takahoko M, Goto M, Sassa T, Amo R,

Shiraki T, Kawakami K, Hosoya T, Higashijima S-I, Okamoto H (2010) The habenula

is crucial for experience-dependent modification of fear responses in zebrafish. Nat

Neurosci 13:1354–1356.

Aizawa H, Yanagihara S, Kobayashi M, Niisato K, Takekawa T, Harukuni R, McHugh TJ,

Fukai T, Isomura Y, Okamoto H (2013) The synchronous activity of lateral habenular

neurons is essential for regulating hippocampal theta oscillation. J Neurosci

33:8909–8921.

Aizenberg M, Schuman EM (2011) Cerebellar-Dependent Learning in Larval Zebrafish. J

Neurosci 31:8708–8712.

Amo R et al. (2014) The habenulo-raphe serotonergic circuit encodes an aversive

expectation value essential for adaptive active avoidance of danger. Neuron

84:1034–1048.

Appelbaum L, Wang GX, Maro GS, Mori R, Tovin A, Marin W, Yokogawa T, Kawakami K,

Smith SJ, Gothilf Y, Mignot E, Mourrain P (2009) Sleep-wake regulation and

hypocretin-melatonin interaction in zebrafish. Proceedings of the National Academy

of Sciences 106:21942–21947.

Arrenberg A, Arrenberg AB, Del Bene F, Del Bene F, Baier H, Baier H (2009) Optical

control of zebrafish behavior with halorhodopsin. Proceedings of the National

Academy of Sciences 106:17968–17973.

Aschoff J (1960) Exogenous and endogenous components in circadian rhythms. Cold

Spring Harb Symp Quant Biol 25:11–28.

Badia P, Myers B, Boecker M, Culpepper J, Harsh JR (1991) Bright light effects on body

temperature, alertness, EEG and behavior. Physiol Behav 50:583–588.

Burgess HA, Granato M (2007) Modulation of locomotor activity in larval zebrafish during

light adaptation. J Exp Biol 210:2526–2539.

Burgess HA, Schoch H, Granato M (2010) Distinct retinal pathways drive spatial

orientation behaviors in zebrafish navigation. Curr Biol 20:381–386.

Burrill JD, Easter SS (1994) Development of the retinofugal projections in the embryonic

and larval zebrafish (Brachydanio rerio). J Comp Neurol 346:583–600.

Butler AB, Saidel WM (2003) Clustered phylogenetic distribution of nucleus

rostrolateralis among ray-finned fishes. Brain Behav Evol 62:152–167.

Chen T-W, Wardill TJ, Sun Y, Pulver SR, Renninger SL, Baohan A, Schreiter ER, Kerr

RA, Orger MB, Jayaraman V, Looger LL, Svoboda K, Kim DS (2013) Ultrasensitive

fluorescent proteins for imaging neuronal activity. Nature 499:295–300.

Cheng R-K, Krishnan S, Jesuthasan S (2016) Activation and inhibition of tph2

serotonergic neurons operate in tandem to influence larval zebrafish preference for

light over darkness. Sci Rep 6:20788.

Chou M-Y, Amo R, Kinoshita M, Cherng B-W, Shimazaki H, Agetsuma M, Shiraki T, Aoki

T, Takahoko M, Yamazaki M, Higashijima S-I, Okamoto H (2016) Social conflict

resolution regulated by two dorsal habenular subregions in zebrafish. Science

352:87–90.

Cragg BG (1961) The connections of the habenula in the rabbit. Exp Neurol 3:388–409.

deCarvalho TN, Subedi A, Rock J, Harfe BD, Thisse C, Thisse B, Halpern ME, Hong E

(2014) Neurotransmitter map of the asymmetric dorsal habenular nuclei of zebrafish.

Genesis 52:636–655.

Díaz C, Puelles L (1992) Afferent connections of the habenular complex in the lizard

Gallotia galloti. Brain Behav Evol.

Dreosti E, Vendrell-Llopis N, Carl M, Yaksi E, Wilson SW (2014) Left-right asymmetry is

required for the habenulae to respond to both visual and olfactory stimuli. Curr Biol

24:440–445.

Fernandes AM, Fero K, Driever W, Burgess HA (2013) Enlightening the brain: Linking

deep brain photoreception with behavior and physiology. Bioessays 35:775–779.

Fite KV, Wu PS, Bellemer A (2005) Photostimulation alters c-Fos expression in the

dorsal raphe nucleus. Brain Res 1031:245–252.

Fowler CD, Lu Q, Johnson PM, Marks MJ, Kenny PJ (2011) Habenular α5 nicotinic

receptor subunit signalling controls nicotine intake. Nature 471:597–601.

Golden SA et al. (2016) Basal forebrain projections to the lateral habenula modulate
aggression reward. Nature 534:688–692.

Hattar S, Kumar M, Park A, Tong P, Tung J, Yau K-W, Berson DM (2006) Central
projections of melanopsin-expressing retinal ganglion cells in the mouse. J Comp
Neurol 497:326–349.

Hendricks M, Jesuthasan S (2007) Asymmetric innervation of the habenula in zebrafish.
J Comp Neurol 502:611–619.

Higashijima S-I, Masino MA, Mandel G, Fetcho JR (2003) Imaging neuronal activity
during zebrafish behavior with a genetically encoded calcium indicator. J
Neurophysiol 90:3986–3997.

Hong S, Hikosaka O (2008) The globus pallidus sends reward-related signals to the
lateral habenula. Neuron 60:720–729.

Jhou TC, Fields HL, Baxter MG, Saper CB, Holland PC (2009) The rostromedial
tegmental nucleus (RMTg), a GABAergic afferent to midbrain dopamine neurons,
encodes aversive stimuli and inhibits motor responses. Neuron 61:786–800.

Krishnan S, Mathuru AS, Kibat C, Rahman M, Lupton CE, Stewart J, Claridge-Chang A,
Yen S-C, Jesuthasan S (2014) The Right Dorsal Habenula Limits Attraction to an
Odor in Zebrafish. Curr Biol.

Lee A, Mathuru AS, Teh C, Kibat C, Korzh V, Penney TB, Jesuthasan S (2010) The
habenula prevents helpless behavior in larval zebrafish. Curr Biol 20:2211–2216.

LeGates TA, Altimus CM, Wang H, Lee H-K, Yang S, Zhao H, Kirkwood A, Weber ET,
Hattar S (2012) Aberrant light directly impairs mood and learning through

melanopsin-expressing neurons. Nature 491:594–598.

Marburg O (1944) The structure and fiber connections of the human habenula. Journal

of Comparative Neurology 80:211–233.

Matos-Cruz V, Blasic J, Nickle B, Robinson PR, Hattar S, Halpern ME (2011)

Unexpected diversity and photoperiod dependence of the zebrafish melanopsin

system. PLoS ONE 6:e25111.

Matsumoto M, Hikosaka O (2007) Lateral habenula as a source of negative reward

signals in dopamine neurons. Nature 447:1111–1115.

Matsumoto M, Hikosaka O (2009) Representation of negative motivational value in the

primate lateral habenula. Nat Neurosci 12:77–84.

Miyasaka N, Morimoto K, Tsubokawa T, Higashijima S-I, Okamoto H, Yoshihara Y

(2009) From the olfactory bulb to higher brain centers: genetic visualization of

secondary olfactory pathways in zebrafish. J Neurosci 29:4756–4767.

Moore RY, Weis R, Moga MM (2000) Efferent projections of the intergeniculate leaflet

and the ventral lateral geniculate nucleus in the rat. J Comp Neurol 420:398–418.

Morley BJ, Spangler KM, Javel E (1985) The development of somatostatin

immunoreactivity in the interpeduncular nucleus of the cat. Brain Res 352:241–248.

Mueller T (2012) What is the Thalamus in Zebrafish? Front Neurosci 6:64.

Oh SW et al. (2014) A mesoscale connectome of the mouse brain. Nature 508:207–214.

Quina LA, Tempest L, Ng L, Harris JA, Ferguson S, Jhou TC, Turner EE (2014) Efferent

Pathways of the Mouse Lateral Habenula. J Comp Neurol.

Robles E, Laurell E, Baier H (2014) The retinal projectome reveals brain-area-specific

visual representations generated by ganglion cell diversity. Curr Biol 24:2085–2096.

Saidel WM (2013) Nucleus Rostrolateralis: An Expansion of the Epithalamus in Some

Actinopterygii. The Anatomical Record 296:1594–1602.

Satou C, Kimura Y, Hirata H, Suster ML, Kawakami K, Higashijima S-I (2013)

Transgenic tools to characterize neuronal properties of discrete populations of

zebrafish neurons. Development 140:3927–3931.

Scott EK, Baier H (2009) The cellular architecture of the larval zebrafish tectum, as

revealed by gal4 enhancer trap lines. Front Neural Circuits 3:13.

Semm P, Demaine C (1984) Electrophysiology of the pigeon's habenular nuclei:

evidence for pineal connections and input from the visual system. Brain Res Bull

12:115–121.

Semmelhack JL, Donovan JC, Thiele TR, Kuehn E, Laurell E, Baier H (2014) A

dedicated visual pathway for prey detection in larval zebrafish. Elife 3:17968.

Stephenson-Jones M, Floros O, Robertson B, Grillner S (2012) Evolutionary

conservation of the habenular nuclei and their circuitry controlling the dopamine and

5-hydroxytryptophan (5-HT) systems. Proc Natl Acad Sci USA 109:E164–E173.

Thornton EW, Davies C (1991) A water-maze discrimination learning deficit in the rat

following lesion of the habenula. Physiol Behav 49:819–822.

Torrisi S, Nord CL, Balderston NL, Roiser JP, Grillon C, Ernst M (2016) Resting State

Connectivity of the Human Habenula at Ultra-High Field. Neuroimage:1–21.

Turner KJ, Hawkins TA, Yáñez J, Anadón R, Wilson SW, Folgueira M (2016) Afferent

Connectivity of the Zebrafish Habenulae. Front Neural Circuits 10:3512.

Vandewalle G, Schwartz S, Grandjean D, Wuillaume C, Balteau E, Degueldre C,

Schabus M, Phillips C, Luxen A, Dijk DJ, Maquet P (2010) Spectral quality of light

modulates emotional brain responses in humans. Proceedings of the National

Academy of Sciences 107:19549–19554.

Venugopalan V, Guerra A, Nahen K, Vogel A (2002) Role of Laser-Induced Plasma

Formation in Pulsed Cellular Microsurgery and Micromanipulation. Phys Rev Lett

88:078103.

Wang RY, Aghajanian GK (1977) Physiological evidence for habenula as major link

between forebrain and midbrain raphe. Science 197:89–91.

Wyart C, Del Bene F, Warp E, Scott EK, Trauner D, Baier H, Isacoff EY (2009)

Optogenetic dissection of a behavioural module in the vertebrate spinal cord. Nature

461:407–410.

Yáñez J, Busch J, Anadón R, Meissl H (2009) Pineal projections in the zebrafish (Danio

rerio): overlap with retinal and cerebellar projections. Neuroscience 164:1712–1720.

Yañez J, Anadon R (1994) Afferent and efferent connections of the habenula in the

larval sea lamprey (Petromyzon marinus L.): an experimental study. J Comp Neurol

345:148–160.

Zhang J, Tan L, Ren Y, Liang J, Lin R, Feng Q, Zhou J, Hu F, Ren J, Wei C, Yu T,

Zhuang Y, Bettler B, Wang F, Luo M (2016) Presynaptic Excitation via GABAB

Receptors in Habenula Cholinergic Neurons Regulates Fear Memory Expression.

Cell 166:716–728.

Zhao H, Rusak B (2005) Circadian firing-rate rhythms and light responses of rat

habenular nucleus neurons in vivo and in vitro. Neuroscience 132:519–528.

Figure legends

Figure 1. The larval zebrafish habenula has a broad and complex response to

change in irradiance. (A) Dorsal view of the head of a live 7 day-old fish, with GCaMP3

expression in the habenula (arrows) under the control of the *s1011t* GAL4 driver. (B) A

single two-photon slice through the dorsal habenula of the fish in panel A (boxed region).

(C) A yz reconstruction at the point indicated by the yellow line in panel B, showing a

transverse view of the habenula. The dotted lines indicate imaging planes separated by

10 µm. The yellow line indicates the plane imaged in B. Dashed lines show the border

of the habenula. (D) Spatial distribution of responses in the habenula of one fish (7 dpf)

to pulses of light. 5 planes are shown here. The colors are coded according to the

temporal pattern of response, as indicated in (E). Images were collected at a rate of 1

stack/second, and four pulses of light were delivered for 20 seconds each, with variable

inter-stimulus interval. (E) Centers of *k*-means clusters corresponding to colors of pixels

in (D). Cluster centers in (D) and the corresponding pixels in (E) indicating responses to

light ON are colored in shades of blue and light OFF in shades of red. The horizontal

black line represents Z-score of 0. (F) Trajectory of the habenula response in two-

dimensional state space, using the first two principal components (PC1 and PC2).

Traces are color-coded according to the wedges in panel E, to represent direction in

which change in irradiance drives the neural state. In panels E and F, the bold lines

correspond to light onset while the dashed lines indicate offset. The presence of light is

also indicated by the blue bars. lHb: left habenula; rHb: right habenula. a: anterior; p:

posterior. d: dorsal; v: ventral Scale bar = 100 µm in panel A, 25 µm elsewhere.

Figure 2. Habenula response to irradiance change is reproducible. (A-C)

Segmentation of habenula neurons using a semi-automated algorithm (see Methods).

(D-F) *K*-means cluster analysis of segmented habenula neuron responses to pulses of

blue light in 6 fish (7-8 dpf). Traces show cluster center, with shaded regions indicating

standard error of the mean. Clusters were grouped by their temporal activity pattern and

clusters with excitation to light ON (D) or OFF (E), or inhibition to light (Inhibitory, F) were

seen. Clusters without evoked responses are not shown. (G) Activity traces of each cell

from the 6 fish grouped into ON, OFF and Inhibitory (Inh) categories based on their

membership to the clusters shown in D-F. Horizontal black lines divide the categories.

(H) Correlation between activity of cells belonging to ON, OFF and Inhibitory (Inh)

clusters shown in D-F. In general, the ON and OFF responding cells were uncorrelated

(correlation coefficient < 0). Vertical and horizontal black lines divide ON, OFF and

Inhibitory categories. (I) Activity traces of cells in ON and OFF clusters that showed high

correlation with the other category (313 of 1767 cells). The traces showed that this

correlation may be due to OFF cells showing slow decay in fluorescence following light

ON. Manual inspection of the traces did not reveal any cells that responded reliably to

both light ON and OFF. Colorbar for panels G and I is shown below panel I. Scale bar =

25 µm.

Figure 3. The response of different habenula subdomains to change in irradiance.

(A-C) A *narp:GAL4, UAS:DsRed, Brn3a:eGFP* larva, with label in the dorsal habenula

(arrows) and projection to the IPN (arrowhead). A coronal (B) and reconstructed sagittal

(C) section through the left habenula, with dHbM in green and dHbL in magenta. (D)

Spatial distribution of responses in the dorsal habenula of a *narp:GAL4, UAS:GCaMP6s*

fish (8dpf). (E). Responses of non-DsRed cells in the dorsal habenula of a

elavl3:GCaMP6f, narp:GAL4, UAS:DsRed fish (6dpf). (F) Responses in the ventral

habenula of a *dao:GAL4, UAS:GCaMP6s* fish (6dpf). Pixels in panels D-F are coloured

by their membership to *k*-means cluster centers. (G) Cluster centers obtained from

running *k*-means on data in D-F. (H-J) Heatmaps plotting temporal traces from

segmented cells in dHbL (H; n = 5 fish), dHbM (I; n = 5 fish) and vHb (J; n = 8 fish). The

cells are sorted into ON, OFF and Inhibitory categories that are separated by a

horizontal black line. Mean traces of each category are plotted below the heatmaps. (K-

L) Activity in axons of habenula neurons innervating the interpeduncular nucleus, in a

s1011tGAL4, UAS:GCaMP3 fish. Four different planes are shown, in lateral view. All

domains show evoked activity, and responses appear to be organized according to

region. Pixels are colour-coded according to the *k*-means clusters (L). In G-J and L, blue

bars indicate the presence of light and vertical bold lines correspond to light onset while

the dashed lines indicate light offset. a: anterior, p: posterior; l: lateral, m: medial. Scale

bar = 25 μm.

Figure 4. Spatio-temporal characterization of light-evoked activity using high

speed imaging. (A-F) Widefield imaging at 200 Hz. (A) Average of all frames in the

time-lapse, to show morphology. This is a dorsal view of a 5 day old fish *elavl3:CaMP6f*

fish. The habenula has been outlined in black. (B-F) Change in fluorescence relative to

the preceding frame. An increase in fluorescence is seen in the thalamus (arrows) and in

the left habenula (arrowhead). (G-I) Two photon imaging of the habenula in a

s1011tGAL4, UAS:GCaMP6s fish, at 13 Hz. (G) Average of the time-lapse sequence,

showing anatomy. The neuropil is indicated by the arrowhead. (H) Responses to pulses

of light. Pixels are color-coded according to the traces in panel I. rHb: right habenula;

lHb: left habenula. Panels A-E were smoothened using Gaussian blur with sigma = 1.7.

The time indicated is time since start of illumination with the excitation blue LED.

Figure 5. The thalamus projection to the habenula. (A) Dorsal view of the forebrain of

a *Et(SqKR11)* larva, in which habenula afferents from the entopeduncular nucleus are

labeled. The dorsal left neuropil (arrow) is weakly labeled. (B) Dorsal view of the

habenula of a *Et(SqKR11)* larva, following DiD injection into the dorsal neuropil of the left

habenula. The parapineal (arrow) has been retrogradely labeled. Habenula afferents

from the entopeduncular nucleus are labeled in red. (C) 80 μm deeper in the same fish,

showing label in the thalamus (white arrows). (D) Lateral view of another larva, in which

the dorsal left neuropil had been injected with DiD. The retrogradely labeled thalamic

neuropil is indicated (white arrow). The entopeduncular nucleus is indicated by the

yellow arrow. The inset shows a higher magnification labeled thalamic neuropil. Cell

bodies are labeled (white arrowhead). (E) A close up view of the neuropil retrogradely

labeled by DiD (cyan), in a fish where retinal ganglion cells had been labeled with DiI

(yellow). RGC terminals intermingle with fibers from DiD-labeled cells innervating the

neuropil (arrow). The arrowhead indicates a thalamic neuron labeled retrogradely with

DiD. (F, G) Dorsal view of the thalamus (F) and habenula (G) of a fish expressing Kaede

(red) under the control of the *GAL4s1020t* driver. Labelled cells are visible in the

thalamus (F, white arrowheads). Labelled neurites are visible in the thalamic neuropil (F,

yellow arrowheads) and in the habenula neuropils (G, arrowheads). GCaMP6f (green) is

broadly expressed in this fish. (H) Dorsal view of a 6-day-old fish, labeled with an anti-

vGlut1/2 antibody, which marks glutamatergic pre-synapses. All neuropils, including the

dorsal left (arrowhead), are labelled. (I) Dorsal view showing label with an anti-

GAD65/67 antibody. Labeled puncta are visible in the habenula neuropil (arrowhead).

No labeled cell bodies were detected in the habenula. (J) A *gad1b:RFP, vGlut2:GAL4,*

UAS:eGFP fish, with GABAergic cells indicated in magenta and glutamatergic cells

shown in green. Both cell types can be detected in the thalamus. The arrowhead

indicates the neuropil of the putative nucleus rostrolateralis. (K) RFP expression in the

thalamus of a *gad1b:RFP* fish. Arrowheads indicate neurites extending to the neuropil of

the putative nucleus rostrolateralis. All panels except (A) and (H) are single optical

sections. Pa: pallium; rHb: right habenula; lHb: left habenula; Th: thalamus. EN:

entopeduncular nucleus; OT: optic tectum; *: auto-fluorescing pigment cell. Scale bar =

25 µm. Anterior is to the left in all cases.

Figure 6. Optogenetic stimulation of the thalamus triggers habenula activity. (A)

Expression of ChR2-eYFP in the thalamus (arrowheads) of a 5 day old *s1020tGAL4,*

UAS:ChR2-eYFP, elavl3:GCaMP6f fish. (B, C) Activity in the habenula of a ChR2-

expressing fish, with (B) and without (C) blue LED stimulation of the thalamus. The

images show the maximum projections of F/F_0 images for a 25-second period after blue

LED illumination, following subtraction of maximum projections of the period before

illumination (i.e. difference in activity before and after stimulation). (D-F) Heatmaps

showing temporal activity from habenula neurons segmented in fish with (D, E) and

without (F) ChR2. In D (n = 5 fish) and F (n = 2 fish), blue light pulse was given at the

time indicated by the black dashed line. No blue light stimulation was given in E (n = 4

fish). Z-scores were calculated by subtracting each time traces by the total mean and

dividing by the standard deviation. (G) Mean amplitude of z-scores before and after

optogenetic stimulation. Each square stands for a stimulus trial. Amplitude difference

before and after stimulation in ChR2-expressing fish: mean ± 95% CI: 1 Hz: 0.43 ± 0.56,

2 Hz: 0.72 ± 0.35, 4 Hz: 0.89 ± 0.28 and 8 Hz : 1.05 ± 0.18; in siblings: 0.21 ± 0.51.

Scale bar = 25 µm.

Figure 7. Light-evoked activity in the thalamus. (A-E) Evoked activity in five different

focal planes, from dorsal to ventral, of a 5-day-old fish expressing GCaMP6s in thalamic

neurons. Responses are seen in cell bodies (arrows) and in the thalamic neuropil

(arrowheads). The colours represent k-means cluster center (F). (G-J) Lateral view of

light-evoked activity in the thalamus. (G) The region imaged. (H) Average projection of a

lateral view of an *elavl3:GCaMP6f* fish, showing the thalamic neuropil (arrowhead).

(I) The response in this fish to four pulses of blue light. Pixels are colored according to

the *k*-means cluster center (J). (K-P) The effect of eye removal on light-evoked activity in

the thalamus. (K-M) Response in a control (K-M) and eye lesioned fish (N-P), color-

coded according to the k-means cluster centers in (L) and (O) respectively. (M, P)

Trajectory of the thalamic response using the first two principal components (PC1 and

PC2) in control (M) and lesioned (P) fish, colored according to the wedges in panels L

and O to indicate the direction of the trajectories in Principal Component space following

light ON and OFF. Controls, but not lesioned fish, show a reproducible response to light.

Insets in M and P show the correlation coefficient (CC) of the first 20 PCs with a trace

mimicking light evoked activity (see Methods). Unlike controls, the top 20 PCs in the eye

lesioned fish showed weak correlation and did not pick up any light evoked response.

(Q) Correlation coefficients between response of each thalamic pixel and a trace

mimicking response to blue light in multiple *s1020tGAL4, UAS:GCamp6s* fish. All

positive correlation coefficients are plotted here. Each boxplot represents one fish. The

black diamonds are outliers (see Methods). (R-S) Heatmaps showing the number of

pixels with correlation coefficient > 0.5 in control (R) and fish lacking eyes (S). Traces

below show the mean of all the pixels (black traces) and standard error of mean (shaded

region). In panels F, J, L, O and R-S, light onset is indicated by the solid line, while light

offset is indicated by the dashed line. Presence of light is indicated by the blue bars. a:

anterior; p: posterior; d:dorsal, v: ventral. Th: thalamus; Hb:habenula. PC: Principal

component, PC1: First principal component, PC2: Second principal component, CC:

Correlation Coefficient. Scale bar = 25 µm. The drawing in panel G was obtained from

www.uoneuro.uoregon.edu

Figure 8. Asymmetric light-evoked activity in thalamic axon terminals in the dorsal

habenula. (A) Standard deviation projection of a time-series recording of the dorsal

habenula of a 6-day-old *s1020tGAL4,UAS:GCaMP6s* fish. The bright pixels are those

with large change in activity. The surrounding skin, which was auto-fluorescent, has

been masked. The dorsal left neuropil is indicated with a yellow arrowhead while the

dorsal right neuropil is indicated with a green arrowhead. Habenula neurons are dimly

auto-fluorescent. (B) Pixels within the dorsal neuropils with activity above 1 standard

deviation, colour-coded yellow for left habenula and green for right habenula. This

criteria included all terminals in the neuropils. (C) Pixels colour-coded according to

whether they responded to light ON (cyan) or OFF (magenta). A relatively large

proportion of pixels in the left neuropil responded, compared to the right (compare C with

B). Pixels were selected by correlating their activity to a square wave form that was 1

during light ON (for ON pixels) and 1 during light OFF (for OFF pixels). Pixels with

correlation coefficient greater than 0.5 were selected. (D-I) Analysis of multiple fish

(n=7). (D) Boxplot of total number of pixels present in the left and right habenula. The

number of terminals in left and right habenula were comparable across fish. (E) Number

of terminals responding to light ON or light OFF. In D and E, each circle represent data

from a single fish. p-value was obtained using non parametric paired Wilcoxon signed

rank test. W is sum of the ranks, Z is the test statistic, and r is the effect size. (F-G) Heat

maps of activity from all fish, in all thresholded pixels in the left habenula (F), the right

habenula (G), and in pixels corresponding to light ON (H) and light OFF (I). Each line

corresponds to a single pixel. Panels below show the average of the heat maps above.

The shaded region is standard error of mean. Blue boxes indicate when light was

delivered. Light onset is indicated by the solid line, while light offset is indicated by the

dashed line. Anterior is to the top in panels A-C. Scale bar = 25 µm.

Figure 9. The effect of lesioning the thalamus on habenula response to light. (A-C)

Dorsal view of an 8-day-old *elavl3:GCaMP6f* fish, showing neural responses before (A)

and after (C) lesioning the region of the thalamic neuropil that responds to light (yellow

arrowheads in panel A). Pixels in panels A and C are colored according to their activity,

as indicated by traces in panel B. The prominent sustained response to light (cyan

pixels) is reduced after lesion. (D) The extent of lesion, shown in magenta. (E-G)

Habenula activity before (E) and after (G) lesion of the thalamic neuropil. Pixels are

colored according to the traces in (F). There is a reduction in the sustained response to

light, but some activity that is not stimulus-locked can be seen. (H) The habenula after

lesion of AF9, with pixels colored according to the traces in panel (I). (J-M) Heatmaps

showing activity in segmented cells before (J) and after (K) AF9 lesion, and before (L)

and after (M) thalamic neuropil lesion in one fish. Panels below show mean (black trace)

and standard error of mean (shaded region). Light evoked activity is missing following

this lesion. (N) Boxplot showing number of cells in one plane of the dorsal left habenula

that are excited by blue light, following lesion of the thalamic neuropil (n = 12 fish), or

AF7 (n = 2 fish) or AF9 (n = 3 fish), or before lesion (n = 5 fish). P-value was obtained

using non parametric Wilcoxon rank sum test. Z is the test statistic, and r is the effect

size. The statistical comparisons were made between before lesion and after lesion. a:

anterior; p: posterior; Pa: pallium; rHb: right habenula. Images are all single optical

sections. Scale bar = 25 μm.

Figure 10. A potential projection from the ventral lateral geniculate nucleus to the

habenula in mouse. Anterograde label was performed by injecting recombinant adeno

associated virus into the ventral lateral geniculate nucleus (Oh et al., 2014). 0.33 mm^3

was injected into Bregma (-2.46, 2.6, 2.36, 0) of a p56 slc32a1-IRES-Cre mouse, which

expresses Cre in GABAergic neurons. (A) Coronal view, showing an overview of the

label. (B) A high magnification of the area boxed in panel A, showing labeled fibers in the

habenula. mHb: medial habenula. These images are from http://connectivity.brain-

map.org/projection/experiment/siv/267538006?imageId=267538231&imageType=TWO_

PHOTON,SEGMENTATION&initImage=TWO_PHOTON&x=14704&y=7847&z=3.

Movie 1. Habenula neurons do not project to the thalamus. 3D rendition of habenula

projection in a zebrafish larva, visualized by expression of RFP under the *narp* promoter

(red) and eGFP under the *brn3a* promoter (green). There is a clear projection to the

interpeduncular nucleus (IPN), but not to the thalamus.

Movie2. Thalamic neurons project to the habenula. Z-stack of a *s1020t:GAL4, UAS:*

Kaede, elavl3:GCaMP6f fish. Thalamic neurons are shown in red, and they can be seen

to project to the neuropils of the habenula. Red label also appears in streaks in the

lateral habenula. Anterior is to the top.

Movie 3. GAD65/67 label in a zebrafish larva. z-stack of a s*1011t:GAL4,*

UAS:GCaMP3 transgenic fish, after immuno-labelling with an antibody to GAD65/67

(magenta). The stack goes from dorsal to ventral. GAD65/67 label is visible in neuropils

of the habenula; puncta can be seen between cells in the lateral regions of the habenula

in more ventral planes. GAD65/67 labeled cells are visible in the deep focal planes, but

these do not express GCaMP3. The location of GAD65/67 expressing cells correlates

with the thalamus. S1011Et drives GAL4 expression in the habenula, medial pallium and

anterior-lateral pallium. This is a dorsal view, with anterior to the left.

Movie 4. Z-stack of 6 day old *gad1b:RFP, elavl3:GCaMP6f* fish. GABAergic neurons

(magenta) are visible in the thalamus, below the habenula. Anterior is to the left. The

stack goes from dorsal to ventral.

Figure 1. The larval zebrafish habenula has a broad and complex response to change in irradiance. (A) Dorsal view of the head of a live 7 day-old fish, with GCaMP3 expression in the habenula (arrows) under the control of the s1011t GAL4 driver. (B) A single two-photon slice through the dorsal habenula of the fish in panel A (boxed region). (C) A yz reconstruction at the point indicated by the yellow line in panel B, showing a transverse view of the habenula. The dotted lines indicate imaging planes separated by 10 μm. The yellow line indicates the plane imaged in B. Dashed lines show the border of the habenula. (D) Spatial distribution of responses in the habenula of one fish (7 dpf) to pulses of light. 5 planes are shown here. The colors are coded according to the temporal pattern of response, as indicated in (E). Images were collected at a rate of 1 stack/second, and four pulses of light were delivered for 20 seconds each, with variable inter-stimulus interval. (E) Centers of k-means clusters corresponding to colors of pixels in (D). Cluster centers in (D) and the corresponding pixels in (E) indicating responses to light ON are colored in shades of blue and light OFF in shades of red. The horizontal black line represents Z-score of 0. (F) Trajectory of the habenula response in two-dimensional state space, using the first two principal components (PC1 and PC2). Traces are color-coded according to the wedges in panel E, to represent direction in which change in irradiance drives the neural state. In panels E and F, the bold lines correspond to light onset while the dashed lines indicate offset. The presence of light is also indicated by the blue bars. lHb: left habenula; rHb: right habenula. a: anterior; p: posterior. d: dorsal; v: ventral Scale bar = 100 μm in panel A, 25 μm elsewhere.

Figure 2. Habenula response to irradiance change is reproducible. (A-C) Segmentation of habenula neurons using a semi-automated algorithm (see Methods). (D-F) K-means cluster analysis of segmented habenula neuron responses to pulses of blue light in 6 fish (7-8 dpf). Traces show cluster centroids, with shaded regions indicating standard error of the mean. Clusters were grouped by their temporal activity pattern and clusters with excitation to light ON (D) or OFF (E), or inhibition to light (Inhibitory, F) were seen. Clusters without evoked responses are not shown. (G) Activity traces of each cell from the 6 fish grouped into ON, OFF and Inhibitory (Inh) categories based on their membership to the clusters shown in D-F. Horizontal black lines divide the categories. (H) Correlation between activity of cells belonging to ON, OFF and Inhibitory (Inh) clusters shown in D-F. In general, the ON and OFF responding cells were uncorrelated (correlation coefficient < 0). Vertical and horizontal black lines divide ON, OFF and Inhibitory categories. (I) Activity traces of cells in ON and OFF clusters that showed high correlation with the other category (313 of 1767 cells). The traces showed that this correlation may be due to OFF cells showing slow decay in fluorescence following light ON. Manual inspection of the traces did not reveal any cells that responded reliably to both light ON and OFF. Colorbar for panels G and I is shown below panel I. Scale bar = 25 μm.

Figure 3. The response of different habenula subdomains to change in irradiance. (A-C) A *narp:GAL4, UAS:DsRed, Brn3a:eGFP* larva, with label in the dorsal habenula (arrows) and projection to the IPN (arrowhead). A coronal (B) and reconstructed sagittal (C) section through the left habenula, with dHbM in green and dHbL in magenta. (D) Spatial distribution of responses in the dorsal habenula of a *narp:GAL4, UAS:GCaMP6s* fish (8dpf). (E). Responses of non-DsRed cells in the dorsal habenula of a *elavl3:GCaMP6f, narp:GAL4, UAS:DsRed* fish (6dpf). (F) Responses in the ventral habenula of a *dao:GAL4, UAS:GCaMP6s* fish (6dpf). Pixels in panels D-F are coloured by their membership to k-means cluster centers. (G) Cluster centers obtained from running k-means on data in D-F. (H-J) Heatmaps plotting temporal traces from segmented cells in dHbL (H; n = 5 fish), dHbM (I; n = 5 fish) and vHb (J; n = 8 fish). The cells are sorted into ON, OFF and Inhibitory categories that are separated by a horizontal black line. Mean traces of each category are plotted below the heatmaps. (K-L) Activity in axons of habenula neurons innervating the interpeduncular nucleus, in a *s1011tGAL4, UAS:GCaMP3* fish. Four different planes are shown, in lateral view. All domains show evoked activity, and responses appear to be organized according to region of the IPN. Pixels are colour-coded according to the k-means clusters (L). In G-J and L, blue bars indicate the presence of light and vertical bold lines correspond to light onset while the dashed lines indicate light offset. a: anterior, p: posterior; l: lateral, m: medial. Scale bar = 25 μm.

Figure 4. Spatio-temporal characterization of light-evoked activity using high speed imaging. (A-F) Widefield imaging at 200 Hz. (A) Average of all frames in the time-lapse, to show morphology. This is a dorsal view of a 5 day old fish *elavl3:CaMP6f* fish. The habenula has been outlined in black. (B-F) Change in fluorescence relative to the preceding frame. An increase in fluorescence is seen in the thalamus (arrows) and in the left habenula (arrowhead). (G-I) Two photon imaging of the habenula in a *s1011tGAL4, UAS:GCaMP6s* fish, at 13 Hz. (G) Average of the time-lapse sequence, showing anatomy. The neuropil is indicated by the arrowhead. (H) Responses to pulses of light. Pixels are color-coded according to the traces in panel I. rHb: right habenula; lHb: left habenula. Panels A-E were smoothened using Gaussian blur with sigma = 1.7. The time indicated is time since start of illumination with the excitation blue LED.

Figure 5. The thalamus projection to the habenula. (A) Dorsal view of the forebrain of a *Et(SqKR11)* larva, in which habenula afferents from the entopeduncular nucleus are labeled. The dorsal left neuropil (arrow) is weakly labeled. (B) Dorsal view of the habenula of a *Et(SqKR11)* larva, following DiD injection into the dorsal neuropil of the left habenula. The parapineal (arrow) has been retrogradely labeled. Habenula afferents from the entopeduncular nucleus are labeled in red. (C) 80 μm deeper in the same fish, showing label in the thalamus (white arrows). One retrogradely cell in the entopeduncular nucleus is visible (yellow arrowhead). (D) Lateral view of another larva, in which the dorsal left neuropil had been injected with DiD. The retrogradely labeled thalamic neuropil is indicated (white arrow). The entopeduncular nucleus is indicated by the yellow arrow. The inset shows a higher magnification labeled thalamic neuropil. Cell bodies are labeled (white arrowhead). (E) A close up view of the neuropil retrogradely labeled by DiD (cyan), in a fish where retinal ganglion cells had been labeled with DiI (yellow). RGC terminals intermingle with fibers from DiD-labeled cells innervating the neuropil (arrow). The arrowhead indicates a thalamic neuron labeled retrogradely with DiD. (F, G) Dorsal view of the thalamus (F) and habenula (G) of a fish expressing Kaede (red) under the control of the *s1020tGAL4* driver. Labelled cells are visible in the thalamus (F, white arrowheads). Labelled neurites are visible in the thalamic neuropil (F, yellow arrowheads) and in the habenula neuropils (G, arrowheads). GCaMP6f (green) is broadly expressed in this fish. (H) Dorsal view of a 6-day-old fish, labeled with an anti-vGlut1/2 antibody, which marks glutamatergic pre-synapses. All neuropils, including the dorsal left (arrowhead), are labelled. (I) Dorsal view showing label with an anti-GAD65/67 antibody. Labeled puncta are visible in the habenula neuropil (arrowhead). No labeled cell bodies were detected in the habenula. (J) A *gad1b:RFP, vGlut2:GAL4, UAS:eGFP* fish, with GABAergic cells indicated in magenta and glutamatergic cells shown in green. Both cell types can be detected in the thalamus. The arrowhead indicates the neuropil of the putative nucleus rostrolateralis. (K) RFP expression in the thalamus of a *gad1b:RFP* fish. Arrowheads indicate neurites extending to the neuropil of the putative nucleus rostrolateralis. All panels except (A) and (H) are single optical sections. Pa: pallium; rHb: right habenula; lHb: left habenula; Th: thalamus. EN: entopeduncular nucleus; OT: optic tectum; *: auto-fluorescing pigment cell. Scale bar = 25 μm. Anterior is to the left in all cases.

Figure 6. Optogenetic stimulation of the thalamus triggers habenula activity. (A) Expression of ChR2-eYFP in the thalamus (arrowheads) of a 5 day old *s1020t:GAL4, UAS:ChR2-eYFP, elavl3:GCaMP6f* fish. (B, C) Activity in the habenula of a ChR2-expressing fish, with (B) and without (C) blue LED stimulation of the thalamus. The images show the maximum projections of F/F_0 images for a 25-second period after blue LED illumination, following subtraction of maximum projections of the period before illumination (i.e. difference in activity before and after stimulation). (D-F) Heatmaps showing temporal activity from cells segmented in fish with (D, E) and without (F) ChR2. In D (n = 5 fish) and F (n = 2 fish), blue light pulse was given at the time indicated by the black dashed line. No blue light stimulation was given in E (n = 4 fish). Z-scores were calculated by subtracting each time traces by the total mean and dividing by the standard deviation. (G) Mean amplitude of z-scores before and after optogenetic stimulation. Each square stands for a stimulus trial. Amplitude difference before and after stimulation in ChR2-expressing fish: mean ± 95% CI: 1 Hz: 0.43 ± 0.56, 2 Hz: 0.72 ± 0.35, 4 Hz: 0.89 ± 0.28 and 8 Hz : 1.05 ± 0.18; in siblings: 0.21 ± 0.51. Scale bar = 25 μm.

Figure 7. Light-evoked activity in the thalamus. (A-E) Evoked activity in five different focal planes, from dorsal to ventral, of a 5-day-old fish expressing GCaMP6s in thalamic neurons under the s1020t driver. Responses are seen in cell bodies (arrows) and in the thalamic neuropil (arrowheads). The colours represent k-means cluster centers (F). (G-J) Lateral view of light-evoked activity in the thalamus. (G) The region imaged. (H) Average projection of a lateral view of an *elavl3:GCaMP6f* fish, showing the thalamic neuropil (arrowhead). (I) The response in this fish to four pulses of blue light. Pixels are colored according to the k-means cluster centers (J). (K-P) The effect of eye removal on light-evoked activity in the thalamus. (K-M) Response in a control (K-M) and eye lesioned fish (N-P), color-coded according to the k-means cluster centers in (L) and (O) respectively. (M, P) Trajectory of the thalamic response using the first two principal components (PC1 and PC2) in control (M) and lesioned (P) fish, colored according to the wedges in panels L and O to indicate the direction of the trajectories in Principal Component space following light ON and OFF. Controls, but not lesioned fish, show a reproducible response to light. Insets in M and P show the correlation coefficient (CC) of the first 20 PCs with a trace mimicking light evoked activity (see Methods). Unlike controls, the top 20 PCs in the eye lesioned fish showed weak correlation and did not pick up any light evoked response. (Q) Correlation coefficients between response of each thalamic pixel and a trace mimicking response to blue light in multiple *s1020tGAL4, UAS:GCamp6s* fish. All positive correlation coefficients are plotted here. Each boxplot represents one fish. The black diamonds are outliers (see Methods). (R-S) Heatmaps showing the number of pixels with correlation coefficient > 0.5 in control (R) and fish lacking eyes (S). Traces below show the mean of all the pixels (black traces) and standard error of mean (shaded region). In panels F, J, L, O and R-S, light onset is indicated by the solid line, while light offset is indicated by the dashed line. Presence of light is indicated by the blue bars. a: anterior; p: posterior; d:dorsal, v: ventral. Th: thalamus; Hb:habenula. PC: Principal component, PC1: First principal component, PC2: Second principal component, CC: Correlation Coefficient. Scale bar = 25 μm. The drawing in panel G was obtained from www.uoneuro.uoregon.edu

Figure 8. Asymmetric light-evoked activity in thalamic axon terminals in the dorsal habenula. (A) Standard deviation projection of a time-series recording of the dorsal habenula of a 6-day-old *s1020tGAL4,UAS:GCaMP6s* fish. The bright pixels are those with large change in activity. The surrounding skin, which was auto-fluorescent, has been masked. The dorsal left neuropil is indicated with a yellow arrowhead while the dorsal right neuropil is indicated with a green arrowhead. Habenula neurons are dimly auto-fluorescent. (B) Pixels within the dorsal neuropils with activity above 1 standard deviation, colour-coded yellow for left habenula and green for right habenula. This criteria included all terminals in the neuropils. (C) Pixels colour-coded according to whether they responded to light ON (cyan) or OFF (magenta). A relatively large proportion of pixels in the left neuropil responded, compared to the right (compare C with B). Pixels were selected by correlating their activity to a square wave form that was 1 during light ON (for ON pixels) and 1 during light OFF (for OFF pixels). Pixels with correlation coefficient greater than 0.5 were selected. (D-I) Analysis of multiple fish (n=7). (D) Boxplot of total number of pixels present in the left and right habenula. The number of terminals in left and right habenula were comparable across fish. (E) Number of terminals responding to light ON or light OFF. In D and E, each circle represent data from a single fish. p-value was obtained using non parametric paired Wilcoxon signed rank test. W is sum of the ranks, Z is the test statistic, and r is the effect size. (F-G) Heat maps of activity from all fish, in all thresholded pixels in the left habenula (F), the right habenula (G), and in pixels corresponding to light ON (H) and light OFF (I). Each line corresponds to a single pixel. Panels below show the average of the heat maps above. The shaded region is standard error of mean. Blue boxes indicate when light was delivered. Light onset is indicated by the solid line, while light offset is indicated by the dashed line. Anterior is to the top in panels A-C. Scale bar = 25 μm.

Figure 9. The effect of lesioning the thalamus on habenula response to light. (A-C) Dorsal view of an 8-day-old elavl3:GCaMP6f fish, showing neural responses before (A) and after (C) lesioning the region of the thalamic neuropil that responds to light (yellow arrowheads in panel A). Pixels in panels A and C are colored according to their activity, as indicated by traces in panel B. The prominent sustained response to light (cyan pixels) is reduced after lesion. (D) The extent of lesion, shown in magenta. (E-G) Habenula activity before (E) and after (G) lesion of the thalamic neuropil. Pixels are colored according to the traces in (F). There is a reduction in the sustained response to light, but some activity that is not stimulus-locked can be seen. (H) The habenula after lesion of AF9, with pixels colored according to the traces in panel (I). (J-M) Heatmaps showing activity in segmented cells before (J) and after (K) AF9 lesion, and before (L) and after (M) thalamic neuropil lesion in one fish. Panels below show mean (black trace) and standard error of mean (shaded region). Light evoked activity is missing following this lesion. (N) Boxplot showing number of cells in one plane of the dorsal left habenula that are excited by blue light, following lesion of the thalamic neuropil (n = 12 fish), or AF7 (n = 2 fish) or AF9 (n = 3 fish), or before lesion (n = 5 fish). P-value was obtained using non parametric Wilcoxon rank sum test. Z is the test statistic, and r is the effect size. The statistical comparisons were made between before lesion and after lesion. a: anterior; p: posterior; Pa: pallium; rHb: right habenula. Images are all single optical sections. Scale bar = 25 μm.

Figure 10. A potential projection from the ventral lateral geniculate nucleus to the habenula in a mouse. Anterograde label was performed by injecting recombinant adeno associated virus into the ventral lateral geniculate nucleus. 0.33 mm³ was injected into Bregma (-2.46, 2.6, 2.36, 0) of a p56 slc32a1-IRES-Cre mouse, which expresses Cre in GABAergic neurons (Oh et al., 2014). (A) Coronal view, showing an overview of the label. (B) A high magnification of the area boxed in panel A, showing labeled fibers in the habenula. mHb: medial habenula. These images are from http://connectivity.brain-map.org/projection/experiment/siv/267538006?imageId=267538231&imageType=TWO_PHOTON,SEGMENTATION&initImage=TWO_PHOTON&x=14704&y=7847&z=3.