1 **Wild-type U2AF1 antagonizes the splicing program characteristic of U2AF1-**
2 **mutant tumors and is required for cell survival**
3
4
5 Short title: Interplay of mutant and wild-type U2AF1
6
7

8 Dennis Liang Fei[1,2,*], Hayley Motowski[1], Rakesh Chatrikhi[3], Sameer Prasad[1], Jovian
9 Yu[1], Shaojian Gao[4], Clara L. Kielkopf[3,*], Robert K. Bradley[5,6,*], Harold Varmus[1,2*]
10
11
1. Cancer Biology Section, Cancer Genetics Branch, National Human Genome
Research Institute, Bethesda, MD 20892, United States
2. Department of Medicine, Meyer Cancer Center, Weill Cornell Medicine, New York,
NY 10065, United States
3. Department of Biochemistry and Biophysics, Center for RNA Biology, University of
Rochester Medical Center, Rochester, NY 14642, United States
4. Genetics Branch, National Cancer Institute, Bethesda, Maryland, 20892, United
States
5. Computational Biology Program, Public Health Sciences Division, Fred Hutchinson
Cancer Research Center, Seattle, WA 98109, United States
6. Basic Sciences Division, Fred Hutchinson Cancer Research Center, Seattle,
Washington 98109, United States
24
25
26 * Correspondence: D.L.F. dlf2002@med.cornell.edu; C.L.K.
27 clara_kielkopf@urmc.rochester.edu; R.K.B. rbradley@fredhutch.org; H.V.
28 varmus@med.cornell.edu
29
30

31 Keywords: U2AF1, splicing factor, S34F, RNA binding, lung adenocarcinoma

**Abstract**

We have asked how the common S34F mutation in the splicing factor U2AF1 regulates alternative splicing in lung cancer, and why wild-type U2AF1 is retained in cancers with this mutation. A human lung epithelial cell line was genetically modified so that *U2AF1*S34F is expressed from one of the two endogenous *U2AF1* loci. By altering levels of mutant or wild-type U2AF1 in this cell line and by analyzing published data on human lung adenocarcinomas, we show that S34F-associated changes in alternative splicing are proportional to the ratio of S34F:wild-type gene products and not to absolute levels of either the mutant or wild-type factor. Preferential recognition of specific 3' splice sites in S34F-expressing cells is largely explained by differential *in vitro* RNA-binding affinities of mutant versus wild-type U2AF1 for those same 3' splice sites. Finally, we show that lung adenocarcinoma cell lines bearing *U2AF1* mutations do not require the mutant protein for growth *in vitro* or *in vivo*. In contrast, wild-type U2AF1 is required for survival, regardless of whether cells carry the *U2AF1*S34F allele. Our results provide mechanistic explanations of the magnitude of splicing changes observed in *U2AF1*-mutant cells and why tumors harboring *U2AF1* mutations always retain an expressed copy of the wild-type allele.

**<u>Author Summary</u>**

Large-scale genomics studies have identified recurrent mutations in many genes

that fall outside the conventional domain of proto-oncogenes. They include genes

encoding factors that mediate RNA splicing; mutations affecting four of these genes are

present in up to half of proliferative myeloid disorders and in a significant number of

solid tumors, including lung adenocarcinoma. Here we have characterized several

properties of a common mutant version of the U2AF1 splicing factor, a component of

the U2 auxiliary factor complex, in lung cells. We have found that mutant-associated

changes in splice site selection are primarily influenced by the ratio of mutant and wild-

type U2AF1 gene products; thus increasing wild-type U2AF1 levels represses the

mutant-induced splicing program. We show that the altered splice site preferences of

mutant U2AF1 can be attributed to changes in its binding to relevant 3' splice sites. We

also show that mutant U2AF1 is different from some oncogenes: the growth properties

of lung cancer cell lines carrying the mutant allele are unaffected by loss of the mutant

gene, while the wild-type allele is absolutely required for survival. These results

advance our understanding of the molecular determinants of the mutant-associated

splicing program, and they highlight previously unappreciated roles of wild-type U2AF1

in the presence of the recurrent *U2AF1*S34F mutation.

## Introduction

Somatic mutations in genes encoding four splicing factors (*U2AF1*, *SF3B1*, *SRSF2* and *ZRSR2*) have recently been reported in up to 50% of myelodysplastic syndromes (MDS) and related neoplasms and at lower frequencies in a variety of solid tumors [1-9]. Among these factors, only *U2AF1* is known to be recurrently mutated in lung adenocarcinomas (LUADs) [3,9]. The only recurrent missense mutation of *U2AF1* in LUAD affects codon 34 and always changes the conserved serine in a zinc knuckle motif to phenylalanine (p.Ser34Phe, or S34F). This striking mutational consistency suggests a critical, yet unknown, role for *U2AF1*S34F during lung carcinogenesis. In addition, the wild-type (WT) *U2AF1* allele is always retained in cancers with common *U2AF1* mutations, including *U2AF1*S34F [2]. However, the functional significance of the wild-type allele in cells with mutant *U2AF1* is not known.

U2AF1 is a component of the U2 small nuclear ribonucleoprotein auxiliary factor complex (U2AF) [10,11]. During early spliceosome assembly, U2AF recognizes sequences at the 3' ends of introns to facilitate the recruitment of the U2 small nuclear ribonucleoprotein (snRNP) complex to the 3' splice site; the recruitment occurs in conjunction with recognition of the intronic branch point by splicing factor 1 (SF1) [12,13]. *In vitro* crosslinking assays showed that U2AF1 contacts the AG dinucleotide at the intron-exon boundary and flanking sequences [14-16].

Consistent with the critical role that U2AF1 plays in RNA splicing, *U2AF1* mutations are known to cause specific alterations in RNA splicing, most notably affecting the inclusion of cassette exons in mRNA [17-20]. However, the precise molecular basis of these splicing alterations, as well as how they are quantitatively

regulated, is unknown. One possibility is that U2AF1 mutations cause altered RNA-

binding affinity, resulting in altered splice site recognition. A computational model of the

structure of the U2AF1:RNA complex suggested that Ser34 is a critical residue that

contacts RNA [17]. Another study reported that U2AF1S34F exhibited altered affinity

relative to the wild-type protein for RNA oligonucleotides derived from a cassette exon

whose recognition is repressed in S34F-expressing cells [18]. Finally, the S34F

mutation reportedly prevented a minimal fission yeast U2AF heterodimer from binding to

a particular 3' splice site RNA sequence [21]. However, it is not known whether altered

RNA binding accounts for most S34F-associated splicing alterations and whether

mechanisms other than altered binding control S34F-associated splicing.

Here, we combine genetic and biochemical approaches to show that wild-type

U2AF1 antagonizes the S34F-associated splicing program in lung epithelial cells.

Analyses of the transcriptomes of primary LUAD samples as well as isogenic lung cells

in culture indicate that the ratio of mutant to wild-type U2AF1 gene products is a critical

determinant of the magnitude of S34F-associated changes in alternative splicing. S34F-

associated splicing alterations can be largely explained by differences in the relative

affinities of U2AF-SF1 complexes containing mutant versus wild-type U2AF1 for RNA

containing the relevant 3' splice sites. Moreover, we show that proliferation of cancer

cells with *U2AF1*S34F is critically dependent on expression of the wild-type, but not the

mutant, allele of *U2AF1*.

**Results**

**S34F-associated splicing correlated with S34F:WT mRNA ratios in LUAD**

Before undertaking experiments to study the effects of mutant U2AF1 in cultured

cells, we first examined 512 transcriptomes from primary human LUADs published by

The Cancer Genome Atlas (TCGA) [9]. Thirteen of these tumors harbor the most

common *U2AF1* mutation, S34F*.* (Two others carry rare mutations of unknown

significance, S65L and G216R, and were not considered further.) We identified cassette

exons whose inclusion was increased or decreased by ten percent or more in each

tumor with the S34F mutation relative to the median inclusion of each cassette exon

across all tumors without a *U2AF1* mutation. We then identified consensus sequence

motifs for the 3' splice sites lying immediately upstream of the cassette exons that were

promoted or repressed in association with *U2AF1*S34F, represented by "sequence

logos" as shown in Figs. 1A and S1. The same analysis was performed on 19 random

tumors lacking a *U2AF1* mutation as controls.

As illustrated by data from patient 7903 in Fig. 1A, over two hundred cassette

exons were included more frequently and a similar number were included less

frequently in this *U2AF1*-mutant tumor. Notably, as illustrated by the sequence logos,

the nucleotide distribution at the -3 position (boxed) of promoted and repressed exons

was different from that observed upstream of the much larger number of unaffected

exons: A replaced T as the second most common nucleotide preceding the promoted

exons, while T was more common than C in the sequence preceding the repressed

exons. These patterns were observed in nine of the thirteen tumors with the

*U2AF1*S34F allele (Supplemental Fig. S1A). They have been observed previously in

comparisons between transcriptomes carrying the *U2AF1*S34F allele with wild-type

transcriptomes [17-20], and are therefore henceforth referred to as the "typical S34F"

consensus 3' splice sites. In the other four tumors with the *U2AF1*S34F mutation, these

"typical S34F" consensus 3' splice sites were partially or completely absent (Figs. 1A,

S1B). Those four tumors exhibited consensus 3' splice sites that were similar to the

consensus 3' splice sites identified in tumors lacking a *U2AF1* mutation, where

variations in inclusion relative to the median wild-type sample are presumably stochastic

(Supplemental Fig. S1C). Thus, we henceforth refer to these sequence patterns, as

shown for tumor from patient 7727 in Fig. 1A, as "quasi-WT".

To explain why transcriptomes of some tumors with *U2AF1* mutations showed a

typical S34F-associated consensus 3' splice sites, while others exhibited quasi-WT

patterns, we estimated the levels of mutant and total *U2AF1* mRNA based on available

data from the tumors to determine the ratio of mutant to wild-type (S34F:WT) mRNA.

Tumors with quasi-WT patterns had low S34F:WT mRNA ratios (ranging from 0.27 to

0.31), whereas all but one tumor with the S34F-associated pattern had higher ratios

(0.43 or more) (Fig. 1B). In contrast, absolute levels of *U2AF1*S34F mRNA or total

*U2AF1* mRNA levels were not different between these two groups of tumors

(Supplemental Figs. S2A, B).

We next sought to understand the origin of the wide range of S34F:WT ratios that

we observed. These ratios, ranging from 0.26 to 0.82, could not be explained by

contamination of tumor cells with non-tumor cells, since the proportion of tumor nuclei

reported for these samples did not correlate with S34F:WT mRNA ratios (Fig. 1C) or

with the presence or absence of the expected S34F-associated consensus 3' splice

sites (Supplemental Fig. S2C). Conversely, *U2AF1* DNA copy number correlated with

the estimated levels of total *U2AF1* mRNA (Supplemental Fig. S2D). Seven of the 13

*U2AF1*S34F mutant samples, including two of the four samples displaying quasi-WT

consensus 3' splice site sequences, showed either copy number loss or gain at the

*U2AF1* locus, suggesting that copy number variation (CNV) might account, at least in

part, for the varying S34F:WT mRNA ratios in LUAD samples (Fig. 1B). Unbalanced

allelic expression or proportions of tumor subclones might also contribute to the variable

S34F:WT mRNA ratios, although these possibilities could not be readily tested using the

available LUAD data.

We next tested whether the *U2AF1* S34F:WT ratio was associated with

quantitative changes in splicing (versus the qualitative differences in consensus 3' splice

sites described above). We correlated S34F:WT ratios with the quantitative inclusion of

specific S34F-responsive cassette exons and 5' extensions of exons resulting from

competing 3' splice sites that were reported previously [17,19,20]. We studied three

cassette exons that exhibited less (*ASUN* and *STRAP*) or more (*ATR*) inclusion, and

two 5' extensions of exons (*FMR1* and *CASP8*) that were used less frequently in the

presence of *U2AF1*S34F. Tumors with the highest S34F:WT mRNA ratios showed the

lowest inclusion levels of the cassette exon in *STRAP* mRNA, whereas tumors without a

*U2AF1* mutation had the highest level of inclusion (Fig. 1D; Pearson correlation of -

0.79). Similar correlations were observed between inclusion levels of other

representative exons and S34F:WT mRNA ratios (Supplemental Fig. S3, panels E, I, M

and Q). As controls, we tested for correlations between the inclusion of these cassette

exons and levels of *U2AF1*S34F mRNA, total *U2AF1* mRNA, or percent tumor nuclei.

None of these analyses, with the exception of the *U2AF1*S34F mRNA level versus the

inclusion of the 5' extension of the *FMR1* exon, showed a relationship as strong as that

observed for the S34F:WT mRNA ratio (Supplemental Figs. S3 panels B-D, F-H, J-L, N-

P and R-T). These results indicate that the S34F:WT mRNA ratio predicts the

magnitude of S34F-associated splicing in human LUAD.

**Creation of isogenic lung cell lines that recapitulated features of S34F-associated**

**splicing in LUAD.**

The results presented in the preceding section, based on analyses of LUAD

tumors with the *U2AF1*S34F mutation, suggest that the magnitude of S34F-associated

splicing is a function of the S34F:WT mRNA ratio. To directly test this hypothesis, we

developed a cell line that allows manipulation of WT and mutant U2AF1 gene product

levels and measurement of the corresponding effects on RNA splicing.

The human bronchial epithelial cell line (HBEC3kt) was previously derived from

normal human bronchial tissue and immortalized by introduction of expression vectors

encoding human telomerase reverse transcriptase (*hTERT*) and cyclin-dependent

kinase-4 (*CDK4*) [22]. To knock in a *U2AF1*S34F allele at an endogenous locus in

HBEC3kt cells, we adopted a published genomic DNA editing approach [23], using the

*PiggyBac* transposon that leaves no traces of exogenous DNA at the locus (Fig. 2A; see

Supplemental Materials and Methods for details). We identified three cell clones at

intermediate stages of gene editing after screening more than 50 primary transfectants

(Supplemental Fig. S4A, B). Sanger sequencing of these intermediate clones revealed

that one of the three clones carried the desired S34F missense sequence, while two

clones were wild-type (Supplemental Fig. S4C). Wild-type intermediates are expected

because a homologous sequence between the S34F point mutation and the drug

cassette in the vector can serve as the 5' homology arm for recombination (designated

as 5' HA#2 in Fig. 2A). From the final clones derived from these intermediates (after

transposition to remove the drug cassette flanked by the *PiggyBac* elements), we chose

one subclone from each of the two wild-type intermediate clones (referred to as WT1

and WT2 cells) and two subclones from the sole mutant intermediate clone (referred to

as MUT1a and MUT1b cells) for all subsequent experiments with isogenic HBEC3kt

cells (Supplemental Fig. S4D). The MUT and WT cells all expressed similar levels of

*U2AF1* mRNA and protein (Supplemental Fig. S4E). Using high-throughput mRNA

sequencing (RNA-seq) and allele-specific RT-qPCR, we observed similar levels of wild-

type and mutant *U2AF1* mRNAs in the MUT cells (Figs. 2B, S5C), consistent with

heterozygosity at the *U2AF1* locus.

To determine how the engineered *U2AF1 S34F* allele affects mRNA splicing, we

first assayed the inclusion levels of 20 cassette exons that were previously reported to

be associated with mutant *U2AF1* in both LUAD and AML (acute myeloid leukemia)

[19]. We confirmed that all 20 of these cassette exons, which included the previously

studies *ASUN* and *STRAP* cassette exons, were indeed S34F-dependent in our

engineered cells using RT-qPCR with isoform-specific primers (Figs. 2C, S6).

We next evaluated the global difference in cassette exon recognition in MUT

versus WT cells using RNA-seq (Supplemental Table S1). MUT and WT cells were

grouped separately in an unsupervised cluster analysis based on cassette exon

inclusion (Fig. 2D). We observed the expected consensus 3' splice sites of cassette

exons that were promoted or repressed in MUT versus WT cells (Fig. 2E). Overall,

these results indicate that we successfully created clonal HBEC3kt cells isogenic for

*U2AF1*S34F and that the MUT cells exhibited similar alterations in splicing relative to

their WT counterparts as do primary LUAD transcriptomes.

**The ratio of S34F:WT gene products controlled S34F-associated splicing in**

**isogenic lung cells.**

We next used our isogenic cell model with the *U2AF1*S34F mutation to test the

hypothesis that the S34F:WT ratio, rather than absolute levels of the mutant or wild-type

gene products, controls S34F-associated splicing. We tested two specific predictions.

First, changing the levels of *U2AF1*S34F while keeping the S34F:WT ratio constant

should not affect S34F-associated splicing. Second, changing the level of wild-type

*U2AF1* while keeping the level of *U2AF1*S34F constant (e.g., allowing the S34F:WT

ratio to change) should alter the inclusion of S34F-dependent cassette exons. We

tested these predictions in the isogenic HBEC3kt cells by manipulating levels of wild-

type or mutant *U2AF1* gene products and measuring the subsequent changes in S34F-

associated splicing.

We first reduced the amounts of both mutant and wild-type *U2AF1* mRNA

concordantly in MUT1a cells, keeping the S34F:WT mRNA ratio constant. This was

achieved by transducing MUT1a cells with short hairpin RNAs (shRNAs) that target

regions of the *U2AF1* transcripts distant from the S34F missense mutation. The same

shRNAs were also introduced in WT1 cells as a control. Allele-sensitive RT-qPCR

confirmed that the mRNA ratio of the mutant and wild-type *U2AF1* remained constant

(Supplemental Fig. S7A), while the overall U2AF1 mRNA and protein levels were

reduced by more than 90% (Figs. 3A, bottom panel, and S7B). Knockdown of total

*U2AF1* in both MUT1a and WT1 cells did not cause significant changes in recognition of

the *ASUN* or *STRAP* cassette exons, two splicing events that are strongly associated

with *U2AF1*S34F, in either cell line (Fig. 3A, upper panels). Similar results were

obtained for two additional S34F-associated cassette exons in *USP25* and *AXL* that

exhibit increased inclusion in cells expressing *U2AF1*S34F (Supplemental Fig. S7C, D).

We next confirmed that knockdown of *U2AF1* was sufficient to alter splicing

events known to be dependent on wild-type U2AF1. We studied a competing 3' splice

site event in *SLC35C2*, in which the use of an intron-proximal over an intron-distal 3'

splice site depends on the level of U2AF1 independent of a *U2AF1* mutation [24].

Knockdown of total *U2AF1* in either WT1 or MUT1a cell lines reduced the use of the

U2AF1-dependent intron-proximal 3' splice site (Supplemental Fig. S7E). Thus,

reduction of *U2AF1* to levels sufficient to affect U2AF1-dependent alternative splicing

did not affect S34F-associated splicing in MUT1a cells when the S34F:WT ratio was

maintained.

We next altered the S34F:WT ratio by separately overexpressing mutant or wild-

type *U2AF1* in WT1 and MUT1a cells and examining the subsequent changes in the

recognition of the *ASUN* and *STRAP* cassette exons. These cassette exons are

preferentially excluded in cells expressing *U2AF1*S34F. Increasing the amount of

U2AF1S34F protein—hence increasing the S34F:WT ratio in either cell type—further

enhanced skipping of these cassette exons (Fig. 3B). Conversely, decreasing the

S34F:WT ratio in MUT1a cells by increasing the production of wild-type U2AF1 protein

reduced the extent of exon skipping to approximately the same levels seen in WT1 cells

(Fig. 3B).

We also altered the production of wild-type U2AF1 mRNA and protein in MUT1a

cells by disrupting the endogenous wild-type *U2AF1* locus with the CRISPR-Cas9

system. Single-guide RNAs (sgRNAs) designed to match either the WT or mutant

*U2AF1* sequences were shown to disrupt either reading frame selectively, generating

indels (insertions and deletions) at the *U2AF1* locus and thereby changing the S34F:WT

ratios (Supplemental Figs. S8, S9). Since WT *U2AF1* is required for the growth of these

cells (as shown below; Fig. 6), we extracted RNA and protein from cells six days after

transduction with Cas9 and sgRNA-WT, when depletion of wild-type U2AF1 was

incomplete (Fig. 3C). Selective disruption of wild-type *U2AF1* increased the S34F:WT

mRNA ratio in MUT1a cells; as predicted, the extent of exon skipping was further

increased in *ASUN* and *STRAP* mRNAs (Fig. 3C). Notably, the degree of exon skipping

induced by mutant cDNA was similar to that caused by disrupting the wild-type *U2AF1*

allele (compare Figs. 3C with 3B), even though the absolute protein levels of

U2AF1S34F were different in the two experiments. These results show that wild-type

U2AF1 antagonizes the activity of U2AF1S34F by a competitive mechanism, such that

the S34F:WT ratio controls the magnitude of S34F-associated splicing changes

independent of levels of either mutant or wild-type protein.

**Disruption of WT U2AF1 globally enhanced S34F-associated splicing.**

Results in the preceding sections are based on studies of a few well-documented

S34F-responsive cassette exons that likely serve as surrogates for the global effects of

U2AF1S34F on splice site recognition. To determine whether these results reflect

general rules governing S34F-associated splicing, we used RNA-seq to measure the

consequences of altering S34F:WT ratios on global recognition of cassette exons.

When wild-type *U2AF1* was diminished by CRISPR-Cas9-mediated disruption in

MUT1a cells (see Fig. 3C), the S34F-associated changes in inclusion (Fig. 4A) or

exclusion (Fig. 4B) of cassette exons were enhanced. These global effects are

consistent with our measurements of individual cassette exons by RT-qPCR (Fig. 3C).

An unsupervised cluster analysis suggests that disruption of wild-type *U2AF1* in MUT1a

cells primarily enhances the magnitude of changes for S34F-associated cassette exons

(Fig. 4C). We also observed a modest increase in the preference for C versus T at the -

3 position of the consensus 3' splice sites of cassette exons that were promoted versus

repressed in association with *U2AF1*S34F (Fig. 4D) following reduction of wild-type

*U2AF1* levels, consistent with the observed association between the S34F:WT mRNA

ratio and typical S34F-associated consensus 3' splice sites identified in LUAD tumor

transcriptomes (Fig. 1).

We observed similar results when we extended our analyses to include other

types of alternative splicing beyond cassette exon recognition (Supplemental Fig. S10).

We used RT-qPCR to validate five splicing alterations that are sensitive to ablation of

wild-type *U2AF1* in the presence of *U2AF1*S34F (Supplemental Fig. S11). Two of the

five events involve incorporation of cassette exons (in *ATR* and *MED15*), two involve

competition between different 3' splice sites (in *CASP8* and *SRP19*), and one involves a

choice between two mutually exclusive exons (in *H2AFY*). We conclude that the

importance of the S34F:WT ratio for S34F-dependent splicing changes extends from

cassette exon recognition to other types of alternative splicing.

**Changes in RNA splicing correlated with relative binding affinities of mutant and**

**WT U2AF1 complexes.**

Based on the role of U2AF1 in 3' splice site recognition, we hypothesized that

differential RNA binding by mutant and WT U2AF1 could contribute to the observed

dependence of S34F-associated splicing on the WT:S34F ratios. It has previously been

shown that the S34F mutation reduces the binding affinity of the U2AF1-containing

complex for a representative skipped splice site [18]. However, whether changes in

RNA binding could account for exon inclusion, as well as the generality of this

observation, were unknown. We therefore tested whether altered RNA-binding affinity

could explain mutation-dependent increases (*ZFAND1*, *FXR1, ATR, MED15*) and

decrease (*CEP164*) in exon inclusion, as well as competing 3' splice site recognition

(*FMR1*). These splicing events exhibited S34F-associated alterations in both our

isogenic cell lines (Supplemental Table S1) and in LUAD transcriptomes (Supplemental

Table S2).

We determined the RNA binding affinities of purified U2AF1-containing protein

complexes using fluorescence anisotropy assays, in which the anisotropy increases of

fluorescein-labeled RNA oligonucleotides following protein titration were fit to obtain the

apparent equilibrium binding affinities (Supplemental Fig. S12; Supplemental Materials

and Methods). The recombinant proteins comprised either WT or S34F-mutant U2AF1

as ternary complexes with the U2AF2 and SF1 subunits that recognize the adjoining 3'

splice site consensus sequences. The constructs were nearly full length and included

the relevant domains for 3' splice site recognition [15,25,26] (Fig. 5A). The six pairs of

tested RNA oligonucleotides (33 - 35 nucleotides in length) were derived from the

proximal or distal 3' splice sites of the six genes listed above (Figs. 5B, S12). Combined

with prior results for sequence variants derived from the S34F-skipped *DEK* cassette

exon [18], we have in total measured binding affinities for 16 RNA oligonucleotides,

which consist of five sequences with a U at the -3 position of the 3' splice site ("UAG"

splice sites), seven "CAG" splice sites, and four "AAG" splice sites.

The trends among S34F-altered RNA binding affinities of U2AF1 complexes (Fig.

5C - H) for the tested splice site sequences generally agreed with the nucleotide

distributions observed in consensus 3' splice site that are promoted or repressed by

*U2AF1*S34F (Figs. 1A, S1A). The S34F mutation reduced the affinities of U2AF1-

containing complexes for four out of five tested "UAG" splice sites, consistent with T as

the most common nucleotide at the -3 position of the 3' splice site (henceforth referred

to as -3T) for S34F-repressed exons. Likewise, the S34F mutation often increased the

affinities of U2AF1 complexes for "CAG" splice sites (for three out of seven tested

sequences), consistent with -3C as the most common nucleotide preceding S34F-

promoted exons. The remaining tested "UAG" or "CAG" splice sites showed no

significant difference between S34F and WT protein binding. The "AAG" splice sites

lacked a consistent relationship to the S34F-induced RNA affinity changes of U2AF1

complexes *per se*. However, the S34F mutation increased the binding of the U2AF1-

containing complex for one "AAG" splice site for an S34F-promoted exon (*ZFAND1*),

which is consistent with the greater prevalence of -3A than -3T preceding S34F-

promoted exons.

Overall, the altered binding affinities of U2AF1-containing complexes for the

proximal 3' splice site could account for four of the six S34F-associated alternative

splicing events (*CEP164*, *FMR1*, *ZFAND1*, *FXR1*) (Figs. 5C - F, Supplementary Table

S3). Similar to the previously-tested S34F-skipped splice site in *DEK* [18], the S34F

mutation decreased affinities of the U2AF1-containing complexes for the skipped 3'

splice sites of the *CEP164* and *FMR1* exons (Fig. 5C, D). Remarkably, the S34F

mutation enhanced binding of the U2AF1-containing complexes to the proximal 3' splice

sites of *ZFAND1* and *FXR1* (Fig. 5E, F), which could explain the enhanced inclusion of

these exons in cell lines and LUAD (Supplementary Table S3). In agreement with the

observed splicing changes, the S34F mutation had no significant effect on the distal 3'

splice sites of these exons.

The affinities of the mutant U2AF1 complexes for the proximal splice site

oligonucleotides of the remaining two S34F-promoted exons (*ATR* and *MED15*) were

either similar to wild-type or decreased (Fig. 5G, H). These results differed from the

S34F-dependent increase in U2AF1 binding to the proximal 3' splice site that was

observed for *ZFAND1* and *FXR1* (Fig. 5E, F), which could readily explain the enhanced

exon inclusions. However, for the *ATR* pre-mRNA, the *U2AF1* mutation reduced binding

to the distal more than to the proximal 3' splice site (Fig. 5G, third and fourth columns).

Given co-transcriptional splicing in the 5'-to-3' direction, the downstream (as opposed to

upstream) 3' splice sites could compete as splicing acceptors for a given 5' donor splice

site when transcription is relatively rapid as compared to splicing. (Such 3' splice site

competition likely occurs relatively frequently, as the *ATR* cassette exon is alternatively

spliced even in wild-type cells). As such, a "net gain" in affinity for the proximal relative

to distal 3' splice site could explain the observed S34F-associated exon inclusion in

*ATR* mRNAs. For the *MED15* pre-mRNA, deviation of the S34F-associated splicing

changes and a simple RNA affinity model suggested that additional mechanisms control

selection of the *MED15* 3' splice sites. In total, our measurements of *in vitro* RNA-

binding affinities are sufficient to explain six of seven tested alterations in splice site

recognition driven by *U2AF1*S34F (Fig 5C-H and in [18]).

**HBEC3kt and LUAD cells were not dependent on *U2AF1*S34F for growth, but wild-**

**type *U2AF1* was absolutely required.**

Other than its effect on RNA splicing, the consequences of the *U2AF1*S34F

mutation on cell behavior are largely unknown. Recurrent mutations, such as *U2AF1*

*S34F*, are considered likely to confer a selective advantage to cells in which they occur

when expressed at physiologically relevant levels. However, mutant HBEC3kt cells

(MUT1a and MUT1b) do not exhibit obvious phenotypic properties of neoplastic

transformation—such as a growth advantage over wild-type cells (Supplemental Fig.

S13) or an ability to grow in an anchorage-independent manner—that are frequently

observed in cultured cells expressing well-known oncogenes, like mutant *RAS* genes.

Another attribute of some well-known oncogenes, such as *BCR-ABL* fusion in

chronic myeloid leukemia or mutant *EGFR* or *KRAS* in LUAD, is the dependence on

sustained expression of those oncogenes for the maintenance of cell growth or viability.

To determine whether LUAD cells harboring a pre-existing S34F mutation are

dependent upon (or "addicted to") the mutant allele, we searched the COSMIC

database for LUAD cell lines with the *U2AF1*S34F allele [27]. Two cell lines (H441 and

HCC78) were found and both these cells exhibit copy number gains at the *U2AF1* locus

(three copies for H441 cells; four copies for HCC78 cells) and one copy of a variant

allele. We confirmed that *U2AF1*S34F was the minor allele in these cells by Sanger

sequencing and allele-specific RT-qPCR (Supplemental Fig. S14A, B). We further used

the CRISPR-Cas9 system to selectively disrupt the wild-type or mutant *U2AF1*

sequences and then assessed the impact of inactivating the *U2AF1* alleles on the

clonogenic growth of the two LUAD lines with the *U2AF1* mutation. In addition, we

performed similar experiments with the LUAD cell line A549 (wild-type for U2AF1) and

the HBEC3kt-derived MUT1a cell line.

In all instances, loss of the mutant allele did not impair cell growth. Only one line

(H441) exhibited altered growth, in the form of a two-fold increase in clonogenicity (Fig.

6A). Successful disruption of the *U2AF1*S3F allele was confirmed by restoration of a

normal RNA splicing profile in subclonal cells derived from the clonogenic assay

colonies (see Supplemental Figs. S14 – S17, Tables S4 and S5, and text below). In

contrast, loss of the wild-type allele completely inhibited clonogenic growth in all tested

cell lines, regardless of whether the line carried the *U2AF1*S34F allele or not. A rescue

experiment confirmed that loss of cell growth was due to loss of wild-type *U2AF1*

expression. The loss of clonogenic capacity after disrupting endogenous *U2AF1* in

A549 cells was prevented by first transducing them with a form of wild-type *U2AF1*

cDNA (Fig. 6B) that is not predicted to be a target for sgRNA-WT (Supplemental Fig.

S8). Overall, these findings indicate that wild-type *U2AF1* is required for the clonogenic

growth of cells, including lung cancer cell lines, that the S34F mutant is unable to

compensate for loss of the wild-type allele, and that LUAD cells with the S34F mutation

are not dependent on the mutant allele for growth *in vitro*.

To examine the effect of *U2AF1*S34F on tumor growth *in vivo*, we derived H441

and HCC78 cells transduced with Cas9 and sgRNA-S34F or sgRNA-GFP as polyclonal

pools or as clones (Supplemental Figs. S14, S15). The cell lines were verified to either

carry or not carry the *U2AF1*S34F allele, and we confirmed that the Cas9 and sgRNA-

S34F-transduced cells lost the S34F-associated splicing program (Supplemental Figs.

S16, S17).

We inoculated these subclonal cell lines subcutaneously in nude mice and

monitored xenograft tumor growth. The H441-derived cell lines, in which the

*U2AF1*S34F allele was disrupted, were able to establish tumors *in vivo* at rates similar

to those observed for tumor cells carrying the mutant allele (Supplemental Table S6).

The HCC78-derived cell lines did not grow palpable tumors after xenografting within the

observation period, so the requirement for *U2AF1*S34F *in vivo* could not be tested in

that line. These experiments show that *U2AF1*S34F is dispensable for growth of these

LUAD cell lines *in vivo*, a result consistent with the clonogenicity assays shown in Fig. 6.

We conclude that *U2AF1*S34F appears to be neither sufficient nor necessary for lung

cell transformation in these assays. In contrast, wild-type *U2AF1* is required for cell

viability, consistent with the retention of a wild-type allele in human cancers carrying

common *U2AF1* mutations.

454
455

## Discussion

In this report, we have examined the mechanistic and phenotypic consequences

of the common *U2AF1*S34F mutation. Our data demonstrate that the S34F:WT ratio

controls the quantitative consequences of the *U2AF1* mutation for splice site

recognition, and suggest that differential RNA-binding affinities of the mutant and wild-

type protein result in preferential recognition of specific 3' splice sites. Moreover, our

finding that wild-type *U2AF1* is required for cell survival irrespective of the presence or

absence of *U2AF1*S34F explains the genetic observation that tumors always retain an

expressed copy of the wild-type allele. Finally, we expect that the genetic models of

*U2AF1*S34F that we derived from immortalized lung epithelial cells, as well as cell lines

derived from lung adenocarcinomas with the mutation, will prove useful for future

functional studies of this common mutation.

**469  The S34F:WT ratio controls S34F-associated splicing**

U2AF1S34F is known to induce specific splicing alterations, but it is not known

how these changes are regulated. We show that the ratio of S34F:WT U2AF1 gene

products is a critical determinant of the magnitude of S34F-associated splicing. This

conclusion was demonstrated in an isogenic lung epithelial cell line engineered to

express *U2AF1*S34F from one of the two endogenous *U2AF1* loci, and was further

supported by analyses of human LUAD transcriptomes carrying the *U2AF1*S34F allele.

These results suggest that wild-type U2AF1 antagonizes the splicing program

associated with the S34F mutation.

**Altered RNA binding affinities often explain S34F-associated splicing changes**

We find that a major functional difference between purified S34F mutant and

wild-type U2AF1 proteins resides in altered binding affinities for a subset of 3' splice

sites. The trends in the RNA sequence preferences of S34F-mutant U2AF1 are

consistent with the preferred 3' splice sites of S34F-affected transcripts (here and in [17-

20,28]), which we use as the signature of S34F-associated differential splicing (Fig. 1A).

For oligonucleotides that showed significant changes in binding affinities, the S34F

mutation typically reduced or enhanced respective binding of the U2AF1 splicing factor

complexes to 3' splice sites preceded by a -3U or -3C (Fig. 5 and [18]). In support of our

findings for the relevant ternary complex of human U2AF1, U2AF2 and SF1 subunits,

recent studies confirmed that the corresponding S34F mutation inhibited binding of the

minimal fission yeast U2AF heterodimer to a "UAG" splice site RNA [21]. These S34F-

altered RNA affinities are consistent with the location of the substituted amino acid in a

zinc knuckle that may directly contact RNA [17]. Although the effects of the S34F

mutation on binding 3' splice sites preceded by -3A are variable, extrusion or alternative

U2AF1 binding sites for disfavored nucleotides could occur in different sequence

contexts by analogy with other RNA binding proteins [29,30].

Several lines of evidence support the idea that U2AF1S34F is capable of

initiating pre-mRNA splicing once it binds to RNA. Nuclear extracts of cells

overexpressing mutant *U2AF1* can support *in vitro* splicing reactions more efficiently

than nuclear extracts derived from cells overexpressing wild-type *U2AF1* for a minigene

with a specific 3' splice site sequence [17]. In addition, mutant U2AF1 can compensate

for loss of the wild-type factor for the inclusion of some U2AF1-dependent cassette

exons [31]. Our observations that the altered RNA-binding affinities correlate well with

S34F-associated splicing for the majority of splice sites that we tested further suggest

that mutant and wild-type U2AF1 are functionally equivalent for downstream steps of

splicing (Fig. 5 and [18]).

**A working model of S34F-associated Splicing**

In light of our findings and existing evidence from the literature, we propose a

model wherein mutant U2AF1 drives differential splicing by favoring the recognition of

one of two competing 3' splice sites (Supplemental Fig. S18). This model is motivated

by three key facts. First, alternative splicing, in contrast to constitutive splicing,

necessarily results from implicit or explicit competition between splice sites. (For

example, cassette exon recognition can involve competition between the 3' splice sites

of the cassette exon itself and a downstream constitutive exon.) Second, mutant and

wild-type U2AF1 complexes have different binding specificities, largely due to their

preferences for distinct nucleotides at the -3 position, that lead them to preferentially

bind distinct 3' splice sites. Third, mutant and wild-type U2AF1 are likely functionally

equivalent once they bind to RNA. Therefore, altering the cellular ratio of mutant and

wild-type U2AF1 changes the amount of total U2AF1 bound to a given 3' splice site in a

sequence-specific manner, thereby promoting or repressing that splice site relative to a

competing 3' splice site.

Our proposed model is not exclusive of other possible contributing effects, such

as competitive binding of mutant and wild-type U2AF1 to a factor with low stoichiometry

(Supplemental Fig. S18), or effects of U2AF1S34F on the kinetics of co-transcriptional

splicing as suggested recently [32]. Future studies are needed to resolve these

possibilities.

**How are cells with the U2AF1 S34F mutation selected during oncogenesis?**

*U2AF1*S34F is recurrently found in LUAD, other solid tumors, and myeloid

disorders, suggesting that the mutant allele confers a physiological property that

provides a selective advantage during neoplasia. A gene ontology (GO) analysis for

genes that show S34F-associated alterative splicing in HBEC3kt-derived isogenic cells

shows significant alterations in many biological processes such as mRNA processing,

RNA splicing, G2/M transition of mitotic cell cycle, double-strand break repair and

organelle assembly (FDR-adjusted p-values < 0.001). However, we did not observe

signs of neoplastic transformation or changes in cell proliferation after introducing

*U2AF1*S34F into the endogenous *U2AF1* locus in HBEC3kt cells (Supplemental Fig.

S13). Moreover, targeted inactivation of *U2AF1*S34F in LUAD cell lines does not

diminish, and in one case even increases, clonogenic growth in culture (Fig. 6).

The lack of a testable cellular phenotype has been a major hindrance to

understanding the functional significance of mutant *U2AF1* in carcinogenesis. Cell

proliferation is only one of the many hallmarks of cancer, so careful examination of other

cell properties in the isogenic cells may be needed to establish the presumptive role of

*U2AF1*S34F in carcinogenesis.

More recently, after completion of our study [33], Park *et al* reported that

tumorigenic cells emerge after *U2AF1*S34F is ectopically produced in Ba/F3 pro-B cells

or in an immortalized line of small airway epithelial cells [34]. The authors attributed the

transformation events by mutant U2AF1 to the consequences of altered 3' processing of

mRNA's. In particular, they observed an increase in the length of the 3' untranslated

region of *ATG7* mRNA and a reduction in the amount of ATG7 protein, proposing that

the anticipated defect in autophagy predisposes cells to mutations, some of which are

transforming. This "hit-and-run" type of mechanism is consistent with our observation

and theirs that mutant U2AF1 is dispensable for maintenance of the transformed

phenotype in LUAD cell lines (Figure 6) and in their cell lines [34]. Their observations do

not, moreover, exclude a role for S34F-associated splicing in the oncogenic

mechanism.

**Can cancer cells carrying the S34F mutation be targeted therapeutically?**

We have shown that the wild-type *U2AF1* allele is absolutely required for the

growth of lung epithelial and LUAD cells that carry a mutant *U2AF1* allele (Fig. 6). This

result indicates that mutant U2AF1 cannot complement a deficiency of wild-type U2AF1;

it may also explain why tumors homozygous for the *U2AF1*S34F mutation have not

been observed, although the number of tumors found to have even a heterozygous

mutation is still relatively small, so the analysis may not be adequately powered. Still,

the frequent occurrence of a low ratio of S34F:WT *U2AF1* mRNA, accompanied by

increased copy number of wild-type *U2AF1* alleles in lung adenocarcinoma cell lines

and possibly LUADs, suggests that there may be selection for a lower ratio of S34F:WT

in addition to the likely selection, perhaps at an earlier stage of tumorigenesis, for the

S34F mutant.

These results are consistent with a study of mutations affecting the splicing factor

SF3B1, which reported that cancer cells harboring recurrent *SF3B1* mutations also

depend on wild-type SF3B1 for growth [35]. Finally, a recent study similarly found that a

wild-type copy of *SRSF2* is required for leukemic cell survival, and that *SRSF2*

mutations generated a therapeutic index for treatment with a small molecule that inhibits

3' splice site recognition [36]. Together, our results and these recent studies provide a

genetic rationale for targeting wild-type splicing factors (or the splicing machinery more

generally) in cancers harboring spliceosomal mutations.

## Materials and Methods

### *Cell culture, reagents, and assays*

The HBEC3kt cells (a gift from Dr. John Minna, UT Southwestern Medical

Center), H441, A549 (ATCC) and HCC78 cells (DMSZ) were cultured as previously

described [22] or according to vendors' instructions. The primary antibodies for

immunoblots are: rabbit anti U2AF1 (1:5000, # NBP1-19121, Novus), rabbit anti GFP

(1:5000, #A-11122, Invitrogen), mouse anti ACTB (1:5000, Clone 8H10D10, Cell

Signaling). Lentiviruses were produced and titered in HEK293T cells as previously

described [37]. An MOI (multiplicity of infection) of 1 – 5 were used for all assays.

Total RNA was extracted and reverse transcribed as previously described [38].

Splicing alterations were measured by quantitative PCR on reverse-transcribed cDNA

(RT-qPCR), using isoform-specific primers (Supplemental Table S7). These primers

were designed following a previously described method [39]. The PCR efficiency and

specificity of each primer set were determined before they were used for measuring

splicing changes (See Supplemental Materials and Methods for details).

Clonogenic assay was performed by infecting cells with lentiviruses two days

before seeding them into 100 mm dishes (1000 live cells per dish) to grow colonies.

Growth media were supplemented with puromycin (1 µlg/ml) for selecting infected cells

and were changed once a week for up to three weeks. Cell colonies were stained with

0.03% methylene blue (in 20% methanol) for 5 min. Clonogenicity is defined as colony

numbers formed as a percent of those in control cells.

600   Details of all DNA constructs used in the study and the genome editing

601 approaches for creating the *U2AF1*S34F allele and allelic-specific disruption of *U2AF1*

602 are described in the Supplemental Materials and Methods.

604 ***mRNA sequencing and analysis***

605   High throughput mRNA sequencing (RNA-seq) was conducted in the Sequencing

606 Facility of the National Cancer Institute. RNA quality was assessed by 2100 Bioanalyzer

607 (Agilent). Total RNA with good integrity values (RIN > 9.0) was used for poly A selection

608 and library preparation using the Illumina TruSeq RNA library prep kit. Two or four

609 samples were pooled per lane and ran on the HiSeq2000 or HiSeq2500 instrument

610 using TruSeq V3 chemistry. All samples were sequenced to the depth of 100 million

611 pair-end 101 bp reads per sample.

612   Splicing analysis of RNA-seq data from the TCGA LUAD cohort as well as

613 engineered HBEC3kt, H441, and HCC78 cell lines was performed as previously

614 described [17]. A brief description of the method was provided in the Supplemental

615 Materials and Methods.

617 ***Purification of U2AF1 protein complexes and RNA affinity measurement***

618   Purification of U2AF1 complexes, as illustrated in Fig.5, was explained in

619 Supplemental Materials and Methods. Sequences of synthetic 5′-labeled fluorescein

620 RNAs (GE Healthcare Dharmacon) and binding curves are given in the Supplementary

621 Fig. S12. Apparent equilibrium affinity constant of the purified U2AF1 complexes with

622 RNA was measured based on changes in fluorescence anisotropy as previously

described [18]. A brief description of this method was provided in the Supplemental

Materials and Methods.

***Statistics***

All experiments were independently performed at least three times unless

otherwise stated. Statistical significance was determined by two-tailed Student's *t* test or

otherwise stated. In all analyses, *p* values ≤ 0.05 are considered statistically significant.

## Acknowledgements

**Acknowledgements**

We thank Ms. Jackie Idol, Ursula Harper, Danielle Miller-O'Mard and the

transgenic mouse core at NHGRI for technical assistance, Dr. Heidi Dvinge (FHCRC)

for help with the LUAD data analysis, Dr. Matthew J. Walter (Wash.U.) for sharing the

*CEP164* and *FMR1* splice site sequences prior to publication. We thank members of the

Varmus lab, Drs. Janine Ilagan (FHCRC) and Paul Liu (NHGRI) for helpful discussions

during the course of the study. HV was supported by the Intramural Program at the

National Institutes of Health and is now supported by the Meyer Cancer Center at Weill

Cornell Medicine. RKB is supported by the Edward P. Evans Foundation, Ellison

Medical Foundation (AG-NS-1030-13), NIH/NHLBI (R01 HL128239), and NIH/NIDDK

(R01 DK103854). CLK is supported by the Edward P. Evans Foundation and

NIH/NIGMS (R01 GM070503). The results published here are in part based upon data

generated by the TCGA Research Network: http://cancergenome.nih.gov/.

## **References**

1. Yoshida K, Sanada M, Shiraishi Y, Nowak D, Nagata Y, Yamamoto R, et al. Frequent pathway mutations of splicing machinery in myelodysplasia. Nature. 2011;478: 64–69. doi:10.1038/nature10496

2. Graubert TA, Shen D, Ding L, Okeyo-Owuor T, Lunn CL, Shao J, et al. Recurrent mutations in the U2AF1 splicing factor in myelodysplastic syndromes. Nature Genetics. 2012;44: 53–57. doi:10.1038/ng.1031

3. Imielinski M, Berger AH, Hammerman PS, Hernandez B, Pugh TJ, Hodis E, et al. Mapping the hallmarks of lung adenocarcinoma with massively parallel sequencing. Cell. 2012;150: 1107–1120. doi:10.1016/j.cell.2012.08.029

4. Furney SJ, Pedersen M, Gentien D, Dumont AG, Rapinat A, Desjardins L, et al. SF3B1 Mutations Are Associated with Alternative Splicing in Uveal Melanoma. Cancer Discovery. 2013;3: 1122–1129. doi:10.1158/2159-8290.CD-13-0330

5. Wang L, Lawrence MS, Wan Y, Stojanov P, Sougnez C, Stevenson K, et al. SF3B1and Other Novel Cancer Genes in Chronic Lymphocytic Leukemia. N Engl J Med. 2011;365: 2497–2506. doi:10.1056/NEJMoa1109016

6. Waterfall JJ, Arons E, Walker RL, Pineda M, Roth L, Killian JK, et al. High prevalence of MAP2K1 mutations in variant and IGHV4-34-expressing hairy-cell leukemias. Nature Genetics. 2014;46: 8–10. doi:10.1038/ng.2828

7. Cancer Genome Atlas Network. Comprehensive molecular portraits of human breast tumours. Nature. 2012;490: 61–70. doi:10.1038/nature11412

8. Biankin AV, Waddell N, Kassahn KS, Gingras M-C, Muthuswamy LB, Johns AL, et al. Pancreatic cancer genomes reveal aberrations in axon guidance pathway genes. Nature. 2012;491: 399–405. doi:10.1038/nature11547

9. Cancer Genome Atlas Research Network. Comprehensive molecular profiling of lung adenocarcinoma. Nature. 2014;511: 543–550. doi:10.1038/nature13385

10. Ruskin B, Zamore PD, Green MR. A factor, U2AF, is required for U2 snRNP binding and splicing complex assembly. Cell. 1988;52: 207–219.

11. Zamore PD, Green MR. Identification, purification, and biochemical characterization of U2 small nuclear ribonucleoprotein auxiliary factor. Proc Natl Acad Sci U S A. 1989;86: 9243–9247.

12. Krämer A, Utans U. Three protein factors (SF1, SF3 and U2AF) function in pre-splicing complex formation in addition to snRNPs. The EMBO Journal. 1991;10: 1503–1509.

13. Berglund JA, Abovich N, Rosbash M. A cooperative interaction between U2AF65

and mBBP/SF1 facilitates branchpoint region recognition. Genes & Development.
1998;12: 858–867.

14.   Merendino L, Guth S, Bilbao D, Martínez C, Valcárcel J. Inhibition of msl-2
splicing by Sex-lethal reveals interaction between U2AF35 and the 3' splice site
AG. Nature. 1999;402: 838–841. doi:10.1038/45602

15.   Wu S, Romfo CM, Nilsen TW, Green MR. Functional recognition of the 3' splice
site AG by the splicing factor U2AF35. Nature. 1999;402: 832–835.
doi:10.1038/45590

16.   Zorio DA, Blumenthal T. Both subunits of U2AF recognize the 3' splice site in
Caenorhabditis elegans. Nature. 1999;402: 835–838. doi:10.1038/45597

17.   Ilagan JO, Ramakrishnan A, Hayes B, Murphy ME, Zebari AS, Bradley P, et al.
U2AF1 mutations alter splice site recognition in hematological malignances.
Genome Res. 2015;25: 14–26. doi:10.1101/gr.181016.114

18.   Okeyo-Owuor T, White BS, Chatrikhi R, Mohan DR, Kim S, Griffith M, et al.
U2AF1 mutations alter sequence specificity of pre-mRNA binding and splicing.
Leukemia. 2015;29: 909–917. doi:10.1038/leu.2014.303

19.   Brooks AN, Choi PS, de Waal L, Sharifnia T, Imielinski M, Saksena G, et al. A
Pan-Cancer Analysis of Transcriptome Changes Associated with Somatic
Mutations in U2AF1 Reveals Commonly Altered Splicing Events. Ast G, editor.
PLoS ONE. 2014;9: e87361. doi:10.1371/journal.pone.0087361.s008

20.   Przychodzen B, Jerez A, Guinta K, Sekeres MA, Padgett R, Maciejewski JP, et al.
Patterns of missplicing due to somatic U2AF1 mutations in myeloid neoplasms.
Blood. 2013;122: 999–1006. doi:10.1182/blood-2013-01-480970

21.   Yoshida H, Park S-Y, Oda T, Akiyoshi T, Sato M, Shirouzu M, et al. A novel 3′
splice site recognition by the two zinc fingers in the U2AF small subunit. Genes &
Development. 2015;29: 1649–1660. doi:10.1101/gad.267104.115

22.   Ramirez RD, Sheridan S, Girard L, Sato M, Kim Y, Pollack J, et al.
Immortalization of human bronchial epithelial cells in the absence of viral
oncoproteins. Cancer Res. 2004;64: 9027–9034. doi:10.1158/0008-5472.CAN-04-
3703

23.   Yusa K. Seamless genome editing in human pluripotent stem cells using custom
endonuclease-based gene targeting and the piggyBac transposon. Nat Protoc.
2013;8: 2061–2078. doi:10.1038/nprot.2013.126

24.   Kralovicova J, Vorechovsky I. Allele-specific recognition of the 3' splice site of INS
intron 1. Hum Genet. 2010;128: 383–400. doi:10.1007/s00439-010-0860-1

25.   Liu Z, Luyten I, Bottomley MJ, Messias AC, Houngninou-Molango S, Sprangers

R, et al. Structural basis for recognition of the intron branch site RNA by splicing
factor 1. Science. 2001;294: 1098–1102. doi:10.1126/science.1064719

26.  Agrawal AA, Salsi E, Chatrikhi R, Henderson S, Jenkins JL, Green MR, et al. An
extended U2AF(65)-RNA-binding domain recognizes the 3' splice site signal. Nat
Commun. 2016;7: 10950. doi:10.1038/ncomms10950

27.  Forbes SA, Beare D, Gunasekaran P, Leung K, Bindal N, Boutselakis H, et al.
COSMIC: exploring the world's knowledge of somatic mutations in human cancer.
Nucleic Acids Res. 2015;43: D805–11. doi:10.1093/nar/gku1075

28.  Shirai CL, Ley JN, White BS, Kim S, Tibbitts J, Shao J, et al. Mutant U2AF1
Expression Alters Hematopoiesis and Pre-mRNA Splicing In Vivo. Cancer Cell.
Elsevier Inc; 2015;27: 631–643. doi:10.1016/j.ccell.2015.04.008

29.  Lu G, Hall TMT. Alternate modes of cognate RNA recognition by human PUMILIO
proteins. Structure. 2011;19: 361–367. doi:10.1016/j.str.2010.12.019

30.  Miller MT, Higgin JJ, Hall TMT. Basis of altered RNA-binding specificity by PUF
proteins revealed by crystal structures of yeast Puf4p. Nat Struct Mol Biol.
2008;15: 397–402. doi:10.1038/nsmb.1390

31.  Shao C, Yang B, Wu T, Huang J, Tang P, Zhou Y, et al. Mechanisms for U2AF to
define 3' splice sites and regulate alternative splicing in the human genome. Nat
Struct Mol Biol. 2014;21: 997–1005. doi:10.1038/nsmb.2906

32.  Coulon A, Ferguson ML, de Turris V, Palangat M, Chow CC, Larson DR. Kinetic
competition during the transcription cycle results in stochastic RNA processing.
Elife. 2014;3. doi:10.7554/eLife.03939

33.  Fei DL, Motowski H, Chatrikhi R, Prasad S, Yu J, Gao S, et al. Wild-type U2AF1
antagonizes the splicing program characteristic of U2AF1-mutant tumors and is
required for cell survival. bioRxiv. 2016 Apr. doi:10.1101/048553

34.  Park SM, Ou J, Chamberlain L, Simone TM, Yang H, Virbasius C-M, et al.
U2AF35(S34F) Promotes Transformation by Directing Aberrant ATG7 Pre-mRNA
3. Mol Cell. Elsevier Inc; 2016;: 1–34. doi:10.1016/j.molcel.2016.04.011

35.  Zhou Q, Derti A, Ruddy D, Rakiec D, Kao I, Lira M, et al. A Chemical Genetics
Approach for the Functional Assessment of Novel Cancer Genes. Cancer Res.
2015;75: 1949–1958. doi:10.1158/0008-5472.CAN-14-2930

36.  Lee SC-W, Dvinge H, Kim E, Cho H, Micol J-B, Chung YR, et al. Modulation of
splicing catalysis for therapeutic targeting of leukemia with mutations in genes
encoding spliceosomal proteins. Nat Med. 2016. doi:10.1038/nm.4097

37.  Fei DL, Sanchez-Mejias A, Wang Z, Flaveny C, Long J, Singh S, et al. Hedgehog
Signaling Regulates Bladder Cancer Growth and Tumorigenicity. Cancer Res.

2012;72: 4449–4458. doi:10.1158/0008-5472.CAN-11-4123

38.   Fei DL, Li H, Kozul CD, Black KE, Singh S, Gosse JA, et al. Activation of
Hedgehog signaling by the environmental toxicant arsenic may contribute to the
etiology of arsenic-induced tumors. Cancer Res. 2010;70: 1981–1988.
doi:10.1158/0008-5472.CAN-09-2898

39.   Brosseau JP, Lucier JF, Lapointe E, Durand M, Gendron D, Gervais-Bird J, et al.
High-throughput quantification of splicing isoforms. RNA. 2010;16: 442–449.
doi:10.1261/rna.1877010

 **Figure Legends**

**Fig. 1. S34F-associated splicing program correlates with S34F:WT mRNA ratios in**

**LUAD.**

**(A)**. Different consensus 3' splice site sequences preceding cassette exons for two

representative LUADs with the S34F mutation computed from published TCGA

transcriptomes. In both cases, changes in the use of cassette exons were determined

by comparisons with an average transcriptome from LUADs without *U2AF1* mutations.

Boxes highlight nucleotides found preferentially at the -3 position. The nucleotide

frequencies preceding exons that are more often included or more often excluded in

tumors with the S34F mutation differ from the genomic consensus in the tumor from

patient #7903, which represent "typical S34F" consensus 3' splice sites, but not in the

tumor from patient #7727, which represent "quasi-WT" consensus 3' splice sites.

(Consensus 3' splice sites from transcriptomes of every S34F-mutant LUAD are

presented in Supplemental Fig. S1.) The analysis was restricted to introns with

canonical GT-AG U2-type splice sites. The invariable AG at the 3' splice site was not

plotted to scale in order to highlight the consensus sequences at the -3 position. The

vertical axis represents the information content in bits (the maximal value is two. Zero to

one bit is shown). n is the number of cassette exon sequences used to construct the

logo. A cartoon illustrating alternative splicing of a cassette exon (black box) is shown

on the left side of the sequence logo. Black lines over splice junctions illustrate the

S34F-promoted isoform for each comparison. **(B)**. *U2AF1*-mutant LUAD transcriptomes

harboring "typical S34F" consensus 3' splice sites have relatively high S34F:WT mRNA

ratios. *U2AF1*-mutant LUAD samples were grouped based on the nature of the

consensus 3' splice sites. The asterisk represents a statistically significant change by

Student's t test. **(C)**. S34F:WT *U2AF1* mRNA ratios do not correlate with tumor purity in

LUAD tumors with the S34F mutation. Tumor purity is represented by the percent of

tumor nuclei in each LUAD sample (derived from TCGA clinical data) and plotted

against the S34F:WT mRNA ratios. **(D)**. Inclusion of the *STRAP* cassette exon

correlates with the S34F:WT mRNA ratio. Same as Panel **C** but the S34F:WT mRNA

ratio is plotted against the inclusion frequency for the *STRAP* cassette exon. The

median inclusion level of the same cassette exon for all transcriptomes from tumors

without a *U2AF1* mutation (S34F:WT mRNA ratio = zero) is shown as a triangle. r,

Pearson's correlation coefficient. In panels B – D, circles represent samples with typical-

S34F consensus 3' splice site sequences; squares represent samples with quasi-WT

consensus 3' splice site sequences. Colors indicate *U2AF1* copy number status as

calculated by GISTIC 2.0 (See details in the Supplemental Materials and Methods):

black, diploid; blue, shallow deletion; red, gain.

**Fig. 2. Creation of isogenic lung cell lines that recapitulate features of S34F-**

**associated splicing.**

**(A)**. Strategy to create a TCT to TTT point mutation (S34F) at the endogenous *U2AF1*

locus in HBEC3kt cells. TALEN, transcription activator-like effector nuclease; E, exon;

mE, mutant (S34F) exon; ITR, inverted terminal repeat; HA, homology arm. See Results

and Supplemental Materials and Methods for details. **(B)**. MUT1a and MUT1b cells

contain similar levels of mutant and wild-type *U2AF1* mRNA. The number of reads

supporting mutant or wild-type *U2AF1* was obtained from RNA-seq, using poly(A)-

selected RNA from the four cell lines. **(C).** The S34F-associated cassette exons in

*ASUN* and *STRAP* mRNAs show decreased inclusion in MUT cell lines. (Top) Scheme

of alternative splicing with a cassette exon (black box) to generate short and long

isoforms in which the cassette exon is excluded or included. (Bottom) Alternative

splicing of cassette exons in *ASUN* and *STRAP* mRNAs, measured by RT-qPCR using

isoform-specific primers. The short/long isoform ratio in WT1 cells was arbitrarily set to

1 for comparison. Asterisks represent statistical significant changes as compared to that

in WT1 cells. Error bars represent s.e.m. (standard error of the mean) (n = 4). **(D).** Heat

map depicting the inclusion levels of all cassette exons that showed at least a 10%

change in use among the cell lines. The dendrogram was constructed from an

unsupervised cluster analysis. **(E).** Sequence logos from 3' splice sites preceding

cassette exons with altered inclusion in MUT1a cells display typical S34F consensus 3'

splice sites. Logos were constructed as in Fig. 1A based on the transcriptome of MUT1a

cells in comparison with that of WT1 cells. Other comparisons of transcriptomes from

MUT and WT cell lines yielded similar sequence logos.

**Fig. 3. The ratio of S34F:WT U2AF1 gene products controls S34F-associated**

**splicing in isogenic HBEC3kt cell lines.**

**(A)**. Reduction of both mutant and wild-type U2AF1 RNA and protein does not affect

S34F-associated splicing. WT1 and MUT1a cells were transduced with shRNAs against

*U2AF1* (shU2AF1#1 and #4) or two control shRNAs (shScbr, an scrambled shRNA;

shGFP, an shRNA against *GFP*). Total RNA and protein were harvested 4 days later.

The frequencies of incorporation of cassette exon sequences in *ASUN* and *STRAP*

mRNAs (top and middle panels) were determined by the relative short/long isoform

ratios by RT-qPCR, as represented in Fig. 2C. Immunoblots for U2AF1 and ACTB in

total cell lysates are shown in the bottom panel. Asterisks represent statistical significant

changes as compared to shScbr-transduced condition in each cell line. Error bars

represent s.e.m (n = 3). **(B)**. Overexpression of wild-type or mutant *U2AF1* to change

S34F:WT ratios alters S34F-sensitive splicing.  WT1 and MUT1a cells were transduced

with expression vectors encoding *GFP*, wild-type (WT) or mutant (S34F) *U2AF1* for 3

days before harvesting cells to quantify the level of splicing changes and proteins as in

panel A. Asterisks represent statistical significant changes as compared to GFP-

transduced condition in each cell line. Error bars represent s.e.m (n = 3). **(C)**. Disruption

of WT *U2AF1* by gene editing to increase S34F:WT ratios enhances S34F-sensitive

splicing. WT1 and MUT1a cells were transduced with Cas9 and either sgRNA-GFP or

sgRNA-WT. Total RNA and protein were harvested 6 days later for assays as in panel

**A**. Asterisks represent statistical significant changes as compared to Cas9 and sgRNA-

GFP-transduced condition in each cell line. Error bars represent s.e.m. (n = 3).

**Fig. 4. Increasing the ratio of S34F:WT gene products by disrupting the wild-type**

***U2AF1* locus enhances S34F-associated splicing of cassette exons.**

**(A)**. RNA-seq was performed for WT1 and MUT1a cells transduced with Cas9 and

either sgRNA-GFP or sgRNA-WT.  Upper panel: The Venn diagrams show overlap of

290 cassette exons that display at least a ten percent increase in inclusion levels in

MUT1a cells relative to levels in WT1 cells, with or without CRISPR-Cas9-mediated

disruption of wild-type *U2AF1*. Lower panel: Increasing the S34F:WT ratio by disrupting

wild-type *U2AF1* enhances the magnitude of S34F-associated splicing. The "waterfall"

plot depicts changes in percent inclusion levels for all shared cassette exons identified

from the Venn diagram when the wild-type *U2AF1* locus was disrupted. Each vertical

bar represents one shared cassette exon. **(B)**. The analysis shown in Panel A was

repeated for cassette exons showing ten percent or more decreased inclusion in

MUT1a cells. **(C)**. Heat map depicting the inclusion levels of all cassette exons that

showed at least a 10% change in use among the treatment conditions. The dendrogram

was constructed from an unsupervised cluster analysis. (**D**). Enhanced features of

S34F-associated logos at 3' splice acceptor sites after disruption of wild-type *U2AF1*.

Sequence logos were constructed as in Fig. 1A, based on the indicated comparisons.

**Fig. 5. Differential binding of mutant and wild-type U2AF1 complexes to RNA**

**oligonucleotides explains most S34F-associated alterations in RNA splicing.**

**(A)**. Cartoons illustrate components of recombinant U2AF1 complexes used in the

binding assay. Full-length proteins are shown but only partial sequences (denoted by bi-

directional arrows) were used to make recombinant protein complexes. KH-QUA2, K-

Homology Quaking motif; RRM, RNA recognition motif domain; RS, arginine/serine-rich

domain; UHM, U2AF homology motif; ULM, U2AF ligand motif; ZnF, zinc finger domain.

**(B)**. Scheme of alternative splicing patterns for cassette exons (top diagram) and 5'

extended exons from competing 3' splice site selection (bottom). The brackets indicate

the positions of RNA oligonucleotides used for the binding assays. Exons are shown as

boxes: white boxes indicate invariant exonic sequences and black boxes denote

sequences that are incorporated into mRNA (exonic) only when the proximal 3' splice

sites are used. Introns are shown as solid lines. The grey lines represent possible

splices. The names of characterized genes that conform to the patterns shown in the

upper and lower cartoons are indicated. **(C - H)**. Mutant and wild-type U2AF1

complexes have different affinities ($K_A$'s) for relevant 3' splice site oligonucleotides. To

accomplish the binding assays, the wild-type or mutant U2AF1 protein complexes were

titrated into 5' fluorescein-tagged RNA oligonucleotides over a range of concentrations

as described in the Supplemental Materials and Methods. RNA sequences from -4 to +3

relative to the 3' splice sites (vertical lines) in proximal and distal positions are shown.

The nucleotide at the -3 position is bolded and underlined. Empty bars, $K_A$ for WT

U2AF1 complex; grey bar, $K_A$ for mutant U2AF1 complex. For ease of comparison

between the affinity binding results with S34F-associated splicing, S34F-promoted

splices, as determined from RNA-seq data, are shown on top of each bar graph in black

lines. The fitted binding curves, full oligonucleotide sequences, and apparent equilibrium

dissociation constants are shown in Supplemental Fig. S12. The relative changes in

affinity and use of proximal *versus* distal splice sites are summarized in Supplemental

Table S3. The asterisk represents a statistically significant change by unpaired t-tests

with Welch's correction. ns, not statistically significant.

**Fig. 6. Wild-type but not mutant *U2AF1* is required for the clonogenic growth of**

**the isogenic HBEC3kt cells and LUAD cell lines.**

**(A)**. Clonogenic growth assays after selective disruption of the wild-type or mutant

*U2AF1* allele. Left panel: The indicated cell lines were transduced with lentiviruses

expressing Cas9 and sgRNA-GFP, sgRNA-S34F or sgRNA-WT, followed by clonogenic

assays. Cell colonies were stained with methylene blue and counted three weeks later.

Right panel: Quantification of the clonogenic assay. The results are shown as percent

clonogenicity by setting the number of control cell colonies (cells transduced with Cas9

and sgRNA-GFP) as 100%. Asterisks represent statistical significant changes as

compared to Cas9 and sgRNA-GFP-transduced condition in each cell line. Error bars

represent s.e.m (n = 3). **(B)**. Rescue of growth inhibition by Cas9 and sgRNA-WT by

overexpressing a form of wild-type *U2AF1* cDNA that is not predicted to be the target for

sgRNA-WT (See Supplemental Fig. S8 and Materials and Methods). Left panel: A549

cells were transduced with a control (DsRed-Express 2) or a wild-type *U2AF1* cDNA

that is not predicted to be the target for sgRNA-WT. Increased expression of wild-type

U2AF1 was confirmed by immunoblot (left bottom panel).  These cells were

subsequently transduced with Cas9 and either sgRNA-GFP or sgRNA-WT followed by

clonogenic assays as in panel A (left upper panel).  Right panel: Quantification as in

Panel A. The asterisk represents a statistical significant change for the indicated

comparison. Error bars represent s.e.m. (n = 3).

**Fei_Fig.2**

**A** Knockdown of total *U2AF1*

**B** Overexpression of WT or mutant *U2AF1*

**C** Disruption of WT *U2AF1*

## A

**Cassette Exons with Increased % Inclusion by *U2AF1*S34F**

## B

**Cassette Exons with Decreased % Inclusion by *U2AF1*S34F**

## C

## D

**A**

**B**