The Genetic Architecture of a Complex Trait is more Sensitive to Genetic Model than Population Growth

Jaleal S. Sanjak[1,2] *, Anthony D. Long[1,2], Kevin R. Thornton[1,2] *,

1 Department of Ecology and Evolutionary Biology, University of California, Irvine, California 92697, USA
2 Center for Complex Biological Systems, University of California Irvine, Irvine, CA 92697, USA

* E-mail: jsanjak@uci.edu (JSS) , krthornt@uci.edu (KRT)

Abstract

The genetic component of complex disease risk in humans remains largely unexplained. A corollary is that the allelic spectrum of genetic variants contributing to complex disease risk is unknown. Theoretical models that relate population genetic processes to the maintenance of genetic variation for quantitative traits may suggest profitable avenues for future experimental design. Here we use forward simulation to model a genomic region evolving under a balance between recurrent deleterious mutation and Gaussian stabilizing selection. We consider three different genetic models, two different population growth models, and several different methods for identifying genomic regions harboring variants associated with complex disease risk. We demonstrate that the genetic model, relating genotype to phenotype, has a qualitative effect on the genetic architecture of a complex trait. In particular, the variance component partitioning across the allele frequency spectrum and the power of statistical tests is more affected by the assumed genetic model than by population growth. Models with incomplete recessivity most closely match the minor allele frequency distribution of significant hits from empirical genome-wide association studies. Such models show little dominance variance, which is consistent with recent empirical estimates of heritability explained by typed markers. We highlight a particular model of incomplete recessivity that is appealing from first principles. Under that model, deleterious mutations at the same gene partially fail to complement one another. Interestingly this gene-based model predicts considerable levels of unexplained variance associated with within locus epistasis. Our results suggest a need for improvement of statistical tools for region based genetic association and heritability estimation.

Author Summary

Gene action determines how mutations affect phenotype. When placed in an evolutionary context, the details of the gene action model can impact the maintenance of genetic variation for complex traits. Likewise, population size changes affect the relative strength of different evolutionary forces influencing patterns of genetic variation. Here, we explore the impact of genetic model and population growth on distribution of

genetic variance across the allele frequency spectrum underlying risk fora complex disease. Using forward-in-time population genetic simulations, we show that the genetic model has a drastic impact on the genetic architecture of a complex disease-trait. We explicitly simulate genome-wide association studies (GWAS) and perform heritability estimation. Each genetic model makes qualitatively distinct predictions for GWAS and heritability mapping experiments. In many cases, the effect of the genetic model has a larger effect on GWAS outcomes than does population expansion. A particular model of gene-based partial recessivity, based on allelic non-complementation, aligns well with empirical results from GWAS. This model is appealing from first principles and shows peculiar behavior with respect to intralocus epistatic variance.

Introduction

Risk for complex diseases in humans, such as diabetes and hypertension, is highly heritable yet the causal DNA sequence variants responsible for that risk remain largely unknown. Genome-wide association studies (GWAS) have found many genetic markers associated with disease risk [1]. However, follow-up studies have shown that these markers explain only a small portion of the total heritability for most traits [2,3].

There are many hypotheses which attempt to explain the 'missing heritability' problem [2–5]. Genetic variance due to epistatic or gene-by-environment interactions is difficult to identify statistically because of, among other reasons, increased multiple hypothesis testing burden [6,7], and could artificially inflate estimates of broad-sense heritability [8]. Well-tagged intermediate frequency variants may not reach genome-wide significance in an association study if they have smaller effect sizes [9,10]. One appealing verbal hypothesis for this 'missing heritability' is that rare alleles of large effect are the causal factors underlying these traits and are difficult to detect [4,11,12]. These hypotheses are not mutually exclusive, and it is probable that a combination of models will be needed to explain all heritable disease risk [13].

Further, the performance of statistical association methods has been shown to be sensitive to population genetic parameters [14,15]. Thus, it is necessary to use theoretical disease-trait models in experimental design and the development of methods of analysis for genetic association studies. However, the field has neither a comprehensive theoretical framework for the simulation of disease-traits in populations nor a firm understanding of the relationship between model structure and predictions. Typically, specific parameters such as effect-size and frequency of causal mutations or degree of linkage disequilibrium are explored under a single modeling framework.

Many attempts to simulate GWAS datasets have been based on sampling from arbitrary statistical models of the relationships between allele frequencies and mutational effect sizes [16–22]. These approaches make large simulated samples feasible to obtain, but they are not the outcome of an evolutionary process, so they may or may not represent a situation that occurs with any frequency in actual disease samples. An empirical approach, sampling haplotypes from known sequences [23], might provide more realism. Yet, it was demonstrated in [24] that such an approach fails to capture the spectrum of variation in a population subjected to recent population growth.

Population genetics approaches to modeling the genetic architecture of diseases are either based in a coalescent or forward simulation framework. In simulating the genetics of a complex trait under a coalescent framework the phenotypic effects of mutations are typically chosen based on hypothetical relationships

between allele frequencies and effect sizes [25–28]. Although, recent work has developed rigorous formulations ₃₁ of quantitative trait distributions under a neutral coalescent framework [29], those approaches have not been ₃₂ applied in the context of GWAS yet. Forward-in-time simulations, as in [24,30–36], explicitly model the ₃₃ allele frequencies and effect sizes of mutations in a genomic region with neutral sites, selected sites and ₃₄ recombination. ₃₅

Within the forward simulation framework, methods differ in their approach to assigning phenotypes to ₃₆ particular genotypes. One approach, based on the work of [37], involves the assumption that a site has an ₃₇ effect on fitness which is pleiotropically related to its effect on phenotype. In that model, by specifying a ₃₈ parameter τ, the user establishes a degree of correlation between fitness and phenotype [15,33–35]. A related ₃₉ approach that builds off of much earlier work by [38], models trait and fitness effects as coming from a ₄₀ bivariate gamma distribution with a specified correlation parameter ρ [36]. In both approaches the ₄₁ disease-trait itself is not a component of fitness and thus standing variation may be dominated by occasional ₄₂ large trait effect mutations with small fitness effects that can reach intermediate frequency. The extent to ₄₃ which this occurs is dependent on the degree of correlation. Furthermore, both approaches indicate that an ₄₄ intermediate degree of correlation between complex disease traits and fitness is most ₄₅ plausible [33,36,39](although [39] is a non-forward simulation based implementation of [37], a similar ₄₆ conclusion was reached.) ₄₇

Another set of studies were mixed in applying versions of explicit quantitative genetic models [24,31], or ₄₈ more arbitrary models designed to obtain certain desirable behaviors [30,32]. As pointed out in [14], allowing ₄₉ the disease-trait to be a component of an individuals phenotype ensures that the distributions of phenotypic ₅₀ values and the frequency of underlying causal variants are shaped by selection. In reality, a particular variant ₅₁ may impact fitness both through the disease-trait and pleiotropically through other traits. However, if those ₅₂ other traits are also under stabilizing selection then their distribution and the frequency of underlying causal ₅₃ variants will be shaped by that process. At this time we do not have a model where each variant contributes ₅₄ a different effect to each of a subset of traits that each contribute to phenotype at a different stage of life ₅₅ history. The approach in [14] is similar to classical models [40–45] of selection on quantitative traits where ₅₆ phenotype is the sum of genetic and environmental components and is subjected to Gaussian stabilizing ₅₇ selection. A key difference between [14] and the classical quantitative trait models is that causal mutations ₅₈ are unconditionally deleterious and the gene action model involved gene-based recessivity,i.e, allelic ₅₉ non-complementation. ₆₀

While the work in [14] presented a new genetic model, it was limited because only that model was ₆₁ explored under a single demographic scenario (constant sized population). To our knowledge, there has not ₆₂ been a joint analysis of the effect of genetic model and demography on the predicted outcomes of GWAS. ₆₃ Thus, we extend the approach of [14] by including a model of recent population expansion and a set of ₆₄ genetic models. As in [14], we simulate a 100 kilobase region of human genome, contributing to a complex ₆₅ disease phenotype and fitness. The region evolves forward in time subject to neutral and deleterious ₆₆ mutation, recombination, selection, and drift. We allow deleterious mutations to contribute to phenotype in ₆₇ one of three ways: additivity (co-dominance), multiplicative recessivity, or gene-based partial recessivity. The ₆₈ two former models are classical models and the latter is that of [14]. The critical conceptual difference ₆₉ between recessive models is whether dominance is a property of a locus (nucleotide/SNP) in a gene or the ₇₀

gene overall. Mathematically this amounts to whether one performs the first operation across alleles at a site or across sites on a haplotype. For completely co-dominant models, this distinction is irrelevant, however for a model with arbitrary dominance one needs to be more specific. As an example, imagine a compound heterozygote for two biallelic loci, i.e. genotype Ab/aB. In the case of traditional multiplicative recessivity the compound heterozygote is wild type for both loci and therefore wild-type over all; this implies that these loci are in different genes (or independent functional units of the same gene) because the mutations are complementary. However, In the case of gene-based recessivity, neither haplotype is wild-type and so the individual is not wild-type; the failure of mutant alleles to complement defines these loci as being in the same gene [46]. Here we show that different models make different predictions about GWAS results and these predictions have strong implications for future research on the genetics of complex traits.

The degree to which demographic history of a population influences the expected architecture of a complex trait remains unclear as various authors have made only partially overlapping conclusions on the subject [34, 47, 48]. One common thread is that recent population expansion, under a genetic model where the disease-trait significantly contributes to fitness, will increase the importance of rare large effect variants. This hypothesis is well supported by empirical studies which suggest that the great majority of variation($>$70%) in the human exome is due to rare mutations (minor allele frequencies (MAF) $<$0.5%) and that the observed patterns of nucleotide diversity are most consistent with a model of recent population expansion with many sites under weak purifying selection [49–55]. These studies also indicate that rare variants are enriched at putatively deleterious sites, implicating a role for rare variants in disease.

The case for the importance of rare variation in the genetics of complex disease is strong. Yet, the construction of a theoretical model which simultaneously predicts the observed patterns of nucleotide diversity and the observed results of GWAS has remained elusive. It was noted by [56] and summarized in [4], that the MAF distribution of SNP markers that reach genome-wide significance in GWAS is not consistent with existing models in which rare variants drive the heritability of disease. Specifically, we do not observe an enrichment of low frequency to rare (MAF$<$ 5%) significant hits in GWAS. These observations were made in comparison to simulations under an additive co-dominant genetic model. This incongruence between theory and empirical data is one major motivation for exploring other genetic models. Each model imposes a different strength of selection on causal sites of different magnitudes. The genetic model also influences the relationship between the distribution of phenotypic and genetic variation in the population. In addition, we were motivated to explore alternative approaches to simulating a gene region in which mutations are recessive.

We find that a model of gene-based recessivity in a recently expanded population reconciles the predicted MAF distribution of significant SNP's with the observed patterns from human GWAS. The gene-based recessive model under recent population expansion predicts that a substantial portion of heritability will be due to rare alleles of large effect; but, not as much as predicted by a co-dominant genetic model. In addition, this model is consistent with recent estimates of total heritability [57], and estimates of additive and dominance heritability [58]. Lastly, we explore the power of several association tests to detect a causal gene region under each genetic and demographic model. We find significant heterogeneity in performance of burden tests [14, 28, 59] across models of the trait and demographic history. The behavior of these tests under our models provides us with insight as to the circumstances in which they are best suited and potentially the

nature of the empirical associations which are revealed by each test. Overall, we demonstrate that there is a   111
need to consider how predictions about the genetic architecture of complex traits are altered by changes in   112
the structure of population genetic models.   113

Results and Discussion   114

To perform genetic association and heritability estimation studies *in silico*, we need to impose a trait onto   115
simulated populations. By explicitly modeling a quantitative trait in a population genetic setting, we   116
introduce assumptions about the molecular basis of a trait. Yet, we do not know how that molecular genetic   117
basis influences population genetic signatures in the genome. To begin addressing this question, it was   118
necessary to restrict ourselves to a small subset of molecular and evolutionary scenarios. We analyzed a set   119
of approaches to modeling a single gene region experiencing recurrent unconditionally-deleterious mutation   120
contributing to a quantitative trait subject to weak Gaussian stabilizing selection. Specifically, we studied   121
three different genetic models and two different demographic models. The details of how the models are   122
constructed are described in the Materials and Methods section. The genetic models are named the additive   123
co-dominant (AC) model, multiplicative recessive (Mult. recessive; MR) model and the gene-based recessive   124
(GBR) model. The MR model has a parameter, h, that controls the degree of recessivity; we call this model   125
the complete MR (cMR) when $h = 0$ and the incomplete MR (iMR) when $0 \leq h \leq 1$. The AC and MR   126
models represent the traditional models used in simulations of population-genetic processes and the evolution   127
of quantitative traits. Further, such models draw no distinction between a "mutation" and a "gene" (as   128
discussed in [14]). The GBR is also a recessive model, but recessivity is at the level of a *haplotype* and is not   129
an inherent property of individual mutations (see [14] for motivation of this model). Viewed in light of the   130
traditional AC and MR models, the recessivity of a *site* in the GBR model is a function of the genetic   131
background on which it is found. Based on several qualitative comparisons we find that the GBR model is   132
approximated by iMR models with $0.1 \leq h \leq 0.25$. However, no specific iMR model seems to match well in   133
all aspects. The demographic models are that of a constant sized population (no growth) and rapid   134
population expansion (growth).   135

The use of the MR model is inspired by Risch's work [60,61], linking a classic evolutionary model of   136
multiple loci interacting multiplicatively [62,63] to the the genetic epidemiological parameter relative risk.   137
Risch's approach was pushed forward in an important study by Risch and Merikangas [64] which determined   138
the power to detect casual risk variants as a function of allele frequency and relative risk under their model.   139
Pritchard extended Risch's model to consider a trait explicitly as a product of the evolutionary process [65].   140
Pritchard's work demonstrated that the equilibrium frequency distribution suggested an important role for   141
rare deleterious mutations when a trait evolves in a constant sized, randomly mating population where   142
mutations of constant effect sizes are recurrent at multiple loci. However, until now, an explicit exploration   143
of the relationship between model structure and predicted outcomes of GWAS had not been done.   144

Heritability and genetic load under population growth   145

Before deeply exploring the predicted genetic architecture of a trait under each model, we looked at two key   146
mean values: total genetic variance and load. The genetic variance underlying a trait is, in part, determined   147

by the outcome of mutation-selection balance. Approximations for the expected genetic variance under $\quad$ 148
models of stabilizing selection with Gaussian mutational effect sizes and additive gene action have been $\quad$ 149
derived previously based on Kingman's "house-of-cards" model [43, 45]. According to those approximations, $\quad$ 150
when the variance of mutational effect sizes is large compared to total genetic variance, the genetic variance $\quad$ 151
will be dependent only on the mutation rate, μ, and the intensity of stabilizing selection, $1/V_s$. Here, μ refers $\quad$ 152
to the total mutation rate in the "gene region" (per gamete, per generation), with mutations arising $\quad$ 153
according to an infinitely-many sites scheme [66]. In a diploid species, $V_G \approx 4\mu_d V_s$ for an additive trait, and $\quad$ 154
$V_G \approx 2\mu V_s$ for a recessive trait [45, 67]. By keeping V_s and μ constant, we can modulate the broad sense $\quad$ 155
heritability ($H^2 = (V_G)/(V_G + V_E)$) by changing the environmental variance, $V_E = \sigma_E^2$. These $\quad$ 156
approximations are expected to hold for any well-behaved probability distribution of mutational effect $\quad$ 157
sizes [68]. Here, as in [14], we draw the effect sizes of causal mutation from an exponential distribution, $\quad$ 158
modeling unconditionally deleterious mutations. As previously shown in [45], for a Gaussian distribution of $\quad$ 159
effect sizes, and in [14] for an exponential distribution, heritability plateaus at the value expected under the $\quad$ 160
HOC approximation when the mean effect size (λ) is large (Fig S1). $\quad$ 161

Previous work, under additive genetic models, on the impact of population growth on the genetic $\quad$ 162
architecture of complex traits suggests that mean heritability is constant under growth [34, 48]. We confirm $\quad$ 163
this in Fig S1, showing that mean broad-sense heritability, $H^2 = (V_G)/(V_P)$, initially increases as λ, the $\quad$ 164
mean effect of a new deleterious mutation, increases and then plateaus to approximately the same level as $\quad$ 165
models with constant population size. This general trend is observed under each genetic model, but the MR $\quad$ 166
model is qualitatively different in its behavior under population growth. The MR model predicts a broad $\quad$ 167
sense heritability under growth of about 90% of constant sized population levels when $\lambda = 0.01$, and 50% $\quad$ 168
when $\lambda = 0.5$ (Fig S2). $\quad$ 169

Genetic burden (load), the average relative deviation from optimum fitness, of the population is also $\quad$ 170
unaffected by recent population history under the AC model (Fig S3), as shown in [34, 48]. We find this same $\quad$ 171
behavior under the GBR model, but not under the MR model. Under rapid population expansion, the load $\quad$ 172
decreases slightly (at most 2%). As λ increases the load increases under the additive model in both $\quad$ 173
demographic scenarios. Load is effectively constant over the range of λ under the GBR model. Increasing the $\quad$ 174
heritability of the trait decreased the magnitude of the genetic load, but had no interaction with the effects $\quad$ 175
of demography or increasing mean effect size. $\quad$ 176

From these results we can conclude that the AC and GBR models are fairly comparable with respect to $\quad$ 177
total genetic variance and genetic load. Therefore, the remaining differences, which we highlight in later $\quad$ 178
sections, between the AC and GBR model can be attributed to the fine scale composition of the genetic $\quad$ 179
variance in the population. In other words, the AC and GBR model differ in how the genetic variance and $\quad$ 180
load are accounted for, despite the total amounts being roughly equivalent. However, the MR model is $\quad$ 181
qualitatively different in it's behavior under population growth. This makes comparison to the MR model $\quad$ 182
somewhat difficult. However, there are important reasons to explore it further. Based on first principles, the $\quad$ 183
application of the MR model in simulation of a single gene region is inappropriate. However, it would be $\quad$ 184
appropriate for simulating each mutation as a variant of a distinct functional genomic unit. It is also the $\quad$ 185
most analytically tractable model of recessivity in population genetics, and as such it is our best reference $\quad$ 186
point for comparison to the GBR model. $\quad$ 187

Choice of genetic model effects key population genetic signatures

It is unlikely that there is a single correct model for all traits. Rather, we must determine how the structure of a model influences certain predetermined important measures. We find that modifying the function that determines how genotype contributes to phenotype, the genetic model, impacts the genetic load and the site frequency spectrum for risk variants. Mean genetic load decreases with degree of dominance (Fig S3). Recessive deleterious mutations segregate to higher frequencies in the population without increasing genetic load (Fig S5). The MR model demonstrates a slight decrease in load under growth (Fig S3). Similarly, the additive model has greater skew and kurtosis for both the number of mutations and the genetic value of a gamete over the range of λ and demographic models (Fig S10 and Fig S11).The increase skew and kurtosis implies that total genetic load in the additive models is dominated by rare large excursions from the population mean.

Population expansion has been shown to impact the site frequency spectrum [50, 52, 69, 70]. In general, we expect to find an increase in rare private mutations under a rapid population expansion scenario. We find all of our models showcase the expected pattern (Fig S7), but there are consistent and important differences between models. In Fig S7, the site frequency spectrum of risk variants from a population sample (n=100) shows a dependency on population growth, mean effect size λ, and genetic model. Population growth increases the proportion of singletons in for all genetic models and values of λ. Increasing the value λ increases the proportion of singletons in each genetic model and demographic scenario, but the increase is qualitatively dependent on genetic model and independent of demography. The recessive models show the strongest dependence on λ. When λ is small the recessive models show fewer singletons as compared to the additive model, but as λ increases the relative proportion of singletons between recessive and additive genetic model increases. When the value of λ is large, the recessive models show more singletons than the additive model. The GBR model shows more singletons than the MR model in all cases. Fig S8 shows the site frequency spectrum for non-risk variants, which demonstrates a dependence on population growth, shifting towards low-frequency sites, but shows no dependence on genetic model or λ. Since the neutral variant site frequency spectrum is consistent across models we determine that the difference in linked selection between models is not important.

Additive and dominance genetic variance in the population

The amount of narrow sense heritability, $h^2 = (V_A)/(V_P)$, explained by variants across the frequency spectrum is directly related to the effect sizes of those variants [71]. Thus, this measure is an important predictor of statistical power of GWAS and should inform decisions about study design and analysis [72]. Empirically, there appears to be very little dominance variance underlying most quantitative traits [58]. The results in [58] are relevant to our aims and any plausible model for quantitative trait variation should be consistent with their work. Thus, it is important that we explore additive and dominance variance across the allelic frequency spectrum in our models. We have a particular interest in the amount of additive variance, V_A, that is due to rare alleles and how much of genetic variance, V_G, is attributable to V_A under different recessive models. Neither of these questions were addressed in [14].

We follow the approach of [48], by calculating the cumulative percent of V_G explained by the additive effects of variants less than or equal to frequency x, $(V_{A;q\leq x})/(V_G)$. The product of this ratio and

broad-sense heritability is an estimate of the narrow-sense heritability of the trait ($h^2 = (V_A)/(V_P)$). In   227
addition we calculate the same distribution for dominance effects ($V_{D;q \leq x}$)/(V_G) using the orthogonal model   228
of [58]. Methods based on summing effect sizes [71] or the site frequency spectrum [48] would not apply to   229
the GBR model, because the effect of a variant is not independent of other variants. Therefore, we resort to   230
a regression based approach, where we regress the genotypes of the population onto the total genetic value as   231
defined in our disease trait models (see METHODS). For consistency, we applied the regression approach to   232
all models. Overall, these distributions are substantially different across genetic models, demographic   233
scenarios and model parameters (Fig 1).   234

Under the AC model, all of V_G is explained by additive effects if all variants are included in the   235
calculation; in Fig 1 the solid variance curves reach unity in the AC panel. Low frequency and rare variants   236
($q < 0.01$) explain a large portion of narrow sense heritability, 26% - 95%, even in models without rapid   237
population expansion. Fig S4 demonstrates that, for the AC model, the variance explained at any given   238
frequency threshold increases asymptotically to unity as a function of increasing λ. While the total   239
heritability of a trait in the population is generally insensitive to population size changes (Fig S1, see   240
also [34, 48, 73]), rapid population growth increases the fraction of additive genetic variation due to rare   241
alleles (Fig 1).   242

Here, increasing λ corresponds to stronger selection against causative mutations, due to their increased   243
average effect size. As the strength of selection increases, we expect rare variants to make larger   244
contributions to V_G [48], which is exactly what we observe. Recent work by Zuk et al., takes a similar   245
approach and relates the allele frequency distribution directly to design of studies for detecting the role of   246
rare variants [47]. However, our findings contrast with those of Zuk [47] and agree with those of   247
Lohmueller [34], in that we predict that population expansion will substantially increase the heritability, or   248
portion of genetic variance, that is due to rare variants. Our results under the AC model agree with those of   249
Simon's et al. in that we find that increasing strength of selection, increasing λ in our work, increases the   250
contribution to heritability of rare variants [48]. However, under the GBR model the distribution of variance   251
over allele frequency as function λ is non-linear (Fig 1 and Fig S4).   252

For all recessive models, $V_A < V_G$ when including all mutations (Fig 1). For the MR models, all   253
additional genetic variation is explained by the dominance component; in Fig 1 the dotted variance curves   254
reach unity in the MR panels. As expected, genetic variation under the MR model with partial recessivity   255
($h = 0.25$) is primarily additive [71, 74], whereas V_G under the cMR model ($h = 0$) is primarily due to   256
dominance. The MR model differ from the third recessive model, the GBR model. This latter model shows   257
little dominance variance and is the only model considered here for which the total V_G explained by $V_A + V_D$ 258
is less than the true V_G for all λ. This can be clearly seen in Fig 1 because the variance curves do not reach   259
unity in the GBR panel. Under the GBR model, large trait values are usually due to compound heterozygote   260
genotypes (*e.g.*, Ab/aB) [14]. Therefore, the recessivity is at the level *of the gene region* while the typical   261
approach to estimating V_A and V_D assigns effect sizes and dominance to individual mutations. Thus,   262
compound heterozygosity, which is commonly observed for Mendelian diseases (see [14] and references   263
therein) would be interpreted as variation due to *interactions* (epistasis) between risk variants. Importantly,   264
the GBR model predicts that such interactions should be local, occurring amongst causal mutations in the   265
same gene. These observations concerning the GBR model are consistent with the finding of [58] that there   266

is little dominance variance underlying complex traits. These observations warrant future investigation, as the GBR model is reflective of the classic definition of a gene in which recessive mutations fail to complement. Because of allelic heterogeneity and the frequency of large effect mutations, it is unclear how to effectively reformulate approaches to estimating heritability to focus on haplotypes rather than sites. If we partitioned haplotypes into two classes, unaffected (A) and affected (a), we might see dominance between classes, but this exercise would not be particularly helpful because we have no way of doing so in practice.

The increase in the number of rare alleles due to population growth is a well established theoretical and empirical result [50, 52, 70, 75–85]. The exact relationship between rare alleles [4, 26, 56, 65, 86], and the demographic and/or selective scenarios from which they arose [34, 48, 87], and the genetic architecture of common complex diseases in humans is an active area research. An important parameter dictating the relationships between demography, natural selection, and complex disease risk is the degree of correlation between a variant's effect on disease trait and it's effect on fitness [33, 34, 37, 48]. In our simulations, we do not impose an explicit degree of correlation between the phenotypic and fitness effects of a variant. Rather, this correlation is context dependent, varying according to the current genetic burden of the population, the genetic background in which the variant is present and random environmental noise. However, if we re-parameterized our model in terms of that of [37], then we would have $0 << \tau << 1$, which is consistent with recent attempts at estimating that parameter [33, 39]. Our approach is reflective of weak selection acting directly on the complex disease phenotype, but the degree to which selection acts on genotype is an outcome of the model. While the recent demographic history has little effect on key mean values such has broad-sense heritability of a trait or population genetic burden (Fig S1 and Fig S3), the structure of the individual components of the population which add up to those mean values varies considerably. The specific predictions with respect to the composition of the populations varies drastically across different modeling approaches. It is therefore necessary to carefully consider the meaning of a genetic model in a simulation study.

Estimating additive and dominance variance from population samples

Based on our results (Fig 1), we expected that approaches for estimating dominance variance that are implicitly based on a site-by-site model should not find significant dominance variance even when the underlying model is recessive at the level of the gene, in agreement with the results in [58]. We hypothesized that the GBR model and intermediate iMR model would not necessarily show dominance variance when calculated using GREMLd. We also wished to explore, in a more general sense, the behavior of techniques for estimating heritability due to a genomic region on our simulated data. We tried three different approaches to estimating heritability from population samples. We tried GREML/GREMLd, minor allele frequency stratified(MS) GREML/GREMLd and MS Haseman-Elston (HE) regression. In all cases we used GCTA [88], to make genetic relatedness matrices (GRM) for both additive and dominance components. For MS estimates we stratified the additive and dominance GRM's into two bins $MAF \leq 0.01$ and $MAF > 0.01$. HE-MS regression was carried out by regressing the off diagonal elements of each GRM onto the cross product of the scaled and centered phenotypes in a multiple regression setting using R [89]. Unfortunately, due to the nature of our simulated data GREML/GREMLd-MS did not result in sufficiently reliable results. Under GREML/GREMLd-MS, many replicates resulted in numerical errors in GCTA. These problems were present at a rate of less than 1/100 replicates using non-MS GREML/GREMLd, but were exacerbated by

splitting the data into multiple GRMs. This is likely a result of some samples having one or more of the ⁣ 306
following problems: low numbers of SNPS, low total genetic variance, or extremely uneven distribution of ⁣ 307
SNP effect sizes. At large values of λ, especially under non-AC models, the realized heritability in sample ⁣ 308
often does not reflect the heritability in the population (Fig S17). Despite known bias induced in non-MS ⁣ 309
GREML/GREMLd due to differences in underlying allele frequency distribution between typed and causal ⁣ 310
SNPs, we present non-MS GREML/GREMLd and HE-MS results. ⁣ 311

Earlier, we showed how genetic variance is distributed over allele frequency in our simulated populations. ⁣ 312
Here we want to understand how methods for estimating total genetic variance using population samples ⁣ 313
perform on our simulated data. Recent analysis of a large pool of quantitative trait variation data suggests ⁣ 314
that dominance variance is rarely a significant contributor to total heritability [58]. Our analysis of additive ⁣ 315
genetic variance explained as function of allele frequency suggested a significant presence of non-additive ⁣ 316
heritability in some recessive models (Fig 1); this observation called into question the relevance of our ⁣ 317
simulations to natural variation in quantitative traits. Other recent work has found that when genotyping is ⁣ 318
sufficiently thorough the typed markers account for almost all heritability for human height and body mass ⁣ 319
index [57]. Importantly, [57] found that one must include nearly all variants to get accurate estimates and ⁣ 320
that significant portion of the genetic variance is due to rare alleles. We calculated GREMLd estimates of ⁣ 321
additive and dominance heritability in random population sample panels under the AC, MR ($h = 0, 0.25$) ⁣ 322
and GBR models. ⁣ 323

In Fig S12 we show the GREMLd additive and dominance heritability estimates, as a percent of the true ⁣ 324
broad sense heritability, over λ. Models with high degree of dominance $h = 0, 0.1$, show a significant ⁣ 325
heritability due to dominance effects (Fig S12). Neither the GBR nor the iMR models with $h \geq 0.25$ show ⁣ 326
significant dominance heritability. Filtering out rare alleles($MAF < 0.01$), as was done in [58], can make the ⁣ 327
estimates more or less accurate depending on the model and parameters. When λ is low, filtering out low ⁣ 328
MAF variants makes the GREMLd estimates more accurate. If λ is high the, unfiltered estimates tend to be ⁣ 329
better. The difference between filtered and unfiltered estimates is exacerbated by population growth. In the ⁣ 330
cMR model($h = 0$) filtering out rare alleles causes GREMLd to be much more accurate, because the ⁣ 331
unfiltered estimate has substantial upward bias. However, under the GBR and iMR models, filtering tends to ⁣ 332
create substantial downward bias. While filtering out rare alleles may be viewed as necessary given the noise ⁣ 333
present in many current genotype datasets, there may be a significant impact on the accuracy of the ⁣ 334
estimates. ⁣ 335

As an alternative to GREML estimates, Fig S13-Fig S16 show that HE-MS estimates of additive and ⁣ 336
dominance heritability. For intermediate values of λ, HE-MS regression is an accurate estimate of broad-sense ⁣ 337
heritability due to the analyzed genomic region in the population. When λ is small under the AC or GBR ⁣ 338
models there is a tendency to over-estimate total heritability. However, when λ is large there is a tendency to ⁣ 339
underestimate total heritability. Under the cMR model, the dominance component is massively overestimated ⁣ 340
and should be viewed as entirely unreliable. Although this issue is less pronounced under the iMR model, the ⁣ 341
dominance component tends to be negative and should also be viewed as unreliable. In general we find that ⁣ 342
intermediate iMR $h = 0.25$ and GBR models are not inconsistent with the findings of [58] and [57]. However, ⁣ 343
we caution against the reliability of these results. Our results suggest that estimating heritability due to ⁣ 344
regions as small as 100Kb should be done with careful quality control of data and results. ⁣ 345

The genetic model affects the outcomes of GWAS   346

Both demography and the model of gene action affect the degree to which rare variants contribute to the   347
genetic architecture of a trait (Figure 1). However, the different mappings of genotype to phenotype from   348
model to model make it difficult to predict *a priori* the outcomes of GWAS under each model. Further, in   349
order for models based on explicit population-genetic principles to inform the design and/or the   350
interpretation of GWAS studies, it is essential to understand if GWAS outcomes under different models are   351
consistent with current GWAS results [4, 14, 56]. Therefore, we sought to examine the performance of   352
statistical methods for GWAS under each genetic and demographic model. We assessed the power of a single   353
marker logistic regression to detect the gene region by calculating the proportion of model replicates in which   354
at least one variant reached genome wide significance at $\alpha \leq 10^{-8}$ (Fig 2A). The basic logistic regression is   355
equivalent to testing for association under the AC model and simulated both perfect "genotyping chip" (all   356
markers with $MAF \geq 0.05$) and complete re-sequencing including all markers (Fig 2B).   357

   Although rare alleles contribute relatively more to genetic variance under population growth (Fig 1),   358
there is less power to detect associations under all models when populations have experienced recent, rapid   359
growth (Fig 2A). The loss of power is attributable to a combination of rapid growth resulting in an excess of   360
rare variants overall [50, 52, 70, 75–85], and the increasing strength of selection against causal variants in   361
growing populations [48]. While complete resequencing is more powerful than a gene-chip design, the relative   362
power gained is modest under growth (Figure 2A). Region-based rare association tests behave similarly with   363
respect to population growth (Fig 2B).   364

   There are important differences in the behavior of the examined statistical methods across genetic models.   365
We focus first on the single marker tests (Fig 2A). For gene-chip strategies, power increases for "site-based"   366
models as recessivity of risk variants increases (compare power for AC, iMR, and cMR models in Fig 2B).   367
This increase in power is due to the well-known fact that recessive risk mutations are shielded from selection   368
when rare (due to being mostly present as heterozyogtes), thus reaching higher frequencies on average (Fig   369
S5), and that the single-marker test is most powerful when risk variants are common [90]. Further, for the   370
complete multiplicative-recessive model (cMR), the majority of V_G is due to common variants (Fig 1),   371
explaining why resequencing does not increase power for this model (Fig 2A).   372

   The GBR model predicts large gains in power under re-sequencing for intermediate λ (the mean   373
trait-effect size of newly arising causal mutations), similar to the AC or iMR model. But, when λ is larger   374
these gains are significantly diminished. The GBR model is effectively a model of complete recessivity when   375
λ is high because the average number of mutations per gamete (Fig S6) is much less than 0.5. This means   376
that an average diploid is likely to have at most one causal mutation. Under the GBR model one completely   377
wild-type gamete results in a wild-type phenotype. However, the GBR does not show a pattern of genetic   378
variance in the population consistent with either cMR or iMR models (1). While the behavior of the number   379
of mutations per gamete is consistent across models (Fig S6), only the GBR model demonstrates this effect   380
reflecting a qualitative difference between the GBR and other models. Surely, the overall shift toward rare   381
alleles at larger values of λ also contributes to the decrease in power, as we see such a decrease under the AC   382
model. However, if the drop in power were only as a result of frequency, then we would expect the decrease   383
to be largest in the AC model. Yet, the largest drop in power as a function of λ is seen in the GBR model.   384
We can conclude that there are significant non-linearities in the observed power of logistic regression as a   385

function of λ, model, and demography. ₃₈₆

Region-based rare variant association tests show many of the same patterns across genetic model and ₃₈₇ effect size distribution as single marker tests, but there are some interesting differences. The ESM test is the ₃₈₈ most powerful method tested for the AC, iMR, and GBR models (Fig 2B), with the c-Alpha test as a close ₃₈₉ second in some cases. For those models, the power of naive SKAT, linear kernel SKAT and SKAT-O, is ₃₉₀ always lower than the ESM and c-Alpha tests. This is peculiar since the c-Alpha test statistic is the same as ₃₉₁ the linear kernel SKAT test. The major difference between SKAT and ESM/c-Alpha is in the evaluation of ₃₉₂ statistical significance. SKAT uses an analytical approach to determining p-values while the ESM/c-Alpha ₃₉₃ tests use an explicit permutation approach. This implies that using permutation based p-values results in ₃₉₄ greater power. Yet, under the cMR model the linear kernel SKAT is the most powerful, followed by c-alpha. ₃₉₅ The cMR model does not predict a significant burden of rare alleles and so the default beta weights of SKAT ₃₉₆ are not appropriate, and the linear kernel is superior. The ESM test does poorly on this model because there ₃₉₇ are not many marginally significant low-frequency markers. It is logical to think that these tests would all ₃₉₈ perform better if all variants were included. The massive heterogeneity in the performance of region-based ₃₉₉ rare variant tests across models strongly suggests that multiple methods should be used when prior ₄₀₀ knowledge of underlying parameters is not available. In agreement with [24, 34], we predict that population ₄₀₁ growth reduces the power to associate variants in a causal gene region with disease status (Fig 2) when the ₄₀₂ disease also impacts evolutionary fitness. We also show that there is potential for more powerful burden tests, ₄₀₃ such as the ESM test, to be implemented in practice. ₄₀₄

The distribution of the allele frequencies of GWAS hits ₄₀₅

The distribution of the minor allele frequency of the most statistically significant single marker in a region is ₄₀₆ an interesting statistical signature of GWAS. This signature captures information regarding the underlying ₄₀₇ effect sizes and frequencies of typed variants, and statistical power. We investigate this by pooling model ₄₀₈ replicates across values of λ to reflect a uniform prior on this parameter (Fig 3). ₄₀₉

It was noted by [4, 56], that an excess of rare significant hits, relative to empirical data, is predicted by ₄₁₀ AC models where the effects of mutations contribute directly to fitness and the disease trait. We confirm ₄₁₁ that AC models are inconsistent with the empirical data (Fig 3). The empirical data represents (Fig 3) a ₄₁₂ pooled data set with the same diseases and quality filters as in [56], but updated to include more recent data. ₄₁₃ Close to half of the data comes from GWAS studies uploaded to the NHGRI database after 2011, yet the ₄₁₄ same qualitative pattern is observed. This contradicts the hypothesis that the initial observation was simply ₄₁₅ due to the relatively small sample sizes and low marker density in early GWAS. This observation is a robust ₄₁₆ and important one to consider when evaluating the parsimony of a theoretical model with empirical results. ₄₁₇

The GBR model agrees extremely well with empirical data, it predicts few rare significant hits and an ₄₁₈ approximately uniform distribution across the remainder of MAF domain (Fig 3). The more uniform ₄₁₉ distribution of significant single markers seen under the GBR is consistent with the flatter distribution of ₄₂₀ genetic variance (Fig 1). The cMR model shows an excess of intermediate frequency variants. If one ₄₂₁ considers trying to determine an approximate dominance coefficient in the GBR model, it would be found ₄₂₂ that there is a distribution of coefficients across sites. Yet, when simulating iMR model, we find that an ₄₂₃ intermediate degree of dominance, $h = 0.25 - 0.5$ results in distribution of significant hits which agrees well ₄₂₄

with empirical data as well (Fig 3 and Fig S9). In future studies, with more extensive simulation and careful pruning of empirical data, it might be possible to use the distribution of significant hits as part of a more rigorous model-based parameter estimation procedure.

In consideration of the rare allele of large effect hypothesis, [26] proposed a models where multiple rare alleles dominate disease risk and create synthetic associations with common SNPS. However, shortly after it was shown that this particular model was inconsistent with GWAS theoretically and empirically [4, 56, 91]. This inconsistency remained un-reconciled in the literature until now. We find that the MAF distribution of significant hits in a GWAS varies widely with choice of genetic model. In particular, we confirm the results of Wray et al., that AC evolutionary models predict an excess of low frequency significant hits. Also, the cMR model predicts an excess of intermediate and common significant hits. Utilizing a GBR model or an intermediate iMR model with $h = 0.25 - 0.5$, reconciles this inconsistency by simultaneously predicting the importance of rare alleles of large effect and the correct allele frequency distribution among statistically significant single markers.

Conclusion

The genetic model may be more important than the demographic model with respect to designing association studies. While changes in population size do affect the relationship between effect size and mutation frequency [50, 52, 70, 75–85] (Figs 1 and Fig S5), different mappings of genotype to trait value do this in radically different ways for the same demographic history (Fig 1). Further, the fact that models with strong recessive components are more compatible with GWAS results (Fig 3) implies that the development of methods with explicit consideration of gene-based dominance or recessivity might be an important next step. From an empirical perspective, our findings suggest that re-sequencing approaches in large samples is likely the best way forward in the face of the allelic heterogeneity imposed by the presence of rare alleles of large effect. Re-sequencing of candidate genes [92–95] and exomes [54, 55, 96–100] in case-control panels have observed an abundance of rare variants associated with case status. Here we show that under a model of mutation-selection balance at genes, neither current single-marker nor popular multi-marker tests are especially powerful at detecting large genomic regions harboring multiple risk variants (Fig 2). The extent to which the conclusions of this work can be extended to exome resequencing studies is unclear. Exons likely have a large fraction of sites mutable to, and larger effect size at, causative alleles than a large genomic region, and physical clustering of causal mutations may be more block-like in exome resequencing datasets given exon/intron structure. It will be of value in future work to more explicitly explore the properties of exome resequencing case-control studies under a gene-based mutation-selection balance model.

Here, the total heritability due to the simulated locus is only either 8% or 4% with the remaining phenotypic variation due to the un-simulated portions of the genome, and/or the environment. Passing the trait value through the Gaussian function to determine fitness has the effect of decreasing the strength of selection on variants which occur on predominantly unaffected genetic backgrounds, and only creates strong selection against variants which are on genetic backgrounds with multiple mutations. Thus, selection is inherently frequency-dependent [101, 102]. This model may also apply to traits with much higher heritability due to multiple unlinked loci with effects similar to that of our simulated region.

While a complete exploration of the impact of arbitrary dominance on the predictions of a model with

site-based recessivity remains to be done, we argue that it is unnecessary in this context. The use of multi-locus evolutionary models in simulating single nucleotide variants in a gene region is an incorrect application of the theory. Each locus in a multi-locus model is best suited to represent a gene, not a polymorphic site within a gene [103]. This distinction is critical as the alleles in these classic evolutionary models represent a class of haplotypes which have similar (idealized as identical) contributions to fitness. Like those models, our GBR model adheres to the classical definition of a gene in which mutant recessive alleles fail to complement [46]. It is therefore compelling to observe that a simple change in perspective, a reversion to a classical perspective, in the modeling of a trait appears to reconcile an inconsistency between theory and experiment (compare Fig 3 to the results of [4, 56]).

Materials and Methods

Forward simulation

Using the fwdpp template library v0.2.8 [104], we implemented a forward in time individual-based simulation of a Wright-Fisher population with mutation under the infinitely many sites model [66], recombination, and selection occurring each generation. We simulated populations of size $N = 2e4$ individuals for a time of $8N$ generations with a neutral mutation rate of $\mu = 0.00125$ per gamete per generation and a per diploid per generation recombination rate of $r = 0.00125$. Deleterious mutations occurred at a rate of $\mu_d = 0.1\mu$ per gamete per generation. These parameters correspond to $\theta = 4N\mu = \rho = 4Nr = 100$ and thus our simulation approximates a 100Kb region of the human genome. Population growth has been suggested to change the architecture of complex traits while keeping genetic load and genetic variance constant [34, 48]. Therefore, it was important for us to explore how the population growth impacted traits evolving under different genetic models. For simulations of population growth we extended the constant population simulations for 500 generations during which the population size grows exponentially from $N_i = 2e4$ to $N_{final} = 1e6$. This demographic model is much simpler than current models fit to empirical data [83]. However, this simple model allows us to more easily get a sense of the impact of population expansion [34, 48].

Models of disease and fitness

We implemented three disease-trait models with the phenotypic form $P = G + E$. G is the genetic component, and $E = N(0, \sigma_e)$ is the environmental noise expressed as a Gaussian random variable with mean 0 and standard deviation $\sigma_e = 0.075$. The additive co-dominant (AC) genetic model is $G = \sum_{i=1}^{n} \sum_{j=1}^{2} (c_{ij})$, where c_{ij} is the effect of the allele at the i-th site on the j-th haplotype, and is effectively the sum of the effect sizes of all mutations in a diploid. The classic population genetic multiplicative site-based recessive viability model is implemented as $G = \left(\prod_{i=1}^{n} \left(1 + \sum_{j=1}^{2} (h_{ij}c_{ij}) \right) \right) - 1$, where c_{ij} is as above and h_{ij} is the dominance coefficient. This model is referred to as the SBR model and is classified as either complete or incomplete. For example, when an individual has $k_i = \{0, 1, 2\}$ risk mutations at the i-th locus the complete site-based recessive (cSBR) model has $h_{ij} = 0, 0, 1$ and a incomplete site-based recessive (iSBR) model might be $h_{ij} = 0, 0.25, 1$. The gene-based partially recessive (GBR) model of [14] is implemented as

$$G = \sqrt{\sum_{i=1}^{n} c_{i,1} \times \sum_{i=1}^{n} c_{i,2}}.$$

In all cases, the phenotype's are subject to Gaussian stabilizing selection with an optimum of zero and standard deviation of $\sigma_s = 1$ such that the fitness, w, of a diploid is proportional to $e^{-\frac{P^2}{2\sigma_s^2}}$. The effect sizes of a causative mutations were sampled from an exponential distribution with means of $\lambda = 0.01, 0.025, 0.05, 0.1, 0.125, 0.25,$ or 0.5. 250 simulation trials were performed for each parameter/model combination. The parameters chosen here reflect a trait under weak "real" stabilizing selection [105].

Exploring the gene region's contribution to heritability

Broad-sense heritability can be calculated directly from our simulated data as $H^2 = \frac{Var_G}{Var_P}$. We explored broad-sense heritability as a function of mean causative effect size under each model. We used the house-of-cards (HOC) approximations for total genetic variance under stabilizing selection to predict broad-sense heritability [45,67]. Using these predictions we analyzed models where $H^2 \sim 0.04$ or 0.08. The total locus heritability under a co-dominant model is expected to follow the HOC approximation in certain parameter regimes. The HOC approximations to the expected genetic variance for an additive and recessive genetic model are $V_G \approx 4\mu_d V_s$ and $V_G \approx 2\mu V_s$ respectively [45,67]. In our simulations, $V_s = 1$, because $\sigma_s = 1$, and we varied $V_e = \sigma_e^2$ to adjust the total heritability.

Determining the genetic load of the population

Genetic load is defined as the relative deviation in a populations fitness from the fitness optimum, $L = (w_{max} - \bar{w})/(w_{max})$. We set the phenotypic optimum to be zero; $P_{opt} = 0$. When determining fitness for the SBR models, we subtract one from all phenotypes. This implies that $w_{max} = e^{-\frac{P_{opt}^2}{2\sigma_s^2}} = 1$ and that load is a simple function of the phenotypes of the population, $L = 1 - e^{-\frac{P^2}{2\sigma_s^2}}$. We also used the mean number of mutations per individual, and the mean frequency and effect sizes of segregating risk variants as proxies for the genetic load.

Additive and dominance genetic variance over allele frequency

We used an approach based sequential (type-1) regression sums of squares to estimate the contribution of the additive and dominance effects of variants to the total genetic variation due to a locus. Given a genotype matrix (rows are individuals and columns are risk variants) of $(0,1,$ or $2)$ copies of a risk allele (*e.g.* all mutations affecting phenotype), we sort the columns by decreasing risk mutation frequency. Then, within frequency classes, columns were sorted by decreasing effect sizes. For each variant a dominance component was also coded as 0, 2p, or 4p-2 according to the orthogonal model of [58], where p is the frequency of the variant in the population. We then used the R package `biglm` [106] to regress the individual genetic values (G in the previous section) onto this matrix. The variance explained by the additive and dominance effects of the m markers with $p \le x$ is then approximately $r^2 = (\sum_{i=1}^{m} SS_{reg,i})/(SS_{tot})$. Averaging results across replicates, this procedure results in a Monte-Carlo estimate of the fraction of V_G that is due to additive and dominance effects of variants with population frequency less than or equal to x is $(V_{A;p\le x} + V_{D;p\le x})/(V_{G;p\le 1})$ [48]. This fraction can be easily partitioned into strictly additive and dominance components.

Additive and dominance heritability in population samples $_{534}$

Using raw individual phenotypic values as quantitative trait values, random samples from simulated $_{535}$
populations (n=6000) were converted to .bed format using PLINK 1.90a [107]. PLINK was also used to test $_{536}$
for HWE ($p < 1e-6$) and filter on minor allele frequency. We tested the effect of filtering on MAF by $_{537}$
performing the analysis on unfiltered datasets and with markers with $MAF < 0.01$ removed. For $_{538}$
comparison, we calculated the true total heritability in the sample as $_{539}$
$H^2_{sample} = (VarG; sample)/(Var_{P;sample})$. Here we can not rely on the direct regression methods used in $_{540}$
the previous section on the whole populations. Instead, we used both GREMLd of [58], as implemented in $_{541}$
GCTA 1.24.4 [88], and Haseman-Elston (HE) regression [108, 109], to estimate additive and dominance $_{542}$
contributions to broad sense heritability in our simulated sample panels. GCTA was run with the Fisher $_{543}$
scoring, no variance component constraint and a max of 200 iterations. HE regressions were performed on $_{544}$
MAF stratified [110] genetic relationship matrices (GRMs). More specifically, we regress the off diagonal of $_{545}$
each GRM onto the cross-product of the centered and scaled sample phenotypes in a multiple linear $_{546}$
regression setting. The GRMs were constructed using GCTA with the flags –autosome and –make-grm(-d). $_{547}$
For MAF stratification, we use two bins $MAF \leq 0.01$ and $MAF > 0.01$ for each component. $_{548}$

Case-control studies $_{549}$

Following [14], we sampled 3000 cases and 3000 controls from each simulated population. Cases were $_{550}$
randomly sampled from the upper 15% of phenotypic values in the population, and controls were randomly $_{551}$
sampled from within 1 standard deviation of the population mean. This is the classic liability scale model $_{552}$
(see [71]). We define a "GWAS" to be a study including all markers with MAF $\geq 5\%$ and a re-sequencing $_{553}$
study to include all markers. In all cases we used a minor allele count logistic regression as the single marker $_{554}$
test. For single marker tests, the p-value cut off for significance is $p \leq 1e-08$ which is common in current $_{555}$
GWAS [26, 111]. Power is determined by the percentage of simulation replicates in which at least one marker $_{556}$
reaches genome wide significance. $_{557}$

Region-based tests of association due to rare alleles $_{558}$

We applied multiple region-based tests to our simulated data, ESM_K [14], several variations SKAT [28] and $_{559}$
c-Alpha [59]. We used the `R` package from the SKAT authors to implement their test $_{560}$
(`http://cran.r-project.org/web/packages/SKAT/index.html`). The remaining tests were implemented $_{561}$
in a custom `R` package (see SOFTWARE AVAILABILITY below). For the ESM_K and c-Alpha we performed up $_{562}$
to $2e6$ permutations of case-control labels to determine empirical p-values. All region-based tests were run on $_{563}$
low-frequency variants ($q < 0.05$). $_{564}$

Distribution of Significant GWAS Hits $_{565}$

Following [4, 56], we calculated distribution of the minor allele frequency (MAF) of the most significant SNPs $_{566}$
in a GWAS in empirical and simulated data. The empirical data was obtained from the NHGRI GWAS $_{567}$
database (`http://www.genome.gov/gwastudies`) on 02/05/2015. We considered the same diseases and $_{568}$
applied the same filters as in Table 3 of [56], $_{569}$

In order to mimic ascertained SNP data, we sampled markers from our case/control panels according to their minor allele frequencies [69], as done in [14] . The minor allele frequency of the most significant marker with a single-marker score $-log_{10}(p) \geq 8$, for all replicates where significant markers were present.

Software availability

Our simulation code and code for downstream analyses are freely available at

- `http://github.com/molpopgen/disease_sims`

- `http://github.com/molpopgen/buRden`

Acknowledgments

We are thankful toJ oseph Farran, Harry Mangalam, , Adam Brenner, Garr Updegraff, and Edward Xia for administering the University of California, Irvine HighPerformance Computing cluster. We are thankful to Kirk Lohmueller for helpful detailed comments throgought this project. We would like to thank Peter Andolfatto, Bogdan Pasaniuc and Nick Mancuso for helpful discussion. This work was supported by NIH grant R01-GM115564 to KRT, and NIH grant R01-GM115562 to ADL. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship Program under Grant No. DGE-1321846. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.

Figures

Fig 1. Variance explained over allele frequency. The cumulative additive and dominance genetic variance which can be explained by markers whose frequencies, q, are $\leq x$. Each color represents a different value of λ: the mean effects size of a new deleterious mutation. Shown here are the gene-based (GBR), additive co-dominant (AC), incomplete multiplicative recessive (Mult. recessive ($h = 0.25$); iMR) and complete multiplicative recessive (Mult. recessive ($h = 0$);cMR) models. Solid lines show the additive variance alone and dotted lines show the combined additive and dominance variance. All data shown are for models where $H^2 \sim 0.08$, although it was confirmed (data not shown) that these particular results are independent of total H^2. The additive and dominance genetic variance is estimated by the adjusted r^2 of the regression of all markers (and their corresponding dominance encoding) with $q \leq x$ onto total genotypic value (see methods for details); data are displayed as the mean of 250 simulation replicates. The vertical dotted and dashed lines correspond to the $q = 0.01$ and $q = 0.05$, respectively. The curves under no growth appear to be truncated with respect to rapid growth because the range of the x-axis differs between growth and no growth (minimum $q = 1/2N$).

Fig 2. Power of association tests. (A) The power of a single marker logistic regression, at significance threshold of $\alpha \leq 10^{-8}$, as a function of λ: the mean effect size of a new deleterious mutation. For single marker tests we define power as the number of simulation replicates in which any single marker reaches genome wide significance. Two study designs were emulated. For the gene chip design only markers with $MAF > 0.05$ were considered and all markers were considered for the resequencing design. Genetic models shown here are the additive co-dominant (AC), gene-based (GBR), complete multiplicative site-based recessive (Mult. recessive ($h = 0$); cMR) and incomplete multiplicative site-based recessive models (Mult. recessive ($h = 0.25$); iMR) (B) The power of region-based rare variant to detect association with the simulated causal gene region at significance threshold of $\alpha \leq 10^{-6}$. For region-based tests, we define power as the percent of simulation replicates in which the p-value of the test was less than α. The p-values for the ESM, c-Alpha were evaluated using 2×10^6 permutations. SKAT p-values were determined by the SKAT R package and represent numerical approximations to the presumed analytical p-value.

Fig 3. Distribution of significant hits. The distribution of minor allele frequency (MAF) of the most associated single marker in a GWAS. Each panel contains simulated data pooled across model replicates and λ (reflecting a uniform prior on the value of that parameter) overlayed onto empirical data. Empirical data are described in Materials and Methods. In cases where more than one marker was tied for the lowest p-value, one was chosen at random. Shown here are the additive co-dominant (AC), gene-based (GBR), incomplete multiplicative recessive (Mult. recessive ($h = 0.25$); iMR) and complete multiplicative recessive (Mult. recessive ($h = 0$);cMR) models. All data shown are for models where $H^2 \sim 0.08$, because single marker test power was too low under $H^2 \sim 0.04$ to make informative density plots. Simulated data were subjected to ascertainment sampling such that the MAF distribution of all markers on the simulated genotyping chip was uniform. The multiplicative recessive model is the cMR.

References

1. Welter D, MacArthur J, Morales J, Burdett T, Hall P, Junkins H, et al. The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. Nucleic Acids Research. 2014;42(December 2013):1001–1006.

2. Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, et al. Finding the missing heritability of complex diseases. Nature. 2009 oct;461(7265):747–53. Available from: `http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2831613{&}tool=pmcentrez{&}rendertype=abstract`.

3. Visscher PM, Brown MA, McCarthy MI, Yang J. Five years of GWAS discovery. American journal of human genetics. 2012 jan;90(1):7–24. Available from: `http://www.sciencedirect.com/science/article/pii/S0002929711005337`.

4. Gibson G. Rare and common variants: twenty arguments. Nature Reviews Genetics. 2012;13(2):135–145. Available from: `http://dx.doi.org/10.1038/nrg3118`.

5. Robinson MR, Wray NR, Visscher PM. Explaining additional genetic variation in complex traits. Trends in Genetics. 2014;30:124–132. Available from: `http://dx.doi.org/10.1016/j.tig.2014.02.003`.

6. Eichler EE, Flint J, Gibson G, Kong A, Leal SM, Moore JH, et al. Missing heritability and strategies for finding the underlying causes of complex disease. Nature reviews Genetics. 2010 jun;11(6):446–50. Available from: `http://dx.doi.org/10.1038/nrg2809`.

7. Wei WH, Hemani G, Haley CS. Detecting epistasis in human complex traits. Nature Reviews Genetics. 2014;15(11):722–733. Available from: `http://dx.doi.org/10.1038/nrg3747`.

8. Zuk O, Hechter E, Sunyaev SR, Lander ES. The mystery of missing heritability: Genetic interactions create phantom heritability. Proceedings of the National Academy of Sciences. 2012;109(4):1193–1198.

9. Fisher RA. The Genetical Theory Of Natural Selection : Fisher, R. A : Free Download & Streaming : Internet Archive. Oxford, UK; 1930. Available from: `https://archive.org/details/geneticaltheoryo031631mbp`.

10. Visscher PM, Hill WG, Wray NR. Heritability in the genomics era–concepts and misconceptions. Nature reviews Genetics. 2008 apr;9(4):255–66. Available from: `http://dx.doi.org/10.1038/nrg2322`.

11. McClellan J, King MC. Genetic heterogeneity in human disease. Cell. 2010 apr;141(2):210–7. Available from: `http://www.sciencedirect.com/science/article/pii/S009286741000320X`.

12. Cirulli ET, Goldstein DB. Uncovering the roles of rare variants in common disease through whole-genome sequencing. Nature reviews Genetics. 2010 jun;11(6):415–25. Available from: `http://dx.doi.org/10.1038/nrg2779`.

13. Visscher PM, Goddard ME, Derks EM, Wray NR. Evidence-based psychiatric genetics, AKA the false dichotomy between common and rare variant hypotheses. Molecular psychiatry. 2012 may;17(5):474–85. Available from: `http://dx.doi.org/10.1038/mp.2011.65`.

14. Thornton KR, Foran AJ, Long AD. Properties and Modeling of GWAS when Complex Disease Risk Is Due to Non-Complementing, Deleterious Mutations in Genes of Large Effect. PLoS Genetics. 2013;9(2).

15. Moutsianas L, Agarwala V, Fuchsberger C, Flannick J, Rivas MA, Gaulton KJ, et al. The power of gene-based rare variant methods to detect disease-associated variation and test hypotheses about complex disease. PLoS genetics. 2015 apr;11(4):e1005165. Available from: `http://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1005165`.

16. Wang T, Elston RC. Improved power by use of a weighted score test for linkage disequilibrium mapping. American journal of human genetics. 2007 feb;80(2):353–60. Available from: `http://www.sciencedirect.com/science/article/pii/S0002929707626937`.

17. Li B, Leal S. Methods for detecting associations with rare variants for common diseases: application to analysis of sequence data. The American Journal of Human Genetics. 2008;83:311–321. Available from: `http://www.sciencedirect.com/science/article/pii/S0002929708004084`.

18. Madsen BE, Browning SR. A groupwise association test for rare mutations using a weighted sum statistic. PLoS Genetics. 2009;5(2).

19. Basu S, Pan W. Comparison of statistical tests for disease association with rare variants. Genetic epidemiology. 2011 nov;35(7):606–19. Available from: `http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3197766{&}tool=pmcentrez{&}rendertype=abstract`.

20. Urbanowicz RJ, Kiralis J, Sinnott-Armstrong NA, Heberling T, Fisher JM, Moore JH. GAMETES: a fast, direct algorithm for generating pure, strict, epistatic models with random architectures. BioData mining. 2012 jan;5(1):16. Available from: `http://www.biodatamining.org/content/5/1/16`.

21. Xu Y, Wu Y, Song C, Zhang H. Simulating realistic genomic data with rare variants. Genetic epidemiology. 2013 feb;37(2):163–72. Available from: `http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3543480{&}tool=pmcentrez{&}rendertype=abstract`.

22. Pan W, Kim J, Zhang Y, Shen X, Wei P. A powerful and adaptive association test for rare variants. Genetics. 2014 aug;197(4):1081–95. Available from: `http://www.genetics.org/content/197/4/1081.long{#}ref-43`.

23. Su Z, Marchini J, Donnelly P. HAPGEN2: simulation of multiple disease SNPs. Bioinformatics (Oxford, England). 2011 aug;27(16):2304–5. Available from: `http://bioinformatics.oxfordjournals.org/content/27/16/2304.long`.

24. Uricchio LH, Torres R, Witte JS, Hernandez RD. Population genetic simulations of complex phenotypes with implications for rare variant association tests. Genetic epidemiology. 2015 jan;39(1):35–44. Available from: `http://www.ncbi.nlm.nih.gov/pubmed/25417809`.

25. Hudson RR. Generating samples under a Wright-Fisher neutral model of genetic variation. Bioinformatics (Oxford, England). 2002;18(2):337–338.

26. Dickson SP, Wang K, Krantz I, Hakonarson H, Goldstein DB. Rare Variants Create Synthetic Genome-Wide Associations. PLoS Biology. 2010;8(1).

27. Luo L, Boerwinkle E, Xiong M. Association studies for next-generation sequencing. Genome research. 2011 jul;21(7):1099–108. Available from: `http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3129252{&}tool=pmcentrez{&}rendertype=abstract`.

28. Wu MC, Lee S, Cai T, Li Y, Boehnke M, Lin X. Rare-variant association testing for sequencing data with the sequence kernel association test. American Journal of Human Genetics. 2011;89(1):82–93. Available from: `http://dx.doi.org/10.1016/j.ajhg.2011.05.029`.

29. Schraiber JG, Landis MJ. Sensitivity of quantitative traits to mutational effects and number of loci. Theoretical population biology. 2015 jun;102:85–93. Available from: `http://www.ncbi.nlm.nih.gov/pubmed/25840144`.

30. Price AL, Kryukov GV, de Bakker PIW, Purcell SM, Staples J, Wei LJ, et al. Pooled association tests for rare variants in exon-resequencing studies. American journal of human genetics. 2010 jun;86(6):832–8. Available from: `http://www.cell.com/article/S0002929710002077/fulltext`.

31. King CR, Rathouz PJ, Nicolae DL. An evolutionary framework for association testing in resequencing studies. PLoS genetics. 2010 nov;6(11):e1001202. Available from: `http://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1001202`.

32. Browning SR, Thompson EA. Detecting rare variant associations by identity-by-descent mapping in case-control studies. Genetics. 2012 apr;190(4):1521–31. Available from: `http://www.genetics.org/content/early/2012/01/18/genetics.111.136937`.

33. Agarwala V, Flannick J, Sunyaev S, Altshuler D. Evaluating empirical bounds on complex disease genetic architecture. Nature genetics. 2013;45(12):1418–27. Available from: `http://www.ncbi.nlm.nih.gov/pubmed/24141362`.

34. Lohmueller KE. The impact of population demography and selection on the genetic architecture of complex traits. PLoS genetics. 2014 may;10(5):e1004379. Available from: `http://dx.plos.org/10.1371/journal.pgen.1004379`.

35. North TL, Beaumont MA. Complex trait architecture: the pleiotropic model revisited. Scientific reports. 2015 jan;5:9351. Available from: `http://www.nature.com/srep/2015/150320/srep09351/full/srep09351.html`.

36. Caballero A, Tenesa A, Keightley PD. The Nature of Genetic Variation for Complex Traits Revealed by GWAS and Regional Heritability Mapping Analyses. Genetics. 2015 oct;201(4):1601–13. Available from: `http://genetics.org/content/201/4/1601.abstract`.

37. Eyre-Walker A. Evolution in health and medicine Sackler colloquium: Genetic architecture of a complex trait and its implications for fitness and genome-wide association studies. Proceedings of the National Academy of Sciences of the United States of America. 2010;107 Suppl:1752–1756. ⁠692 ⁠693 ⁠694

38. Keightley PD, Hill WG. Variation Maintained in Quantitative Traits with Mutation-Selection Balance: Pleiotropic Side-Effects on Fitness Traits. Proceedings of the Royal Society B: Biological Sciences. 1990 nov;242(1304):95–100. Available from: http://rspb.royalsocietypublishing.org/content/242/1304/95. ⁠695 ⁠696 ⁠697 ⁠698

39. Mancuso N, Rohland N, Rand KA, Tandon A, Allen A, Quinque D, et al. The contribution of rare variation to prostate cancer heritability. Nature genetics. 2015 nov;advance on. Available from: http://dx.doi.org/10.1038/ng.3446. ⁠699 ⁠700 ⁠701

40. Haldane JBS. The measurement of natural selection. Proc 9th Int Congr Genet. 1954;1:480–487. ⁠702

41. Latter BDH. Natural selection for an intermediate optimum. Australian Journal of Biological Sciences. 1960;13:30–35. Available from: http://www.cabdirect.org/abstracts/19600101674.html;jsessionid=3B1562EB69031455E4C7BB05F3D8B699?freeview=true. ⁠703 ⁠704 ⁠705

42. Crow JF, Kimura M. The theory of genetic loads. Proc XI Int Congr Genet. 1964;p. 495–505. ⁠706

43. Kingman JFC. A Simple Model for the Balance between Selection and Mutation. Journal of Applied Probability. 1978 mar;15(1):1–12. Available from: http://www.jstor.org/stable/3213231. ⁠707 ⁠708

44. Fleming WH. Equilibrium Distributions of Continuous Polygenic Traits. SIAM Journal on Applied Mathematics. 1979 jul;Available from: http://epubs.siam.org/doi/abs/10.1137/0136014. ⁠709 ⁠710

45. Turelli M. Heritable genetic variation via mutation-selection balance: Lerch's zeta meets the abdominal bristle. Theoretical Population Biology. 1984 apr;25(2):138–193. Available from: http://www.sciencedirect.com/science/article/pii/0040580984900170. ⁠711 ⁠712 ⁠713

46. Benzer S. FINE STRUCTURE OF A GENETIC REGION IN BACTERIOPHAGE. Proceedings of the National Academy of Sciences of the United States of America. 1955 jun;41(6):344–54. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=528093{&}tool=pmcentrez{&}rendertype=abstract. ⁠714 ⁠715 ⁠716 ⁠717

47. Zuk O, Schaffner SF, Samocha K, Do R, Hechter E, Kathiresan S, et al. Searching for missing heritability: designing rare variant association studies. Proceedings of the National Academy of Sciences of the United States of America. 2014;111:E455–64. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3910587{&}tool=pmcentrez{&}rendertype=abstract. ⁠718 ⁠719 ⁠720 ⁠721 ⁠722

48. Simons YB, Turchin MC, Pritchard JK, Sella G. The deleterious mutation load is insensitive to recent population history. Nature genetics. 2014;46(August 2013):220–4. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24509481. ⁠723 ⁠724 ⁠725

49. Kryukov GV, Shpunt A, Stamatoyannopoulos JA, Sunyaev SR. Power of deep, all-exon resequencing   726
    for discovery of human trait genes. Proceedings of the National Academy of Sciences of the United   727
    States of America. 2009 mar;106(10):3871–6. Available from: 728
    `http://www.pnas.org/content/106/10/3871.abstract`.   729

50. Coventry A, Bull-Otterson LM, Liu X, Clark AG, Maxwell TJ, Crosby J, et al. Deep resequencing   730
    reveals excess rare recent variants consistent with explosive population growth. Nature   731
    communications. 2010 jan;1:131. Available from: `http://dx.doi.org/10.1038/ncomms1130`.   732

51. Fu W, O'Connor TD, Jun G, Kang HM, Abecasis G, Leal SM, et al. Analysis of 6,515 exomes reveals   733
    the recent origin of most human protein-coding variants. Nature. 2013 jan;493(7431):216–20.   734
    Available from: `http://dx.doi.org/10.1038/nature11690`.   735

52. Keinan a, Clark aG. Recent Explosive Human Population Growth Has Resulted in an Excess of Rare   736
    Genetic Variants. Science. 2012;336(6082):740–743.   737

53. Marth GT, Yu F, Indap AR, Garimella K, Gravel S, Leong WF, et al. The functional spectrum of   738
    low-frequency coding variation. Genome biology. 2011 jan;12(9):R84. Available from: 739
    `http://genomebiology.com/2011/12/9/R84`.   740

54. Nelson MR, Wegmann D, Ehm MG, Kessner D, St Jean P, Verzilli C, et al. An abundance of rare   741
    functional variants in 202 drug target genes sequenced in 14,002 people. Science (New York, NY).   742
    2012 jul;337(6090):100–4. Available from: 743
    `http://www.sciencemag.org/content/337/6090/100.abstract`.   744

55. Tennessen Ja, Bigham aW, O'Connor TD, Fu W, Kenny EE, Gravel S, et al. Evolution and   745
    Functional Impact of Rare Coding Variation from Deep Sequencing of Human Exomes. Science.   746
    2012;337(6090):64–69.   747

56. Wray NR, Purcell SM, Visscher PM. Synthetic associations created by rare variants do not explain   748
    most GWAS results. PLoS Biology. 2011;9(1).   749

57. Yang J, Bakshi A, Zhu Z, Hemani G, Vinkhuyzen AAE, Lee SH, et al. Genetic variance estimation   750
    with imputed variants finds negligible missing heritability for human height and body mass index.   751
    Nature Genetics. 2015 aug;advance on. Available from: `http://dx.doi.org/10.1038/ng.3390`.   752

58. Zhu Z, Bakshi A, Vinkhuyzen AE, Hemani G, Lee S, Nolte I, et al. Dominance Genetic Variation   753
    Contributes Little to the Missing Heritability for Human Complex Traits. The American Journal of   754
    Human Genetics. 2015;p. 377–385. Available from: 755
    `http://linkinghub.elsevier.com/retrieve/pii/S0002929715000099`.   756

59. Neale BM, Rivas Ma, Voight BF, Altshuler D, Devlin B, Orho-Melander M, et al. Testing for an   757
    unusual distribution of rare variants. PLoS Genetics. 2011;7(3).   758

60. Risch N. Linkage strategies for genetically complex traits. III. The effect of marker polymorphism on   759
    analysis of affected relative pairs. American journal of human genetics. 1990;46:242–253.   760

61. Risch N. Linkage strategies for genetically complex traits. I. Multilocus models. American journal of human genetics. 1990;46:222–228.

62. Haldane JBS. The cost of natural selection; 1957.

63. Charlesworth B, Morgan MT, Charlesworth D. The effect of deleterious mutations on neutral molecular variation. Genetics. 1993 aug;134(4):1289–1303. Available from: http://www.genetics.org/content/134/4/1289.abstract.

64. Risch N, Merikangas K. The future of genetic studies of complex human diseases. Science (New York, NY). 1996 sep;273(5281):1516–7. Available from: http://www.ncbi.nlm.nih.gov/pubmed/8801636.

65. Pritchard JK. Are rare variants responsible for susceptibility to complex diseases? American journal of human genetics. 2001 jul;69(1):124–37. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1226027{&}tool=pmcentrez{&}rendertype=abstract.

66. Kimura M. The number of heterozygous nucleotide sites maintained in a finite population due to steady flux of mutations. Genetics. 1969;61(694):893–903.

67. Slatkin M. Heritable variation and heterozygosity under a balance between mutations and stabilizing selection. Genetical research. 1987;50:53–62.

68. Burger R, Hofbauer J. Mutation load and mutation-selection-balance in quantitative genetic traits. Journal of Mathematical Biology. 1994 feb;32(3):193–218. Available from: http://link.springer.com/10.1007/BF00163878.

69. Marth GT, Czabarka E, Murvai J, Sherry ST. The Allele Frequency Spectrum in Genome-Wide Human Variation Data Reveals Signals of Differential Demographic History in Three Large World Populations. Genetics. 2004;166(1):351–372.

70. Gao F, Keinan A. High burden of private mutations due to explosive human population growth and purifying selection. BMC genomics. 2014;15 Suppl 4(Suppl 4):S3. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25056720.

71. Falconer D, Mackay TFC. Introduction to Quantitative Genetics, Ed. 2. Harlow, Essex, UK: Longmans Green; 1996.

72. Sham PC, Purcell SM. Statistical power and significance testing in large-scale genetic studies. Nature reviews Genetics. 2014 may;15(5):335–46. Available from: http://dx.doi.org/10.1038/nrg3706.

73. Uricchio LH, Witte JS, Hernandez RD. Selection and explosive growth may hamper the performance of rare variant association tests; 2015. Available from: http://www.biorxiv.org/content/early/2015/03/01/015917.abstract.

74. Hill WG, Goddard ME, Visscher PM. Data and theory point to mainly additive genetic variance for complex traits. PLoS genetics. 2008 feb;4(2):e1000008. Available from: http://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1000008.

75. Griffiths RC, Tavaré S. Sampling theory for neutral alleles in a varying environment. Philosophical transactions of the Royal Society of London Series B, Biological sciences. 1994 jun;344(1310):403–10. Available from: http://www.ncbi.nlm.nih.gov/pubmed/7800710.

76. Griffiths RC, Tavare S. Ancestral Inference in Population Genetics. Statistical Science. 1994 aug;9(3):307–319. Available from: http://projecteuclid.org/euclid.ss/1177010378.

77. Rogers A, Harpending H. Population growth makes waves in the distribution of pairwise genetic differences. Mol Biol Evol. 1992 may;9(3):552–569. Available from: http://mbe.oxfordjournals.org/content/9/3/552.

78. Fu YX. Statistical Tests of Neutrality of Mutations Against Population Growth, Hitchhiking and Background Selection. Genetics. 1997 oct;147(2):915–925. Available from: http://www.genetics.org/content/147/2/915.short.

79. Beaumont MA. Detecting Population Expansion and Decline Using Microsatellites. Genetics. 1999 dec;153(4):2013–2029. Available from: http://www.genetics.org/content/153/4/2013.short.

80. Aris-Brosou S, Excoffier L. The impact of population expansion and mutation rate heterogeneity on DNA sequence polymorphism. Molecular Biology and Evolution. 1996 mar;13(3):494–504. Available from: http://mbe.oxfordjournals.org/content/13/3/494.short.

81. Gazave E, Chang D, Clark AG, Keinan A. Population growth inflates the per-individual number of deleterious mutations and reduces their mean effect. Genetics. 2013;195(3):969–978.

82. Gazave E, Ma L, Chang D, Coventry A, Gao F, Muzny D, et al. Neutral genomic regions refine models of recent rapid human population growth. Proceedings of the National Academy of Sciences of the United States of America. 2014;111(2):757–62. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3896169{&}tool=pmcentrez{&}rendertype=abstract.

83. Gravel S, Henn BM, Gutenkunst RN, Indap AR, Marth GT, Clark AG, et al. Demographic history and rare allele sharing among human populations. Proceedings of the National Academy of Sciences. 2011 jul;108(29):11983–11988. Available from: http://www.pnas.org/content/108/29/11983.short.

84. Kimmel M, Chakraborty R, King JP, Bamshad M, Watkins WS, Jorde LB. Signatures of Population Expansion in Microsatellite Repeat Data. Genetics. 1998 apr;148(4):1921–1930. Available from: http://www.genetics.org/content/148/4/1921.short.

85. Reich DE, Goldstein DB. Genetic evidence for a Paleolithic human population expansion in Africa. Proceedings of the National Academy of Sciences. 1998 jul;95(14):8119–8123. Available from: http://www.pnas.org/content/95/14/8119.full.

86. Maher MC, Uricchio LH, Torgerson DG, Hernandez RD. Population genetics of rare variants and complex diseases. Human heredity. 2012 jan;74(3-4):118–28. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3698246{&}tool=pmcentrez{&}rendertype=abstract.

87. Reich DE, Lander ES. On the allelic spectrum of human disease. Trends in Genetics. 2001   831
sep;17(9):502–510. Available from: 832
`http://www.sciencedirect.com/science/article/pii/S0168952501024106`.   833

88. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis.   834
American journal of human genetics. 2011 jan;88(1):76–82. Available from: 835
`http://www.sciencedirect.com/science/article/pii/S0002929710005987`.   836

89. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria; 2014.   837
Available from: `http://www.R-project.org/`.   838

90. Spencer CCA, Su Z, Donnelly P, Marchini J. Designing Genome-Wide Association Studies: Sample   839
Size, Power, Imputation, and the Choice of Genotyping Chip. PLoS Genetics. 2009   840
may;5(5):e1000477. Available from: 841
`http://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1000477`.   842

91. Orozco G, Barrett JC, Zeggini E. Synthetic associations in the context of genome-wide association   843
scan signals. Human molecular genetics. 2010 oct;19(R2):R137–44. Available from: 844
`http://hmg.oxfordjournals.org/content/19/R2/R137.long`.   845

92. Johansen CT, Wang J, Lanktree MB, Cao H, McIntyre AD, Ban MR, et al. Excess of rare variants in   846
genes identified by genome-wide association study of hypertriglyceridemia. Nature genetics. 2010   847
aug;42(8):684–7. Available from: `http://www.pubmedcentral.nih.gov/articlerender.fcgi?`   848
`artid=3017369{&}tool=pmcentrez{&}rendertype=abstract`.   849

93. Kotowski IK, Pertsemlidis A, Luke A, Cooper RS, Vega GL, Cohen JC, et al. A spectrum of PCSK9   850
alleles contributes to plasma levels of low-density lipoprotein cholesterol. American journal of human   851
genetics. 2006;78(3):410–422.   852

94. Marini NJ, Gin J, Ziegle J, Keho KH, Ginzinger D, Gilbert Da, et al. The prevalence of   853
folate-remedial MTHFR enzyme variants in humans. Proceedings of the National Academy of   854
Sciences of the United States of America. 2008;105(23):8055–8060.   855

95. Romeo S, Pennacchio LA, Fu Y, Boerwinkle E, Tybjaerg-Hansen A, Hobbs HH, et al.   856
Population-based resequencing of ANGPTL4 uncovers variations that reduce triglycerides and   857
increase HDL. Nature genetics. 2007 apr;39(4):513–6. Available from: `http://www.pubmedcentral.`   858
`nih.gov/articlerender.fcgi?artid=2762948{&}tool=pmcentrez{&}rendertype=abstract`.   859

96. Huyghe JR, Jackson AU, Fogarty MP, Buchkovich ML, Stančáková A, Stringham HM, et al. Exome   860
array analysis identifies new loci and low-frequency variants influencing insulin processing and   861
secretion. Nature genetics. 2013;45(2):197–201. Available from: 862
`http://dx.doi.org/10.1038/ng.2507`.   863

97. Wessel J, Goodarzi MO. Low-frequency and rare exome chip variants associate with fasting glucose   864
and type 2 diabetes susceptibility. Nature Communications. 2015;6:5897. Available from: 865
`http://www.nature.com/doifinder/10.1038/ncomms6897`.   866

98. Purcell SM, Moran JL, Fromer M, Ruderfer D, Solovieff N, Roussos P, et al. A polygenic burden of rare disruptive mutations in schizophrenia. Nature. 2014;506(7487):185–90. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24463508.

99. Luciano M, Svinti V, Campbell A, Marioni RE, Hayward C, Wright AF, et al. Exome Sequencing to Detect Rare Variants Associated With General Cognitive Ability: A Pilot Study. Twin research and human genetics : the official journal of the International Society for Twin Studies. 2015 mar;p. 1–9. Available from: http://journals.cambridge.org/abstract{_}S1832427415000109.

100. Cirulli ET, Lasseigne BN, Petrovski S, Sapp PC, Dion PA, Leblond CS, et al. Exome sequencing in amyotrophic lateral sclerosis identifies risk genes and pathways. Science. 2015 feb;347(6229):1436–1441. Available from: http://www.sciencemag.org/content/347/6229/1436.full.

101. Zhang XS, Hill WG. Joint Effects of Pleiotropic Selection and Stabilizing Selection on the Maintenance of Quantitative Genetic Variation at Mutation-Selection Balance. Genetics. 2002 sep;162(1):459–471. Available from: http://www.genetics.org/content/162/1/459.long.

102. Zhang XS. Influence of Dominance, Leptokurtosis and Pleiotropy of Deleterious Mutations on Quantitative Genetic Variation at Mutation-Selection Balance. Genetics. 2004 jan;166(1):597–610. Available from: http://www.genetics.org/content/166/1/597.long.

103. Karlin S, Liberman U. Central equilibria in multilocus systems. I. Generalized nonepistatic selection regimes. Genetics. 1979;91:777–798.

104. Thornton KR. A C++ Template Library for Efficient Forward-Time Population Genetic Simulation of Large Populations. Genetics. 2014;(2013):1–21. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24950894.

105. Johnson T, Barton N. Theoretical models of selection and mutation on quantitative traits. Philosophical transactions of the Royal Society of London Series B, Biological sciences. 2005 jul;360(1459):1411–25. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1569515{&}tool=pmcentrez{&}rendertype=abstract.

106. Lumley T. biglm: bounded memory linear and generalized linear models; 2013. R package version 0.9-1. Available from: http://CRAN.R-project.org/package=biglm.

107. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. American journal of human genetics. 2007 sep;81(3):559–75. Available from: http://www.sciencedirect.com/science/article/pii/S0002929707613524.

108. Haseman JK, Elston RC. The investigation of linkage between a quantitative trait and a marker locus. Behavior Genetics. 1972 mar;2(1):3–19. Available from: http://link.springer.com/10.1007/BF01066731.

109. Sham PC, Purcell S. Equivalence between Haseman-Elston and variance-components linkage analyses for sib pairs. American journal of human genetics. 2001 jun;68(6):1527–32. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1226141{&}tool=pmcentrez{&}rendertype=abstract.

110. Lee SH, Yang J, Chen GB, Ripke S, Stahl EA, Hultman CM, et al. Estimation of SNP heritability from dense genotype data. American journal of human genetics. 2013 dec;93(6):1151–5. Available from: http://www.sciencedirect.com/science/article/pii/S0002929713004722.

111. McCarthy MI, Hirschhorn JN. Genome-wide association studies: Potential next steps on a genetic journey. Human Molecular Genetics. 2008;17(2):156–165.