A model of compound heterozygous, loss-of-function alleles is broadly consistent with observations from complex-disease GWAS datasets

Jaleal S. Sanjak[1,2] *, Anthony D. Long[1,2], Kevin R. Thornton[1,2] *,

1 Department of Ecology and Evolutionary Biology, University of California, Irvine, California 92697, USA
2 Center for Complex Biological Systems, University of California Irvine, Irvine, CA 92697, USA

* E-mail: jsanjak@uci.edu (JSS) , krthornt@uci.edu (KRT)

Abstract

The genetic component of complex disease risk in humans remains largely unexplained. A corollary is that the allelic spectrum of genetic variants contributing to complex disease risk is unknown. Theoretical models that relate population genetic processes to the maintenance of genetic variation for quantitative traits may suggest profitable avenues for future experimental design. Here we use forward simulation to model a genomic region evolving under a balance between recurrent deleterious mutation and Gaussian stabilizing selection. We consider multiple genetic and demographic models, and several different methods for identifying genomic regions harboring variants associated with complex disease risk. We demonstrate that the model of gene action, relating genotype to phenotype, has a qualitative effect on several relevant aspects of the population genetic architecture of a complex trait. In particular, the genetic model impacts genetic variance component partitioning across the allele frequency spectrum and the power of statistical tests. Models with partial recessivity closely match the minor allele frequency distribution of significant hits from empirical genome-wide association studies without requiring homozygous effect-sizes to be small. We highlight a particular gene-based model of incomplete recessivity that is appealing from first principles. Under that model, deleterious mutations in a genomic region partially fail to complement one another. This model of gene-based recessivity predicts the empirically observed inconsistency between twin and SNP based estimated of dominance heritability. Furthermore, this model predicts considerable levels of unexplained variance associated with intralocus epistasis. Our results suggest a need for improved statistical tools for region based genetic association and heritability estimation.

Author Summary

Gene action determines how mutations affect phenotype. When placed in an evolutionary context, the details of the genotype-to-phenotype model can impact the maintenance of genetic variation for complex traits. Likewise, non-equilibrium demographic history may affect patterns of genetic variation. Here, we explore the impact of genetic model and population growth on distribution of genetic variance across the allele frequency

spectrum underlying risk for a complex disease. Using forward-in-time population genetic simulations, we show that the genetic model has important impacts on the composition of variation for complex disease risk in a population. We explicitly simulate genome-wide association studies (GWAS) and perform heritability estimation on population samples. A particular model of gene-based partial recessivity, based on allelic non-complementation, aligns well with empirical results. This model is congruent with the dominance variance estimates from both SNPs and twins, and the minor allele frequency distribution of GWAS hits.

Introduction

Risk for complex diseases in humans, such as diabetes and hypertension, is highly heritable yet the causal DNA sequence variants responsible for that risk remain largely unknown. Genome-wide association studies (GWAS) have found many genetic markers associated with disease risk [1]. However, follow-up studies have shown that these markers explain only a small portion of the total heritability for most traits [2,3].

There are many hypotheses which attempt to explain the 'missing heritability' problem [2–5]. Genetic variance due to epistatic or gene-by-environment interactions is difficult to identify statistically because of, among other reasons, increased multiple hypothesis testing burden [6,7], and could artificially inflate estimates of broad-sense heritability [8]. Well-tagged intermediate frequency variants may not reach genome-wide significance in an association study if they have smaller effect sizes [9,10]. One appealing verbal hypothesis for this 'missing heritability' is that there are rare causal alleles of large effect that are difficult to detect [4,11,12]. These hypotheses are not mutually exclusive, and it is probable that a combination of models will be needed to explain all heritable disease risk [13].

The standard GWAS attempts to identify genetic polymorphisms that differ in frequency between cases and controls. A complementary approach is to estimate the heritability explained by genotyped (and imputed) markers (SNPs) under different population sampling schemes [14,15]. Stratifying markers by minor allele frequency (MAF) prior to performing SNP-based heritability estimation allows the partitioning of genetic variation across the allele frequency spectrum to be estimated [16], which is an important summary of the genetic architecture of a complex trait [16–23]. This approach has inferred a contribution of rare alleles to genetic variance in both human height and body mass index (BMI) [16], consistent with theoretical work showing that rare alleles will have large effect sizes if fitness effects and trait effects are correlated [18,20–25]. Yet, simulations of causal loci harboring multiple rare variants with large additive effects predict an excess of low-frequency significant markers relative to empirical findings [4,26].

SNP-based heritability estimates have concluded that there is little missing heritability for height and BMI, and that the causal loci simply have effect sizes that are too small to reach genome-wide significance under current GWAS sample sizes [14,16]. Further, extensions to these methods decompose genetic variance into additive and dominance components and find that dominance variance is approximately one fifth of the additive genetic variance on average across seventy-nine complex traits [27]. When taken into account together with results from GWAS, these observations can be interpreted as evidence that the genetic architecture of human traits is best-explained by a model of small additive effects. However, a recent large twin study found a substantial contribution of dominance variance for fourteen out of eighteen traits [28]. The reason for this discrepancy in results remains unclear. One possibility is a statistical artifact; for

example, twin studies may be prone to mistakenly infer non-additive affects when none exist. Another   33
possibility, which we return to later, is that this apparently contradictory results are expected under a   34
different model of gene action.   35

The design, analysis, and interpretation of GWAS are heavily influenced by the "standard model" of   36
quantitative genetics [29]. This model assigns an effect size to a mutant allele, but formally makes no   37
concrete statement regarding the molecular nature of the allele. Early applications of this model to the   38
problem of human complex traits include Risch's work on the power to detect causal mutations [30, 31] and   39
Pritchard's work showing that rare alleles under purifying selection may contribute to heritable variation in   40
complex traits [17]. When applied to molecular data, such as SNP genotypes in a GWAS, these models treat   41
the SNPs themselves as the loci of interest. For example, influential power studies informing the design of   42
GWAS assign effect sizes directly to SNPs and assume Risch's model of multiplicative epistasis [32].   43
Similarly, the single-marker logistic regression used as the primary analysis of GWAS data typically assumes   44
an additive or recessive model at the level of individual SNPs [33]. Finally, recent methods designed to   45
estimate the heritability of a trait explained by genotyped markers assigns additive and dominance effects   46
directly to SNPs [14, 16, 27, 34]. Naturally, the results of such analyses are interpreted in light of assumed   47
model of gene action.   48

A weakness of the multiplicative epistasis model [30, 31] when applied to SNPs is that the concept of a   49
gene, defined as a physical region where loss-of-function mutations have the same phenotype [35], is lost.   50
Specifically, under the standard model, the genetic concept of a failure to complement is a property of SNPs   51
and not "gene regions" (see [36] for a detailed discussion of this issue). We have recently introduced an   52
alternative model of gene action, one in which risk mutations are unconditionally deleterious and fail to   53
complement at the level of a "gene region" [36]. This model, influenced by the standard operational   54
definition of a gene [35], gives rise to the sort of allelic heterogeneity typically observed for human Mendelian   55
diseases [37], and to a distribution of GWAS "hit" minor allele frequencies [4, 26] consistent with empirical   56
results [36]. In this article, we explore this "gene-based" model under more complex demographic scenarios   57
as well as its properties with respect to the estimation of variance components using SNP-based   58
approaches [34] and twin studies. We also compare this model to the standard models of strictly additive   59
co-dominant effects, and multiplicative epistasis with dominance.   60

We further explore the power of several association tests to detect a causal gene region under each genetic   61
and demographic model. We find significant heterogeneity in the performance of burden tests [36, 38, 39]   62
across models of the trait and demographic history. We find that population expansion reduces the power to   63
detect causal gene-regions due to an increase in rare variation, in agreement with work by [22, 23]. The   64
behavior of the tests under different models provides us with insight as to the circumstances in which each   65
test is best suited.   66

In total, our results show that modeling gene action is key to modeling GWAS, and thus plays an   67
important role in both the design and interpretation of such studies. Further, the model of gene-based   68
recessivity best explains the differences between estimates of additive and dominance variance components   69
from SNP-based methods [27] and from twin studies [28] and is consistent with the distribution of   70
frequencies of significant associations in GWAS [4, 26]. Further, the genetic model plays a much more   71
important role than the demographic model, which is expected based on previous work on additive models   72

showing that the genetic load is approximately unaffected by changes in population size over time, [21, 22]. ₇₃
Consistent with recent work by [23], we find that rapid population growth in the recent past increases the ₇₄
contribution of rare variants to total genetic variance. However, we show here that different models of gene ₇₅
action are qualitatively different with respect to the partitioning of genetic variance across the allele ₇₆
frequency spectrum. We also show that these conclusions hold under the more complex demographic models ₇₇
that have been proposed for human populations [21, 40]. ₇₈

Results and Discussion ₇₉

The Models ₈₀

As in [36],we simulate a 100 kilobase region of human genome, contributing to a complex disease phenotype ₈₁
and fitness. The region evolves forward in time subject to neutral and deleterious mutation, recombination, ₈₂
selection, and drift. To perform genetic association and heritability estimation studies *in silico*, we need to ₈₃
impose a trait onto simulated individuals. In doing so, we introduce strong assumptions about the molecular ₈₄
underpinnings of a trait and its evolutionary context. ₈₅

How does the molecular genetic basis of a trait under natural selection influence population genetic ₈₆
signatures in the genome? This question is very broad, and therefore it was necessary to restrict ourselves to ₈₇
a small subset of molecular and evolutionary scenarios. We analyzed a set of approaches to modeling a single ₈₈
gene region experiencing recurrent unconditionally-deleterious mutation contributing to a quantitative trait ₈₉
subject to Gaussian stabilizing selection. Specifically, we studied three different genetic models and two ₉₀
different demographic models, holding the fitness model as a constant. Parameters are briefly described in ₉₁
Table 1. ₉₂

We implemented three disease-trait models of the phenotypic form $P = G + E$. G is the genetic ₉₃
component, and $E = N(0, \sigma_e^2)$ is the environmental noise expressed as a Gaussian random variable with ₉₄
mean 0 and standard deviation σ_e^2. In this context, σ_e^2 should be thought of as both the contribution from ₉₅
the environment and from the remaining genetic variance at loci in linkage equilibrium with the simulated ₉₆
100kb region. The genetic models are named the additive co-dominant (AC) model, multiplicative recessive ₉₇
(Mult. recessive; MR) model and the gene-based recessive (GBR) model. The MR model has a parameter, h, ₉₈
that controls the degree of recessivity; we call this model the complete MR (cMR) when $h = 0$ and the ₉₉
incomplete MR (iMR) when $0 \leq h \leq 1$. It is important to note that here recessivity is being defined in terms ₁₀₀
of phenotypic effects; this may be unusual for those more accustomed to dealing directly with recessivity for ₁₀₁
fitness effects. An idealized relationship between dominance for fitness effects and trait effects of a mutation ₁₀₂
on an unaffected genetic background is shown in S15 Fig. ₁₀₃

The critical conceptual difference between recessive models is whether dominance is a property of a locus ₁₀₄
(nucleotide/SNP) in a gene or the gene overall. Mathematically, this amounts to whether one first determines ₁₀₅
diploid genotypes at sites (and then multiplies across sites to get a total genetic effect) or calculates a score ₁₀₆
for each haplotype (the maternal and paternal alleles). For completely co-dominant models, this distinction ₁₀₇
is irrelevant, however for a model with arbitrary dominance one needs to be more specific. As an example, ₁₀₈
imagine a compound heterozygote for two biallelic loci, i.e. genotype Ab/aB. In the case of traditional ₁₀₉
multiplicative recessivity the compound heterozygote is wild type for both loci and therefore wild-type over ₁₁₀

all; this implies that these loci are in different genes (or independent functional units of the same gene) because the mutations are complementary. However, in the case of gene-based recessivity [36], neither haplotype is wild-type and so the individual is not wild-type; the failure of mutant alleles to complement defines these loci as being in the same gene [35].

For a diploid with m_i causative mutations on the i^{th} haplotype, we may define the additive model as

$$G_{AC} = \sum_{i=1}^{2} \sum_{j=1}^{m_i} c_{i,j}, \tag{1}$$

where $c_{i,j}$ is the effect size of the j^{th} mutation on the i^{th} haplotype. Each $c_{i,j}$ is sampled from an exponential distribution with mean of λ, to reflect unconditionally deleterious mutation. In other words, when a new mutation arises it's effect c is drawn from an exponential distribution, and remains constant throughout it's entire sojourn in the population.

The GBR model is the geometric mean of the sum of effect sizes on each haplotype [36]. We sum the causal mutation effects on each allele (paternal and maternal) to obtain a haplotype score. We then take the square root of the product of the haplotype scores to determine the total genetic value of the diploid.

$$G_{GBR} = \sqrt{\sum_{j=1}^{m_1} c_{1,j} \times \sum_{j=1}^{m_2} c_{2,j}} \tag{2}$$

Finally, the MR model depends on the number of positions for which a diploid is heterozygous (m_{Aa}) or homozygous (m_{aa}) for causative mutations,

$$G_{MR} = \left(\prod_{j=1}^{m_{Aa}} (1 + hc_j) \right) \left(\prod_{j=1}^{m_{aa}} (1 + 2c_j) \right) - 1. \tag{3}$$

Thus, $h = 0$ is a model of multiplicative epistasis with complete recessivity (cMR), and $h = 1$ closely approximates the additive model when effect sizes are small.

Here, phenotypes are subject to Gaussian stabilizing selection with an optimum at zero and standard deviation of $\sigma_s = 1$ such that the fitness, w, of a diploid is proportional to a Gaussian function [41].

$$w = e^{-\frac{P^2}{2\sigma_s^2}} \tag{4}$$

The AC and MR models draw no distinction between a "mutation" and a "gene" (as discussed in [36]). The GBR is also a recessive model, but recessivity is at the level of a *haplotype* (or allele) and is not an inherent property of individual mutations (see [36] for motivation of this model). Viewed in light of the traditional AC and MR models, the recessivity of a *site* in the GBR model is a function of the local genetic background on which it is found. Based on several qualitative comparisons we find that the GBR model is approximated by iMR models with $0.1 \leq h \leq 0.25$. However, no specific iMR model seems to match well in all aspects. The demographic models are that of a constant sized population (no growth) and rapid population expansion (growth).

The use of the MR model is inspired by Risch's work [30, 31], linking a classic evolutionary model of multiple loci interacting multiplicatively [42, 43] to the the genetic epidemiological parameter relative risk. Risch and Merikangas [44] used this model to calculate the power detect causal risk variants as a function of

their frequency and effect size. Pritchard extended Risch's model to consider a trait explicitly as a product of the evolutionary process [17]. Pritchard's work demonstrated that the equilibrium frequency distribution suggested an important role for rare deleterious mutations when a trait evolves in a constant sized, randomly mating population with recurrent mutation and constant effect sizes. However, multiplicative epistasis is only one model of gene action, and the effect of different genotype-to-phenotype models on the genetic architecture of traits and GWAS outcomes remains unknown. Exploring this issue is the focus of the current work.

Additive and dominance genetic variance in the population

The amount of narrow sense heritability, $h^2 = (V_A)/(V_P)$, explained by variants across the frequency spectrum is directly related to the effect sizes of those variants [29]. Thus, this measure is an important predictor of statistical power of GWAS and should inform decisions about study design and analysis [45]. Empirically, SNP-based estimates of heritability have inferred negligible dominance variance underlying most quantitative traits [27]. We have a particular interest in the amount of additive variance, V_A, that is due to rare alleles and how much of genetic variance, V_G, is attributable to V_A under different recessive models.

We follow the approach of [21], by calculating the cumulative percent of V_G explained by the additive effects of variants less than or equal to frequency x, $(V_{A;q \leq x})/(V_G)$. The product of this ratio and broad-sense heritability is an estimate of the narrow-sense heritability, h^2. This calculation is a population-wide equivalent to a SNP-based estimate of heritability in a population sample. In addition we calculate the same distribution for dominance effects $(V_{D;q \leq x})/(V_G)$ using the orthogonal model of [27]. Methods based on summing effect sizes [29] or the site frequency spectrum [21] would not apply to the GBR model, because the effect of a variant is not independent of other variants (*e.g.*, there is intralocus epistasis). Therefore, we resort to a regression-based approach, where we regress the genotypes of the population onto the total genetic value as defined in our disease trait models (see Material and Methods). In the limit of Hardy-Weinberg and linkage equilibrium, the regression estimates are equivalent to standard quantitative genetic estimates [29] (S14 Fig). For consistency, we applied the regression approach to all models. Overall, these distributions are substantially different across genetic models, demographic scenarios and model parameters (Fig 1).

Under the AC model, all of V_G is explained by additive effects if all variants are included in the calculation; in Fig 1 the solid variance curves reach unity in the AC panel. Low frequency and rare variants ($q < 0.01$) explain a large portion of narrow sense heritability (26% - 95%) even in models without rapid population expansion. Further, the variance explained at any given frequency threshold increases asymptotically to unity as a function of increasing λ (S4 Fig). While the total heritability of a trait in the population is generally insensitive to population size changes (S1 Fig, see also [21, 22, 46]), rapid population growth increases the fraction of additive genetic variation due to rare alleles (Fig 1).

Here, increasing λ corresponds to stronger selection against causative mutations, due to their increased average effect size. Recent work by Zuk et al. [24], takes a similar approach and relates the allele frequency distribution directly to design of studies for detecting the role of rare variants. However, our findings contrast with those of Zuk [24] and agree with those of Lohmueller [22], in that we predict that population expansion will substantially increase the heritability, or portion of genetic variance, that is due to rare variants. Our results under the AC model agree with those of Simons et al. [21], in that we find that increasing strength of selection, increasing λ in our work, increases the contribution to heritability of rare

variants. However, under the GBR model and the cMR model the distribution of genetic variance over risk ₁₇₉
allele frequency as function λ is non-monotonic (Fig 1 and S4 Fig). ₁₈₀

For all recessive models, we find that total V_A is less than V_G (Fig 1). For the MR models, all additional ₁₈₁
genetic variation is explained by the dominance variance component; in Fig 1 the dotted variance curves ₁₈₂
reach unity in the MR panels. As expected, genetic variation under the MR model with partial recessivity ₁₈₃
($h = 0.25$) is primarily additive [29, 47], whereas V_G under the cMR model ($h = 0$) is primarily due to ₁₈₄
dominance. The GBR model shows little dominance variance and is the only model considered here for ₁₈₅
which the total V_G explained by $V_A + V_D$ is less than the true V_G for all λ. This can be clearly seen in Fig 1 ₁₈₆
where the dotted curves do not reach unity in the GBR panel. These observations concerning the GBR ₁₈₇
model are consistent with the finding of [27] that dominance effects of SNPs do not contribute significantly ₁₈₈
to the heritability for complex traits. ₁₈₉

Under the GBR model, large trait values are usually due to compound heterozygote genotypes (*e.g.*, ₁₉₀
Ab/aB, where A and B represent different sites in the same gene) [36]. Therefore, the recessivity is at the ₁₉₁
level *of the gene region* while the typical approach to estimating V_A and V_D assigns effect sizes and ₁₉₂
dominance to individual mutations. Thus, compound heterozygosity, which is commonly observed for ₁₉₃
Mendelian diseases (see [36] and references therein) would be interpreted as variation due to *interactions* ₁₉₄
(epistasis) between risk variants. Importantly, the GBR model assumes that such interactions should be local, ₁₉₅
occurring amongst causal mutations in the same locus. While the GBR model is reflective of the original ₁₉₆
definition of a gene in which recessive mutations fail to complement, we emphasize that this does not imply ₁₉₇
that mutations are necessarily exomic. The GBR model is of a general genomic region in which mutations ₁₉₈
act locally in *cis* to disrupt the function of that region with respect to a phenotype. ₁₉₉

The increase in the number of rare alleles due to population growth is a well established theoretical and ₂₀₀
empirical result [48–61]. The exact relationship between rare alleles [4, 17, 26, 62, 63], and the demographic ₂₀₁
and/or selective scenarios from which they arose [21, 22, 64], and the genetic architecture of common complex ₂₀₂
diseases in humans is an active area of research. An important parameter dictating the relationships between ₂₀₃
demography, natural selection, and complex disease risk is the degree of correlation between a variant's effect ₂₀₄
on disease trait and its effect on fitness [18, 20–22]. In our simulations, we do not impose an explicit degree of ₂₀₅
correlation between the phenotypic and fitness effects of a variant. Rather, this correlation is context ₂₀₆
dependent, varying according to the current genetic burden of the population, the genetic background in ₂₀₇
which the variant is present and random environmental noise. However, if we re-parameterized our model in ₂₀₈
terms of [18], then we would have $\tau \leq 0.5$ (Gaussian function is greater than or equal to its quadratic ₂₀₉
approximation), which is consistent with recent attempts at estimating that parameter [20, 65]. Our ₂₁₀
approach is reflective of weak selection acting directly on the complex disease phenotype, but the degree to ₂₁₁
which selection acts on genotype is an outcome of the model. While the recent demographic history has little ₂₁₂
effect on key mean values such as broad-sense heritability of a trait or population genetic burden (S1 Fig and ₂₁₃
S3 Fig), the structure of the individual components in the population which add up to those mean values ₂₁₄
varies considerably. The specific predictions with respect to the composition of the populations varies ₂₁₅
drastically across different modeling approaches. It is therefore necessary to carefully consider the structure ₂₁₆
of a genetic model in a simulation study. ₂₁₇

The conclusions reached here also hold when we consider more complex demographic scenarios relevant to ₂₁₈

human populations. Under the demographic model for European populations from [40], the additive and $_{219}$
GBR models show the same behavior as in Fig 1 (S17 Fig). At all key time points where population size $_{220}$
changes, $V_A = V_G$ for the additive model, and the variance explained by rare mutations depends primarily on $_{221}$
λ (S17 Fig). For the GBR model, $V_A < V_G$ (as in Fig 1), and plateaus at the same ratio V_A/V_G for all time $_{222}$
points except immediately after the bottleneck, which results in a short-lived increase in V_A/V_G that is $_{223}$
undetectable by the time growth begins (S17 Fig). All recessive models (GBR, iMR and cMR) may show a $_{224}$
transient increase in total V_G after the bottleneck, depending on the value of λ (S18 Fig). However, the GBR $_{225}$
and iMR models with $h > 0.25$ showed a return to constant population size levels by the final time point. $_{226}$
The changes in V_A and V_G under recessive models is likely due to the transfer of non-additive variation into $_{227}$
V_A during a bottleneck, which has been studied thoroughly in the theoretical literature [66,67]. As in Fig 1, $_{228}$
the genetic model, and not the demographic details, drive the relationship between mutation frequency and $_{229}$
additive genetic variance. $_{230}$

Estimating additive and dominance variance from population samples $_{231}$

The previous section shows that the relationship between genetic variance and allele frequency in the entire $_{232}$
population strongly depends on the genetic model. Recent estimates of variance components from large $_{233}$
population samples of unrelated individuals have inferred that dominance variance (V_D) is negligible for most $_{234}$
traits [27]. However, a recent study of more than 10^4 Swedish twins and 18 traits obtained a contradictory $_{235}$
result, inferring significant non-additive variance for most traits, which was interpreted as V_D [68]. In this $_{236}$
section, we show that this apparent inconsistency is expected under certain models of gene action. $_{237}$

We applied GREMLd, MAF-stratified GREMLd (MS-GREMLd), and MAF-stratified Haseman-Elston $_{238}$
regression (see Methods for details). We found MS-GREMLd to be numerically unstable on our simulated $_{239}$
data, and thus we present results for non-MS-stratified GREMLd. The numerical stability issues likely $_{240}$
resulted from some combination of small number of SNPs per region ($\mathcal{O}(1000)$), low total V_G in a region, or $_{241}$
high variance in effect sizes across causal mutations [69]. Further, for large λ, where V_G is primarily due to $_{242}$
rare alleles (Fig 1), heritability in a sample may not reflect heritability in the entire population (S13 Fig). $_{243}$

Fig 2 shows the GREMLd additive and dominance heritability estimates, as compared to the respective $_{244}$
population value, over λ. Under the cMR model ($h = 0$), the dominance component is much larger than the $_{245}$
additive component as predicted from Fig 1. When GREMLd is performed on data after removing variants $_{246}$
with $MAF \leq 0.01$, as done in [27], the total heritability estimate (AD) is quite accurate. As anticipated, $_{247}$
GREMLd using unfiltered data yields results with a slight upward bias [70]. However, for the iMR ($h = 0.25$) $_{248}$
model the filtered GREMLd estimates are only accurate for $\lambda < 0.1$ reflecting the preponderance of rare $_{249}$
causal variants for larger values of λ. Unfiltered GREMLd estimates under the iMR ($h = 0.25$) model show a $_{250}$
slight upward bias for small values of λ, but are otherwise accurate. This shows that GREMLd is performing $_{251}$
as expected under the site-based model for which it is designed. The MS-HE regression results are generally $_{252}$
consistent with the GREMLd results. $_{253}$

The GREMLd and MS-HE estimates are accurate under the GBR model when λ is small, because most $_{254}$
heritability is additive in that case(Fig 1). However, under the GBR model, both filtered and unfiltered $_{255}$
GREMLd heritability estimates show downward bias when λ is large (Fig 2). The MS-HE regression results $_{256}$
reveal a similar pattern, which indicates that the downward bias for large values of λ is not strictly due to $_{257}$

removal of rare variants in the filtered GREMLd analysis. Instead, the bias shown for large values of λ is   258
likely due to the presence of substantial non-additive heritability, which is not captured by the dominance   259
effects of SNPs.   260

In contrast to the variance component methods, our simulated large twin studies provide approximately   261
unbiased estimates of total heritability for large values of λ, but were biased upward for small effect sizes   262
under the AC and GBR models (Fig 2). The variance in twin-study estimates was large enough that   263
study-to-study disagreement would be expected. Formally, twin studies estimate an additive and a   264
non-additive component of variance and interpreting the non-additive component as epistatic or dominance   265
variance is a matter of perspective. However, the GBR model is inspired by the definition of a gene as a   266
physical region in which recessive mutations leading to the same phenotypic outcome fail to complement [35],   267
consistent with the allelic heterogeneity observed for human Mendelian disorders (see [36] for further   268
discussion). Thus, the model of recessivity at the level of the gene region is picked up as non-additive   269
variance in twin studies, but missed by variance component methods (GREML and HE regression) because   270
the dominance in the GBR model is due to Ab/aB (compound heterozygotes) genotypes rather than a/a 271
genotypes (heterozygotes for a specific loss of function variant) assumed by variance component methods.   272
Thus the contradictory results of applying variance component methods [27] and analysis of large twin   273
studies [68] in order to estimate V_A and V_D may be interpreted as evidence for a model of gene action such   274
as the GBR, which may be viewed as either recessivity at the haplotype/gene level or intralocus epistasis at   275
the level of causative mutations in a single gene region. Both interpretations are valid. The alternative   276
explanation is that we must assert that one of the study designs is generating artifacts.   277

The genetic model affects the outcomes of GWAS   278

Both demography and the model of gene action affect the degree to which rare variants contribute to the   279
genetic architecture of a trait (Fig 1). However, the different mappings of genotype to phenotype from model   280
to model make it difficult to predict *a priori* the outcomes of GWAS under each model. Therefore, we sought   281
to explicitly examine the performance of statistical methods for GWAS under each genetic and demographic   282
model. We assessed the power of a single marker logistic regression to detect the gene region by calculating   283
the proportion of model replicates in which at least one variant reached genome wide significance at   284
$\alpha \leq 10^{-8}$ (Fig 3A). The basic logistic regression is equivalent to testing for association under the AC model.   285
We simulated both a perfect "genotyping chip" (all markers with $MAF \geq 0.05$) and complete re-sequencing   286
including all markers (Fig 3B).   287

Across all genetic models, the single marker logistic regression has less power under population expansion   288
(Fig 3A). The loss of power is attributable to a combination of rapid growth resulting in an excess of rare   289
variants overall [48–61], and the increasing efficacy of selection against causal variants in growing   290
populations [21]. While complete resequencing is more powerful than a gene-chip design, the relative power   291
gained is modest under growth (Fig 3A). Region-based rare variant association tests behave similarly with   292
respect to population growth (Fig 3B).   293

There are important differences in the behavior of the examined statistical methods across genetic models.   294
We focus first on the single marker tests (Fig 3A). For gene-chip strategies, power increases for "site-based"   295
models as recessivity of risk variants increases (compare power for AC, iMR, and cMR models in Fig 3B).   296

This increase in power is due to the well-known fact that recessive risk mutations are shielded from selection [297] when rare (due to being mostly present as heterozyogtes), thus reaching higher frequencies on average (S5 [298] Fig), and that the single-marker test is most powerful when risk variants are common [32]. Further, for the [299] complete multiplicative-recessive model (cMR), the majority of V_G is due to common variants (Fig 1), [300] explaining why resequencing does not increase power for this model (Fig 3A). [301]

For single-marker tests, the GBR model predicts large gains in power under re-sequencing for intermediate [302] λ (the mean trait-effect size of newly arising causal mutations), similar to the AC or iMR model. But, when [303] λ is larger power may actually be less under the GBR model than under AC or iMR. For all models, causal [304] mutations are more rare with increasing lambda (S7 Fig). However, as a function of frequency, all V_G may be [305] attributed to V_A or V_D in the site-specific models whereas there is increasing intralocus epistasis in the GBR [306] model as a function of λ (Fig 1). It is well-known that the single marker test has lower power when causal [307] mutations have low frequencies, are poorly tagged by more common SNPs, or have small main effects [32,71]. [308]

Region-based rare variant association tests show many of the same patterns across genetic model and [309] effect size distribution as single marker tests, but there are some interesting differences. The ESM [310] test [36,72] is the most powerful method tested for the AC, iMR, and GBR models (Fig 3b), with the [311] c-Alpha test as a close second in some cases. For those models, the power of naive SKAT, linear kernel SKAT [312] and SKAT-O, is always lower than the ESM and c-Alpha tests. This is peculiar since the c-Alpha test [313] statistic is the same as the linear kernel SKAT test. The major difference between SKAT and ESM/c-Alpha [314] is in the evaluation of statistical significance. SKAT uses an analytical approach to determine p-values while [315] the ESM/c-Alpha tests use an explicit permutation approach. This implies that using permutation based [316] p-values results in greater power. Yet, under the cMR model the linear kernel SKAT is the most powerful, [317] followed by c-alpha. The cMR model does not predict a significant burden of rare alleles and so the default [318] beta weights of SKAT are not appropriate, and the linear kernel is superior. The ESM test does poorly on [319] this model because there are not many marginally significant low-frequency markers. It is logical to think [320] that these tests would all perform better if all variants were included. The massive heterogeneity in the [321] performance of region-based rare variant tests across models strongly suggests that multiple methods should [322] be used when prior knowledge of underlying parameters is not available. In agreement with [22,73], we [323] predict that population growth reduces the power to associate variants in a causal gene region with disease [324] status (Fig 3) when the disease also impacts evolutionary fitness. We have recently released software to apply [325] the ESM test to case control data [72] in order to facilitate applying this test to real data. [326]

The distribution of minor allele frequencies of GWAS hits [327]

It was noted by [4,26], that an excess of rare significant hits, relative to empirical data, is predicted by AC [328] models where large effect mutations contribute directly to fitness and the disease trait. We confirm that AC [329] models are inconsistent with the empirical data (Fig 4), except when $\lambda \leq 0.01$. The empirical data in Fig 4 [330] represent a pooled data set with the same diseases and quality filters as in [26], but updated to include more [331] recent data. The data are described in S1 Table, and can be visualized alone more clearly in S16 Fig. Close [332] to half of the data comes from GWAS studies uploaded to the NHGRI database after 2011, yet the same [333] qualitative pattern is observed. This contradicts the hypothesis that the initial observation of an excess of [334] common significant hits relative to the prediction under an AC model was simply due to small sample sizes [335]

and low marker density in early GWAS previously analyzed in [4, 26]. Yet the initial observation is in fact ₃₃₆
robust and the meta-pattern provides an appropriate point of comparison when considering the compatibility ₃₃₇
of explicit population-genetic models with existing GWAS data. ₃₃₈

The GBR model predicts few rare significant hits and an approximately uniform distribution across the ₃₃₉
remainder of MAF domain (Fig 4), even for intermediate and large values of λ. For smaller values of λ, the ₃₄₀
GBR predicts an excess of common significant hits. The more uniform distribution of significant single ₃₄₁
markers seen under the GBR is consistent with the flatter distribution of genetic variance (Fig 1). Under the ₃₄₂
GBR model and population growth, a KS test (marginal $\alpha = 0.05$) cannot reject identical distributions of ₃₄₃
MAF's of significant hits for all values of λ (S19 Fig). The cMR ($h = 0$) model shows an excess of ₃₄₄
intermediate frequency variants, although this does not result in rejection under the KS test (S19 Fig). If one ₃₄₅
considers trying to determine an approximate dominance coefficient in the GBR model, it would be found ₃₄₆
that there is a distribution of coefficients across sites. Yet, when simulating iMR model, we find that an ₃₄₇
intermediate degree of dominance, $h = 0.25$, results in distribution of significant hits which is similar to the ₃₄₈
GBR results (Fig 4). ₃₄₉

We note that there is no compelling reason to expect any specific value of λ to be a particularly good fit ₃₅₀
to the empirical data. The empirical data are composed of genome-wide data for multiple traits. We feel that ₃₅₁
the mutational parameters, λ and mutation rate to causal variants, are likely to vary across the genome and ₃₅₂
across traits. Thus, the empirical data reflect a mixture of different underlying models and ascertainment ₃₅₃
schemes. The reason we emphasize this feature of the data is to demonstrate that models with rare alleles of ₃₅₄
large effect do not necessarily imply an observed excess of rare significant GWAS hits. ₃₅₅

In consideration of the rare allele of large effect hypothesis, [62] proposed a model where multiple rare ₃₅₆
alleles dominate disease risk and create synthetic associations with common SNPs. However, later it was ₃₅₇
shown that this particular model was inconsistent with GWAS theoretically and empirically [4, 26, 74]. This ₃₅₈
inconsistency remained un-reconciled in the literature until now. We find that the MAF distribution of ₃₅₉
significant hits in a GWAS varies widely with choice of genetic model. In particular, we confirm the results of ₃₆₀
Wray et al. [26], that AC evolutionary models predict an excess of low frequency significant hits unless trait ₃₆₁
effect sizes are quite small. Also, the cMR model predicts an excess of intermediate and common significant ₃₆₂
hits. Utilizing a GBR model or an intermediate iMR model with $h = 0.25 - 0.5$, reconciles this inconsistency ₃₆₃
by simultaneously predicting the importance of rare alleles of large effect and the correct allele frequency ₃₆₄
distribution among statistically significant single markers. ₃₆₅

Conclusion ₃₆₆

Several empirical observations provide support for the presence of gene-based recessivity underlying variation ₃₆₇
for some complex traits in humans. The minor allele frequency distribution of significant GWAS hits is ₃₆₈
relatively flat [4, 75], which our results show is consistent with either the presence of small additive effect loci ₃₆₉
or gene-/site-based partially-recessive loci with intermediate to large effects (Fig 4). Models with loci of large ₃₇₀
additive effects predict an excess of rare significant hits. Oppositely, models with complete site-based ₃₇₁
recessivity predict an excess of common significant hits for all simulated mutation effect size distributions. ₃₇₂

SNP based estimates of dominance heritability are much lower than estimates of dominance from ₃₇₃
twins [27, 68]. Of the models we explored, only the gene-based recessive model with intermediate to large ₃₇₄

effects is consistent with difference between twin and SNP based estimates of dominance variance (Fig 2). ₃₇₅
Under a site-based recessive model of partial recessivity (*e.g.* h =0.25), there should be no significant ₃₇₆
difference between estimates of dominance variance from SNP and twin studies, provided that the statistical ₃₇₇
assumptions are met for both approaches (Fig 2). Our findings support a more thorough investigation into ₃₇₈
the importance of compound heterozygosity in the genetics complex-traits. However, it may be difficult to ₃₇₉
directly observe non-additive gene-level effects through analysis of individual SNP markers. ₃₈₀

Additionally, the genetic model appears to be important in the design and analysis of association studies. ₃₈₁
While changes in population size do affect the relationship between effect size and mutation frequency [48–61] ₃₈₂
(Fig 1 and S5 Fig), different mappings of genotype to trait value do this in radically different ways for the ₃₈₃
same demographic history (Fig 1). From an empirical perspective, our findings suggest that re-sequencing in ₃₈₄
large samples is likely the best way forward in the face of the allelic heterogeneity imposed by the presence of ₃₈₅
rare alleles of large effect. Re-sequencing of candidate genes [76–79] and exomes [40, 80–85] in case-control ₃₈₆
panels have observed an abundance of rare variants associated with case status. Here we show that under a ₃₈₇
model of mutation-selection balance at genes, neither current single-marker nor popular multi-marker tests ₃₈₈
are especially powerful at detecting large genomic regions harboring multiple risk variants (Fig 3). However, ₃₈₉
we show that using permutations to derive p-values improves the power of SKAT [69] with a linear kernel ₃₉₀
(equivalent test statistic to c-Alpha [38]). Similarly, another permutation based test, the ESM test [72], has ₃₉₁
more robust power across demographic and genetic models (Fig 3). ₃₉₂

Conceptually, *cis*-effects arise naturally from the original definition of a gene in which mutant recessive ₃₉₃
alleles fail to complement [35]. We show that *cis*-effects within a locus, represented by the GBR model, can ₃₉₄
have an important impact on the population level architecture of a complex trait. This conclusion is ₃₉₅
important for future simulation studies as well as the interpretation of empirical data. It is important to note ₃₉₆
that despite our use of the term "gene-based" this model may apply to any functional genomic element in ₃₉₇
which there are multiple mutable sites affecting a trait in in *cis*, not just to genes. From a theoretical ₃₉₈
perspective, our work motivates the development of a more generalized gene-based model to include arbitrary ₃₉₉
dominance and arbitrary locus size. Empirically, we find that the GBR model is broadly consistent with a ₄₀₀
variety of observations from the human statistical genetics literature. Thus, there is an evident need for ₄₀₁
improved region-based association tests and the development of genetic variance component methods for ₄₀₂
haplotypes. ₄₀₃

Materials and Methods ₄₀₄

Forward simulation ₄₀₅

Using the fwdpp template library v0.2.8 [86], we implemented a forward in time individual-based simulation ₄₀₆
of a Wright-Fisher population with mutation under the infinitely many sites model [87], recombination, and ₄₀₇
selection occurring each generation. We simulated populations of size $N = 2e4$ individuals for a time of $8N$ ₄₀₈
generations with a neutral mutation rate of $\mu = 0.00125$ per gamete per generation and a per diploid per ₄₀₉
generation recombination rate of $r = 0.00125$. Deleterious mutations occurred at a rate of $\mu_d = 0.1\mu$ per ₄₁₀
gamete per generation. These parameters correspond to $\theta = 4N\mu = \rho = 4Nr = 100$ and thus our simulation ₄₁₁
approximates a 100Kb region of the human genome. For simulations with growth, we simulated an additional ₄₁₂

500 generations of exponential growth from $N_i = 2e4$ to $N_{final} = 1e6$. This demographic model is much simpler than current models fit to empirical data [58]. However, this simple model allows us to more easily get a sense of the impact of population expansion [21, 22]. 250 simulation trials were performed for each parameter/model combination.

Exploring the gene region's contribution to heritability

Broad-sense heritability can be calculated directly from our simulated data as $H^2 = \frac{V_G}{V_P}$. We explored broad-sense heritability as a function of mean causative effect size under each model. We compare our simulation results to $H^2 \sim 4\mu_d V_s$ for additive models and $H^2 \sim 2\mu_d V_s$ for recessive models [88, 89]. In our simulations, $V_s = 1$, and we tuned the environmental variance V_e to generate simulations for which $E[H^2] \sim 0.04$ or ~ 0.08.

Determining the genetic load of the population

Genetic load is defined as the relative deviation in a populations fitness from the fitness optimum, $L = (w_{max} - \bar{w})/(w_{max})$. We set the phenotypic optimum to be zero; $P_{opt} = 0$. When determining fitness for the SBR models, we subtract one from all phenotypes. This implies that $w_{max} = e^{-\frac{P_{opt}^2}{2\sigma_s^2}} = 1$ and that load is a simple function of the phenotypes of the population, $L = 1 - e^{-\frac{P^2}{2\sigma_s^2}}$. We also used the mean number of mutations per individual, and the mean frequency and effect sizes of segregating risk variants as proxies for the genetic load [21, 90].

Additive and dominance genetic variance over allele frequency

We used an approach based sequential (type-1) regression sums of squares to estimate the contribution of the additive and dominance effects of variants to the total genetic variation due to a locus. Given a genotype matrix (rows are individuals and columns are risk variants) of (0,1, or 2) copies of a risk allele (*e.g.* all mutations affecting phenotype), we sort the columns by decreasing risk mutation frequency. Then, within frequency classes, columns were sorted by decreasing effect sizes. For each variant a dominance component was also coded as 0, 2q, or 4q-2 according to the orthogonal model of [27], where q is the frequency of the variant in the population. We then used the R package `biglm` [91] to regress the individual genetic values (G in the previous section) onto this matrix. The variance explained by the additive and dominance effects of the m markers with $q \leq x$ is then approximately $r^2 = (\sum_{i=1}^{m} \Sigma SS_{reg,i})/(SS_{tot})$. Averaging results across replicates, this procedure results in a Monte-Carlo estimate of the fraction of V_G that is due to additive and dominance effects of variants with population frequency less than or equal to x is $(V_{A;q \leq x} + V_{D;q \leq x})/(V_{G;q \leq 1})$ [21]. This fraction can be easily partitioned into strictly additive and dominance components.

Additive and dominance heritability in random population samples

We employed three different SNP-based approaches to estimating heritability from population samples: GREMLd, minor allele frequency stratified(MS) GREMLd [27], and MS-Haseman-Elston (HE)

regression [92,93]. For comparison, we calculated the true total heritability in the sample as $H^2_{sample} = (V_{G;sample})/(V_{P;sample})$. Unfortunately, due to the nature of our simulated data MS-GREMLd did not result in sufficiently reliable results. Under MS-GREMLd, many replicates resulted in numerical errors in GCTA. These problems were present at a rate of less than 1/100 replicates using non-MS GREMLd, but were increased by splitting the data into multiple GRMs.

Using raw individual phenotypic values as quantitative trait values, random samples from simulated populations (n=6000) were converted to .bed format using PLINK 1.90a [94]. PLINK was also used to test for HWE ($p < 1e - 6$) and filter on minor allele frequency. GCTA 1.24.4 [34] was used to make genetic relatedness matrices (GRM) for both additive and dominance components with the flags –autosome and –make-grm(-d).

For non-MS runs, we tested the effect of filtering on MAF by performing the analysis on unfiltered datasets and with markers with $MAF < 0.01$ removed. For MS estimates we stratified the additive and dominance GRM's into two bins $MAF \leq 0.01$ and $MAF > 0.01$. GREMLd analysis was performed in GCTA with Fisher scoring, no variance component constraint and a max of 200 iterations. MS-HE regression was carried out by regressing the off diagonal elements of each GRM onto the cross product of the scaled and centered phenotypes in a multiple linear regression setting in R [95].

Twin studies

To simulate twin studies we sampled 2000 monozygotic (MZ) and 2000 dizygotic (DZ) twins pairs from the final generation of the simulations. Parents were sampled randomly without replacement. MZ twin pairs were formed by sampling a single gamete pair, one recombinant from each parent, and two environmental random deviates. DZ twin pairs were formed by sampling two gamete pairs, two recombinant gametes from each parent, and two environmental random deviates. Our simulated studies are ideal in that there are no correlated environmental effects, but potentially problematic due to low total heritability. We explored the use of structural equation modeling (SEM) using the package OpenMx [96], but chose to rely strictly on estimates of twin correlation obtained directly from the data. For monozygotic (MZ) twins, we used only a single child gamete pair with two unique environmental deviates. For dizygotic (DZ) twins we used two child gamete pairs, each with a unique environmental deviate. Broad sense heritability is the correlation between MZ twin pairs; $H^2 = r_{MZ}$. Under a purely additive model, the DZ twin correlation should be half of the MZ twin correlation. Non-additive genetic components of phenotypic variance reduce the DZ twin correlation. If all non-additive heritability is due to dominance, then the dominance heritability can be calculated as twice the difference between the MZ twin correlation and two-times the DZ twin correlation: $\delta^2 = 2 * (r_{MZ} - 2 * r_{DZ})$. The additive heritability can then be calculated as the difference between the broad-sense and non-additive component: $h^2 = H^2 - \delta^2 = 4 * r_{DZ} - r_{MZ}$ [29].

These direct estimates of MZ and DZ twin correlations in our simulations are reliable as we have no measurement error, shared environmental effects, gene-by-environment effects, or gene-by-gene interactions. Additionally, we only simulate a single genomic region contributing $H^2 \sim 0.04$, which made use of SEM difficult numerically. This creates a limitation in that we can not discuss when a model with dominance is a better fit to the data than the additive only model. But, the benefit of using direct estimates is that we can clearly see what signals are present in the data. To further clarify the data visualization, we pooled our 512

twin-study replicates into groups of 8, creating 64 sets of MZ-DZ twin phenotypes. This did not have an ₄₈₆ effect on the central tendencies of our estimates, but it reduced the variance. The twin study error bars in ₄₈₇ Fig 2 are based on 64 sets of 64,000 individuals, which is larger than a typical twin study. However, one ₄₈₈ reason our results have high variance is because we only simulate a single locus, rather than a whole trait. ₄₈₉

Case-control studies ₄₉₀

Following [36], we sampled 3000 cases and 3000 controls from each simulated population. Cases were ₄₉₁ randomly sampled from the upper 15% of phenotypic values in the population, and controls were randomly ₄₉₂ sampled from within 1 standard deviation of the population mean. This is the liability scale model (see [29]). ₄₉₃ We define a "GWAS" to be a study including all markers with MAF $\geq 5\%$ and a re-sequencing study to ₄₉₄ include all markers. In all cases we used a minor allele count logistic regression as the single marker test. For ₄₉₅ single marker tests, the p-value cut off for significance is $p \leq 1e-08$ which is common in current ₄₉₆ GWAS [62,97]. Power is determined by the percentage of simulation replicates in which at least one marker ₄₉₇ reaches genome wide significance. ₄₉₈

Region-based tests of association due to rare alleles ₄₉₉

We applied multiple region-based tests to our simulated data, ESM_K [36], several variations of SKAT [39] ₅₀₀ and c-Alpha [38]. We used the `R` package from the SKAT authors to implement their test ₅₀₁ (`http://cran.r-project.org/web/packages/SKAT/index.html`). The remaining tests were implemented ₅₀₂ in a custom `R` package (see SOFTWARE AVAILABILITY below). For the ESM_K and c-Alpha we performed up ₅₀₃ to $2e6$ permutations of case-control labels to determine empirical p-values. Common variants ($q \geq 0.05$) were ₅₀₄ removed prior to performing region-based rare variant association tests. ₅₀₅

Distribution of Significant GWAS Hits ₅₀₆

Following [4,26], we calculated the distribution of the minor allele frequency (MAF) of the most significant ₅₀₇ SNPs in a GWAS in empirical and simulated data. The empirical data was obtained from the NHGRI-EBI ₅₀₈ GWAS database (`http://www.ebi.ac.uk/gwas/`) on 02/05/2015. We considered the same diseases and ₅₀₉ applied the same filters as in Table 3 of [26]. Specific information regarding the empirical data can be ₅₁₀ obtained in S1 Table. ₅₁₁

In order to mimic ascertained SNP data, we sampled markers from our case/control panels according to ₅₁₂ their minor allele frequencies [98], as done in [36]. Additionally, we removed all markers with MAF < 0.01 to ₅₁₃ reflect common quality controls used in GWAS. The simulated data were grouped by genetic model, ₅₁₄ demographic scenario, heritability level, and mutation effect distribution. We then plotted the minor allele ₅₁₅ frequency of the most significant marker with a single-marker score $-log_{10}(p) \geq 8$, for all replicates where ₅₁₆ significant markers were present.Finally, we performed a two-sample KS test in `R` between each group of ₅₁₇ simulated GWAS hit allele frequencies and the empirical data. ₅₁₈

Human demography

We simulated a demographic model for Europeans based on [40] as described in [21]. For simplicity, we ignored migration between the European (EA) and African American (AA) populations. The model was implemented using the Python package `fwdpy` version 0.0.4, which uses `fwdpp` [86] version 0.5.1 as a C++ back-end. During the evolution of the EA population, we recorded the genetic variance in the population, V_G, and the number of deleterious mutations per diploid (a measure of genetic load [21]) every 50 generations. In a separate set of simulations, we applied the regression method described above to calculate cumulative additive genetic variance as a function of allele frequency. Because the regressions are computationally demanding, we applied the method in the generation immediately before, and at the start of, any changes in population size.

These simulations were run with no neutral mutations, and the recombination rate and mutation rate to causative mutations were the same as in the simulations described above.

The Python scripts for these simulations and iPython/Jupyter notebooks used for generating figures are available online (see Software availability section below).

Software availability

Our simulation code and code for downstream analyses are freely available at

- `http://github.com/ThorntonLab/disease_sims`

- `http://github.com/molpopgen/buRden`

- `http://github.com/molpopgen/fwdpy`

- `http://github.com/molpopgen/TennessenEAonly`

References

1. Welter D, MacArthur J, Morales J, Burdett T, Hall P, Junkins H, et al. The NHGRI GWAS Catalog, a curated resource of SNP-trait associations. Nucleic Acids Research. 2014;42(December 2013):1001–1006.

2. Manolio TA, Collins FS, Cox NJ, Goldstein DB, Hindorff LA, Hunter DJ, et al. Finding the missing heritability of complex diseases. Nature. 2009 oct;461(7265):747–53. Available from: `http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2831613{&}tool= pmcentrez{&}rendertype=abstract`.

3. Visscher PM, Brown MA, McCarthy MI, Yang J. Five years of GWAS discovery. American journal of human genetics. 2012 jan;90(1):7–24. Available from: `http://www.sciencedirect.com/science/article/pii/S0002929711005337`.

4. Gibson G. Rare and common variants: twenty arguments. Nature Reviews Genetics. 2012;13(2):135–145. Available from: `http://dx.doi.org/10.1038/nrg3118`.

5. Robinson MR, Wray NR, Visscher PM. Explaining additional genetic variation in complex traits. Trends in Genetics. 2014;30:124–132. Available from: `http://dx.doi.org/10.1016/j.tig.2014.02.003`.

6. Eichler EE, Flint J, Gibson G, Kong A, Leal SM, Moore JH, et al. Missing heritability and strategies for finding the underlying causes of complex disease. Nature reviews Genetics. 2010 jun;11(6):446–50. Available from: `http://dx.doi.org/10.1038/nrg2809`.

7. Wei WH, Hemani G, Haley CS. Detecting epistasis in human complex traits. Nature Reviews Genetics. 2014;15(11):722–733. Available from: `http://dx.doi.org/10.1038/nrg3747`.

8. Zuk O, Hechter E, Sunyaev SR, Lander ES. The mystery of missing heritability: Genetic interactions create phantom heritability. Proceedings of the National Academy of Sciences. 2012;109(4):1193–1198.

9. Fisher RA. The Genetical Theory Of Natural Selection : Fisher, R. A : Free Download & Streaming : Internet Archive. Oxford, UK; 1930. Available from: `https://archive.org/details/geneticaltheoryo031631mbp`.

10. Visscher PM, Hill WG, Wray NR. Heritability in the genomics era–concepts and misconceptions. Nature reviews Genetics. 2008 apr;9(4):255–66. Available from: `http://dx.doi.org/10.1038/nrg2322`.

11. McClellan J, King MC. Genetic heterogeneity in human disease. Cell. 2010 apr;141(2):210–7. Available from: `http://www.sciencedirect.com/science/article/pii/S009286741000320X`.

12. Cirulli ET, Goldstein DB. Uncovering the roles of rare variants in common disease through whole-genome sequencing. Nature reviews Genetics. 2010 jun;11(6):415–25. Available from: `http://dx.doi.org/10.1038/nrg2779`.

13. Visscher PM, Goddard ME, Derks EM, Wray NR. Evidence-based psychiatric genetics, AKA the false dichotomy between common and rare variant hypotheses. Molecular psychiatry. 2012 may;17(5):474–85. Available from: http://dx.doi.org/10.1038/mp.2011.65.

14. Yang J, Benyamin B, McEvoy BP, Gordon S, Henders AK, Nyholt DR, et al. Common SNPs explain a large proportion of the heritability for human height. Nature Genetics. 2010 jul;42(7):565–569. Available from: http://www.nature.com/doifinder/10.1038/ng.608.

15. Golan D, Lander ES, Rosset S. Measuring missing heritability: Inferring the contribution of common variants. Proceedings of the National Academy of Sciences of the United States of America. 2014;.

16. Yang J, Bakshi A, Zhu Z, Hemani G, Vinkhuyzen AAE, Lee SH, et al. Genetic variance estimation with imputed variants finds negligible missing heritability for human height and body mass index. Nature Genetics. 2015 aug;advance on. Available from: http://dx.doi.org/10.1038/ng.3390.

17. Pritchard JK. Are rare variants responsible for susceptibility to complex diseases? American journal of human genetics. 2001 jul;69(1):124–37. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1226027{&}tool=pmcentrez{&}rendertype=abstract.

18. Eyre-Walker A. Evolution in health and medicine Sackler colloquium: Genetic architecture of a complex trait and its implications for fitness and genome-wide association studies. Proceedings of the National Academy of Sciences of the United States of America. 2010;107 Suppl:1752–1756.

19. Park JH, Wacholder S, Gail MH, Peters U, Jacobs KB, Chanock SJ, et al. Estimation of effect size distribution from genome-wide association studies and implications for future discoveries. Nature genetics. 2010 jul;42(7):570–5. Available from: http://dx.doi.org/10.1038/ng.610.

20. Agarwala V, Flannick J, Sunyaev S, Altshuler D. Evaluating empirical bounds on complex disease genetic architecture. Nature genetics. 2013;45(12):1418–27. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24141362.

21. Simons YB, Turchin MC, Pritchard JK, Sella G. The deleterious mutation load is insensitive to recent population history. Nature genetics. 2014;46(August 2013):220–4. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24509481.

22. Lohmueller KE. The impact of population demography and selection on the genetic architecture of complex traits. PLoS genetics. 2014 may;10(5):e1004379. Available from: http://dx.plos.org/10.1371/journal.pgen.1004379.

23. Uricchio LH, Zaitlen NA, Ye CJ, Witte JS, Hernandez RD. Selection and explosive growth alter genetic architecture and hamper the detection of causal rare variants. Genome Research. 2016 may;Available from: http://genome.cshlp.org/lookup/doi/10.1101/gr.202440.115.

24. Zuk O, Schaffner SF, Samocha K, Do R, Hechter E, Kathiresan S, et al. Searching for missing heritability: designing rare variant association studies. Proceedings of the National Academy of Sciences of the United States of America. 2014;111:E455–64. Available from:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3910587{&}tool=
pmcentrez{&}rendertype=abstract.

25. Caballero A, Tenesa A, Keightley PD. The Nature of Genetic Variation for Complex Traits Revealed by GWAS and Regional Heritability Mapping Analyses. Genetics. 2015 oct;201(4):1601–13. Available from: http://genetics.org/content/201/4/1601.abstract.

26. Wray NR, Purcell SM, Visscher PM. Synthetic associations created by rare variants do not explain most GWAS results. PLoS Biology. 2011;9(1).

27. Zhu Z, Bakshi A, Vinkhuyzen AE, Hemani G, Lee S, Nolte I, et al. Dominance Genetic Variation Contributes Little to the Missing Heritability for Human Complex Traits. The American Journal of Human Genetics. 2015;p. 377–385. Available from: http://linkinghub.elsevier.com/retrieve/pii/S0002929715000099.

28. Chen HS, Hutter CM, Mechanic LE, Amos CI, Bafna V, Hauser ER, et al. Genetic simulation tools for post-genome wide association studies of complex diseases. Genetic epidemiology. 2015 jan;39(1):11–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25371374.

29. Falconer D, Mackay TFC. Introduction to Quantitative Genetics, Ed. 2. Harlow, Essex, UK: Longmans Green; 1996.

30. Risch N. Linkage strategies for genetically complex traits. III. The effect of marker polymorphism on analysis of affected relative pairs. American journal of human genetics. 1990;46:242–253.

31. Risch N. Linkage strategies for genetically complex traits. I. Multilocus models. American journal of human genetics. 1990;46:222–228.

32. Spencer CCA, Su Z, Donnelly P, Marchini J. Designing Genome-Wide Association Studies: Sample Size, Power, Imputation, and the Choice of Genotyping Chip. PLoS Genetics. 2009 may;5(5):e1000477. Available from: http://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1000477.

33. Wellcome T, Case T, Consortium C. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. Nature. 2007;447(June):661–678.

34. Yang J, Lee SH, Goddard ME, Visscher PM. GCTA: a tool for genome-wide complex trait analysis. American journal of human genetics. 2011 jan;88(1):76–82. Available from: http://www.sciencedirect.com/science/article/pii/S0002929710005987.

35. Benzer S. Fine Structure of a Genetic Region in Bacteriophage. Proceedings of the National Academy of Sciences of the United States of America. 1955 jun;41(6):344–354. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC528093/?tool=pmcentrez.

36. Thornton KR, Foran AJ, Long AD. Properties and Modeling of GWAS when Complex Disease Risk Is Due to Non-Complementing, Deleterious Mutations in Genes of Large Effect. PLoS Genetics. 2013;9(2).

37. Strachan, T and Read A. Human Molecular Genetics (New York: Garland Science, Taylor \& Francis Group). 2011;.

38. Neale BM, Rivas Ma, Voight BF, Altshuler D, Devlin B, Orho-Melander M, et al. Testing for an unusual distribution of rare variants. PLoS Genetics. 2011;7(3).

39. Wu MC, Lee S, Cai T, Li Y, Boehnke M, Lin X. Rare-variant association testing for sequencing data with the sequence kernel association test. American Journal of Human Genetics. 2011;89(1):82–93. Available from: `http://dx.doi.org/10.1016/j.ajhg.2011.05.029`.

40. Tennessen Ja, Bigham aW, O'Connor TD, Fu W, Kenny EE, Gravel S, et al. Evolution and Functional Impact of Rare Coding Variation from Deep Sequencing of Human Exomes. Science. 2012;337(6090):64–69.

41. Bürger RR. The mathematical theory of selection, recombination, and mutation. Wiley; 2000.

42. Haldane JBS. The cost of natural selection; 1957.

43. Charlesworth B, Morgan MT, Charlesworth D. The effect of deleterious mutations on neutral molecular variation. Genetics. 1993 aug;134(4):1289–1303. Available from: `http://www.genetics.org/content/134/4/1289.abstract`.

44. Risch N, Merikangas K. The future of genetic studies of complex human diseases. Science (New York, NY). 1996 sep;273(5281):1516–7. Available from: `http://www.ncbi.nlm.nih.gov/pubmed/8801636`.

45. Sham PC, Purcell SM. Statistical power and significance testing in large-scale genetic studies. Nature reviews Genetics. 2014 may;15(5):335–46. Available from: `http://dx.doi.org/10.1038/nrg3706`.

46. Uricchio LH, Witte JS, Hernandez RD. Selection and explosive growth may hamper the performance of rare variant association tests; 2015. Available from: `http://www.biorxiv.org/content/early/2015/03/01/015917.abstract`.

47. Hill WG, Goddard ME, Visscher PM. Data and theory point to mainly additive genetic variance for complex traits. PLoS genetics. 2008 feb;4(2):e1000008. Available from: `http://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1000008`.

48. Griffiths RC, Tavaré S. Sampling theory for neutral alleles in a varying environment. Philosophical transactions of the Royal Society of London Series B, Biological sciences. 1994 jun;344(1310):403–10. Available from: `http://www.ncbi.nlm.nih.gov/pubmed/7800710`.

49. Griffiths RC, Tavare S. Ancestral Inference in Population Genetics. Statistical Science. 1994 aug;9(3):307–319. Available from: `http://projecteuclid.org/euclid.ss/1177010378`.

50. Rogers A, Harpending H. Population growth makes waves in the distribution of pairwise genetic differences. Mol Biol Evol. 1992 may;9(3):552–569. Available from: `http://mbe.oxfordjournals.org/content/9/3/552`.

51. Fu YX. Statistical Tests of Neutrality of Mutations Against Population Growth, Hitchhiking and Background Selection. Genetics. 1997 oct;147(2):915–925. Available from: `http://www.genetics.org/content/147/2/915.short`.

52. Beaumont MA. Detecting Population Expansion and Decline Using Microsatellites. Genetics. 1999 dec;153(4):2013–2029. Available from: `http://www.genetics.org/content/153/4/2013.short`.

53. Aris-Brosou S, Excoffier L. The impact of population expansion and mutation rate heterogeneity on DNA sequence polymorphism. Molecular Biology and Evolution. 1996 mar;13(3):494–504. Available from: `http://mbe.oxfordjournals.org/content/13/3/494.short`.

54. Coventry A, Bull-Otterson LM, Liu X, Clark AG, Maxwell TJ, Crosby J, et al. Deep resequencing reveals excess rare recent variants consistent with explosive population growth. Nature communications. 2010 jan;1:131. Available from: `http://dx.doi.org/10.1038/ncomms1130`.

55. Gao F, Keinan A. High burden of private mutations due to explosive human population growth and purifying selection. BMC genomics. 2014;15 Suppl 4(Suppl 4):S3. Available from: `http://www.ncbi.nlm.nih.gov/pubmed/25056720`.

56. Gazave E, Chang D, Clark AG, Keinan A. Population growth inflates the per-individual number of deleterious mutations and reduces their mean effect. Genetics. 2013;195(3):969–978.

57. Gazave E, Ma L, Chang D, Coventry A, Gao F, Muzny D, et al. Neutral genomic regions refine models of recent rapid human population growth. Proceedings of the National Academy of Sciences of the United States of America. 2014;111(2):757–62. Available from: `http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3896169{&}tool=pmcentrez{&}rendertype=abstract`.

58. Gravel S, Henn BM, Gutenkunst RN, Indap AR, Marth GT, Clark AG, et al. Demographic history and rare allele sharing among human populations. Proceedings of the National Academy of Sciences. 2011 jul;108(29):11983–11988. Available from: `http://www.pnas.org/content/108/29/11983.short`.

59. Keinan a, Clark aG. Recent Explosive Human Population Growth Has Resulted in an Excess of Rare Genetic Variants. Science. 2012;336(6082):740–743.

60. Kimmel M, Chakraborty R, King JP, Bamshad M, Watkins WS, Jorde LB. Signatures of Population Expansion in Microsatellite Repeat Data. Genetics. 1998 apr;148(4):1921–1930. Available from: `http://www.genetics.org/content/148/4/1921.short`.

61. Reich DE, Goldstein DB. Genetic evidence for a Paleolithic human population expansion in Africa. Proceedings of the National Academy of Sciences. 1998 jul;95(14):8119–8123. Available from: `http://www.pnas.org/content/95/14/8119.full`.

62. Dickson SP, Wang K, Krantz I, Hakonarson H, Goldstein DB. Rare Variants Create Synthetic Genome-Wide Associations. PLoS Biology. 2010;8(1).

63. Maher MC, Uricchio LH, Torgerson DG, Hernandez RD. Population genetics of rare variants and   711
    complex diseases. Human heredity. 2012 jan;74(3-4):118–28. Available from: 712
    `http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3698246{&}tool=`   713
    `pmcentrez{&}rendertype=abstract`.   714

64. Reich DE, Lander ES. On the allelic spectrum of human disease. Trends in Genetics. 2001   715
    sep;17(9):502–510. Available from: 716
    `http://www.sciencedirect.com/science/article/pii/S0168952501024106`.   717

65. Mancuso N, Rohland N, Rand KA, Tandon A, Allen A, Quinque D, et al. The contribution of rare   718
    variation to prostate cancer heritability. Nature genetics. 2015 nov;advance on. Available from: 719
    `http://dx.doi.org/10.1038/ng.3446`.   720

66. Naciri-Graven Y, Goudet2 J. THE ADDITIVE GENETIC VARIANCE AFTER BOTTLENECKS IS   721
    AFFECTED BY THE NUMBER OF LOCI INVOLVED IN EPISTATIC INTERACTIONS.   722
    Evolution. 2003;57(574):706–716. Available from: 723
    `http://www.jstor.org/stable/3094609http://about.jstor.org/terms`.   724

67. Barton NH, Turelli M, Barton ” NH, Turelli3 M. Effects of Genetic Drift on Variance Components   725
    under a General Model of Epistasis. Source: Evolution INTERNATIONAL JOURNAL OF   726
    ORGANIC EVOLUTION PUBLISHED BY THE SOCIETY FOR THE STUDY OF EVOLUTION   727
    Evolution. 2004;58(5810):2111–2132. Available from: 728
    `http://www.jstor.org/stable/3449464http://about.jstor.org/terms`.   729

68. Chen X, Kuja-Halkola R, Rahman I, Arpegård J, Viktorin A, Karlsson R, et al. Dominant Genetic   730
    Variation and Missing Heritability for Human Complex Traits: Insights from Twin versus   731
    Genome-wide Common SNP Models. The American Journal of Human Genetics. 2015   732
    nov;97(5):708–714. Available from: 733
    `http://linkinghub.elsevier.com/retrieve/pii/S0002929715004061`.   734

69. Lee JJ, Chow CC. Conditions for the validity of SNP-based heritability estimation. Human genetics.   735
    2014 aug;133(8):1011–22. Available from: `http://www.ncbi.nlm.nih.gov/pubmed/24744256http:`   736
    `//www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4159725`.   737

70. Lee SH, Yang J, Chen GB, Ripke S, Stahl EA, Hultman CM, et al. Estimation of SNP heritability   738
    from dense genotype data. American journal of human genetics. 2013 dec;93(6):1151–5. Available   739
    from: `http://www.sciencedirect.com/science/article/pii/S0002929713004722`.   740

71. Chapman JM, Cooper JD, Todd JA, Clayton DG. Detecting disease associations due to linkage   741
    disequilibrium using haplotype tags: a class of tests and the determinants of statistical power. Human   742
    heredity. 2003;56(1-3):18–31. Available from: `http://www.ncbi.nlm.nih.gov/pubmed/14614235`.   743

72. Sanjak JS, Long AD, Thornton KR. Efficient Software for Multi-marker, Region-Based Analysis of   744
    GWAS Data. G3 (Bethesda, Md). 2016 jan;6(4):1023–30. Available from: 745
    `http://www.g3journal.org/content/early/2016/02/08/g3.115.026013`.   746

73. Uricchio LH, Torres R, Witte JS, Hernandez RD. Population genetic simulations of complex   747
    phenotypes with implications for rare variant association tests. Genetic epidemiology. 2015   748
    jan;39(1):35–44. Available from: `http://www.ncbi.nlm.nih.gov/pubmed/25417809`.   749

74. Orozco G, Barrett JC, Zeggini E. Synthetic associations in the context of genome-wide association   750
    scan signals. Human molecular genetics. 2010 oct;19(R2):R137–44. Available from: 751
    `http://hmg.oxfordjournals.org/content/19/R2/R137.long`.   752

75. Wray NR, Purcell SM, Visscher PM. Synthetic associations created by rare variants do not explain   753
    most GWAS results. PLoS biology. 2011;9(1):e1000579. Available from: 754
    `http://www.ncbi.nlm.nih.gov/pubmed/21267061http:`   755
    `//www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC3022526`.   756

76. Johansen CT, Wang J, Lanktree MB, Cao H, McIntyre AD, Ban MR, et al. Excess of rare variants in   757
    genes identified by genome-wide association study of hypertriglyceridemia. Nature genetics. 2010   758
    aug;42(8):684–7. Available from: `http://www.pubmedcentral.nih.gov/articlerender.fcgi?`   759
    `artid=3017369{&}tool=pmcentrez{&}rendertype=abstract`.   760

77. Kotowski IK, Pertsemlidis A, Luke A, Cooper RS, Vega GL, Cohen JC, et al. A spectrum of PCSK9   761
    alleles contributes to plasma levels of low-density lipoprotein cholesterol. American journal of human   762
    genetics. 2006;78(3):410–422.   763

78. Marini NJ, Gin J, Ziegle J, Keho KH, Ginzinger D, Gilbert Da, et al. The prevalence of   764
    folate-remedial MTHFR enzyme variants in humans. Proceedings of the National Academy of   765
    Sciences of the United States of America. 2008;105(23):8055–8060.   766

79. Romeo S, Pennacchio LA, Fu Y, Boerwinkle E, Tybjaerg-Hansen A, Hobbs HH, et al.   767
    Population-based resequencing of ANGPTL4 uncovers variations that reduce triglycerides and   768
    increase HDL. Nature genetics. 2007 apr;39(4):513–6. Available from: `http://www.pubmedcentral.`   769
    `nih.gov/articlerender.fcgi?artid=2762948{&}tool=pmcentrez{&}rendertype=abstract`.   770

80. Huyghe JR, Jackson AU, Fogarty MP, Buchkovich ML, Stančáková A, Stringham HM, et al. Exome   771
    array analysis identifies new loci and low-frequency variants influencing insulin processing and   772
    secretion. Nature genetics. 2013;45(2):197–201. Available from: 773
    `http://dx.doi.org/10.1038/ng.2507`.   774

81. Wessel J, Goodarzi MO. Low-frequency and rare exome chip variants associate with fasting glucose   775
    and type 2 diabetes susceptibility. Nature Communications. 2015;6:5897. Available from: 776
    `http://www.nature.com/doifinder/10.1038/ncomms6897`.   777

82. Purcell SM, Moran JL, Fromer M, Ruderfer D, Solovieff N, Roussos P, et al. A polygenic burden of   778
    rare disruptive mutations in schizophrenia. Nature. 2014;506(7487):185–90. Available from: 779
    `http://www.ncbi.nlm.nih.gov/pubmed/24463508`.   780

83. Nelson MR, Wegmann D, Ehm MG, Kessner D, St Jean P, Verzilli C, et al. An abundance of rare   781
    functional variants in 202 drug target genes sequenced in 14,002 people. Science (New York, NY).   782

2012 jul;337(6090):100–4. Available from: ₇₈₃
`http://www.sciencemag.org/content/337/6090/100.abstract`. ₇₈₄

84. Luciano M, Svinti V, Campbell A, Marioni RE, Hayward C, Wright AF, et al. Exome Sequencing to ₇₈₅
Detect Rare Variants Associated With General Cognitive Ability: A Pilot Study. Twin research and ₇₈₆
human genetics : the official journal of the International Society for Twin Studies. 2015 mar;p. 1–9. ₇₈₇
Available from: `http://journals.cambridge.org/abstract{_}S1832427415000109`. ₇₈₈

85. Cirulli ET, Lasseigne BN, Petrovski S, Sapp PC, Dion PA, Leblond CS, et al. Exome sequencing in ₇₈₉
amyotrophic lateral sclerosis identifies risk genes and pathways. Science. 2015 ₇₉₀
feb;347(6229):1436–1441. Available from: ₇₉₁
`http://www.sciencemag.org/content/347/6229/1436.full`. ₇₉₂

86. Thornton KR. A C++ Template Library for Efficient Forward-Time Population Genetic Simulation ₇₉₃
of Large Populations. Genetics. 2014;(2013):1–21. Available from: ₇₉₄
`http://www.ncbi.nlm.nih.gov/pubmed/24950894`. ₇₉₅

87. Kimura M. The number of heterozygous nucleotide sites maintained in a finite population due to ₇₉₆
steady flux of mutations. Genetics. 1969;61(694):893–903. ₇₉₇

88. Turelli M. Heritable genetic variation via mutation-selection balance: Lerch's zeta meets the ₇₉₈
abdominal bristle. Theoretical Population Biology. 1984 apr;25(2):138–193. Available from: ₇₉₉
`http://www.sciencedirect.com/science/article/pii/0040580984900170`. ₈₀₀

89. Slatkin M. Heritable variation and heterozygosity under a balance between mutations and stabilizing ₈₀₁
selection. Genetical research. 1987;50:53–62. ₈₀₂

90. Gravel S. When Is Selection Effective? Genetics. 2016 may;203(1):451–62. Available from: ₈₀₃
`http://www.ncbi.nlm.nih.gov/pubmed/27010021http:` ₈₀₄
`//www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC4858791`. ₈₀₅

91. Lumley T. biglm: bounded memory linear and generalized linear models; 2013. R package version ₈₀₆
0.9-1. Available from: `http://CRAN.R-project.org/package=biglm`. ₈₀₇

92. Haseman JK, Elston RC. The investigation of linkage between a quantitative trait and a marker ₈₀₈
locus. Behavior Genetics. 1972 mar;2(1):3–19. Available from: ₈₀₉
`http://link.springer.com/10.1007/BF01066731`. ₈₁₀

93. Sham PC, Purcell S. Equivalence between Haseman-Elston and variance-components linkage analyses ₈₁₁
for sib pairs. American journal of human genetics. 2001 jun;68(6):1527–32. Available from: ₈₁₂
`http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1226141{&}tool=` ₈₁₃
`pmcentrez{&}rendertype=abstract`. ₈₁₄

94. Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, et al. PLINK: a tool set for ₈₁₅
whole-genome association and population-based linkage analyses. American journal of human ₈₁₆
genetics. 2007 sep;81(3):559–75. Available from: ₈₁₇
`http://www.sciencedirect.com/science/article/pii/S0002929707613524`. ₈₁₈

95. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria; 2014.   ₈₁₉
    Available from: `http://www.R-project.org/`.   ₈₂₀

96. Neale MC, Hunter MD, Pritikin JN, Zahery M, Brick TR, Kickpatrick RM, et al. OpenMx 2.0:   ₈₂₁
    Extended structural equation and statistical modeling. Psychometrika. in press;.   ₈₂₂

97. McCarthy MI, Hirschhorn JN. Genome-wide association studies: Potential next steps on a genetic   ₈₂₃
    journey. Human Molecular Genetics. 2008;17(2):156–165.   ₈₂₄

98. Marth GT, Czabarka E, Murvai J, Sherry ST. The Allele Frequency Spectrum in Genome-Wide   ₈₂₅
    Human Variation Data Reveals Signals of Differential Demographic History in Three Large World   ₈₂₆
    Populations. Genetics. 2004;166(1):351–372.   ₈₂₇

99. Price AL, Kryukov GV, de Bakker PIW, Purcell SM, Staples J, Wei LJ, et al. Pooled association   ₈₂₈
    tests for rare variants in exon-resequencing studies. American journal of human genetics. 2010   ₈₂₉
    jun;86(6):832–8. Available from: `http://www.cell.com/article/S0002929710002077/fulltext`.   ₈₃₀

100. King CR, Rathouz PJ, Nicolae DL. An evolutionary framework for association testing in   ₈₃₁
    resequencing studies. PLoS genetics. 2010 nov;6(11):e1001202. Available from: ₈₃₂
    `http://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1001202`.   ₈₃₃

101. Browning SR, Thompson EA. Detecting rare variant associations by identity-by-descent mapping in   ₈₃₄
    case-control studies. Genetics. 2012 apr;190(4):1521–31. Available from: ₈₃₅
    `http://www.genetics.org/content/early/2012/01/18/genetics.111.136937`.   ₈₃₆

102. North TL, Beaumont MA. Complex trait architecture: the pleiotropic model revisited. Scientific   ₈₃₇
    reports. 2015 jan;5:9351. Available from: ₈₃₈
    `http://www.nature.com/srep/2015/150320/srep09351/full/srep09351.html`.   ₈₃₉

103. Moutsianas L, Agarwala V, Fuchsberger C, Flannick J, Rivas MA, Gaulton KJ, et al. The power of   ₈₄₀
    gene-based rare variant methods to detect disease-associated variation and test hypotheses about   ₈₄₁
    complex disease. PLoS genetics. 2015 apr;11(4):e1005165. Available from: ₈₄₂
    `http://journals.plos.org/plosgenetics/article?id=10.1371/journal.pgen.1005165`.   ₈₄₃

104. Keightley PD, Hill WG. Variation Maintained in Quantitative Traits with Mutation-Selection   ₈₄₄
    Balance: Pleiotropic Side-Effects on Fitness Traits. Proceedings of the Royal Society B: Biological   ₈₄₅
    Sciences. 1990 nov;242(1304):95–100. Available from: ₈₄₆
    `http://rspb.royalsocietypublishing.org/content/242/1304/95`.   ₈₄₇

105. Haldane JBS. The measurement of natural selection. Proc 9th Int Congr Genet. 1954;1:480–487.   ₈₄₈

106. Latter BDH. Natural selection for an intermediate optimum. Australian Journal of Biological   ₈₄₉
    Sciences. 1960;13:30–35. Available from: `http://www.cabdirect.org/abstracts/19600101674.`   ₈₅₀
    `html;jsessionid=3B1562EB69031455E4C7BB05F3D8B699?freeview=true`.   ₈₅₁

107. Crow JF, Kimura M. The theory of genetic loads. Proc XI Int Congr Genet. 1964;p. 495–505.   ₈₅₂

108. Kingman JFC. A Simple Model for the Balance between Selection and Mutation. Journal of Applied Probability. 1978 mar;15(1):1–12. Available from: `http://www.jstor.org/stable/3213231`.

109. Fleming WH. Equilibrium Distributions of Continuous Polygenic Traits. SIAM Journal on Applied Mathematics. 1979 jul;Available from: `http://epubs.siam.org/doi/abs/10.1137/0136014`.

110. Kimura M. A stochastic model concerning the maintenance of genetic variability in quantitative characters. Proceedings of the National Academy of Sciences of the United States of America. 1965 sep;54(3):731–6. Available from: `http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=219735{&}tool=pmcentrez{&}rendertype=abstract`.

111. Burger R, Hofbauer J. Mutation load and mutation-selection-balance in quantitative genetic traits. Journal of Mathematical Biology. 1994 feb;32(3):193–218. Available from: `http://link.springer.com/10.1007/BF00163878`.

112. Lohmueller KE, Indap AR, Schmidt S, Boyko AR, Hernandez RD, Hubisz MJ, et al. Proportionally more deleterious genetic variation in European than in African populations. Nature. 2008;451(7181):994–997.

113. Fu W, Gittelman R, Bamshad M, Akey J. Characteristics of Neutral and Deleterious Protein-Coding Variation among Individuals and Populations. The American Journal of Human Genetics. 2014;95(4):421–436.

114. Do R, Balick D, Li H, Adzhubei I, Sunyaev S, Reich D. No evidence that selection has been less effective at removing deleterious mutations in Europeans than in Africans. Nature Genetics. 2015 jan;47(2):126–131. Available from: `http://www.nature.com/doifinder/10.1038/ng.3186`.

115. Peischl S, Excoffier L. Expansion load: recessive mutations and the role of standing genetic variation. Molecular Ecology. 2015 may;24(9):2084–2094. Available from: `http://doi.wiley.com/10.1111/mec.13154`.

116. Henn BM, Botigué LR, Peischl S, Dupanloup I, Lipatov M, Maples BK, et al. Distance from sub-Saharan Africa predicts mutational load in diverse human genomes. Proceedings of the National Academy of Sciences. 2016 jan;113(4):E440–E449. Available from: `http://www.pnas.org/lookup/doi/10.1073/pnas.1510805112`.

Figures

Parameter	Description
N	Population size
V_S	The total inverse selection intensity
V_E	Variance of environmental gaussian deviates
λ	Mean and standard deviation of trait effects
h	Dominance of trait effects
G	Genetic contribution to phenotype
E	Environmental contribution to phenotype
P	Phenotype
w	Fitness, based on gaussian function

Table 1. Description of parameters used in the models.

Fig 1. Variance explained over allele frequency. The cumulative additive and dominance genetic variance which can be explained by markers whose frequencies, q, are $\leq x$. Each color represents a different value of λ: the mean effects size of a new deleterious mutation. Shown here are the gene-based (GBR), additive co-dominant (AC), incomplete multiplicative recessive (Mult. recessive ($h = 0.25$); iMR) and complete multiplicative recessive (Mult. recessive ($h = 0$);cMR) models. Solid lines show the additive variance alone and dotted lines show the combined additive and dominance variance. All data shown are for models where $H^2 \sim 0.08$, although it was confirmed (data not shown) that these particular results are independent of total H^2. The additive and dominance genetic variance is estimated by the adjusted r^2 of the regression of all markers (and their corresponding dominance encoding) with $q \leq x$ onto total genotypic value (see methods for details); data are displayed as the mean of 250 simulation replicates. The vertical dotted and dashed lines correspond to the $q = 0.001$ and $q = 0.05$, respectively. The curves under no growth appear to be truncated with respect to rapid growth because the range of the x-axis differs between growth and no growth (minimum $q = 1/2N$).

Fig 2. Heritability estimates compared to population heritability. Heritability estimates and population heritability as a function of λ: the mean effect size of a new deleterious mutation. Additive (A; orange) component of true heritability is calculated by multiplying the end point($q = 1$) of the variance curves in Fig 1 by the broad-sense heritability values summarized in S1 Fig. HE-regression and GREMLd estimates were obtained from random population samples (n = 6000). GREMLd analysis was performed in GCTA using genotype data that was either unfiltered or filtered to remove variants with MAF<0.01. Twin study estimates are directly calculated using MZ and DZ twin correlations from 64 sets of twin studies. Each study consisted of pooling 2000 MZ twin pairs and 2000 DZ twin pairs from each of 8 model replicates for a total of 64,000 individual phenotypes. Data are plotted as the median across replicate sets ± half the interquartile range. Shown are the additive co-dominant (AC), gene-based (GBR) incomplete multiplicative recessive (Mult. recessive ($h = 0.25$); iMR) and complete multiplicative recessive (Mult. recessive ($h = 0$); cMR) models.

Fig 3. Power of association tests. (A) The power of a single marker logistic regression, at significance threshold of $\alpha \leq 10^{-8}$, as a function of λ: the mean effect size of a new deleterious mutation. For single marker tests we define power as the number of simulation replicates in which any single marker reaches genome wide significance. Two study designs were emulated. For the gene chip design only markers with $MAF > 0.05$ were considered and all markers were considered for the resequencing design. Genetic models shown here are the additive co-dominant (AC), gene-based (GBR), complete multiplicative site-based recessive (Mult. recessive ($h = 0$); cMR) and incomplete multiplicative site-based recessive models (Mult. recessive ($h = 0.25$); iMR) (B) The power of region-based rare variant association tests to detect association with the simulated causal gene region at significance threshold of $\alpha \leq 10^{-6}$. For region-based tests, we define power as the percent of simulation replicates in which the p-value of the test was less than α. The p-values for the ESM, c-Alpha were evaluated using 2×10^6 permutations. SKAT p-values were determined by the SKAT R package and represent numerical approximations to the presumed analytical p-value.

Fig 4. Distribution of significant GWAS hits. Horizontal violin plots depict the distribution of minor allele frequencies (MAF) of the most strongly associated single marker in a GWAS. Individual hits are plotted as translucent points and jittered to provide a sense of the total number and density of hits. Each panel contains simulated data pooled across model replicates for each value of λ with empirical data for comparison. Empirical data are described in Materials and Methods. In cases where more than one marker was tied for the lowest p-value, one was chosen at random. Shown here are the additive co-dominant (AC), gene-based (GBR), incomplete multiplicative recessive (Mult. recessive ($h = 0.25$); iMR) and complete multiplicative recessive (Mult. recessive ($h = 0$);cMR) models. All data shown are for models where $H^2 \sim 0.08$, because single marker test power was too low under $H^2 \sim 0.04$ to make informative density plots. Simulated data were subjected to ascertainment sampling such that the MAF distribution of all markers on the simulated genotyping chip was uniform. Specific information regarding the empirical data can be obtained in S1 Table.