# $C_4$ photosynthesis and climate through the lens of optimality

Haoran Zhou[*][†], Brent R. Helliker[*] and Erol Akçay[*]

[*]Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA

[†]*Correspondence:* Haoran Zhou

*Phone: 1-215-808-7042*

*Email: haoranzh@sas.upenn.edu*

Brent R. Helliker*: helliker@sas.upenn.edu*

Erol Akçay: *eakcay@sas.upenn.edu*

***Statement of authorship****:* HZ, BH and EA conceptualized the study. HZ and EA built the model,

HZ and BH put the idea in a general evolutionary context, HZ performed the modeling work and

analyzed output data. HZ wrote the first draft, BH and EA contributed substantially to revisions.

***Runing title:*** Selective pressures in $C_4$ grasses evolution

***Conflict of interest:*** None declared.

***The type of article:*** Regular research articles (research report)

***Classification:*** biological science (major), evolution/ecology/plant biology (minor)

**Word count limitation:** Five and a half page (1 column for Fig. 1, 2, 3, 5 and 1.5 column for
Fig.4) using Article Sizing Tool.

**Abstract**

$CO_2$, temperature, water availability and light intensity were all potential selective pressures to

propel the initial evolution and global expansion of $C_4$ photosynthesis over the last 30 million

years. To tease apart how the primary selective pressures varied along this evolutionary

trajectory, we coupled photosynthesis and hydraulics models while optimizing photosynthesis

over stomatal resistance and leaf/fine-root allocation. We further examined the importance of

resource (e.g. nitrogen) reallocation from the dark to the light reactions during and after the

initial formation of $C_4$ syndrome. We show here that the primary selective pressures— all acting

upon photorespiration in $C_3$ progenitors— changed through the course of $C_4$ evolution. The

higher stomatal resistance and leaf-to-root allocation ratio enabled by the $C_4$

carbon-concentrating mechanism led to a $C_4$ advantage without any change in hydraulic

properties, but selection on nitrogen reallocation varied. Water limitation was the primary driver

for the initial evolution of $C_4$ 25-32 million years ago, and could positively select for $C_4$

evolution with atmospheric $CO_2$ as high as 600 ppm. Under these high $CO_2$ conditions, nitrogen

reallocation was necessary. Low $CO_2$ and light intensity, but not nitrogen reallocation, were the

primary drivers during the global radiation of $C_4$ 5-10 MYA. Finally, our results suggest that

identifying the predominate selective pressures at the time $C_4$ first evolved within a lineage

should help explain current biogeographical distributions.

*Keywords* $C_4$ evolution, optimal stomatal conductance, resource allocation, water limitation,

selective pressure, dark/light reaction

**Significance Statement**

$C_4$ photosynthesis pathway had evolved more than 60 times independently across the terrestrial

plants through mid-Oligocene (~30 MYA) and diversified at late Miocene (5 to 10 MYA). We

use an optimal physiology model to examine the primary selective pressures along the

evolutionary history. Water limitation was the primary driver for $C_4$ evolution from the initial

evolutionary events 25 -32 MYA until $CO_2$ became low enough to, along with light intensity,

drive the global radiation of $C_4$ 5-10 MYA. This modeling framework can be used to investigate

evolution of other physiological traits (e.g. N reallocation, hydraulics) after the initial formation

of $C_4$ syndrome, which contributed to further increasing productivity of $C_4$ in historical and

current environmental conditions.

# 1 **Introduction**

The evolution of the $C_4$ photosynthetic pathway enabled the concentration of $CO_2$ around

Rubisco, the enzyme responsible for the first major step of carbon fixation in the $C_3$

photosynthetic pathway, thus eliminating photorespiration. $C_3$ photosynthesis is present in all

plants, and within $C_4$ plants the $C_3$ pathway is typically ensconced within specialized bundle

sheath cells that surround leaf veins. $CO_2$ that diffuses into a leaf is shuttled from adjacent

mesophyll cells to the bundle sheath via a four-carbon pump, the energetic cost of which is

remunerated by ATP derived from the light reactions (1, 2). As a whole, the $C_4$ pathway

eliminates photorespiration, a process that can dramatically reduce photosynthesis and begins

with the assimilation of $O_2$, instead of $CO_2$, by Rubisco. Over the last 30 million years, the

reduction in $C_3$ photosynthesis by photorespiration was large and broad enough to select for the

independent evolution of the $C_4$ pathway more than 60 times across the terrestrial plants (3). The

diversity of plant families with $C_4$ is greatest in the eudicots (1200 species) and the Poaceae, the

monocot family containing the grasses (4500 species) (2), account for nearly 25% of terrestrial

plant productivity and several important agricultural species (4).

While increased photorespiration in $C_3$ progenitors was central to the evolution of the $C_4$ carbon

concentrating mechanism (CCM), the relative importance of different environmental drivers of

the photorespiratory increase has been the subject of much debate (5, 6, 7, 8). Lower $CO_2$ leads

to higher rates of photorespiration, as does higher temperature. Past physiological models

therefore focused on examining temperature and $CO_2$ concentration as selective pressures for $C_4$

evolution and expansion (5, 9, 10). Under warmer temperatures and low $CO_2$, $C_4$ photosynthesis

has greater carbon gain than $C_3$, but under cooler temperatures and high $CO_2$, the metabolic costs

of the $C_4$ pathway and lower photorespiration in $C_3$ leads to greater carbon gain in $C_3$.

Alternatively, water availability has been proposed as the impetus for $C_4$ evolution in eudicots

(2), and recent phylogenetic analyses have suggested the same in grasses (7, 13). Water

availability should have an impact on $C_4$ evolution that could work independently or in concert

with changes in $CO_2$ and temperature. First, water deficits indirectly increase photorespiration in

$C_3$ plants by forcing stomatal closure to reduce leaf water loss; consequently decreasing the flux

of $CO_2$ into the leaf and the availability of $CO_2$ for Rubisco. Second, the $C_4$ carbon concentrating

mechanism allows for the maintenance of lower stomatal conductance, and therefore lower water

loss, for a given assimilation rate; leading to a higher water-use-efficiency (WUE) than $C_3$ (11,

12).

The different environmental drivers of the photorespiratory increase in $C_3$ plants—atmospheric

$CO_2$ concentration, temperature and water availability— have changed dramatically as $C_4$

photosynthesis has evolved over the last 30 million years. Atmospheric $CO_2$ decreased

monotonically from the mid-Oligocene ($680 \pm 200$ ppm) to the early Miocene ($357 \pm 108$ ppm)

down to the Pleistocene minima, where $CO_2$ oscillated between approximately 180 and 280 ppm

through glacial/interglacial cycles (14). Physiological models that focused on temperature and

$CO_2$ implied that $C_4$ evolved, in both grasses and eudicots, at the low end of this $CO_2$ range in

the mid- Miocene to the Pleistocene (2, 5, 9, 10, 15). $C_4$ grasses did become a major component

of grassland biomes— in terms of biomass, $C_4$ lineage diversity, or herbivore dietary

components— in the late Miocene (5 to 10 MYA), but molecular evidence suggests that $C_4$

photosynthesis arose in the grasses in the mid-Oligocene (~30 MYA) (16). Similarly,

phylogenetic reconstructions provide evidence that eudicots of the Chenopodiaceae evolved $C_4$

as early as the monocots, but saw the greatest rate of $C_4$ evolution and diversification in the late

Miocene (17, 18, 19). Along with $CO_2$, precipitation declined over the period of $C_4$ evolution and

diversification, leading to vast terrestrial areas where low or highly seasonal precipitation inputs

led to the loss of forests and consequently, the evolution of the world's first grasslands (20, 21).

The spread of grasslands indicate a habitat change with larger surface radiation loads, higher

surface temperatures and increased potential for plant water loss (6, 22). Therefore, the early

evolution of $C_4$ suggested by molecular phylogenies indicates that water availability played an

important role for both $C_4$ grasses and eudicots while $CO_2$ was still relatively high (6, 18, 23, 24,

22, 25). The potentially interacting roles of water availability, changes in radiation and $CO_2$

along the evolutionary trajectory of $C_4$ photosynthesis have not been fully investigated within a

comprehensive physiological model.

A related but largely unstudied evolutionary change during the divergence of $C_4$ photosynthesis

from $C_3$ is the allocation of nutrients/resources (e.g. N considering enzymes and proteins)

between the dark reactions and the light reactions. $C_4$ plants might allocate a greater proportion

of N to light reactions than to dark reactions as compared to $C_3$ because of the extra ATP costs of

the CCM (26, 27). We propose that the reallocation of N between dark and light reactions

provides a further advantage for $C_4$ above the CCM alone, and that different environmental

conditions can select for a shift in the degree of reallocation both through evolutionary time and

across species in extant plants.

Changes in the environmental controls on photorespiration suggest that multiple environmental

drivers interacted to differing degrees along the trajectory of $C_4$ evolution. Our goal here is to

tease apart the selective pressures that led to the evolution of $C_4$ photosynthesis initially and the

global expansion 5-10 MYA through to the current day. We use the framework of an optimality

model in which the plant makes allocation "decisions" in order to maximize photosynthetic

assimilation rate. Our approach advances our understanding of $C_4$ evolution in four important

ways. First, we revisit the temperature-$CO_2$ crossover approach and integrate the effects of water

limitation, light, optimal allocation decisions, and the interactions between these in a single

model. Second, the hypothesis that $C_4$ photosynthesis has a higher WUE than $C_3$ implicitly relies

on an optimality argument to balance carbon gain and water loss (28), yet the role of optimal

stomatal conductance in mediating selective pressures due to water limitation during the

evolution of $C_4$ plants remains largely unexplored (but see 15). Most previous models assume *a*

*priori* that $C_4$ grasses have lower stomatal conductance. Instead, we let both stomatal

conductance and leaf/fine-root allocation emerge endogenously from the model. Third, we use

the $C_4$ model including cost of the $C_4$ pathway in the light reactions (2 additional ATP per $CO_2$

fixed; 1, 29), which previous modeling analysis did not explicitly consider (9, 22, 30, 31).

Finally, we consider reallocation of nitrogen from the dark reactions to the light reactions, which

can change the tradeoffs between photosynthesis and water use by $C_4$ grasses.

**Results**

Assimilation-based crossover temperatures, defined as the temperature at which assimilation by

the $C_4$ pathway exceeds that of the $C_3$ pathway, decrease as water limitation increases and light

intensity increases across all $CO_2$ concentrations (Fig. 1, Fig. S1). Without water stress (solid

black line in Fig 1), our model predicts a $C_3/C_4$ crossover temperature of 23°C under 380 ppm; a

result similar to previous data and/or models that did not explicitly account for water stress (9,

10, 32). The model results in Fig. 1 were all under saturated light and with a $C_4$ $J_{max}/V_{cmax}$ ratio of

4.5, which corresponds to a full reallocation of nitrogen from light to dark reactions. Model

results for a $C_4$ $J_{max}/V_{cmax}$ ratio of 2.1 (corresponding to no reallocation) were similar (Fig. S1a)

with the primary exception being that under low $CO_2$ and low water availability (e.g. $CO_2$=300

ppm, VPD = 3 kPa and $\Psi_S$ = -1.5 MPa or all $CO_2$ concentrations with higher VPD and lower

$\Psi_S$), crossover temperatures are higher with $J_{max}/V_{cmax}$= 4.5, showing that nitrogen reallocation

decreases the $C_4$ advantage under water limitation and low $CO_2$. Under saturated soil and low

VPD, crossover temperatures decrease along with increasing light intensity (Fig. S1c, d). An

increase in light intensity provides a larger relative benefit for $C_4$ at low $CO_2$, because $C_3$

photosynthesis remains $CO_2$ limited throughout while $C_4$ light limitations lessen as light

increases.

Fig. 1. Crossover temperatures of photosynthesis for $C_3$ and $C_4$ with the change of $CO_2$ concentration

under different water conditions. Light intensity was 1400 μmol photons $m^{-2}s^{-1}$ for all model runs.

$J_{max}/V_{cmax}$=2.1 for $C_3$ and $J_{max}/V_{cmax}$=4.5 for $C_4$. Solid black line: VPD=0.15kPa, $\Psi_S$=0 MPa; dashed black

line: VPD=1kPa, $\Psi_S$=-0.5 MPa; dot-dashed black line: VPD=2kPa, $\Psi_S$=-1 MPa; dotted black line:

VPD=3 kPa, $\Psi_S$=-1.5 MPa.

While crossover temperatures allow for a clear diagnostic of comparative assimilation, they do

not demonstrate the degree of $C_4$ photosynthetic advantage. To this end, we calculated the net

assimilation rate difference between $C_4$ and $C_3$, $\Delta A_n$ (net assimilation of $C_4$ minus that of $C_3$), to

comprehensively examine the whole suite of environmental conditions (Fig. 2, 3). The positive

contour space ($\Delta A_n > 0$) means that $C_4$ outcompetes $C_3$ within given environmental dimensions,

and the higher the $\Delta A_n$, the greater the advantage of $C_4$. Under a $CO_2$ concentration of 200 ppm

and saturated light, $\Delta A_n$ is higher under moist conditions than water-limited conditions (Fig. 2a,

b). In contrast, under higher $CO_2$ concentrations (400 and 600 ppm), $C_4$ has the greatest

advantage only in water-limited conditions, which leaves a relatively small environmental

envelope for $C_4$ to evolve (areas where $\Delta A_n > 0$ in Fig. 2c-f). This result is due to the fact that $C_3$

photosynthesis has a greater proportional increase in assimilation from 200 to 400 and 600 ppm

$CO_2$. Across all scenarios, increasing $J_{max}/V_{cmax}$ increases both the $\Delta A_n$ and space for $C_4$

advantage (Fig. 2 b, d, f). At 200 ppm and saturated soils, $\Delta A_n$ is highest under saturated light,

and decreases as light intensity decreases (Fig. 3a, b). At 400 ppm $CO_2$ and higher, $\Delta A_n$

indicated a $C_4$ advantage ($\Delta A_n > 0$) only when light intensities were above 1000 mmol m-2 s-1,

temperature was above 30 °C and $J_{max}/V_{cmax}$ was high (Fig. 3 c, d). Finally, we calculated the

photosynthesis rates of the two pathways under conditions often encountered in today's

grasslands to look at the effect of nitrogen reallocation (Fig. 4). With $J_{max}/V_{cmax}$ =2.1 for both $C_3$

(solid black line) and $C_4$ (dashed line), the $C_4$ assimilation rate is rarely higher than $C_3$, which

indicates $C_4$ does not have an obvious advantage under current $CO_2$ from saturated soils down to

VPD = 2 kPa and $\Psi_S$ = -1 MPa. However, with $J_{max}/V_{cmax}$ =4.5 for $C_4$ (dotted line), $C_4$ does have

an advantage over $C_3$ at temperatures above 25 $^o$C.

No reallocation

Reallocation

200 ppm

400 ppm

600 ppm

Water limitation intensity

Water limitation intensity

$\Delta An$

−5.0    −2.5    0    2.5    5.0

2    Fig. 2. The total difference in $CO_2$ assimilation between $C_4$ and $C_3$ ($A_n(C_4)$-$A_n(C_3)$) under various $CO_2$

1    (200 ppm, 400 ppm and 600 ppm) and water conditions under saturated light intensity (1400 μmol

2    photons m$^{-2}$s$^{-1}$). $J_{max}/V_{cmax}$=2.1 for C$_3$ and C$_4$ (a, c, e) and $J_{max}/V_{cmax}$=2.1 for C$_3$ and $J_{max}/V_{cmax}$=4.5 for C$_4$

3    (b, d, f). Water limitation intensity is: 1: VPD =0.15 kPa, $\Psi_S$=0 MPa; 2: 1.5 kPa, -0.5MPa; 3: 2kPa, -1

4    MPa; 4: 3kPa, -1.5 MPa; 5: 4kPa, -2 MPa.

Fig. 3. The total difference in $CO_2$ assimilation between $C_4$ and $C_3$ ($A_n(C_4)$-$A_n(C_3)$) with $J_{max}/V_{cmax}$=2.1 for

$C_3$ and $C_4$ under various $CO_2$ (200 ppm, 400 ppm) and different light intensities (from 200 to 1400 μmol

photons m$^{-2}$s$^{-1}$) with saturated water condition (VPD=0.15kPa, $\Psi_S$=0 MPa) (a, c) and with $J_{max}/V_{cmax}$=2.1

for $C_3$ and $J_{max}/V_{cmax}$=4.5 for $C_4$ (b, d).

Fig. 4. Assimilation rates of $C_3$ with $J_{max}/V_{cmax}$=2.1 (solid black line), C4 with $J_{max}/V_{cmax}$=2.1 (dashed

black line) and $C_4$ with $J_{max}/V_{cmax}$=4.5 (dotted black line) under light intensity of 1400 μmol photons

m$^{-2}$s$^{-1}$, $CO_2$ of 400 ppm and different water limitation conditions. (a) saturated soils; (b) VPD=1kPa and

$\Psi_S$=-0.5 MPa; (c) VPD=2kPa and $\Psi_S$=-1 MPa.

Under all environmental and nitrogen allocation scenarios, optimal stomatal resistance ($r_s$) and

leaf biomass/total biomass allocation (f) are higher in $C_4$ plants than $C_3$ plants. The general

pattern of response was similar across $CO_2$ concentrations, so only 400 ppm is presented with

both highest and lowest water availability (Fig. 5 and Fig. S2). Optimal f and $r_s$ for $C_3$ is always

lower than that for $C_4$ under different water availability and $CO_2$ (Fig. 5). In addition, f decreases

and $r_s$ *increases* as intensity of water limitation increases. Results are the consistent for $C_4$ with a

$J_{max}/V_{cmax}$ of 2.1 and $J_{max}/V_{cmax}$ of 4.5. The jumps in $r_s$ and f in Fig. 5 correspond to the transition

from RuBP carboxylation limited assimilation to RuBP regeneration limited assimilation ($A_j$).

The transition temperature decreases as $CO_2$ increases. In the Fig. S3, the various limitation

states are plotted together under multiple environmental scenarios, using both $J_{max}/V_{cmax}$=2.1 and

4.5 for $C_4$. With $J_{max}/V_{cmax}$=2.1, $C_4$ is light limited in most of the environmental conditions. With

$J_{max}/V_{cmax}$=4.5, C$_4$ starts to be limited by CO$_2$ under low temperatures and to be limited by light

under high temperatures.

Fig. 5. Stomatal resistance (r$_s$) and leaf/fine-root allocation (f) as a function of temperature, with

$J_{max}/V_{cmax}$=2.1 for both C$_3$ and C$_4$ with saturated light under CO$_2$ of 400 ppm and different water

conditions. Solid black line: C$_3$ with VPD=0.15kPa, $\Psi_S$=0 MPa; dashed black line: C$_4$ with

VPD=0.15kPa, $\Psi_S$=0; solid grey line: C$_3$ with VPD=4 kPa, $\Psi_S$=-2 MPa; dashed grey line: C$_4$ with VPD=4

kPa, $\Psi_S$=-2 MPa. Vertical lines indicate transition from RuBP carboxylation limited condition to RuBP

regeneration limited condition for C$_3$ and C$_4$.

**Discussion**

Our modeling results imply that the environmental selection for C$_4$ evolution was in place during

the mid-Oligocene at warm, arid sites as water limitation acted as the primary selective pressure

to increase photorespiration when CO$_2$ is above 400 ppm, and even up to 600 ppm. Under

saturated water conditions, there was little room for C$_4$ to evolve 20-30 MYA as CO$_2$ was likely

above 400 ppm (14, 33), and the ATP costs of the C$_4$ mechanism is too high and photorespiration

in C$_3$ plants too low. As water becomes more limited, however, the C$_4$ advantage becomes

increasingly larger. Enhanced carbon gain under water limited conditions has been believed to be

the selective force behind the evolution of $C_4$ in dicotyledonous plants (2) and more recently, in

accordance with phylogenetic evidence, in grasses (7, 16). Molecular-based evidence supports a

mid-Oligocene evolution of $C_4$ in some grass and eudicot lineages, and our results suggest the

same (8, 18, 34). Lineages that evolved in $C_4$ during this period may have been predisposed to

exist as arid-land, saline, or disturbed-site specialists as we see today in most $C_4$ eudicots and the

earliest known grass subfamily in which $C_4$ evolved, the Chloridoideae (8, 35).

As $CO_2$ decreased through the Miocene, warm temperatures remained a strong selective force,

but the main selective force for $C_4$ evolution shifted from water limitation to low $CO_2$ and, to a

lesser extent, light intensity. Since increased light intensity alone could not lead to advantage of

$C_4$ under high $CO_2$ (Fig. 3c), it seems likely that $C_4$ grasses could not dominate open grasslands,

except in locally arid areas, while $CO_2$ was still high. So between the initial evolutionary events

leading to the emergence of $C_4$ and the large-scale expansion within the grasses 5-10 MYA, $C_4$

grass radiation likely idled in small pockets of selective favorability as $CO_2$ concentrations

declined through the Miocene (2, 8). As $CO_2$ declined, the high light levels inherent to grassland

systems gave $C_4$ photosynthesis an increasing advantage, leading to broader geographic and

evolutionary radiation. Our results are consistent with previous studies showing that low $CO_2$

(200-300 ppm) strongly favors $C_4$ over $C_3$ photosynthesis (e.g. 9, 15). And we further show that

low $CO_2$ provides a clear $C_4$ advantage under a large range of water availability and light

intensity regimes. The greatest $C_4$ advantage occurs, however, in relatively moist and mildly

water-limited conditions; opposite to that which is seen under high $CO_2$ (Fig. 2c). The

environmental conditions that lead to the largest $C_4$ advantage within our model therefore

perfectly parallels those documented in the 20[th] century, $C_4$-dominated grasslands: highly

seasonal precipitation that occurs chiefly within a warm growing season (36, 37, 38). These are

also similar to the conditions that led to the large-scale expansion of $C_4$ grasslands in the

Miocene, for example the onset of summer monsoons and subsequent $C_4$ grassland expansion in

the Indian subcontinent (39).

The potential role of water limitation to play a central role $C_4$ grass evolution has sparked interest

in grass hydraulics and the anatomical shifts in $C_3$ grasses that were prerequisites to $C_4$ evolution

(24, 22, 40). The modeling effort of Osborne and Sack (22) suggests a hydrological underpinning

to the evolution of $C_4$ grasses, but found a much smaller environmental window for $C_4$ evolution

than we did. At 400 ppm $CO_2$ and soil water potential of -1 MPa— a common occurrence in

grasslands (41)— they showed that $C_4$ hydraulic conductance must be twice that of $C_3$ grasses

for $C_4$ grasses to achieve slightly greater carbon uptake. In contrast, we find a clear $C_4$ advantage

under these, and even drier, conditions by allowing for optimal solutions of stomatal

conductance and leaf/fine-root ratio to maximize photosynthesis. Plant hydraulic conductance

was kept equal across $C_3$ and $C_4$ throughout simulations, and increasing hydraulic conductance

had little impact on our major results and conclusion (Fig. S4, S5), the implication being that the

$C_4$ pathway itself is enough to result in greater carbon gain under water stress without any

required increase in hydraulic conductance. These results do not necessarily contradict the idea

that larger bundle sheaths and smaller interveinal distance— which were clear prerequisites for

$C_4$ evolution (24, 42) — led to greater hydraulic conductance and drought tolerance among $C_3$

grass progenitors (24), but they do suggest that greater hydraulic conductance is not necessary to

give $C_4$ plants an advantage once the carbon-concentrating mechanism evolved. In fact, we

hypothesize that once $C_4$ evolves in a lineage, selection on increased hydraulic conductance

would not only lessen, but invert, leading to the development of even narrower xylem conduits

and greater drought resistance. There is some empirical support for such a prediction in eudicots

(43).

We assumed that during the early evolution of $C_4$, both $C_3$ and $C_4$ plants had a similar balance of

nitrogen across the light and dark reactions, and that the allocation of nitrogen could be treated

separately from the evolution of the $C_4$ CCM as a target of selection. We propose that different

environmental conditions can select for a shift in the degree of reallocation (assessed here by a

change in $J_{max}/V_{cmax}$) both through evolutionary time and across species in extant plants. In

general, CCMs allow for less investment in nitrogen-rich Rubisco (44), and the nitrogen not used

for Rubisco could be either reinvested in light harvesting machinery, or simply not used at all;

thus reducing the plant nitrogen requirement. Modeling studies have long assumed a high

$J_{max}/V_{cmax}$ for $C_4$ photosynthesis (22, 31, 45) and measurements show lower Rubisco content and

higher chlorophyll and thylakoid content, giving evidence of reallocation in extant $C_4$ species

(26, 27, 46). Empirical estimates of $J_{max}/V_{cmax}$, in $C_4$ plants paint a more variable picture, ranging

from 2 to above 6, with a mean of around 4.5 (47, 48, 49, 50, 51, 52), which is higher than the

mean $J_{max}/V_{cmax}$ estimates for $C_3$ plants of 2.1 (53). Increasing $J_{max}/V_{cmax}$ almost always increases

the photosynthesis rate of $C_4$ grasses (Fig. 4, Fig. S6), and therefore could lead to a competitive

advantage over $C_3$ grasses as well as $C_4$ grasses that do not reallocate. Assuming there is little

cost or no genetic constraints for reallocation, the selection pressure to reallocate would have

been strongest when $CO_2$ was high, e.g. during the initial evolutionary events in the

Oligocene/Miocene, when the CCM alone does not give $C_4$ a large advantage (Fig. 2 c, e, Fig. 3

c). When $CO_2$ was low during the $C_4$ radiation 5-10 MYA, however, the CCM alone would give

$C_4$ an advantage and reallocation would not change the competitive balance between $C_3$ and $C_4$

(Fig. 2 a and Fig. 3a). As $CO_2$ remained low through to the Pleistocene, selection for nitrogen

reallocation to the light reactions would lessen further, especially during the $CO_2$ minima of the

Pleistocene glacial periods (~ 180 ppm). In this context, an interesting question is how the

evolutionary picture of $J_{max}$ and $V_{cmax}$ allocation was coordinated with the formation of $C_4$

pathway in response to the high $CO_2$ in Oligocene, to the $CO_2$ decrease through Pleistocene and,

further, to the increase of $CO_2$ in the last 150 years.

$C_4$ photosynthesis first evolved 25 – 32 MYA, and many subsequent and independent

evolutionary origins occurred through to the Pleistocene 2.8 MYA. Each evolutionary origin

represents both different selective pressures and taxonomic (genetic) constraints as climate and

$CO_2$ changed. Taking the Chloridoideae as an example, we can use our model to develop

hypotheses along the evolutionary trajectory of $C_4$ in this grass subfamily. The initial evolution

of $C_4$ photosynthesis $25 - 32$ MYA while $CO_2$ was high was driven by aridity, acting to decrease

stomatal conductance that increased photorespiration in $C_3$ progenitors initially, and led to higher

water use efficiency upon the evolution of the CCM. There would have been enough of a

reduction in water use that selection for increased hydraulic conductance would relax, allowing

for the development of more resilient— and less conductive— xylem. Also at this point, there

would have been strong selection for reallocation of nitrogen from the dark reactions to the light

reactions. The large radiation of $C_4$ within the Chloridoideae that occurred $5 - 10$ MYA was

likely driven by low $CO_2$ and high light, and the previously-evolved hydraulic resilience would

lead to this subfamily becoming dry-site specialists observed in current-day distributions (35).

There would have been much less selective pressure to reallocate N during the large radiation,

but such a reorganization was likely already in place within the clade. In contrast, for the

lineages that first evolved $C_4$ in the late Miocene (e.g. *Stipagrostis*, *Eriachne*, *Neurachne*), $CO_2$

would have been the primary impetus for $C_4$ evolution, but for these lineages there would have

been little selection to reallocate nitrogen until the dawn of the industrial revolution. We would

also expect these more recently evolved lineages to have greater hydraulic conductance than

those of the Chloridoideae. By optimizing carbon gain over water loss, we developed a plausible

physiological explanation for the early evolution of $C_4$ and further proposed hypotheses about

how the variety of traits that comprise the $C_4$ syndrome developed in concert with the climate

changes that occurred through the evolutionary trajectory (54). By selecting extant species within

select lineages, these hypotheses can be examined empirically, ultimately providing an

integrative view of the selection pressures that led to the current physiologies and distribution of

$C_4$ plants.

**Materials and Methods**

*Overview of the model*

We used different modeling scenarios to examine the advantage of $C_4$ photosynthesis for the

initial origin, expansion and current distribution. Initially, we assume that the CCM is the only

difference between $C_3$ and $C_4$. This comparison corresponds to two closely related species whose

other traits have not had time to diverge in response to differential selection pressures. We then

examine shifts in N allocation between the light and dark reactions of $C_4$, which may have

happened in subsequent divergence of $C_3$ and $C_4$ after the CCM evolved.

The soil-plant-air water continuum was incorporated in $C_3$ photosynthesis models (55) and $C_4$

models (29) to examine interactions of $CO_2$, water availability, light and temperature. We used

the optimality approach of Givnish (1986) (56), where $C_3$ and $C_4$ plants optimize stomatal

resistance and leaf/fine-root allocation to balance carbon gain and water loss. A full model

description is in SI in supporting information with Table S1 for parameter abbreviation and

Table S2 for input parameters. The model derivation using Mathematica (Wolfram Research,

Inc.) and methods for numerical solutions are in SII.

*Photosynthesis model*

We are using the traditional $C_3$ photosynthesis models (55) and $C_4$ models (29) for the

photosynthesis modeling (SI in supporting information).

*Hydraulic system*

At equilibrium, the rate of water loss through transpiration equals the rate of water absorption by

the roots (56):

$$\frac{EfN}{\rho} = k(1 - f)N(\Psi_l - \Psi_s) \qquad (1)$$

where $\Psi_s$ is soil water potential, $k$ is the effective root hydraulic conductivity, $N$ is the total

biomass of fine root and leaves, $\rho$ is the leaf mass density ($gcm^{-2}$) and $E$ is the transpiration rate

per leaf area. $E$ could be written as $\delta/r_s$, where $\delta$ is the water partial pressure deficit between

saturated leaf surface and the atmosphere. Thus, leaf water potential ($\Psi_l$) is a function of $r_s$ and

leaf/fine-root allocation ($f$, defined as investment into leaves/total investment in leaves and fine

root)).

$$\Psi_l = \Psi_s - \frac{f\delta}{\rho k r_s (1-f)} \qquad (2)$$

### Inhibition of photosynthesis by water stress

7 Reduced leaf water potential inhibits photosynthesis (57, 58, 59). We model this cost of

8 transpiration as Weibull-type vulnerability curves relating leaf $\Psi_l$ and photosynthetic parameters

9 (45):

$$V_{cmax,\Psi_l} = V_{cmax} e^{-(\frac{\Psi_l}{d_v})^{b_v}} \qquad (3)$$

$$J_{max,\Psi_l} = J_{max} e^{-(\frac{\Psi_l}{d_j})^{b_j}} \qquad (4)$$

$$V_{pmax,\Psi_l} = V_{pmax} e^{-(\frac{\Psi_l}{d_p})^{b_p}}, \qquad (5)$$

13 where $b$ and $d$ are curve fitting parameters. Since $\Psi_l$ is a function of $r_s$ and $f$, all those parameters

14 are functions of $r_s$ and $f$.

### Optimal stomatal resistance and optimal allocation of energy between leaves and fine roots

17 We assume that the plant adjusts the $r_s$ and $f$ to optimize the total carbon gain by

$$A_{total} = \frac{fNA_n}{\rho} \qquad (6)$$

where $\rho$ is the leaf mass density (g cm$^{-2}$). As a simplifying assumption, we assume $N$ and $\rho$ are

fixed (similar to 56). Effectively, we consider the optimization problem faced by the plant in a

given instance during its growth, where its size (of which $N$ is a proxy) can be regarded as a

constant. Clearly, during plant growth, the assimilate will be turned into plant biomass, but the

instantaneous optimization problem will still yield the optimal growth path as the growth rate is

maximized at any given time. Finally, we regard $\rho$ as a species-specific trait that changes at a

slower time-scale than $r_s$ and $f$. The first order optimality conditions for $r_s$ and $f$ are given by (56):

$$\frac{\partial(fA_n)}{\partial r_s} = f\frac{\partial A_n}{\partial r_s} = 0 \qquad (7)$$

$$\frac{\partial(fA_n)}{\partial f} = A_n + f\frac{\partial A_n}{\partial f} = 0. \qquad (8)$$

We checked the second order derivative to ensure that the numerical solutions to the first order

conditions were maxima.

*Allocation of nitrogen*

We examine how nitrogen allocation between RuBP carboxylation and RuBP regeneration in $C_4$

grasses affect competitive advantage over $C_3$ grasses. Despite great variation in $V_{cmax}$ and $J_{max}$

based on the total leaf nitrogen content within $C_3$ plants, Wullschleger (1993) (53) found a mean

of $J_{max}/V_{cmax}$ =2.1 across 109 $C_3$ species, which we use as a baseline for $C_3$ and $C_4$ pathways in

analyzing the initial evolution of $C_4$. Then, we used $J_{max}/V_{cmax}$ =4.5 for $C_4$ (22, 45) to analyze the

role that nitrogen reallocation played in the evolutionary trajectory of $C_4$ plants. In determining

the values of $J_{max}$ and $V_{cmax}$, we used a simplified stoichiometry: we consider the total of $J_{max}$ and

$V_{cmax}$ as a constant to hold nitrogen concentration constant (22, 45). Two assumptions are

underlying this simplified stoichiometry: (1) investing one molecule of nitrogen to the dark

reactions will increase of $V_{cmax}$ equal to the increase of $J_{max}$ by investing one molecule of

nitrogen to the light reactions; (2) nitrogen allocation to enzymes involved in photorespiration

and the CCM balanced each other.

*Modeling scenarios*

We modeled the photosynthesis rates of $C_3$ and $C_4$ under temperature range from 10 °C to 40 °C

with an interval of 5 °C, under $CO_2$ mixing ratios ranging from 200 ppm to 600 ppm with an

interval of 50 ppm, under different water conditions (VPD=0.001, 1, 2, 3, 4kPa corresponding to

soil water potential ($\Psi_S$) =0, -0.5, -1, -1.5, -2 MPa) and under different light intensities (1400,

1000, 600, 200, 100 μmol photons $m^{-2}s^{-1}$). We consider VPD=0.001 kPa and $\Psi_S$ =0 MPa as

saturated water condition and light intensity of 1400 μmol photons $m^{-2}s^{-1}$ as an average saturated

light intensity of a day.

**Acknowledgements**

We are grateful for support from the University of Pennsylvania.

**References**

1. Hatch MD (1987) $C_4$ photosynthesis: a unique blend of modified biochemistry, anatomy and ultrastructure. Biochim. *Biophys Acta* 895(2):81-106.

2. Sage RF (2004) The evolution of C4 photosynthesis. *New Phytol* 161(2):341-370.

3. Grass Phylogeny Work. Group II (2012) New grass phylogeny resolves deep evolutionary relationships and discovers $C_4$ origins. New Phytol 193:304–12.

4. Still CJ, Berry JA, Collatz GJ, DeFries RS (2003) Global distribution of $C_3$ and $C_4$ vegetation: Carbon cycle implications. Global Biogeochem Cycles 17(1):1006.

5. Ehleringer JR, Monson RK (1993) Evolutionary and ecological aspects of photosynthesis pathway variation. *Annu Rev Ecol Syst* 24:411-439. DOI: 10.1146/annurev.es.24.110193.002211.

6. Edwards EJ, Smith SA (2010) Phylogenetic analyses reveal the shady history of $C_4$ grasses. *Proc Natl Acad Sci USA* 107(6):2532-2537.

7. Pau S, Edwards EJ, Still CJ (2013) Improving our understanding of environmental controls on the distribution of $C_3$ and $C_4$ grasses. *Glob Chang Biol* 19(1):184-196.

8. Christin PA, Besnard G, Samaritani E, Duvall MR, Hodkinson TR, Savolainen V, Salamin N (2008) Oligocene $CO_2$ decline promoted $C_4$ photosynthesis in grasses. Curr Biol 18(1):37-43.

9. Ehleringer JR, Cerling TE, Helliker BR (1997) $C_4$ photosynthesis, atmospheric $CO_2$, and climate. *Oecologia* 112(3):285-299.

10. Collatz GJ, Berry JA, Clark JS (1998) Effects of climate and atmospheric $CO_2$ partial pressure on the global distribution of $C_4$ grasses: present, past, and future. *Oecologia* 114(4):441-454.

11. Ghannoum O (2009) $C_4$ photosynthesis and water stress. Ann Bot 103(4):635–644.

12. Taylor SH, Ripley BS, Martin T, De-Wet L-A, Woodward FI, Osborne CP (2014) Physiological advantages of $C_4$ grasses in the field: a comparative experiment demonstrating the importance of drought. *Glob Change Biol* 20(6):1922-2003.

13. Edwards EJ, Still CJ (2008) Climate, phylogeny and the ecological distribution of $C_4$ grasses. *Ecol Lett* 11(3):266-276.

14. Royer DL (2014) Atmosperic $CO_2$ and $O_2$ during the Phanerozoic: tools, patterns and impacts. In: Treatise on Geochemistry, 2nd edition, Eds: H. Holland and K. Turekian. http://dx.doi.org/10.1016/B978-0-08-095975-7.01311-5.

15. Way DA, Katul GG, Manzoni S, Vico G (2014) Increasing water use efficiency along the $C_3$ to $C_4$ evolutionary pathway: a stomatal optimization perspective. *J Exp Bot* 65(13):3683-3693.

16. Edwards EJ, Osborne CP, Strömberg CAE, Smith SA, C4 Grasses Consortium (2010) The Origins of $C_4$ Grasslands: Integrating Evolutionary and Ecosystem Science. *Science* 328(5978):587-591.

17. Kadereit G, Ackerly D, Pirie MD (2012) A broader model for $C_4$ photosynthesis evolution in plants inferred from the goosefoot family (Chenopodiaceae ss). Proc R Soc Lond B Biol Sci 279(1741): 3304-3311.

18. Vicentini A, Barber JC, Aliscioni SS, Giussani LM, Kellogg EA (2008) The age of the grasses and clusters of origins of $C_4$ photosynthesis. Glob Change Biol 14(12):2963-2977.

19. Pagani M, Freeman KH, Arthur MA (1999) Late Miocene atmospheric $CO_2$ concentrations and the expansion of C4 grasses. Science 285(5429):876-879.

20. Osborne CP (2008) Atmosphere, ecology and evolution: what drove the Miocene expansion of $C_4$ grasslands? *J Ecol* 96(1):35-45.

21. Strömberg CAE (2011) Evolution of grasses and grassland system. *Annu Rev Earth Planet Sci* 39:517-544. DOI:10.1146/annurev-earth-040809-152402.

22. Osborne CP, Sack L (2012) Evolution of $C_4$ plants: a new hypothesis for an interaction of $CO_2$ and water relations mediated by plant hydraulics. *Philos Trans R Soc Lond B Biol Sci* 367(1588):583-600.

23. Sage RF, Sage TL, Kocacinar F (2012) Photorespiration and the evolution of $C_4$ photosynthesis. *Annu Rev Plant Biol* 63:19-47. DOI:10.1146/annurev-arplant-042811-105511.

24. Griffiths H, Weller G, Toy L, Dennis RJ (2012) You're so vein: bundle sheath physiology, phylogeny and evolution in $C_3$ and $C_4$ plants. *Plant Cell Environ* 36(2):249–261.

25. Osborne CP, Freckleton RP (2009) Ecological selection pressures for $C_4$ photosynthesis in the grasses. *Proc Biol Sci* 276(1663):1753-1760.

26. Tissue DT, Griffin KL, Thomas RB, Strain BR (1995) Effects of low and elevated $CO_2$ on $C_3$ and $C_4$ annuals. II. Photosynthesis and leaf biochemistry. *Oecologia* 101(1):21-28.

27. Ghannoum O, Evans JR, von Caemmerer S (2010) Nitrogen and water use efficiency of $C_4$ plants. In: *$C_4$ photosynthesis and related $CO_2$ concentrating mechanisms* (eds. Raghavendra, A.S. & Sage, R.F.). Springer Science, The Netherlands, Dordrecht, pp. 129–146.

28. Prentice IC, Dong N, Gleason SM, Maire V, Wright IJ (2014) Balancing the costs of carbon gain and water transport: testing a new theoretical framework for plant functional ecology. *Ecol Lett* 17(1):82-91.

29. von Caemmerer S (2000) *Biochemical models of photosynthesis. In Techniques in Plant Sciences*. CSIRO Publishing, Colingwood, Australia, pp 91–122.

30. Chen D-X, Coughenour MB, Knapp AK, Owensby CE (1994) Mathematical simulation of $C_4$ grass photosynthesis in ambient and elevated $CO_2$. *Ecol Model* 73(1-2):63–80.

31. Collatz GJ, Ball JT, Grivet C, Berry JA (1992) Coupled photosynthesis-stomatal conductance model for leaves of $C_4$ plants. *Aust J Plant Physiol* 19(5):519-538.

32. Ehleringer J, Björkman O (1977) Quantum yields for $CO_2$ uptake in $C_3$ and $C_4$ plants. Plant Physiol 59(1): 86-90.

33. Cerling TE, Harris JM, MacFadden BJ, Leakey MG, Quade J, Eisenmann V, Ehleringer JR (1997). Global vegetation change through the Miocene/Pliocene boundary. *Nature* 389(6647):153-158.

34. Christin PA, Osborne CP, Sage RF, Arakaki M, Edwards EJ (2011) $C_4$ eudicots are not younger than $C_4$ monocots. J Exp Bot 62(9):3171-3181.

35. Liu H, Edwards EJ, Freckleton RP, Osborne CP (2012) Phylogenetic niche conservatism in $C_4$ grasses. Oecologia 170(3):835-845.

36. Hattersley PW (1983) The distribution of $C_3$ and $C_4$ grasses in Australia in relation to climate. Oecologia 57(1):113–128.

37. Vogel JC, Fuls A, Danin A (1986) Geographical and environmental distribution of $C_3$ and $C_4$ grasses in the Sinai, Negev, and Judean deserts. Oecologia 70(2): 258–265.

38. Paruelo JM, Lauenroth WK (1996) Relative abundance of plant functional types in grasslands and shrublands of North America. Ecol Appl 6(4):1212–1224.

39. Quade J, Cerling TE, Bowman JR (1989) Development of Asian monsoon revealed by marked ecological shift during the latest Miocene in northern Pakistan. Nature 342(6246):163-166.

40. Taylor SH, Franks PJ, Hulme SP, Spriggs E, Christin PA, Edwards EJ, Woodward FI, Osborne CP (2012) Photosynthetic pathway and ecological adaptation explain stomatal trait diversity amongst grasses. *New Phytol* 193(2):387-396.

41. Knapp AK, Briggs JM, Hartnett DC, L. Collins L (1998) Patterns and controls of aboveground net primary production in tallgrass prairie. Grassland Dynamics: Long-Term Ecological Research in Tallgrass Prairie eds Knapp AK, Briggs JM, Hartnett DC, L. Collins L (Oxford University Press, New York), pp 193-221.

42. Christin PA, Osborne CP, Chatelet DS, Columbus JT, Besnard G, Hodkinson TR, Garrison LM, Vorontsova MS, Edwards EJ (2013) Anatomical enablers and the evolution of $C_4$ photosynthesis in grasses. *Proc Natl Acad Sci USA* 110(4):1381-1386.

43. Kocacinar F, Sage RF (2003) Photosynthetic pathway alters xylem structure and hydraulic function in herbaceous plants. Plant Cell Environ 26(12):2015-2026.

44. Ku MSB, Schmitt MR, Edwards GE (1979) Quantitative determination of RuBP carboxylase-orygenase protein in leaves of several $C_3$ and $C_4$ plants. *J Exp Bot* 114(1)89–98.

45. Vico G, Porporato A (2008) Modelling $C_3$ and $C_4$ photosynthesis under water-stressed conditions. *Plant and Soil* 313(1):187-203.

46. Vogan PJ, Sage RF (2012) Effects of low atmospheric $CO_2$ and elevated temperature during growth on the gas exchange responses of $C_3$, $C_3$–$C_4$ intermediate, and $C_4$ species from three evolutionary lineages of $C_4$ photosynthesis. *Oecologia* 169(2):341-352.

47. Domingues TF, Martinelli LA, Ehleringer JR (2007) Ecophysiological traits of plant functional groups in forest and pasture ecosystems from eastern Amazônia, Brazil. *Plant Ecol* 193(1)101-112.

48. Massad RS, Tuzet A, Bethenod O (2007) The effect of temperature on $C_4$-type leaf photosynthesis parameters. *Plant Cell Environ*, 30(9):1191-1204.

49. Grant RF, Flanagan LB (2007) Modeling stomatal and nonstomatal effects of water deficits on $CO_2$ fixation in a semiarid grassland. *J Geophys Res 112*, G03011, doi:10.1029/2006JG000302.

50. Kathilankal JC, Mozdzer TJ, Fuentes JD, McGlathery KJ, D'Odorico P, Zieman JC (2011) Physiological responses of Spartina alterniflora to varying environmental conditions in Virginia marshes. *Hydrobiologia* 669(1):167-181.

51. Ye Z, Suggett DJ, Robakowski P, Kang H (2013) A mechanistic model for the photosynthesis-light response based on the photosynthetic electron transport of photosystem II in $C_3$ and $C_4$ species. *New Phytol* 199(1):110–120.

52. Ge ZM, Zhang LQ, Yuan L, Zhang C (2014) Effects of salinity on temperature-dependent photosynthetic parameters of a native $C_3$ and a non-native $C_4$ marsh grass in the Yangtze Estuary, China. *Photosynthetica*, 52(4):484–492.

53. Wullschleger SD (1993) Biochemical limitations to carbon assimilation in $C_3$ plants - A retrospective analysis of the A/C$_i$ curves from 109 species. *J Exp Bot* 44(5):907-920.

54. Christin PA, Osborne CP (2014) Tansley Review. The evolutionary ecology of $C_4$ photosynthesis. *New Phytol*, 204(4):765-781.

55. Farquhar GD, Von Caemmerer S, Berry JA (1980) A biochemical model of photosynthetic carbon dioxide assimilation in leaves of 3-carbon pathway species. *Planta* 149(1):78–90.

56. Givnish TJ (1986) Optimal stomatal conductance, allocation of energy between leaves and roots, and the marginal cost of transpiration. In: *On the Economy of Plant Form and Function* (ed. Givnish TJ). Cambridge University Press, Cambridge, pp. 171–213.

57. Tezara W, Mitchell VJ, Driscoll SD, Lawlor DW (1999) Water stress inhibits plant photosynthesis by decreasing coupling factor and ATP. *Nature* 401(6756):914-917.

58. Lawlor DW, Cornic G (2002) Photosynthetic carbon assimilation and associated metabolism in relation to water deficits in higher plants. *Plant Cell Environ* 25(2):275–294.

59. Tang AC (2002) Photosynthetic oxygen evolution at low water potential in leaf discs lacking an epidermis. *Ann of Bot* 89(7):861-870.

Figure Legends

Fig. 1. Crossover temperatures of photosynthesis for $C_3$ and $C_4$ with the change of $CO_2$

concentration under different water conditions. Light intensity was 1400 μmol photons $m^{-2}s^{-1}$ for

all model runs. $J_{max}/V_{cmax}$=2.1 for $C_3$ and $J_{max}/V_{cmax}$=4.5 for $C_4$. Solid black line: VPD=0.15kPa,

$\Psi_S$=0 MPa; dashed black line: VPD=1kPa, $\Psi_S$=-0.5 MPa; dot-dashed black line: VPD=2kPa,

$\Psi_S$=-1 MPa; dotted black line: VPD=3 kPa, $\Psi_S$=-1.5 MPa.

Fig. 2. The total difference in $CO_2$ assimilation between $C_4$ and $C_3$ ($A_n(C_4)$-$A_n(C_3)$) under various

$CO_2$ (200 ppm, 400 ppm and 600 ppm) and water conditions under saturated light intensity (1400

μmol photons $m^{-2}s^{-1}$). $J_{max}/V_{cmax}$=2.1 for $C_3$ and $C_4$ (a, c, e) and $J_{max}/V_{cmax}$=2.1 for $C_3$ and

$J_{max}/V_{cmax}$=4.5 for $C_4$ (b, d, f). Water limitation intensity is: 1: VPD =0.15 kPa, $\Psi_S$=0 MPa; 2: 1.5

kPa, -0.5MPa; 3: 2kPa, -1 MPa; 4: 3kPa, -1.5 MPa; 5: 4kPa, -2 MPa.

Fig. 3. The total difference in $CO_2$ assimilation between $C_4$ and $C_3$ ($A_n(C_4)$-$A_n(C_3)$) with

$J_{max}/V_{cmax}$=2.1 for $C_3$ and $C_4$ under various $CO_2$ (200 ppm, 400 ppm) and different light

intensities (from 200 to 1400 μmol photons $m^{-2}s^{-1}$) with saturated water condition

(VPD=0.15kPa, $\Psi_S$=0 MPa) (a, c) and with $J_{max}/V_{cmax}$=2.1 for $C_3$ and $J_{max}/V_{cmax}$=4.5 for $C_4$ (b,

d).

Fig. 4. Assimilation rates of $C_3$ with $J_{max}/V_{cmax}$=2.1 (solid black line), C4 with $J_{max}/V_{cmax}$=2.1

(dashed black line) and $C_4$ with $J_{max}/V_{cmax}$=4.5 (dotted black line) under light intensity of 1400

μmol photons $m^{-2}s^{-1}$, $CO_2$ of 400 ppm and different water limitation conditions. (a) saturated

soils; (b) VPD=1kPa and $\Psi_S$=-0.5 MPa; (c) VPD=2kPa and $\Psi_S$=-1 MPa.

Fig. 5. Stomatal resistance ($r_s$) and leaf/fine-root allocation (f) as a function of temperature, with

$J_{max}/V_{cmax}$=2.1 for both $C_3$ and $C_4$ with saturated light under $CO_2$ of 400 ppm and different water

conditions. Solid black line: $C_3$ with VPD=0.15kPa, $\Psi_S$=0 MPa; dashed black line: $C_4$ with

VPD=0.15kPa, $\Psi_S$=0; solid grey line: $C_3$ with VPD=4 kPa, $\Psi_S$=-2 MPa; dashed grey line: $C_4$

with VPD=4 kPa, $\Psi_S$=-2 MPa. Vertical lines indicate transition from RuBP carboxylation

limited condition to RuBP regeneration limited condition for $C_3$ and $C_4$.