**High-Throughput Sequencing of Transposable Elements Insertions**

**Provides Evidence for Adaptive Evolution of the Invasive Asian Tiger**

**Mosquito Towards Temperate Environments**

Clément Goubert[1], Hélène Henri[1], Guillaume Minard[2,3], Claire Valiente Moro[2], Patrick

Mavingui[2,4], Cristina Vieira[1] and Matthieu Boulesteix[1]

[1]Université de Lyon, F-69622, Lyon, France; Université Claude Bernard Lyon 1, CNRS,

Laboratoire de Biométrie et Biologie Evolutive, UMR5558, F-69100 Villeurbanne, France

[2]Université de Lyon, F-69622, Lyon, France; Université Lyon 1, Villeurbanne, France;

CNRS, UMR 5557, Ecologie Microbienne, Villeurbanne, France; INRA, UMR 1418,

Villeurbanne, France

[3]Metapopulation Research Center, Department of Biosciences, University of Helsinki,

Helsinki, Finland

[4]Université de La Réunion, UMR PIMIT, INSERM 1187, CNRS 9192, IRD 249,

Plateforme Technologique CYROI, Sainte-Clotilde, La Réunion.

**Keywords:** Invasive species, *Aedes albopictus*, local adaptation, genome scan,

Transposable Elements, diapause

**Corresponding author:** Matthieu Boulesteix, Université de Lyon, F-69622, Lyon,

France; Université Claude Bernard Lyon 1, CNRS, Laboratoire de Biométrie et Biologie

Evolutive, UMR5558, F-69100 Villeurbanne, France; phone: +334 72 43 29 16, fax:

+334 72 43 13 88; email: matthieu.boulesteix@univ-lyon1.fr

**Running title:** Molecular adaptation in *Ae. albopictus*

# Abstract

Invasive species represent unique opportunities to evaluate the role of local

adaptation during colonization of new environments. Among these, the Asian

tiger mosquito, *Aedes albopictus*, is a threatening vector of several human viral

diseases, including dengue, chikungunya and the emerging Zika fevers. Its broad

presence in both temperate and tropical environments has sometimes been

considered as the reflect of a great "ecological plasticity". However, no study has

been conducted to assess the role of adaptive evolution in the ecological success

of *Ae. albopictu*s at the molecular level. In the present study we performed a

genomic scan to search for potential signatures of selection leading to local

adaptation in a hundred of field collected mosquitoes from native populations of

Vietnam and temperate invasive populations of Europe. High throughput

genotyping of transposable element insertions generated more than 120 000

polymorphic loci, which in their great majority revealed a virtual absence of

structure between bio-geographic areas. Nevertheless, 92 outlier loci show a high

level of differentiation between temperate and tropical populations. The majority

of these loci segregates at high insertion frequencies among European

populations, indicating that this pattern could have been caused by recent events

of adaptive evolution in temperate areas. Six outliers were located near putative

diapause effector genes, suggesting fine-tuning of this critical pathway during

42    local adaptation.

# Introduction

Biological invasions represent unique opportunities to study fast evolutionary changes such as adaptive evolution. Indeed, settlement in a novel area represents a biological challenge that invasive species have successfully overcome. The underlying processes could be studied at the molecular level, particularly to gather empirical knowledge about the genetics of invasions, a field of study that has produced extensive theoretical predictions, but for which there is still little evidence in nature (Colautti & Lau 2015). Some of the main concerns are to disentangle the effects of neutral processes during colonization, such as founder events or allele surfing at the migration front, from adaptive evolution (i.e. local adaptation, Lande 2015; Peischl & Excoffier 2015; Colautti & Lau 2015).

Adaptation can arise either through the appearance and spread of a new beneficial mutation, the spread of a favorable allele from standing genetic variation, or from hybridization in the introduction area (Handley *et al.* 2011; Colautti & Lau 2015; Bock *et al.* 2015). Detection of the footprint of natural selection is however dependent on the availability of informative genetic markers, which should provide a substantial coverage of the genome to allow selection scans and be easily and confidently scored across many individuals. Unfortunately, invasive organisms are rarely model species, making the development of a reliable and efficient marker challenging.

The Asian tiger mosquito, *Aedes* (*Stegomya*) *albopictus* (Diptera:Culicidae) is currently one of the most threatening invasive species (Invasive Species Specialist Group).

Originating from South-Eastern Asia, this species is one of the primary vectors of

Dengue and Chikungunya viruses, and is also involved in the transmission of other

threatening arboviruses (Paupy *et al.* 2009), in particular the newly emerging Zika virus

(Grard *et al.* 2014; Marcondes & Ximenes 2015; Chouin-Carneiro *et al.* 2016). *Ae.*

*albopictus* has now settled in every continent except Antarctica, and is found both under

tropical and temperate climates (Bonizzoni *et al.* 2013). While this species is supposed

to originate from rain forests of South-Eastern Asia (Hawley 1988), the native area of

*Ae. albopictus* encompasses contrasted environments including temperate regions of

Japan and China, offering a large potential of fit towards newly colonized environments.

For example, the induction of photoperiodic diapause in temperate areas, that has a

genetic basis in *Ae. albopictus* (Hawley *et al.* 1987; Hanson & Craig 1994; Urbanski *et*

*al.* 2010), is decisive to ensure invasive success in Europe or Northern America. Indeed

it allows the susceptible populations to survive through winter at the larval stage into the

eggs. Such a trait appears governed by a "genetic toolkit" involving numerous genes

and metabolic networks, for which however the genetic polymorphism between

diapausing and non-diapausing strains remains to be elucidated (Poelchau *et al.*

2013b). In addition, the colonization of new areas that look similar at first glance can still

involve *de novo* adaptation: indeed, even environment sharing climatic variables are not

necessarily similar regarding edaphic and biotic interactions (Colautti & Lau 2015).

Hence, this suggests that whatever are the native and settled environments, it might be

possible to find evidence of adaptive evolution in invasive populations of *Ae. albopictus*.

To better understand the invasive success of this species, we genotyped 140 field

individuals, collected from three Vietnamese (native tropical area) and five European

(invasive temperate area) populations, aiming to identify genomic regions involved in

local adaptation. To do so, we developed new genetic markers, based on high

throughput genotyping of the insertion of Transposable Elements (TEs) that represent at

least one third of the genome of *Ae. albopictus* and include recently active families that

could reach thousands of copies in one genome (Goubert *et al.* 2015).

Amplification of TE insertions is particularly efficient to obtain a large number of genetic

markers throughout one genome (Bonin *et al.* 2008), especially if few genomic

resources are available (Monden *et al.* 2014), which was case until recently for the

Asian tiger mosquito. In addition, such markers represent a seducing alternative to other

methods of diversity reduction, such as RAD-sequencing (Miller et al. 2007), that could

be less efficient in species with high TE load (Davey et al. 2012) and did not produce

satisfying results in *Ae. albopictus* (Goubert *et al.* 2016). In mosquitoes, TEs have been

shown to be powerful markers both for population structure analysis (Biedler *et al.* 2003;

Boulesteix *et al.* 2007; Santolamazza *et al.* 2008; Esnault *et al.* 2008) and genome

scans (Bonin *et al.* 2009). We hypothesized that some TE insertion sites could be

located at the neighborhood of targets of natural selection and thus could reach high

level of differentiation between native and invasive populations if selective sweeps

occurred during local adaptation. In addition, some TEs could also insert near or inside

coding regions and thus could be directly involved in environmental adaptation

(Casacuberta & González 2013), eventually contributing to the success of invasive

species (Stapley *et al.* 2015).

To distinguish between neutral demographic effects and adaptive evolution, we first

performed population genetic analyses to reveal the global genetic structure of the

studied populations. We then performed a genomic scan for selection and identified 92

candidate loci under directional selection, among which several can be located in the

neighborhood of diapause related genes.

# Material and Methods

### Biological samples

115 A total number of 140 flying adult females *Ae. albopictus* were collected in the field at

116 eight sampling sites in Europe and Vietnam during the summers of 2012 and 2013

117 (Figure 1 and Table S1). Individuals were either sampled using a single trap or using

118 aspirators through the sampling site within a 50 meters radius. When traps were used,

119 live mosquitoes were collected after a maximum of two days.

### High throughput Transposon Display (TD) genotyping.

Insertion polymorphism of five transposable elements families: I Loner Ele1 (IL1), Loa

Ele2B (L2B), RTE4, RTE5 and Lian 1 identified by Goubert *et al.* (2015) in *Ae.*

*albopictus* was characterized using Transposon Display (TD), a TE insertion specific

PCR method, combined with Illumina sequencing of all TD amplification products

(Figure S1). These TE families were chosen according to their high estimate of copy

number (from 513 to 4203 copies), high identity between copies, and a "copy and paste"

mode of transposition (all these TEs are non-LTR Class I retrotransposons).

**DNA Extraction and TD adapter ligation.** Total DNA was extracted from whole adult

bodies following the Phenol-Chloroform protocol described by Minard *et al.* (2015). TD

was then set up combining methods from previous studies (Munroe *et al.* 1994; Roy *et*

*al.* 1999; Akkouche *et al.* 2012; Carnelossi *et al.* 2014). First, individual extracted DNA

($\approx$ 75ng) was used for enzymatic digestion in a total volume of 20 $\mu$L, with HindIII

enzyme (10U/$\mu$L) and buffer R (Thermo Scientific) for 3 hours at 37C. The enzyme was

inactivated at 80C for 20 minutes. TD adapters were then build hybridizing Hindlink with

MSEB oligonucleotides (100$\mu$M, see Table S2) in 20X SSC and 1M Tris in a total

volume of 333$\mu$L after 5mn of initial denaturation at 92C and 1h at room temperature for

hybridization of the two parts. Once ready, TD adapters were then ligated to 20 $\mu$L of the

digested DNA mixing 2$\mu$L of TD adapter with 10U T4 ligase and 5X buffer (Fermentas)

in a final volume of 50$\mu$L for 3 hours at 23C.

**Library construction.** For each individual, and for each of the five TE families, TE

insertions were amplified by PCR (PCR 1) in a Biorad Thermal Cycler (either C1000 or

S1000), in a final volume of 25$\mu$L. Mixture contained 2$\mu$L of digested-ligated DNA with

1$\mu$L dNTPs (10mM), 0.5$\mu$L TD-adapter specific primer (LNP, 10$\mu$M, see S3 Table) and

0.5 $\mu$L of TE specific primer (10$\mu$M), 1U AccuTaq polymerase (5U/$\mu$L) with 10X buffer

and Dimethyl-Sulfoxyde (Sigma). Amplification was performed as follows: denaturation

at 98C for 30 seconds then 30 cycles including 94C for 15 seconds, hybridization at 60C

for 20 seconds and elongation at 68C for 1 minute; final elongation was performed for 5

minutes at 68C. For L2B and RTE5 TEs, a nested PCR was performed in order to

increase specificity in the same PCR conditions using internal forward TE primers and

LNP (Table S2). PCR 1 primers include a shared tag sequence that was used for

hybridization of the individual indexes by PCR 2.

For each TE, three independents PCR 1 were performed from the same digestion

product. PCR 1 products (3 PCR * 5 TE per individual) were then purified using volume-

to-volume Agencout AMPure XP beads (20$\mu$L PCR 1 + 20$\mu$L beads) and eluted in 30$\mu$L

Resuspension buffer. After nanodrop quantification, equimolar pools containing the 3*5

PCR products per individual were made using Tecan EVO200 robot. Individual pools

were then size selected for fragment ranging from 300 to 600 bp using Agencout

AMPure XP beads as follow: first magnetic beads were diluted in H2O with a ratio of

1:0.68 then add to 0.625 X PCR products in order to exclude long fragments. A second

purification was performed using a non-diluted bead: DNA ratio of 1:8.3 to exclude small

fragments.

Samples' multiplexing was performed using home made 6 bp index (included in SRA

individual name), which were added to the R primer (Table S2) during a second PCR

(PCR 2) with 12 cycles in ABI 2720 Thermal Cycler. Mixture contained 15ng PCR

products, 1$\mu$l of dNTPs (10mM), 0.5$\mu$l MTP Taq DNA Polymerase (5U/$\mu$l, Sigma), 5$\mu$l

10X MTP Taq Buffer and 1.25$\mu$l of each tagged-primer (20$\mu$m) in a final volume of 50$\mu$l.

Amplification was performed as follows: denaturation at 94C for 60 seconds then 12

cycles including denaturation at 94C for 60 seconds, hybridization at 65C for 60 seconds

and elongation at 72C for 60 seconds; final elongation was performed for 10 minutes at

72C. PCR 2 products were purified using Agencout AMPure XP beads: DNA ratio of

1:1.25 to obtain libraries. Finally, TD products were paired-end sequenced on an

Illumina Hiseq 2000 (1 lane) at the GeT-PlaGe core facility (Genome and

Transcriptome, Toulouse) using TruSeq PE Cluster Kit v3 (2x100 bp) and TruSeq SBS

Kit v3.

**Bioinformatic treatment of TD sequencing.**

The different steps of the informatics treatment from the raw sequencing dataset to

population binary matrices for presence/absence of TE insertions per individual are

described in Figure S2. First, the paired-end reads of each individual were quality

checked and trimmed using UrQt v. 1.0.17 (Modolo & Lerat 2015) with standard

parameters and a *t* quality threshold of 10. Reads pairs were then checked and trimmed

for Illumina adapter contamination using cutadapt (Martin 2011). Specific amplification of

TE insertions was controlled by checking for the expected 3' TE sequence on the R1

read using Blat (Altschul *et al.* 1990) with an identity threshold of 0.90. Only reads with

an alignment-length/read-length ratio ≥ 0.90 were then retained. R2 reads for which

the R1 mate passed this filter were then selected for the insertion loci construction, after

the removal of the TD adapter on the 5' start using cutadapt and the removal of reads

under 30 bp. Selected reads were separated in each individual according to the TE

families for loci construction.

In order to correct the inter-individual coverage variations, we performed a sampling of

the cleaned reads. First, for each TE family, distribution of the number of read per

individual was drown, and individuals with less reads than the first decile of this

distribution were removed; then cleaned reads of the remaining individuals were

randomly sampled at the value of the first decile of coverage (this value varies among

TE families). For each TE, the sampled reads of each retained individual were clustered

together using the CD-HIT-EST program (Li & Godzik 2006) to recover insertion loci.

During this all-to-all reads comparison, the alignments must had a minimum of 90

percent identity, and the shortest sequence should be 95% length of the longest, global

identity was used and each read was assigned at its best cluster (instead of the first that

meet the threshold). In a second step, the reference reads of each locus within

individual, given by CD-HIT-EST, were clustered with all the reference reads of all

individuals, using the same threshold, in order to build the locus catalog including list of

loci of all individuals and the coverage for each locus in each individual. After this step,

insertion loci that matched known repeats of the Asian tiger mosquito (Goubert *et al.*

2015) were discarded; alignments were performed with Blastn using default parameters.

Since the quality control removed a substantial number of reads for the construction of

TE insertions catalog, the raw R2 reads (with their TD adapter removed), that could

have been discarded in a first attempt were then mapped over the catalog in order to

increase the scoring sensibility. Before mapping, the raw R2 reads were also sampled at

the first decile of individual coverage (as described previously). At this step, individuals

that have been removed from at least two TE families for loci construction were

definitively removed from the whole analysis. Mapping was performed over all insertion

loci of all TE families in a single run in order to prevent multiple hits. Blat was used with

an identity threshold of 90 percent. Visual inspection of alignment quality over 30

sampled loci per TE family was performed in order to ensure the quality of scoring.

In order to check if the sampling procedure would affect our results, the read sampling

procedures and subsequent analysis were performed independently 3 times (replicates

M1, M2 and M3).

**Genetic analyses and Genomic scan.**

Population structure analyses were performed independently for each TE family.

Principal Coordinate Analysis (PCoAs) were performed to identify genetic clusters using

the ade4 package (Dray & Dufour 2007) of R 3.2.1 (R development core team 2015). S7

coefficient of Gower and Legendre was used as a genetic distance since it gives more

weight to shared insertions. Shared absences were not used because they do not give

information about the genetic distance between individuals. Pairwise populations $F_{ST}$

were computed using Arlequin 3.5 (Excoffier & Lischer 2010); significance of the index

was assessed over 1,000 permutations using a significance threshold of 0.05.

The genomic scan was performed in two steps for each of the sampling replicates of

each TE. First, Bayescan 2.1 (Foll & Gaggiotti 2008) was used to test for each locus

deviation from neutrality. Bayescan consider a fission/island model where all

subpopulations derive from a unique ancestral population. In this model, variance in

allele frequencies between subpopulations is expected to be due either to the genetic

drift that occurred independently in each subpopulation or to selection that is a locus-

specific parameter. The differentiation at each locus in each subpopulation from the

ancestral population is thus decomposed into a $\beta$ component (shared by all loci in a

subpopulation) and is related to genetic drift, and a $\alpha$ component (shared for a locus by

all subpopulations) due to selection. Using a Bayesian framework, Bayescan tests for

each locus the significance of the $\alpha$ component. Rejection of the neutral model at one

locus is done using posterior Bayesian probabilities and controlled for multiple testing

using false discovery rate. In addition, Bayescan manage uncertainty about allele

frequency from dominant data such as the TD polymorphism, leaving the $F_{IS}$ to freely

vary during the estimation of parameters. Bayescan was used with default values except

for the prior odds that were set to 100 (more compatible with datasets with a large

number of loci, see Bayescan manual), and a significance $q$-value threshold of 0.05 was

used to retain outlier loci. In a second step, only outliers suggesting divergent directional

selection between, Europe and Vietnam were considered. To identify them, locus-by-

locus Analyses of Molecular Variance (AMOVAs) were performed using Arlequin 3.5 for

each TE family. Significance of the $F_{CT}$ (inter group differentiation) between Vietnamese

and European populations was assessed performing 10,000 permutations with a

significance threshold of 0.05. For each dataset, Bayescan outliers were crossed with

significant $F_{CT}$ loci to retain candidate loci.

To identify the genomic environment of the candidate loci, the outlier sequences

(reference R2 read) were mapped onto the assembled genome of *Ae. albopictus* (Chen

*et al.* 2015) using Blastn. Blastn alignments were performed with default parameters and

sorted according to alignment score and after visual inspection of each alignment.

Outlier loci with multiple identical hits were discarded. To identify genes surrounding the

mapped outliers, the complete transcriptome of *Ae. albopictus* (including eggs, larvae

and adult females, downloaded at www.albopictusexpression.org Armbruster *et al.*) was

mapped over the reference genome using blat with default parameters; after alignment,

one best hit was retained per transcript according to the best alignment score. When a

transcript had multiple best hits, all positions for the transcript were considered.

**PCR validation and Outlier analyses.**

Pairs of primers were designed for each outlier locus in order to be used in standardized

conditions. Forward primer was located in the TE end of the concerned family and

reverse primer was set from the outlier locus (pairs of primers which successfully

amplified insertions are provided in Table S2). All primer pairs were first tested on a set

of 10 individuals in order to assess their specificity using 1/50 dilution of starting DNA

from the TD experiment. Validated primers were then used to check the insertions

polymorphism in 47 representative individuals from the 8 populations studied in the TD

experiment using 1/50 dilutions of the starting DNA (not all individuals could be used

because of DNA limitations). All PCRs were conducted in a final volume of 25$\mu$L using

0.5$\mu$L of diluted DNA, 0.5$\mu$L of each primer (10$\mu$M), 1$\mu$l of dNTPs (10mM) and 1U of

DreamTaq Polymerase with 1X green buffer (ThermoFisher Scientific). Amplification

was performed as follows: denaturation at 94C for 2 minutes then 34 cycles including

denaturation at 94C for 30 seconds, hybridization at 60C for 45 seconds and elongation

at 72C for 45 seconds; final elongation was performed for 10 minutes at 72C. After 45

minutes migration of the PCR product on 1X electrophoresis agarose gel, CG and MB

assessed insertion polymorphism independently.

# Results

**High throughput TE insertion genotyping.**

Sequencing produced a total of 102,319,300 paired-end reads (2x101bp). After quality

and specificity filtering 24,332,715 reads were suitable for analyses. On average, a total

number of 128,491 polymorphic insertion loci were available for each of the three

sampling replicates. The mean number of loci per individual and per TE family ranged

from 1025 ± 290 s.d. (IL1 family, mean and s.d. averaged over the three replicates) to

3266 ± 766 s.d. (RTE5 family). Details are given in Table S1. While our read sampling

procedure could have artificially lowered the mean insertion frequency of the loci, this

effect should be small because in our final datasets, the TE insertion frequencies (*i. e.*

the number of individuals that share an insertion) are not correlated with the mean

number of read per individual at the considered locus (Figure S3).

**Population structure.**

Principal Coordinate Analyses (PcoAs) were performed independently for each of the

five TEs (Figure 2). Among the three main Principal Coordinates (PCs), individuals tend

to be grouped according to their respective populations with little overlap between

groups. However, the three main PCs represent only a small fraction of the total

variance (< 10%), suggesting a weak genetic structuring between the populations.

Overall, individuals from Vietnamese populations (HCM, TA, VT) tend to be grouped

together in a single cluster, at the exception of 13 to 14 individuals from HCM when

using L2B and RTE5 TE families (S2 Figure), along with six individuals of VT with the

RTE4 TE family (Figure 1) that can not be clearly distinguished from European samples.

BCN individuals (Spain) represent the most homogeneous group,  well differentiated

from Vietnamese and French individuals (SP, CGN, NCE and PLV).

In agreement with PCoAs, Analyses of Molecular Variance (AMOVAs) attributed very

few genetic variances among groups (Vietnam-Europe) and between populations within

groups (Table 1). In the studied populations, most of the genetic variance was

distributed among individuals within groups.

Measures of genetic differentiation among pairs of populations were consistent with

PCoAs and AMOVAs (File S1): the BCN population shows the highest $F_{ST}$ values with

the other populations for each of the five TEs (0.051 < $F_{ST}$ < 0.148), while Vietnamese

populations were the most closely related (0.011 < $F_{ST}$ < 0.032). While VT is located

100 km away from TA and HCM (both sampled in the same city, Hô Chi Minh, Vietnam)

the $F_{ST}$ values are very similar between the three Vietnamese populations, suggesting

no influence of geography at this scale. CGN and NCE, sampled in the same urban area

(Nice agglomeration), are also little or not significantly differentiated depending on the

TE family. The previously identified intermediate pattern of HCM with some European

populations at L2B and RTE5 loci (PCoAs analyses) is also found at the $F_{ST}$ level,

especially regarding the low differentiation with the PLV population for these markers

(0.011 < $F_{ST}$ < 0.020).

**Genomic scan.**

Research of outlier loci for both selection signature using Bayescan (non-hierarchical

island model), and for significant $F_{CT}$ (between Europe-Vietnam group differentiation)

identified 92 candidate insertion loci (Figure 3). Most of these insertions are found in

both areas (no private allele), except for RTE4_6 and RTE4_7 that were not found in

Vietnam. In addition, a majority of outliers corresponds to high frequency insertions in

Europe, while the same trend is not observed at 92 randomly chosen loci among those

having the same minimum insertion frequency (≥ 20 individuals/locus) between Europe

and Vietnam (Figure 4). PCR amplification of the outlier loci were carried out on a

representative panel of 47 individuals to validate the insertion pattern detected by TD

(see Material and Method). For loci where the amplification was successful, the insertion

pattern observed by PCR always confirmed the results from TD (Figure S5).

From 92 outlier loci, 21 could be attributed to a unique position on the *Ae. albopictus*

genome. Annotation and distance to surrounding genes are reported on S2 File. We

found that six outliers (S2 File, sheet 3, highlighted) loci are located on contigs that

harbor genes previously identified in *Ae. albopictus* as being differentially expressed

between diapause-induced and non diapause-induced samples , and for which

orthologs in *Drosophila melanogaster* are known to be part of well-identified functional

networks (Poelchau *et al.* 2013b). All these six loci are found to be outliers because of

their high insertion frequencies in Europe compared with Vietnam.

# Discussion

The goal of our study was to identify genomic regions involved in adaptive evolution of *Ae. albopictus* thanks to the development of new genetic markers. Through high-throughput genotyping of five TE families' insertion polymorphisms, we identified up to 128,617 polymorphic loci among a hundred of individuals from eight sampling sites. The estimated genome size of *Ae. albopictus* exceeds one billion base-pairs (Goubert *et al.* 2015; Dritsou *et al.* 2015; Chen *et al.* 2015). Accordingly, the amount of markers scored in this study offers a comfortable genomic density of one marker every 10 kb.

We provide here a new and cost efficient method to quickly generate a large amount of polymorphic markers without extensive knowledge about one species genome. Specifically, this strategy appears extremely valuable for species with a large genome size, where TE density could severely compromise the development of more classical approaches, such as the very popular RAD-sequencing.

The genetic structure of the studied populations showed strong consistency between sampling replicates of individuals' reads, demonstrating the robustness of the method in spite of an initial substantial coverage variation among individuals. Population genetics analyses revealed a very low level of genetic structuring between European and Vietnamese populations. Among the studied populations, AMOVAs showed that most of the genetic variation is distributed between individuals within populations (> 90%), and as suggested by pairwise $F_{ST}$ and PCoAs, only a small part (< 10%) of the genetic variance is due to differentiation between populations. The genetic differentiation we

measured is indeed as high among European populations as it is between populations

from Europe and Vietnam.

This singular population structure is in agreement with previous results gathered in *Ae.*

*albopictus* using different collections of allozymes, mtDNA or microsatellites markers

(Black *et al.* 1988; Kambhampati *et al.* 1991; Zhong *et al.* 2013; Gupta & Preet 2014;

Manni *et al.* 2015). Moreover, a recent analysis performed with a set of 11

microsatellites on individuals from the same populations (at the exception of BCN)

showed a similar distribution of genetic variation among hierarchical levels (Minard *et al.*

2015). These results demonstrate the reliability of our markers and confirm that a non-

hierarchical island model can likely fit the global genetic structure. The genetic diversity

observed in Europe is compatible with a scenario of multiple and independent

introductions, as already suggested for *Ae. albopictus* (Urbanelli *et al.* 2000; Birungi &

Munstermann 2002; Takumi *et al.* 2009; Becker *et al.* 2013). However, as previously

suggested, this pattern could also be the result of founder events that may occur during

colonization and/or a restriction of gene flow between populations consecutive to their

introduction. Answering such a question would require an extended sampling all over

the native area.

Outlier analysis revealed 92 loci with high posterior probabilities of being under positive

selection between European and Vietnamese populations. When possible, the PCR

amplification of the outlier loci, using a set of representative individuals, always

confirmed a shift of insertion frequencies toward either the European or the Vietnamese

sampling sites. This suggests that in spite of a reduced coverage, introduced by

sampling in the dataset, the scored insertion polymorphisms are reliable. In addition, our

method of analysis is likely to be conservative: the Bayescan outliers were selected for

their consistency with a significant $F_{CT}$ between European temperate and Vietnamese

tropical populations, which avoid retaining outliers that we were not looking for, for

example those due to a population specific event.

We were able to assign a unique position for 21 of the outlier loci on the *Ae. albopictus*

genome. As expected by the *Ae. albopictus* genomic composition (Goubert *et al.* 2015;

Dritsou *et al.* 2015; Chen *et al.* 2015), an important part of the other outlier loci were

located in repeated regions (44,6% of total), despite our efforts to remove *a priori* loci

occurring in known transposable elements. Since the *Ae albopictus* genome publication

is very recent, no gene set or other genome annotation are currently available.  We thus

took advantage of the *Ae. albopictus* transcriptome data (Poelchau *et al.* 2013a) to

annotate regions surrounding the detected ouliers. We found six outliers located in

contigs which also harbor genes differentially expressed between individuals induced for

diapause and controls (Poelchau *et al.* 2013b), two of them, being located either in an

intron (RTE4_7442) or within 3kb (RTE4_17015) of these candidate genes. It is worth

mentioning that these two genes belong to the same functional group (GO:0005576

extracellular region). Diapause is a critical developmental stage found only in temperate

populations of the Asian tiger mosquito. Interestingly, this functional pathway has been

shown to benefit from fast adjustments thanks to local adaptation. For instance,

Urbanski *et al.* (2012) showed that invasive American populations originating from

Japan have rapidly evolved a new adaptive clinal response to diapause induction,

independent from that observed in the native area. Thus, adaption in the temperate

regions could have led to several selective sweeps on gene or regulatory sequences

involved in this critical pathway, allowing the settlement of the mosquito in new

temperate areas.

Interestingly, and as it is the case for these six outliers, we found significantly more

outlier loci with a high frequency in Europe and low frequency in Vietnam than the

opposite pattern. This was unexpected regarding our initial assumptions: a favored allele

selected in one or another environment has *a priori* no reason to be more often

associated with the presence or the absence of a TE insertion at linked sites. However,

we found that the majority of the sequenced TE insertions segregates at low frequencies

(around 10% of all individuals). When considering the linked region of one polymorphic

TE insertion, if a favorable mutation appears in an individual where the insertion is

absent, the increase of frequency of this "absence" haplotype will thus, most of the time,

have a modest effect on the genetic differentiation at this marker, since it is already

segregating at high frequency. By contrast, if a favorable mutation appears in a TE

"presence" haplotype, the increase in frequency of the linked TE insertion would lead to

high differentiation ($F_{CT}$) values. In absence of an alternative explanation, our outlier loci

could thus indicate in which subset of populations the adaptive mutation occurred, and

in the present case, this would have happened more frequently in the temperate

populations.

Two scenarios, not mutually exclusive, could be invoked in the light of our data. A simple

case would be a direct adaptive evolution in European invasive population that

originated from tropical regions of the native area. A second hypothesis, could be that

invasive temperate populations came from Northernmost territories of the native area

such as northern China or Japan where *Ae. albopictus* populations are already cold-

adapted. It would be thus interesting to know whether the observed signature of

selection results from more "ancient" adaptations in the native area, or if it originates

from more recent fine tuning of cold-related traits in the invasive areas. A recent study

(Porretta *et al.* 2012) suggested, using new variable *COI* mtDNA sequences and

historical species range modeling, that Northern areas of the native range of *Ae.*

*albopictus* would be the latest to have been colonized after a range expansion from

Southern refugia following the last glacial around 21,000 years ago. The authors

suggested that *Ae. albopictus* may have followed the human populations during their

expansion from South to North in this area, that began approximately 15,000 years ago.

Thus wherever the origin of the invasive individuals sampled in Europe, it is likely that

they are representatives of populations that had recently undergone a shift of selective

pressure from tropical to temperate climatic conditions. This could explain why so many

outliers are associated with high insertion frequency in Europe, and that candidate

genes in the diapause pathway are found in the neighborhood of some of these outliers.

An easy way to distinguish between these possibilities would be to search if the same

outlier insertions are present in several temperate populations from the native area.

It is important to note that the results presented here are only restricted to a subset of

the Asian tiger mosquito populations located in temperate and tropical environments. It

is thus probable that some of the outliers detected could be specific to this particular

comparison and do not reflect the global pattern of differentiation between tropical and

temperate populations. Research of the same outliers between other tropical and

temperate populations from the native and non-native areas would be extremely

valuable to extrapolate our results at a larger scale. Should the same outlier insertions

be found at high frequencies in temperate locations – such as in USA, Japan or China –,

extended investigations about the origin of invasive populations would help clarify if

those similarities are due to an ancestral sweep or parallel sweeps that occurred

independently in several populations. This study already provides for some candidate

loci a set of functional primers that could be directly used to answer this question in any

DNA sample of *Ae. albopictus*.

We report here the first leads supporting adaptive evolution at the molecular level in the

Asian tiger mosquito. Progress in the annotation of published genomes, and the looming

availability of supplementary genomic resources will allow gaining the most from these

results. We hope that this work will contribute to unravel the implication of adaptive

processes during the invasion of disease vectors.

## Acknowledgements

We thank Van Tran-Van, Christophe Bellet, Grégory Lambert, Huynh Kim Ly Khanh and

Trang Huynh who made possible and contributed to the samplings in France and

Vietnam. We are grateful to Valèria Romero Soriano and her family for their help during

sampling in Sant Cugat dèl Vallès. Library construction and sequencing was made in

collaboration with Clémence Genthon and Olivier Bouchez. We thank Manon Vigneron

for PCR validation experiments. We also thank Rita Rebollo who provided insightful

comments and English revision of the manuscript. This work was performed using the

computing facilities of the CC LBBE/PRABI. C.G. received a grant from the French

Ministry of Superior Education. This work was supported by the Centre National de la

Recherche Scientifique, the Institut Universitaire de France, and preliminary

experiments benefited from of grant of the Federation de Recherche 41 "Bio-

Environnement et Santé". ". Funding for mosquito sampling in Vietnam was provided by

grants from EC2CO CNRS and occurred within the framework of GDRI "Biodiversity and

Infectious Diseases in Southeast Asia". Original maps used to describe sampling were

freely downloaded from http://d-maps.com/carte.php?num_car=4719&lang=en and

http://d-maps.com/carte.php?num_car=708&lang=en

# References

487
Akkouche A, Rebollo R, Burlet N *et al.* (2012) tirant, a newly discovered active
endogenous retrovirus in Drosophila simulans. *Journal of virology*, **86**, 3675–81.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment
search tool. *Journal of molecular biology*, **215**, 403–10.

Becker N, Geier M, Balczun C *et al.* (2013) Repeated introduction of Aedes albopictus
into Germany, July to October 2012. *Parasitology research*, **112**, 1787–90.

Biedler J, Qi Y, Holligan D *et al.* (2003) Transposable element (TE) display and rapid
detection of TE insertion polymorphism in the Anopheles gambiae species complex.
*Insect Molecular Biology*, **12**, 211–216.

Birungi J, Munstermann LE (2002) Genetic Structure of Aedes albopictus (Diptera:
Culicidae) Populations Based on Mitochondrial ND5 Sequences: Evidence for an

Independent Invasion into Brazil and United States. *Annals of the Entomological*
*Society of America*, **95**, 125–132.

Black WC, Ferrari JA, Sprengert D (1988) Breeding structure of a colonising species :
Aedes albopictus ( Skuse ) in the United States. , **60**.

Bock DG, Caseys C, Cousens RD *et al.* (2015) What we still don't know about invasion
genetics. *Molecular Ecology*, n/a–n/a.

Bonin A, Paris M, Després L *et al.* (2008) A MITE-based genotyping method to reveal
hundreds of DNA polymorphisms in an animal genome after a few generations of
artificial selection. *BMC genomics*, **9**, 459.

Bonin A, Paris M, Tetreau G, David J-P, Després L (2009) Candidate genes revealed by
a genome scan for mosquito resistance to a bacterial insecticide: sequence and
gene expression variations. *BMC genomics*, **10**, 551.

Bonizzoni M, Gasperi G, Chen X, James AA (2013) The invasive mosquito species
Aedes albopictus: current knowledge and future perspectives. *Trends in*
*parasitology*, **29**, 460–468.

Boulesteix M, Simard F, Antonio-Nkondjio C *et al.* (2007) Insertion polymorphism of
transposable elements and population structure of Anopheles gambiae M and S
molecular forms in Cameroon. *Molecular ecology*, **16**, 441–452.

Carnelossi EAG, Lerat E, Henri H *et al.* (2014) Specific activation of an I-like element in
Drosophila interspecific hybrids. *Genome biology and evolution*, **6**, 1806–17.

Casacuberta E, González J (2013) The impact of transposable elements in
environmental adaptation. *Molecular ecology*, 1503–1517.

Chen X-G, Jiang X, Gu J *et al.* (2015) Genome sequence of the Asian Tiger mosquito,
Aedes albopictus, reveals insights into its biology, genetics, and evolution.
*Proceedings of the National Academy of Sciences of the United States of America*,
**112**, E5907–5915.

Chouin-Carneiro T, Vega-Rua A, Vazeille M *et al.* (2016) Differential Susceptibilies of
Aedes aegypti and Aedes albopictus from the Americas to Zika Virus (MJ Turell,
Ed,). *PLOS Neglected Tropical Diseases*, **10**, e0004543.

Colautti RI, Lau JA (2015) Contemporary evolution during invasion: evidence for
differentiation, natural selection, and local adaptation. *Molecular Ecology*, **24**, 1999–
2017.

Davey JW, Cezard T, Fuentes-Utrilla P *et al.* (2012) Special features of RAD
Sequencing data: implications for genotyping. *Molecular ecology*.

Dray S, Dufour A-B (2007) The ade4 Package: Implementing the Duality Diagram for
Ecologists. *Journal of Statistical Software*, **22**, 1–20.

Dritsou V, Topalis P, Windbichler N *et al.* (2015) A draft genome sequence of an
invasive mosquito: an Italian Aedes albopictus. *Pathogens and global health*,
2047773215Y0000000031.

Esnault C, Boulesteix M, Duchemin JB *et al.* (2008) High genetic differentiation between
the M and S molecular forms of Anopheles gambiae in Africa. (P Awadalla, Ed,).
*PloS one*, **3**, e1968.

Excoffier L, Lischer HEL (2010) Arlequin suite ver 3.5: a new series of programs to
perform population genetics analyses under Linux and Windows. *Molecular ecology*
*resources*, **10**, 564–7.

Foll M, Gaggiotti O (2008) A genome-scan method to identify selected loci appropriate
for both dominant and codominant markers: a Bayesian perspective. *Genetics*, **180**,
977–93.

Goubert C, Modolo L, Vieira C *et al.* (2015) De novo assembly and annotation of the
Asian tiger mosquito (Aedes albopictus) repeatome with dnaPipeTE from raw
genomic reads and comparative analysis with the yellow fever mosquito (Aedes
aegypti). *Genome Biology and Evolution*, evv050–.

Goubert C, Minard G, Vieira C, Boulesteix M (2016) Population genetics of the Asian
tiger mosquito *Aedes albopictus*, an invasive vector of human diseases. *Heredity*,
10.1038/hdy.2016.35.

Grard G, Caron M, Mombo IM *et al.* (2014) Zika Virus in Gabon (Central Africa) – 2007:
A New Threat from Aedes albopictus? (R Charrel, Ed,). *PLoS Neglected Tropical*
*Diseases*, **8**, e2681.

Gupta S, Preet S (2014) Genetic differentiation of invasive Aedes albopictus by RAPD-
PCR: Implications for effective vector control. *Parasitology Research*, **113**, 2137–
2142.

Handley L-J, Estoup A, Evans DM *et al.* (2011) Ecological genetics of invasive alien
species. *BioControl*, **56**, 409–428.

Hanson SM, Craig GB (1994) Cold Acclimation, Diapause, and Geographic Origin Affect
Cold Hardiness in Eggs of Aedes albopictus (Diptera: Culicidae). *Journal of Medical*
*Entomology*, **31**, 192–201.

Hawley WA (1988) The biology of Aedes albopictus. *Journal of the American Mosquito*
*Control Association. Supplement*, **1**, 1–39.

Hawley W, Reiter P, Copeland R, Pumpuni C, Craig G (1987) Aedes albopictus in North
America: probable introduction in used tires from northern Asia. *Science*, **236**,
1114–1116.

Kambhampati S, Black WC, Rai KS (1991) Geographic origin of the US and Brazilian
Aedes albopictus inferred from allozyme analysis. *Heredity*, **67 ( Pt 1)**, 85–93.

Lande R (2015) Evolution of phenotypic plasticity in colonizing species. *Molecular*
*ecology*.

Li W, Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of
protein or nucleotide sequences. *Bioinformatics (Oxford, England)*, **22**, 1658–9.

Manni M, Gomulski LM, Aketarawong N *et al.* (2015) Molecular markers for analyses of
intraspecific genetic diversity in the Asian Tiger mosquito, Aedes albopictus.
*Parasites & vectors*, **8**, 188.

Marcondes CB, Ximenes M de FF de M (2015) Zika virus in Brazil and the danger of
infestation by Aedes (Stegomyia) mosquitoes. *Revista da Sociedade Brasileira de*
*Medicina Tropical*.

Martin M (2011) Cutadapt removes adapter sequences from high-throughput
sequencing reads. *EMBnet.journal*, **17**, 10.

Miller MR, Dunham JP, Amores A, Cresko WA, Johnson EA (2007) Rapid and cost-
effective polymorphism identification and genotyping using restriction site
associated DNA (RAD) markers. *Genome research*, **17**, 240–8.

Minard G, Tran F-H, Tran-van V *et al.* (2015) French invasive Asian tiger mosquito
populations harbor reduced bacterial microbiota and genetic diversity compared to
Vietnamese autochthonous relatives. *Frontiers in Microbiology*, **6**.

Modolo L, Lerat E (2015) UrQt: an efficient software for the Unsupervised Quality
trimming of NGS data. *BMC Bioinformatics*, **16**, 137.

Monden Y, Yamamoto A, Shindo A, Tahara M (2014) Efficient DNA fingerprinting based
on the targeted sequencing of active retrotransposon insertion sites using a bench-
top high-throughput sequencing platform. *DNA research : an international journal for*
*rapid publication of reports on genes and genomes*, **21**, 491–8.

Munroe DJ, Haas M, Bric E *et al.* (1994) IRE-bubble PCR: a rapid method for efficient
and representative amplification of human genomic DNA sequences from complex
sources. *Genomics*, **19**, 506–14.

Paupy C, Delatte H, Bagny L, Corbel V, Fontenille D (2009) Aedes albopictus, an
arbovirus vector: from the darkness to the light. *Microbes and infection / Institut*
*Pasteur*, **11**, 1177–85.

Peischl S, Excoffier L (2015) Expansion load: recessive mutations and the role of
standing genetic variation. *Molecular ecology*.

Poelchau MF, Reynolds J a., Denlinger DL, Elsik CG, Armbruster P a. (2013a)
Transcriptome sequencing as a platform to elucidate molecular components of the
diapause response in the Asian tiger mosquito, Aedes albopictus. *Physiological*
*entomology*, **38**, 173–181.

Poelchau MF, Reynolds J a, Elsik CG, Denlinger DL, Armbruster P a (2013b) Deep
sequencing reveals complex mechanisms of diapause preparation in the invasive
mosquito, Aedes albopictus. *Proceedings. Biological sciences / The Royal Society*,
**280**, 20130143.

Porretta D, Mastrantonio V, Bellini R, Somboon P, Urbanelli S (2012) Glacial history of a
modern invader: phylogeography and species distribution modelling of the Asian
tiger mosquito Aedes albopictus. *PloS one*, **7**, e44515.

Roy AM, Carroll ML, Kass DH *et al.* (1999) Recently integrated human Alu repeats:
finding needles in the haystack. *Genetica*, **107**, 149–161.

Santolamazza F, Mancini E, Simard F *et al.* (2008) Insertion polymorphisms of SINE200
retrotransposons within speciation islands of Anopheles gambiae molecular forms.
*Malaria journal*, **7**, 163.

Stapley J, Santure AW, Dennis SR (2015) Transposable elements as agents of rapid
adaptation may explain the genetic paradox of invasive species. *Molecular ecology*,
**24**, 2241–52.

Takumi K, Scholte E-J, Braks M *et al.* (2009) Introduction, scenarios for establishment
and seasonal activity of Aedes albopictus in The Netherlands. *Vector borne and*
*zoonotic diseases (Larchmont, N.Y.)*, **9**, 191–6.

Urbanelli S, Bellini R, Carrieri M, Sallicandro P, Celli G (2000) Population structure of
Aedes albopictus (Skuse): the mosquito which is colonizing Mediterranean
countries. *Heredity*, **84 ( Pt 3)**, 331–337.

Urbanski JM, Benoit JB, Michaud MR, Denlinger DL, Armbruster P (2010) The
molecular physiology of increased egg desiccation resistance during diapause in
the invasive mosquito, Aedes albopictus. *Proceedings. Biological sciences / The*
*Royal Society*, **277**, 2683–92.

Urbanski J, Mogi M, O'Donnell D *et al.* (2012) Rapid adaptive evolution of photoperiodic
response during invasion and range expansion across a climatic gradient. *The*
*American naturalist*, **179**, 490–500.

Zhong D, Lo E, Hu R *et al.* (2013) Genetic analysis of invasive Aedes albopictus
populations in Los Angeles County, California and its potential public health impact.
*PloS one*, **8**, e68586.

# 640 Data Accessibility

641 Paired-end raw sequences are available through SRA at NCBI under SRP070185

642    (Bioproject PRJNA312147)

643    Final presence/absence matrices (including replicates) are available at Dryad

644    (doi:10.5061/dryad.9p925) at http://datadryad.org/review?doi=doi:10.5061/dryad.9p925

## 645    Author contributions

646    CG, CV and MB conceived the experiments and conducted the analyses. CG and HH

647    developed and performed the molecular experiments. GM, CVM and PM conducted the

648    sampling in France and Vietnam. All authors contributed to the final version of the

649    manuscript.

## Tables and Figures

**Figure 1.** Sampling sites (with abbreviations) of *Ae. albopictus* in Europe and Vietnam. Red numbers correspond to the total number of individual sampled. Supporting information about samples are available in Table S1.

**Figure 2. Principal Coordinate Analyses (PCoAs).** Projection of individuals over the three first principal coordinates (PC) of PCoAs for each of the 5 TE families and for the first replicate (M1, see Material and Methods). Proportion of inertia represented by each axes is noted in %. circles: European populations; triangles: Vietnamese populations. Results for other sampling replicate can be found in Figure S4.

661 **Table 1. Analyses of Molecular Variance (AMOVAs).** Results for the three replicates
662 (M1, M2, M3) of read sampling for the five TE families (IL1, L2B, RTE5, RTE4, Lian1).
663 Values are given in percentage of the total genetic variance

| | IL1 | L2B | RTE5 | RTE4 | Lian1 |
|---|---|---|---|---|---|
| Among groups | [0.59-0.70] | [1.22-1.29] | [1.08-1.10] | [1.97-2.04] | [0.67-0.74] |
| Among populations within groups | [5.15-5.37] | [3.58-3.63] | [3.36-3.40] | [6.67-6.78] | [4.47-4.56] |
| Within populations | [94.04-94.16] | [95.08-95.18] | [95.51-95.55] | [91.18-91.30] | [94.77-94.81] |

intervals reports min and max values among the 3 sampling replicates

**Figure 3. Insertion frequencies in Europe and Vietnam for the 92 outlier loci**. Bars
represent the median value from the three reads sampling replicates and dots the
values from the other replicates (if outlier found in replicate). Colors correspond to each
of the 5 TE families.

**Figure 4. Comparison of outliers frequencies with randomly selected loci.** Insertion frequencies of 92 randomly chosen loci among those having the same minimum insertion frequency (≥20 individuals) as outliers compared to the 92 outlier loci. Random loci were taken from the first replicate (M1) and values for outliers are median values obtained among the three replicates. Non-overlapping notches indicate a significant difference between the true medians (thick dark horizontal bars).