Induction of Kanizsa contours requires awareness

Theodora Banica[1] & Dietrich Samuel Schwarzkopf[1,2] *

1. Experimental Psychology, University College London, 26 Bedford Way, London, UK
2. Institute of Cognitive Neuroscience, University College London, 17 Queen Square, London, UK

* To whom all correspondence should be directed: s.schwarzkopf@ucl.ac.uk

Abstract

It remains unknown to what extent the human visual system interprets information about complex scenes without conscious analysis. Here we used visual masking techniques to assess whether illusory contours (Kanizsa shapes) are perceived when the inducing context creating this illusion does not reach awareness. In experiment 1 we tested perception directly by having participants discriminate the orientation of an illusory contour. In experiment 2, we probed perception indirectly by using Kanizsa shapes to prime performance on a visual search task. In experiment 3, we exploited the fact that the presence of an illusory contour enhances performance on a spatial localization task. Moreover, in experiment 3 we also used a different masking method to rule out the effect of stimulus duration. Our results suggest that participants do not perceive illusory contours when they are unaware of the inducing context. While perceptual grouping and surface segmentation may occur without awareness of the stimuli, the induction of the phenomenal percept of an illusory shape does not. This is consistent with theories of a multistage, recurrent process of perceptual integration. Our findings thus challenge some reports, including those from neurophysiological experiments in anaesthetized animals. We discuss the importance to test the presence of the phenomenal percept directly with appropriate methods.

Keywords

Illusory contours, modal completion, awareness, masking

Introduction

What role does conscious processing of the environment fulfill and how much processing occurs in the absence of awareness? It is self-evident that much of the internal bodily functions and controlling learned motor behaviors, like walking or driving, operate mostly without awareness. But for processing by the classical senses, like vision, there have been widely discrepant findings on how much stimulus processing can occur and how much it affects decision-making when the subject is unaware of the stimulus. The approach with which unconscious stimulus processing is studied has also been subject to controversy [1].

Several experiments suggest that the effect of a contextual stimulus on a target, such as in the tilt illusion or adaptation effects, persists even when participants are unaware of the presented contextual information [2–5]. In recent years, the use of continuous flash suppression (CFS), in which a dynamic, high-contrast stimulus is presented to one eye to suppress the stimulus in the other eye from awareness, has become a popular way to probe unconscious stimulus processing [6]. Using this procedure it has been claimed that the perception of physical facial attributes [7], the complex analysis of naturalistic scenes [8], and even linguistic processing and arithmetic can be performed without awareness [9].

Neuroimaging experiments showed that while both simple and more complex stimuli have a neural signature in the visual cortex under masking conditions [10–12], the encoding of unconscious stimuli appears to be qualitatively different. Not only is the overall response to unconscious visual stimuli weaker [11] but coupling between different stages in the visual processing hierarchy is also reduced [13] and the information content differs [12,14,15]. In particular, the response to these stimuli is more variable, and also localized in more posterior regions than to conscious stimuli [14].

These findings corroborate behavioral experiments suggesting that unconscious stimulus representations are more variable [16]. Further, it is consistent with our proposal that only simple positional or geometric information may be processed in the absence of awareness, but that more complex abstraction and perceptual integration requires consciousness [17]. Specifically, we used shape stimuli that were either defined by the position or the orientation of simple image elements. We demonstrated that when such stimuli were rendered invisible using fast counter-phase flicker (at 120Hz) they could speed up performance on a shape discrimination task of conscious stimuli. Critically, this priming effect was only present for stimuli sharing the *same* positions, or – if oriented elements were used for priming – the positions along the path implied by the elements. We observed no priming by

invisible primes if the test stimulus was smaller than the prime stimulus. This
suggests that invisible priming operated locally, possibly in retinotopic space in early
visual cortex, but no abstract integration of individual elements into the concept of a
shape occurred without awareness.

We further tested whether two brightness illusions manifest when the inducing
context is rendered invisible by means of CFS [18]. We found that masking the
context (a smooth gradient in luminance) had little impact on simultaneous
brightness contrast of two targets that were unmasked. In a stark dissociation,
participants could not discriminate the orientation of an illusory contour, defined by
a Kanizsa triangle, when the inducing context (the 'Pacman' shapes whose open
segments define the corners of the triangle) were masked selectively by CFS. This
could indicate that the generation of the illusory contour percept occurs at a later
stage of visual processing than simultaneous brightness contrast, either in terms of
the visual hierarchy or in the latency of processing. These findings were also
consistent with previous reports that illusory contours are not perceived when the
inducers are suppressed from awareness during binocular rivalry [19].

These findings challenge some previous reports, using psychophysical tests in
healthy volunteers [20] and even neurophysiological experiments in anesthetized
animals [21] that suggested that illusory contours could be formed in the absence of
awareness. However, none of these previous studies specifically tested whether the
phenomenological percept was indeed formed in these conditions. Similarly,
neuropsychological studies in neglect patients [22–25] suggested that illusory
contour processing occurs when part of the inducing context is placed in the blind
hemifield. However, this does not conclusively support the assertion that contours
are formed in the absence of any contextual awareness. The absence of illusory
contours under dichoptic masking [18,19] therefore highlights that experiments on
Kanizsa stimuli must use appropriate methods and should be interpreted with
caution.

However, two issues confound the interpretation of these findings. First, there is
evidence that illusory contours are processed by binocular neurons in early visual
cortex [26–29]. It does in fact seem unsurprising that the mechanisms inducing
illusory contours at least partially overlap with those for segmenting surfaces in
depth [30]. When retinal disparity implies that inducers are at different depths from
the background, the visual system not only produces the percept of illusory contours
but the surface bounded by illusory contours is also perceived in stereoscopic depth
[27]. Thus when binocular processing is disrupted or overwhelmed by a dichoptic
mask or binocular rivalry, the illusory contour percept is also broken. Another recent

study further complicated this situation by finding that Kanizsa shapes broke through
CFS faster than control stimuli [31]. Leaving aside conceptual issues with the time-to-
emergence paradigm, it is imperative that the dependence of illusory contours on
awareness must be confirmed using masking methods other than CFS.
122
A second confound with these previous studies [18,19] is that even if participants
perceived an illusory contour, this percept was far less salient than real luminance
contours, as it may have been obscured by the dominant masking stimulus. The
addition of a simple control condition could remedy this problem: One experimental
stimulus should be a *real contour* defined by a subtle luminance contrast that mimics
the illusory contour percept as closely as possible. If participants can detect and
discriminate this stimulus but are unable to do so for the illusory contour condition,
this indicates that the *illusory percept* is indeed disrupted specifically, rather than the
more general ability to detect subtle stimuli.
132
Here we carried out three experiments to address these questions. In experiment 1,
we used a similar design as in our previous study [18] by asking participants to
discriminate the orientation of a Kanizsa triangle. However, instead of CFS we
employed a temporal masking method to render the inducers invisible. Moreover,
we included a control condition in which a real, luminance-defined contour was
present. Previous experiments used illusory contours to capture attention in a visual
search task [32] or for priming [33,34]. In experiment 2, we therefore further tested
if Kanizsa triangles rendered invisible using a similar masking procedure could be
used as primes in a visual search task on a subsequent, visible stimulus array. Finally,
because both these masking methods rely on very brief stimulus durations, in
experiment 3 we presented long (500ms) stimuli rendered invisible by means of fast
counter-phase flicker [3,17]. This is critical because the formation of illusory
contours arise comparably slowly [35–37] and thus may be disrupted by the fast
temporal masking techniques in our first two experiments. Previous research
demonstrated that the presence of illusory contours boosts participants' ability to
discriminate the position of a tiny target [19,38] proving a more objective measure
of whether the participant in fact perceives an illusory contour. We therefore
measured the ability of a group of participants, who were well trained at
psychophysical tasks, to discriminate the position of a dot target for Kanizsa and
control stimuli presented with or without masking.
153
154 **Materials and Methods**
155
156 All three experiments were carried out at the UCL Department of Experimental
157 Psychology. Procedures adhered to the Declaration of Helsinki. Ethical approval for

158 this study was obtained from the UCL Research Ethics Committee and all participants
159 gave written, informed consent.
160
161 Participants in experiments 1 and 2 were recruited among the UCL student
162 population. In experiment 3 we recruited participants who were familiar with
163 psychophysical tasks. All Participants had normal or corrected-to-normal visual
164 acuity.
165
166 All experiments were conducted in a dark, sound-attenuated room. For the duration
167 of the experimental sessions, participants were asked to stabilize their head on a
168 chin rest located at a fixed distance of 48cm from the stimulus presentation screen
169 where stimuli were presented to them binocularly.
170
171 Stimuli were generated by a computer and presented on a 22-inch Samsung
172 SM2233RZ LCD monitor at a resolution of 1680 x 1050 pixels. Screen refresh rate in
173 experiments 1 and 2 was set to 60Hz. In experiment 3 it was set to 120Hz. The
174 experiment was controlled and behavioral responses were recorded using MATLAB
175 (The Mathworks, Inc.) and Psychtoolbox 3 [39] using a standard keyboard.
176
177 Experiment 1
178
179 The experiment comprised two tasks: the first, henceforth called 'Kanizsa' task,
180 investigated whether participants can perceive illusory contours without awareness
181 of their inducers. The second task, the 'Visibility' task, assessed the effectiveness of
182 the masking technique directly.
183
184 *Study design*
185
186 Like our earlier experiment using continuous flash suppression, this experiment
187 aimed to measure the perception of illusory contours in a direct manner. We
188 implemented a 2x3 design with visibility (invisible, visible) and type of stimulus (real,
189 illusory, control) as within subject factors.
190
191 Every participant completed two different tasks. Each task comprised 25 blocks of
192 trials, where one block consisted of 24 trials in the Kanizsa task and 8 trials in the
193 visibility task. Across one task, each condition appeared 100 times. The visibility task
194 only comprised the visible and invisible control conditions.
195
196 Participants made behavioral responses in a forced-choice design by button press
197 (left or right arrow) on a standard computer keyboard. Each trial required either a
198 left or a right response and each response type appeared twice per block for each

199 condition. Conditions were selected pseudo-randomly for every trial but were
200 counterbalanced over each block.
201
202 *Participants*
203
204 Seventeen (13 female; age range: 18-29, mean age: 23.8±2.5) normal, healthy
205 participants took part in the experiment. An additional two participants were tested
206 but they failed to discriminate the real luminance contour under masking conditions
207 and were therefore excluded from further analysis (see more details below). All
208 participants were unaware of the experimental hypothesis.
209
210 *Stimuli*
211
Stimuli were created by placing four discs (inducer elements, diameter=2.2°) in the
configuration of a square (width=4.3°). The configuration of these discs was centered
on fixation as both standard edge type and a number of line-end inducing elements
have comparable efficacy in the clarity with which illusory contours are perceived by
participants [40,41]. Here, the discs were defined by partial concentric circles. Each
of the four lines forming the circles had a width of approximately 0.07° with a
luminance of 0.6 cd/m^2. The positioning of the gap in the circles gave rise to the
percept of a Kanizsa triangle (Figure 1A). Thus, a number of line-end inducing
elements gave rise to perception of illusory contours [42].
221
We also included a real luminance condition. Here, the stimuli did not contain any
discs but instead there was a triangle defined by a real but subtle luminance contrast
at the exact location where the Kanizsa triangle would be perceived in the illusory
condition (Figure 1B). We reasoned that if participants were unable to discriminate
the orientation of this subtle luminance edge when the inducers were masked, this
implied that they would be unable to detect any illusory contour that could have
formed without awareness of the inducers. Therefore, we removed two participants
whose discrimination performance for this condition was at chance levels from any
further analyses.
231
232 Control stimuli did not form any triangle and were created by altering the
233 orientations of the inducers by a systematic rotation of 180º (Figure 1C). These
234 stimuli were useful in that they provided information on whether participants could
235 indeed perceive the triangle contours or whether they adopted a strategy of
236 matching the inducer orientation to feedback.
237
The background had a luminance of 76.6 cd/m^2 and real triangles were defined by a
subtly greater luminance of 86.9 cd/m^2. The mask was only used in the invisible

conditions and was designed as a square configuration of four black discs (0.6
cd/m^2), each containing four white concentric circles (230 cd/m^2). It was wide
enough to cover all four Pacmen in the illusory and control conditions (Figure 1D).
The masking technique consisted of a sequence of three frames which was repeated
three consecutive times: At first the mask appeared on the screen, followed by the
appearance of the stimulus and a final blank screen. In the visible conditions, a blank
grey screen replaced the mask frame.

248
Figure 1. Illustration of the stimuli. A) (i) Kanizsa triangle pointing right (ii) Kanizsa triangle
pointing left. B) (i) Real triangle pointing right (ii) Real triangle pointing left. C) Control stimuli

were created by a systematic 180° rotation of the individual Pacmen. D) Illustration of the
masking procedure in the invisible conditions: A mask-stimulus-blank screen frame sequence
was repeated three times. In the visible conditions a blank screen replaced the mask.

Procedure

Kanizsa task: At first, participants were instructed that they would see a triangle
appear on the screen. They were asked to judge whether its hypotenuse was tilted
clockwise or counter-clockwise from vertical by pressing the corresponding response
key. We explained this task to them as a decision whether the right angle of the
triangle was pointing to the left or to the right but they were told explicitly to *judge*
the contours of the triangle, in particular the long hypotenuse extending through the
center of the stimulus display. Because we wished to keep the contrast between real
triangle and the background as low as possible, we trained participants on the visible
and invisible real triangle condition until they were able to detect them correctly.

Participants were instructed to fixate a small black dot (0.2° wide) that was present
in the center of the screen throughout the experiment. On each trial, the fixation dot
was displayed alone for 500ms. This was followed by a sequence of three frames
that defined whether the condition was a visible or an invisible one. In the invisible
condition a 300ms mask followed the fixation period. Subsequently, the stimulus
appeared on the screen for one frame of approx. 16.7 milliseconds (ms),
immediately followed by a blank screen that was shown for one frame as well. This
mask-stimulus-blank sequence was repeated three times before a second and final
post-stimulus blank screen was presented until participants gave their response.
Figure 2A shows the general paradigm for the experimental procedure.

Visibility task: We further tested whether participants indeed did not consciously
perceive any contextual information of the stimuli in the invisible condition. For this
purpose, the visibility task assessed the effectiveness of the masking technique by
measuring whether participants could consciously discriminate the inducer
elements. Participants were asked to judge whether a right angle was presented in
the left or right bottom inducer. The timing of the trial sequence in this task was
identical to the Kanizsa task, with one exception: only control stimuli were used
(Figure 2B).

286
Figure 2. Illustration of the trial sequence in experiment 1. A) Kanizsa task: Each trial was
composed of five frames: fixation dot, mask, stimulus, blank, and post-stimulus blank. After
the participant's response, the fixation point provided feedback for 100ms (green: correct;
red: incorrect). The duration of each frame is shown on the time-line. Note that in the visible
conditions, a blank screen replaced the mask. B) Visibility task: Trials proceeded in the same
way as in the Kanizsa task except that only control stimuli were being shown and
participants judged whether the right-angle gap was in the bottom left or right inducer
(indicated by the green circle, which was not present in the actual stimuli).

295
296
297 *Data analysis*
298
299 Performance in both tasks was defined in terms of mean proportion of accurate
300 responses in each condition. Initially, we conducted binomial tests at the individual

level to quantify how many participants performed significantly above chance. A condition for any participant's data to be considered in the group analysis of this experiment was that their performance to the invisible real stimulus condition was significantly above chance (0.5). This is because the features necessary to distinguish the contours of the real triangle were not masked and if a participant could not perform the task for this particular stimulus, any test of the perception of illusory contours would be redundant.

One-sample t-tests were carried out at the group level for each condition individually to assess whether the participants' level of performance was significantly above chance (>0.5). In addition to traditional frequentist statistical tests, we also quantified the evidence for or against the hypothesis that participants could discriminate the stimuli by calculating a Bayes Factor using a default Cauchy prior with scaling factor 0.707 for the alternative hypothesis [43]. For conditions with performance near chance levels, this enabled us to also quantify how strongly the evidence supported the null hypothesis that participants were actually guessing.

Experiment 2

This experiment comprised two consecutive tasks as well. The first, again called 'Kanizsa' task, was an indirect test that investigated whether priming with a Kanizsa triangle in the location of the target improves detection of the target. A second task consisted of a 'Visibility' task that directly tested the effectiveness of the masking technique.

Study design

In the context of this experiment, priming refers to the attentional cue provided by the prime shape about the location of the subsequently presented target in one of four possible locations that in turn affords a boost in behavioral performance on an orthogonal discrimination task. In the primed conditions the location of the target was always identical with the location of the prime, whereas in the unprimed conditions the prime shape was replaced by a control stimulus, which looked exactly like the distractors in the other three locations. Therefore, the unprimed condition provided a baseline that allowed the assessment of whether the triangle prime boosted the participants' performance by capturing their attention.

We tested priming with a 2x2x2 design with attentional priming (primed vs. unprimed), the type of priming triangle (real vs. illusory), and prime visibility

(invisible vs. visible) as within subject factors. There were two different prime shapes: a real triangle defined by a subtle luminance contrast or an illusory (Kanizsa) one. The visibility of the cue was manipulated by varying presentation times of the priming array that followed the mask. A short presentation time of 16.7ms was selected for the invisible conditions to render the prime invisible. In the visible conditions, however, the priming array was presented for 300ms, which left enough time for the prime to enter awareness.

Every participant completed two different tasks. Each task comprised 20 blocks of trials, where one block consisted of 32 trials. Across one task, the unprimed visible and invisible conditions appeared for a total of 160 times each, whereas the four primed conditions (real vs illusory and visible vs invisible) occurred 80 times each. Thus, half the block consisted of unprimed trials and the other half of primed ones.

Behavioral responses were given on a forced-choice discrimination task by button press (left or right arrow) on a standard computer keyboard. Each trial required either a left or a right response (see experimental procedure) and each response type appeared twice per block for each condition, except for the unprimed ones where it appeared four times per block. Conditions were selected pseudo-randomly for every trial but were counterbalanced within each block.

Participants

Seventeen (13 female, age range: 21-32 years, mean age 24.7±2.5 years) normal, healthy participants took part in the experiment, including the two authors. All other participants were unaware of the experimental hypothesis.

Stimuli

In half the trials, participants were cued with one of four possible cues. We created shape stimuli by placing four black discs (inducer elements, diameter=1.96°) in the configuration of a square (width=2.8°). In order to produce the illusory shape of a right isosceles triangle sitting on top of the square, a wedge was removed from each inducer element (Pacman). Wedges started at the center of their respective discs and extended to their edges. One wedge measured 90° and was always positioned either in the bottom left or bottom right inducer. The two wedges that formed the apexes of the triangle measured 45°. Finally, the wedge of the fourth inducer element measured 90°. This Pacman did not form part of the triangle and was positioned opposite the hypotenuse of the right isosceles triangle. It always faced the outside of the disc configuration. When the inducers were oriented with the

gaps forming the three corners of a triangle, an illusory light grey Kanizsa triangle was visible on top of the black discs (Figure 3A, i and ii).

Real triangles were identical to the corresponding illusory shapes except that their contour was defined by a luminance contrast with a black line connecting the corners of the triangle (Figure 3A, iii and iv).

The control stimuli were created by altering the orientations of the three Pacmen by a systematic rotation of 180º. This modification broke the link between these inducers and thus no illusory triangle was perceived (Figure 3B). Control stimuli could also be horizontal mirror images of the illustrated examples.

Stimuli in the search array were right isosceles triangles with a contour defined by luminance (Figure 3C, i and ii). Target stimuli were triangles with the right angle pointing down, either to the left or to the right, while in distractors the right angle pointed up (Figure 3C, iii and iv). Participants were asked to detect the target and respond via button press whether its right angle was pointing to the left or to the right.

The mask consisted of a square of black lines connecting four discs, one at each corner. It was wide enough to cover all four Pacmen, as well as the location of the triangle within them (Figure 3D, i). In the 'Visibility' task, there was no search array. Instead, the stimulus array following the prime was a square containing a cross that gave no clue about the location and orientation of the target triangle (Figure 3D, ii).

The luminance of the background was 156 cd/m^2. The luminance of inducer elements, the contour of the real triangle and the black of the mask was 2 cd/m^2. Finally, the grey of the mask in the first task was 58.2 cd/m^2.

Note that although there were differences between stimuli, the individual inducers remained the same and there was no change in the local image properties (i.e. square configuration of the four Pacmen and their position among the other 3 configurations). This enabled an assessment of the different effects of priming with the Kanizsa shape, as compared to the corresponding real triangle and ungrouped Pacmen or distractors in the unprimed conditions.

Procedure

Kanizsa task, indirect test assessing priming:
At first, participants were instructed that they would see an array of four triangles of which only one, the target, had a right angle pointing down and were asked to decide whether this angle was pointing to the left or to the right by pressing the respective response keys.

The 20 experimental blocks were preceded by as many practice blocks as needed in
order to fully familiarize the participants with the task requirements. Participants
could initiate a new block by pressing any button on the keyboard and, if needed,
take small breaks between blocks.

Participants were instructed to fixate a small black dot (0.2° wide) that was present
in the center of the screen throughout the experiment. On each trial, the fixation dot
was displayed alone for 500ms. Subsequently, a mask array was displayed for 100ms
and was immediately followed by one of the priming arrays (see conditions in study
design), which was presented for either 16.7ms or 300ms. This array could contain
one of the two possible prime shapes (real or illusory triangle) or a simple control
stimulus among the other three control stimuli. Immediately after this, a target array
containing one target among three distractors was displayed and remained on the
screen until the participant's response.

The response was self-paced; however, participants were instructed to respond as
quickly as possible. Responses were made with the index (for left) and the middle or
ring finger (for right), depending on how participants felt comfortable. For every
response, the fixation dot provided feedback (100ms) by changing its color (green:
correct; red: incorrect). Then, the next trial started without any delay. Figure 4A
shows the general paradigm for the experimental procedure.

Visibility task, direct test assessing prime discrimination:
The second task assessed the effectiveness of the mask by measuring the conscious
discriminability of the primes in a direct manner. For this purpose, participants were
asked to make the same decision as in the previous task, but this time with respect
to the prime. Thus, instead of discriminating whether the right angle of the target
was pointing to the left or to the right, participants discriminated this aspect of the
prime shape.

Procedure, stimuli and timing of the trial sequence in this task were kept identical to
the previous indirect test, with a single exception: Instead of a target array, an array
comprising new identical mask shapes covered the four stimuli locations (Figure 4B).

475
Figure 4. Illustration of an example trial sequence in experiment 2. A) Kanizsa task: Each
trial was composed of four frames: fixation period, mask, cueing array and target array.
After the participant's response, the fixation point provided feedback for 100ms (green:
correct; red: incorrect). The duration of each frame is shown on the time-line. The duration
of the cueing array was varied between conditions: 16.7ms in the invisible conditions and
300ms in the visible ones. Note that in the unprimed conditions, the cueing array did not
contain any prime shape and was composed exclusively of control stimuli. B) Visibility task:
Trial procedure was identical to the one in the Kanizsa task except for the fourth frame in
which the search array was replaced by an array of four identical mask shapes.
485
486
487
488

Data analysis

Response times (in milliseconds) and performance accuracy (in proportion correct)
were measured for each of the six conditions. To analyze priming effects, response
times were calculated on correct trials only.

For the Kanizsa task, accuracy of responses was less informative, as the target array
was present until the participant's response. We reasoned that priming in the
indirect Kanizsa task would be reflected by faster mean response times to primed
than unprimed conditions. Therefore the priming effect was calculated by
subtracting response times under priming from those without priming.

Priming effects from all participants for all conditions were entered into a two-way
repeated-measures analysis of variance (ANOVA) with prime visibility (visible vs
invisible) and prime stimulus (real vs illusory triangle) as within subject factors. In
addition to this, further comparisons between experimental conditions (visible or
invisible/real or illusory) and control conditions (unprimed) were assessed with
paired t-tests. We also determined the significance of the priming effect by
comparing it against zero with one-sample t-tests.

In the visibility task, the variable of interest was the accuracy of responses as this
allowed us to test whether participants could discriminate the orientation of the
primes. One-sample t-tests were carried out at the group level for each condition
individually in order to assess whether the participants' level of performance was
significantly above chance (>0.5) and paired-tests were used to compare
performance in different experimental conditions. As in experiment 1, for all t-tests
we also calculated the default Bayes Factor in addition to frequentist statistics.

<u>Experiment 3</u>

This experiment only comprised one task that indirectly tested the percept of illusory
contours by measuring participants' threshold on a spatial localization task. Such a
manipulation has been used successfully in previous experiments [19,38]. While it is
an indirect test, the contour aids performance on the spatial localization task. This
provides independent evidence about whether participants perceived any illusory
contours and therefore helps to address confounds with measuring the percept
directly as we did in experiment 1. Pilot experiments using a task directly measuring
the illusory percept (the same task as experiment 1 but with the long-lasting masking
technique employed in experiment 3) could encourage participants to pay close
attention to the masked context and thus reduce the effectiveness of the masking
procedure. This would mean that participants use residual awareness of the inducers

to perform the task instead of actually making a perceptual judgment of the illusory
contour.
532
Moreover, because the masking procedure in experiments 1 and 2 used very brief
presentations of the invisible Kanizsa stimuli (16.7ms) while these were much longer
(300ms) for the visible ones, in experiment 3 we used a different masking procedure:
stimuli were defined by sinusoidal gratings which reversed contrast polarity at
120Hz. This method can effectively render stimuli invisible for prolonged periods so
that participants only perceive a grey screen. Previous research suggests that stimuli
masked in this way are processed in early visual cortex [3]. Moreover, we showed
that stimuli rendered invisible by this method could induce local priming effects on a
shape discrimination task [17].
542
543 *Participants*
544
Five normal, healthy participants (3 female; age range: 24-37; mean age: 30.2)
completed this experiment, including one of the authors (DSS). All participants were
experienced with psychophysical experiments. While the results from a larger
subject base might generalize more to a broad population, we reasoned that only
precise measurements of participants' visual performance would afford
interpretable data. Unlike in the first two experiments, in experiment 3 we
measured how position discrimination thresholds changed between experimental
conditions. Naïve, untrained participants from the general population would be
more likely to contribute noisy data and this might obscure potential subtle
perception of illusory contours in the masked condition.
555
556 *Stimuli*
557
We generated a Kanizsa shape by presenting four Gabor patches (sinusoidal gratings
with wavelength 0.33° visual angle convolved with a Gaussian with standard
deviation 0.8° at 30% contrast) in the locations of the four corners of a square with a
side-length of 8.2°. We turned these patches into Pacmen by setting a right-angled
region of each patch to zero contrast (uniform background grey). This stimulus thus
described an illusory square (Figure 5A). In the control conditions, the Pacmen were
rotated by 180° so that the corners faced outward breaking the illusory percept
(Figure 5B). In the real luminance control, we did not present any Gabor patches but
instead a square region with a subtle luminance contrast (54 cd/m^2) relative to the
background (Figure 5C). The orientation and phase of each Gabor patch was
randomized in each trial. Finally, we created a mask stimulus by overlaying 24 Gabor
patches in the four locations. These patches were presented at 5% contrast and
covered the full range of orientations in equal steps of 15° but their phases were

randomized. This resulted in a patchy pattern without any obvious orientation cue
(Figure 5D). The mask pattern was also generated anew in every trial.
573
574 These stimuli were presented in the center of the screen. The target stimulus was a
575 tiny dark grey dot (diameter: ~0.1°) which could appear somewhere along the
576 horizontal meridian, either near the right or the left vertical boundary of the square.
577
578 The square stimuli were either invisible or visible. In the invisible condition, the
579 gratings reversed contrast polarity at 120Hz. In the visible conditions and the real
580 luminance control, every even-numbered frame was uniform grey (except for the
581 fixation dot).
582

583
Figure 5. Stimuli (A-D) and trial sequence (E) in experiment 3. During the stimulus interval
either an illusory Kanizsa square (A), a control stimulus with rotated inducers (B), or a real
luminance square without any inducers (C) was presented. In visible trials every odd-
numbered video frame at 120Hz contained the stimulus while every even-numbered frame
contained only a blank screen with the blue fixation dot. In invisible trials, both frames
contained the Gabor patches for A and B but their contrast polarity alternated between
frames. For real luminance stimuli the even-numbered frames always contained a blank
screen. D. A mask stimulus preceded and followed each stimulus interval. E. Each trial
started with 500ms of fixation, followed by a 100ms mask, the 500ms stimulus interval,
another 100ms mask, and finally a blank screen without a fixation dot that remained until
participants made their response. Their task was to locate the small target dot in the

stimulus display and respond whether it appeared to the left or the right of the vertical
boundary of the square (here, the correct response is right).

Procedure

Figure 5E depicts the sequence of an invisible Kanizsa trial in experiment 3. Each trial
started with 500ms of a blank grey screen in which only a black fixation dot
(diameter 0.26°) was present. This was followed by a 100ms presentation of the
mask and then 500ms of the stimulus, after which another 100ms mask interval was
presented. Then the screen turned grey and the fixation dot was removed, indicating
that participants could make their behavioral response.

During the stimulus interval the fixation dot was blue instead of black to denote that
this was the task-relevant interval. As described above, the stimuli in this interval
either reversed in contrast polarity at 120Hz (invisible Kanizsa and control
conditions) or the frames were interleaved with blank frames (visible Kanizsa, real
luminance stimuli and control). During the stimulus presentation the small dark
target dot also appeared at some location along the horizontal meridian. Its position
was randomized to be either near the left or the right vertical boundary of the
square region. The dot could either appear inside or outside the square region.

The dependent variable in this experiment was the distance between the target dot
and the boundary of the square region and it was controlled by a 2-down, 1-up
staircase procedure that converged on the threshold distance in each of the five
experimental conditions at which performance was approximately 70.7% correct.
That is, after every consecutive two correct trials the distance would decrease by
one pixel (~0.03°) while after every incorrect trial it increased by one pixel. All five
staircases started at a distance of 15 pixels (0.49°). The minimum and maximum that
they could reach were one pixel and 25 pixels (0.82°), respectively.

Before the actual experiment we showed participants static examples of the Kanizsa
and the control stimuli. Participants were instructed to judge whether the target dot
was left or right of the (imaginary) vertical boundary of the square region. While
showing them still images of the stimuli, we specifically explained to them that this
boundary was defined by the exact center of the Gabor patches (i.e. the corner of
the Pacman's mouth) and that this was identical in both the Kanizsa and the control
conditions. We further informed them that there would be a third condition in which
they should only see a subtly lighter grey square against the background but no
Gabor patches. In order to become acquainted with the task, they then performed 1-
2 blocks of the experiment with only the visible Kanizsa, the control and the real
luminance condition. Finally, the actual experiment would commence. Participants

were informed that the task was largely always the same as the familiarization run,
although during this experiment there would be many more trials in which they
would either only see a light grey square but no Gabor patches or that they might
even only see a grey blank screen. Except for the author, all participants were
unaware of the experimental hypothesis. In debriefing none of the participants
reported seeing any Gabor patches during the invisible condition.
642
643 Participants performed two consecutive runs comprising 500 trials each. These runs
644 were further subdivided into 25 blocks. In every block each of the five conditions
645 appeared 4 times in a pseudo-randomly interleaved order. Blocks were initiated by
646 button-press. During block breaks a message on the screen reminded the participant
647 of the task instructions and of the number of blocks they had already completed.
648
649 *Data analysis*
650
We determined the threshold distance in each of the five experimental conditions by
calculating the mean distance across the final 15 reversals in the staircase
procedure. Then we conducted a group level analysis in which we compared the
average thresholds for conditions across participants. For this we used a two-way
repeated-measures ANOVA with the factors visibility (visible vs invisible) and
stimulus type (illusory vs control stimulus). As there was only one condition with a
real luminance contour, the thresholds for this condition were not included in the
ANOVA. However, we used paired t-tests to compare results for the illusory and
control stimuli directly to the real luminance condition.
660
661 **Results**
662
663 <u>Experiment 1</u>
664
665 In this experiment, 17 participants judged the orientation (left vs right) of a triangle
666 that was either defined by an illusory contour or a subtle real luminance edge. In a
667 third control condition the Pacmen stimuli inducing illusory contours were rotated so
668 that no triangle shape could be perceived.
669
670 *Kanizsa Task*
671
672 Accuracy across participants for discriminating the orientation of the triangle for the
673 six conditions is displayed in Figure 6. We tested whether discrimination
674 performance in each condition was above chance at the group level.
675

In the visible condition, the mean proportion of correct responses for the group was close to ceiling and clearly better than chance both for the real (M=0.97, t(16)=83.5, p< 0.001, BF_{10}>9.8*10^{18}) and the illusory condition (M=0.99, t(16)=120.8, p<0.0001, BF_{10}>2.4*10^{21}). In the control condition, however, the group performed significantly below chance level (M=0.30, t(16)=-3.87, p=0.001, BF_{10}=29.3). There was considerable variability in performance for this condition ranging from 0 to 0.74.

In the invisible condition, only performance on the real triangle condition was significantly above chance level (M=0.71, t(16)=6.4, p<0.001, BF_{10}=2608.7). Performance on both the illusory triangle condition (M=0.51, t(16)=1, p=0.332, BF_{10}=0.385) and the control condition (M=0.50, t(16)=0.2, p=0.859, BF_{10}=0.253) were at chance level. Importantly, performance on the real triangle condition was also significantly greater than for either the illusory contour (t(16)=6.1, p<0.0001, BF_{10}=1475.4) or the control condition (t(16)=5.6, p<0.0001, BF_{10}=688). In contrast, performance for the illusory contour did not differ from the control condition (t(16)=0.7, p=0.525, BF_{10}=0.3).

Visibility Task

In a supplementary control task we asked participants to make a decision directly based on the inducers by reporting which one of the two bottom inducers contained the right angle. This assessed any residual perception of the inducer shapes under masking conditions. In the visible condition, the mean proportion of correct responses across participants was again close to ceiling and far above chance level (M=0.96, t(16)=22.2, p<0.001, BF_{10}>2.9*10^{10}). In contrast, participants' mean performance in the invisible condition did not significantly differ from chance level (M=0.48, t(16)=-1.3, p=0.227, BF_{10}=0.489), suggesting that participants could generally not perceive the inducers under masking conditions. Note, however, that there was only anecdotal support for the null hypothesis (BF_{10}>1/3).

The first experiment thus suggested that if participants were unaware of the inducers because they had been masked, they were unable to judge the orientation of the Kanizsa triangle. This supports the interpretation that illusory contours are not formed under these conditions.

Figure 6. Results of experiment 1. Accuracy for discriminating the orientation of a triangle stimulus in visible or invisible (masked) trials. Each dot represents the performance of an individual participant in each of the conditions. The large symbols and error bars denote the mean ±1 standard error for each condition. Black: real luminance contour. Red: illusory (Kanizsa) contour. Blue: control stimuli.

Experiment 2

In the second experiment we asked whether masked Kanizsa shapes could afford an attentional priming effect. We tested whether a Kanizsa stimulus presented as part of an array of four stimuli could prime participants to the location of a target in a visual search array presented subsequently. The prime could either be a Kanizsa triangle or a triangle defined by a real luminance contour. In the unprimed condition all four stimuli in the priming array were control stimuli without a triangle shape.

Kanizsa Task

Overall participants made very few errors in determining whether the target pointed to the left or to the right. Indeed, the mean accuracy in all conditions was above 0.94. Therefore, for each experimental condition we quantified the effect of priming by the difference in response time with priming subtracted from that without priming (Figure 7A).

Figure 7. Results of experiment 2. Priming effect (response time for unprimed trials minus primed trials) on the Kanizsa task (A) and accuracy for discriminating the inducer location in the Visibility task (B) for visible or invisible (masked) trials. Each dot represents the mean performance of an individual participant in each of the conditions. The large symbols and error bars denote the mean ±1 standard error for each condition. Black: real luminance contour. Red: illusory (Kanizsa) contour. In B) Grey: unprimed trials dummy coded for real contour prime. Orange: unprimed trials dummy coded for illusory contour prime.

We conducted a two-way repeated-measures analysis of variance, with visibility (visible vs invisible) and stimulus (illusory vs real triangle) as within subject factors, to compare the priming effects (response time on unprimed minus primed trials) between conditions. There was a significant main effect of visibility of the prime ($F(1,16)=60.06$, p<0.001), and a significant main effect of stimulus type ($F(1,16)=13.82$, p=0.002). The interaction between visibility and the stimulus type was not significant ($F(1,16=3.52$, p=0.079).

There was a clear difference in the strength of priming between the visible and invisible condition. However, the critical test is whether the priming effect was significant, that is, if it differed from zero (i.e. a paired t-test between primed and unprimed response times). Unsurprisingly, the very pronounced priming effect for visible trials was extremely significant for both conditions (real: M=336ms, $t(16)$=8.2, p<0.001, BF_{10}>4.1*10^4; illusory: M=275ms, $t(16)$=6.9, p<0.001, BF_{10}>5.8*10^3). However, there was also a much more modest but nonetheless very significant priming effect for both stimulus types in the invisible condition when inducers were masked (real: M=43ms, $t(16)$=3, p=0.01, BF_{10}=5.6; illusory: M=36ms, $t(16)$=4.4, p<0.001, BF_{10}=80.8).

Visibility Task

As in experiment 1 we used a direct Visibility test to assess whether participants might have had residual awareness of the priming stimuli under masking. The task

procedure was the same as in the Kanizsa task except that the search array was
replaced with a foil in which all four stimuli were identical and non-informative
about the correct answer. We quantified the accuracy (proportion correct) with
which participants could discriminate the orientation of the priming triangle (Figure
7B).

Unsurprisingly, in the visible condition performance for discriminating the
orientation of a real (M=0.88, t(16)=14.9, $p<0.001$, BF_{10}=9.5*10^7) and an illusory
triangle (M=0.87, t(16)=14, $p<0.001$, BF_{10}=4.0*10^7) was significantly above chance.
Conversely, performance for the unprimed conditions, that is, stimulus arrays that
contained no triangle shape, performance was consistently at chance ("real":
M=0.52, t(16)=1.3, p=0.21, BF_{10}=0.52; "illusory": M=0.53, t(16)=1.3, p=0.219,
BF_{10}=0.5; but note that these two conditions were dummy coded in this case as they
contained the same stimuli).

Critically, performance in the invisible condition showed that discrimination in the
primed with an illusory triangle condition was also significantly above chance (M=
0.55, t(16)=3.24, p=0.005, BF_{10}=9.4), and discrimination of a real triangle showed a
similar result even though it did not reach significance (M=0.53, t(16)=1.9, p=0.075,
BF_{10}=1.1). Performance for the two invisible triangles also did not differ significantly
(t(16)=-0.9, p=0.385, BF_{10}=0.35). Again, as expected both of the unprimed conditions
were at chance level ("real": M=0.51, t(16)=0.6, p=0.534, BF_{10}=0.3; "illusory":
M=0.51, t(16)=0.7, p=0.494, BF_{10}=0.31).

Taken together, the results of the second experiment contradicted those of the first,
as they suggested that Kanizsa triangles rendered invisible by masking could afford
an attentional priming effect in a subsequent visual search task. However, a control
experiment testing participants' ability to discriminate the prime shape orientation
directly suggested that this small priming effect for invisible trials could have been
due to some residual awareness of the masked triangle stimuli.

Experiment 3

In experiment 3 we therefore changed the approach in a number of ways. First,
instead of the brief stimulus presentations used in the first two experiments we used
counter-phase flicker to render stimuli invisible for prolonged periods. As in
experiment 2 we used an indirect measure of illusory contour processing: we took
advantage of previous reports that the presence of an illusory contour stimulus
enhanced participants' thresholds at discriminating the position of a small dot
[19,38]. Here we tested whether this also occurred when the inducers generating
the illusory contour were invisible. Because this task was more challenging than

those in previous experiments, and to rule out that our previous results might have
been due to insufficient practice or familiarity with psychophysical experiments, in
this final experiment we only tested a small group of well-trained psychophysics
participants.
811
Figure 8 plots position discrimination thresholds in the different conditions for all
individual participants and the group averages. Overall, thresholds measured while a
Kanizsa stimulus was presented were significantly lower than those measured for
control stimuli ($F(1,4)=34.6$, p=0.004). There was also a non-significant trend of
lower thresholds during visible than invisible trials ($F(1,4)=7.5$, p=0.052). Importantly,
there was a significant interaction between visibility and stimulus type ($F(1,4)=15.9$,
p=0.016). In the visible condition thresholds measured while an illusory Kanizsa
contour was present were significantly lower ($M=0.08°$, $t(4)=-5.27$, $p=0.006$,
$BF_{10}=9.95$) compared to the control condition without a contour ($M=0.21°$). In
contrast, in the invisible condition the difference in thresholds for illusory contours
($M=0.20°$) was not significantly different from that for the control stimulus ($M=0.21°$,
$t(4)=-0.16$, $p=0.882$, $BF_{10}= 0.4$).
824
Thresholds measured for the real luminance contour were of a similar magnitude as
those measured for the visible illusory contour ($M=0.10°$, $t(4)=-0.92$, $p=0.409$,
$BF_{10}=0.55$). In contrast, thresholds for the real luminance contour were significantly
lower than for invisible Kanizsa stimuli ($t(4)=3.84$, $p=0.019$, $BF_{10}=4.53$). Thresholds
for the real luminance contour were also significantly lower than for either control
stimulus (visible: $t(4)=4.4$, $p=0.012$, $BF_{10}=6.19$; invisible: $t(4)=5.3$, $p=0.006$, $BF_{10}=9.9$).
831

Figure 8. Results of experiment 3. Spatial discrimination thresholds per condition (real luminance vs illusory vs control) on a dot localization task while participants were presented with visible (blue squares), invisible (orange diamonds), or real luminance stimuli (black circles). A-E. Plots for the five individual participants. F. Thresholds averaged across participants ±1 standard error of the mean.

Discussion

In three psychophysical experiments we tested whether the visual system forms illusory (Kanizsa) contours in the absence of awareness of the inducing context. None of the experiments showed any conclusive evidence that illusory contours are perceived when the inducers are masked.

In the first experiment, we directly measured illusory contour perception by asking participants to discriminate the orientation of a Kanizsa triangle. This procedure was akin to our earlier experiments [18] that showed no evidence either that illusory contours are formed when inducers are suppressed from awareness. In those experiments we employed CFS in which a dynamic, high-contrast mask presented to one eye suppresses awareness of a stimulus viewed by the fellow eye. Similarly, previous studies showed that illusory contours are not formed when the inducers are suppressed from awareness during binocular rivalry [19]. However, because previous reports indicate that illusory contours are processed by binocular neurons the use of dichoptic stimuli may simply disrupt their processing.

Therefore here we used normal binocular viewing conditions and a different method to render the inducers invisible. In addition, we included a real luminance contour condition as a baseline check. This helped to rule out another trivial explanation for our earlier findings: The presence of the bright masks could possibly have obscured the detection of the subtle illusory boundary. Our real luminance contour was never masked because the masks only overlapped with the corners of the triangle; therefore, if participants were able to discriminate the subtle real luminance contour they should also be able to do so if an illusory contour were indeed formed. We only included participants for whom discrimination of this real luminance contour was significantly above chance levels. Nonetheless, participants were unable to discriminate the orientation of the illusory contour when inducers were masked. The phenomenological experience of the real luminance and the illusory contour is not perfectly identical (and it obviously is not for the visible conditions). However, the real luminance conditions were very faint while illusory contours tend to be subjectively quite salient. Thus it seems unlikely that participants should be able to detect only the real luminance contour but not the illusory contour.

Experiment 1 also contained control conditions in which the individual inducers had been rotated by 180° and thus no illusory triangle should be formed. This condition was somewhat unnecessary because the dependent variable, accuracy for discriminating the orientation of the triangle, was orthogonal to the condition and for these control stimuli there was no strictly correct answer in this discrimination task. We included this condition as catch trials – participants should be guessing here because there was no triangle to discriminate. This was clearly the case for control stimuli when inducers were invisible. However, for visible control stimuli discrimination performance varied widely from 0-0.74 proportion correct responses and was even significantly below chance level. The "correct" responses in this condition were dummy coded so that the stimulus-response mapping matched that for the equivalent illusory triangle stimuli before rotating the inducers. Therefore one inducer in the top row had a right-angled cutout that pointed inwards (Figure 1) and this would be consistent with the presence of a triangle oriented in the opposite direction (even though no actual triangle should be perceived). Some participants may have adopted this stimulus response mapping while others did not. It would therefore have been better to have all inducers point outwards in this control and randomize their locations. However, our actual results serendipitously support the fact that masking was effective in this experiment: The fact that participants performed at chance level when the control stimuli were masked rules out that they had any residual awareness of the inducers. The results of our explicit Visibility task after the main experiment also corroborated this conclusion.

The results of experiment 1 contradict previous claims that a "ghostly triangle" could
be perceived when the inducers of a Kanizsa triangle were masked [20]. However,
follow-up experiments failed to replicate these experiments but instead suggested
that conscious processing of the inducers precedes awareness of the illusory
contours and that these earlier results were in fact due to residual awareness of the
inducers [44]. Another critical issue with these experiments is that the perception of
illusory contours is based on a Yes or No judgment of whether a triangle shape was
present. Such a task could theoretically be performed based on any residual
awareness of the corners in the inducers. Our task required an explicit orientation
discrimination of the illusory contour. If participants had been able to perform this
task in spite of being unable to perceive the inducers, this would have been more
conclusive evidence that illusory contours are indeed formed when inducers are
masked.

In our second experiment, we used an indirect approach to infer whether illusory
contours were processed without awareness of the inducers. We measured whether
the presentation of a Kanizsa triangle or one with a real luminance edge afforded
participants with a response time advantage on a subsequent visual search task
compared to the situation where no triangle was presented. Previous research has
shown that Kanizsa shapes can capture attention in this manner when visible [32]
and illusory contours can also prime performance on shape or orientation
discrimination tasks [33,34]. As expected, there was a very strong priming effect for
both the real luminance and the illusory triangle when priming stimuli were visible as
the two shapes produce a salient "pop-out" effect. Interestingly, even when the
primes were removed from awareness by forward masking, they nonetheless
produced a weak but significant response time advantage for either stimulus type.
However, our subsequent Visibility task, in which participants judged the orientation
of the priming triangle directly, suggested that masking may not have been entirely
effective because participants performed significantly above chance level even for
the masked stimuli. Therefore, the most parsimonious interpretation of the priming
results is that participants had some residual awareness of the primes and that
Kanizsa contours were probably not processed when inducers were masked.

In both experiments we also included an explicit Visibility task in which we modified
the task to test whether participants had any residual awareness of the inducers.
When conducting such tests of awareness, it is of paramount importance to ensure
that trials with masked and unmasked conditions are interspersed. When only
masked conditions are tested for awareness, it is possible that the participant's
performance is at chance even though there is in fact residual awareness in the main
task due to what has been referred to as "priming of awareness" [45]. Conversely,
the ability to correctly identify purportedly subliminal prime stimuli (as in our

experiment 2) may have been enhanced by the inclusion of trials with clearly visible
primes [46].
940

Naturally, our design that split the main task from a test of awareness did not allow
us to measure of the awareness on each actual trial of the main experiment – only a
dual-task design in which visibility is probed directly in the main experiment would
permit this. However, a dual-task design cannot classify the awareness of individual
trials perfectly as the participant's judgment of their own awareness is subject to
variability. Dual-tasks also entail a division of attentional resources across the
different task components. In experiment 2, where the Visibility task involved a
simple change of the search array and the instruction to now judge directly based on
the priming array, the conditions between the Kanizsa and the Visibility tasks were
very closely matched. In experiment 1, it could be argued that the Visibility task was
more difficult than the Kanizsa task because the former required the judgment of
one a small, peripheral feature of the stimulus. This may complicate the
interpretation of chance performance on this task.

Both experiments 1 and 2 used very brief stimulus presentations and powerful, high-
contrast masks to render stimuli invisible. Rather than the absence of awareness, the
reason why invisible contours were not formed could be the brevity of the stimuli or
because the mask fundamentally disrupted stimulus processing. This is certainly a
possibility because previous research indicated that the processing of illusory
contours occurs relatively slowly [35–37]. Therefore, in our third experiment, we
used counter-phase flicker to mask the inducers instead of the masking methods
employed in the earlier experiments. This allowed us to present stimuli for
prolonged periods. While this is also a temporal masking procedure, the stimulus
energy is constant during the entire presentation because only the polarity changes
between frames. Furthermore, we exploited the fact that the presence of an illusory
contour enhances performance on a dot localization task because it provides a visual
aid for determining its spatial location [19,38]. We confirmed this advantage when
inducers were not masked and performance was comparable to when we presented
real luminance contours only. However, when inducers were rendered invisible this
advantage for illusory contours disappeared. This task is a more objective test of the
induction of the illusory contours than directly testing the contour itself as in
experiment 1. Pilot experiments had suggested that with the long-lasting counter-
phase flicker masking, participants might in occasional trials have had some residual
awareness of the corners in the inducers. Thus they might still have performed
above chance at a shape/orientation discrimination on the purported illusory
contour even though they did not in fact perceive any illusory contour. This problem
also plagues many previous experiments that directly tested the presence of an
illusory shape [35,36]. Only tasks that specifically rely on a fine discrimination of a

feature of the illusory contour such as its curvature [37] or that exploit the presence
of the illusory contour to modulate performance on an orthogonal task, like the dot
localization task we used, can provide conclusive evidence that an illusory contour
was in fact perceived.

In all tests of performance against chance levels we used Bayesian hypothesis tests
that can quantify the strength of evidence for the null hypothesis that participants
were guessing [43]. None of these tests revealed *strong* evidence for the null
hypothesis as typical Bayes Factors fell between 0.3-0.5. To establish more
compelling support for the null hypothesis in those cases much larger samples would
be required. Crucially, the Bayes Factor indicates by how much the observed
evidence should update one's prior belief in the null or alternative hypothesis. Even
if the evidence is relatively weak, any Bayes Factor below 1 is evidence *in favor of*
the null and not for the alternative hypothesis. Unless one starts with a prior belief
that people are clearly able to discriminate masked stimuli, even these modest Bayes
Factors suggest that participants were probably guessing.

Taken together, our present findings and previous experiments using dichoptic
stimulation [18,19] all indicate that awareness of the inducers is required to form a
percept of illusory contours. Even if we interpret the somewhat inconclusive results
of the second experiment as showing that masked triangles produced priming
effects, this does not provide direct evidence that this advantage is actually caused
by the presence of illusory contours. A Kanizsa triangle can clearly provide a salient
attentional cue but this may be due to shared features, such as the collinearity of the
edges in the inducers or the implication of a surface. This is also consistent with
previous reports that priming of a shape discrimination task by subliminal stimuli
depends on the strength of the salient region [47] and that salient regions of such
stimuli are detected efficiently regardless of whether they are bounded by illusory
contours [38,48].

Several previous studies used stimulus manipulations that seek to disentangle these
factors [38,49,50]. Rounding the corners of the Pacman inducers results in a notable
reduction in the illusory contour percept and abolishes the concordant improvement
on a spatial localization task. However, stimuli like this nonetheless activated higher
extrastriate cortex to a similar degree as Kanizsa stimuli [38]. This is consistent with
the theory that later stages of visual processing, presumably mediated by higher
visual areas, segment surfaces and assign boundaries to objects. These
segmentations are then fed back to early visual cortex to generate signals that are
interpreted as illusory contours [49]. Practical support for this idea comes from
transcranial magnetic stimulation experiments that disrupted neural processing
either in object-sensitive lateral occipital (LO) cortex or in early visual areas V1 and

V2 [51]. Critically, the disruption of LO cortex only abolished the illusory contour
percept early after stimulus onset while disruption of early visual areas only did so at
a later stage – presumably affecting feed-back signals rather than the early feed-
forward response.
1024
Possibly the attentional priming effect we observed in our second experiment was
therefore not due to the illusory contours but could have also been observed with
other salient shapes that match the conditions of the Kanizsa triangle in all other
ways except for forming the illusory contour percept. This confound could also
explain previous reports that Kanizsa stimuli are faster to break through CFS masking
than control stimuli [31]. The collinearity of edges and the thereby inferred surface
may be processed even while the stimulus is suppressed from awareness – this may
in turn produce an attentional signal that causes the stimulus to break suppression.
Recent experiments that tested a range of visual control stimuli under CFS suggest
that low-level properties of the stimulus determine the time it takes to break
suppression [52]. Such stimulus-dependent effects are also plausible because
attentional processing can occur without awareness of the stimulus [53–55].
However, this does not prove that any percept of an illusory contour was actually
formed. This process may also explain why crowding interferes with discrimination
of the inducer orientation but not with illusory contour formation [56].
1040
While we did not manipulate the presence of illusory contours in this way in our
experiments, we nonetheless controlled this factor by including real luminance
contour conditions. In experiment 3 the presence of a contour should afford an
improvement on an orthogonal spatial localization task. Such an improvement in
localization thresholds only occurred for the real luminance contour or when Kanizsa
inducers were visible. Improvements like this are not observed for stimuli that match
the global characteristics but which do not produce illusory contours [38]. Therefore
our results from this experiment strongly support the conclusion that illusory
contours were simply not formed when inducers were masked.
1050
1051 Our findings also agree with previous findings that only local processing occurs in the
1052 absence of awareness but that more complex analysis of scene geometry requires
1053 conscious processing [17]. This would also explain why the nature of stimulus
1054 representations in higher visual areas differs depending on awareness.
1055
1056 However, our findings do not accord with a number of other studies that suggest
1057 that illusory contours are processed unconsciously. Experiments on a patient with
1058 extinction due to a parietal lesion suggest that perception of a Kanizsa shape occurs
1059 even when some inducers are placed into parts of the visual field where the patient's
1060 conscious perception is impaired [22–25]. These findings indicate that visual

processing operates at the surface- or object-based level. Segmenting and grouping the local features of an image into a coherent, global shape may only require awareness of some component features but it then spreads to the whole object. However, this is not sufficient evidence to demonstrate that these processes would occur when the participant is unaware of all components. More importantly, it also does not demonstrate that the percept of illusory contours was actually formed under these conditions but only that some processing of the features producing illusory contours under normal viewing conditions still occurred.

Another study used functional magnetic resonance imaging (fMRI) and an inattentional blindness paradigm to study illusory contours processing [50]. Participants were presented with a sequence of images, some of which contained Kanizsa shapes whilst others were various types of control stimuli. Simultaneously, participants were engaged in a demanding attentional task at fixation. A sub-group of participants subsequently reported not to have seen any Kanizsa stimuli. These participants nonetheless showed stronger fMRI responses to Kanizsa than control stimuli. Multivariate classification methods further demonstrated that the activation patterns produced by unseen Kanizsa stimuli were more reliable than those produced by control stimuli. The authors suggested that the neural signature of illusory contours differs from that of other, carefully matched stimuli and thus argued that illusory contours are processed even without awareness of the stimuli.

This finding is interesting because it tests the consequences of awareness without any experimental manipulation of the stimuli. The distinction of what is or is not processed without awareness only depends on the contents of the participant's consciousness. However, this also makes it difficult to interpret these results. First, it is unclear whether participants apparently oblivious to the presence of the stimuli really did not perceive any illusory shapes. Due to the design of the experiment, awareness could only be assessed after the main fMRI experiment rather than on a trial-by-trial basis (but see above our discussion why such trial-by-trial judgments of awareness are complicated). Only participants who reported having seen the actual Kanizsa stimulus during the experiment were classified as having had awareness of the stimuli. However, many of the candidate stimuli were similar Kanizsa shapes. Therefore it is possible that participants had some awareness of the stimuli, even if an imprecise one.

Second, while the control stimuli in the main fMRI experiment were very carefully matched to rule out the influence of global characteristics this can by definition only be an approximation: if conditions were perfectly matched, the stimulus would be identical and thus an illusory contour would be perceived. It is possible that the conditions resulting in an illusory contour percept are also particularly effective in

producing discriminable activation patterns in visual cortex. For instance, the contrast energy along the mouths of the Pacmen inducers differed considerably between the Kanizsa and the global control stimuli. Thus, surface segmentation or collinear interactions may have also differed between these conditions.

All these issues highlight an important point: The only way one can truly infer that illusory contours are formed is by direct report. This is indeed what we did in our previous CFS experiment [18], our present experiment 1, or what was done in several other previous experiments [19,30,51,56]. Another useful manipulation is the spatial localization task used by other studies [19,38] as well as our experiment 3. This is because the performance enhancement only occurs in the presence of an actual contour helping the participant to localize the target. Many previous studies suggesting that illusory contours are processed without awareness of the inducers are confounded by this problem, including our experiment 2 (if one assumes that those results are not trivially explained by residual awareness).

Neurophysiological and neuroimaging experiments have shown that illusory contours are encoded by neurons even in the early visual cortex [21,29,57–68]. The overwhelming majority of these experiments were conducted on awake participants. However, some neurophysiological studies reported neuronal tuning to illusory contours even in anesthetized animals [21,59,68]. Such findings seem to superficially contradict our conclusion that awareness of the inducing stimuli is necessary for the formation of illusory contours.

However, these experiments are in fact a perfect illustration of the importance to distinguish between *perceptual experience* and correlated processing. Because animals are anesthetized in these experiments and thus by definition unaware of the stimuli, it is impossible to determine whether illusory contours were formed. The neural correlates of these stimuli could be related to the contextual processing of the inducing stimuli, such as the discontinuities detected by "end-stopped cells" or the detection of collinearity in the image. Such processing may indeed occur in the absence of awareness as is supported by our finding of collinear priming [17] or the induction of the tilt illusion under masking conditions [2]. Furthermore, it is quite likely that such stimulus processing is a necessary prerequisite for the formation of illusory contours. However, they do not conclusively prove that illusory contours are formed under anesthesia.

A related issue is that all of these studies use illusory contours induced by abutting lines (offset gratings) rather than Kanizsa shapes. While the two share a similar phenomenology, contours induced by abutting lines are arguably simpler and may be based mostly on local processing while Kanizsa shapes are likely to involve more

complex inferences of surface depth and boundary ownership. Our experiments did not explicitly test illusory contours generated with abutting lines and therefore do not speak to the question whether such simpler illusory contours are in fact perceived when the inducers are masked.

We conclude that there is little evidence that illusory contours are processed when participants are not aware of the inducing context. This appears to be the case for a range of different methods to render the inducers invisible. However, all of these experiments nonetheless employed physical stimulus manipulations. Only an experimental design that allows the use of the participant's own report to determine awareness on a trial-by-trial basis whilst also testing directly whether an illusory contour was in fact formed, either behaviorally or through physiological measures, can answer this question conclusively. Such a design will however require great care to ensure that misclassifications of aware trials does not skew the results.

Acknowledgements

This research was supported by an ERC Starting Grant (310829) to DSS. We thank Pieter Moors, Jacob Jolij, and David Shanks as well as two anonymous reviewers for comments on a previous manuscript and Jonas Larsson for inspiring this study in the first place.

Data and Materials

All stimulus code and data for this study are available for download at:
https://dx.doi.org/10.6084/m9.figshare.2009952.

References

1. Newell BR, Shanks DR. Unconscious influences on decision making: a critical review. Behav Brain Sci. 2014;37: 1–19. doi:10.1017/S0140525X12003214

2. Clifford CWG, Harris JA. Contextual modulation outside of awareness. Curr Biol. 2005;15: 574–8. doi:10.1016/j.cub.2005.01.055

3. Falconbridge M, Ware A, MacLeod DIA. Imperceptibly rapid contrast modulations processed in cortex: Evidence from psychophysics. J Vis. 2010;10. doi:10.1167/10.8.21

4. Pearson J, Clifford CWG. Suppressed patterns alter vision during binocular rivalry. Curr Biol. 2005;15: 2142–8. doi:10.1016/j.cub.2005.10.066

5. Vul E, MacLeod DIA. Contingent aftereffects distinguish conscious and preconscious color processing. Nat Neurosci. 2006;9: 873–874. doi:10.1038/nn1723

6.   Tsuchiya N, Koch C. Continuous flash suppression reduces negative afterimages. Nat
Neurosci. 2005;8: 1096–1101. doi:10.1038/nn1500

7.   Stewart LH, Ajina S, Getov S, Bahrami B, Todorov A, Rees G. Unconscious evaluation of
faces on social dimensions. J Exp Psychol Gen. 2012;141: 715–727.
doi:10.1037/a0027950

8.   Mudrik L, Breska A, Lamy D, Deouell LY. Integration without awareness: expanding the
limits of unconscious processing. Psychol Sci. 2011;22: 764–770.
doi:10.1177/0956797611408736

9.   Sklar AY, Levy N, Goldstein A, Mandel R, Maril A, Hassin RR. Reading and doing
arithmetic nonconsciously. Proc Natl Acad Sci U S A. 2012;109: 19614–19619.
doi:10.1073/pnas.1211645109

10.   Haynes J-D, Rees G. Predicting the orientation of invisible stimuli from activity in
human primary visual cortex. Nat Neurosci. 2005;8: 686–91. doi:10.1038/nn1445

11.   Moutoussis K, Zeki S. The relationship between cortical activation and perception
investigated with invisible stimuli. Proc Natl Acad Sci U S A. 2002;99: 9527–9532.
doi:10.1073/pnas.142305699

12.   Sterzer P, Haynes J-D, Rees G. Fine-scale activity patterns in high-level visual areas
encode the category of invisible objects. J Vis. 2008;8: 10.1-12. doi:10.1167/8.15.10

13.   Haynes J-D, Driver J, Rees G. Visibility reflects dynamic changes of effective
connectivity between V1 and fusiform cortex. Neuron. 2005;46: 811–821.
doi:10.1016/j.neuron.2005.05.012

14.   Schurger A, Pereira F, Treisman A, Cohen JD. Reproducibility distinguishes conscious
from nonconscious neural representations. Science. 2010;327: 97–99.
doi:10.1126/science.1180029

15.   Schurger A, Sarigiannidis I, Naccache L, Sitt JD, Dehaene S. Cortical activity is more
stable when sensory stimuli are consciously perceived. Proc Natl Acad Sci U S A.
2015;112: E2083-2092. doi:10.1073/pnas.1418730112

16.   Ling S, Blake R. Suppression During Binocular Rivalry Broadens Orientation Tuning.
Psychol Sci J Am Psychol Soc APS. 2009; doi:10.1111/j.1467-9280.2009.02446.x

17.   Schwarzkopf DS, Rees G. Interpreting local visual features as a global shape requires
awareness. Proc Biol Sci. 2011;278: 2207–2215. doi:10.1098/rspb.2010.1909

18.   Harris JJ, Schwarzkopf DS, Song C, Bahrami B, Rees G. Contextual illusions reveal the
limit of unconscious visual processing. Psychol Sci J Am Psychol Soc APS. 2011;22: 399–
405. doi:10.1177/0956797611399293

19.   Sobel KV, Blake R. Subjective contours and binocular rivalry suppression. Vision Res.
2003;43: 1533–1540.

20.   Gellatly AR. Perception of an illusory triangle with masked inducing figure. Perception.
1980;9: 599–602.

21.   Ramsden BM, Hung CP, Roe AW. Real and illusory contour processing in area V1 of the
1221   primate: a cortical balancing act. Cereb Cortex N Y N 1991. 2001;11: 648–665.

1222   22.   Conci M, Böbel E, Matthias E, Keller I, Müller HJ, Finke K. Preattentive surface and
1223   contour grouping in Kanizsa figures: evidence from parietal extinction.
1224   Neuropsychologia. 2009;47: 726–732. doi:10.1016/j.neuropsychologia.2008.11.029

1225   23.   Mattingley JB, Davis G, Driver J. Preattentive filling-in of visual surfaces in parietal
1226   extinction. Science. 1997;275: 671–674.

1227   24.   Vuilleumier P, Landis T. Illusory contours and spatial neglect. Neuroreport. 1998;9:
1228   2481–2484.

1229   25.   Vuilleumier P, Valenza N, Landis T. Explicit and implicit perception of illusory contours
1230   in unilateral spatial neglect: behavioural and anatomical correlates of preattentive
1231   grouping mechanisms. Neuropsychologia. 2001;39: 597–610.

1232   26.   Gillam B, Nakayama K. Quantitative depth for a phantom surface can be based on
1233   cyclopean occlusion cues alone. Vision Res. 1999;39: 109–112.

1234   27.   Häkkinen J, Nyman G. Phantom surface captures stereopsis. Vision Res. 2001;41: 187–
1235   199.

1236   28.   Liu L, Stevenson SB, Schor CM. Quantitative stereoscopic depth without binocular
1237   correspondence. Nature. 1994;367: 66–69. doi:10.1038/367066a0

1238   29.   von der Heydt R, Peterhans E, Baumgartner G. Illusory contours and cortical neuron
1239   responses. Science. 1984;224: 1260–1262.

1240   30.   Rubin N. The role of junctions in surface completion and contour matching. Perception.
1241   2001;30: 339–66.

1242   31.   Wang L, Weng X, He S. Perceptual grouping without awareness: superiority of Kanizsa
1243   triangle in breaking interocular suppression. PloS One. 2012;7: e40106.
1244   doi:10.1371/journal.pone.0040106

1245   32.   Davis G, Driver J. Parallel detection of Kanizsa subjective figures in the human visual
1246   system. Nature. 1994;371: 791–793. doi:10.1038/371791a0

1247   33.   Barlasov-Ioffe A, Hochstein S. Illusory-contour figures prime matching of real shapes.
1248   Perception. 2009;38: 1118–1131.

1249   34.   Seydell-Greenwald A, Schmidt T. Rapid activation of motor responses by illusory
1250   contours. J Exp Psychol Hum Percept Perform. 2012;38: 1168–1182.
1251   doi:10.1037/a0028767

1252   35.   Parks TE. On the microgenesis of illusory figures: a failure to replicate. Perception.
1253   1994;23: 857–862.

1254   36.   Reynolds RI. Perception of an illusory contour as a function of processing time.
1255   Perception. 1981;10: 107–115.

37. Ringach DL, Shapley R. Spatial and temporal properties of illusory contours and amodal
boundary completion. Vision Res. 1996;36: 3037–3050.

38. Stanley DA, Rubin N. fMRI activation in response to illusory contours and salient
regions in the human lateral occipital complex. Neuron. 2003;37: 323–331.

39. Brainard DH. The Psychophysics Toolbox. Spat Vis. 1997;10: 433–6.

40. Frisby JP, Clatworthy JL. Illusory Contours: Curious Cases of Simultaneous Brightness
Contrast? Perception. 1975;4: 349–357. doi:10.1068/p040349

41. Halpern DF. The determinants of illusory-contour perception. Perception. 1981;10:
199–213.

42. Varin D. Fenomeni di contrasto e diffusione cromatica nell'organizzazione spaziale del
campo percettivo. [Contrast and chromatic diffusion phenomena in the organization of
the perceptive field]. Riv Psicol. 1971;65: 101–128.

43. Rouder JN, Speckman PL, Sun D, Morey RD, Iverson G. Bayesian t tests for accepting
and rejecting the null hypothesis. Psychon Bull Rev. 2009;16: 225–237.
doi:10.3758/PBR.16.2.225

44. Muise JG, LeBlanc RS, Blanchard LC, de Warnaffe A. Discrimination of the shape of the
masked inducing figure precedes perception of the illusory triangle. Perception.
1993;22: 623–628.

45. Lin Z, Murray SO. Priming of awareness or how not to measure visual awareness. J Vis.
2014;14. doi:10.1167/14.1.27

46. Pratte MS, Rouder JN. A task-difficulty artifact in subliminal priming. Atten Percept
Psychophys. 2009;71: 1276–1283. doi:10.3758/APP.71.6.1276

47. Poscoliero T, Marzi CA, Girelli M. Unconscious priming by illusory figures: the role of
the salient region. J Vis. 2013;13: 27. doi:10.1167/13.5.27

48. Stanley DA, Rubin N. Rapid detection of salient regions: evidence from apparent
motion. J Vis. 2005;5: 690–701. doi:10.1167/5.9.4

49. Kogo N, Strecha C, Van Gool L, Wagemans J. Surface construction by a 2-D
differentiation-integration process: a neurocomputational model for perceived border
ownership, depth, and lightness in Kanizsa figures. Psychol Rev. 2010;117: 406–439.
doi:10.1037/a0019076

50. Vandenbroucke ARE, Fahrenfort JJ, Sligte IG, Lamme VAF. Seeing without knowing:
neural signatures of perceptual inference in the absence of report. J Cogn Neurosci.
2014;26: 955–969. doi:10.1162/jocn_a_00530

51. Wokke ME, Vandenbroucke ARE, Scholte HS, Lamme VAF. Confuse your illusion:
feedback to early visual cortex contributes to perceptual completion. Psychol Sci.
2013;24: 63–71. doi:10.1177/0956797612449175

52.   Moors P, Wagemans J, van Ee R, de-Wit L. No evidence for surface organization in
1293   Kanizsa configurations during continuous flash suppression. Atten Percept Psychophys.
1294   2015; doi:10.3758/s13414-015-1043-x

1295   53.   Bahrami B, Carmel D, Walsh V, Rees G, Lavie N. Spatial attention can modulate
1296   unconscious orientation processing. Perception. 2008;37: 1520–8.

1297   54.   Koch C, Tsuchiya N. Attention and consciousness: two distinct brain processes. Trends
1298   Cogn Sci. 2007;11: 16–22. doi:10.1016/j.tics.2006.10.012

1299   55.   Zhaoping L. Attention capture by eye of origin singletons even without awareness--a
1300   hallmark of a bottom-up saliency map in the primary visual cortex. J Vis. 2008;8: 1.1-
1301   18. doi:10.1167/8.5.1

1302   56.   Lau JSF, Cheung S-H. Illusory contour formation survives crowding. J Vis. 2012;12: 15.
1303   doi:10.1167/12.6.15

1304   57.   Cox MA, Schmid MC, Peters AJ, Saunders RC, Leopold DA, Maier A. Receptive field
1305   focus of visual area V4 neurons determines responses to illusory surfaces. Proc Natl
1306   Acad Sci U S A. 2013;110: 17095–17100. doi:10.1073/pnas.1310806110

1307   58.   Ffytche DH, Zeki S. Brain activity related to the perception of illusory contours.
1308   NeuroImage. 1996;3: 104–108. doi:10.1006/nimg.1996.0012

1309   59.   Grosof DH, Shapley RM, Hawken MJ. Macaque V1 neurons can signal "illusory"
1310   contours. Nature. 1993;365: 550–552. doi:10.1038/365550a0

1311   60.   Heider B, Meskenaite V, Peterhans E. Anatomy and physiology of a neural mechanism
1312   defining depth order and contrast polarity at illusory contours. Eur J Neurosci. 2000;12:
1313   4117–4130.

1314   61.   Kok P, de Lange FP. Shape perception simultaneously up- and downregulates neural
1315   activity in the primary visual cortex. Curr Biol CB. 2014;24: 1531–1535.
1316   doi:10.1016/j.cub.2014.05.042

1317   62.   Lee TS, Nguyen M. Dynamics of subjective contour formation in the early visual cortex.
1318   Proc Natl Acad Sci U S A. 2001;98: 1907–1911. doi:10.1073/pnas.031579998

1319   63.   Mendola JD, Dale AM, Fischl B, Liu AK, Tootell RB. The representation of illusory and
1320   real contours in human cortical visual areas revealed by functional magnetic resonance
1321   imaging. J Neurosci Off J Soc Neurosci. 1999;19: 8560–8572.

1322   64.   Montaser-Kouhsari L, Landy MS, Heeger DJ, Larsson J. Orientation-selective adaptation
1323   to illusory contours in human visual cortex. J Neurosci Off J Soc Neurosci. 2007;27:
1324   2186–2195. doi:10.1523/JNEUROSCI.4173-06.2007

1325   65.   Pan Y, Chen M, Yin J, An X, Zhang X, Lu Y, et al. Equivalent representation of real and
1326   illusory contours in macaque V4. J Neurosci Off J Soc Neurosci. 2012;32: 6760–6770.
1327   doi:10.1523/JNEUROSCI.6140-11.2012

1328   66.   Peterhans E, von der Heydt R. Subjective contours--bridging the gap between
1329   psychophysics and physiology. Trends Neurosci. 1991;14: 112–9.

67.   von der Heydt R, Peterhans E. Mechanisms of contour perception in monkey visual
cortex. I. Lines of pattern discontinuity. J Neurosci Off J Soc Neurosci. 1989;9: 1731–
1748.

68.   Zhan CA, Baker CL. Critical spatial frequencies for illusory contour processing in early
visual cortex. Cereb Cortex N Y N 1991. 2008;18: 1029–1041.
doi:10.1093/cercor/bhm139

1336