Elisabetta Versace[1*], Jana Schill[1,2], Andrea Maria Nencini[1], Giorgio Vallortigara[1]

Naïve chicks prefer hollow objects

[1] Center for Mind/Brain Sciences, University of Trento (Italy)

[2] Universität Osnabrück, Germany

Abstract

Cognitive predispositions can influence approaching and avoid responses since the early

stages of life. Young individuals of species that require parental care (e.g. human babies

and chicks of the domestic fowl) are attracted by stimuli that contain features present in

social partners such as face-like configurations, biological motion and self-propulsion.

Studies on human infants showed that 8-month-old babies might possess expectations

about the biological properties of animate entities. It is not clear though whether

previous experience with animate entities had generated those expectations, or whether

they arose spontaneously. We reasoned that naïve chicks of the domestic fowl (*Gallus*

gallus) might be a convenient subject to investigate whether the mere property of being

filled vs. hollow triggers unlearned preferences. To this aim we tested preferences of

naïve and imprinted chicks for hollow and closed cylinders of the size and colour that

elicit affiliative responses. We documented an unlearned attraction for hollow stimuli,

showing that the property of being filled is not sufficient to elicit affiliative responses in

chicks. The preference for hollow stimuli could be decreased through filial imprinting, by

exposing naïve chicks to filled stimuli. When chicks were imprinted on stimuli that could

be either filled or hollow, the preference for hollow stimuli emerged again. Further

experiments revealed that hollow objects were mainly attractive by means of depth cues

such as darker innards, more than as places to hide or as objects with high contrast.

Introduction

Sensory and cognitive predispositions can help living beings to cope with their environment

by influencing approach and avoid responses (Vallortigara 2012a; Rosa Salva et al. 2015;

Versace and Vallortigara 2015). It is not surprising, hence, to observe in different species

unlearned preferences for specific colours (Schaefer and Hess 1951; Giurfa et al. 1995; Raine

and Chittka 2007; Ham and Osorio 2007), shapes and sizes (Fabricius and Boyd; Schulman et

al. 1970; Ham and Osorio 2007), configurations (Rosa-Salva et al. 2010), dynamics

(Vallortigara et al. 2005; Mascalzoni et al. 2010), and odours (Knaden et al. 2012; Dekker et al.

2015; Keesey et al. 2015; Versace and Reisenberger 2015).

Precocial species – in which individuals are mobile from the moment of birth, and can be tested

with little if any experience with their environment– are very convenient to investigate the

unlearned predispositions that can orient living beings since their early stages of life (Versace

and Vallortigara 2015). Soon after hatching, chicks of the domestic fowl (*Gallus gallus*), which

is a nidifugal species, possess some unlearned preferences to approach stimuli that are more

similar to animate social partners (Vallortigara 2012b).

Evolved mechanisms to approach objects that possess the features of animate partners would

provide a critical advantage for survival in species that rely on parental care. In line with this,

when given a choice between a stuffed hen *vs.* a stuffed scrambled hen, chicks prefer to

approach the hen (Johnson and Horn 1988). Behavioural studies have found that this

preference is driven by an unlearned attraction towards the face configuration contained in

the stuffed hen (Johnson and Horn 1988; Rosa-Salva et al. 2010). Moreover, between the

biological movement of a hen or a cat and the rigid motion of a hen rotated on its vertical axis,

chicks prefer to approach the biologically moving object (Vallortigara et al. 2005; Miura and

Matsushima 2016); and between a self-propelled object and an object propelled by another

one, naïve chicks prefer the self-propelled object (Mascalzoni et al. 2010). Overall, chicks

prefer to approach objects which are endowed with more animate features (Rosa Salva et al.

2015; Versace and Vallortigara 2015).

Observations on infants (Gelman 1990) suggest that 3-year-old children have a

representation of the insides of animate beings as more likely to be filled than those of

inanimate objects. Studies on human infants (Setoh et al. 2013) have shown that 8-month-old

babies might possess expectations about the biological properties of animate and agentive

entities. In this study infants were more surprised to see that self-propelled and agentive

objects were hollow than when there was no evidence that those objects were hollow. It is not

clear though whether previous experience with animate entities with innards (e.g. the

parents) had generated infants' expectations, or whether they arose spontaneously. We

reasoned that chicks of the domestic fowl (*Gallus gallus*), who are spontaneously attracted by

entities which show cues associated with animacy in the absence of previous experience

(Rosa Salva et al. 2015; Versace and Vallortigara 2015), might be a convenient subject to

identify whether the property of being filled/hollow triggers unlearned preferences.

To this aim we tested preferences of naïve chicks (**Experiment 1**) maintained in darkness

(Experiment 1a) or exposed to light (Experiment 1b) for hollow and closed cylinders of the

size and colour that elicit filial responses. Moreover, since chicks rapidly learn features of

their social partners by mere exposure through filial imprinting (Horn 1985; McCabe 2013)

(Horn 1985; McCabe 2013), they are a valuable model to study the role of experience in

modifying spontaneous preferences. To this aim we investigated how imprinting modified

unlearned preferences for hollow and filled objects (**Experiment 2**) after imprinting on

hollow objects (Experiment 2a), filled objects (Experiment 2b) and objects who could not be

perceived hollow or filled because their sides were occluded (Experiment 2c).

Since we noticed an overall preference for hollow objects, we investigated whether this

behaviour was elicited by a preference for the stimulus that could better hide the chick

(chicks could enter the hollow stimulus). In **Experiment 3** we checked whether the

preference for hollow stimuli was still present when the stimuli were too small to host and

hide chicks. We tested both dark-reared chicks and chicks exposed to light that had never

seen the test stimuli or any other object of similar size, shape and colour. We observed a

preference for hollow objects. In **Experiment 4** we checked whether the size of the hollow

object was important in determining the preference for hollow objects comparing the

preference for the large and the narrow hollow objects. In **Experiment 5** we checked whether

the darker colour of the shadows present in the innards of hollow objects has a role in driving

preferences for hollow stimuli by comparing preferences for filled objects with a white *vs.* a

black stopper (Experiment 5a). Since chicks preferred the object with the black stopper, we

tested whether the preference for a hollow stimulus was stronger or weaker than the

preference for a black cap (Experiment 5b). The observed preference for the black cap

stimulus could be explained both by brightness (chicks preferred lower brightness) and by

contrast (chicks preferred greater contrast). To clarify the importance of contrast and

brightness in determining the preference for hollow objects, in **Experiment 6** we used two-

dimensional stimuli with different colour and identical contrast, i.e. a white disk on a black

background *vs.* a black disk on a white background. If the preference of chicks for hollow *vs.*

Filled and for Black *vs.* Hollow was driven by the darker colour (innards or cap), in this

contrast chicks should have chosen the white disk on a black background. If the preference

was driven by contrast, chicks were expected to have no preference. A preference for the

black disk on a white background would be consistent with a preference for darker

objects/innards, possibly a cue of depth.

 MATERIALS AND METHODS

Ethical note

This study was approved by the ethical committee of the University of Trento (Organismo

preposto al benessere degli Animali) prot. N. 14-2015 and was licensed by the Ministero della

Salute, authorization n. 1138/2015. The research adhered to the ASAB/ABS Guidelines for the

Use of Animals in Research.

Subjects

The subjects were 24-hour old chicks of the domestic fowl (*Gallus gallus*) of the Hybro strain

(a local hybrid variety of the White Leghorn breed). This breed is sexually dimorphic at the

moment of hatching. The eggs were obtained from a commercial hatchery (Agricola Berica,

Montegalda, Italy), then incubated in complete darkness at 37.7 °C until hatching. Three days

before hatching humidity was increased from 40% to 60%. Eggs hatched in individual boxes

(11 x 8.5 x 14 cm) and chicks could hear their conspecifics but had no visual or tactile contact

with conspecifics until the moment of test. The exact number of chicks used in each

experiment, divided by Sex, is presented in Table 1.

Experiment	Experience	Test stimuli	Females	Males
Experiment 1a	Naïve: Dark-reared	Hollow – Filled (large)	57	52
Experiment 1b	Naïve: Light-reared	Hollow – Filled (large)	27	26
Experiment 2a	Imprinted: Occluded	Hollow – Filled (large)	37	27
Experiment 2b	Imprinted: Filled	Hollow – Filled (large)	59	64
Experiment 2c	Imprinted: Hollow	Hollow – Filled (large)	57	67
Experiment 3	Naïve: Light-reared	Hollow – Filled (narrow)	17	14
Experiment 4	Naïve: Dark-reared	Large – Narrow (hollow)	47	36
Experiment 5a	Naïve: Dark-reared	White – Black (large filled)	11	16
Experiment 5b	Naïve: Light-reared	Hollow – Black filled (large)	19	21
Experiment 6	Naïve: Dark-reared	White disk – Black disk	19	17

Test stimuli

Test stimuli are shown in Figure 1. In Experiment 1 and 2 test stimuli were large plastic tubes

(12 cm, ø 4 cm) left open (Hollow, Fig. 1A) or closed with a white cap (Filled, Fig. 1B), with an

orange external surface and a white internal surface. In Experiment 3 we used the same

stimuli with the only difference that the diameter was 2.5 cm (Narrow stimuli are shown in

Fig. 1C and 1D). In Experiment 4 we used Large and Narrow hollow stimuli (Fig. 1A and 1C).

In Experiment 5 we used stimuli similar to those used in Experiments 1 and 2 with the only

difference that one cap was black (Fig. 1E). In Experiment 6 we used a white disk on a black

background, and a black disk on a white background (Fig. 1F) with a diameter of 4 cm located

at 4.5 cm from the ground.

Figure 1. Stimuli used in Experiment 1 and 2 (**A** and **B**), Experiment 3 (**C** and **D**), Experiment 4 (**A** and **C**),

Experiment 5 (**A** and **E**), and Experiment 6 (Panel **F** shows the Black disk on the white background. The other

stimulus was a White disk on a black background).

Imprinting stimuli

In experiment 2 chicks were individually imprinted to orange cylinders (12 cm, ø 4 cm), that

were presented through a 7.5 x 10 cm transparent plastic window. Imprinting lasted 24 ± 3

and was immediately followed by the test. Chicks had no direct interaction with the stimulus

during imprinting and the only interaction with conspecifics was auditory. In the Occluded

condition the cylinder was presented horizontally and the chicks could not see whether it was

hollow or filled because the edges were covered. In the Hollow and Filled condition the hollow

and or filled cylinder were presented perpendicular to the transparent window and the chick

could see whether it was hollow or not.

Test apparatus

The experiment took place in a 100 x 30 x 31 cm white arena open on the top (see Figure 2).

Test stimuli were located in the middle of each short side on a white plastic platform that was

4.5 cm high. The box was virtually divided into three areas: a left area (41 cm), a central area

(18 cm) and a right area (41 cm). The white platforms occupied 15 cm in each side area. In

Experiment 4, the platforms were removed and the stimuli were placed directly on the walls

of the apparatus.

153
154
Figure 2. Illustration of the testing apparatus.

PROCEDURE

Imprinting

Soon after hatching, in the imprinting experiments chicks were individually exposed to the

imprinting stimulus for 24 hours before testing under constant light. Imprinting cages were

28 x 38 x 32 cm and the stimulus was presented through a transparent partition (7.5 x 10 cm).

In this way chicks had no direct interaction with the stimuli before testing, similarly to naïve

chicks that had never experienced stimuli like those used during the test.

Test

Procedure and data analysis

We followed the same procedure in all experiments. Each chick was individually located in the

centre area facing the long side of the box opposite to the experimenter and video recorded

for 360 seconds. We recorded which side area was entered first (First choice) and the seconds

spent in each side area. The chick was considered to have entered a new sector as soon as it

crossed the borderline with both feet. After the testing phase chicks were not used in any

other experiment.

We checked whether the first choice was significantly different from the 0.5 chance level

using a Chi-squared test, with alpha = 0.05.

For each chick that left the central area we calculated an index of preference for the Hollow

stimulus (Experiments 1, 2 and 3) or an index of preference for the Narrow (Experiment 4) or

Black stimulus (5) in this way:

Hollow preference = (seconds in the Hollow stimulus area) / (seconds in the Hollow stimulus area+seconds in the

Filled stimulus area)

Narrow preference = (seconds in the Narrow stimulus area) / (seconds in the Narrow stimulus area+seconds in

the Large stimulus area)

Black preference = (seconds in the Black stimulus area) / (seconds in the Black stimulus area+seconds in the

White stimulus area)

For all indices, 1 indicates a full preference for the respective stimulus (Hollow, Narrow,

Black), 0.5 no preference and 0 a full preference for the opposite stimulus (Filled, Large,

White). Since all data had a bimodal distribution with peaks on the extremes (0 and 1) we

used non-parametric statistics to test for significance: the Kruskal-Wallis test to test for

differences between conditions and sexes, and the Mann-Whitney-Wilcoxon one-sample test

vs. the 0.5 chance level.

RESULTS

Experiment 1: naïve chicks (dark-reared and light-reared) chicks tested with Hollow *vs.* Filled stimuli.

We assessed the preference for the hollow/filled object in naïve chicks, namely dark-reared and light-reared chicks that had never experiences any of the test stimuli before the test.

First choice. There was no significant difference between dark- and light-reared chicks (Chi-square test: $\chi = 0.073$, df = 1, $P = 0.79$), and in both conditions chicks had the same trend, therefore we collapsed the two naïve conditions for further analyses. The number of chicks that approached the Hollow *vs.* Filled stimulus was significantly different from chance (Chi-square test: $\chi = 8.91$, df = 1, $P = 0.003$) with an overall preference for the Hollow stimulus (Figure 3A).

Hollow preference. Considering the Hollow preference index we did not observe any significant Sex difference (Kruskal-Wallis test: $H = 0.32$, df = 1, $P = 0.57$) or Exposure (dark- *vs.* light-rearing) difference (Kruskal-Wallis test: $H = 0.07$, df = 1, $P = 0.79$), therefore we collapsed the two naïve conditions for further analyses. We documented a significant preference for the Hollow stimulus (Mann-Whitney test: $V = 8053$, df = 1, $P = 0.01$), see Figure 3B.

Figure 3. **A.** Number of naïve chicks that first approached the Hollow or Filled stimulus in the dark-reared and light-reared condition. **B.** Proportion of time spent at the Hollow stimulus by dark-reared and light-reared chicks.

Experiment 2: chicks imprinted with Hollow, Filled or Occluded and tested with Hollow *vs.* Filled stimuli.

To investigate the role of experience in determining the preferences for hollow objects we investigated the preference for the hollow/filled object in imprinted chicks, namely chicks that had been exposed to the filled or hollow object, or to an object located horizontally the sides of which were occluded, so that it did not show whether it was filled or hollow.

First choice. The number of chicks that approached the Hollow *vs.* Filled stimulus was significantly different between imprinting conditions (Chi-square test: χ = 7.15, df = 2, *P* = 0.028). Chicks imprinted on the Occluded object showed a significant preference for the hollow object (Chi-square test: χ = 7.56, df = 1, *P* = 0.006), whereas chicks imprinted on the Filled (Chi-square test: χ = 0.40, df = 1, *P* = 0.53) and Hollow object (Chi-square test: χ = 2.61, df = 1, *P* = 0.11) did not. While the first choice of chicks imprinted on the Occluded object did not differ from the first choice of chicks imprinted on the Hollow object (Chi-square test: χ = 1.35, df = 1, *P* = 0.24), there was a significant difference between the first choice of chicks imprinted on the Occluded object and the first choice of chicks imprinted on the Filled object (Chi-square test: χ = 6.02, df = 1, *P* = 0.014). Only chicks imprinted on the Filled object had a tendency to choose the Filled object (Figure 4A).

Figure 4. **A.** Number of imprinted chicks that first approached the Hollow or Filled stimulus after being exposed to Occluded, Hollow or Filled imprinting stimuli. **B.** Proportion of time spent at the Hollow stimulus for chicks exposed to Occluded, Hollow or Filled imprinting stimuli.

Hollow preference. Considering the Hollow preference index we did not observe any

significant Sex difference (Kruskal-Wallis test: $H = 1.60$, df = 1, $P = 0.21$) or Exposure

difference (Kruskal-Wallis test: $H = 3.66$, df = 2, $P = 0.161$). We observed an overall trend for

preferring the the Hollow stimulus (Mann-Whitney test: $V = 27016.5$, df = 1, $P = 0.063$), that

turned out highly significant when considering only the chicks imprinted on the occluded and

hollow objects (Mann-Whitney test: $V = 10721.5$, df = 1, $P = 0.009$), see Figure 4B.

Experiment 3: naïve chicks tested with narrow Hollow *vs.* narrow Filled stimuli.

To investigate the extent and consistency of the hollow preference, we tested the preference

for the hollow/filled object in naïve chicks, using smaller stimuli than those used in

Experiment 1.

First choice. Chicks confirmed the preference for hollow stimuli (Chi-square test: $\chi = 17.06$, df

= 1, $P < 0.001$).

Hollow preference. Considering the Hollow preference index we did not observe any

significant Sex difference (Kruskal-Wallis test: $H = 1.46$, df = 1, $P = 0.23$) but an overall

preference for the Hollow stimulus (Mann-Whitney test: $V = 461$, df = 1, $P < 0.001$).

Experiment 4: naïve chicks tested with Large hollow *vs.* Narrow hollow stimuli.

To investigate whether the preference of young chicks for hollow objects was driven by the

possibility to hide inside hollow objects, we presented naïve dark-reared chicks with a choice

between Large (4 cm in diameter, large enough to hide a chick) and Narrow hollow stimuli

(2.5 cm in diameter, too small to hide a chick).

First choice. The number of chicks that approached the Large *vs.* Narrow stimulus was not

significantly different between Sexes (Chi-square test: $\chi = 0.14$, df = 1, $P = 0.71$), therefore we

collapsed the data from males and females together. There was no significant preference for

the Large or Narrow stimulus (Chi-square test: $\chi = 0.108$, df = 1, $P = 0.74$), suggesting that the

possibility to hide inside the Large hollow stimuli is not the main drive of the preference for

hollow stimuli.

Narrow preference. Considering the Narrow preference index we did not observe any

significant Sex difference (Kruskal-Wallis test: $H = 0.11$, df = 1, $P = 0.74$). Overall we observed

no significant preference for Large or narrow stimuli (Mann-Whitney test: $V = 1583.5$, df = 1, P

= 0.56).

Experiment 5a: naïve chicks tested with filled White *vs.* filled Black stimuli.

First choice. The number of chicks that approached the White *vs.* Black stimulus was not significantly different between Sexes (Chi-squared test: $\chi = 0.12$, df = 1, $P = 0.73$), therefore we collapsed the data from males and females together. We observed a significant preference for the Black stimulus (Chi-squared test: $\chi = 16.33$, df = 1, $P < 0.001$), see Figure 5A.

Black preference. Considering the Black preference index, we did not observe any significant Sex difference (Kruskal-Wallis test: $H = 0.066$, df = 1, $P = 0.80$). Overall we observed a significant preference for the Black stimulus (Mann-Whitney test: $V = 354$, df = 1, $P < 0.001$), see Figure 5B.

Figure 5. **A.** Number of chicks that first approached the Black or White stimulus. **B.** Proportion of time spent at the Black stimulus.

Experiment 5b: naïve chicks tested with filled Black *vs.* Hollow stimuli.

First choice. The number of chicks that approached the Hollow *vs.* Black stimulus was not significantly different between Sexes (Chi-squared test: $\chi = 0.307$, df = 1, $P = 0.58$), therefore we collapsed the data from males and females together. We observed a significant preference for the Black stimulus (Chi-squared test: $\chi = 14.4$, df = 1, $P < 0.001$), see Figure 6A.

Black preference. Considering the Black preference index we did not observe any significant Sex difference (Chi-squared test: $\chi = 0.818$, df = 1, $P = 0.366$). Overall we observed a significant preference for the Black stimulus (Mann-Whitney test: $V = 174$, $P < 0.001$), see Figure 6B.

Figure 6. **A.** Number of chicks that first approached the Black or Hollow stimulus. **B.** Proportion of time spent at the Black stimulus.

Experiment 6: naïve chicks tested with a White disk on a black background *vs.* a Black disk on a white background.

First choice. The number of chicks that approached the White *vs.* Black disk was not significantly different between Sexes (Chi-squared test: χ = 0.166, df = 1, *P* = 0.68). Overall we observed a significant preference for the Black stimulus (Chi-squared test: χ = 19.882, df = 1, *P* < 0.001), see Figure 7A.

Black preference. Considering the Black preference index we did not observe any significant Sex difference (Kruskal-Wallis χ^2 = 0.65, df = 1, *P* = 0.42). Overall we observed a significant preference for the Black stimulus (Mann-Whitney test: *V* = 519, df = 1, *P* < 0.001), see Figure 7B.

Figure 7. **A.** Number of chicks that first approached the Black disk on a white background or the White disk on a black background. **B.** Proportion of time spent at the Black disk on a white background.

Discussion

Sensory and cognitive predispositions can help naïve individuals in deciding whether to approach or avoid novel objects (Versace and Vallortigara 2015). Chicks of the domestic fowl, which belong to a precocial social species, appear to be endowed with predispositions to approach animate social partners (Rosa Salva et al. 2015), given that in the absence of previous experience, young individuals prefer to approach face-like configurations (Rosa-Salva et al. 2010), self-propelled objects (Mascalzoni et al. 2010) and biologically-moving objects (Vallortigara et al. 2005). For young chicks, approaching choices are particularly important as they can influence imprinting. Filial imprinting is a process through which young chicks develop a strong social attachment, including following responses, to the first conspicuous objects they encounter in their life (for general reviews on chick's development and imprinting see Rogers (1995), Bolhuis (1991) and McCabe (2013)). Although chicks can imprint on a variety of objects – including both natural and artificial objects –, specific colours, shapes, size and motion types induce stronger imprinting (see Introduction). Chicks' predispositions produce a bias in favour of naturalistic objects compared to artificial objects, as shown by the fact that once imprinted on a naturalistic object chicks cannot reverse their preference for an artificial object (Boakes and Panter 1985; Bolhuis and Trooster 1988) or have a delayed reversal (de Vos and van Kampen 1993), although the opposite seems to be easier.

Little is known though on the spontaneous preferences of chicks for approaching hollow or filled objects. This property can be particularly relevant to orient filial responses, because the presence of innards is associated with animate objects (Gelman 1990), that in the wild include social partners. Moreover, it has been observed that preschool children can reason about inside and outside features of objects (Gelman and Wellman 1991), 14-month-old babies associate an object's behaviour more with internal than with external features (Newman et al. 2008), and preverbal infants (8-month-old) expect animate objects to possess insides (Setoh et al. 2013). In the case of human babies, spontaneous preferences in the absence of previous experience with hollow or filled objects have not been assayed. We wondered whether the mere presence/absence of visible innards might trigger spontaneous approach preferences of young chicks for the first conspicuous objects encountered in their life, or whether experience might bias chicks preferences about the innards of social partners. To this aim we tested naïve and imprinted chicks using as hollow or filled objects orange cylinders of the size that can elicit filial responses.

In our experiments we consistently observed a preference of naïve chicks for approaching

hollow objects. The same preference held for chicks that during imprinting had been exposed

to objects occluded on their sides, that therefore were not explicitly filled or hollow, but were

clearly approached. The preference for hollow objects was reduced when chicks were

imprinted for 24 hours on filled objects, suggesting that chicks are sensitive to this feature of

social partners, and that even a brief experience can modify preferences for hollow/filled

objects. Yet, we did not observe an increase of the preference for hollow objects after

imprinting on hollow objects, and difference in performance between chicks imprinted on

hollow and filled objects was not strong. This suggests that after imprinting takes place, the

difference between hollow and filled imprinting objects is less salient, namely that objects

that differ only for the property of being filled of hollow are not categorized as markedly

different.

To establish which property of hollow objects was attracting chicks we ran a series of

subsequent experiments to investigate whether chicks were attracted by hollow objects as

hiding cavities, and/or whether the brightness and contrast of hollow objects were attractive

cues that triggered exploration. Although inexperienced chicks spontaneously recognize the

properties of occluding objects, and search objects behind barriers that completely occlude

them (Chiandetti and Vallortigara 2011), in our experiments chicks did not prefer larger

hollow objects, in which they could more easily hide, to smaller hollow objects. This suggests

that the preference for hollow objects is not mainly driven by the possibility to hide into them.

On the contrary, chicks were more attracted by darker insides or darker "caps". The attractive

feature of hollow objects could be either the darker part inside the object (its shadows, which

are a depth cue), or the higher contrast introduced by the presence of shadows. If the contrast

but not the lower brightness was attracting the chicks, we expected them to have no

preference when facing a choice between two scenes with the same (but opposite) contrast: a

white disk on a black background and a black disk on a white background. Instead, in this

setting chicks strongly preferred the black disk on a white background, suggesting that lower

brightness of an object but not the contrast per se is attractive for chicks.

To sum up, naïve chicks exhibited a consistent preference for hollow objects, which was

mainly mediated by the lower brightness on the insides, probably interpreted as a depth cue.

This preference could be modified by imprinting experience, simply by exposing chicks to a

filled object for 24 hours. At least for still objects such as the stimuli used in our experiments,

the property of being "filled" does not make objects more attractive as social partners for

chicks of the domestic fowl. Further experiments should clarify whether the preference for

hollow *vs.* filled objects is modified introducing cues of animacy, such as the presence of

movement or face configurations in the presented objects.

REFERENCES

Boakes R, Panter D (1985) Secondary imprinting in the domestic chick blocked by previous

exposure to a live hen. Anim Behav 33:353–365. doi: 10.1016/S0003-3472(85)80059-2

Bolhuis JJ (1991) Mechanisms of avian imprinting: a review. Biol Rev 66:303–345. doi:

10.1111/j.1469-185X.1991.tb01145.x

Bolhuis JJ, Trooster WJ (1988) Reversibility revisited: stimulus-dependent stability of filial

preference in the chick. Anim Behav 36:668–674. doi: 10.1016/S0003-3472(88)80149-0

Chiandetti C, Vallortigara G (2011) Intuitive physical reasoning about occluded objects by

inexperienced chicks. Proc R Soc B Biol Sci 278:2621–7. doi: 10.1098/rspb.2010.2381

de Vos GJ, van Kampen HS (1993) Effects of primary imprinting on the subsequent

development of secondary filial attachments in the chick. Behaviour 3-4:245–263. doi:

10.1017/CBO9781107415324.004

Dekker T, Revadi S, Mansourian S, et al (2015) Loss of Drosophila pheromone reverses its role

in sexual communication in *Drosophila suzukii*. Proc Biol Sci 282:20143018. doi:

10.1098/rspb.2014.3018

Fabricius E, Boyd H Experiments on the following-reaction of ducklings. Wildfowl Trust Annu

Rep 84–89.

Gelman R (1990) First principles organize attention to and learning about relevant data:

Number and the animate-inanimate distinction as examples. Cogn Sci 14:79–106. doi:

10.1016/0364-0213(90)90027-T

Gelman SA, Wellman HM (1991) Insides and essences: Early understandings of the non-

obvious. Cognition 38:213–244.

Giurfa M, Nunez J, Chittka L, Menzel R (1995) Colour preferences of flower-naive honeybees. J

Comp Physiol A. doi: 10.1007/BF00192415

Ham AD, Osorio D (2007) Colour preferences and colour vision in poultry chicks. Proc R Soc B

Biol Sci 274:1941–8. doi: 10.1098/rspb.2007.0538

Horn G (1985) Memory, imprinting, and the brain an inquiry into mechanisms. Clarendon

Press, Oxford

Johnson MH, Horn G (1988) Development of filial preferences in dark-reared chicks. Anim

Behav 36:675–683. doi: 10.1016/S0003-3472(88)80150-7

Keesey IW, Knaden M, Hansson BS (2015) Olfactory Specialization in *Drosophila suzukii*

Supports an Ecological Shift in Host Preference from Rotten to Fresh Fruit. J Chem Ecol.

doi: 10.1007/s10886-015-0544-3

Knaden M, Strutz A, Ahsan J, et al (2012) Spatial Representation of Odorant Valence in an

Insect Brain. Cell Rep 1:392–399. doi: 10.1016/j.celrep.2012.03.002

Mascalzoni E, Regolin L, Vallortigara G (2010) Innate sensitivity for self-propelled causal

agency in newly hatched chicks. Proc Natl Acad Sci U S A 107:4483–5. doi:

10.1073/pnas.0908792107

McCabe BJ (2013) Imprinting. Wiley Interdiscip Rev Cogn Sci 4:375–390. doi:

10.1002/wcs.1231

Miura M, Matsushima T (2016) Biological motion facilitates imprinting.

Newman GE, Herrmann P, Wynn K, Keil FC (2008) Biases towards internal features in infants'

reasoning about objects. Cognition 107:420–32. doi: 10.1016/j.cognition.2007.10.006

Raine NE, Chittka L (2007) The Adaptive Significance of Sensory Bias in a Foraging Context:

Floral Colour Preferences in the Bumblebee Bombus terrestris. PLoS One 2:e556. doi:

10.1371/journal.pone.0000556

Rogers LJ (1995) The Development of Brain and Behaviour in the Chicken. CAD International,

Wallingford

Rosa Salva O, Mayer U, Vallortigara G (2015) Roots of a social brain: Developmental models of

emerging animacy-detection mechanisms. Neurosci Biobehav Rev 50:150–168. doi:

10.1016/j.neubiorev.2014.12.015

Rosa-Salva O, Regolin L, Vallortigara G (2010) Faces are special for newly hatched chicks:

evidence for inborn domain-specific mechanisms underlying spontaneous preferences

for face-like stimuli. Dev Sci 13:565–77. doi: 10.1111/j.1467-7687.2009.00914.x

Schaefer H, Hess H (1951) Color Preferences in Imprinting Objects. Z Tierpsychol 16:161–172.

Schulman AH, Hale BE, Graves H. (1970) Visual stimulus characteristics for initial approach

response in chicks (*Gallus domesticus*). Anim Behav 18:461–466.

Setoh P, Wu D, Baillargeon R, Gelman R (2013) Young infants have biological expectations

about animals. Proc Natl Acad Sci U S A 110:15937–42. doi: 10.1073/pnas.1314075110

Vallortigara G (2012a) Core knowledge of object, number, and geometry: A comparative and

neural approach. Cogn Neuropsychol 29:213–236. doi: 10.1080/02643294.2012.654772

Vallortigara G (2012b) Aristotle and the chicken: Animacy and the origins of beliefs. In: Fasolo

A (ed) The Theory of Evolution and its Impact. Springer International Publishing, New

York, pp 189–200

Vallortigara G, Regolin L, Marconato F (2005) Visually inexperienced chicks exhibit

spontaneous preference for biological motion patterns. PLoS Biol 3:e208. doi:

10.1371/journal.pbio.0030208

Versace E, Reisenberger J (2015) Large-scale assessment of olfactory preferences and

learning in *Drosophila melanogaster*: behavioural and genetic components. PeerJ

3:e1214. doi: 10.7717/peerj.1214

Versace E, Vallortigara G (2015) Origins of knowledge: Insights from precocial species. Front

Behav Neurosci 9:338. doi: 10.3389/fnbeh.2015.00338

Acknowledgements

G.V. was funded by the ERC Advanced Grant ERC-2011-ADG_20110406, Project No:

295517, PREMESOR