Coherent synthesis of genomic associations with phenotypes and home environments

Jesse R. Lasky[1,2], Brenna R. Forester[3,4], and Matthew Reimherr[5]

5   [1]Department of Biology
Pennsylvania State University
408 Life Sciences Building
University Park PA 16802-5301

10   [2]Correspondence: jrl35@psu.edu, Phone: (814) 863-5318, Fax: (814) 865-9131

[3]Nicholas School of the Environment
Duke University
450 Research Drive
15   Durham, NC 27708-0001

[4]Current address: Department of Biology
Colorado State University
1878 Campus Delivery
20   Fort Collins, CO 80523

[4]Department of Statistics
Pennsylvania State University
411 Thomas Building
25   University Park PA 16802-5301

Keywords: landscape genomics, reciprocal transplant, genome-wide association study, fitness tradeoff, flowering time, plant defense, plasticity

30   Running title: Synthesis of genome, environment, & phenotype

Abstract

35　Local adaptation is often studied via 1) multiple common garden experiments comparing

performance of genotypes in different environments and 2) sequencing genotypes from

multiple locations and characterizing geographic patterns in allele frequency. Both

approaches aim to characterize the same pattern (local adaptation), yet the

complementary information from each has not yet been coherently integrated. Here, we

40　develop a genome-wide association model of genotype interactions with continuous

environmental gradients (G×E), *i.e.* reaction norms. We present an approach to impute

relative fitness, allowing us to coherently synthesize evidence from common garden and

genome-environment associations. Our approach identifies loci exhibiting environmental

clines where alleles are associated with higher fitness in home environments. Simulations

45　show our approach can increase power to detect loci causing local adaptation. In a case

study on *Arabidopsis thaliana*, most identified SNPs exhibited home allele advantage and

fitness tradeoffs along climate gradients, suggesting selective gradients can maintain

allelic clines. SNPs exhibiting G×E associations with fitness were enriched in genic

regions, putative partial selective sweeps, and associations with an adaptive phenotype

50　(flowering time plasticity). We discuss extensions for situations where only adaptive

phenotypes other than fitness are available. Many types of data may point toward the loci

underlying G×E and local adaptation; coherent models of diverse data provide a

principled basis for synthesis.

Introduction

55   Populations commonly exhibit phenotypic differences, often due to local adaptation to

environment (Leimu & Fischer 2008; Hereford 2009). Local adaptation is defined as a

genotype-by-environment interaction (G×E) for fitness that favors home genotypes

(Kawecki & Ebert 2004). Local adaptation has long interested empirical and theoretical

biologists (Clausen *et al.* 1940, 1948; Levene 1953; Slatkin 1973). However, little is

60   known about the genomic basis of local adaptation, such as genetic architecture, major

molecular mechanisms, and the extent to which genomic divergence among populations

is driven by local adaptation. Because local adaptation involves organismal responses to

environmental gradients, understanding the mechanisms of local adaptation has important

applications in agriculture and biodiversity conservation (Aitken & Whitlock 2013; van

65   Oppen *et al.* 2015; Lasky *et al.* 2015). Additionally, G×E are important in human

phenotypes like disease (Anastasi 1958; Hunter 2005; Gage *et al.* 2016). Understanding

the genomic basis of G×E is an emerging area of biomedical research (Thomas 2010;

Keller 2014) as are the genomics of local adaptation (reviewed by Des Marais *et al.* 2013;

Manel & Holderegger 2013; Tiffin & Ross-Ibarra 2014; Adrion *et al.* 2015; Bragg *et al.*

70   2015; Hoban *et al.* 2016).

A central question in local adaptation is whether selective gradients alone can

maintain allelic clines at individual loci, or whether stochastic processes, like limited

dispersal, are required to explain clines at individual loci causing local adaptation

(Mitchell-Olds *et al.* 2007; Anderson *et al.* 2011b). If selective gradients cause rank

75   changes in alleles with the highest relative fitness at an individual locus, selection may

maintain a cline, a pattern known as genetic tradeoff or antagonistic pleiotropy (Ågren *et*

al. 2013). Detecting loci that exhibit antagonistic pleiotropy in locally adapted systems has been challenging, partly due to limited statistical power of approaches that conduct multiple tests of significance for opposing fitness effects in different environments

80   (Fournier-Level *et al.* 2011a; Anderson *et al.* 2013).

Common garden experiments have long been employed to characterize genetic variation in phenotypes (Langlet 1971). In particular, reciprocal common gardens at multiple positions along environmental gradients are a powerful tool to reveal local adaptation (Clausen *et al.* 1940, 1948). One approach to identifying the loci underlying

85   local adaptation is to combine fitness data from multiple common garden experiments with genomic data (Lowry & Willis 2010; Fournier-Level *et al.* 2011a; Anderson *et al.* 2011a; Ågren *et al.* 2013). However, common gardens are logistically challenging and it is unclear how the typically small spatiotemporal scales of common gardens relate to the scales of processes that generate local adaptation in the wild (Weigel & Nordborg 2015).

90   An alternative to discovering genetic and ecological mechanisms of local adaptation is to study changes in allele frequency along environmental gradients (Hedrick *et al.* 1976; Tiffin & Ross-Ibarra 2014; Adrion *et al.* 2015; Bragg *et al.* 2015; Rellstab *et al.* 2015; Hoban *et al.* 2016). In this approach, known as a genome-environment association study, individuals are sequenced from multiple locations along environmental

95   gradients. Genetic markers and environmental gradients showing the strongest correlations are then considered as potentially involved in local adaptation (*e.g.* Hancock *et al.* 2008, 2011; Eckert *et al.* 2010; Turner *et al.* 2010; Coop *et al.* 2010; Lasky *et al.* 2012; Jones *et al.* 2012; Fitzpatrick and Keller 2015). A challenge of both traditional association studies (genome-phenotype) and genome-environment association studies is

100   that the genomic variation is observational and is not experimentally randomized (as

opposed to linkage mapping with experimental crosses, Devlin & Roeder 1999; Hancock

et al. 2008; Kang *et al.* 2008; Nordborg & Weigel 2008). As a result, many loci may

show spurious associations with phenotypes or with environment (Price *et al.* 2010;

Schoville *et al.* 2012; Bragg *et al.* 2015). Spurious associations are particularly

105   problematic for environmental gradients that are spatially autocorrelated due to

confounding with population structure (Schaffer & Johnson 1974). A technique for

dealing with this confounding is to control for putative population structure when testing

associations (Coop *et al.* 2010) by controlling for genome-wide relatedness (*e.g.*

estimated by identity-in-state) among accessions (Yoder *et al.* 2014; Lasky *et al.* 2014).

110   Understanding the genomic basis of adaptation may benefit from synthesizing

lines of evidence, for example by combining multiple types of genome scans to

strengthen the evidence that a locus is under selection (Lasky *et al.* 2014; Evans *et al.*

2014). For example, researchers have identified overlap between outliers for selection

statistics and markers associated with putatively adaptive phenotypes (Horton *et al.* 2012)

115   or between SNPs associated with phenotypes and those associated with climate gradients

(Berg & Coop 2014). Lasky *et al.* (2015) used a Bayesian approach to combine

associations with phenotype and environment, first calculating climate associations and

then using each marker's association to determine the prior probability it was associated

with G×E for adaptive phenotypes, yielding a posterior. Although combining multiple

120   lines of evidence is potentially useful, the quantitative approaches in past studies have

often been *ad hoc* and lacked reasoned principles. Here we develop a modeling

framework to conduct genome-wide association scans for G×E while coherently

synthesizing multiple data types. Existing approaches to genome-wide association studies

(GWAS) with G×E (sometimes referred to as genome-wide interaction studies, GWIS)

125   have dealt with categorical nominal environments (Murcray *et al.* 2009; Thomas 2010;

Korte *et al.* 2012; Gauderman *et al.* 2013; Marigorta & Gibson 2014), benefiting from the

statistical convenience of modeling phenotypes in different environments as correlated

traits (Falconer 1952). Association models have not been applied to G×E along

continuous environmental gradients, such as modeling SNP associations with reaction

130   norms (Jarquín *et al.* 2014; Tiezzi *et al.* 2017). Despite the existence of studies where

fitness was measured in multiple common gardens for diverse genotyped accessions

(Fournier-Level *et al.* 2011a), studies where linkage mapping was conducted for fitness at

multiple sites (Ågren *et al.* 2013), and studies where authors conducted association

mapping for G×E effects on phenotypes (Li *et al.* 2014), we found no example of

135   association studies of G×E for fitness, which is the basis of local adaptation.

The underlying processes generating local adaptation are the same regardless of

the approach used for inference, be it genome-environment association or common

gardens. Thus it is natural to synthesize data from multiple approaches. Furthermore, by

combining data types into a single inferential framework we may increase our power to

140   find causal loci. Here, we simultaneously leverage data from multiple common gardens

and genome-environment associations by using a new approach to imputing relative

fitness when common gardens are missing (Figure 1). In the remainder, we describe our

approach, present test cases using simulations and published data on *Arabidopsis thaliana*

(hereafter Arabidopsis), and discuss extensions.

Methods

Genome-wide association study of G×E effects on fitness

In common garden experiments, environment is often treated as a factor. When more than two gardens are conducted, variation among them may be considered in a more general fashion. For a given environmental gradient, each common garden may be located along the gradient according to its conditions. Describing common gardens as such may be informative about the specific ecological mechanisms driving selective gradients, taking advantage of the ordered nature of gardens' environments. We leverage multiple common garden experiments to identify markers (single nucleotide polymorphisms, SNPs) that show the strongest G×E effects, loci where allelic state shows the strongest interaction with environment in its association with fitness.

Local adaptation requires a genotype by environment interaction for fitness at the whole genome-level. Variation in individual phenotypes from multiple environments can be separated into components determined by genotype, environment, and G×E (Yates & Cochran 1938; Falconer 1952). To assess this interaction at an individual locus, one can assume that the relative fitness of individual i in a single location, w_i, satisfies the linear model

$$w_i = \alpha + \beta_E E_i + \beta_G G_{i,l} + \beta_{G \times E} G_{i,l} E_i + \varepsilon_i,$$

(eqn 1)

where $G_{i,l}$ is the genotype of individual i at locus l and E_i is the value of a single environmental variable at the location where w_i was measured. The β_E parameter gives the effect of environment on fitness and β_G gives the effect of genotype on fitness. Our primary interest lies in the $\beta_{G \times E}$ parameter, which gives the strength and direction of G×E

effects; $\beta_{G \times E}$ determines how responses to environmental gradients are mediated by

170   genotype. The term α gives the fitness intercept. We assume that the vector of errors, $\boldsymbol{\varepsilon}$, can be expressed as

$$\boldsymbol{\varepsilon} = \boldsymbol{E}\boldsymbol{v} + \boldsymbol{e}$$

where $\boldsymbol{E}$ is a diagonal matrix of the environmental values, and

$$\boldsymbol{v} \sim N_n(0, \sigma^2_{G \times E} \boldsymbol{K}) \quad \boldsymbol{e} \sim N_n(0, \sigma^2_e \boldsymbol{I}).$$

175   (eqn 2)

Here $\boldsymbol{v}$ and $\boldsymbol{e}$ are independent. The $n \times n$ matrix $\mathbf{K}$ is calculated as the genome-wide identity in state for each pair of the sampled n accessions (Kang *et al.* 2008). Random effects $\boldsymbol{v}$ are included because a substantial portion of G×E may be associated with population structure (Lasky *et al.* 2015); naively applying standard F-tests to assess the

180   interaction effects can result in a dramatic increase in false positive rates. To ameliorate this issue, the random effect $\boldsymbol{v}$ represents the genetic background interactions with environment (G×E, magnitude of their variance determined by $\sigma^2_{G \times E}$), while $\boldsymbol{e}$ represents the independent and identically distributed error in the model (variance determined by σ^2_e). However, incorporating random effects may also decrease power when causal loci

185   covary with genomic background.

Coherent synthesis of common gardens and genome-environment associations via imputation

We now tackle the goal of synthesizing genome-environment associations and G×E

190   observed across common gardens. A challenge in synthesizing these approaches is that genome-environment studies are purely observational and lack common garden

experiments. However, an implicit assumption in studies of genome-environment associations is that local adaptation occurs; if a common garden were conducted at each location where wild genotypes are collected, the home genotype would tend to be most

195 fit. Our imputation approach makes this assumption explicit. A formal consequence of this assumption is an (imputed) observation of highest relative fitness for genotypes at home. We combine this imputation with two observed datasets: genomic markers and environment of origin (Figure 1). Next, we scale relative fitness within each common garden so that the maximum observed fitness is given a relative fitness of unity, yielding

200 a metric that can be directly observed or imputed in each type of study (common garden or genome-environment association). For imputation, we then assume that each genotype collected in the wild is locally adapted at its home and thus has a relative fitness = 1 (Figure 1). After imputation, we can calculate marker associations with G×E for fitness, where each fitness observation arises from either (A) observations on a given genotype

205 by common garden combination or (B) imputation on a given genotype collected from its natural home and subsequently sequenced.

Fitting models and comparing approaches

We compared four reaction norm approaches to genome-wide G×E association studies in

210 addition to the more common approach of genome-environment associations. In Approach 1, we ignored potential confounding of population structure, using least squares to fit the model in eqn 1 where $\boldsymbol{\varepsilon}$ is normal, independent, and identically distributed (excluding random effects $\boldsymbol{v}$), but only including observed fitness data from common gardens and excluding imputed fitness data. In Approach 2, we again used least

215   squares but included imputed fitness data; these imputations using information from the

ancillary geographic data could possibly reduce false positives. In Approach 3, we fit the

full mixed-effects model (including random effects v), but excluded imputed fitness. In

Approach 4, we fit the full mixed-effects model while including imputed fitness data. To

test a genome-environment association approach (Approach 5), we also compared

220   associations between SNPs and home environments used a mixed-effects model in an

approach akin to traditional association mapping but where environment is substituted for

phenotype (Yoder *et al.* 2014; Lasky *et al.* 2014).

To improve computation time for the mixed-model approaches (3-5), we used the

method of Kang *et al.* (2010) and first fit the random effects with covariance determined

225   by kinship, and then fixed these effects while testing the effects of each SNP on the

phenotype. We included the environmental covariate effect in this initial step, following

the recommendation of Kang *et al.* (2010) for fitting additional non-SNP covariates. In

other words, we first fit the model:

$$w_i = \alpha + \beta_E E_i + \varepsilon_i,$$

230   (eqn 3).

Eqn. 3 is the same as Eqn. 1 but with genetic effects omitted. We obtained parameter

estimates $\hat{\alpha}, \widehat{\beta_E}, \hat{\sigma}_e^2, \hat{\sigma}_{G \times E}^2$. We then take the variance parameter estimates and use them to

estimate the remaining slope coefficients in eqn 1 using generalized least squares.

Because inclusion of the β_E term in ordinary least squares regression (Approaches 1-2)

235   led to poor model fit, we excluded the term from those approaches. We fit the discussed

mixed-model using Minimum Norm Quadratic Unbiased Estimation, MINQUE (Rao

1971; Brown 1976; Reimherr & Nicolae 2016). This approach is equivalent to restricted

maximum likelihood, REML, but rephrased in a way that more fully exploits the linearity

of the model, resulting in a flexible framework that can be quickly computed.

Simulation

We used simulations to demonstrate how our imputation technique can improve power to

identify loci causing G×E for fitness. To assess scenarios with varying strength of local

adaptation, we tested simulations of varying dispersal distances. We used results of

245   Forester *et al.* (2016), who previously simulated local adaptation along a continuous

environmental gradient (Landguth & Cushman 2010). Specifically, we used genome and

environment data from Forester *et al.* (2016) and here simulated new common garden

data. Forester *et al.* (2016) simulated 100 bi-allelic loci, one of which was under selection

(more detail is found in Forester *et al.* 2016 and the Supplement for this paper). In the

250   simulation, selection changed linearly along an environmental gradient, with *AA* and *aa*

genotypes favored at different ends of the gradient and *Aa* genotypes under uniform

selection across the gradient (Fig. S1). We tested three values of dispersal parameters

(low to high), resulting in varying strength of local adaptation, with the Pearson's

correlation between selected locus and selective gradient equal to 0.28, 0.24, and 0.11,

255   respectively.

Here, we sampled 250 individuals randomly. We then located four common

gardens at equal intervals along the gradient, encompassing the extremes of the selection

surface (Fig. S1). For the moderate dispersal and local adaptation scenario, we tested the

effect of common gardens that sample only half the environmental gradient (Fig. S1). For

260   the gardens, we subsampled 100 individuals from the full 250, and then averaged fitness

for 25 clones of each individual (each with the identical adaptive genotype of their parent clone) in each common garden using the selection model of Forester *et al.* (2016). After imputing fitness for the 250 individuals in their home environments, we had a total of 650 observations of fitness × location (250 imputed observations from individuals sampled across the landscape + 400 real observations arising from 100 clones in each of four common gardens).

For both the simulations and the *Arabidopsis* case study, we focus on the 1% of SNPs with the lowest p-values and their role in local adaptation. In simulations with 100 SNPs, this was equivalent to the lowest p-value SNP. To determine false positive rates in simulations, we calculated the proportion of simulations for a given scenario where a neutral (as opposed to a causal) SNP had the lowest p-value for $\beta_{G\times E}$.

Model extensions

Two extensions to our approach could increase its generality. First, one could treat unobserved fitness of a genotype in its home environment as a free parameter. To constrain estimates of unobserved fitness one could use informative priors, such that the prior probability of relative fitness at home for each genotype would be monotonically increasing, *i.e.* local adaptation is the most likely state, but minor maladaptation is common. Inferences about unobserved fitness could be further constrained using hierarchical models, such that home fitness parameters for multiple genotypes arise from a common distribution (Gelman & Hill 2007). Relaxing the assumption of perfect local adaptation would also generate less biased, if less precise, parameter estimates for $\beta_{G\times E}$,

which are currently conservative when imputation of local adaptation for maladapted

genotypes pushes $\beta_{G \times E}$ toward zero and weakens estimates for selective gradients.

285   Second, when fitness is not measured, components of fitness (*e.g.* survival) or

traits thought to be locally adaptive (*e.g.* physiological) can be measured and used to infer

the genomic basis of local adaptation. For example, instead of modeling

SNP×environment associations with fitness, one could model SNP×environment

associations with components of fitness measured in common gardens, and estimate

290   unobserved traits for sequenced genotypes using informative priors. To be clear, in our

case study of Arabidopsis, we had near but not complete lifetime fitness data (missing

germination stage). Here we do not fit these model extensions to data, given the current

computational challenge of fitting many more parameters in a Bayesian framework.

295   *Case study: local adaptation to climate in* Arabidopsis thaliana

We applied these approaches to published data from studies of *Arabidopsis thaliana* in its

native Eurasian range. Fournier-Level *et al.* (2011a) conducted replicated common

gardens at four sites across Europe: Spain, England, Germany, and Finland (Figure 2).

With these data, Fournier-Level *et al.* (2011a) and Wilczek *et al.* (2014) showed evidence

300   that genotypes are locally adapted to their home temperature and moisture regimes and

that alleles associated with high fitness in a given garden tended to be found nearer to that

garden than alternate alleles, suggesting these loci were involved in local adaptation. At

each site the authors transplanted 157 accessions (59 in the case of Finland) on a date in

the fall matched to germination of local winter-annual natural populations (Fournier-

305   Level *et al.* 2011a). The authors calculated survival (out of individuals surviving

transplant) and average fecundity (where individuals that died before reproducing had fecundity zero) giving an estimate of absolute fitness (excluding the seed to seedling transition, Fournier-Level *et al.* 2011a).

These accessions were part of a panel of 1,307 accessions from around the globe that were genotyped at ~250k SNPs using a custom Affymetrix SNP tiling array (AtSNPtile1), with 214,051 SNPs remaining after quality control (Figure 2, Horton *et al.* 2012). This array was generated by resequencing 19 diverse ecotypes from across the range of Arabidopsis (Kim *et al.* 2007). Of the 1,307 genotyped accessions, we used 1,001 accessions that were georeferenced and likely not contaminant lines (Anastasio *et al.* 2011), in addition to being from the native range in Eurasia (Hoffmann 2002; Lasky *et al.* 2012b), and excluding potentially inaccurate high altitude outliers (i.e. > 2000 m). After imputing fitness for accessions in their home environments we had a total of 1,531 observations of fitness × location (1001 imputed observations + 530 real observations arising from the four common gardens). We removed from association tests SNPs having minor allele frequency (MAF) below 0.01, though we also considered a more conservative threshold of MAF = 0.1.

We used climate and derived data from a previous study (Lasky *et al.* 2012b) using published global climate datasets (Hijmans *et al.* 2005; Zomer *et al.* 2008). Here we focus on four climate variables that differ among common gardens, are not strongly correlated, and may be involved in local adaptation: minimum temperature of the coldest month, average monthly minimum temperature in the growing season, coefficient of variation of monthly growing season precipitation, and aridity index.

Genome-wide G×E associations

330 We separately tested for each SNP's interaction with each of the four climate variables. For each of the four approaches using common garden data we fit a model for each combination of SNP and environmental variable. To characterize the types of patterns identified by our approach, we studied variation in the SNPs in the 0.01 lower tail of p-values for the hypothesis test that $\beta_{G×E} = 0$ (the coefficient for SNP×environment effects

335 on fitness) for each climate gradient. For these SNPs, we calculated whether the direction of allelic differentiation along environmental gradients was consistent with the sign of $\beta_{G×E}$. For example, if one allele was more common in accessions from warmer locations, we assessed whether that same allele showed an increase in relative fitness in warmer common gardens.

340 Next, we assessed whether our model predicted that different alleles were most fit in the two common gardens at either extreme of a climate gradient, *i.e.* whether the SNP was associated with rank changes in fitness that are consistent with genetic tradeoffs, versus a pattern where $\beta_{G×E}$ merely involved changes in fitness difference between alleles (such as conditional neutrality or variance changing G×E), the latter of which cannot

345 stably maintain local adaptation. For example, if one allele was estimated to be most fit in the coldest common garden, we determined whether a different allele was estimated to be most fit in the warmest common garden. Furthermore, we also quantified similarity (rank correlation in SNP scores and proportion of SNPs common to the strongest 0.01 tail of associations) between results from our G×E approach versus those from other recent

350 studies of association with home climate in Arabidopsis (Hancock *et al.* 2011; Lasky *et al.* 2012b, 2014).

Enrichment of strong SNP×environment associations across the genome

We studied whether loci we identified as likely being involved in local adaptation

355   exhibited supportive patterns in ancillary datasets. First, to assess whether our association

approach is capable of identifying the signal of local adaptation rather than spurious

background associations, we tested for enrichment of SNPs in genic (from transcription

start to stop sites) versus intergenic regions. These tests are based on the hypothesis that

loci involved in adaptation are on average more likely to be found near genes and linked

360   to genic variation, in comparison with loci evolving neutrally (Hancock *et al.* 2011;

Lasky *et al.* 2012b). For a test statistic, we calculated the portion of SNPs in the 0.01

lower p-value tail that were genic versus intergenic.

Second, we hypothesized that locally-adaptive alleles may have been subject to

partial (local) selective sweeps, especially given that much of Arabidopsis' Eurasian

365   range was recently colonized following the last glacial maximum. We tested for an

enrichment of significant (alpha = 0.05) pairwise haplotype sharing (PHS, Toomajian *et*

al. 2006) in the SNPs (using PHS calculated by Horton et al. 2012) showing the greatest

evidence of G×E for fitness. We also tested evidence that these SNPs are enriched for

significant (alpha = 0.05) integrated extended haplotype homozygosity (standardized,

370   iHS, Voight *et al.* 2006), an additional metric of partial sweeps. We used ancestral SNP

allele determinations from Horton *et al.* (2012, based on alignment with the *Arabidopsis*

lyrata genome) and the R package 'rehh' to calculate iHS (Gautier *et al.* 2012).

Third, we also studied whether loci we identified were associated with plasticity

in flowering time, a trait that plays a major role in local adaptation to climate in plants

375 (Hall & Willis 2006; Franks *et al.* 2007; Keller *et al.* 2012; Lowry *et al.* 2014). Recently

Li *et al.* (2014) tested the flowering time response of 417 natural accessions to simulated

warming (up to ~4ºC), and then identified SNP associations with changes in flowering

time across treatments, G×E for flowering time. We tested whether SNPs we identified as

having SNP×environment interactions for fitness (0.01 lower p-value tail) were enriched

380 in nominally significant associations (alpha = 0.05) with G×E for flowering time.

To generate a null expectation for each enrichment while maintaining a signal of

linkage disequilibrium in the null model, we circularly permuted SNP categories (*e.g.* as

genic versus intergenic, having significant iHS or not) along the genome and recalculated

the test statistics 10,000 times (Hancock *et al.* 2011; Lasky *et al.* 2012b).

Results

We compared four approaches to genome-wide G×E association study and one approach

for genotype-environment association. Approach 1 used only observed (excluding

imputed) fitness data, but no correction for population structure. Approach 2 used

390 observed and imputed fitness data, and no correction for population structure. Approach 3

fitted the full mixed-effects model, but only including observed fitness data from

common gardens, excluding imputed fitness. Approach 4 fitted the full mixed-effects

model while including both observed and imputed fitness data. Approach 5 used a mixed

model of genotype associations with environment (no common garden data).

Simulations

Across dispersal scenarios, we found that mixed models decreased false positive rates

and increased accuracy of inference as to the SNPs driving G×E for fitness (Figure 3).

When dispersal was highest and local adaptation weakest, all approaches exhibited an

400   increase in false positives compared to the moderate dispersal scenario. Among the

approaches using common garden data (Approaches 1-4), the mixed models (Approaches

3-4) generally had low false positive rates and thus high true positive rates compared to

models excluding random effects (Approaches 1-2). Based on the low false positive rates

and low p-values for causal SNPs in Approach 3, common garden data were a clear

405   source of statistical power to identify causal SNPs (Figure 3). Including imputed data

(Approach 4) further reduced false positive rates and resulted in no false positives under

the two lower dispersal scenarios. Genotype-environment associations that did not use

common garden data (Approach 5) had similarly low false positive rates (Table S5).

However, common garden data combined with imputations (Approach 4) yielded

410   stronger inference for SNPs driving G×E; causal SNPs had lower p-values (Figure 3,

median causal SNP p for low, medium, high dispersal scenarios: 3.2×10^{-6}, 4.5×10^{-13},

7.4×10^{-7}) compared to ignoring common garden data (Approach 5, median causal SNP p

for low, med., high dispersal scenarios: 2.3×10^{-7}, 1.9×10^{-10}, 1.2×10^{-5}). Under a scenario

of medium dispersal and common gardens that only covered half the gradient, false

415   positive rates were elevated for approaches excluding random effects (Approaches 1-2)

or excluding imputations (Approach 3) but not when imputations and random effects

were included (Approach 4, Figure S4).

Case study on Arabidopsis

420   We found that simple linear model tests (Approaches 1-2) of SNP×environment

interactions were highly enriched in very low p-values (Figure S5) relative to the

theoretical expectation. After incorporating the kinship×environment random effects (but

excluding imputed fitness observations, Approach 3), we found that SNP×environment

associations with fitness were closer to the theoretical expectation but still highly

425   enriched in low p-values for three climate variables. After incorporating imputed fitness

observations into the mixed model (Approach 4, right column, Figure S5), we found p-

value distributions hewed closer to the theoretical expectation and were slightly

conservative (under-enriched in low p-values) for two climate variables. These

approaches tended to identify different SNPs as having the strongest SNP×environment

430   associations with fitness (Table S6).

      Based on the results of our simulations and the p-value distributions noted above,

we focus the remainder of analyses on results from mixed models with imputed fitness

included (Approach 4). We found that climate variables differed in the importance of

kinship-climate interaction associations with fitness (proportion of variance in fitness

435   explained by random effects v), suggesting that population structure in Arabidopsis is

more strongly correlated with some climatic axes of local adaptation (G×E for fitness)

compared to other climate gradients. For growing season minimum temperatures,

kinship×environment interactions explained most of the variation in fitness ($R^2 = 0.78$,

Table 1, row 3). By contrast, kinship×environment interactions for fitness were weaker

440   along a gradient in winter minimum temperature ($R^2 = 0.07$).

      Approach 4 tended to identify SNPs where SNP×environment interactions

favored alleles in climates where they were relatively more common, that is, the sign of

allelic differences in home climates were matched by the sign of fitted mixed model

SNP×environment associations with relative fitness (Table 1, row 1, and see outlier

445   examples in Figure 4). In addition to characterizing SNP×environment associations, we

tended to identify SNPs where we estimated a rank change in relative fitness for alternate

alleles along the environmental gradient between the two extreme common gardens

(where the fitted model expectation was that the allele with higher fitness at one extreme

common garden differed from the allele with higher fitness at the other extreme, Table 1,

450   row 2). It appeared that the proportion of SNPs expected to show rank changes in relative

fitness among the common gardens was related to how much of each climate variable's

range was covered by gardens (Table 1, row 4). Thus the common gardens may have

been limited in their ability to capture rank changing of alleles at some loci involved in

local adaptation to aridity and growing season cold.

455   We found non-random, but very weak overlap between the SNPs we identified

and those outliers in previous analyses (Hancock *et al.* 2011; Lasky *et al.* 2012b, 2014).

When considering mixed model (Lasky *et al.* 2014) or partial Mantel (Hancock *et al.*

2011) SNP associations with the same climate variables (genome-environment

associations with no common garden data), we found significant overlap among the

460   previously identified SNPs in the 0.01 lower tail of p-values versus those in the 0.01 tail

identified here (permutation test, all p<0.05, Table S7). However, rank correlations

among SNP scores from previous approaches versus our current approach were very

weak (all rho < 0.2, Table S7, Figure S6).

SNP×environment associations with fitness are enriched in regions suggestive of local adaptation

We found that SNP×environment interactions for fitness were significantly enriched in genic regions (again focusing on Approach 4: mixed model including imputations, Table 2; for reference, SNPs identified via Approach 2, including imputation but without random effects, were not significantly enriched in genic regions). Additionally, we found that SNP×environment interactions for fitness were enriched for high pairwise haplotype sharing (PHS) and high integrated extended haplotype homozygosity (iHS, Table 2). Finally, SNPs associated with G×E for flowering time response to growing temperature (Li *et al.* 2014) tended to also have strong SNP×growing season minimum temperature interactions for fitness (p < 0.0002) but not for other climate variables. (Table 2) Enrichments reported above did not change qualitatively (with respect to statistical significance) when we only considered SNPs with MAF > 0.1.

SNP×environment associations with fitness identify genes potentially involved in local adaptation

Our approach identified a number of strong candidates for local adaptation at the top of lists of SNPs with the strongest SNP×environment associations with relative fitness (Tables S1-S4). For example, the top SNP associated with aridity interaction effects on fitness (chr. 4, position 11005059) fell within LESION SIMULATING DISEASE 1 (LSD1), which affects a number of traits in Arabidopsis, including survival and fecundity under drought (Wituszyńska *et al.* 2013; Szechyńska-Hebda *et al.* 2016, Figure 4), while the third SNP (chr. 2, position 7592008) fell within ATMLO8, MILDEW RESISTANCE

LOCUS O 8, homologous with barley MLO which controls resistance to the fungal

pathogen powdery mildew (Büschges *et al.* 1997). The top SNP associated with winter

490   cold interaction effects on fitness (chr. 5, position 7496047) falls within coding region of

WRKY38, involved in the salicylic acid pathway and pathogen defense (Kim *et al.*

2008), and was the same locus identified as most strongly associated with multivariate

climate in Lasky et al. (2012, Figure 4). The top SNP associated with variability in

growing season precipitation interaction effects on fitness (chr. 2, position 18504858)

495   falls 380 bp from ABA HYPERSENSITIVE GERMINATION 11, AHG11, which

mediates the effect of abscisic acid (ABA), a major hormone of abiotic stress response,

on germination (Murayama *et al.* 2012). The fifth highest SNP (and second highest locus)

associated with growing season cold interaction effects on fitness (chr. 3, position

8454439) fell within ABERRANT LATERAL ROOT FORMATION 5, ALF5, a gene

500   that confers resistance to toxins (Diener *et al.* 2001) belonging to the MATE gene family,

which play a variety of roles responding to environment (Shoji 2014).

Discussion

Genetic variation in environmental responses (G×E) is ubiquitous but its genetic and

505   physiological basis and role in local adaptation is poorly understood. Replicated common

garden experiments and genome scans for loci exhibiting evidence for local adaptation

have been important in understanding the genetic basis of G×E and local adaptation

(Hancock *et al.* 2008; Eckert *et al.* 2010; Turner *et al.* 2010; Fournier-Level *et al.* 2011a;

Lasky *et al.* 2012b; Ågren *et al.* 2013; Evans *et al.* 2014; Lasky *et al.* 2015). However,

510   the complementary information in common gardens and geographic variation in allele

frequency have not been coherently synthesized. Previous association studies of G×E

have modeled discrete, categorical environmental effects (Murcray *et al.* 2009; Thomas 2010; Korte *et al.* 2012; Marigorta & Gibson 2014). The modeling of G×E across discrete, categorical environments is often employed for mathematical convenience, as such a treatment allows the use of models designed for multiple phenotypes, where the same phenotype in different environments is considered as multiple phenotypes (Falconer 1952).

We demonstrated an approach to association study of G×E for fitness and an imputation technique that allowed us to coherently synthesize evidence from common gardens and genome-environment associations. Our imputation method relied on making explicit the implicit assumption of local adaptation that underlies genome-environment association studies (Coop *et al.* 2010; Hancock *et al.* 2011; Lasky *et al.* 2012b). Using simulation, we demonstrated that this imputation can increase power to identify SNPs causing G×E for fitness and local adaptation. Our approach also identified strong candidate genes in Arabidopsis associated with SNPs that exhibit fitness tradeoffs along climate gradients such that locally common alleles had greater relative fitness.

The relative information on selective and adaptive genetic mechanisms contained in the two datasets (common garden, geographical genomic) for a given system will be determined by several factors. First, the power of common gardens depends on the range of sampled covariates (genotype and environment). We found evidence with both our simulations and empirical case study that greater coverage of environmental gradients can increase power to detect causal loci. Similarly, power may be enhanced by including in common gardens a range of variation at locally adaptive loci using diverse germplasm from across gradients. However, alternate mechanisms of local adaptation across regions

535   and confounding between population structure and adaptive loci suggest that regional

stratification in scans for local adaptation may be more powerful (Horton *et al.* 2016).

Additionally, the power of common gardens is influenced by the match between

conditions in gardens and long-term natural selective gradients that give rise to local

adaptation (Weigel & Nordborg 2015), and the heritability of adaptive traits and fitness.

540   The information contained in genome-environment associations (and hence imputed

fitness data here), is influenced by the strength of local adaptation in sampled populations

(Figure 3), which itself is determined by steepness of selective gradients, the level of

gene flow, and time populations have had to evolve toward equilibrium allele frequencies

(Yeaman & Whitlock 2011; Lotterhos & Whitlock 2014; Forester *et al.* 2016). It is

545   important to recognize that our simulations covered a limited range of the parameter

space relevant in nature (genetic architecture of local adaptation, dimensionality of

environmental selective gradients, *etc.*). Here, populations were given time to reach

equilibrium (Forester *et al.* 2016), which likely enhanced the power of genotype-

environment associations compared to scenarios common in nature where populations are

550   still responding to environmental changes. Apart from information on genetic

mechanisms of G×E for fitness, common gardens afford a more direct opportunity to

study phenotypes under selection, as opposed to genotype–environment associations

where information on phenotype is limited to gene annotations.

Above we described a method of imputation based on the assumption of local

555   adaptation, *i.e.* home genotypes had greater fitness than away genotypes. However, local

adaptation in nature is typically imperfect, such that the optimal genotype for a given

location might not be the home genotype (Leimu & Fischer 2008; Hereford 2009). Local

adaptation may not occur for several reasons, such as gene flow across environmental gradients (Slatkin 1973), limited genetic variation (Barton 2001), temporal environmental

560   shifts, and other processes (Bridle & Vines 2007). The utility of our imputation is dependent on local adaptation occurring, just as in genome-environment association studies, thus both approaches are useful for identifying loci subject to selective gradients when local adaptation occurs. Our imputation can be considered a heuristic to be improved by further development.

Genotype-by-environment interactions in genome-wide association studies
Recent advances in association models have included explicit modeling of categorical G×E (Murcray *et al.* 2009; Thomas 2010; Korte *et al.* 2012; Marigorta & Gibson 2014; Li *et al.* 2014; Kooperberg *et al.* 2016; Windle 2016), but to our knowledge there are no

570   published genome-wide association studies accounting for genotype interactions with continuous environmental gradients (a reaction norm approach, cf. Jarquín *et al.* 2014; Tiezzi *et al.* 2017). By employing a reaction norm approach to G×E (as we did here), models can be applied to prediction at new sites, which is not possible using categorical, correlated trait approaches to G×E (Falconer 1952; Korte *et al.* 2012) where sites are

575   treated as idiosyncratic. Some of the aforementioned categorical treatments of SNP×environment interactions were used in association studies for human disease. However, many of the environmental variables that mediate genetic risk of disease are continuous in nature, such as exposure to ultraviolet radiation and tobacco smoke. Future research on local adaptation and human disease may benefit from exchange of

580   approaches given the shared importance across disciplines of understanding the genomic

basis of G×E.

Case study on Arabidopsis thaliana

Our approach identified many SNPs where allelic variation was associated with rank-

585   changing relative fitness tradeoffs along climate gradients (*e.g.* all 214 of the SNPs with

strongest interaction with winter minimum temperature association for fitness, *i.e.* 0.001

quantile), loci where selective gradients may stably maintain population differentiation

(Anderson *et al.* 2011b; Ågren *et al.* 2013). Studies of local adaptation genomics often

find limited evidence for loci with such fitness tradeoffs (antagonistic pleiotropy). A

590   previous study of the common garden data used here (Fournier-Level *et al.* 2011a) found

that the SNPs with the strongest association with fitness in one common garden were

rarely among those with the strongest associations in another garden, which the authors

interpreted as evidence for conditional neutrality. However, the fact that a locus is not

among the strongest associated with fitness at an individual site does not indicate the

595   locus is neutral at that site, it may simply be under relatively weaker selection (see Figure

S7 for example illustration). By contrast with previous approaches that model phenotypes

at a single site, our model was explicitly focused on detecting alleles with the strongest

evidence for SNP×climate interactions favoring home alleles. Thus our explicit model of

G×E is more likely to detect loci with patterns indicative of antagonistic pleiotropy

600   compared with approaches that model fitness in a single common garden at a time,

approaches that do not model G×E. Nevertheless, apparent tradeoffs at the level of

individual QTL or SNPs may be driven by complementary conditionally neutral mechanisms at tightly linked loci.

Local adaptation may often involve complex traits governed by many loci. Loci exhibiting antagonistic pleiotropy and loci exhibiting G×E but no tradeoffs (variance changing or conditionally neutral) may both underlie genome-level local adaptation. Note that our study, like that of Fournier-Level *et al.* (2011a) is based on association mapping, which may suffer from identification of more false positives compared with linkage mapping approaches (Hall *et al.* 2010; Anderson *et al.* 2013; Ågren *et al.* 2013). Follow-up experimental study of phenotypic effects of variation at individual loci is required to confirm the results of association mapping (Verslues *et al.* 2014; Broekgaarden *et al.* 2015). The SNPs that exhibited the strongest evidence for SNP×climate interaction effects on fitness often fell within the coding regions of strong candidate genes based on known roles in environmental responses, suggesting our approach is a useful for identifying loci underlying local adaptation.

We found evidence that SNP×climate interaction effects on fitness were enriched in genic regions, suggesting that our model captured a signal of local adaptation rather than population structure. We found that enrichments in genic SNPs only emerged after using a mixed model to control for the putative effects of population structure (genome-wide similarity), suggesting that the genic-enriched patterns of divergence we modeled were not simply associated with overall patterns of among-population divergence. This enrichment is consistent with other findings in Arabidopsis (Hancock *et al.* 2011; Lasky *et al.* 2012b) and other species (Coop *et al.* 2009; Fumagalli *et al.* 2011; Lasky *et al.* 2015, but see Pyhäjärvi *et al.* 2013). We do not interpret this enrichment as indicating

625   that changes in amino acid sequences are more important than regulatory evolution in local adaptation, but rather as supporting the hypothesis that local adaptation is more likely to involve sequence evolution near genes as opposed to locations farther from genes, where many intergenic SNPs are found.

We found evidence that loci we identified as candidates were enriched in

630   evidence for partial selective sweeps (PHS and iHS statistics), suggesting that recent sweeps in particular environments are an important mode of local adaptation (Voight *et al.* 2006; Toomajian *et al.* 2006). These local sweeps may be expected based on the range dynamics of Arabidopsis, which has colonized much of its Eurasian range following the retreat of glaciers (Sharbel *et al.* 2000), a process that likely involved recent local

635   adaptation. It is important to note that extended haplotype patterns suggestive of partial sweeps may occur at the shoulders (away from causal loci) of complete sweeps (Schrider *et al.* 2015), thus caution is warranted in attributing our observed PHS and iHS enrichment to localized sweeps versus global sweeps at nearby loci.

We found significant overlap between SNPs associated with G×E for fitness

640   along growing season cold gradients and SNPs associated with G×E for flowering time across growing season temperature treatments (Li *et al.* 2014). This concordance suggests that variants causing flowering time plasticity drive changes in fitness across temperature gradients. Thus the evolution of temperature-responsive plasticity in flowering time may be a mechanism of local adaptation to

645   environments that differ in growing season temperatures. For organisms inhabiting seasonal environments, timing of the life cycle can have large impacts on fitness. Previous common garden experiments have provided strong evidence that

flowering time is a central trait involved in local adaptation (Hall & Willis 2006; Franks *et al.* 2007; Keller *et al.* 2012; Lowry *et al.* 2014) with molecular study further
650   supporting the role of flowering time (Stinchcombe *et al.* 2004; Caicedo *et al.* 2004; Shindo *et al.* 2005; Lovell *et al.* 2013) and the role of plasticity (Fraser 2013; Lasky *et al.* 2014) in local adaptation to climate.

Though there was overlap with signal identified by previous approaches using the same data (Hancock *et al.* 2011; Lasky *et al.* 2012b, 2014), overlap was generally
655   weak, indicating our approach identified distinct loci. This weak overlap is likely due to the different statistical approaches used (partial Mantel, linear mixed models, and redundancy analysis) and the additional information contained in fitness data. Furthermore, the common garden data contain signatures of processes that differ to an unknown degree from processes generating genome-environment associations. Here, the
660   common garden data represented one generation of fitness variation among sites, whereas spatial genomic patterns accumulated due to multiple evolutionary mechanisms acting over thousands of generations.

Conclusions
665   Genome-wide association studies are a promising approach for identifying the genomic basis of local adaptation and G×E. Given that many selective gradients driving local adaptation are continuous in nature, reaction norm models of G×E across multiple common gardens are a key tool for quantifying mechanisms of local adaptation. Various approaches such as genome-wide expression profiling (Des
670   Marais *et al.* 2013), metabolomics (Meijón *et al.* 2016), and ecophysiology (Keller *et*

al. 2011) are useful for uncovering integrated mechanisms of local adaptation

Future approaches that use a principled basis for quantitative synthesis of patterns

in multiple data types (e.g. Levy Karin *et al.* 2017) may enhance our ability to

characterize mechanisms of adaptation across levels of biological organization.

Acknowledgements

We thank David Lowry, Thomas Juenger, and six anonymous reviewers for comments on

earlier versions of this manuscript.

680 **References**

Adrion, J. R., Hahn, M. W., & Cooper, B. S. (2015). Revisiting classic clines in
Drosophila melanogaster in the age of genomics. *Trends in Genetics*.
doi:10.1016/j.tig.2015.05.006

685 Ågren, J., Oakley, C. G., McKay, J. K., Lovell, J. T., & Schemske, D. W. (2013). Genetic
mapping of adaptation reveals fitness tradeoffs in Arabidopsis thaliana.
Proceedings of the National Academy of Sciences, *110*(52), 21077–21082.
doi:10.1073/pnas.1316773110

Aitken, S. N., & Whitlock, M. C. (2013). Assisted Gene Flow to Facilitate Local

690 Adaptation to Climate Change. *Annual Review of Ecology, Evolution, and
Systematics*, *44*(1), 367–388. doi:10.1146/annurev-ecolsys-110512-135747

Anastasi, A. (1958). Heredity, environment, and the question" How?". *Psychological
Review*, *65*(4), 197.

Anastasio, A. E., Platt, A., Horton, M., Grotewold, E., Scholl, R., Borevitz, J. O., …

695   Bergelson, J. (2011). Source verification of mis-identified Arabidopsis thaliana

accessions. *The Plant Journal*, *67*(3), 554–566. doi:10.1111/j.1365-

313X.2011.04606.x

Anderson, J. T., Lee, C.-R., & Mitchell-Olds, T. (2011). Life history QTLs and natural

selection on flowering time in Boechera stricta, a perennial relative of

700   Arabidopsis. *Evolution; International Journal of Organic Evolution*, *65*(3), 771–

787. doi:10.1111/j.1558-5646.2010.01175.x

Anderson, J. T., LEE, C.-R., RUSHWORTH, C. A., COLAUTTI, R. I., & MITCHELL-

OLDS, T. (2013). Genetic trade-offs and conditional neutrality contribute to local

adaptation. *Molecular Ecology*, *22*(3), 699–708. doi:10.1111/j.1365-

705   294X.2012.05522.x

Anderson, J. T., Willis, J. H., & Mitchell-Olds, T. (2011). Evolutionary genetics of plant

adaptation. *Trends in Genetics*, *27*(7), 258–266. doi:10.1016/j.tig.2011.04.001

Barton, N. H. (2001). Adaptation at the edge of a species' range. *SPECIAL*

PUBLICATION-BRITISH ECOLOGICAL SOCIETY, *14*, 365–392.

710   Berg, J. J., & Coop, G. (2014). A Population Genetic Signal of Polygenic Adaptation.

PLoS Genet, *10*(8), e1004412. doi:10.1371/journal.pgen.1004412

Bragg, J. G., Supple, M. A., Andrew, R. L., & Borevitz, J. O. (2015). Genomic variation

across landscapes: insights and applications. *New Phytologist*, n/a-n/a.

doi:10.1111/nph.13410

715   Bridle, J. R., & Vines, T. H. (2007). Limits to evolution at range margins: when and why does adaptation fail? *Trends in Ecology & Evolution*, *22*(3), 140–147. doi:10.1016/j.tree.2006.11.002

Broekgaarden, C., Bucher, J., Bac-Molenaar, J., Keurentjes, J. J. B., Kruijer, W., Voorrips, R. E., & Vosman, B. (2015). Novel Genes Affecting the Interaction

720   between the Cabbage Whitefly and Arabidopsis Uncovered by Genome-Wide Association Mapping. *PLoS ONE*, *10*(12), e0145124. doi:10.1371/journal.pone.0145124

Brown, K. G. (1976). Asymptotic Behavior of Minque-Type Estimators of Variance Components. *Ann. Statist.*, (4), 746–754. doi:10.1214/aos/1176343546

725   Büschges, R., Hollricher, K., Panstruga, R., Simons, G., Wolter, M., Frijters, A., … Schulze-Lefert, P. (1997). The Barley Mlo Gene: A Novel Control Element of Plant Pathogen Resistance. *Cell*, *88*(5), 695–705. doi:10.1016/S0092-8674(00)81912-1

Caicedo, A. L., Stinchcombe, J. R., Olsen, K. M., Schmitt, J., & Purugganan, M. D.

730   (2004). Epistatic interaction between Arabidopsis FRI and FLC flowering time genes generates a latitudinal cline in a life history trait. *PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA*, *101*(44), 15670–15675. doi:10.1073/pnas.0406232101

Clausen, J., Keck, D. D., & Hiesey, W. M. (1940). Experimental Studies on the Nature of

735   Species. I. Effect of Varied Environments on Western North American Plants. Carnegie Institution of Washington Publication No. 520. *Washington, DC.*

Clausen, J., Keck, D. D., & HIESEY, W. M. (1948). Experimental studies on the nature

of species. III. Environresponses of climatic races of Achillea. *Experimental*

Studies on the Nature of Species. III. Environresponses of Climatic Races of

740 *Achillea.*, (Publ. 581).

Coop, G., Pickrell, J. K., Novembre, J., Kudaravalli, S., Li, J., Absher, D., … Pritchard, J.

K. (2009). The Role of Geography in Human Adaptation. *PLoS Genet*, *5*(6),

e1000500.

Coop, G., Witonsky, D., Di Rienzo, A., & Pritchard, J. K. (2010). Using Environmental

745 Correlations to Identify Loci Underlying Local Adaptation. *Genetics*, *185*(4),

1411–1423. doi:10.1534/genetics.110.114819

Des Marais, D. L., Hernandez, K. M., & Juenger, T. E. (2013). Genotype-by-

Environment Interaction and Plasticity: Exploring Genomic Responses of Plants

to the Abiotic Environment. *Annual Review of Ecology, Evolution, and*

750 *Systematics*, *44*(1), 5–29.

Devlin, B., & Roeder, K. (1999). Genomic Control for Association Studies. *Biometrics*,

55(4), 997–1004. doi:10.1111/j.0006-341X.1999.00997.x

Diener, A. C., Gaxiola, R. A., & Fink, G. R. (2001). Arabidopsis ALF5, a Multidrug

Efflux Transporter Gene Family Member, Confers Resistance to Toxins. *The*

755 *Plant Cell*, *13*(7), 1625–1637. doi:10.2307/3871390

Eckert, A. J., BOWER, A. D., GONZÁLEZ-MARTÍNEZ, S. C., WEGRZYN, J. L.,

COOP, G., & NEALE, D. B. (2010). Back to nature: ecological genomics of

loblolly pine (Pinus taeda, Pinaceae). *Molecular Ecology*, *19*(17), 3789–3805.

doi:10.1111/j.1365-294X.2010.04698.x

760   Evans, L. M., Slavov, G. T., Rodgers-Melnick, E., Martin, J., Ranjan, P., Muchero, W.,

        … DiFazio, S. P. (2014). Population genomics of Populus trichocarpa identifies

        signatures of selection and adaptive trait associations. *Nat Genet*, *46*(10), 1089–

        1096.

        Falconer, D. S. (1952). The Problem of Environment and Selection. *The American*

765   *Naturalist*, *86*(830), 293–298.

        Fitzpatrick, M. C., & Keller, S. R. (2015). Ecological genomics meets community-level

        modelling of biodiversity: mapping the genomic landscape of current and future

        environmental adaptation. *Ecology Letters*, *18*(1), 1–16. doi:10.1111/ele.12376

        Forester, B. R., Jones, M. R., Joost, S., Landguth, E. L., & Lasky, J. R. (2016). Detecting

770   spatial genetic signatures of local adaptation in heterogeneous landscapes.

        Molecular Ecology, *25*(1), 104–120. doi:10.1111/mec.13476

        Forester BR, Jones MR, Joost S, Landguth EL, & Lasky JR. (2015, November 17). Data

        from: Detecting spatial genetic signatures of local adaptation in heterogeneous

        landscapes. Dryad Data Repository. doi:10.5061/dryad.v0c77

775   Fournier-Level, A., Korte, A., Cooper, M. D., Nordborg, M., Schmitt, J., & Wilczek, A.

        M. (2011). A Map of Local Adaptation in Arabidopsis thaliana. *Science*,

        334(6052), 86–89.

        Fournier-Level, A., Korte, A., Cooper MD, Nordborg M, Schmitt J, & Wilczek AM.

        (2011, October 7). Data from: A map of local adaptation in Arabidopsis thaliana.

780   Dryad Data Repository. Retrieved from http://dx.doi.org/10.5061/dryad.37f9t

Franks, S. J., Sim, S., & Weis, A. E. (2007). Rapid evolution of flowering time by an

annual plant in response to a climate fluctuation. *Proceedings of the National*

Academy of Sciences, *104*(4), 1278–1282. doi:10.1073/pnas.0608379104

Fraser, H. B. (2013). Gene expression drives local adaptation in humans. *Genome*

785   *Research*, *23*(7), 1089–1096. doi:10.1101/gr.152710.112

Fumagalli, M., Sironi, M., Pozzoli, U., Ferrer-Admettla, A., Pattini, L., & Nielsen, R.

(2011). Signatures of Environmental Genetic Adaptation Pinpoint Pathogens as

the Main Selective Pressure through Human Evolution. *PLoS Genet*, *7*(11),

e1002355. doi:10.1371/journal.pgen.1002355

790   Gage, S. H., Davey Smith, G., Ware, J. J., Flint, J., & Munaf�, M. R. (2016). G = E:

What GWAS Can Tell Us about the Environment. *PLoS Genet*, *12*(2), e1005765.

doi:10.1371/journal.pgen.1005765

Gauderman, W. J., Zhang, P., Morrison, J. L., & Lewinger, J. P. (2013). Finding Novel

Genes by Testing G×E Interactions in a Genomewide Association Study. *Genetic*

795   *Epidemiology*, *37*(6), 603–613. doi:10.1002/gepi.21748

Gautier, M., Vitalis, & R. (2012). rehh: An R package to detect footprints of selection in

genome-wide SNP data from haplotype structure. *Bioinformatics*, *0*(0), accepted.

Gelman, A., & HILL, J. (2007). *Data Analysis Using Regression and*

Multilevel/Hierarchical Models. Cambridge: Cambridge University Press.

800   Hall, M. C., LOWRY, D. B., & WILLIS, J. H. (2010). Is local adaptation in Mimulus

guttatus caused by trade-offs at individual loci? *Molecular Ecology*, *19*(13),

2739–2753. doi:10.1111/j.1365-294X.2010.04680.x

Hall, M. C., & Willis, J. H. (2006). DIVERGENT SELECTION ON FLOWERING TIME CONTRIBUTES TO LOCAL ADAPTATION IN MIMULUS

805   GUTTATUS POPULATIONS. *Evolution*, *60*(12), 2466–2477. doi:10.1111/j.0014-3820.2006.tb01882.x

Hancock, A. M., Brachi, B., Faure, N., Horton, M. W., Jarymowycz, L. B., Sperone, F. G., … Bergelson, J. (2011). Adaptation to Climate Across the Arabidopsis thaliana Genome. *Science*, *334*(6052), 83–86.

810 Hancock, A. M., Witonsky, D. B., Gordon, A. S., Eshel, G., Pritchard, J. K., Coop, G., & Di Rienzo, A. (2008). Adaptations to Climate in Candidate Genes for Common Metabolic Disorders. *PLoS Genet*, *4*(2), e32. doi:10.1371/journal.pgen.0040032

Hedrick, P. W., Ginevan, M. E., & Ewing, E. P. (1976). Genetic Polymorphism in Heterogeneous Environments. *Annual Review of Ecology and Systematics*, *7*, 1–

815   32.

Hereford, J. (2009). A Quantitative Survey of Local Adaptation and Fitness Trade-Offs. *The American Naturalist*, *173*(5), 579–588.

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high resolution interpolated climate surfaces for global land areas. *International*

820   *Journal of Climatology*, *25*(15), 1965–1978.

Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., … Whitlock, M. C. (2016). Finding the Genomic Basis of Local Adaptation: Pitfalls, Practical Solutions, and Future Directions. *The American Naturalist*, *188*(4), 379– 397. doi:10.1086/688018

825   Hoffmann, M. H. (2002). Biogeography of Arabidopsis thaliana (L.) Heynh.

        (Brassicaceae). *Journal of Biogeography*, *29*(1), 125–134.

        Horton, M. W., Hancock, A. M., Huang, Y. S., Toomajian, C., Atwell, S., Auton, A., …

        Bergelson, J. (2012). Genome-wide patterns of genetic variation in worldwide

        Arabidopsis thaliana accessions from the RegMap panel. *Nat Genet*, *44*, 212–216.

830   doi:10.1038/ng.1042

        Horton, M. W., Willems, G., Sasaki, E., Koornneef, M., & Nordborg, M. (2016). The

        genetic architecture of freezing tolerance varies across the range of Arabidopsis

        thaliana. *Plant, Cell & Environment*, *39*(11), 2570–2579. doi:10.1111/pce.12812

        Hunter, D. J. (2005). Gene-environment interactions in human diseases. *Nat Rev Genet*,

835   *6*(4), 287–298. doi:10.1038/nrg1578

        Jarquín, D., Crossa, J., Lacaze, X., Du Cheyron, P., Daucourt, J., Lorgeou, J., … de los

        Campos, G. (2014). A reaction norm model for genomic selection using high-

        dimensional genomic and environmental data. *Theoretical and Applied Genetics*,

        127(3), 595–607. doi:10.1007/s00122-013-2243-1

840   Jones, F. C., Grabherr, M. G., Chan, Y. F., Russell, P., Mauceli, E., Johnson, J., …

        Kingsley, D. M. (2012). The genomic basis of adaptive evolution in threespine

        sticklebacks. *Nature*, *484*(7392), 55–61. doi:10.1038/nature10944

        Kang, H. M., Sul, J. H., Service, S. K., Zaitlen, N. A., Kong, S., Freimer, N. B., … Eskin,

        E. (2010). Variance component model to account for sample structure in genome-

845   wide association studies. *Nat Genet*, *42*(4), 348–354. doi:10.1038/ng.548

        Kang, H. M., Zaitlen, N. A., Wade, C. M., Kirby, A., Heckerman, D., Daly, M. J., &

        Eskin, E. (2008). Efficient Control of Population Structure in Model Organism

Association Mapping. *Genetics*, *178*(3), 1709–1723.

doi:10.1534/genetics.107.080101

850   Kawecki, T. J., & Ebert, D. (2004). Conceptual issues in local adaptation. *Ecology*

Letters, *7*(12), 1225–1241. doi:10.1111/j.1461-0248.2004.00684.x

Keller, M. C. (2014). Gene × Environment Interaction Studies Have Not Properly

Controlled for Potential Confounders: The Problem and the (Simple) Solution.

Temperament: Genetic and Environmental Factors, *75*(1), 18–24.

855   doi:10.1016/j.biopsych.2013.09.006

Keller, S. R., Levsen, N., Olson, M. S., & Tiffin, P. (2012). Local Adaptation in the

Flowering-Time Gene Network of Balsam Poplar, Populus balsamifera L.

Molecular Biology and Evolution, *29*(10), 3143–3152.

doi:10.1093/molbev/mss121

860   Keller, S. R., Soolanayakanahally, R. Y., Guy, R. D., Silim, S. N., Olson, M. S., & Tiffin,

P. (2011). Climate-driven local adaptation of ecophysiology and phenology in

balsam poplar, Populus balsamifera L. (Salicaceae). *American Journal of Botany*,

98(1), 99–108. doi:10.3732/ajb.1000317

Kim, K.-C., Lai, Z., Fan, B., & Chen, Z. (2008). Arabidopsis WRKY38 and WRKY62

865   Transcription Factors Interact with Histone Deacetylase 19 in Basal Defense. *The*

Plant Cell Online, *20*(9), 2357–2371.

Kim, S., Plagnol, V., Hu, T. T., Toomajian, C., Clark, R. M., Ossowski, S., … Nordborg,

M. (2007). Recombination and linkage disequilibrium in Arabidopsis thaliana.

Nat Genet, *39*(9), 1151–1155. doi:10.1038/ng2115

870   Kooperberg, C., Dai, J. Y., & Hsu, L. (2016). Two-stage procedures for the identification

of gene x environment and gene x gene interactions in genome-wide association

studies. In *Statistical approaches to gene x environment interactions for complex*

phenotypes (pp. 11–34). Cambridge, Massachusetts: MIT Press.

Korte, A., Vilhjalmsson, B. J., Segura, V., Platt, A., Long, Q., & Nordborg, M. (2012). A

875   mixed-model approach for genome-wide association studies of correlated traits in

structured populations. *Nat Genet*, *44*(9), 1066–1071. doi:10.1038/ng.2376

Landguth, E. L., & Cushman, S. A. (2010). cdpop: A spatially explicit cost distance

population genetics program. *Molecular Ecology Resources*, *10*(1), 156–161.

doi:10.1111/j.1755-0998.2009.02719.x

880   Langlet, O. (1971). Two hundred years genecology. *Taxon*, 653–721.

Lasky, J. R., Des Marais, D. L., Lowry, D. B., Povolotskaya, I., McKay, J. K., Richards,

J. H., … Juenger, T. E. (2014). Natural variation in abiotic stress responsive gene

expression and local adaptation to climate in Arabidopsis thaliana. *Molecular*

Biology and Evolution, *31*(9), 2283–2296. doi:10.1093/molbev/msu170

885   Lasky, J. R., Des Marais, D. L., McKay, J. K., Richards, J. H., Juenger, T. E., & Keitt, T.

H. (2012). Characterizing genomic variation of Arabidopsis thaliana: the roles of

geography and climate. *Molecular Ecology*, *21*(22), 5512–5529.

doi:10.1111/j.1365-294X.2012.05709.x

Lasky, J. R., Des Marais DL, McKay JK, Richards JH, Juenger TE, & Keitt TH. (2012,

890   October 29). Data from: Characterizing genomic variation of Arabidopsis

thaliana: the roles of geography and climate. Dryad Data Repository. Retrieved

from http://dx.doi.org/10.5061/dryad.2gp18

Lasky, J. R., Upadhyaya, H. D., Ramu, P., Deshpande, S., Hash, C. T., Bonnette, J., …

Morris, G. P. (2015). Genome-environment associations in sorghum landraces

predict adaptive traits. *Science Advances*, *1*(6), e1400218.

doi:10.1126/sciadv.1400218

Leimu, R., & Fischer, M. (2008). A Meta-Analysis of Local Adaptation in Plants. *PLoS ONE*, *3*(12), e4010. doi:10.1371/journal.pone.0004010

Levene, H. (1953). Genetic Equilibrium When More Than One Ecological Niche is

Available. *The American Naturalist*, *87*(836), 331–333. doi:10.1086/281792

Levy Karin, E., Wicke, S., Pupko, T., & Mayrose, I. (2017). An Integrated Model of

Phenotypic Trait Changes and Site-Specific Sequence Evolution. *Systematic Biology*. doi:10.1093/sysbio/syx032

Li, Y., Cheng, R., Spokas, K. A., Palmer, A. A., & Borevitz, J. O. (2014). Genetic

Variation for Life History Sensitivity to Seasonal Warming in Arabidopsis

thaliana. *Genetics*, *196*(2), 569–577. doi:10.1534/genetics.113.157628

Lotterhos, K. E., & Whitlock, M. C. (2014). Evaluation of demographic history and

neutral parameterization on the performance of FST outlier tests. *Molecular Ecology*, *23*(9), 2178–2192. doi:10.1111/mec.12725

Lovell, J. T., Juenger, T. E., Michaels, S. D., Lasky, J. R., Platt, A., Richards, J. H., …

McKay, J. K. (2013). Pleiotropy of FRIGIDA enhances the potential for

multivariate adaptation. *Proceedings of the Royal Society B: Biological Sciences*,

280(1763), 20131043. doi:10.1098/rspb.2013.1043

Lowry, D. B., & Willis, J. H. (2010). A Widespread Chromosomal Inversion

Polymorphism Contributes to a Major Life-History Transition, Local Adaptation,

and Reproductive Isolation. *PLoS Biol*, *8*(9), e1000500.

doi:10.1371/journal.pbio.1000500

Lowry, D., Behrman, K., Grabowski, P., Morris, G., Kiniry, J., & Juenger, T. (2014).

Adaptations between ecotypes and along environmental gradients in Panicum

920   virgatum. *The American Naturalist*.

Manel, S., & Holderegger, R. (2013). Ten years of landscape genetics. *Trends in Ecology*

& Evolution, *28*(10), 614–621. doi:10.1016/j.tree.2013.05.012

Marigorta, U. M., & Gibson, G. (2014). A simulation study of gene-by-environment

interactions in GWAS implies ample hidden effects. *Frontiers in Genetics*,

925   *5*(225). doi:10.3389/fgene.2014.00225

Meijón, M., Feito, I., Oravec, M., Delatorre, C., Weckwerth, W., Majada, J., & Valledor,

L. (2016). Exploring natural variation of Pinus pinaster Aiton using

metabolomics: Is it possible to identify the region of origin of a pine from its

metabolites? *Molecular Ecology*, *25*(4), 959–976. doi:10.1111/mec.13525

930   Mitchell-Olds, T., Willis, J. H., & Goldstein, D. B. (2007). Which evolutionary processes

influence natural genetic variation for phenotypic traits? *Nat Rev Genet*, *8*(11),

845–856. doi:10.1038/nrg2207

Murayama, M., Hayashi, S., Nishimura, N., Ishide, M., Kobayashi, K., Yagi, Y., …

Hirayama, T. (2012). Isolation of Arabidopsis ahg11, a weak ABA hypersensitive

935   mutant defective in nad4 RNA editing. *Journal of Experimental Botany*, *63*(14),

5301–5310. doi:10.1093/jxb/ers188

Murcray, C. E., Lewinger, J. P., & Gauderman, W. J. (2009). Gene-Environment

    Interaction in Genome-Wide Association Studies. *American Journal of*

    Epidemiology, *169*(2), 219–226. doi:10.1093/aje/kwn353

940   Nordborg, M., & Weigel, D. (2008). Next-generation genetics in plants. *Nature*,

    456(7223), 720–723. doi:10.1038/nature07629

Price, A. L., Zaitlen, N. A., Reich, D., & Patterson, N. (2010). New approaches to

    population stratification in genome-wide association studies. *Nat Rev Genet*,

    11(7), 459–463.

945   Pyhäjärvi, T., Hufford, M. B., Mezmouk, S., & Ross-Ibarra, J. (2013). Complex Patterns

    of Local Adaptation in Teosinte. *Genome Biology and Evolution*, *5*(9), 1594–

    1609. doi:10.1093/gbe/evt109

Rao, C. R. (1971). Minimum variance quadratic unbiased estimation of variance

    components. *Journal of Multivariate Analysis*, *1*(4), 445–456. doi:10.1016/0047-

950   259X(71)90019-4

Reimherr, M., & Nicolae, D. (2015). Estimating Variance Components in Functional

    Linear Models with Applications to Genetic Heritability. *Journal of the American*

    Statistical Association, 00–00. doi:10.1080/01621459.2015.1016224

Rellstab, C., Gugerli, F., Eckert, A. J., Hancock, A. M., & Holderegger, R. (2015). A

955   practical guide to environmental association analysis in landscape genomics.

    Molecular Ecology, *24*(17), 4348–4370. doi:10.1111/mec.13322

Schaffer, H. E., & Johnson, F. M. (1974). ISOZYME ALLELIC FREQUENCIES

    RELATED TO SELECTION AND GENE-FLOW HYPOTHESES. *Genetics*,

    77(1), 163–168.

960   Schoville, S. D., Bonin, A., François, O., Lobreaux, S., Melodelima, C., & Manel, S.

           (2012). Adaptive Genetic Variation on the Landscape: Methods and Cases.

           Annual Review of Ecology, Evolution, and Systematics, *43*(1), 23–43.

           doi:10.1146/annurev-ecolsys-110411-160248

       Schrider, D. R., Mendes, F. K., Hahn, M. W., & Kern, A. D. (2015). Soft Shoulders

965   Ahead: Spurious Signatures of Soft and Partial Selective Sweeps Result from

           Linked Hard Sweeps. *Genetics*, *200*(1), 267–284.

           doi:10.1534/genetics.115.174912

       Sharbel, T. F., Haubold, B., & Mitchell-Olds, T. (2000). Genetic isolation by distance in

           Arabidopsis thaliana: biogeography and postglacial colonization of Europe.

970   *Molecular Ecology*, *9*(12), 2109–2118.

       Shindo, C., Aranzana, M. J., Lister, C., Baxter, C., Nicholls, C., Nordborg, M., & Dean,

           C. (2005). Role of FRIGIDA and FLOWERING LOCUS C in Determining

           Variation in Flowering Time of Arabidopsis. *Plant Physiol.*, *138*(2), 1163–1173.

       Shoji, T. (2014). ATP-binding cassette and multidrug and toxic compound extrusion

975   transporters in plants: A common theme among diverse detoxification

           mechanisms. *Int Rev Cell Mol Biol*, *309*, 303–346.

       Slatkin, M. (1973). GENE FLOW AND SELECTION IN A CLINE. *Genetics*, *75*(4),

           733–756.

       Stinchcombe, J. R., Weinig, C., Ungerer, M., Olsen, K. M., Mays, C., Halldorsdottir, S.

980   S., … Schmitt, J. (2004). A latitudinal cline in flowering time in Arabidopsis

           thaliana modulated by the flowering time gene FRIGIDA. *Proceedings of the*

National Academy of Sciences of the United States of America, *101*(13), 4712–4717. doi:10.1073/pnas.0306401101

Szechyńska-Hebda, M., Czarnocka, W., Hebda, M., & Karpiński, S. (2016). PAD4, LSD1 and EDS1 regulate drought tolerance, plant biomass production, and cell wall properties. *Plant Cell Reports*, 1–13. doi:10.1007/s00299-015-1901-y

Thomas, D. (2010). Methods for Investigating Gene-Environment Interactions in Candidate Pathway and Genome-Wide Association Studies. *Annual Review of Public Health*, *31*, 21–36. doi:10.1146/annurev.publhealth.012809.103619

Tiezzi, F., de los Campos, G., Parker Gaddis, K. L., & Maltecca, C. (2017). Genotype by environment (climate) interaction improves genomic prediction for production traits in US Holstein cattle. *Journal of Dairy Science*, *100*(3), 2042–2056. doi:10.3168/jds.2016-11543

Tiffin, P., & Ross-Ibarra, J. (2014). Advances and limits of using population genetics to understand local adaptation. *Trends in Ecology & Evolution*, *29*(12), 673–680. doi:10.1016/j.tree.2014.10.004

Toomajian, C., Hu, T. T., Aranzana, M. J., Lister, C., Tang, C., Zheng, H., … Nordborg, M. (2006). A Nonparametric Test Reveals Selection for Rapid Flowering in the Arabidopsis Genome. *PLoS Biol*, *4*(5), e137. doi:10.1371/journal.pbio.0040137

Turner, T. L., Bourne, E. C., Von Wettberg, E. J., Hu, T. T., & Nuzhdin, S. V. (2010). Population resequencing reveals local adaptation of Arabidopsis lyrata to serpentine soils. *Nat Genet*, *42*(3), 260–263. doi:10.1038/ng.515

van Oppen, M. J. H., Oliver, J. K., Putnam, H. M., & Gates, R. D. (2015). Building coral reef resilience through assisted evolution. *Proceedings of the National Academy of Sciences*, *112*(8), 2307–2313. doi:10.1073/pnas.1422301112

Verslues, P. E., Lasky, J. R., Juenger, T. E., Liu, T.-W., & Kumar, M. N. (2014). Genome-Wide Association Mapping Combined with Reverse Genetics Identifies New Effectors of Low Water Potential-Induced Proline Accumulation in Arabidopsis. *Plant Physiology*, *164*(1), 144–159. doi:10.1104/pp.113.224014

Voight, B. F., Kudaravalli, S., Wen, X., & Pritchard, J. K. (2006). A Map of Recent Positive Selection in the Human Genome. *PLoS Biology*, *4*(3), e72. doi:10.1371/journal.pbio.0040072

Weigel, D., & Nordborg, M. (2015). Population Genomics for Understanding Adaptation in Wild Plant Species. *Annual Review of Genetics*, *49*(1), 315–338. doi:10.1146/annurev-genet-120213-092110

Wilczek, A. M., Cooper, M. D., Korves, T. M., & Schmitt, J. (2014). Lagging adaptation to warming climate in Arabidopsis thaliana. *Proceedings of the National Academy of Sciences*, *111*(22), 7906–7913. doi:10.1073/pnas.1406314111

Windle, M. (2016). *Statistical Approaches to Gene X Environment Interactions for Complex Phenotypes*. MIT Press.

Wituszyńska, W., Ślesak, I., Vanderauwera, S., Szechyńska-Hebda, M., Kornaś, A., Van Der Kelen, K., … Karpiński, S. (2013). LESION SIMULATING DISEASE1, ENHANCED DISEASE SUSCEPTIBILITY1, and PHYTOALEXIN DEFICIENT4 Conditionally Regulate Cellular Signaling Homeostasis,

1025   Photosynthesis, Water Use Efficiency, and Seed Yield in Arabidopsis. *Plant Physiology*, *161*(4), 1795–1805. doi:10.1104/pp.112.208116

Yates, F., & Cochran, W. G. (1938). The analysis of groups of experiments. *The Journal of Agricultural Science*, *28*(04), 556–580.

Yeaman, S., & Whitlock, M. C. (2011). THE GENETIC ARCHITECTURE OF

1030   ADAPTATION UNDER MIGRATION–SELECTION BALANCE. *Evolution*, *65*(7), 1897–1911. doi:10.1111/j.1558-5646.2011.01269.x

Yoder, J. B., Stanton-Geddes, J., Zhou, P., Briskine, R., Young, N. D., & Tiffin, P. (2014). Genomic Signature of Adaptation to Climate in Medicago truncatula. *Genetics*. doi:10.1534/genetics.113.159319

1035   Zomer, R. J., Trabucco, A., Bossio, D. A., & Verchot, L. V. (2008). Climate change mitigation: A spatial analysis of global land suitability for clean development mechanism afforestation and reforestation. *Agriculture, Ecosystems & Environment*, *126*(1–2), 67–80. doi:10.1016/j.agee.2008.01.014

Data Accessibility

1040   All data were previously published, including fitness (Data Dryad package Fournier-Level *et al.* 2011b), environment (Data Dryad package Lasky *et al.* 2012a), and SNPs (Horton *et al.* 2012, available at http://bergelson.uchicago.edu/wp-content/uploads/2015/04/call_method_75.tar.gz). Simulation data previously published in Forester *et al.* (2016) are archived in Data Dryad package (Forester *et al.* 2015). New

1045   simulation data, and code for simulations and GWAS are included in an online supplemental file with this manuscript.

Author contributions

JRL designed research and led paper writing. All authors conducted analyses and

1050   contributed to writing.

Table 1. Characterization of patterns in *Arabidopsis* case study identified by Approach 4 (mixed model including imputations) for SNPs in 0.01 lower tail of p-values for SNP×environment interactions for fitness (first two rows of table) and for kinship×environment interactions for fitness (third row). SNPs with home allele advantage are defined as those where the sign of allelic differences in home climates were matched by the sign of fitted mixed model SNP×environment associations with relative fitness. Rank changing SNPs are those where we estimated a rank change in relative fitness for alternate alleles along the environmental gradient between the two extreme common gardens. The final row gives the proportion of total observed climate gradient (among ecotypes) captured by the two most extreme common gardens.

Statistic	*Aridity*	*CV grow. seas. prec.*	*Min. temp.*	*Min. temp. grow. seas.*
Proportion SNP×E with home allele advantage	0.46	0.89	0.92	>0.99
Proportion SNP×E rank changing	0.31	0.92	>0.99	0.73
Kinship×E R^2 for fitness	0.08	0.58	0.07	0.43
Proportion climate gradient covered by gardens	0.13	0.65	0.78	0.31

1055

1060

1065

Table 2. Permutation tests of enrichment p-values (Approach 4) for various signals suggestive of local adaptation to climate in case study on Arabidopsis. For each statistic, we tested for enrichment of signal in the SNPs in the 0.01 lower tail of p-values for SNP×environment associations with relative fitness. "Genic" tests enrichment of genic versus non-genic SNPs, "PHS" and "iHS" test for enrichment with significant (alpha = 0.05) pairwise haplotype sharing and standardized integrated extended haplotype homozygosity, respectively. The final row shows enrichment with SNPs having significant (alpha = 0.05) associations with change in flowering time (G×E) in response to warming during growth.

Statistic	*Aridity enrichment*	*CV grow. seas. prec. enrichment*	*Min. temp. enrichment*	*Min. temp. grow. seas. enrichment*
Genic	0.0036	<0.0002	0.0054	0.0006
PHS	0.0272	0.0122	0.0068	0.0272
iHS	0.2840	0.0006	0.0020	0.0008
Flowering time under warming, G×E	0.8582	0.0680	0.5844	<0.0002

Figure 1. Illustration of our imputation technique and stereotypical patterns captured by our approach for neutral (top panels) and selected (bottom panels) loci. Here we show hypothetical data from four common gardens along an environmental gradient (solid circles in four vertical streaks in right panels, with small amount of noise added to environmental values for visualization) that have fitness scaled to a maximum of 1. We also show accessions (or ecotypes) collected in home environments and sequenced (mapped in two-dimensional geographical space in left panels) having imputed relative fitness of 1 in their home environment (environment-of-origin, open circles at top of right panels). The selected locus (bottom right panel) in question shows strong G×E for fitness, such that allele *B* (blue) is more fit (observed, solid circles) and more common (imputed, open circles) at the right side of the environmental gradient while allele *b* (orange) is more fit and more common at the left side of the gradient.

Neutral locus

- ● Observed fitness in common garden
- ○ Imputed fitness at environment–of–origin

Selected locus

1090

Figure 2. Data used in case study on Arabidopsis. The location of common gardens, natural accessions in common gardens, and all other sequenced natural accessions are shown.

1095

Figure 3. Comparison of causal (red) and neutral (black) SNP associations with G×E for fitness across three different levels of dispersal and 10 replicate simulations for each level. Approaches used (row 1) no imputation and no random effects, (row 2) imputation but no random effects, (row 3) mixed models that used only observations from four common gardens, (row 4) mixed models combining imputed observations of relative fitness in home environments with common garden observations, or (row 5) a genotype-environment association approach. False positive rate (FPR) is indicated, calculated as the proportion of simulations where a neutral SNP had the lowest p-value. Each simulation had 1 causal SNP and 99 neutral SNPs; plots show aggregate distributions for all SNP by simulation combinations (*i.e.* total of 10 causal and 990 neutral SNPs). Y-axes show the proportion of SNPs in each category (causal or neutral) falling into a given p-value bin. For reference, dashed line indicates a strict Bonferroni cutoff for alpha = 0.05, $-\log(0.05/100) = 7.6$.

Figure 4. Example SNPs with the strongest associations (lowest p-values) with cold winter temperatures (A) and aridity (B). Top subpanels show the climate distribution of alleles in home genotypes (natural geographic patterns), known as genotype-environment associations. Bottom subpanels show relative fitness of alleles in four common gardens, where common gardens' climates determine position on x-axes. Each SNP falls within the coding region of indicated genes (WRKY38 and LSD1). Box widths are scaled to relative number of accessions having each allele. In both (A) and (B), the allele with the greatest relative fitness in common gardens changes along the environmental gradient consistent with change in allele frequency in native accessions (*i.e.* ecotypes).

A. WRKY38 (Chr. 5, pos. 7496047)

B. LSD1 (Chr. 4, pos. 11005059)

Supporting information

Simulations

1125 Forester *et al.* (2016) previously simulated local adaptation in a square two-dimensional

1024 x 1024 grid-cell landscape along a continuous environmental gradient, using the

program CDPOP v1.2 (Landguth & Cushman 2010). Forester *et al.* (2016) simulated

5,000 diploid individuals with 100 bi-allelic loci, with one locus under selection and 99

loci evolving neutrally. All loci had a 0.0005 mutation rate per generation, free

1130 recombination, and no physical linkage. The authors ran 10 Monte Carlo replicates of the

simulation for 1,250 generations, using the first 250 generations as a burn-in, with no

selection imposed, to establish a spatial genetic pattern.

Selection varied linearly along an environmental gradient, such that *AA* and *aa*

genotypes were favored at different ends of the gradient (North and South, Fig. S1). The

1135 selection strength of *s*=0.10 at extreme ends of the gradient was mediated through

density-independent mortality determined by an individual's genotype at the selected

locus, where *AA* experienced 0% mortality at the North extreme and *s* mortality at the

South extreme, while *aa* genotypes experienced the opposite selection gradient. *Aa*

genotypes experienced uniform selection of *s*/2 across the gradient. The number of

1140 offspring produced from mating (fitness) was determined from a Poisson distribution (λ =

4), These same demographic parameters were used for common garden simulations

conducted for the present paper.

In Forester *et al.* (2016), this level of fecundity produced an excess of individuals

each generation. Carrying capacity of the landscape surface was 5,000 individuals, and

1145 excess new individuals were discarded once all 5,000 locations became occupied.

Mating pairs of hermaphroditic individuals and dispersal locations of offspring were chosen using a random draw from the inverse-square probability function of distance, truncated at a distance equal to *d,* the proportion of a landscape edge. We tested three values of *d*: 0.03, 0.1, and 0.25 (*i.e.* truncated at 31, 102, and 256 pixels, respectively). Individuals near landscape edges were unable to disperse or mate with individuals beyond the edge such that boundaries were not periodic.

1150

1155

Tables S1-S4 (attached csv files). List of Arabidopsis genes within 1 kb of SNPs in the lower 0.001 quantile for p-values for SNP×environment interactions for each climate variable, including imputed observations and accounting for kinship.

1160

Table S5. Comparison of scenarios under which five approaches studied in simulation identify true causal SNPs. Cells represent the correlation coefficient between each approach's success at identifying the causal SNP as having the lowest p-value, across 10 different simulations across 3 dispersal levels (i.e. correlation between success for 30 total simulations).

	A2 Imputation, no random effect	A3 No imputation, random effect	A4 Imputation, random effect	A5 Gene-env. asscn., random effect
A1 No imputation, no random effect	0.71	0.24	0.38	0.26
A2 Imputation, no random effect		0.39	0.53	0.37
A3 No imputation, random effect			0.80	0.56
A4 Imputation, random effect				0.69

Table S6. Rank correlation among SNPs for SNP×environment effect p-values

comparing three tested approaches (Approaches 2-4) with Arabidopsis.

	A3 Random effect, no imputation	*A4 Random effect, imputation*
Min. temp. growing season		
A2 No random effect, imputation	0.396	-0.001
A3 Random effect, no imputation		0.041
Aridity		
A2 No random effect, imputation	0.318	0.008
A3 Random effect, no imputation		-0.032
Min. temp. coldest month		
A2 No random effect, imputation	-0.021	-0.012
A3 Random effect, no imputation		0.104
CV growing season prec.		
A2 No random effect, imputation	0.178	0.090
A3 Random effect, no imputation		-0.006

1170

Table S7. Overlap (one-tailed permutation tests) and rank correlations (Spearman's rho) between our approach (Approach 4, mixed-model with imputation, climate variables give row labels at left) and published approaches to identifying SNPs associated with local adaptation in Arabidopsis. Citations are given for the original publication of previous approaches. For partial Mantel and RDA, we used absolute value of SNP scores to rank SNPs. Empty cells indicate climate variables not tested in Hancock *et al*. (2011).

1175

Climate var.	Mixed model, overlap in 0.01 tail strongest associations	Mixed model, rank correlation between associations	First axis of RDA, overlap in 0.01 tail strongest associations	First axis of RDA, rank correlation between associations	First axis of RDA after removing spatial structure, overlap in 0.01 tail strongest associations	First axis of RDA after removing spatial structure, rank correlation between associations	Partial Mantel overlap in 0.01 tail strongest associations
	Lasky et al. 2014	Lasky et al. 2014	Lasky et al. 2012	Lasky et al. 2012	Lasky et al. 2012	Lasky et al. 2012	Hancock et al 2011
Min. temp. coldest month	<0.0001	0.0416	<0.0001	-0.2126	<0.0001	-0.0575	0.015
Aridity	<0.0001	0.0573	<0.0001	-0.0153	0.1137	0.0127	<0.000
CV growing season prec.	<0.0001	0.1871	0.9623	-0.0282	0.9959	0.0303	
Min. temp. growing season	<0.0001	0.0794	0.0556	-0.0928	<0.0001	-0.1319	

Figure S1. Example outcome of selection on a simulated landscape under moderate

1180 dispersal (maximum dispersal of 10% of the landscape surface per generation), with

selective gradient (grayscale background), 250 randomly sampled accessions (small

circles), and four common gardens (green squares). 100 accessions were simulated in the

four common gardens (i.e. reciprocal transplant) covering the full gradient (large light

green squares) and in four gardens covering only half the gradient (smaller dark green

1185 squares).

Figure S2. Example outcome of selection on a simulated landscape under low dispersal (maximum dispersal of 3% of the landscape surface per generation), with selective gradient (grayscale background), 250 randomly sampled accessions (small circles), and four common gardens (green squares). 100 accessions were simulated in the four common gardens (i.e. reciprocal transplant) covering the full gradient (large light green squares).

1195

Figure S3. Example outcome of selection on a simulated landscape under high dispersal (maximum of 25% of the landscape surface per generation), with selective gradient (grayscale background), 250 randomly sampled accessions (small circles), and four common gardens (green squares). 100 accessions were simulated in the four common

1200 gardens (i.e. reciprocal transplant) covering the full gradient (large light green squares).

Figure S4. Comparison of inferred causal (red) and neutral (black) SNP associations with G×E for fitness across 10 replicate simulations for moderate dispersal with common gardens covering only half the gradient. Approaches used (row 1) no imputation and no random effects, (row 2) imputation but no random effects, (row 3) mixed models that used only observations from four common gardens, or (row 4) mixed models combining imputed observations of relative fitness in home environments with common garden observations. False positive rate (FPR) is indicated, calculated as the proportion of simulations where a neutral SNP had the lowest p-value. Each simulation had 1 causal SNP and 99 neutral SNPs; plots show aggregate distributions for all SNP by simulation combinations (*i.e.* total of 10 causal and 990 neutral SNPs). Y-axes show the proportion of SNPs in each category (causal or neutral) falling into a given p-value bin. For reference, dashed line indicates a strict Bonferroni cutoff for alpha = 0.05, $-\log(0.05/100)$ = 7.6.

1215

Figure S5. Quantile-quantile plots of p-value distributions for four approaches to calculating genome-wide SNP×environment associations with fitness (columns 1-4) and one genome-environment approach (column 5, results from Lasky *et al.* 2014), using published data on *Arabidopsis thaliana*. X-axes show expected $-\log_{10}(p)$ and y-axes show observed $-\log_{10}(p)$. Approaches used (column 1) no imputation and no random effects, (column 2) imputation but no random effects, (column 3) mixed models that used only observations from four common gardens, or (column 4) mixed models combining imputed observations of relative fitness in home environments with common garden observations. In the fifth column, environment is tested as a function of SNP alleles in a typical mixed model GWAS approach where environment is substituted for phenotype (Lasky *et al.* 2014).

Figure S6. Comparison of published genome-environment association approaches to

1230   identifying loci underlying local adaptation in *Arabidopsis thaliana* (Hancock *et al.* 2011,

Lasky *et al.* 2012, Lasky *et al.* 2014). Each *x*-axis shows results from the imputation

approach developed in this paper, using a mixed model accounting for kinship.

1235   **Figure S7.** Example of fitted effects on fitness of allelic variation at two SNPs (arbitrary

labeled 1 and 2) across two common gardens (arbitrary labeled x and y). In this scenario,

both SNPs exhibit antagonistic pleiotropy (rank changing fitness tradeoffs) across the

environmental gradient. However, each SNP has relatively weaker effects on fitness in

one environment. A test of whether an individual SNP shows the strongest associations

1240   with fitness in an individual garden (*e.g.* SNP 1 at site x) might incorrectly infer

conditional neutrality because other markers show stronger associations (SNP 2 at site x).