<u>Title:</u>

QTL mapping reveals complex genetic architecture of quantitative virulence in the

wheat pathogen *Zymoseptoria tritici*

<u>Authors:</u>

Ethan L. Stewart, Daniel Croll, Mark H. Lendenmann, Andrea Sanchez-Vallet, Fanny E.

Hartmann, Javier Palma-Guerrero, Xin Ma, Bruce A. McDonald.

<u>Authors Address:</u>

Plant Pathology Group, ETH Zürich, Universitätstrasse 2, 8092, Zürich, Switzerland.

<u>Corresponding Author:</u>

Ethan Stewart

ethan.stewart@usys.ethz.ch

+41 44 632 38 64

<u>Key Words</u>

QTL mapping, quantitative virulence, *Zymoseptoria tritici*, *Mycosphaerella graminicola*,

image analysis.

Word count: 7180

Summary

We conducted a comprehensive analysis of virulence in the fungal wheat pathogen

Zymoseptoria tritici using QTL mapping. High throughput phenotyping based on

automated image analysis allowed measurement of pathogen virulence on a scale and

with a precision that was not previously possible. Across two mapping populations

encompassing more than 520 progeny, 540,710 pycnidia were counted and their sizes

and grey values were measured, yielding over 1.6 million phenotypes associated with

pathogen reproduction. Large pycnidia were shown to produce more numerous and

larger spores than small pycnidia. Precise measures of percent leaf area covered by

lesions provided a quantitative measure of host damage. Combining these large and

accurate phenotype datasets with a dense panel of RADseq genetic markers enabled us

to genetically dissect pathogen virulence into components related to host damage and

components related to pathogen reproduction. We show that different components of

virulence can be under separate genetic control. Large- and small-effect QTLs were

identified for all traits, with some QTLs specific to mapping populations, cultivars and

traits and other QTLs shared among traits within the same mapping population. We

associated the presence or absence of accessory chromosomes with several virulence

traits, providing the first evidence for an important function associated with accessory

chromosomes in this organism. A large-effect QTL involved in host specialization was

identified on chromosome 7, leading to identification of candidate genes having a large

effect on virulence.

Introduction

Much plant pathology research has been oriented around understanding the classic

gene-for-gene (GFG) interaction (Flor, 1955) between the pathogen and its host, largely

because early plant breeding programs focused on the introgression of major,

qualitative resistance genes (R-genes) to control plant pathogens. Because most

pathogens evolved quickly to defeat major R-genes, and quantitative resistance

appeared more durable, the genetic basis of quantitative resistance became an area of

intensive investigation, leading eventually to identification of host genes encoding

quantitative resistance (Krattinger et al, 2009, St. Clair, 2010). The term aggressiveness

is typically used by plant pathologists to describe the quantitative degree of damage

caused by a pathogen strain. In the broader literature, virulence is defined as the

quantitative degree of damage caused by a pathogen to its host. We will use the latter

definition to be consistent with the wider field of life sciences. In contrast to

quantitative resistance in the host, quantitative virulence in pathogens is much less

studied and remains poorly understood (Lannou, 2012, Pariaud et al, 2009), though

quantitative virulence appears to be common in plant pathogens (Pariaud et al, 2009).

Given the broad spectrum in observed virulence phenotypes, the genetic architecture of

virulence may be mediated both by single genes of large effect that follow the GFG

model (Flor, 1955) and by many genes that make smaller contributions to quantitative

virulence. Quantitative traits often exhibit a continuous distribution in a population and

are typically governed by independent assortment of multiple alleles with small

individual effects, with different combinations of alleles responsible for the range of

observed phenotypes in recombining populations (Mackay 2001). Numerous

confounding factors such as dominance, pleiotropy and additive allele effects also

contribute to the genetic architecture of a trait (Hansen 2006). In order to disentangle

individual loci contributing to overall virulence phenotypes, we utilized quantitative

trait locus (QTL) mapping, an established technique for determining the genetic

architecture of quantitative traits. QTL mapping was successfully used to identify

candidate genes underlying traits of interest in plants (Alonso-Blanco and Méndez-Vigo,

2014), animals (Solberg Woods, 2014) and humans (Almasy and Blangero, 2009). It has

also been employed in fungi (Larraya et al, 2003), though its use in fungi remains rare

compared to other eukaryotes (Foulongne-Oriol, 2012).

Virulence in the wheat pathogen *Zymoseptoria tritici* (formerly *Mycosphaerella*

graminicola), causal agent of Septoria tritici blotch (STB), is a predominantly

quantitative trait (Stewart and McDonald 2014, Zhan et al, 2007). *Z. tritici* is the most

damaging wheat pathogen in Europe (Jorgensen et al, 2014; O'Driscoll et al, 2014) and

is considered an important fungal pathogen worldwide (Dean et al, 2012). Under

optimal conditions, yield losses can reach 50% (Eyal 1987). Even with the use of

resistant cultivars and regular fungicide treatments, yield losses of 5–10% can be

expected (Fones and Gurr, 2015). Ten to fourteen days after infection chlorotic areas

begin to appear that later become necrotic lesions. Pycnidia containing asexual

pycnidiospores develop mostly within these necrotic lesions in the sub-stomatal

cavities. Pycnidiospores are exuded from the pycnidia in a gelatinous cirrhus during

periods of high humidity and are spread throughout the plant canopy by rain splash.

Numerous asexual infection cycles occurring during a growing season are the main

cause of STB epidemics in the field.

Despite its agricultural importance, the mechanisms responsible for virulence in *Z.*

tritici and their underlying genetics remain poorly understood. Twenty genes affecting

virulence in *Z. tritici* have been described and functionally characterized (Figure S1).

Most of these genes have developmental regulatory functions or affect inherent fitness

without having a specific role in virulence. Host resistance is also quantitative, with 21

genes and 89 genomic regions implicated in resistance to *Z. tritici* (Brown et al, 2015).

The quantitative nature of virulence in *Z. tritici* makes QTL mapping an ideal technique

to elucidate the genetic determinants of quantitative virulence, as already validated

using several other quantitative traits in *Z. tritici* (Lendenmann et al, 2014, Lendenmann

et al, 2015, Lendenmann et al, 2016).

Epidemic development is strongly affected by the ability of a pathogen to reproduce and

cause subsequent infections (Parlevliet, 1979), hence the reproductive output of a

pathogen is an important determinant of the total damage it can cause during an

epidemic. Damage caused by plant pathogens is frequently assessed by quantifying the

area covered by disease lesions on the host. However, a pathogen's ability to induce

lesions does not necessarily reflect its ability to reproduce. Indeed, several studies have

shown that pathogen damage, as indicated by the plant area covered by disease lesions,

can be independent of pathogen reproduction, as indicated by the number of spores

produced during an infection (Burns et al, 2014, Habgood, 1977, Halama et al, 1999,

Pariaud et al, 2009, Twizeyimana et al, 2014), suggesting that these traits may be under

separate genetic control. The contribution of each virulence component combines to

produce quantitative disease phenotypes.

The success of a QTL mapping project depends on the availability of genetic markers

and accurate phenotypic data. With current availability of genomic tools that can

generate very large numbers of genetic markers comparatively easily and cheaply, the

limiting factor in most studies is accurate phenotyping (Furbank and Tester, 2011).

Using image analysis to phenotype *Z. tritici* infection allows for measurement of both

leaf damage, based on the percentage of the leaf covered by necrotic lesions, and

pathogen reproduction, based on the number and size of the pycnidia formed on the

infected leaves (Stewart and McDonald 2014, Stewart et al, 2016).

The aim of this study was to investigate the genetic architecture of virulence traits in *Z.*

tritici. We conducted a QTL mapping study using two mapping populations derived

from parental isolates exhibiting varying degrees of virulence. Combining digital image

analysis with a large number of RADseq SNP markers in a large number of offspring

allowed us to genetically separate traits related to host damage and pathogen

reproduction. We also identified novel candidate genes involved in host specialization

and virulence.

<u>Results</u>

<u>Phenotyping</u>

A phenotyping method based on digital image analysis (Stewart and McDonald, 2014)

was used to measure % leaf area covered by lesions (PLACL), pycnidia density (# of

pycnidia per cm^2 leaf), pycnidia size and pycnidia melanization in two *Z. tritici* mapping

populations (3D1x3D7 and 1A5x1E4) that have also been used to study melanization

(Lendenmann et al, 2014), fungicide sensitivity (Lendenmann et al, 2015) and

temperature sensitivity (Lendenmann et al, 2016). The pairs of parental isolates used to

make the crosses were chosen based on differing levels of virulence observed in

previous work (Zhan et al, 2005). To investigate the effect of host on virulence, we used

cultivars Runal (moderately susceptible to *Z. tritici*) and Titlis (moderately resistant)

(Courvoisier et al, 2015). All phenotypes showed a continuous distribution in both

crosses. Transgressive segregation was evident for all phenotypes, with some progeny

exhibiting more extreme phenotypes than the parent isolates (Figure 1). In cross

3D1x3D7 all phenotypes were significantly higher on Runal than Titlis, including mean

pycnidia size (F=3.93, p=0.048), pycnidia melanization (F=22.7, p<0.001), PLACL

(F=21.4, p<0.001), and pycnidia density (F=19.2, p<0.001). In cross 1A5x1E4, PLACL

(F=22.7, p<0.001) and pycnidia melanization (F=17.7, p<0.001) were significantly

higher in Runal whereas pycnidia size (F=11.5, p<0.001) and pycnidia density (F=73.6,

p<0.001) were significantly higher in Titlis. Across both crosses, a total of 540,710

pycnidia were counted and their sizes and melanization measured, yielding over 1.6

million phenotypic data points associated with fungal fruiting bodies. In order to

evaluate whether counting and measuring pycnidia provides an accurate estimate of

spore number, the spore output of a subset of isolates was measured. The mean number

of spores per pycnidium was 2598. The mean spore size, as measured by length, was

17.6 μm. There were weak but significant positive correlations between pycnidia size

and spore size (r^2=0.111, p=0.001) and between pycnidia size and the number of spores

per pycnidium (r^2= 0.112, p=0.001). These findings indicate that pycnidia size is

correlated with reproductive output of *Z. tritici*.

Quantitative Trait Locus (QTL) mapping

We used QTL mapping to elucidate the genetic architecture of the quantitative virulence

phenotypes. QTLs were identified for all phenotypes. All QTLs in cross 3D1x3D7 were

found only on Runal and mapped to a single large effect QTL on chromosome 7 (Figure

172 2a). In 1A5x1E4 the same QTL for pycnidia density was found on chromosome 5 on

173 both cultivars. A second QTL for pycnidia density was found on chromosome 9 in Titlis

174 and on chromosome 3 in Runal. PLACL in Titlis and pycnidia grey value in Runal both

175 mapped to the same QTL on chromosome 5. Pycnidia size also mapped to the same QTL

176 on chromosome 3 in cv Runal (Figure 2b). No QTLs were shared between the two

177 crosses. The 95% confidence interval for a QTL ranged from 91.3 kb for the

178 chromosome 5 QTL in 1A5x1E4 to 717.7 kb for the chromosome 7 QTL for grey value

179 on Runal in 3D1x3D7.

181 The single, large effect QTL on chromosome 7 in cross 3D1x3D7 explained 54% of the

182 phenotypic variance for PLACL, 57% of the variance for pycnidia density and 18% of the

183 variance for pycnidia grey value. PLACL and pycnidia density shared the same

184 confidence interval containing 35 genes. The wider confidence interval for pycnidia

185 grey value contained 227 genes (Table 1A). The allele from the 3D7 parent was

186 responsible for higher PLACL and pycnidia density and decreased pycnidia

187 melanization.

189 In cross 1A5x1E4 the variance explained by each QTL was smaller and ranged from

190 6.1% for pycnidia density on Titlis to 8.7% for pycnidia size on Runal. The number of

191 genes within each confidence interval was generally higher than in cross 3D1x3D7,

192 ranging from 89 for grey value in Runal to 280 in the QTL on chromosome 5 for

193 pycnidia density in Runal and PLACL in Titlis (Table 1B). The 1A5 allele was responsible

194 for the higher phenotype values in the QTL on chromosome 5 whereas the 1E4 allele

195 was responsible for the higher phenotype values in the chromosome 3 and 9 QTLs.

Additive effects were observed for the two QTLs found for pycnidia density in 1A5x1E4.

On Runal, isolates with the 1A5 allele at the chromosome 5 QTL peak and the 1E4 allele

at the chromosome 3 QTL peak had significantly higher pycnidia density than isolates

with the reverse alleles (F=11.2, p<0.001) (Table 2A). On Titlis, isolates with the 1E4

allele at the chromosome 9 QTL peak and the 1A5 allele at the chromosome 5 QTL peak

had significantly higher phenotypes than isolates with the opposite allele combinations

(F=19.1, p<0.001) (Table 2B). On both cultivars, isolates with the same parental allele

at both QTL peaks were not different from each other and were intermediate to the

isolates with the alleles coming from different parents.

<u>Effect of Accessory Chromosomes on Virulence</u>

Among the parent isolates used to make the crosses, all accessory chromosomes were

present in 3D1 whereas 3D7 was missing chromosomes 14, 15, 18 and 21. In the

1A5x1E4 cross chromosome 17 was absent in 1E4. QTLs could only be mapped on

chromosomes present in both parent isolates. No QTLs were found on the accessory

chromosomes in either cross.

The presence or absence of accessory chromosomes was established for each of the

progeny in cross 3D1x3D7. Chromosome 14 was absent in 67 (26%) progeny,

chromosome 15 in 68 (26%) progeny, chromosome 18 in 77 (30%) progeny and

chromosome 21 in 73 (28%) progeny. All accessory chromosomes were present at a

significantly higher frequency than the expected 1:1 Mendelian inheritance ratio

(chromosome 14: χ^2 =61.1, p<0.001, chromosome 15: χ^2 = 59.1, p<0.001, chromosome

18: χ^2 = 43.1, p <0.001, chromosome 21: χ^2 = 49.9, p<0.001). This is consistent with

previous work that reported skewed inheritance of accessory chromosomes in a subset

of 48 isolates from the same population (Croll et al, 2013).

We studied the effect of accessory chromosome presence-absence on virulence. On

Runal, isolates with chromosome 21 had significantly larger pycnidia (F=6.2, p=0.014)

than isolates without chromosome 21, but the effect size was small (η^2=0.023). On Titlis,

isolates with chromosome 18 had significantly larger pycnidia (F=5.0, p=0.027) and

darker pycnidia (F=5.7, p=0.018) than isolates without chromosome 18. The effect sizes

for pycnidia size (η^2=0.021) and pycnidia melanisation (η^2=0.022) were small. A

significant interaction between chromosomes 15 and 18 was also observed for pycnidia

size (F=7.0, p= 0.009), with isolates having both accessory chromosomes showing larger

pycnidia, but the effect size of the interaction was small (η^2=0.028).

<u>Candidate genes for pathogen virulence within QTLs</u>

In total, 918 candidate genes were identified in the 95% confidence intervals of all

QTLs, including 227 from cross 3D1x3D7 and 691 from cross 1A5x1E4. No candidate

genes were shared between the two crosses. In the literature to date, 20 genes have

been implicated in virulence in *Z. tritici* and functionally characterized (Figure S1). None

of the genes previously shown to play a role in virulence were found within the QTLs.

<u>In depth investigation of the chromosome 7 QTL</u>

The QTL for PLACL and pycnidia density in cross 3D1x3D7 on Runal had a high LOD

score, a low number of candidate genes within the 95% confidence interval, and

explained approximately half of the observed phenotypic variance. Existing gene

models were manually checked for accuracy, sequence polymorphisms between the two

parent isolates were analyzed and gene expression during the infection cycle was

investigated. Population genomic approaches were used to investigate the genomic

features of the region in a natural population.

<u>Re-annotation of the genes in the chromosome 7 QTL</u>

Within the 95% confidence interval, 25 of the 35 gene models from the reference

annotation were not convincingly supported by RNAseq reads. The annotations of 20

genes were replaced with those of Grandaubert et al. (2015). Four genes not present in

any previous annotation were discovered and supported by our RNAseq data. The

annotations of four additional genes were altered slightly from those of Grandaubert et

al. (2015) or the JGI. After re-annotation, 38 genes were found in the 95% confidence

interval (Table 3). Among the newly annotated genes, GO terms could be assigned to 17

genes while the remaining 21 genes had no known function. Six genes within the QTL

encode proteins with a predicted signal peptide but lack a trans-membrane domain and

are therefore predicted to be secreted. Among these six genes, two meet the criteria for

being small secreted effector-like proteins (<300 amino acids, secreted and cysteine

rich).

<u>Identification of high priority candidate genes within the chromosome 7 QTL</u>

Sequence variation within gene coding regions of the chromosome 7 QTL was

investigated in the parental genomes. 82 sequence variants were identified, comprising

51 synonymous and 31 non-synonymous mutations as well as 4 insertions/deletions

(indels). 17 genes were found to have at least one mutation affecting amino acid

sequence. 19 genes showed differential expression *in planta* between the two parental

isolates for at least one time point during the infection process. Several high-priority

candidate genes were identified based on putative function, sequence variation and

expression levels during infection. Candidate gene *Zt_QTL7_5* is not present in any

existing annotation and BLAST searches showed no similarities to other proteins. With

a length of 65 aa, six cysteine residues and a signal peptide domain, it fulfills the criteria

of a typical small secreted effector protein. Its peak expression is between 12 and 14

dpi, during the transition from symptomless growth to the onset of chlorosis (Figure 4).

It is the second most expressed gene in the QTL confidence interval, but shows

significantly lower expression in 3D7 (the more virulent parent) at 14 dpi.

Candidate gene 00558 is also a small secreted protein containing an *Alternaria*

alternata allergen domain. This domain is unique to the Dothideomycete and

Sordariomycete classes of fungi (Chruszcz et al, 2012) and can induce major allergic

reactions in the human respiratory systems (Bush at al, 2001). This gene shows peak

expression during the symptomless phase and lower expression in 3D7 (Figure 4).

Candidate gene 105313 is a fungal-specific transporter from the Major Facilitator

Superfamily (MFS). It contains a trichothecene efflux domain that has been shown to

play a role in toxin secretion in Fusarium (Alexander et al, 1999). The parental alleles

differed by two non-synonymous SNPs. Candidate gene 00579 is a fungal-specific

membrane protein containing 7 trans-membrane domains, a CFEM domain and a signal

peptide. Expression peaks at 7 dpi (Figure 4).

Population genomics of genes in the chromosome 7 QTL

As QTL mapping is highly cross specific, genetic variation in the chromosome 7 QTL was

investigated in a natural population of 28 re-sequenced *Z. tritici* isolates from

Switzerland. The 163 kb confidence interval contained an alternation of gene-rich and

gene-poor regions. A large gene-poor region (found between ca. 1.79 - 1.84 Mb in the

reference genome assembly) contained only two genes (Figure 3E). This gene-poor

region contained all transposable elements (TEs) found within the confidence interval

(Figure 3D). Three classes of TEs were found in this region, including Copia, Gypsy and

Tad1 elements. The gene-rich regions contained a higher density of SNPs than the

average on the chromosome, reaching up to 120 SNPs per kb in some regions (Figure

3C). The genotyping rate was high in the gene-rich region. The gene-poor and TE-rich

regions contained few SNPs and had a low genotyping rate (Figure 3C) and generally

had a lower GC content than the gene-rich regions (Figure 3B).

Genes within the confidence interval were highly variable for the number of SNPs

within the natural population (Figure 3F). Seven genes contained fewer than 10 SNPs

per kb whereas 9 genes contained more then 50 SNPs per kb. On average, the genes

within the confidence interval contained 33 SNPs per kb. The Tajima's D statistic for 27

genes having 10 or more SNPs showed an alternation of clusters of genes with positive

D values and genes with negative D values (Figure 3G). Genes with negative D values are

more likely to be under purifying selection while genes with positive D values are likely

to be under balancing selection. This suggests variability in the selection processes

acting on these genes, including a possibility of hitchhiking effects.

Discussion

Here we report the most comprehensive QTL mapping analysis of fungal virulence to

date, utilizing different mapping populations and different host cultivars as well as

measuring multiple traits associated with virulence. We were able to genetically dissect

pathogen virulence into separate components, showing that different components of

virulence were under separate genetic control. QTLs were identified for all traits,

including QTLs that were specific to mapping populations, cultivars and traits as well as

QTLs that were shared among traits within the same mapping population. We identified

genome regions associated with single large effects as well as regions of smaller

additive effects.

<u>The genetic architecture of virulence in *Z. tritici* is complex</u>

Our results point to a complex genetic architecture involving multiple factors that

combine to produce quantitative virulence. Strong evidence was found for transgressive

segregation, with many progeny showing more extreme phenotypes than the parent

isolates in both crosses. QTLs were specific to each cross and in some cases specific to a

phenotype or cultivar. In all but one case, multiple traits mapped to the same QTL, an

observation also made for other traits in *Z. tritici* (Lendenmann et al. 2014; 2015; 2016).

Both quantitative (Zhan et al, 2005) and large-effect interactions consistent with the

gene-for-gene (GFG) hypothesis have been reported in *Z. tritici* (Brading et al, 2002,

Kema et al, 2000). GFG interactions are frequently reported as conferring

compatible/incompatible interactions (Flor, 1955). The large effect size of the 3D1x3D7

chromosome 7 QTL on one of the cultivars shows similarity to a GFG interaction

between a major resistance gene in Runal and a corresponding avirulence effector gene

in the 3D1 strain of the pathogen. In our case the interaction explains ~50% of the

overall variance for the virulence traits which also show a continuous distribution in the

mapping population. These observations suggest that numerous additional genes of

small effect may also be involved, in addition to the genotype-by-environment

interactions typically observed in quantitative phenotypes.

In contrast to the single large effect QTL in 3D1x3D7, numerous QTLs distributed across

several chromosomes were found for the different phenotypes in cross 1A5x1E4. These

QTLs had relatively small effects on phenotype and contained a high number of

candidate genes. This suggests that the genetic architecture of virulence in 1A5x1E4 is

more complex than that of 3D1x3D7 and demonstrates the contribution of both large

and small effects on virulence.

<u>Additive effects of QTL loci on virulence</u>

The multiple factor hypothesis (Morgan et al 1915) based on additive effects of many

genes is the key principle underlying quantitative traits. Recombination generates

progeny with different combinations of genes than the parents and can contribute to

novel virulence phenotypes (Joseph et al, 2011, Kanvil et al, 2015). Strong additive

effects were observed in cross 1A5x1E4 for the QTL for pycnidia density. For both

cultivars, alternative alleles from the two parents combined to produce a higher

phenotype. Isolates with the same parental allele at both QTLs showed lower virulence

than isolates with the different parental alleles at the two QTLs. Transgressive

segregation was observed for all phenotypes in both crosses. These findings illustrate

the power of recombination to generate progeny phenotypes that are significantly

different from their parents, an observation shared with other plant pathogenic fungi

(Sommerhalder et al, 2010; Stefansson et al. 2014) and show how a range of

quantitative virulence phenotypes can be generated through sexual reproduction in

natural field populations.

<u>The contribution of accessory chromosomes to virulence</u>

By using the novel approach of associating virulence phenotypes with the presence or

absence of specific accessory chromosomes, we discovered a significant correlation for

some phenotypes, with higher virulence observed in offspring carrying certain

accessory chromosomes. Accessory chromosomes were shown to play an important

role in virulence in other fungal pathogens (Akagi et al, 2009, Ma et al, 2010, Miao, et al,

1991) but until now their role in *Z. tritici* was elusive (Stukenbrock et al, 2010). The

increase in virulence was small, around 2-3%, but significant. If this correlation also

exists under natural field conditions, this increase in virulence and the associated

increase in reproduction would represent a significant fitness advantage that could

explain why accessory chromosomes are retained in natural field populations.

<u>Cultivar specificity</u>

Cultivar specificity is well documented in *Z. tritici* (Ahmed et al, 1995, Cowger and

Mundt 2002, Zhan et al, 2002) and is commonly reported as qualitative

compatible/incompatible interactions (e.g. Brading et al, 2002) or as generally

quantitative with isolates being more virulent or less virulent on a particular host (e.g.

Zhan et al, 2005). In cross 3D1x3D7, all phenotypes were higher on the more

susceptible cultivar as expected. However, in cross 1A5x1E4, PLACL was higher in

Runal whereas pycnidia size, density and melanization were higher in Titlis, the more

resistant cultivar. Additionally, in 1A5x1E4 the same chromosome 5 QTL was found for

pycnidia density in both cultivars. However, a second QTL was found on chromosome 3

in Runal and on chromosome 9 in Titlis, suggesting that these QTL loci play a role in

quantitative specialization to these two cultivars. This is in contrast to the findings of

Mirzadi Gohari et al, (2015) who found a QTL at a similar position on chromosome 5

that was implicated in host specificity in a different set of cultivars. The results from the

accessory chromosome presence/absence analysis also point towards cultivar

specificity, with different accessory chromosomes affecting the same phenotype on

different cultivars.

<u>Contribution of pycnidia number and size to epidemic potential</u>

The basic reproductive number (R_0) is the theoretical number of infections arising from

a single infection event and can be useful in predicting the potential development of

plant disease epidemics (van den Bosch et al, 2008). Pycnidia represent the asexual

reproductive output of a *Z. tritici* strain, with larger pycnidia correlated with a greater

number of conidia in earlier work (Gough, 1978). It was previously postulated that the

number and size of pycnidia could be important measures of virulence in *Z. tritici*

(Stewart and McDonald 2014, Suffert et al, 2013). Our results show a significant

correlation indicating that larger pycnidia bear both larger and more numerous spores.

The number and size of pycnidia influence the R_0 and therefore provide a prediction of

the epidemic potential associated with a pathogen strain under field conditions. In

contrast, measuring the leaf area covered by lesions (PLACL) gives an indication of the

amount of damage caused by a pathogen strain on the host plant. The ability to directly

measure both the epidemic potential of an isolate and the amount of damage it causes to

its host represents a powerful improvement in the overall measurement of pathogen

virulence that can also be used to improve measurements of resistance in the host

(Stewart et al. 2016). Furthermore, the identification of a QTL for pycnidia size shows

that this important trait can be under separate genetic control. A separate experiment

oriented around measuring differences in host resistance and conducted using naturally

infected plants under field conditions showed that host genotype could affect pycnidia

formation independently of damage caused by leaf lesions (Stewart et al 2016).

Theoretically, a pathogen strain which causes high levels of leaf necrosis but has a low

R_0 (i.e. producing fewer pycnidia per infected leaf) may appear more damaging at a

small spatial scale but an isolate which has a higher R_0 (i.e. producing more pycnidia per

infected leaf) has a greater potential to cause damage over larger spatial scales. From an

epidemiology perspective, directly measuring pathogen reproduction on a given host

could inform disease management decisions oriented around inhibiting pathogen

reproduction (e.g. through seeking resistance genes that lower pathogen reproduction

in addition to resistance genes that lower host damage), as well as providing a novel

way of selecting for quantitative host resistance. The results from cross 1A5x1E4

illustrate the potential usefulness of this approach with different levels of pycnidia and

PLACL found on the two different cultivars.

Candidate genes affecting pathogen virulence

By combining several disciplines, we were able to narrow the candidate gene list for the

chromosome 7 QTL affecting PLACL, pycnidia density and melanization, but it remains

possible that different genes located within this QTL interact to affect all three traits. It

also is possible that one gene in this QTL is primarily responsible for only one

phenotype, which influences the other two without any genetic or regulatory

connection, a situation referred to as spurious, vertical, relational or reactive pleiotropy

(Paaby and Rockman, 2013). For example, it is plausible that a gene encoding higher

PLACL would lead to greater pycnidia density that could in turn affect pycnidia

melanization. The results from cross 1A5x1E4 offer support for the latter hypothesis,

with multiple traits mapping to the same QTL on chromosome 5, though independent

QTLs were found on different chromosomes for the same traits. In previous work, genes

up-regulated in the biotrophic phase of the infection cycle showed reduced pycnidia

production in knockout mutants (Poppe et al, 2015), providing evidence that pycnidia

development is somewhat dependent on prior processes.

Among the candidate genes within the chromosome 7 QTL, the one we consider most

likely to explain the observed virulence is *Zt_QTL7_5*, a previously un-described small

secreted protein (SSP) that is highly expressed during the switch to necrotrophic

growth. SSPs are common virulence factors in fungi that facilitate infection or elicit a

response in the host (Lo Presti, 2015). SSPs often conform to the GFG paradigm by

interacting with a major R gene in the host. A GFG interaction would explain the large

effect of the chromosome 7 QTL on only one host. Interestingly, no significant sequence

differences exist for this gene between the parent isolates, but the expression levels are

significantly lower in 3D7, the more virulent parent. This is consistent with the theory

of effector triggered immunity (ETI), where recognition of a pathogen effector triggers a

defense response in the host. Another SSP, candidate gene 00558, contains a conserved

domain known to elicit allergic response in humans (Bush et al, 2001). It has

additionally been implicated in virulence in *Alternaria brasicicola* on Arabidopsis

(Cramer and Lawrence, 2004) and inhibits plant antimicrobial proteins (Gomez-Casado

et al, 2014). Given the existing knowledge of this protein, it appears that it could play a

role in *Z. tritici* virulence. An MFS transporter implicated in toxin secretion was also

located in the QTL. MFS transporters contribute to virulence by secretion of fungal

toxins and secondary metabolites as well as efflux of plant-derived antimicrobial

compounds (Coleman and Mylonakis, 2009). Candidate gene 00579 contains a CFEM

domain that is unique to fungi and found at significantly higher frequency in pathogenic

than non-pathogenic fungi (Zhang et al, 2015). Putative functions of the CFEM domain

include cell surface receptors, signal transduction or adhesion of molecules in plant-

pathogen interactions (Kulkarni et al, 2003). None of the genes previously described as

having a role in virulence in *Z. tritici* were identified in this study, highlighting the

usefulness of forward genetic approaches such as QTL mapping to identify novel

virulence genes.

Population genomics of the chromosome 7 QTL

Analysis of the large effect QTL on chromosome 7 in a Swiss field population revealed

several interesting genomic features. The region was highly variable, with genes

exhibiting variable numbers of SNPs as well as evidence for both purifying and

balancing selection. Genes involved in virulence are often under positive selection in

plant pathogens (Lo Presti et al, 2015; Stukenbrock and McDonald, 2009), including in

Z. tritici (Poppe et al, 2015). An island of transposable elements (TEs) was found within

a gene-poor region of the QTL. The *Z. tritici* genome contains 16.7% repetitive elements

(Dhillon et al, 2014). Effector genes show a tendency to cluster in gene-poor regions of

repetitive DNA that are TE-rich in numerous filamentous fungal plant pathogens (Dong

et al, 2015). TEs can influence the expression of effector genes by insertion into

promoter regions (Ali et al 2014) or through epigenetic gene silencing (Shaaban et el,

2010). We hypothesize that these processes may explain the differences in expression

observed in *Zt_QTL7_5.*

Conclusions

This study highlights the complex nature of virulence in the wheat-*Z. tritici*

pathosystem, illustrating that many factors contribute to quantitative phenotypes. We

showed that virulence is comprised of different traits, some affecting host damage and

others affecting pathogen reproduction, that can be under independent genetic control.

In light of this, researchers and breeders should reconsider the best way to measure

virulence in this pathosystem. We propose that more attention should be focused on

resistance that reduces pathogen reproduction in order to decrease R_0 during

epidemics.

Experimental Procedures

Two *Z. tritici* mapping populations (described in Lendenmann et al, 2014) were

phenotyped in a greenhouse-based seedling assay as described in Stewart and

McDonald (2014). The Swiss winter wheat cultivars 'Runal' and 'Titlis' (DSP Ltd, Delly,

Switzerland) were inoculated using each offspring from both crosses along with the

parental isolates. Two plants of each cultivar were inoculated with each *Z. tritici* isolate

to give two technical replicates. Each technical replicate was placed in a separate

greenhouse compartment. This process was repeated three times over three

consecutive weeks to generate three biological replicates, resulting in six replicates

total for each isolate-cultivar combination.

At 23 days post inoculation (dpi) the second leaf from each plant was excised,

photographed and phenotyped using automated image analysis as described previously

(Stewart and McDonald 2014). The method was modified slightly to include a measure

of pycnidia melanization. RGB images were converted to 8-bit greyscale and the mean

grey value for the pixels making up each pycnidium was calculated. The grey scale runs

from 0 (black) to 255 (white). Grey values can be used as a proxy for the degree of

melanisation (Lendenmann et al, 2015). The phenotypes percent leaf area covered by

lesions (PLACL), pycnidia density (pycnidia per cm^2 leaf area), pycnidia size (in mm^2)

and pycnidia melanization were used as phenotypes for QTL mapping (see below).

For the 3D1x3D7 progeny, the mean pycnidia size was calculated for each isolate on

Runal. The 10 isolates showing the largest mean pycnidia sizes and 10 showing the

smallest mean pycnidia sizes were selected for further analysis. The leaves from the six

replicates for each isolate were retrieved from storage. Each leaf was cut into ~30 mm

long sections. All sections from each leaf were placed into a 1.2 ml collection microtube

(QIAGEN) along with a 5 mm x 25 mm strip of filter paper. 100 μl sterile water was

added to each tube to moisten the filter paper. Tubes were capped and placed at 25 °C

for 24 h to provide the humid environment needed to exude the cirrhi containing spores

from pycnidia. 800 μl of water containing 0.001% TWEEN 20 was added to each tube.

Tubes were vortexed for 20 seconds to suspend the released spores. 100 μl of the spore

solution was placed into a black glass-bottomed 96 well plate (Greiner bio one, μClear).

Spores were imaged with an Olympus IX 81 inverted microscope coupled with a

Hamamatsu ORCA-ER camera using transmission illumination. Four images were made

from each well at 20x magnification with a 10% overlap between images. Images from

each well were stitched together using ImageJ (Rasband, 1997–2015). Each spore was

counted and its length was measured using the line selection tool and the measure

command in ImageJ (Figure S2). The mean spore length was used as a proxy for spore

size. The total number of spores per leaf was calculated (Figure S2) and divided by the

number of pycnidia per leaf to derive the mean number of spores per pycnidium.

All analyses were performed in base R (R core team, 2012) unless specified otherwise.

To normalize environmental variation between technical replicates and biological

replicates and account for genotype-by-environment (GxE) interactions, phenotypes

were mean centered (Schielzeth, 2010). For each phenotype, the mean value for each

greenhouse chamber and time point was calculated. This value was subtracted from all

individual values from the corresponding greenhouse chamber and time point, resulting

in a mean value of ~0 for each greenhouse chamber time point. The mean of these

transformed values was calculated for each progeny and used for subsequent analyses.

PLACL and pycnidia density values were log transformed prior to mean centering. Mean

centered values were used for analysis but untransformed values are reported in the

text and figures for clarity.

Phenotype differences between wheat cultivars were calculated using ANOVA. Effects

on phenotypes of accessory chromosome presence or absence were calculated using

ANOVA with the presence or absence of each chromosome as factors. Non-significant

factors were removed from the models. Effect sizes (η^2) were calculated using the lsr

package (Navarro, 2015). Fisher's LSD test was used to calculate the differences

between isolates with different parental alleles at QTL peaks using the Agricolae

package (de Mendiburu, 2014). Correlations between pycnidia size and spore size and

between pycnidia size and number of spores per pycnidia were made with Pearson's

correlation coefficient.

Progeny from both crosses were genotyped using restriction site associated DNA

sequencing (RADseq) based on a method adapted from Etter et al (2011) as described

previously (Lendenmann et al, 2014). The genetic map of Lendenmann et al (2014) was

used to perform QTL mapping. QTL mapping was performed using the r/QTL package

(Arends et al, 2010) following the methods described in Lendenmann et al (2014).

To establish the presence or absence of accessory chromosomes in 3D1x3D7, sequence

reads from the RADseq genotyping were aligned to the IPO323 reference genome

(Goodwin et al, 2011). For each isolate, the mean read depth was calculated for each

chromosome and divided by the mean sequence depth over all chromosomes. Thus,

chromosomes with read depth similar to the genome average have a value of ~1 and

were deemed present whereas chromosomes with low or no read depth have a value of

~0 and were deemed absent. Deviations from Mendelian inheritance among the

accessory chromosomes exhibiting presence/absence polymorphisms in the parents

were tested using a Chi square (χ^2) test.

Parent genome sequences (Croll et al, 2013) were aligned to the IPO323 reference

genome. Sequence variants were identified using the unified genotyper within GATK

(McKenna et al, 2010) and effects of variants found within genes were predicted using

snpEFF (Cingolani et al, 2012a) and snpSIFT (Cingolani et al, 2012b) following the

methods outlined in Lendenmann et al (2014).

The genes within the 95% confidence interval containing a QTL with large effect on

chromosome 7 were manually re-annotated using existing annotations from the IPO323

reference annotation (Goodwin et al, 2011) and Grandaubert et al. (2015) along with

RNAseq data from IPO323 (Rudd et al, 2015) and from the 3D7 parent isolate (Palma-

Guerrero et al, 2016). Genes were annotated using blast2go (Conesa et al. 2005). Signal

peptides were identified using signalP 4.1 (Petersen et al, 2011) and trans-membrane

domains were identified with TMHMM 2.0 (Krogh et al, 2001). The re-annotated genes

were used for all subsequent analyses of this region.

An *in planta* infection assay was performed with the parent isolates 3D1 and 3D7

(Javier Palma-Guerrero, unpublished). RNA was sequenced from leaf samples collected

at 7, 12, 14 and 28 days post inoculation (dpi) as described earlier (Palma-Guerrero et

al. 2016).

The population genomics of the genes in the 95% confidence interval of the

chromosome 7 QTL was characterized using 28 re-sequenced *Z. tritici* isolates sampled

from a single field in Switzerland. Sequences were aligned to the reference genome

using Bowtie 2 version 2.2.3 (Langmead and Salzberg, 2012). SNPs among the isolates

were identified using GATK tools (De Pristo et al. 2011). GC content was calculated in

the reference genome using 1000 bp overlapping windows, with 100 bp overlaps. The

locations of transposable elements within the reference genome were identified using

RepeatMasker (Smit et al. 2013-2015). The repeat library used for annotation was

version 20150807 downloaded from Repbase in September 2015 (Jurka et al. 2005).

The SNP genotyping rate in the 28 re-sequenced isolates was calculated in windows of

1000 bp. The SNP density was calculated in windows of 500 bp for SNPs that were

genotyped in >50% of the re-sequenced isolates. The SNP density per coding sequence

was calculated by dividing the number of SNPs with a genotyping rate > 90% by the

total coding sequence length. Tajima's D statistic (Tajima, 1989) was calculated per

gene using Popgenome in R (Pfeifer, et al, 2014). Only SNPs with a genotyping rate >

90% were included. Tajima's D values for genes with less than 10 SNPs were not

calculated.

<u>Acknowledgements</u>

The research was supported by a grant from the Swiss National Science Foundation

(31003A_134755). RADseq libraries were prepared at the Genetic Diversity Center

Zürich (GDC) and sequenced at the Quantitative Genomics Facility at the Department of

Biosystems Science and Engineering (D-BSSE) at the scientific central facilities of ETH

Zürich. Microscopy was carried out at the Scientific Center for Optical and Electron

Microscopy (ScopeM) of the ETH Zürich. Technical assistance was provided by Renato

Guidon, Thomas Kreienbühl and Leandra Knecht. Cross 3D1x3D7 was made by M.

Lendenmann. Cross 1A5x1E4 was made by Marcello Zala.

References

Ahmed, H. U., Mundt, C. C. and Coakley, S. M. (1995) Host-pathogen relationship of

geographically diverse isolates of Septoria tritici and wheat cultivars. *Plant Pathol.* **44**,

838-847.

Ali, S., Laurie, J. D., Linning, R., Cervantes-Ch·vez, J. A., Gaudet, D. and Bakkeren, G.

(2014) An Immunity-Triggering Effector from the Barley Smut Fungus *Ustilago hordei*

Resides in an Ustilaginaceae-Specific Cluster Bearing Signs of Transposable Element-

Assisted Evolution. *PLoS Pathog.* **10**, e1004223.

Akagi, Y., Akamatsu, H., Otani, H. and Kodama, M. (2009) Horizontal Chromosome

Transfer, a Mechanism for the Evolution and Differentiation of a Plant-Pathogenic

Fungus. *Eukaryotic Cell.* **8**, 1732-1738.

Alexander, J. N., McCormick, P. S. and Hohn, M. T. (1999) TRI12, a trichothecene

efflux pump from *Fusarium sporotrichioides*: gene isolation and expression in yeast. *Mol.*

Genet. Genomics. **261**, 977-984.

Almasy, L. and Blangero, J. (2009) Human QTL Linkage Mapping. *Genetica.* **136**, 333-

340.

Alonso-Blanco, C. and Méndez-Vigo, B. (2014) Genetic architecture of naturally

occurring quantitative traits in plants: an updated synthesis. *Curr. Opin. Plant Biol.* **18**,

37-43.

Arends, D., Prins, P., Jansen, R. C. and Broman, K. W. (2010) R/qtl: high-throughput

multiple QTL mapping. *Bioinformatics.* **26**, 2990-2992.

Bock, C. H., Parker, P. E., Cook, A. Z. and Gottwald, T. R. (2008) Visual Rating and the

Use of Image Analysis for Assessing Different Symptoms of Citrus Canker on Grapefruit

Leaves. *Plant Dis.* **92**, 530-541.

van den Bosch, F., McRoberts, N., van den Berg, F. and Madden, L. V. (2008) The

Basic Reproduction Number of Plant Pathogens: Matrix Approaches to Complex

Dynamics. *Phytopathology.* **98**, 239-249.

Bourras, S., McNally, K. E., Ben-David, R., Parlange, F., Roffler, S., Praz, C. R.,

Oberhaensli, S., Menardo, F., Stirnweis, D., Frenkel, Z., Schaefer, L. K., Flückiger, S.,

Treier, G., Herren, G., Korol, A. B., Wicker, T. and Keller, B. (2015) Multiple

Avirulence Loci and Allele-Specific Effector Recognition Control the Pm3 Race-Specific

Resistance of Wheat to Powdery Mildew. *The Plant Cell*. **10**,2991-3012

Brading, P. A., Verstappen, E.C.P., Kema, G.H.J., Brown, J.K.M. (2002) A gene-for-gene

relationship between wheat and *Mycosphaerella graminicola*, the Septoria tritici blotch

pathogen. *Phytopathology.* **92**, 439-445.

Brown, J. K. M., Chartrain, L., Lasserre-Zuber, P. and Saintenac, C. (2015) Genetics of

resistance to *Zymoseptoria tritici* and applications to wheat breeding. *Fungal Genet. Biol.*

79, 33-41.

Bruns, E., Carson, M. L. and May, G. (2014) The Jack of all trades is master of none: A

pathogen's ability to infect a great number of host genotypes comes at a cost of delayed

reproduction. *Evolution.* **68**, 2453-2466.

Bush, R. K. and Portnoy, J. M. (2001) The role and abatement of fungal allergens in

allergic diseases. *J. Allergy Clin. Immunol.* **107**, S430-S440.

Cowger, C., Hoffer, M. E. and Mundt, C. C. (2000) Specific adaptation by

Mycosphaerella graminicola to a resistant wheat cultivar. *Plant Pathol.* **49**, 445-451.

Cramer, R. A. and Lawrence, C. B. (2004) Identification of *Alternaria brassicicola* genes

expressed *in planta* during pathogenesis of *Arabidopsis thaliana*. *Fungal Genet. Biol.* **41**,

115-128.

Chruszcz, M., Chapman, M. D., Osinski, T., Solberg, R., Demas, M., Porebski, P. J.,

Majorek, K. A., Pomés, A. and Minor, W. (2012) *Alternaria alternata* allergen, Alt a 1 -

a unique, β-barrel protein dimer exclusively found in fungi. *J. Allergy Clin. Immunol.* **130**,

241-247.e249.

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., Land, S. J., Lu, X.

and Ruden, D. M. (2012a) A program for annotating and predicting the effects of single

nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster*

strain w(1118); iso-2; iso-3. *Fly.* **6**, 80-92.

Cingolani, P., Patel, V. M., Coon, M., Nguyen, T., Land, S. J., Ruden, D. M. and Lu, X.

(2012b) Using *Drosophila melanogaster* as a Model for Genotoxic Chemical Mutational

Studies with a New Program, SnpSift. *Front. Genet.* **3**, 35.

Cohen, L. and Eyal, Z. (1993) The histology of processes associated with the infection

of resistant and susceptible wheat cultivars with Septoria tritici. *Plant Pathol.* **42**, 737-

743.

Conesa, A., Götz, S., García-Gómez, J. M., Terol, J., Talón, M. and Robles, M. (2005)

Blast2GO: a universal tool for annotation, visualization and analysis in functional

genomics research. *Bioinformatics.* **21**, 3674-3676.

Cools, H. J., Mullins, J. G. L., Fraaije, B. A., Parker, J. E., Kelly, D. E., Lucas, J. A. and

Kelly, S. L. (2011) Impact of Recently Emerged Sterol 14α-Demethylase (*CYP51*)

Variants of *Mycosphaerella graminicola* on Azole Fungicide Sensitivity. *Appl. Environ.*

Microbiol. **77**, 3830-3837.

Courvoisier, N., Häner, L. L., Bertossa, M.,Thévoz, E., Anders, M., Stoll, P., Weisflog,

P., Dugon, J. and Grünig, K. (2015) Liste recommandée des variétés de céréales pour la

récolte 2016 Recherche Agronomique Suisse. http://www.swissgranum.ch/88-2-Listes-

recommandes.html.

Croll, D., Zala, M., and McDonald, B. A. (2013) Breakage-fusion-bridge Cycles and

Large Insertions Contribute to the Rapid Evolution of Accessory Chromosomes in a

Fungal Pathogen. *PLoS Genet.* **9**(6), e1003567–20.

Croll, D., Lendenmann, M. H., Stewart, E. and McDonald, B. A. (2015) The Impact of

Recombination Hotspots on Genome Evolution of a Fungal Plant Pathogen. *Genetics*.

201(3): 1213-1228.

Cumagun, C. J. R., Bowden, R. L., Jurgenson, J. E., Leslie, J. F. and Miedaner, T. (2004)

Genetic Mapping of Pathogenicity and Aggressiveness of *Gibberella zeae* (*Fusarium*

graminearum) Toward Wheat. *Phytopathology*. **94**, 520-526.

Dean, R., Van Kan, J. A. L., Pretorius, Z. A., Hammond-Kosack, K. E., Di Pietro, A.,

Spanu, P. D., Rudd, J. J., Dickman, M., Kahmann, R., Ellis, J. and Foster, G. D. (2012)

The Top 10 fungal pathogens in molecular plant pathology. *Mol. Plant Pathol.* **13**, 413-

430.

DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C.,

Philippakis, A. A., del Angel, G., Rivas, M. A., Hanna, M., McKenna, A., Fennell, T. J.,

Kernytsky, A. M., Sivachenko, A. Y., Cibulskis, K., Gabriel, S. B., Altshuler, D. and

Daly, M. J. (2011) A framework for variation discovery and genotyping using next-

generation DNA sequencing data. *Nat Genet.* **43**, 491-498.

Dhillon, B., Gill, N., Hamelin, R. C. and Goodwin, S. B. (2014) The landscape of

transposable elements in the finished genome of the fungal wheat pathogen

Mycosphaerella graminicola. BMC Genomics. **15**, 1-17.

Dong, S., Raffaele, S. and Kamoun, S. (2015) The two-speed genomes of filamentous

pathogens: waltz with plants. *Curr. Opin. Genet. Dev.* **35**, 57-65.

Etter, P. D., Bassham, S., Hohenlohe, P. A., Johnson, E. A. and Cresko, W. A. (2011)

SNP Discovery and Genotyping for Evolutionary Genetics Using RAD Sequencing. In

Molecular Methods for Evolutionary Genetics (ed. by V. Orgogozo & V. M. Rockman), pp.

157-178. Humana Press, Totowa, NJ.

Eyal, Z., Scharen, A.L., Prescott, J.M., van Ginkel, M. (1987) The Septoria Diseases of

Wheat. *CYMMIT*.

Flor, H. H. (1955) Host-parasite interaction in flax rust-its genetics and other

implications. *Phytopathology*. **45**:680-85.

Fones, H. and Gurr, S. (2015) The impact of Septoria tritici Blotch disease on wheat: An

EU perspective. *Fungal Genet. Biol.* **79**, 3-7.

Foulongne-Oriol, M. (2012) Genetic linkage mapping in fungi: current state,

applications, and future trends. *Appl. Microbiol. Biotechnol.* **95**, 891-904.

Friesen, T. L. and Faris, J. D. (2010) Characterization of the wheat-*Stagonospora*

nodorum disease system: what is the molecular basis of this quantitative necrotrophic

disease interaction? *Can. J. Plant Pathol.* **32**, 20-28.

781 **Furbank, R. T. and Tester, M.** (2011) Phenomics – technologies to relieve the

782 phenotyping bottleneck. *Trends Plant Sci.* **16**, 635-644.

784 **Gómez-Casado, C., Murua-García, A., Garrido-Arandia, M., González-Melendi, P.,**

785 **Sánchez-Monge, R., Barber, D., Pacios, L. F. and Díaz-Perales, A.** (2014) Alt a 1 from

786 Alternaria interacts with PR5 thaumatin-like proteins. *FEBS Lett.* **588**, 1501-1508.

788 **Goodwin, S. B., M'Barek S, B., Dhillon, B., Wittenberg, A. H., Crane, C. F., Hane, J. K.,**

789 **Foster, A. J., Van der Lee, T. A., Grimwood, J., Aerts, A., Antoniw, J., Bailey, A.,**

790 **Bluhm, B., Bowler, J., Bristow, J., van der Burgt, A., Canto-Canche, B., Churchill, A.**

791 **C., Conde-Ferraez, L., Cools, H. J., Coutinho, P. M., Csukai, M., Dehal, P., De Wit, P.,**

792 **Donzelli, B., van de Geest, H. C., van Ham, R. C., Hammond-Kosack, K. E., Henrissat,**

793 **B., Kilian, A., Kobayashi, A. K., Koopmann, E., Kourmpetis, Y., Kuzniar, A.,**

794 **Lindquist, E., Lombard, V., Maliepaard, C., Martins, N., Mehrabi, R., Nap, J. P.,**

795 **Ponomarenko, A., Rudd, J. J., Salamov, A., Schmutz, J., Schouten, H. J., Shapiro, H.,**

796 **Stergiopoulos, I., Torriani, S. F., Tu, H., de Vries, R. P., Waalwijk, C., Ware, S. B.,**

797 **Wiebenga, A., Zwiers, L. H., Oliver, R. P., Grigoriev, I. V. and Kema, G. H.** (2011)

798 Finished genome of the fungal wheat pathogen *Mycosphaerella graminicola* reveals

799 dispensome structure, chromosome plasticity, and stealth pathogenesis. *PLoS Genet.* **7**,

800 e1002070.

802 **Gough, F. J.** (1978) Effect of Wheat Host Cultivars on Pycnidiospore Production by

803 Septoria tritici. *Phytopathology.* **68**, 1343-1345.

Grandaubert, J., Bhattacharyya, A. and Stukenbrock, E. H. (2015) RNA-seq Based

Gene Annotation and Comparative Genomics of Four Fungal Grass Pathogens in the

Genus Zymoseptoria Identify Novel Orphan Genes and Species-Specific Invasions of

Transposable Elements. *G3*. **5**(7):1323-1333.

Habgood, R. M. (1977) Resistance of barley cultivars to *Rhynchosporium secalis. Trans.*

Br. Mycol. Soc. **69**, 281-286.

Halama, P., Skajennikoff, M. and Dehorter, B. (1999) Tetrad analysis of mating type,

mutations, esterase and aggressiveness in *Phaeosphaeria nodorum. Mycol. Res.* **103**, 43-

49.

Hansen, T. F. (2006) The Evolution of Genetic Architecture. *Annu. Rev. Ecol. Evol. Syst.*

37, 123-157.

Jørgensen, L. N., Hovmøller, M. S., Hansen, J. G., Lassen, P., Clark, B., Bayles, R.,

Rodemann, B., Flath, K., Jahn, M., Goral, T., Jerzy Czembor, J., Cheyron, P.,

Maumene, C., De Pope, C., Ban, R., Nielsen, G. C. and Berg, G. (2014) IPM Strategies

and Their Dilemmas Including an Introduction to www.eurowheat.org. *J. Integr. Agric.*

13, 265-281.

Joseph, B., Schwarz, R. F., Linke, B., Blom, J., Becker, A., Claus, H., Goesmann, A.,

Frosch, M., M¸ller, T., Vogel, U. and Schoen, C. (2011) Virulence Evolution of the

Human Pathogen *Neisseria meningitidis* by Recombination in the Core and Accessory

Genome. *PLoS ONE.* **6**, e18441.

Jurka, J., Kapitonov, V. V., Pavlicek, A., Klonowski, P., Kohany, O. and Walichiewicz, J. (2005) Repbase Update, a database of eukaryotic repetitive elements. *Cytogenet. Genome Res.* **110**, 462-467.

Kanvil, S., Collins, C. M., Powell, G. and Turnbull, C. G. N. (2015) Cryptic Virulence and Avirulence Alleles Revealed by Controlled Sexual Recombination in Pea Aphids. *Genetics.* **199**, 581-593.

Kema, J. G. H., Verstappen, P. E. C., Todorova, M. and Waalwijk, C. (1996) Successful crosses and molecular tetrad and progeny analyses demonstrate heterothallism in *Mycosphaerella graminicola. Curr. Genet.* **30**, 251-258.

Kema, G. H. J., Verstappen, E. C. P. and Waalwijk, C. (2000) Avirulence in the Wheat Septoria tritici Leaf Blotch Fungus *Mycosphaerella graminicola* Is Controlled by a Single Locus. *Mol. Plant Microbe Interact.* **13**, 1375-1379.

Kokko, E. G., Conner, R. L., Lee, B., Kuzyk, A. D. and Kozu, G. C. (2000) Quantification of common root rot symptoms in resistant and susceptible barley by image analysis. *Can. J. Plant Pathol.* **22**, 38-43.

Krattinger, S. G., Lagudah, E. S., Spielmeyer, W., Singh, R. P., Huerta-Espino, J., McFadden, H., Bossolini, E., Selter, L. L. and Keller, B. (2009) A Putative ABC Transporter Confers Durable Resistance to Multiple Fungal Pathogens in Wheat. *Science,* **323**, 1360-1363.

Krogh, A., Larsson, B., von Heijne, G. and Sonnhammer, E. L. L. (2001)

Predicting transmembrane protein topology with a hidden Markov model: Application

to complete genomes. *J of Mol. Bio.* **305**(3):567-580.

Kulkarni, R. D., Kelkar, H. S. and Dean, R. A. (2003) An eight-cysteine-containing

CFEM domain unique to a group of fungal membrane proteins. *Trends Biochem. Sci.* **28**,

118-121.

Langmead B, and Salzberg S. (2012) Fast gapped-read alignment with Bowtie 2.

Nature Methods. **9**:357-359.

Lannou, C. (2011) Variation and Selection of Quantitative Traits in Plant Pathogens.

Annu. Rev. Phytopathol. **50**, 319-338.

Larraya, L. M., Alfonso, M., Pisabarro, A. G. and Ramirez, L. (2003) Mapping of

genomic regions (quantitative trait loci) controlling production and quality in industrial

cultures of the edible basidiomycete *Pleurotus ostreatus. Appl. Environ. Microbiol.*

69(6):3617-3625.

Lendenmann, M. H., Croll, D., Stewart, E. L., & McDonald, B. A. (2014) Quantitative

trait locus mapping of melanization in the plant pathogenic fungus *Zymoseptoria tritici.*

G3. **4**(12), 2519–2533.

Lendenmann, M. H., Croll, D. and McDonald, B. A. (2015) QTL mapping of fungicide

sensitivity reveals novel genes and pleiotropy with melanization in the pathogen

Zymoseptoria tritici. Fungal Genet. Biol. **80**, 53-67.

Lendenmann, M. H., Croll, D., Palma-Guerrero, J., Stewart, E. L. and McDonald, B. A.

(2016) QTL mapping of temperature sensitivity reveals candidate genes for thermal

adaptation and growth morphology in the plant pathogenic fungus *Zymoseptoria tritici.*

Heredity, **116**, 384-394.

Lo Presti, L., Lanver, D., Schweizer, G., Tanaka, S., Liang, L., Tollot, M., Zuccaro, A.,

Reissmann, S. and Kahmann, R. (2015) Fungal Effectors and Plant Susceptibility.

Annu. Rev. Plant. Biol. **66**, 513-545.

Lin, X., Huang, J. C., Mitchell, T. G. and Heitman, J. (2006) Virulence Attributes and

Hyphal Growth of *C. neoformans* Are Quantitative Traits and the *MATα* Allele Enhances

Filamentation. *PLoS Genet.* **2**, e187.

Lind, M., Dalman, K., Stenlid, J., Karlsson, B. and Olson, Å. (2007) Identification of

quantitative trait loci affecting virulence in the basidiomycete *Heterobasidion annosum*

s.l. Curr. Genet. **52**, 35-44.

Liu, Z., Friesen, T. L., Ling, H., Meinhardt, S. W., Oliver, R. P., Rasmussen, J. B. and

Faris, J. D. (2006) The *Tsn1–ToxA* interaction in the wheat–*Stagonospora nodorum*

pathosystem parallels that of the wheat–tan spot system. *Genome,* **49**, 1265-1273.

Ma, L.-J., van der Does, H. C., Borkovich, K. A., Coleman, J. J., Daboussi, M.-J., Di

Pietro, A., Dufresne, M., Freitag, M., Grabherr, M., Henrissat, B., Houterman, P. M.,

Kang, S., Shim, W.-B., Woloshuk, C., Xie, X., Xu, J.-R., Antoniw, J., Baker, S. E., Bluhm,

B. H., Breakspear, A., Brown, D. W., Butchko, R. A. E., Chapman, S., Coulson, R.,

Coutinho, P. M., Danchin, E. G. J., Diener, A., Gale, L. R., Gardiner, D. M., Goff, S.,

Hammond-Kosack, K. E., Hilburn, K., Hua-Van, A., Jonkers, W., Kazan, K., Kodira, C.

D., Koehrsen, M., Kumar, L., Lee, Y.-H., Li, L., Manners, J. M., Miranda-Saavedra, D.,

Mukherjee, M., Park, G., Park, J., Park, S.-Y., Proctor, R. H., Regev, A., Ruiz-Roldan,

M. C., Sain, D., Sakthikumar, S., Sykes, S., Schwartz, D. C., Turgeon, B. G., Wapinski,

I., Yoder, O., Young, S., Zeng, Q., Zhou, S., Galagan, J., Cuomo, C. A., Kistler, H. C. and

Rep, M. (2010) Comparative genomics reveals mobile pathogenicity chromosomes in

Fusarium. *Nature.* **464**, 367-373.

Mackay, T. F. C. (2001) The Genetic Architecture of Quantitative Traits. *Annu. Rev.*

Genet. **35**, 303-339.

Martin, D. P. and Rybicki, E. P. (1998) Microcomputer-Based Quantification of Maize

Streak Virus Symptoms in *Zea mays. Phytopathology.* **88**, 422-427.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A.,

Garimella, K., Altshuler, D., Gabriel, S., Daly, M. and DePristo, M. A. (2010) The

Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA

sequencing data. *Genome Res.* **20**, 1297-1303.

de Mendiburu, F. (2014) Agricolae: Statistical procedures for agricultural research. R

package version 1.1-8. http://CRAN.R-project.org/package=agricolae

Miao, V. P., Covert, S. F. and VanEtten, H. D. (1991) A fungal gene for antibiotic

resistance on a dispensable ("B") chromosome. *Science.* **254**, 1773-1776.

Mirzadi Gohari, A., Ware, S. B., Wittenberg, A. H. J., Mehrabi, R., Ben M'Barek, S.,

Verstappen, E. C. P., van der Lee, T. A. J., Robert, O., Schouten, H. J., de Wit, P. P. J. G.

M. and Kema, G. H. J. (2015) Effector discovery in the fungal wheat pathogen

Zymoseptoria tritici. Mol Plant Pathol. **16**, 931-945.

Morgan, T. H., Sturtevant, A. H., Muller, H. J., and C. B. Bridges (1915) The

Mechanism of Mendelian Heredity. New York: Henry Holt and Company.

Navaro, D. (2015) lsr: Companion to "Learning Statistics with R". https://cran.r-

project.org/web/packages/lsr/

Nosanchuk, J. D. and Casadevall, A. (2003) The contribution of melanin to microbial

pathogenesis. *Cell. Microbiol.* **5**, 203-223.

Nosanchuk, J. D. and Casadevall, A. (2006) Impact of Melanin on Microbial Virulence

and Clinical Resistance to Antimicrobial Compounds. *Antimicrob. Agents Chemother.* **50**,

3519-3528.

O'Driscoll, A., Kildea, S., Doohan, F., Spink, J. and Mullins, E. (2014) The wheat–

Septoria conflict: a new front opening up? *Trends Plant Sci.* **19**, 602-610.

Oliver, R. P., Friesen, T. L., Faris, J. D. and Solomon, P. S. (2012) *Stagonospora*

nodorum: From Pathology to Genomics and Host Resistance. *Annu. Rev. Phytopathol.* **50**,

23-43.

Palma-Guerrero, J., Torriani, S. F. F., Zala, M., Carter, D., Courbot, M., Rudd, J. J.,

McDonald, B. A. and Croll, D. (2016) Comparative transcriptomic analyses of

Zymoseptoria tritici strains show complex lifestyle transitions and intraspecific

variability in transcription profiles. *Mol Plant Pathol.* DOI: 10.1111/mpp.12333

Paaby, A. B. and Rockman, M. V. (2013) The many faces of pleiotropy. *Trends Genet.*

29, 66-73.

Pariaud, B., Robert, C., Goyeau, H. and Lannou, C. (2009) Aggressiveness

Components and Adaptation to a Host Cultivar in Wheat Leaf Rust. *Phytopathology.* **99**,

869-878.

Parlevliet, J. E. (1979) Components of Resistance that Reduce the Rate of Epidemic

Development. *Annu. Rev. Phytopathol.* **17**, 203-222.

Petersen, T. N., Brunak, S., von Heijne, G. and Nielsen, H. (2011) SignalP 4.0:

discriminating signal peptides from transmembrane regions. *Nat. Meth.* **8**, 785-786.

Pfeifer, B., Wittelsbürger, U., Ramos-Onsins, S. E., & Lercher, M. J. (2014)

PopGenome: an efficient Swiss army knife for population genomic analyses in R. *Mol.*

Bio. Evol. **31**(7), 1929–1936.

Poppe, S., Dorsheimer, L., Happel, P. and Stukenbrock, E. H. (2015) Rapidly Evolving

Genes Are Key Players in Host Specialization and Virulence of the Fungal Wheat

Pathogen *Zymoseptoria tritici*(*Mycosphaerella graminicola*). *PLoS Pathog.* **11**, e1005055.

Portnoy, D. A., Auerbuch, V. and Glomski, I. J. (2002) The cell biology of *Listeria*

monocytogenes infection: the intersection of bacterial pathogenesis and cell-mediated

immunity. *J. Cell Biol.* **158**, 409-414.

R Core Team. (2012) R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna. http://www.R-project.org/

Rasband, W.S. (1997-2015) ImageJ; U.S. National Institutes of Health, Bethesda, MD.

(http://imagej.nih.gov/ij/).

Reese, M. L., Zeiner, G. M., Saeij, J. P. J., Boothroyd, J. C. and Boyle, J. P. (2011)

Polymorphic family of injected pseudokinases is paramount in Toxoplasma virulence.

Proc. Natl. Acad. Sci. **108**, 9625-9630.

Schielzeth, H. (2010) Simple means to improve the interpretability of regression

coefficients. *Methods Ecol. Evol.* **1**, 103-113.

1002

1003   **Shaaban, M., Palmer, J. M., El-Naggar, W. A., El-Sokkary, M. A., Habib, E. L. S. E. and**

1004   **Keller, N. P.** (2010) Involvement of transposon-like elements in penicillin gene cluster

1005   regulation. *Fungal Genet. Biol.* **47**, 423-432.

1006

1007   **Smit A.F.A., Hubley R., Green P.** (2013-2015) RepeatMasker Open-4.0.

1008   http://www.repeatmasker.org.

1009

1010   **Solberg Woods, L. C.** (2014) QTL mapping in outbred populations: successes and

1011   challenges. *Physiol. Genomics.* **46**, 81-90.

1012

1013   **Sommerhalder, R. J., McDonald, B. A., Mascher, F. and Zhan, J.** (2010) Sexual

1014   Recombinants Make a Significant Contribution to Epidemics Caused by the Wheat

1015   Pathogen *Phaeosphaeria nodorum. Phytopathology.* **100**, 855-862.

1016

1017   **St.Clair, D. A.** (2010) Quantitative Disease Resistance and Quantitative Resistance Loci

1018   in Breeding. *Annu. Rev. Phytopathol.* **48**, 247-268.

1019

1020   **Stefansson, T. S., McDonald, B. A. and Willi, Y.** (2013) Local adaptation and

1021   evolutionary potential along a temperature gradient in the fungal pathogen

1022   *Rhynchosporium commune. Evol. Appl.* **6**, 524-534.

1023

1024   **Stergiopoulos, I. and de Wit, P. J. G. M.** (2009) Fungal Effector Proteins. *Annu. Rev.*

1025   *Phytopathol.* **47**, 233-263.

1026

1027 **Stewart, E. L. and McDonald, B. A.** (2014) Measuring Quantitative Virulence in the

1028 Wheat Pathogen *Zymoseptoria tritici* Using High-Throughput Automated Image

1029 Analysis. *Phytopathology.* **104**, 985-992.

1030

1031 **Stewart, E. L, Hagerty, C. H., Mikaberidze, A., Mundt, C. C., Zhong, Z. and McDonald,**

1032 **B. A.** (2016) An improved method for measuring quantitative resistance to the wheat

1033 pathogen *Zymoseptoria tritici* using high throughput automated image analysis.

1034 *Phytopathology*. http://dx.doi.org/10.1094/PHYTO-01-16-0018-R

1035

1036 **Stukenbrock, E. H. and McDonald, B. A.** (2009) Population Genetics of Fungal and

1037 Oomycete Effectors Involved in Gene-for-Gene Interactions. *Mol. Plant Microbe Interact.*

1038 **22**, 371-380.

1039

1040 **Stukenbrock, E. H., Jorgensen, F. G., Zala, M., Hansen, T. T., McDonald, B. A. and**

1041 **Schierup, M. H.** (2010) Whole-Genome and Chromosome Evolution Associated with

1042 Host Adaptation and Speciation of the Wheat Pathogen *Mycosphaerella graminicola*.

1043 *PLoS Genet.* **6**, e1001189.

1044

1045 **Su, C., Howe, D. K., Dubey, J. P., Ajioka, J. W. and Sibley, L. D.** (2002) Identification of

1046 quantitative trait loci controlling acute virulence in *Toxoplasma gondii. Proc. Natl. Acad.*

1047 *Sci.* **99**, 10753-10758.

1048

1049 **Suffert, F., Sache, I., Lannou, C.** (2013) Assessment of quantitative traits of

1050 aggressiveness in *Mycosphaerella graminicola* on adult wheat plants. *Plant Pathol.* **62**,

1051 1330-1341.

1052

1053 **Tajima, F.** (1989) Statistical method for testing the neutral mutation hypothesis by DNA

1054 polymorphism. *Genetics.* **123**(3), 585–595.

1055

1056 **Twizeyimana, M., Ojiambo, P. S., Bandyopadhyay, R. and Hartman, G. L.** (2014) Use

1057 of Quantitative Traits to Assess Aggressiveness of *Phakopsora pachyrhizi* Isolates from

1058 Nigeria and the United States. *Plant Dis.* **98**, 1261-1266.

1059

1060 **Wittenberg, A. H. J., van der Lee, T. A. J., Ben M'Barek, S., Ware, S. B., Goodwin, S. B.,**

1061 **Kilian, A., Visser, R. G. F., Kema, G. H. J. and Schouten, H. J.** (2009) Meiosis Drives

1062 Extraordinary Genome Plasticity in the Haploid Fungal Plant Pathogen *Mycosphaerella*

1063 *graminicola. PLoS ONE.* **4**, e5863.

1064

1065 **Xie, W., Yu, K., Pauls, K. P. and Navabi, A.** (2011) Application of Image Analysis in

1066 Studies of Quantitative Disease Resistance, Exemplified Using Common Bacterial Blight–

1067 Common Bean Pathosystem. *Phytopathology.* **102**, 434-442.

1068

1069 **Yang, L., Gao, F., Shang, L., Zhan, J. and McDonald, B. A.** (2013) Association between

1070 Virulence and Triazole Tolerance in the Phytopathogenic Fungus *Mycosphaerella*

1071 *graminicola. PLoS ONE.* **8**, e59568.

1072

1073 **Zhan, J., Mundt, C. C., Hoffer, M. E. and McDonald, B. A.** (2002) Local adaptation and

1074 effect of host genotype on the rate of pathogen evolution: an experimental test in a plant

1075 pathosystem. *Evolutionary Biology.* **15**, 634-647.

1076

1077 **Zhan, J., Linde, C. C., Jurgens, T., Merz, U., Steinebrunner, F. and McDonald, B. A.**

1078 (2005) Variation for neutral markers is correlated with variation for quantitative traits

1079 in the plant pathogenic fungus *Mycosphaerella graminicola*. *Mol. Ecol.* **14**(9):2683-2693.

1080

1081 **Zhan, J., Torriani, S. F. F. and McDonald, B. A.** (2007) Significant difference in

1082 pathogenicity between *MAT1-1* and *MAT1-2* isolates in the wheat pathogen

1083 *Mycosphaerella graminicola. Fungal Genet. Biol.* **44**, 339-346.

1084

1085 **Zhang, Z.-N., Wu, Q.-Y., Zhang, G.-Z., Zhu, Y.-Y., Murphy, R. W., Liu, Z. and Zou, C.-G.**

1086 (2015) Systematic analyses reveal uniqueness and origin of the CFEM domain in fungi.

1087 *Scientific Reports.* **5**, 13032.

1088

1089 **Zwiers, L.-H., Roohparvar, R. and de Waard, M. A.** (2007) *MgAtr7*, a new type of ABC

1090 transporter from *Mycosphaerella graminicola* involved in iron homeostasis. *Fungal*

1091 *Genet. Biol.* **44**, 853-863.

1092

1093

1094

1095   **Tables**

1096

1097

Table 1. Significant QTLs for *in planta* virulence traits from *Z. tritici* mapping populations 3D1x3D7 (A) and 1A5x1E4 (B) on wheat cultivars Runal and Titlis.

A

Cultivar	Phenotype	Chr	LOD Score	Variance Explained (%)	LOD Threshold	p value	Confidence Interval (Mb)	Number of Genes	Mean Phenotype (3D1 allele)	Mean Phenotype (3D7 allele)
Runal	PLACL	7	43.7	53.9	3.34	<0.001	1.75-1.91	35	38.98	78.19
Runal	Pycnidia per cm^2	7	48.2	57.3	3.30	<0.001	1.75-1.91	35	11.26	34.23
Runal	Grey Value[†]	7	11.4	18.1	3.29	<0.001	1.18-1.91	227	71.72	78.44

B

Cultivar	Phenotype	Chr	LOD Score	Variance Explained (%)	LOD Threshold	p value	Confidence Interval (Mb)	Number of Genes	Mean Phenotype (1A5 allele)	Mean Phenotype (1E4 allele)
Runal	Pycnidia per cm^2	3	4.9	8.6	3.38	0.003	1.27-1.87	209	24.79	32.13
Runal	Pycnidia per cm^2	5	3.8	7.0	3.38	0.018	0.14-0.23	280	31.02	26.60
Runal	Pycnidia Size (mm^2 x 10^{-3})	3	5.1	8.7	3.33	0.001	1.27-1.98	143	4.773	4.998
Runal	Grey Value[†]	5	4.7	8.1	3.31	0.001	0.14-0.15	89	88.06	84.74
Titlis	Pycnidia per cm^2	5	4.15	7.2	3.38	0.006	1.36-2.27	280	28.03	22.62
Titlis	Pycnidia per cm^2	9	3.51	6.1	3.38	0.045	3.20-9.19	204	22.37	28.01
Titlis	PLACL	5	4.56	8.0	3.17	0.015	1.36-2.27	280	95.49	90.40

[†]The grey scale ranges from 0 (black) to 255 (white). Lower values indicate darker pycnidia.

Table 2. Additive allele effects for pycnidia density in cross 1A5x1E4 for cultivars Runal (A) and Titlis (B). Chromosome alleles represent the parental allele of the marker at the QTL peak. Phenotypes are the mean pycnidia density of all isolates with the corresponding allele combination. Superscript letters denote significant differences. n represents the number of isolates with each allele combination.

A

Cultivar	Chr 3 allele	Chr 5 allele	Mean Phenotype	n
Runal	1E4	1A5	39.28[a]	43
Runal	1A5	1A5	26.10[b]	32
Runal	1E4	1E4	26.96[b]	44
Runal	1A5	1E4	23.09[c]	46

B

Cultivar	Chr 5 allele	Chr 9 allele	Mean Phenotype	n
Titlis	1A5	1E4	33.38[a]	37
Titlis	1A5	1A5	25.00[b]	41
Titlis	1E4	1E4	25.15[b]	54
Titlis	1E4	1A5	20.11[c]	36

Table 3. Re-annotated candidate genes located in a QTL for pycnidia density and PLACL on chromosome 7 in *Z. tritici*. Superscripts denote the source of the gene annotation: [a] new genes not present in previous annotations, [b] annotations modified from existing annotations, [c] annotation from the JGI reference annotation, [d] annotation from Grandaubert et al. (2015).

Gene Name	Description	Protein Length	Signal Peptide	TM Domain
Zt_QTL7_1[a]	NA	504	N	N
Zt_QTL7_2[a]	NA	228	Y	N
Zt_QTL7_3[a]	NA	357	N	N
Zt_QTL7_4[b]	Hypothetical protein	471	N	Y
Zt_QTL7_5[a]	NA	198	Y	N
Zt_QTL7_6[b]	AMP-binding enzyme family	921	N	Y
Zt_QTL7_7[b]	Serine carboxypeptidase	966	N	N
Zt_QTL7_8[b]	Hypothetical protein	879	N	N
Mycgr3T100839[c]	Fe-containing alcohol dehydrogenase,	1251	N	N
Mycgr3T100840[c]	Vacuolar membrane pq loop repeat	972	N	Y
Mycgr3T105313[c]	Trichothecene efflux pump	1788	N	Y
Mycgr3T44611[c]	Radical s-adenosyl methionine domain-containing 2	1074	N	Y
Mycgr3T44946[c]	Hypothetical protein	666	N	N
Mycgr3T45542[c]	Lipoate- ligase B	720	N	N
Mycgr3T74280[c]	Prolyl 4-hydroxylase	1503	N	N
Mycgr3T94634[c]	Hypothetical protein	525	N	N
Mycgr3T94648[c]	Hypothetical protein	1143	N	N
Mycgr3T94659[c]	Hypothetical protein	972	N	N
Zt09_model_7_00553[d]	Ferric-chelate reductase (Fre2)	2085	N	Y
Zt09_model_7_00555[d]	Hypothetical protein	255	N	N
Zt09_model_7_00556[d]	Hypothetical protein	1038	N	N
Zt09_model_7_00557[d]	Hypothetical protein	378	N	N
Zt09_model_7_00558[d]	Major allergen alt a1, partial	540	Y	N
Zt09_model_7_00559[d]	Hypothetical protein	789	N	Y
Zt09_model_7_00561[d]	Hypothetical protein	1302	N	N
Zt09_model_7_00562[d]	Hypothetical protein	3663	Y	N
Zt09_model_7_00563[d]	Hypothetical protein	1851	N	N
Zt09_model_7_00564[d]	Hypothetical protein	1761	N	Y
Zt09_model_7_00567[d]	Hypothetical protein	651	N	N
Zt09_model_7_00569[d]	Acid phosphatase	456	N	Y
Zt09_model_7_00571[d]	Adenylate kinase	696	N	N
Zt09_model_7_00573[d]	NA	420	N	N
Zt09_model_7_00574[d]	Glycoside hydrolase	933	N	N
Zt09_model_7_00575[d]	Glycoside hydrolase family 36	2238	Y	N
Zt09_model_7_00577[d]	Ca2+-modulated nonselective cation channel polycystin	1917	Y	N
Zt09_model_7_00578[d]	Urease accessory	780	N	N
Zt09_model_7_00579[d]	CFEM domain-containing	1230	Y	Y
Zt09_model_7_00581[d]	Hypothetical protein	294	N	N

Figure Legends

Figure 1. Frequency histograms of virulence phenotypes from the *Z. tritici* mapping populations 3D1x3D7 (upper panels) and 1A5x1E4 (lower panels) on the wheat cultivars Runal (white bars) and Titlis (grey bars). Vertical lines represent phenotype values of the parent isolates, dotted line represents 3D1 and 1A5, dashed line represents 3D7 and 1E4.

Figure 2. LOD (logarithm of the odds) plots from QTL mapping of virulence traits in the *Z. tritici* mapping populations 3D1x 3D7 (A) and 1A5x1E4 (B) for the 13 core chromosomes (x axis). The dashed horizontal line represents the 0.05 significance threshold calculated with 1000 permutations.

Figure 3. Population genomic characterization of the QTL located on chromosome 7 (region 1746 kb to 1909 kb) in the reference isolate IPO323 and a population of 28 *Zymoseptoria tritici* isolates drawn from a single field in Switzerland. (A). GC content in the IPO323 reference genome. (B). SNPs density (bottom) and genotyping rate (top) in the 28 field isolates. Black rectangles represent the frequency of missing SNP genotypes. The horizontal line represents the mean SNP density across chromosome 7 in the 28 isolates. (C). Transposable element locations in the IPO323 reference genome. (D). *Bottom:* Location of coding sequences in the IPO323 reference genome. *Top:* Frequency of the allele carried by 3D7 in the 28 field isolates for the SNPs shared by the 3D1 and 3D7 parents in the gene coding sequences. (E). SNP density per kb in the coding sequences of the genes. Only SNPs with a genotyping rate ≥ 90% were

included. (F). Tajima's D statistic per gene coding sequence. Tajima's D values for genes with less than 10 SNPs are not shown.

Figure 4. Gene expression profiles of *Zymoseptoria tritici* candidate genes at 7, 12, 14 and 28 days post infection (DPI). Solid red lines represent parent isolate 3D1, dashed blue lines represent parent isolate 3D7.

Supplementary Figures

Figure S1. Genes previously implicated in virulence for *Z. tritici*. Adapted from Orton et al, 2011.

Figure S2. Typical microscope images of *Z. tritici* pycnidiospores showing many long, overlapping spores (top left) and short, less dense spores (top right). Yellow lines represent spores counted and measured via a semi-automated ImageJ macro (bottom panels).

Figure S3. GFF file containing details of the re-annotated genes in the chromosome 7 QTL.

Figure S4. blast2go results for the chromosome 7 QTL.