Title: Adaptively introgressed Neandertal haplotype at the OAS locus functionally

impacts innate immune responses in humans.

Authors

Aaron J. Sams[1,*†], Anne Dumaine[2†], Yohann Nédélec[2,3†], Vania Yotova[2], Caroline

Alfieri[2,4], Jerome E. Tanner[2], Philipp W. Messer[1*††], and Luis B. Barreiro[2,5,*††]

Affiliations

[1]Department of Biological Statistics and Computational Biology, Cornell

University, Ithaca, NY

[2]Department of Genetics, Sainte-Justine Hospital Research Centre, University of

Montreal, Montreal, Qc, Canada

[3]Department of Biochemistry, University of Montreal, Montreal, Qc, Canada

[4]Department of Microbiology and Immunology, University of Montreal, Montreal,

Qc, Canada

[5]Department of Pediatrics, University of Montreal, Montreal, Qc, Canada

[†]These authors contributed equally to this work

[††]These authors jointly supervised this work

[*]Correspondence to: (A.J.S.) as2847@cornell.edu, (P.W.M.) messer@cornell.edu,

(L.B.B.) luis.barreiro@umontreal.ca

Abstract

The 2'-5' oligoadenylate synthetase (OAS) locus encodes for three OAS enzymes

(OAS1-3) involved in innate immune response. This region harbors high amounts

of Neandertal ancestry in non-African populations; yet, strong evidence of positive

selection in the OAS region is still lacking. Here we used a combination of neutral

coalescent simulations and neutrality tests to firmly demonstrate a signal of

adaptive introgression at the OAS locus. Furthermore, we characterized the

functional consequences of the Neandertal haplotype in the transcriptional

regulation of OAS genes at baseline and infected conditions. We found that cells

from people with the Neandertal-like haplotype express lower levels of *OAS3*

upon infection, as well as distinct isoforms of *OAS1* and *OAS2*. Notably, the

Neandertal-introgressed haplotype reintroduced an ancestral splice variant of

OAS1 encoding a more active protein, suggesting that adaptive introgression

occurred as a means to resurrect adaptive variation that had been lost outside

Africa.

Introduction

Whole genome sequencing of several archaic human genomes(Green et al., 2010;

Reich et al., 2010; Meyer et al., 2012; Prüfer et al., 2014; Sawyer et al., 2015)

representing Neandertals and an as yet geographically and paleontologically

unknown population referred to as Denisovans has revealed gene flow between

these populations and the ancestors of present-day humans. Neandertal ancestry

makes up approximately 0.5-2 percent of the ancestry of most living humans, with

higher amounts of Neandertal ancestry found outside of Africa(Sankararaman et

al., 2014; Vernot and Akey, 2014; Llorente et al., 2015). While it seems that there

may have been widespread purifying selection against Neandertal ancestry on the

ancestral African genomic background(Sankararaman et al., 2014; Vernot and

Akey, 2014), some positive selection on Neandertal genes (adaptive introgression)

has also been observed(Racimo et al., 2015; 2016). Neandertals and other archaic

populations inhabited Eurasia for several hundred thousand years(Hublin, 2009),

therefore some adaptive introgression is expected, particularly across phenotypes

that are strongly influenced by direct interactions with the surrounding

environment (Racimo et al., 2015), such as our immune response to infectious

agents (Ségurel and Quintana-Murci, 2014).

The OAS locus on chromosome 12, which harbors three genes (*OAS1, OAS2,*

OAS3) encoding the 2'-5' oligoadenylate synthetase enzymes has received

considerable attention due to its clear signs of multiple archaic haplotypes in

populations outside of Africa(Mendez et al., 2012; 2013), and the critical role of

OAS genes in the innate immune response to viruses (Player and Torrence, 1998).

The major Neandertal haplotype at the OAS locus spans a ~190 kilobase region

between two surrounding recombination hotspots. This haplotype is absent from

sub-Saharan African samples in the 1000 Genomes Project data, yet found at

relatively high frequencies outside of Africa, reaching highest frequencies in

European population samples (up to 41%, Figure 1). Indeed, the Neandertal

haplotypes in the OAS region are among the most common Neandertal haplotypes

among Europeans(Sankararaman et al., 2014).

The elevated frequency of the introgressed haplotype along with the key role OAS

genes play in protective immunity against viral infections raises the possibility that

introgressed Neandertal haplotypes at OAS may have been adaptive in modern

humans. While some studies provide suggestive evidence of adaptive introgression

at the OAS locus (Sankararaman et al., 2014; Racimo et al., 2016), strong

evidence of positive selection in the OAS region is still lacking. Indeed, most

studies failed to reject a model of neutral evolution for the Neandertal haplotype

when using standard neutrality tests(Mendez et al., 2013; Deschamps et al., 2016).

We hypothesize that the overall lack of signals of selection in the OAS region

stems from the low power of standard neutrality tests to detect adaptive

introgression(Sankararaman et al., 2016), and the particular genomic architecture

of the region, which is characterized by the presence of two strong recombination

hotspots (Figure 2A). The presence of these recombination hotspots is likely to

significantly reduce power to detect signals of selection, particularly for tests

based on haplotype-lengths or levels of linkage disequilibrium associated with the

selected allele. Here we circumvent these issues by testing the hypothesis of

adaptive introgression using extensive neutral coalescent simulations specifically

tailored to match the genomic features of the OAS region in combination with

ancient DNA data from Eurasia. We firmly demonstrate a population genetic

signal of adaptive introgression at the OAS locus and characterize the functional

consequences of the Neandertal haplotype in the transcriptional regulation of OAS

genes in macrophages and peripheral blood mononuclear cells (PBMCs) at

baseline and infected conditions.

Results

To investigate the hypothesis of non-neutral evolution at the OAS locus we

simulated neutral sequence data using a demographic model based on several

previously inferred parameters of human demographic history(Gravel et al., 2011;

Tennessen et al., 2012; Vernot and Akey, 2015)(Figure S1) and explicitly

incorporated estimated recombination rates in the OAS region (Methods). Our

model included a single pulse of Neandertal introgression occurring over a span of

500-1000 years (sampled from a uniform distribution) into the ancestral Eurasian

population after their population split from Africa. We investigated patterns of

haplotype homozygosity surrounding Neandertal-like sites (NLS), derived alleles

that are shared between Neandertals and a non-African population sample (CEU),

but absent in a sub-Saharan African sample (YRI). This class of sites is frequently

used as a conservative indicator of Neandertal introgression(Yang et al., 2012; Fu

et al., 2014; Sankararaman et al., 2014). In each simulation we chose a NLS at

random to measure two summary statistics, NLS frequency, and $H_{D/A}$, a statistic

which measures the ratio of average pairwise homozygosity tract lengths among

haplotypes carrying a derived versus ancestral allele at each NLS (Methods). This

statistic is based on the H statistic that provides a general measure for haplotype

lengths and has been shown to be robust in detecting both hard and soft selective

sweeps(Schlamp et al., 2016). If Neandertal haplotypes have experienced a greater

rate of positive selection than ancestral haplotypes, this should lead to a relative

increase of H among the subgroup of haplotypes that carry Neandertal alleles

versus those carrying only ancestral alleles, leading to higher values of $H_{D/A}$

compared to a neutrally evolving locus with similar recombination rates (Figure

S2). Additionally, we expected the frequency of NLS at the OAS locus to be

elevated compared to the NLS frequency distribution observed across neutral

simulations.

In practice, we observed both signals. We first examined the probability in our

model that neutrally evolving Neandertal like sites would be segregating at their

current frequencies. We found that the highest frequency NLS (DAF = 38.4%) fall

in the extreme 1% of all simulations (lowest $P = 0.0055$) (Figure 2A). Next, we

compared $H_{D/A}$ at each NLS in the OAS region to distributions of simulated NLS

drawn to match local recombination rate and frequency (Methods), thereby

controlling for associations between frequency and recombination, and haplotype

lengths. We found several NLS in the OAS region that have $H_{D/A}$ values greater

than 99% of comparable simulated NLS (lowest $P = 0.0023$) (Figure 2B). Further,

we measured mean $H_{D/A}$ across all NLS in the OAS region and compared this to

randomly sampled distributions of simulated NLS (again matching frequency and

recombination rate). We found that the true mean $H_{D/A}$ in OAS is greater than four

standard errors from the neutral expectation ($P < 10^{-5}$) (Figure 2D). Lastly, we

considered the joint probability of observing a neutral NLS in our simulations with

frequencies and $H_{D/A}$ values as high as those observed in the OAS region. Only

413 simulated NLS (out of 1,000,079) had both a higher frequency and $H_{D/A}$ value

than the highest frequency/ $H_{D/A}$ pair in our NLS sample ($P < 5 \times 10^{-4}$) (Figure 2E).

Combined, we identified the strongest evidence to date in support of adaptive

introgression in the OAS region.

We further supported and contextualized our simulation results by examining the

OAS region in a dataset of 230 ancient Eurasians(Mathieson et al., 2015).

Assuming neutrality, the expected frequency of an allele in contemporary

European populations can be predicted as a linear combination of allele

frequencies sampled from representative ancient populations that have contributed

ancestry to present-day European populations in different proportions(Lazaridis et

al., 2014; Mathieson et al., 2015) (Methods). Therefore, using ancient allele

frequencies estimated by Mathieson and colleagues, we calculated the expected

allele frequency in four present day European samples from the 1,000 Genomes

Project(Consortium, 2015) at 11 NLS falling within the bounds of the three OAS

genes. To set up our null expectations we performed a similar analysis on a dataset

of approximately one million SNPs scattered around the genome, generated by

Mathieson and colleagues (by merging 213 ancient samples dated between 6500

and 300 BCE with sequencing data from four European samples from the 1,000

Genomes Project). We found that OAS NLS SNPs are outliers in the genome with

respect to deviations from ancient frequencies. More specifically, we found that

the allele frequency of 6 out of the 11 OAS SNPs tested in the *OAS1-OAS3* region

have increased above the frequency predicted by ancient Eurasian samples by

more than 20%, significantly more than what we observed for other SNPs

genome-wide with comparable present-day frequencies (lowest $P = 0.00476$,

Figure 2C, Table S1). Our findings at this single locus are consistent with results

from the genome-wide selection scan performed by Mathieson and

colleagues(2015) where the OAS region was also identified as an outlier.

Our population genetic results provide evidence that Neandertal alleles at the OAS

locus have likely experienced positive selection at some point after their

introduction into the human population, suggesting a possible functional role of

these alleles in human innate immune responses. To study this possibility, we

analyzed RNA-sequencing data collected on primary macrophages from 99

European-descent individuals, before and after *in-vitro* infection with *Salmonella*

typhimurium. After 2 hours of infection, we found that all OAS genes were

strongly up-regulated (up to 19-fold, $P < 1$ x 10^{-10}, Figure S3), confirming the

ability of *Salmonella* to activate the interferon (IFN) production pathway(Nauciel

and Espinasse-Maes, 1992; Gulig et al., 1997; LaRock et al., 2015). Using

genotype data available for the same individuals (673 SNPs spanning the OAS

region, see methods) we tested if NLS were associated with variation in the

expression levels of *OAS1*, *OAS2* or *OAS3,* in either infected or non-infected

macrophages. We found that SNPs linked with the Neandertal haplotype reduced

the expression levels of *OAS3* (i.e., they were expression quantitative trait loci, or

cis eQTL for *OAS3*) (Figure 3A, false discovery rate (FDR) < 5%). Interestingly,

these *cis* eQTL showed a much stronger effect in infected macrophages (best *P*

$_{salmonella} = 3.5$x10^{-3} *vs* best *P* $_{non-infected} = 0.027$), supporting an interaction between

the Neandertal haplotype and the *OAS3* response to *Salmonella* infection.

In addition to overall changes in expression, we took advantage of the power of

RNA-sequencing data to test if NLS in the OAS regions influenced the ratio of

alternative isoforms used for each of the OAS genes (i.e., alternative splicing

QTL: asQTL). We found that SNPs associated with the Neandertal haplotype are

significant asQTL for *OAS1* and *OAS2* in both infected and non-infected

macrophages (FDR<< 1%; Figure 3B-C). The effect of the splice site variant

rs10774671 at determining what isoform is primarily encoded by *OAS1* was

particularly strong ($P \leq 2x10^{-32}$). The ancestral G allele at this SNP (A<u>G</u> at

acceptor site) retains the splice site whereas the derived allele, A, (A<u>A</u> at acceptor

site) disrupts the splice site leading to the usage of a distinct isoform (Figure S4).

The Neandertal haplotype harbors the ancestral allele (encoding the p46 isoform),

which is associated with high enzyme activity(Bonnevie-Nielsen et al., 2005),

whereas the derived allele – present in virtually all non-Neandertal haplotypes

found outside Africa (98.4% in CEU) – is associated with reduced enzymatic

activity.

Because OAS genes are primarily involved in the control of viral infections we

decided to validate our functional findings on peripheral blood mononuclear cells

(PBMCs) from 30 individuals stimulated/infected with viral-ligands (polyI:C and

gardiquimod), and live viruses (Influenza, Herpes simplex virus (HSV) 1 and

HSV2). The individuals were chosen based on their genotype for the NLS

rs1557866, a SNP that is a strong proxy for the presence or absence of the

Neandertal haplotype in the OAS region (9 were homozygous for the Neandertal

haplotype, 9 were heterozygous, and 12 homozygous for the modern human

sequence).

As expected, we found that all viral-associated immune triggers led to a marked

increase in *OAS1-3* gene expression levels, as measured by real-time PCR (up to

30-fold, $P \leq 5.4 \times 10^{-6}$, Figure 4A), concomitantly with the up-regulation of type-I

and type-II interferon genes (Figure S5). Confirming the QTL results obtained in

macrophages, we found that rs10774671 was a strong asQTL for *OAS1* in both

non-infected and infected PBMCs ($P \leq 3.6 \times 10^{-5}$, Figure 4B). Likewise, we found

that the presence of the Neandertal haplotype was associated with reduced

expression levels of *OAS3,* particularly in PBMCs infected with influenza (P =

3.9×10^{-4}) and the synthetic ligand gardiquimod (P = 1.0×10^{-3}), which mimics a

single strand RNA infection (Figure 4B). Interestingly, the Neandertal haplotype

harbors additional regulatory variants that only impact expression levels in a cell-

type and immune stimuli specific fashion. For example, we found that the

Neandertal haplotype is associated with increased expression levels of *OAS2* in

non-infected (*P*=0.009), HSV1-infected (*P*=0.04), and gardiquimod-stimulated

PBMCs (*P*=0.0055), but not in macrophages nor in PBMCs treated with other

viral agents. Collectively, our functional data shows a pervasive impact of the

Neandertal haplotype on the regulation of *OAS* genes. Moreover, they show that

the regulatory impact of the Neandertal haplotype varies depending on the cell

type and the immune stimuli the cells are responding to.

Discussion

The Neandertal lineage was present in Eurasia for at least 400,000 years(Meyer et

al., 2016), providing ample time for Neandertals to adapt to local disease

environments. The admixture process, which likely fostered the transmission of

pathogens between Neandertals and humans migrating out of Africa, could have

also led to the exchange of genes useful in responding to local pathogens. Here,

we have demonstrated that a previously reported case of Neandertal introgression

at the OAS locus(Mendez et al., 2013) displays signatures of positive selection in

the European population. Additionally, we have strengthened the case for adaptive

introgression by providing direct functional evidence of a role for the Neandertal

OAS haplotype to the regulatory responses in innate immune cells to infectious

agents.

Under a model of constant directional selection with codominant fitness effects for

Neandertal alleles we can use present-day frequencies and a plausible

establishment frequency at the time of introgression to estimate the selection

coefficient associated with the Neandertal haplotype at the OAS locus (Methods).

Assuming an establishment frequency of 0.02, and introgression approximately

2000-2400 generations ago the selection coefficient in heterozygotes would be $s \sim$

0.0014 - 0.0017. However, the observed allele frequency shift of 0.26 in only 200-

340 generations (maximum shift in CEU from ancient samples; see Figure 2C)

predicts that the selection coefficient associated with the Neandertal haplotype was

2.6-5.4 times larger during recent human evolution ($s \sim 0.0044 - 0.0075$). These

results suggest that the Neandertal OAS haplotype has not experienced constant

directional selection in humans.

Our results show that the Neandertal haplotype at OAS is associated with several

regulatory variants that reduce expression of *OAS3* in response to infection, as

well as encode alternate isoforms of *OAS1* and *OAS2*. These dramatic functional

implications of the Neandertal OAS haplotype support our case for adaptive

introgression at OAS. Yet, because distinct functional polymorphisms segregate

together in the same haplotype, inferring the exact variant(s) targeted by positive

selection remains a daunting task. We speculate, however, that the direct target of

selection is likely to have been the splice variant identified in *OAS1*.

The Neandertal haplotype carries the ancestral allele (G) of the OAS1 splice

variant (rs10774671), which is common both inside and outside of Africa.

However, outside of Africa, the only haplotypes carrying this ancestral splice site

are closely related to the Neandertal haplotype, with a few exceptions being rare

recombined haplotypes (~2% of all haplotypes with the ancestral allele). This

pattern reflects the possibility that Neandertal introgression, in effect, served as a

means to resurrect the ancestral splice site from local extinction outside of Africa,

probably following the out-of-Africa exodus. The Neandertal-introgressed allele

encodes a protein variant (p46) that is associated with higher enzymatic

activity(Bonnevie-Nielsen et al., 2005). The adaptive potential of this variant is

supported by the observation that this variant (or other variants in strong LD with

it) was shown to be associated with: *(i)* reduced infection and replication rates of

West Nile virus ((Lim et al., 2009), but see (Bigham et al., 2011)), *(ii)* improved

resistance to hepatitis C virus (HCV) infection(Awady et al., 2011; Kwon et al.,

2012), and *(iii)* variable symptomology of Tick-Borne Encephalitis (TBE) Virus-

Induced Disease (homozygous individuals for the Neandertal haplotype show the

most severe symptoms of TBE). Strikingly, West Nile, hepatitis C and TBE are

all members of the *Flaviviridae* family, suggesting that Flaviviruses might have

been the main drivers of selection in *OAS1*.

The differential responses of homozygous carriers of the Neandertal OAS

haplotype to the different viruses described above suggest that the Neandertal

haplotype is not uniformly beneficial in humans. Thus, it is plausible that alleles at

OAS, and particularly the *OAS1* splice variant, might be evolving under balancing

selection. This hypothesis is supported by the observation that the *OAS1* splice

variant (rs10774671) is found at high frequency worldwide (0.11-0.7), and that

OAS1 is among the most diverse genes in both humans and non-human primates.

Indeed, a recent analysis of genome-wide sequence data from a total of 55

individuals from four non-human ape species, chimpanzee (*Pan troglodytes*

ellioti), bonobo (*Pan paniscus*), gorilla (*Gorilla gorilla gorilla*), and orangutan

(*Pongo abelii*), identified *OAS1* as in the top 1% of genes showing the largest

levels of nucleotide diversity among ape species, consistent with a scenario of

long-term balancing selection (*OAS2* and *OAS3* are ranked in the 60th and 36th

percentile of the genome-wide distribution, respectively). Further supporting the

idea of balancing selection on the introgressed haplotypes, our functional data

suggest that the Neandertal haplotype contributes a range of gene expression

responses in a cell-type and stimulus-specific manner.

In conclusion, our study demonstrates that the frequency and haplotype

distribution of Neandertal-like sites can be used in a neutral simulation framework

that accounts for local genomic context to investigate the history of selection at a

candidate locus for which genome-wide tests of selection provide ambiguous

results. When combined with functional data, our results provide the strongest

evidence to date in support of adaptive introgression in the OAS region. More

generally, our study raises the possibility that adaptive introgression might not

necessarily occur to select newly introduced variants but rather as a means to

resurrect adaptive variation into modern humans who had lost it due to

demographic events.

Materials and Methods

1. Genome alignments and identification of Neandertal-like sites

Human/chimpanzee ancestral states were computed by parsimony using

alignments from the UCSC Genome Browser for the human reference (hg19) and

three outgroups chimpanzee (panTro2), orangutan (ponAbe2), and rhesus macaque

(rheMac2)(Kent et al., 2002). Ancestral state was assumed to be the chimpanzee

allele (if available) if its state was confirmed by matching either orangutan or

macaque. All sites with no inferred ancestral state were removed from our

analysis.

We filtered the Altai Neandertal genome(Prüfer et al., 2014) using the map35_50

set of minimum filters provided at

(https://bioinf.eva.mpg.de/altai_minimal_filters/). We combined this filtered

dataset with the CEU (Utah residents (CEPH) with Northern and Western

European Ancestry) and YRI (Yoruba in Ibadan, Nigeria) samples from the 1000

Genomes Project Phase 3(Consortium, 2015), which we downloaded from

(https://mathgen.stats.ox.ac.uk/impute/1000GP%20Phase%203%20haplotypes%2

06%20October%202014.html).

We first extracted all variants that were biallelic in our human (CEU+YRI),

Neandertal, and chimpanzee alignments. We considered as Neandertal-like sites

(NLS) only those variants where the African sample (YRI) had a derived allele

frequency of zero and both CEU and Neandertal carried the derived

allele. Finally, we required the derived allele to be present in CEU in at least two

copies in order to calculate our haplotype-based test statistic ($H_{D/A}$).

2. Demographic model and neutral coalescent simulations

We performed coalescent simulations of the demographic history of the European,

African, and Neandertal populations applied by Vernot and Akey(2015) based on

previously inferred demographic models (Gravel et al., 2011; Tennessen et al.,

2012) with some minor adjustments, including removing the East Asian

component of the model and allowing for Neandertal admixture to occur over a

time span from 500 to 1000 years (Figure S1). Simulations were performed with

macs(Chen et al., 2009) in order to explicitly simulate the genetic map

(downloaded with the 1000 Genomes samples at the link above) of the 2Mb region

centered on OAS (chr12:112400000-114400000). All simulations were performed

with the parameters specified in Figure S1, assuming 25 years per generation and a

mutation rate of 2.5 x 10^{-8} per bp per generation. A sample macs command is

given at the end of this section.

Data were thinned in a manner similar to Sankararaman et al. 2014 (2014) to

account for imperfect SNP ascertainment in the 1000 Genomes dataset, such that

SNPs with minor allele count of 1, 2, 3, 4, 5, 6, 7, 8, 9, and >=10 were accepted

with probabilities 0.25, 0.5, 0.75, 0.8, 0.9, 0.95, 0.96, 0.97, 0.98, and 0.99,

respectively. Additionally, we only kept SNPs that were polymorphic in the

simulated CEU sample. Finally, we performed an additional thinning of SNPs

with uniform probability of 0.05 of removal to account for slightly elevated SNP

density in the simulated data. The resulting simulated datasets had an average

SNP density of 4.9 SNPs per kb compared to 3.6 in the real data. This is a slightly

larger than ideal difference in SNP density, but we note that neither derived allele

frequency, nor our primary haplotype-based test statistic (described below) should

be particularly sensitive to SNP density. In fact, Figure S6 illustrates that our

statistic is conservative with respect to SNP density.

To estimate the probabilities of adaptive introgression conditional on the simulated

locus containing a NLS (SNPs in which the non-African and Neandertal sample

contain a derived allele that is absent from the African sample) we randomly

selected a single NLS from the central fifth of the chromosome (positions 0.4-0.6

in the unit scaled locus) in which we calculated our test statistics. If an NLS was

found, its position on the chromosome was recorded and test statistics were

calculated. If no NLS was found, we repeated the simulation step. This approach

allowed us to estimate under our demographic model, the proportion of

independent loci that will have no obvious signal of introgression under neutrality

as the proportion of successful simulation runs to the total number of attempts. We

obtained an approximate 21 percent chance of observing Neandertal introgression

at the OAS locus under a neutral model. Additionally, we repeated this simulation

approach, setting Neandertal introgression to zero, in a set of 10,000 simulations in

order to estimate the likelihood of observing a NLS in the case of incomplete

lineage sorting (ILS), to estimate the contribution of ILS to our null distribution.

We observed a roughly 0.6% chance that a NLS is produced by ILS at the OAS

locus, suggesting that our null distribution is not significantly biased by ILS.

Sample macs command:

macs 416 2000000 -t 0.000731 -R oas_recrates.txt -I 3 216 2 198 0 -n 1

58.0027359781 -n 2 0.205198358413 -n 3 70.0410396717 -eg 0 1 482.67144247 -

eg 1e-08 3 570.175408787 -em 2e-08 1 3 0.409323665059 -em 3e-08 3 1

0.409323665059 -eg 0.00699726402189 3 16.9958723831 -en 0.00699727402189

1 1.98002735978 -en 0.03146374829 3 0.759313629055 -en 0.0561902522706 3

1.15461721321 -em 0.0561902622706 1 3 4.386 -em 0.0561902722706 3 1 4.386

-em 0.056088019083 3 2 17.3006865925 -em 0.0569023606299 3 2 0.0 -ej

0.08129305892 3 1 -en 0.08129306892 1 1.98002735978 -en 0.202462380301 1

1.0 -ej 0.89014331235 2 1 -h 1e3 1e3

3. Frequency and haplotype-based tests of neutrality

We examined the consistency of genetic variation with our neutral model using

several approaches. First, we examined the likelihood of observing (Neandertal)

allele frequencies as high as the OAS locus. Under neutrality, allele frequency is

not dependent upon recombination rate, therefore, we can estimate the likelihood

of our observed NLS frequency in the OAS region. For example, mean allele

frequency of all NLS observed between chr12:113344000-113450000 is 0.35

(Figure 2A). Frequencies this great or greater were observed in our simulations

less than one percent of the time (0.0085), suggesting that this haplotype

frequency is unlikely under neutrality.

Additionally, we wanted to examine if the haplotypes carrying NLS at the OAS

locus are longer than expected under neutrality when conditioning on the observed

frequencies and the underlying genetic map, which would provide an additional

signature of selection on introgressed haplotypes. For this purpose, we modified a

simple haplotype statistic H(Schlamp et al., 2016), which measures the average

length of pairwise homozygosity tracts in base pairs – a quantity that is very

straightforward to interpret. As selective sweeps are expected to create long

haplotypes around the selected site, the H statistic should be higher in samples

containing positively selected haplotypes compared to samples containing

neutrally evolving haplotypes, when frequency and recombination are properly

controlled, similar to other statistics based on haplotype lengths, such as *EHH,*

iHS, and *nSL* (Sabeti, P.C. et al., 2002; Voight et al., 2006; Ferrer-Admetlla et al.,

2014). However, in contrast to these other statistics, H does not require

specification of analysis parameters such as minimum haplotype homozygosity

levels below which haplotypes are no longer extended.

Under adaptive introgression, we specifically expect the introgressed haplotypes to

be longer than the ancestral haplotypes. We therefore defined our test statistic,

$H_{D/A}$, as:

$H_{D/A} = \ln(H_D/H_A)$;

where H_D and H_A are H calculated across haplotypes carrying derived NLS allele

versus the ancestral allele.

We calculated this statistic for every simulated NLS and for every NLS in our true

sample. We compared our simulation results to the observed data in two

ways. First, for each observed SNP we gathered all simulated NLS within 0.025

frequency and 20 kb. We then calculate the empirical likelihood of observing

each observed NLS under our neutral demographic model (Figure 2A). All NLS

within the bounds of the three OAS genes are tightly linked, therefore, the peak

signal at the center of this locus suggests that the probability of observing these

$H_{D/A}$ values are less than one percent.

However, to examine a single hypothesis test of the null hypothesis, that the true

Neandertal haplotype is not significantly longer than expected under neutrality, we

performed a randomization test. First, we calculated mean $H_{D/A}$ across this central

locus (chr12:113344000-113450000). Next, we randomly chose one control SNP

per true NLS (using the same matching conditions above) and calculated the mean

of the control set. We repeated this procedure 100,000 times to create a normally

distributed null distribution, which we then used to calculate a Z-score (4.5) and

corresponding P (3.35×10^{-6}) for our one-sided hypothesis test (Figure 2B).

Finally, we asked how unusual our observation of NLS frequency and $H_{D/A}$ scores

were across all simulations. Across approximately one million simulations

(1,000,079), we only observed 413 simulations that produced an NLS with a

frequency and $H_{D/A}$ score as great or greater than the highest value observed

among the true NLS (Figure 2C). In other words, the likelihood of our true sample

under the neutral demographic scenario is less than 5×10^{-4}.

4. Analysis of ancient Eurasian data

We utilized supplementary data table 3 from Mathieson et al. (2015). This table

includes maximum likelihood allele frequency estimates for three ancient

population samples (HG- Hunter-gatherer, EF- Early farmer, SA- Steppe ancestry)

and four present day European samples from the 1,000 Genomes Project (see

"Genome-wide scan for selection" section of methods in(Mathieson et al., 2015)).

We intersect this table with allele frequencies for 1,000 Genomes Yorubans (YRI)

and the Altai Neandertal genotypes and only analyze sites for which we have data

for all samples (1,004,612 SNPs).

To calculate the expected allele frequency in modern samples under drift, we used

the estimated proportions (*m*) of (HG, EF, SA) in each of the four present-day

samples estimated by Mathieson et al. (2015): CEU = (0.196, 0.257, 0.547), GBR

= (0.362, 0.229, 0.409), IBS = (0, 0.686, 0.314) and TSI = (0, 0.645, 0.355). We

calculated the expected frequency E[*p*] of site as:

$$E[p] = \sum_{i}^{\{HG,EF,SA\}} (p_i \; x \; m_i)$$

Next, we calculated the absolute difference between observed and expected allele

frequency in all four present-day European samples at all available sites. To test

the null hypothesis that OAS NLS have not changed in frequency more than

expected under neutrality at 11 SNPs in the central OAS region

(chr12:113200000-113600000), we first calculated the fraction of all autosomal

SNPs in the dataset at similar present-day frequency (within 1 percent in the

folded frequency spectrum) of each OAS NLS. Finally, we calculated the fraction

of autosomal SNPs with an absolute observed minus expected frequency

difference greater than or equal to the OAS NLS. These results are given in Table

1 and illustrated in Figure 2A.

This test does not explicitly incorporate variance in estimated ancient allele

frequency. However, any bias in ancient allele frequency estimation should be

distributed randomly across the genome. Therefore, our comparison to a genome-

wide distribution of SNPs at similar present-day frequency should incorporate

most of this error. Nonetheless, the selection test performed by Mathieson et

al.(2015) does incorporate such error, so we can also look to the Ps from that test

to ensure consistency with our results. In fact, when considering OAS as a single

locus, these results are highly significant and consistent with ours (see Table S1).

5. Estimation of selection coefficients

To estimate the selection coefficient *s* under constant positive selection for a given

starting frequency (x_0), final frequency (x_1), and number of generations between

these estimates (Δt), we assumed a model of standard logistic growth of a

codominant allele:

$x_1 = x_0/[x_0 + (1 - x_0)\exp(-s\Delta t)]$.

This equation can be easily solved to obtain *s*, given x_0, x_1, and Δt.

6. Sample collection

Buffy coats from 99 healthy European-descent donors were obtained from Indiana

Blood Center (Indianapolis, IN, USA). Only individuals self-reported as currently

healthy and not under medication were included in the study. The project was

approved by the ethics committee at the CHU Sainte-Justine (protocol #4022). The

individuals recruited in this study were males aged 18 to 55 years old.

7. DNA Extraction and genotyping

DNA from each of the blood donors was extracted using the Gentra Pure Gene

blood kit (Qiagen). Genotyping of each individual was then performed by

Illumina's HumanOmni5Exome bead Chip array and complemented with imputed

data from the 1000 Genomes data using Impute2(Howie et al., 2009). Here, we

only focused on genetic diversity surrounding the OAS region –

chr12:113229549-113574044 (~ 344Kb) spanning from the beginning of *RPH3A*

to the end of *RASAL1* – for a total of 673 SNPs with a MAF above 10%.

8. Isolation of monocytes and differentiation of macrophages

Blood mononuclear cells were isolated by Ficoll-Paque centrifugation. Monocytes

were purified from peripheral blood mononuclear cells (PBMCs) by positive

selection with magnetic CD14 MicroBeads (Miltenyi Biotech) using the

autoMACS Pro Separator. All samples had purity levels above 90%, as measured

by flow cytometry using an antibody against CD14 (BD Biosciences). Monocytes

were then cultured for 7 days in RPMI-1640 (Fisher) supplemented with 10%

heat-inactivated FBS (FBS premium, US origin, Wisent), L-glutamine (Fisher)

and M-CSF (20ng/mL; R&D systems). Cell cultures were fed every 2 days with

complete medium supplemented with the cytokines previously mentioned. Before

infection, we systematically verified that the differentiated macrophages presented

the expected phenotype for non-activated macrophages (CD1a+, CD14+, CD83−,

and HLA-DRlow (BD Biosciences)).

9. Bacterial preparation and infection of macrophages

The day prior to infection, aliquots of *Salmonella typhimurium* (Keller strain) were

thawed and bacteria were grown overnight in Tryptic Soy Broth (TSB) medium.

Bacterial culture was diluted to mid-log phase prior to infection and supernatant

density was checked at OD600. Monocyte-derived macrophages were then

infected with *Salmonella typhimurium* at a multiplicity of infection (MOI) of 10:1.

A control group of non-infected macrophages was treated the same way but using

medium without bacteria. After 2 hours in contact with the bacteria, macrophages

were washed and cultured for another hour in the presence of 50 mg/ml of

gentamycin in order to kill all extracellular bacteria present in the medium. The

cells were then washed a second time and cultured in complete medium with 3

mg/ml gentamycin for an additional 2 hours, the time point to which we refer in

the main text.

10. Infection/stimulation of PBMC

PBMCs from a subset of 30 individuals used to derive macrophages were cultured

in RPMI-1640 (Fisher) supplemented with 10% heat-inactivated FBS (FBS

premium, US origin, Wisent) and 1% L-glutamine (Fisher). The 30 individuals

were chosen based on their genotype for kgp4570197, a SNP which derived allele

is of Neandertal origin and that we used as a proxy to identify individuals

harbouring the Neandertal haplotype in the OAS region. From the 30 individuals,

and based on this SNP, 9 individuals were homozygous for the Neandertal

haplotype, 9 were heterozygous, and 12 homozygous for the modern human

sequence.

For each of the tested individuals, PBMCs (1 million per condition) were

stimulated/infected with one of the following viral-associated immune challenges:

polyI:C (10 μg/ml, TLR3 agonist), gardiquimod (0.5μg/ml, TLR7 and TLR8

agonist), Influenza PR8 WT (multiplicity of infection (MOI) of 0.05:1), Herpes

simplex virus (HSV) 1 (1.55×10^2 CPE), and HSV2 (19.5×10^4 CPE). PBMCs were

stimulated/infected for 4 hours with TLR ligands and Influenza, and 6h with

HSV1 and HSV2. A control group of non-infected PBMC was treated the same

way but with only medium.

11. RNA extraction, RNA-seq library preparation, and sequencing

Total RNA was extracted from the non-infected and infected/stimulated cells

using the miRNeasy kit (Qiagen). RNA quantity was evaluated

spectrophotometrically, and the quality was assessed with the Agilent 2100

Bioanalyzer (Agilent Technologies). Only samples with no evidence of RNA

degradation (RNA integrity number > 8) were kept for further experiments. RNA-

sequencing libraries were prepared using the Illumina TruSeq protocol. Once

prepared, indexed cDNA libraries were pooled (6 libraries per pool) in equimolar

amounts and sequenced with single-end 100bp reads on an Illumina HiSeq2500.

Results based on the entire dataset are described elsewhere (Nédélec *et al*., under

revision). Here, we only studied transcript-level and gene-level expression

estimates for *OAS1*, *OAS2* and *OAS3*.

12. Quantifying gene expression values from RNA-seq data

Adaptor sequences and low quality score bases (Phred score < 20) were first

trimmed using Trim Galore (version 0.2.7). The resulting reads were then mapped

to the human genome reference sequence (Ensembl GRCh37 release 65) using

TopHat (version 2.0.6) and using a hg19 transcript annotation database

downloaded from UCSC. Gene-level expression estimates were calculated using

featureCounts (version 1.4.6-p3) and transcript-level expression values were

obtained using RSEM under default parameters.

13. Quantitative real time PCR

For the PBMC samples we measured the expression levels of OAS and interferon

genes using real time PCR. 100ng of high-quality RNA was reverse-transcribed

into cDNA using the qScript cDNA SuperMix (Quanta Biosciences). Quantitative

real time PCR was performed using 96.96 Dynamic Array™ IFCs and the

BioMark™ HD System from Fluidigm. For the TaqMan gene assays, we used the

following TaqMan Gene Expression Assay (Applied BioSystems) to quantify the

expression levels of interferon genes: *IFNA1* (Hs03044218), *IFNA6*

(Hs00819627), and *IFNG* (Hs00989291). To quantify the overall expression levels

of OAS genes, we used probes that capture all common isoforms of *OAS1*

(Hs00973635), *OAS2* (Hs00942643), and OAS3 (Hs00196324). Custom-made

probes were designed to specifically target the short-isoform of *OAS2* (Forward

Primer Sequence CTGCAGGAACCCGAACAGTT; Reverse Primer Sequence

ACTCATGGCCTAGAGGTTGCA;   Reporter   Sequence

AGAGAAAAGCCAAAGAA). As housekeeping genes we used: *GAPDH*

(Hs02758991), *GUSB* (Hs99999908), *HPRT1* (Hs99999909), and *POLR2A*

(Hs00172187). The results reported in the manuscript used *POLR2A* as a reference

but all conclusions remain unchanged when using any of the other housekeeping

genes.

We start by doing a preamplification of the cDNA using the PreAmp Master Mix

(Fluidigm). Preamplified cDNA was then diluted 2X on a solution of 10 mM Tris–

HCl (pH 8.0) and 0.1 mM EDTA. In order to prepare samples for loading into the

integrated fluid circuit (IFC), a mix was prepared consisting of 360 μL TaqMan

Fast Advanced Master Mix (Applied BioSystems) and 36 μL 20× GE Sample

Loading Reagent (Fluidigm). 2.75 μL of this mix was dispensed to each well of a

96-well assay plate and mixed with 2.25 μL of preamplified cDNA. Following

priming of the IFC in the IFC Controller HX, 5 μL of the mixture of cDNA and

loading reagent were dispensed in each of the sample inlet of the 96.96 IFC. For

the TaqMan gene assays, 5 μL of mixes consisting of 2.5 μL 20× TaqMan Gene

Expression Assay (Applied BioSystems) and 2.5 μL 2X Assay Loading Reagent

(Fluidigm) were dispensed to each detector inlet of the 96.96 IFC. After loading

the assays and samples into the IFC in the IFC Controller HX, the IFC was

transferred to the BioMark HD and PCR was performed using the thermal protocol

GE 96 × 96 Fast v1.pcl. This protocol consists of a Thermal Mix of 70 °C, 30 min;

25 °C, 10 min, Hot Start at 95 °C, 1 min, PCR Cycle of 35 cycles of (96 °C, 5 s;

60 °C, 20 s). Data was analysed using Fluidigm Real-Time PCR Analysis software

using the Linear (Derivative) Baseline Correction Method and the Auto

(Detectors) Ct Threshold Method.

To quantify the expression levels of the *OAS1* isoform associated with the derived

allele at the splicing variant rs10774671 we used SybrGreen and the following

forward (GCTGAGGCCTGGCTGAATTA), and reverse

(CCACTTGTTAGCTGATGTCCTTGA) primers. PCR was performed using the

thermal protocol 50 °C, 2 min; 95 °C, 10 min, PCR Cycle of 40 cycles of (95 °C,

15 s; 60 °C, 1 min). A melting curve was also performed to check for non-specific

amplification.

14. Genotype–Phenotype Association Analysis

eQTL, asQTL were performed against *OAS1*, *OAS2* and *OAS3*. We examined

associations between SNP genotypes and the phenotype of interest using a linear

regression model, in which phenotype was regressed against genotype. In

particular, expression levels were considered as the phenotype when searching for

eQTL and the percentage usage of each isoform in each gene when mapping

asQTL. To avoid low power caused by rare variants, only SNPs in the OAS region

with a minor allele frequency of 10% across all individuals were tested (i.e., 673

SNPs within the region chr12:113229549-113574044). In all cases, we assumed

that alleles affected the phenotype in an additive manner. For the eQTL and

asQTL analyses on macrophages we mapped Salmonella-infected, and non-

infected samples separately. For the non-infected and infected/stimulated PBMCs

we only tested expression levels against the SNPs identified as eQTL or asQTL in

the macrophage data (specifically, the SNPs for which boxplots are shown in

Figure 3).

Author Contributions: Conception and design: AJS, PWM, LBB; Acquisition of

data: AJS, AD, YN, VY, LBB; Analysis and interpretation of data: AJS, AD, YN,

PWM, LBB; Contributed unpublished, essential data, or reagents: CA, JET;

Drafting or revising the article: AJS, PWM, LBB.

Acknowledgements: We thank all members from the Barreiro and Messer lab for

helpful discussions and comments on the manuscript. We thank Dr Silvia Vidal for

the gift of the Influenza PR8 WT used in this study, Dr Hugo Soudeyns and Doris

Ransy for advice with the viral infections, and Marc Montero and George (PJ)

Perry for sharing their analysis on nucleotide diversity levels of OAS genes in

non-human primate species.

Funding Statement: This study was funded by grants from the Canadian

Institutes of Health Research (301538 and 232519), the Human Frontiers Science

Program (CDA-00025/2012) and the Canada Research Chairs Program (950-

228993) (to L.B.B.). Y.N. was supported by a fellowship from the *Réseau de*

Médecine Génétique Appliquée (RMGA).

References:

Awady, El, M.K., Anany, M.A., Esmat, G., Zayed, N., Tabll, A.A., Helmy, A.,
Zayady, El, A.R., Abdalla, M.S., Sharada, H.M., Raziky, El, M., Akel, El, W.,
Abdalla, S., Bader El Din, N.G., 2011. Single nucleotide polymorphism at
exon 7 splice acceptor site of OAS1 gene determines response of hepatitis C
virus patients to interferon therapy. Journal of Gastroenterology and
Hepatology. 26, 843–850.
Bigham, A.W., Buckingham, K.J., Husain, S., Emond, M.J., Bofferding, K.M.,
Gildersleeve, H., Rutherford, A., Astakhova, N.M., Perelygin, A.A., Busch,
M.P., Murray, K.O., Sejvar, J.J., Green, S., Kriesel, J., Brinton, M.A.,
Bamshad, M., 2011. Host Genetic Risk Factors for West Nile Virus Infection
and Disease Progression. PLoS ONE. 6, e24745.
Bonnevie-Nielsen, V., Field, L.L., Lu, S., Zheng, D.-J., Li, M., Martensen, P.M.,
Nielsen, T.B., Beck-Nielsen, H., Lau, Y.L., Pociot, F., 2005. Variation in
antiviral 2',5'-oligoadenylate synthetase (2"5"AS) enzyme activity is
controlled by a single-nucleotide polymorphism at a splice-acceptor site in the
OAS1 gene. The American Journal of Human Genetics. 76, 623–633.
Chen, G.K., Marjoram, P., Wall, J.D., 2009. Fast and flexible simulation of DNA

sequence data. Genome Research. 19, 136–142.

Consortium, T.1.G.P., 2015. A global reference for human genetic variation.
Nature. 526, 68–74.

Deschamps, M., Laval, G., Fagny, M., Itan, Y., Abel, L., Casanova, J.-L., Patin,
E., Quintana-Murci, L., 2016. Genomic Signatures of Selective Pressures and
Introgression from Archaic Hominins at Human Innate Immunity Genes.
American Journal of Human Genetics. 98, 5–21.

Ferrer-Admetlla, A., Liang, M., Korneliussen, T., Nielsen, R., 2014. On detecting
incomplete soft or hard selective sweeps using haplotype structure. Molecular
Biology and Evolution. 31, 1275–1291.

Fu, Q., Li, H., Moorjani, P., Jay, F., Slepchenko, S.M., Bondarev, A.A., Johnson,
P.L.F., Aximu-Petri, A., Prüfer, K., de Filippo, C., Meyer, M., Zwyns, N.,
Salazar-García, D.C., Kuzmin, Y.V., Keates, S.G., Kosintsev, P.A., Razhev,
D.I., Richards, M.P., Peristov, N.V., Lachmann, M., Douka, K., Higham,
T.F.G., Slatkin, M., Hublin, J.-J., Reich, D., Kelso, J., Viola, T.B., Pääbo, S.,
2014. Genome sequence of a 45,000-year-old modern human from western
Siberia. Nature. 514, 445–449.

Gravel, S., Henn, B.M., Gutenkunst, R.N., Indap, A.R., Marth, G.T., Clark, A.G.,
Yu, F., Gibbs, R.A., Bustamante, C.D., Project, 1.G., 2011. Demographic
history and rare allele sharing among human populations. Proceedings of the
National Academy of Sciences of the United States of America. 108, 11983–
11988.

Green, R.E., Krause, J., Briggs, A.W., Maricic, T., Stenzel, U., Kircher, M.,
Patterson, N., Li, H., Zhai, W., Fritz, M.H.-Y., 2010. A draft sequence of the
Neandertal genome. Science. 328, 710–722.

Gulig, P.A., Doyle, T.J., Clare-Salzler, M.J., Maiese, R.L., Matsui, H., 1997.
Systemic infection of mice by wild-type but not Spv- Salmonella typhimurium
is enhanced by neutralization of gamma interferon and tumor necrosis factor
alpha. Infection and Immunity. 65, 5191–5197.

Howie, B.N., Donnelly, P., Marchini, J., 2009. A Flexible and Accurate Genotype
Imputation Method for the Next Generation of Genome-Wide Association
Studies. PLoS Genetics. 5, e1000529.

Hublin, J.J., 2009. The origin of Neandertals. Proceedings of the National
Academy of Sciences. 106, 16022–16027.

Kent, W.J., Sugnet, C.W., Furey, T.S., Roskin, K.M., Pringle, T.H., Zahler, A.M.,
Haussler, D., 2002. The human genome browser at UCSC. Genome Research.
12, 996–1006.

Kwon, Y.-C., Kang, J.-I., Hwang, S.B., Ahn, B.-Y., 2012. The ribonuclease l-
dependent antiviral roles of human 2′,5′-oligoadenylate synthetase family
members against hepatitis C virus. FEBS letters. 587, 156–164.

LaRock, D.L., Chaudhary, A., Miller, S.I., 2015. Salmonellae interactions with
host processes. Nature Reviews Microbiology. 13, 191–205.

Lazaridis, I., Patterson, N., Mittnik, A., Renaud, G., Mallick, S., Kirsanow, K.,
Sudmant, P.H., Schraiber, J.G., Castellano, S., Lipson, M., Berger, B.,
Economou, C., Bollongino, R., Fu, Q., Bos, K.I., Nordenfelt, S., Li, H., de
Filippo, C., Prüfer, K., Sawyer, S., Posth, C., Haak, W., Hallgren, F.,

Fornander, E., Rohland, N., Delsate, D., Francken, M., Guinet, J.-M., Wahl, J.,
Ayodo, G., Babiker, H.A., Bailliet, G., Balanovska, E., Balanovsky, O.,
Barrantes, R., Bedoya, G., Ben-Ami, H., Bene, J., Berrada, F., Bravi, C.M.,
Brisighelli, F., Busby, G.B.J., Cali, F., Churnosov, M., Cole, D.E.C., Corach,
D., Damba, L., van Driem, G., Dryomov, S., Dugoujon, J.-M., Fedorova, S.A.,
Romero, I.G., Gubina, M., Hammer, M., Henn, B.M., Hervig, T., Hodoglugil,
U., Jha, A.R., Karachanak-Yankova, S., Khusainova, R., Khusnutdinova, E.,
Kittles, R., Kivisild, T., Klitz, W., Kucinskas, V., Kushniarevich, A., Laredj,
L., Litvinov, S., Loukidis, T., Mahley, R.W., Melegh, B., Metspalu, E.,
Molina, J., Mountain, J., Näkkäläjärvi, K., Nesheva, D., Nyambo, T., Osipova,
L., Parik, J., Platonov, F., Posukh, O., Romano, V., Rothhammer, F., Rudan,
I., Ruizbakiev, R., Sahakyan, H., Sajantila, A., Salas, A., Starikovskaya, E.B.,
Tarekegn, A., Toncheva, D., Turdikulova, S., Uktveryte, I., Utevska, O.,
Vasquez, R., Villena, M., Voevoda, M., Winkler, C.A., Yepiskoposyan, L.,
Zalloua, P., Zemunik, T., Cooper, A., Capelli, C., Thomas, M.G., Ruiz-
Linares, A., Tishkoff, S.A., Singh, L., Thangaraj, K., Villems, R., Comas, D.,
Sukernik, R., Metspalu, M., Meyer, M., Eichler, E.E., Burger, J., Slatkin, M.,
Pääbo, S., Kelso, J., Reich, D., Krause, J., 2014. Ancient human genomes
suggest three ancestral populations for present-day Europeans. Nature. 513,
409–413.
Lim, J.K., Lisco, A., McDermott, D.H., Huynh, L., Ward, J.M., Johnson, B.,
Johnson, H., Pape, J., Foster, G.A., Krysztof, D., Follmann, D., Stramer, S.L.,
Margolis, L.B., Murphy, P.M., 2009. Genetic Variation in OAS1 Is a Risk
Factor for Initial Infection with West Nile Virus in Man. PLoS Pathogens. 5,
e1000321.
Llorente, M.G., Jones, E.R., Eriksson, A., Siska, V., Arthur, K.W., Arthur, J.W.,
Curtis, M.C., Stock, J.T., Coltorti, M., Pieruccini, P., Stretton, S., Brock, F.,
Higham, T., Park, Y., Hofreiter, M., G, B.D., Bhak, J., Pinhasi, R., Manica,
A., 2015. Ancient Ethiopian genome reveals extensive Eurasian admixture
throughout the African continent. Science.
Mathieson, I., Lazaridis, I., Rohland, N., Mallick, S., Patterson, N., Roodenberg,
S.A., Harney, E., Stewardson, K., Fernandes, D., Novak, M., Sirak, K.,
Gamba, C., Jones, E.R., Llamas, B., Dryomov, S., Pickrell, J., Arsuaga, J.L.,
de Castro, J.M.B., Carbonell, E., Gerritsen, F., Khokhlov, A., Kuznetsov, P.,
Lozano, M., Meller, H., Mochalov, O., Moiseyev, V., Guerra, M.A.R.,
Roodenberg, J., Vergès, J.M., Krause, J., Cooper, A., Alt, K.W., Brown, D.,
Anthony, D., Lalueza-Fox, C., Haak, W., Pinhasi, R., Reich, D., 2015.
Genome-wide patterns of selection in 230 ancient Eurasians. Nature.
Mendez, F.L., Watkins, J.C., Hammer, M.F., 2012. A Haplotype at STAT2
Introgressed from Neanderthals and Serves as a Candidate of Positive
Selection in Papua New Guinea. The American Journal of Human Genetics.
91, 265–274.
Mendez, F.L., Watkins, J.C., Hammer, M.F., 2013. Neandertal origin of genetic
variation at the cluster of OAS immunity genes. Molecular Biology and
Evolution. 30, 798–801.
Meyer, M., Arsuaga, J.L., de Filippo, C., Nagel, S., Aximu-Petri, A., Nickel, B.,

Martínez, I., Gracia, A., de Castro, J.M.B., Carbonell, E., Viola, B., Kelso, J.,
Prüfer, K., Pääbo, S., 2016. Nuclear DNA sequences from the Middle
Pleistocene Sima de los Huesos hominins. Nature.
Meyer, M., Kircher, M., Gansauge, M.T., Li, H., Racimo, F., Mallick, S.,
Schraiber, J.G., Jay, F., Prufer, K., de Filippo, C., 2012. A high-coverage
genome sequence from an archaic denisovan individual. Science. 338, 222–
226.
Nauciel, C., Espinasse-Maes, F., 1992. Role of Gamma Interferon and Tumor-
Necrosis-Factor-Alpha in Resistance to Salmonella-Typhimurium Infection.
Infection and Immunity. 60, 450–454.
Player, M.R., Torrence, P.F., 1998. The 2–5 A system: Modulation of viral and
cellular processes through acceleration of RNA degradation. Pharmacology &
Therapeutics. 78, 55–113.
Prüfer, K., Racimo, F., Patterson, N., Jay, F., Sankararaman, S., Sawyer, S.,
Heinze, A., Renaud, G., Sudmant, P.H., de Filippo, C., Li, H., Mallick, S.,
Dannemann, M., Fu, Q., Kircher, M., Kuhlwilm, M., Lachmann, M., Meyer,
M., Ongyerth, M., Siebauer, M., Theunert, C., Tandon, A., Moorjani, P.,
Pickrell, J., Mullikin, J.C., Vohr, S.H., Green, R.E., Hellmann, I., Johnson,
P.L.F., Blanche, H., Cann, H., Kitzman, J.O., Shendure, J., Eichler, E.E., Lein,
E.S., Bakken, T.E., Golovanova, L.V., Doronichev, V.B., Shunkov, M.V.,
Derevianko, A.P., Viola, B., Slatkin, M., Reich, D., Kelso, J., Pääbo, S., 2014.
The complete genome sequence of a Neanderthal from the Altai Mountains.
Nature. 505, 43–49.
Racimo, F., Marnetto, D., Huerta-Sanchez, E., 2016. The landscape of uniquely
shared archaic alleles in present-day human populations. bioRxiv. 045237.
Racimo, F., Sankararaman, S., Nielsen, R., Huerta-Sanchez, E., 2015. Evidence
for archaic adaptive introgression in humans. Nature Reviews Genetics.
Reich, D., Green, R.E., Kircher, M., Krause, J., Patterson, N., Durand, E.Y., Viola,
B., Briggs, A.W., Stenzel, U., Johnson, P.L.F., 2010. Genetic history of an
archaic hominin group from Denisova Cave in Siberia. Nature. 468, 1053–
1060.
Sabeti, P.C., Reich, D.E., Higgins, J.M., Levine, H.Z.P., Richter, D.J., Schaffner,
S.F., Gabriel, S.B., Platko, J.V., Patterson, N.J., McDonald, G.J., 2002.
Detecting recent positive selection in the human genome from haplotype
structure. Nature. 419, 832–837.
Sankararaman, S., Mallick, S., Dannemann, M., Prüfer, K., Kelso, J., Pääbo, S.,
Patterson, N., Reich, D., 2014. The genomic landscape of Neanderthal
ancestry in present-day humans. Nature. 507, 354–357.
Sankararaman, S., Mallick, S., Patterson, N., Reich, D., 2016. The Combined
Landscape of Denisovan and Neanderthal Ancestry in Present-Day Humans.
Current Biology.
Sawyer, S., Renaud, G., Viola, B., Hublin, J.-J., Gansauge, M.-T., Shunkov, M.V.,
Derevianko, A.P., Prüfer, K., Kelso, J., Pääbo, S., 2015. Nuclear and
mitochondrial DNA sequences from two Denisovan individuals. Proceedings
of the National Academy of Sciences. 201519905.
Schlamp, F., Made, J., Stambler, R., Chesebrough, L., Boyko, A.R., Messer, P.W.,

2016. Evaluating the performance of selection scans to detect selective sweeps
in domestic dogs. Molecular Ecology. 25, 342–356.
Ségurel, L., Quintana-Murci, L., 2014. Preserving immune diversity through
ancient inheritance and admixture. Current opinion in immunology. 30C, 79–
84.
Tennessen, J.A., Bigham, A.W., O'Connor, T.D., Fu, W., Kenny, E.E., Gravel, S.,
McGee, S., Do, R., Liu, X., Jun, G., Kang, H.M., Jordan, D., Leal, S.M.,
Gabriel, S., Rieder, M.J., Abecasis, G., Altshuler, D., Nickerson, D.A.,
Boerwinkle, E., Sunyaev, S., Bustamante, C.D., Bamshad, M.J., Akey, J.M.,
Broad GO, Seattle GO, on behalf of the NHLBI Exome Sequencing Project,
2012. Evolution and Functional Impact of Rare Coding Variation from Deep
Sequencing of Human Exomes. Science. 337, 64–69.
Vernot, B., Akey, J.M., 2014. Resurrecting surviving Neandertal lineages from
modern human genomes. Science. 343, 1021.
Vernot, B., Akey, J.M., 2015. Complex History of Admixture between Modern
Humans and Neandertals. The American Journal of Human Genetics.
Voight, B.F., Kudaravalli, S., Wen, X., Pritchard, J.K., 2006. A map of recent
positive selection in the human genome. PLoS Biology. 4, e72.
Yang, M.A., Malaspinas, A.S., Durand, E.Y., Slatkin, M., 2012. Ancient structure
in Africa unlikely to explain Neanderthal and non-African genetic similarity.
Molecular Biology and Evolution. 29, 2987–2995.

805
806
Figure 1: Neandertal introgresssed haplotypes in the OAS region. (A) Neighbor-
joining tree of 5008 phased haplotypes spanning chr12: 113344739-113449528
(hg19) from phase 3 of the 1000 genomes project. Haplotypes condensed into 12
core haplotypes based on majority allele in clusters with pairwise differences of 85
or less. The figure illustrates that the Altai haplotype is very similar to "cluster 1"
haplotypes found in several human populations. Bootstrap values (1000 replicates)
are provided in blue boxes at each node. (B) Frequencies of the 12 core haplotypes
within each 1000 genomes project population sample. The most common
Neandertal-like haplotype, cluster 1, is found only outside of sub-Saharan African
samples, with the exception of recently admixed populations. Population codes
can be found at http://www.1000genomes.org/category/population/.
818
819
820
821

822
823
Figure 2: OAS-introgressed haplotypes show multiple signatures of positive
selection. (A) Comparison of frequency (y-axis) of NLS in the OAS locus in CEU
sample with respect to neutral expectations (dashed lines) based on coalescent
simulations. (B) The $H_{D/A}$ values associated with several NLS are significantly
larger (-log10 Ps in the y-axis) than expected under a neutral model of evolution.
$H_{D/A}$ at each NLS in the OAS region were compared against the distributions of
simulated NLS drawn to match local recombination rate and allele frequency. (C)
Absolute difference between observed and expected allele frequency in all four
present-day European samples (y-axis) based on ancient DNA data. Dashed lines
represent the 95th percentile of the expected distribution based on similar
deviations calculated on a dataset of approximately one million SNPs scattered
around the genome and with comparable present-day frequencies to those found
for NLS in the OAS region (D) Comparison of mean $H_{D/A}$ across OAS locus to
randomized frequency and genomic position-matched SNPs from simulations.
OAS locus is a strong outlier ($p < 10^{-5}$) compared to simulations. (E) Joint density
distribution of $H_{D/A}$ and Neandertal-like site derived frequency. Cluster of NLS in
OAS are significant outliers to neutral distribution ($p < 5 \times 10^{-4}$).

841
842
843
Figure 3: Pervasive impact of the Neandertal haplotype on the regulation of *OAS*
genes in primary macrophages. (A) –log 10 Ps (y-axis) for the association between
genotypes for SNPs with a MAF > 10% in the OAS region and expression levels
of *OAS3* in non-infected (black) and Salmonella-infected macrophages (orange).
The dashed line shows the P cutoff corresponding to an FDR of 10%. The right
panel shows a boxplot for the association between genotypes at the NLS
rs4767031 (x-axis) and the expression levels of *OAS3* (y-axis)(B) –log 10 Ps (y-
axis) for the association between genotypes in the OAS regions and the percentage
usage of isoform ENST00000202917 (i.e., p46 in the text) in non-infected (black)
and Salmonella-infected macrophages (orange). The dashed line shows the P
cutoff corresponding to an FDR of 1%. The right panel shows a boxplot for the
association between genotypes at the splicing variant rs10774671 (x-axis) and the
percentage usage of isoform p46 (y-axis). (C) Similar to (B) but for the percentage
usage of isoform ENST00000449768 of OAS2. In all the panels NLS are
highlighted by blue dots. The arrows on panel A-C highlight the location of the
SNPs for which the boxplots are shown on the right.

869
870
871
872
Figure 4: The Neandertal haplotype in the OAS regions has a different impact on
the regulation of OAS genes depending on the viral agents PBMCs are exposed to.
(A) Log 2 fold induction (y-axis) of *OAS1*, *OAS2* and *OAS3* in response to
different viral agents or viral-associated immune stimuli (B) –log10 P for the
association between genotype status for the Neandertal haplotype and overall
expression levels of OAS genes and the expression of specific isoforms of *OAS1*
and *OAS2* (those identified in Figure 3 as associated with NLS).
880
881
882

883
884 **Supplementary Figures:**
885

886
887 Supplementary Figure 1: Demographic model for neutral coalescent simulations.

896
897   Supplementary Figure 2: Illustration of $H_{D/A}$ statistic.
898
899
900
901

902
903

904
905
906 Supplementary Figure 3: Expression levels of OAS genes in primary macrophages
907 before and after infection with *Salmonella*.
908
909

910
911

912
913
Supplementary Figure 4: The splicing variant rs10774671 is a strong asQTL for
OAS1. Plotted is the normalized average coverage at which each base was
sequenced along the genomic regions encoding the gene *OAS1*. Individuals were
stratified according to their genotype at rs10774671. Below the figure are gene
models from Ensembl database. Individuals carrying the G allele at rs10444671
(i.e., the Neandertal allele) primarily express the transcript ENST00000202917
(referred to as p46 in the text) whereas individuals carrying the A derived allele
lose the splice site, which leads to the usage of a distinct isoform.
922
923

924
925

926
927 Supplementary Figure 5: Log 2 fold induction (y-axis) of *IFNA1* (type I
928 interferon) and *IFNG* (type II interferon) in PBMCs upon infection/stimulation
929 with several viral agents.

938
939
940
941 Supplementary Figure 6: $H_{D/A}$ does not vary substantially with SNP density.
942 $H_{D/A}$ statistic plotted against number of segregating sites (S) per locus across all
943 neutral coalescent simulations. The slight increase in $H_{D/A}$ with elevated S is
944 conservative with respect to our one-tailed test for elevated $H_{D/A}$.

Supplementary Tables:

	CEU	GBR	IBS	TSI	Mathieson et al. 2015 P
113341598	0.856	0.407	0.967	0.256	0.233
113343563	0.862	0.428	0.529	0.524	0.335
113350796	0.0123	0.0194	0.395	0.0557	0.0249
113360025	0.00850	0.0114	0.0317	0.00633	0.000277
113363284	0.0126	0.0170	0.0288	0.00732	0.0000288
113370966	0.00654	0.00902	0.0190	0.00476	2.29E-08
113371114	0.00735	0.0102	0.0209	0.00518	0.000000315
113375983	0.00947	0.0140	0.0333	0.00787	0.00459
113435450	0.0167	0.0284	0.132	0.0409	0.0000375
113504725	0.0852	0.0797	0.134	0.0797	0.157
113532885	0.354	0.853	0.0112	0.218	0.0236
p <= 0.01					
0.01<p<=0.05					

Supplementary Table 1: Proportion of similar frequency genome-wide SNPs with deviations from expected frequency as great or greater than 11 Neandertal-like sites. Fraction of all non-OAS SNPs for which absolute difference in expected (based on aDNA samples) and observed allele frequency that are as great or greater than NLS derived allele frequency difference. Final column reports P from Mathieson et al. 2015 selection test based on same aDNA samples.