Tumor evolution of glioma intrinsic gene expression subtype associates with immunological changes in the microenvironment

Qianghu Wang[1, 2, 3*], Xin Hu[1, 8*], Florian Muller[1, 4], Hoon Kim[1], Massimo Squatrito[9], Tom Mikkelsen[10], Lisa Scarpace[10], Floris Barthel[11], Yu-Hsi Lin[4], Nikunj Satani[4], Emmanuel Martinez-Ledesma[1], Edward Chang[1], Adriana Olar[5], Baoli Hu[6], Ana C. deCarvalho[10], Eskil Eskilsson[1], Siyuan Zheng[1], Amy B. Heimberger[7], Erik P. Sulman[2†], Do-Hyun Nam[12, 13, 14†], Roel G.W. Verhaak[1, 3†]

[1]Department of Genomic Medicine, [2]Department of Radiation Oncology, [3]Department of Bioinformatics and Computational Biology, [4]Department of Cancer Systems Imaging, [5]Department of Pathology, [6]Department of Cancer Biology, [7]Department of Neurosurgery, The University of Texas MD Anderson Cancer Center, Houston, TX 77030, USA; [8]University of Texas-Houston Graduate School in Biomedical Sciences, Houston, TX 77030, USA; [9]Cancer Cell Biology Programme, Seve Ballesteros Foundation Brain Tumor Group, Centro Nacional de Investigaciones Oncológicas, CNIO, 28029 Madrid, Spain; [10]Departments of Neurology and Neurosurgery, Henry Ford Hospital, Detroit, MI 48202, USA; [11]Oncology Graduate School Amsterdam, VU University Medical Center, 1081 HV Amsterdam, The Netherlands; [12]Institute for Refractory Cancer Research, Samsung Medical Center, Seoul 06351, Korea; [13]Department of Health Sciences and Technology, Samsung Advanced Institute for Health Sciences and Technology, Sungkyunkwan University, Seoul 06351, Korea; [14]Department of Neurosurgery Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, 135-710, Korea.

* These authors contributed equally
[†]Co-senior authors

Correspondence: rverhaak@mdanderson.org

Keywords: glioblastoma, disease recurrence, mesenchymal subtype, proneural to mesenchymal transition, gene expression profiling, tumor microenvironment, macrophages/microglia, immune cells.

ABSTRACT

We leveraged IDH wild type glioblastomas (GBM) and derivative neurosphere models to define the tumor-intrinsic mRNA transcription phenotype. We found that intratumoral heterogeneity is reflected in the transcriptome and associated with increased tumor microenvironment presence. We performed in silico cell sorting to demonstrate that M2 macrophages/microglia are the most frequent type of immune cells in the glioma microenvironment, followed by CD4+ T lymphocytes. Longitudinal transcriptome analysis of 124 pairs of primary and recurrent glioma pairs showed expression subtype is retained in 53% of cases with no proneural to mesenchymal transition being apparent. Inference of the tumor microenvironment through gene signatures revealed a decrease in invading monocytes but a subtype dependent increase in M2 macrophages/microglia cells after disease recurrence. Our longitudinal expression dataset can be accessed at http://ackbar.cnio.es:3838/RecuR/. Our study provides a comprehensive transcriptional and cellular landscape of IDH wild type GBM during treatment modulated tumor evolution.

INTRODUCTION

The intrinsic capacity of glioblastoma (GBM) tumor cells to infiltrate normal brain impedes surgical eradication and predictably results in high rates of early recurrence. To better understand determinants of GBM tumor evolution and treatment resistance, The Cancer Genome Atlas Consortium (TCGA) performed high dimensional profiling and molecular classification of nearly 600 GBM tumors [1-5]. In addition to revealing common mutations in genes such as *TP53*, *EGFR*, *IDH1*, and *PTEN*, as well as the frequent and concurrent presence of abnormalities in the p53, RB and receptor tyrosine kinase pathways. Unsupervised transcriptome analysis identified four clusters, referred to as classical, mesenchymal, neural and proneural, that were tightly associated with genomic abnormalities[6]. The proneural and the mesenchymal expression subtypes have been most consistently described in literature with proneural relating to a more favorable outcome and mesenchymal to unfavorable survival [7-9], but these findings were affected by the relatively favorable outcome of IDH-mutant glioblastoma which are consistently classified as proneural [4,6]. Proneural to mesenchymal switching upon disease recurrence has been described as a source for treatment resistance in GBM relapse [7,10-12], but the relevance of this phenomenon in glioma progress remains ambiguous.

GBM tumor cells along with the tumor microenvironment create a complex milieu that ultimately promotes tumor cell plasticity and disease progression [13]. The presence of tumor-associated stroma results in a mesenchymal tumor gene signature and poor prognosis in colon cancers [14]. Furthermore, the association between a mesenchymal gene expression signature and reduced tumor purity has been identified as a common

theme across cancer [15,16]. Tumor-associated macrophages/microglia in GBM have

been proposed as regulators of proneural-to-mesenchymal transition through NF-kB

activation [10] and may provide growth factor mediated proliferative signals, which could

be therapeutically targeted [17-19].

Here, we explored the properties of the microenvironment in different GBM gene

expression subtypes and characterized the transition between molecular subtypes

before and after therapeutic intervention. In doing so, we improved the robustness of

gene expression subtype classification through revised gene signatures and proposed

analytical methodology. Our results suggested that the tumor microenvironment

interferes with expression based classification of GBM, both at the primary disease

stage as well as at disease recurrence, and suggest a role for the

macrophage/microglia in treatment response.

RESULTS

Harnessing glioma sphere-forming cells identifies GBM specific intertumoral transcriptional heterogeneity

We set out to elucidate the tumor-intrinsic and tumor microenvironment independent transcriptional heterogeneity of GBMs. We performed a pairwise gene expression comparison of independent set of GBMs and the derivative glioma sphere-forming cells (GSCs) (n = 37) [20]. In total, 5,334 genes were found to be significantly higher expressed in parental GBMs relative to derived GSCs that could be attributed by the tumor associated GBM microenvironment (**Figure 1A**). To focus the analysis on the tumor-intrinsic transcriptome, these genes were filtered from further analysis. GBMs with IDH mutations have distinct biological properties and favorable clinical outcomes compared to IDH wild-type GBMs [1,4,5,21]. Using the filtered gene set, we performed consensus non-negative matrix factorization clustering to identify three distinct subgroups amongst 369 IDH wild type GBMs (**Figure 1B; Figure 1C**). When comparing the clustering result with the previously defined proneural (PN), neural (NE), classical (CL) and mesenchymal (MES) classification [1,3], three subgroups were strongly enriched for CL, MES and PN GBMs, respectively (**Figure S1**). Consequently, we labeled the groups as CL, MES and PN. None of the three subgroups was enriched for the NE class, suggesting its neural phenotype is non-tumor specific. The NE group has previously been related to the tumor margin where normal neural tissue is more likely to be present [22] and such contamination might explain why the neural subtype was the only subtype to lack characteristic gene abnormalities [1,23]. To be able to classify external GBM samples, we implemented a single sample gene set enrichment analysis

(ssGSEA) based equivalent distribution resampling classification strategy using 70-gene

signatures for each subgroup (**Table S1**)(**Figure 1D**), to assign each sample three

empirical classification p-values by which we determined the significantly activated

subtype(s) in the samples. We prepared an R-library to facilitate others to evaluate our

approach (**Supplementary File**). Using this method we found that the stability of cluster

assignments of 144 TCGA GBM samples profiled using both RNA sequencing and

Affymetrix U133A microarrays was 95% concordance (**Figure S2, Table S2**). This was

an improvement over the 77% subtype concordance determined using previously

reported methods [3]. We evaluated the distribution of somatic variants across all the

three molecular subtypes (**Figure 1E)(Figure S3)** and confirmed the strong

associations between subtypes and genomic abnormalities in previously reported driver

genes [1,3].

Multi-activation of subtype signatures associated with intratumoral heterogeneity

We observed that 34/369 (9.2%) samples showed significant enrichment of multiple

ssGSEA scores (empirical classification p-value<0.05), suggesting these cases activate

more than one transcriptional subtype (**Figure 2A**). To quantify this phenomenon, a

score ranging from 0 to 1 was defined to quantitatively evaluate the simplicity of subtype

activation based on order statistics of ssGSEA score. Samples with high simplicity

scores activated a single subtype and those with lowest simplicity scores activated

multiple subtypes. All multi-subtype TCGA samples showed simplicity scores of less

than 0.1 (**Figure 2A**). To determine whether transcriptional heterogeneity associated

with genomic intratumoral heterogeneity, we correlated simplicity scores, total mutation

rates and subclonal mutation rates. Included in the analysis were 224 TCGA GBMs with

available whole exome sequencing data [24] and ABSOLUTE [25] determined high tumor

purity (> 0.8) to equalize the mutation detection sensitivity [26]. Although not significant

(Wilcoxon rank test p-value=0.143), the total mutation rate was less in the bottom 30%

with lowest simplicity scores versus the top 30% samples with highest simplicity scores.

The subclonal mutation rate was significantly higher (p-value=0.024) in samples with

lowest simplicity scores (**Figure 2B; Table S3**), suggesting that increased intratumoral

heterogeneity associates with increased transcriptional heterogeneity.

We compared outcomes amongst the three transcriptional groups and observed

no significant differences (**Figure S4**). However, when restricting the analysis to

samples with high simplicity scores, a clear trend of MES showing worst survival and

PN the most favorable outcome became visible. For example, Kaplan-Meier analysis of

88 samples with simplicity scores >0.99 showed a median survival of 11.4, 14.7 and

16.7 months were detected in MES, CL and PN, respectively, which was significantly

different (log rank test, p=0.048)(**Figure 2C)(Figure S4)(Table S4**). Higher simplicity

scores correlated with relative favorable outcome within the PN set, non-significant in

the CL subtype, and correlated with relatively unfavorable survival in the MES class

(**Figure S5**).

Single GBM cell RNA sequencing recently suggested that GBMs are comprised

of a mixture of tumor cells with variable GBM subtype footprints [18]. We used this data to

classify 502 single GBM cells in addition to the bulk tumor derived from five primary

glioblastomas (**Table S5**). All bulk tumor samples showed simplicity scores less than

0.05 suggesting high transcriptional heterogeneity compared to 45 of 369 TCGA GBM

samples with simplicity scores below 0.05 (**Figure 2D**). In four of five cases (MGH26,

MGH28, MGH29 and MGH30), the bulk tumor samples were classified in the same

primary subtype as the majority of their single cells (**Figure 2D**). Our analysis suggests

that the heterogeneity observed at the single cell level is captured in the expression

profile of the bulk tumor, and that the five GBM samples studied at the single cell level

represented samples with relatively high transcriptional heterogeneity.

Transcriptional subtypes differentially activate the immune microenvironment

Despite restricting the cluster analysis to genes exclusively expressed by GBM cells, we

found that tumor purity predictions based on ABSOLUTE were significantly reduced in

GBM classified as MES (Student T-test p-value < 10e-14; **Figure 3A**). This was

corroborated by gene expression based predictions of tumor purity using the

ESTIMATE method (Student T-test p-value < 10e-32; **Figure 3B**)[15] . The ESTIMATE

method has been optimized to quantify tumor-associated fibroblasts and immune cells

[15] and the convergence of a decreased ABSOLUTE and decreased ESTIMATE tumor

purity confirmed previous suggestions on the increased presence of microglial and

neuroglial cells in mesenchymal GBM [12,27,28]. The mean simplicity score of samples

classified as MES was 0.53 which was significantly lower than in PN (Wilcoxon rank test

p-value<0.019) and CL subtypes (Wilcoxon rank test p-value<0.0001), confirming

increased transcriptional heterogeneity. In order to identify genomic determinants of

macrophage/microglia chemoattraction, we compared genomic alterations between

mesenchymal class samples with high (n=51) and low (n=51) ABSOLUTE based tumor

purity. GBM carrying hemyzygous loss of *NF1* or somatic mutations in *NF1* showed

reduced tumor purity compared to GBM with wild type *NF1* (Wilcoxon rank test p-

value=0.0007) and this association was similarly detected when limiting the analysis to

MES samples (Wilcoxon rank test p-value=0.017)(**Figure S6**). Formation of dermal

neurofibromas in the context of *Nf1* loss of heterozygosity has been reported to be

context and microenvironment dependent [29]. Functional studies may clarify whether

NF1 deficient GBM are able to recruit cells that provide them with a proliferative

advantage, or whether *NF1* loss provides that proliferative advantage in a specific

tumor-associated microenvironment context.

To determine the cellular components of the tumor microenvironment across

different transcriptional subtypes, we used the CIBERSORT in silico cytometry [30]

method to evaluate absolute immune cell fractions. We evaluated 22 different immune

cell types in 69 PN, 137 CL and 96 MES samples, after filtering samples with

classification simplicity scores less than 0.1 (**Table S6**). Microglia are the resident

macrophages in the central nervous system. Peripheral blood monocytes also give rise

to tumor associated macrophages. These innate immune cells can be broadly classified

as the proinflammatory M1 type and the alternative tumor promoting M2 type[31]. The M2

macrophage gene signature showed a greater association with the MES subtype

(13.4%) relative to the PN (4.6%) and CL (6.0%)(**Figure 3C**), consistent with previous

analysis of the TCGA database[28,32]. In addition to the M2 macrophage gene signature,

there was also a significantly higher fraction of MES samples that expressed M1

macrophage (Student T-test p-value 3.20E-5) and neutrophil (Student T-test p-value

1.30E-9) gene signatures. In contrast, the activated natural killer T-cell gene signature

(Student T-test p-value 4.91E-2) was significantly reduced in the MES subtype and

resting memory CD4+ T cells (Student T-test p-value 5.40E-7) were less frequently

expressed in the PN subtype.

To confirm the association of macrophages/microglia with the MES GBM subtype,

we assessed protein expression levels of the ITGAM (alternatively known as CD11B)

and IBA1 (also known as AIF1) macrophage/microglial markers in a set of 18 GBM for

which we also characterized the expression subtype (**Figure 3D**) as well as through

immunohistochemistry (**Figure S7**). We confirmed the microenvironment as the main

source for *ITGAM/IBA1* transcription by comparing transcriptional levels in 37 GBM-

neurosphere pairs used for gene filtering, which showed that neurospheres do not

express *ITGAM/IBA1* (**Figure 3E**). The association of the MES GBM subtype with

increased level of M2 microglia/macrophages may suggest that in particular MES GBM

are candidates for therapies directed against tumor-associated macrophages[17].

Activated dendritic cell signatures (Student T-test p-value 7.36E-3)(**Figure 3C**) were

significantly higher in the CL subtype, suggesting this subtype may benefit from

dendritic cell vaccines [33]. Dendritic cells may require an activated phenotype in order to

direct the immune system. A previous study suggested that MES GBM patients treated

with dendritic cells were more likely to benefit [34].

Phenotypic plasticity upon GBM recurrence

Glioblastoma has long been hypothesized to progress along a proneural to

mesenchymal axis[7]. To determine the relevance of this transition process in IDH

wildtype glioma evolution, we performed a longitudinal analysis of the subtype

classification and tumor-associated microenvironment in sample pairs obtained at

diagnosis and first disease recurrence from 124 glioma patients. The cohort included 96

initial GBM and first recurrence, eight pairs of primary low grade glioma and matching

secondary GBM, and 20 pairs of primary and recurrent low grade glioma. Gene

expression profiles of 78 tumor pairs were analyzed through transcriptome sequencing,

and remaining pairs were generated using Affymetrix (n = 31) and Illumina (n = 15)

microarray, respectively. To facilitate exploration of this dataset we have made it

available through the GlioVis portal http://ackbar.cnio.es:3838/RecuR/. We used a gene

expression signature[35] to determine that 33 of 124 cases were IDH-mutant/GCIMP at

presentation and recurrence (**Table S7**). We used the renewed gene signatures and

classification method to determine molecular subtype of the 91 pairs of IDH wild type

cases and found that expression class remained consistent after disease recurrence for

48 of 91 IDH-wildtype cases (52%)(**Figure 4A**). The MES subtype was most stable

(64%) while the CL (47%) and PN (43%) phenotypes were less frequently retained.

Nine, sixteen and eighteen post-treatment tumors switched subtypes to become CL,

MES and PN at disease recurrence, respectively, indicating that PN and MES increased

in frequency after recurrence while the CL subtype was least frequently found (**Figure

4A**). The CL expression class was previously found to be most sensitive to intensive

therapy and it is possible that therapy provides a competitive advantage for non-CL

cells, which could explain the reduced post-treatment incidence[3]. Our results did not

identify enrichment for proneural to mesenchymal transitions.

We observed a significant difference in transcriptional simplicity between primary

GBM retaining their expression class, versus those that switched to a different

phenotype (**Figure 4B**). GBMs with a primary tumor simplicity score greater than 0.5,

242 indicated lower transcriptional heterogeneity, were classified as the same subtype in 31

243 of 48 (64.5%) cases, compared to 15 of 41 (36.6%) cases with primary tumor simplicity

244 scores less than 0.5 (Fisher exact test p-value=0.01).

Microenvironment transitions upon GBM recurrence

Debulking surgery, radiotherapy and chemotherapy provide therapeutic barriers but

nonetheless induce tumor evolution, including influences on the tumor

microenvironment. We explored this possibility by comparing the tumor associated

microenvironment in primary and recurrent GBMs using CIBERSORT (**Table S8**)[30]. A

comparison between 91 primary and recurrent IDH-wild type tumors revealed a

decrease in monocyte gene signature expression at recurrence, suggesting relative

depletion of circulation derived monocytes (**Figure 5A**). Next, we dissected

microenvironment fluctuations between diagnosis and recurrent tumors across different

subtype combinations. Primary non-MES (CL or PN) tumors showed relatively high

tumor purity and consequently, recurrent tumors classified as non-MES demonstrated a

relatively global decrease of immune cells while cases transitioning to MES at

recurrence represented a trend towards increased immune cell fractions (**Figure 5B**).

Gene signatures of immunosuppressive regulatory T cells showed an increase in gene

expression at recurrence across several primary-recurrence subtype combinations

although the inferred cellular fractions are relatively small (**Figure 5B**). In contrast to the

trend of monocyte depletion, the imputed M2 macrophage frequency was significantly

higher at recurrence in cases transitioning to MES (**Figure 5C**). This observation

converges with the higher fraction of M2 macrophages in primary MES GBM relative to

primary non-MES GBM. M1 macrophages and neutrophils also correlated with primary MES GBM, but these associations were not confirmed for recurrent GBM. We validated the increase in macrophages using immunostaining of IBA1 expression in two primary-recurrent GBM pairs which were classified as CL to MES (**Figure 5D**). IBA1 immunoexpression was restricted to macrophages/microglia, cells exhibiting either globular or filamentous/spidery morphology, with no expression in glioma tumor cells (**Figure 5D**). These findings further solidify the association between MES GBM and macrophage/microglia and extend this mutual relationship to disease recurrence. MES tumors at recurrence compared to primary MES tumors showed an increase in transcriptional activity associated with non-polarized M0 macrophages, which has been previously described [28], but also dendritic cells which is potentially motivated by the increased levels of neoantigens at disease recurrence [24]. In contrast, primary PN GBM were found to contain significantly higher fractions of five immune cell categories compared to recurrent PN GBM, indicating a relative absence of immune infiltration in PN GBM upon recurrence.

We evaluated the effect of transcriptional class on patient survival. The analysis was restricted to 50 cases for whom annotation on overall survival (OS) time and time to disease progression (PFS) were available and with high simplicity scores, indicating low transcriptional heterogeneity. We confirmed the worse prognosis for patients whose primary tumor was classified as MES on overall survival (logrank test p=0.029 with HR=1.97)(**Figure 6A; Figure 6B**). This pattern was retained in patients whose secondary glioma was classified as MES (logrank test p=0.09 with HR=1.71)(**Figure 6C; Figure 6D**). Consequently, cases for whom both primary and recurrent tumor were

288   classified as MES subtype showed the least favorable outcome, suggesting an additive

289   effect of transcriptional class at different time points (**Figure 6E, Figure 6F**)(**Figure S8**).

DISCUSSION

Transcriptome profiling of tumor samples is a commonly used modality for interrogating pathway functionality and phenotype based patient classification. The transcriptional footprint left by the tumor microenvironment, which may constitute 10-80% of cells in a tumor biopsy[15], can obscure the true activity of the signaling network[14,36]. Here, we employed in silico methods to integrate mRNA expression profiles from glioma samples and glioma cell culture models to provide insights into glioma-intrinsic pathway activities and classification, and to deconvolute the glioma associated stroma into its immunological cellular components.

GBM expression subtype classification has emerged as an important concept to better understand the biology of this devastating disease [8,37,38]. Robust classification of new GBM tumors is therefore critical to ensure consistency in reporting between different studies. The transcriptional glioma subtypes we discovered using tumor-intrinsic gene expression values strongly overlapped with the proneural, classical and mesenchymal subtypes but identified the neural class as normal neural lineage contamination. Our updated methods, released through a R-library, were found to be highly robust and provide the community with a standardized strategy for classification of gliomas.

Through re-classification of primary GBM samples from TCGA and despite using tumor-only transcripts, we observed that the mesenchymal GBM subtype associated with the presence of tumor-associated glial and microglial cells. Mesenchymal glioma cell differentiation status has been found to correlate with enrichment of macrophages/microglia [10,39]. Through in silico cell type identification we additionally

detected enrichment of various adaptive immunity cell types, including CD4+ T lymphocytes.

Longitudinal analysis of tumor samples is complicated by the lack of tissue collections including such pairs. Through aggregation of existing and novel datasets we compiled a cohort of 124 glioma pairs, including 91 pairs of IDH wild type tumors. Comparison of pairs of initial gliomas and first disease recurrence did not identify the trend of proneural GBM transitioning to a mesenchymal phenotype that has often been suspected [7]. Mesenchymal subtype at diagnosis and at disease recurrent correlated with relatively poor outcome. The recurrent IDH wild type GBM immuno environment showed fewer blood derived monocytes which may reflect lower penetration through the blood brain barrier. While the frequency of M2 macrophage/microglia was increased in recurrent mesenchymal GBM compared to primary non-mesenchymal GBM, the overall fraction of M2 macropage/microglia remained stable. This possibly suggests that the majority of these cells are derived from resident CNS macrophages than through active recruitment from the circulation.

In summary, our study defines a new strategy to determine transcriptional subtype, and associated expression classes to the tumor-associated immuno-environment. Our findings may aid in the implementation of immunotherapy approaches [40] in a disease type with very limited treatment options. Collectively, our results have improved our understanding of determinants of GBM subtype classification, the critical impact of the tumor microenvironment, and provide new handles on the interpretation of transcriptional profiling of glioma.

Acknowledgments

The authors thank Katherine Stemke-Hale for assistance in manuscript editing. The results published here are in part based upon data generated by The Cancer Genome Atlas project established by the National Cancer Institute (NCI) and the National Human Genome Research Institute (NHGRI) of the National Institutes of Health. Information about TCGA and the investigators and institutions that constitute the TCGA research network can be found at http://cancergenome.nih.gov. This work is supported by grants from the National Institutes of Health grants P50 CA127001 (EPS, RGWV), R01 CA190121 (EPS, RGWV), P01 CA085878 (RGWV); RO1 CA120813 (ABH), and P30 CA016672 (MD Anderson Cancer Center Support Grant for the Sequencing and Microarray Facility); Cancer Prevention & Research Institute of Texas (CPRIT) grant number R140606, the University Cancer Foundation via the Institutional Research Grant program at the University of Texas MD Anderson Cancer Center (RGWV); the National Brain Tumor Association Defeat GBM project (EPS, RGWV), the National Brain Tumor Association Oligo Research Fund (RGWV).

References

Brennan, C. W. *et al.* The somatic genomic landscape of glioblastoma. *Cell* **155**, 462-477, doi:10.1016/j.cell.2013.09.034 (2013).

Cancer Genome Atlas Research, N. Comprehensive genomic characterization defines human glioblastoma genes and core pathways. *Nature* **455**, 1061-1068, doi:10.1038/nature07385 (2008).

Verhaak, R. G. *et al.* Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1. *Cancer cell* **17**, 98-110, doi:10.1016/j.ccr.2009.12.020 (2010).

Noushmehr, H. *et al.* Identification of a CpG island methylator phenotype that defines a distinct subgroup of glioma. *Cancer Cell* **17**, 510-522, doi:10.1016/j.ccr.2010.03.017 (2010).

Ceccarelli, M. *et al.* Molecular Profiling Reveals Biologically Discrete Subsets and Pathways of Progression in Diffuse Glioma. *Cell* **164**, 550-563, doi:10.1016/j.cell.2015.12.028 (2016).

Verhaak, C. M., Lintsen, A. M., Evers, A. W. & Braat, D. D. Who is at risk of emotional problems and how do you know? Screening of women going for IVF treatment. *Hum Reprod* **25**, 1234-1240, doi:10.1093/humrep/deq054 (2010).

Phillips, H. S. *et al.* Molecular subclasses of high-grade glioma predict prognosis, delineate a pattern of disease progression, and resemble stages in neurogenesis. *Cancer cell* **9**, 157-173, doi:10.1016/j.ccr.2006.02.019 (2006).

Huse, J. T., Phillips, H. S. & Brennan, C. W. Molecular subclassification of diffuse gliomas: seeing order in the chaos. *Glia* **59**, 1190-1199, doi:10.1002/glia.21165 (2011).

Zheng, S., Chheda, M. G. & Verhaak, R. G. Studying a complex tumor: potential and pitfalls. *Cancer J* **18**, 107-114, doi:10.1097/PPO.0b013e3182431c57 (2012).

Bhat, K. P. *et al.* Mesenchymal differentiation mediated by NF-kappaB promotes radiation resistance in glioblastoma. *Cancer cell* **24**, 331-346, doi:10.1016/j.ccr.2013.08.001 (2013).

Ozawa, T. *et al.* Most human non-GCIMP glioblastoma subtypes evolve from a common proneural-like precursor glioma. *Cancer Cell* **26**, 288-300, doi:10.1016/j.ccr.2014.06.005 (2014).

Bao, S. *et al.* Glioma stem cells promote radioresistance by preferential activation of the DNA damage response. *Nature* **444**, 756-760, doi:10.1038/nature05236 (2006).

Olar, A. & Aldape, K. D. Using the molecular classification of glioblastoma to inform personalized treatment. *The Journal of pathology* **232**, 165-177, doi:10.1002/path.4282 (2014).

Isella, C. *et al.* Stromal contribution to the colorectal cancer transcriptome. *Nat Genet* **47**, 312-319, doi:10.1038/ng.3224 (2015).

Yoshihara, K. *et al.* Inferring tumour purity and stromal and immune cell admixture from expression data. *Nat Commun* **4**, 2612, doi:10.1038/ncomms3612 (2013).

Martinez, E. *et al.* Comparison of gene expression patterns across 12 tumor types identifies a cancer supercluster characterized by TP53 mutations and cell cycle defects. *Oncogene* **34**, 2732-2740, doi:10.1038/onc.2014.216 (2015).

Pyonteck, S. M. *et al.* CSF-1R inhibition alters macrophage polarization and blocks glioma progression. *Nat Med* **19**, 1264-1272, doi:10.1038/nm.3337 (2013).

Patel, A. P. *et al.* Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma. *Science* **344**, 1396-1401, doi:10.1126/science.1254257 (2014).

Yan, J. *et al.* FGL2 as a Multimodality Regulator of Tumor-Mediated Immune Suppression and Therapeutic Target in Gliomas. *J Natl Cancer Inst* **107**, doi:10.1093/jnci/djv137 (2015).

Galli, R. *et al.* Isolation and characterization of tumorigenic, stem-like neural precursors from human glioblastoma. *Cancer Res* **64**, 7011-7021, doi:10.1158/0008-5472.CAN-04-1364 (2004).

Cancer Genome Atlas Research, N. *et al.* Comprehensive, Integrative Genomic Analysis of Diffuse Lower-Grade Gliomas. *N Engl J Med* **372**, 2481-2498, doi:10.1056/NEJMoa1402121 (2015).

Gill, B. J. *et al.* MRI-localized biopsies reveal subtype-specific differences in molecular and cellular composition at the margins of glioblastoma. *Proc Natl Acad Sci U S A* **111**, 12550-12555, doi:10.1073/pnas.1405839111 (2014).

23   Verhaak, R. G. *et al.* Prognostically relevant gene signatures of high-grade serous ovarian
carcinoma. *J Clin Invest* **123**, 517-525, doi:65833/10.1172/JCI65833 (2013).
24   Kim, H. *et al.* Whole-genome and multisector exome sequencing of primary and post-treatment
glioblastoma reveals patterns of tumor evolution. *Genome Res* **25**, 316-327,
doi:10.1101/gr.180612.114 (2015).
25   Carter, S. L. *et al.* Absolute quantification of somatic DNA alterations in human cancer. *Nat*
Biotechnol **30**, 413-421, doi:10.1038/nbt.2203 (2012).
26   Aran, D., Sirota, M. & Butte, A. J. Systematic pan-cancer analysis of tumour purity. *Nat Commun*
6, 8971, doi:10.1038/ncomms9971 (2015).
27   Ye, X. Z. *et al.* Tumor-associated microglia/macrophages enhance the invasion of glioma stem-
like cells via TGF-beta1 signaling pathway. *J Immunol* **189**, 444-453,
doi:10.4049/jimmunol.1103248 (2012).
28   Gabrusiewicz, K. *et al.* Glioblastoma-infiltrated innate immune cells resemble M0 macrophage
phenotype. *JCI insight* **1**, doi:10.1172/jci.insight.85841 (2016).
29   Le, L. Q., Shipman, T., Burns, D. K. & Parada, L. F. Cell of origin and microenvironment
contribution for NF1-associated dermal neurofibromas. *Cell Stem Cell* **4**, 453-463,
doi:10.1016/j.stem.2009.03.017 (2009).
30   Newman, A. M. *et al.* Robust enumeration of cell subsets from tissue expression profiles. *Nat*
Methods **12**, 453-457, doi:10.1038/nmeth.3337 (2015).
31   Hambardzumyan, D., Gutmann, D. H. & Kettenmann, H. The role of microglia and macrophages
in glioma maintenance and progression. *Nature neuroscience* **19**, 20-27, doi:10.1038/nn.4185
(2015).
32   Doucette, T. *et al.* Immune heterogeneity of glioblastoma subtypes: extrapolation from the
cancer genome atlas. *Cancer Immunol Res* **1**, 112-122, doi:10.1158/2326-6066.CIR-13-0028
(2013).
33   Palucka, K. & Banchereau, J. Cancer immunotherapy via dendritic cells. *Nature reviews. Cancer*
12, 265-277, doi:10.1038/nrc3258 (2012).
34   Prins, R. M. *et al.* Gene expression profile correlates with T-cell infiltration and relative survival
in glioblastoma patients vaccinated with dendritic cell immunotherapy. *Clin Cancer Res* **17**,
1603-1615, doi:10.1158/1078-0432.CCR-10-2563 (2011).
35   Baysan, M. *et al.* G-cimp status prediction of glioblastoma samples using mRNA expression data.
PloS one **7**, e47839, doi:10.1371/journal.pone.0047839 (2012).
36   Kim, H. & Verhaak, R. G. Transcriptional mimicry by tumor-associated stroma. *Nat Genet* **47**,
307-309, doi:10.1038/ng.3255 (2015).
37   Dunn, G. P. *et al.* Emerging insights into the molecular and cellular basis of glioblastoma. *Genes*
Dev **26**, 756-784, doi:10.1101/gad.187922.112 (2012).
38   Sturm, D. *et al.* Paediatric and adult glioblastoma: multiform (epi)genomic culprits emerge. *Nat*
Rev Cancer **14**, 92-107, doi:10.1038/nrc3655 (2014).
39   Kreutzberg, G. W. Microglia: a sensor for pathological events in the CNS. *Trends Neurosci* **19**,
312-318 (1996).
40   Blank, C. U., Haanen, J. B., Ribas, A. & Schumacher, T. N. CANCER IMMUNOLOGY. The "cancer
immunogram". *Science* **352**, 658-660, doi:10.1126/science.aaf2834 (2016).

Figures and Legends

Figure 1. Molecular classification of IDHwt GBMs. (A) Filtering tumor associated microenvironment genes. (B) Discarding IDH mutation related GBMs. (C) Overview of NMF clustering. (D) Heatmap of 70-gene signatures by gene expression subtype were developed based on 270 GBMs. Top ten genes are shown for each subtype. (E) Frequency of subtype related somatic genomic alterations.

451
Figure 2. Multi activation of transcriptional subtypes associated with intratumoral
heterogeneity. (**A**) The expression profiles of 369 IDHwt GBMs were analyzed using Affymetrix
U133A. The empirical -log(P-value) of raw ssGSEA enrichment scores at each signature are
shown as heatmaps, with dark blue representing no activation and bright red as highly activated.
Yellow star indicates the secondary activated subtype (empirical p-value<0.05). For each panel,
the first row shows simplicity score, and the second row indicates transcriptional subtype. (**B**)
Comparison of mutation rate, subclonal mutation rate and subclonal mutation fraction between
IDHwt GBMs with high and low simplicity scores. P-values were calculated using Wilcoxon rank
test and shown at the top of each panel. (**C**) Kaplan-Meier survival curver by subtype. (**D**)
Transcriptome classification of five bulk tumor samples and 502 single GBM cells derived from
them. The top two row of each panel show the dominant and secondary subtype of the GBM
tumor bulk. The heatmap of each panel shows the empirical -log(P-value) of the ssGSEA
scores of the derived single GBM cells on each of the three subtype signatures. The bottom row
shows the subtype distribution of derived single GBM cells within the same GBM tumor of origin.

Figure 3. Transcriptional subtypes differentially activate the immune microenvironment. (**A, B**) Tumor purity of 364 respectively 369 TCGA IDHwt GBM samples was determined by ABSOLUTE and ESTIMATE. The difference in tumor purity between subtypes was evaluated using a two-sample heteroscedastic t-test. (**C**) Comparison of immune cell fraction among subtypes. Purple, skyblue and green boxplots indicate PN, CL and MES subtype, respectively. Immune cell fraction was estimated using CIBERSORT and corrected using ABSOLUTE purity scores. Difference of cell fraction between subtypes was evaluated using Mood's test. (**D**) The upper panel shows ssGSEA enrichment scores and associated expression subtype classifications. Bottom panels display protein expression of the microglial markers integrin alpha M (ITGAM) and allograft inflammatory factor 1 (IBA1), astrocyte marker glial fibrillary acidic protein (GFAP) and the loading control tubulin. (**E**) Comparison of *ITGAM* and *IBA1* gene expression levels between GBM and derived neurosphere models.

480
481 **Figure 4. Comparison between transcriptional subtype of primary and paired recurrent**
482 **tumors.** (**A**) Rows and columns of the cross table represents subtype distribution frequency of
483 primary and paired recurrent tumors, respectively. (**B**) Violin plots show the distribution of
484 simplicity scores of pairs with (left) and without (right) subtype transition.

Figure 5. Microenvironment transition between primary and paired recurrent tumors. (**A**) Red and blue boxplots represent the immune cell fraction distribution of each immune cell type. Immune cell fraction was calculated using CIBERSORT and adjusted using ESTIMATE purity scores. Difference between cell fraction of primary and paired recurrent tumors was calculated using Wilcoxon rank test. (**B**) The blue-to-red heatmap represents immune cell fraction changes upon tumor recurrence per subtype transitions which were list on the left of the heatmap. (**C**) Each dot represents a pair of primary and recurrent GBM with axes indicating M2 macrophage cell fraction. (**D**) Immunohistochemical staining of the IBA1 microglia marker in two matched pairs of primary and recurrent GBM. Cellular morphology in combination with the distribution of the staining suggested exclusive microglial expression of monocyte lineage markers IBA1 in GBM tumors.

497
498 **Figure 6. Survival analysis of paired IDH wild type GBM.** (A, B) OS and PFS analyses
499 between samples with different primary subtype. (C) Difference of survival time after secondary
500 surgery between patients with non-MES and MES in primary tumors. (D) Survival analysis of
501 time after secondary surgery between patients with non-MES and MES in recurrent tumors. (E,
502 F) OS and PFS analyses between samples with difference recurrent subtype.

Supplementary Information

Supplementary Figures and Legends

	Proneural	Neural	Classical	Mesenchymal
Class1	0.0%	17.5%	**74.2%**	8.3%
Class2	2.1%	7.5%	5.3%	**85.1%**
Class3	**73.4%**	22.8%	1.3%	2.5%

Figure S1. Comparison between GCIMP- GBM specific classification and TCGA defined GBM subtypes.

Figure S2. Concordance of transcriptional classification of GBMs cross multiple platforms. Through TCGA, the expression profiles of 144 GBM were analyzed using both Affymetrix U133A gene expression arrays and RNA sequencing. The empirical -log(P-value) of raw ssGSEA enrich- ment scores at each signature are shown as heatmaps, with dark blue representing no activation and bright red as highly activated. For each panel, the first row shows U133A based classification, and the second row indicates RNA-seq subtype classification.

Figure S3. Genomic alteration patterns for each subtype. Ten prominent GBM somatic events were manually selected. Simplicity score, primary and secondary subtype were shown by the first three rows.

Figure S4. Patient survival differences between transcriptional subtypes. Samples were
filtered using increased simplicity score as threshold from panel A to D.

Figure S5. Overall and event free survival analysis comparison between samples with high and low simplicity scores in each subtype.

Figure S6. Comparison of tumor purity and immune cell fraction between GBMs with different *NF1* genomic status. P-values were calculated using Wilcoxon rank test between samples carrying *NF1* deletion/mutation and others.

Figure S7. Immunohistochemistry staining of the IBA1 imcroglia marker.

Panel A and B:
Recurrent tumor in
■ N.MES (N=23)
■ MES (N=27)

Panel C:
Pri. -->Rec.
■ N.MES-->N.MES (N=19)   ■ MES-->N.MES (N=4)
■ N.MES-->MES (N=10)   ■ MES-->MES (N=17)

Figure S8 .

Figure S8. (A, B) Overall and progression free survival analysis between samples in different recurrent subtypes. **(C)** Survival after secondary surgery comparison between different transition types.

Supplementary Methods and Experimental Procedures

Collection of pairs of primary and recurrent glioma samples

U133A array profiles for 543 primary GBM, and RNA-Seq data for 166 primary and 13

recurrent GBM were obtained from the TCGA portal https://tcga-data.nci.nih.gov/tcga/.

Mutation calls and DNA copy number profiles were obtained for all samples, where

available. Tissues from 20 additional initial GBM and matched recurrent tumor were

obtained from Henry Ford Hospital (n = 9) in accordance with institutional policies and

all patients provided written consent, with approval from the Institutional Review Boards

(Henry Ford Hospital IRB protocol #402). All RNA samples tested were obtained from

frozen specimens. All of the recurrent GBMs had been previously treated with

chemotherapy and radiation. Three cases had a history of lower grade astrocytoma

prior to the first GBM (HF-2869/HF-3081/HF-3162). Tumors were selected solely on the

basis of availability. RNA-Seq libraries were generated using RNA Truseq reagents

(Illumina, San Diego, CA, USA) and paired-end sequenced using standard Illumina

protocols. Read length was 76 base pairs for cases sequenced by TCGA and from

Henry Ford (processed at MD Anderson). RNA-Seq data on frozen tissue from 44

patients with initial and recurrent GBM that received resection at Samsung Medical

Center and Seoul National University Hospital were provided by Dr. Nam's lab. Surgery

specimens were obtained in accordance to the Institutional Review Board (IRB) of the

Samsung Medical Center (No. 2010-04-004) and Seoul National University Hospital (No.

C-1404-056-572). Affymetrix CEL files of 39 pairs of initial and recurrent glioma were

retrieved from the Gene Expression Omnibus (GEO accession GSE4271, GSE42670,

GSE62153)[1-3]. The expression profiles of the 23 pairs from GSE4271 were determined

using Affymetrix HG-U133 GeneChips, the 1 pairs from GSE42670 were analyzed using

the Affmetrix HuGene-1-0-st platform, the 15 pairs from GSE62153 were analyzed

using Illumina Human HT-12 V4.0 expression BeadChip.

Genome wide DNA copy number profiling and exome sequencing on thirteen

TCGA tumor pairs and nine of ten Henry Ford tumor pairs were performed and data

was analyzed using standard protocols and pipelines as previously described [4].

Data for multiplatform classification comparison

RNA sequencing data was available for 162 primary GBMs[5] for which an Affymetrix HT-

U133A gene expression profile was also available. We observed a low Pearson

Correlation Coefficient (< 0.15) between RNA sequencing based reads per kilo base of

transcript per million reads (RPKM) and Affymetrix HT-U133A profiles in eighteen cases

and these were removed from further analysis. In summary, in order to assess the

concordance between classification results of the new 70-gene signatures and

previously published 210-gene signatures [6], 144 GBMs which were profiled in both RNA

sequencing and Affymetrix U133A platforms were used in our further analyses.

Transcriptome Data processing

The latest version custom CDF files (Version19, http://brainarray.mbni.med.umich.edu)

[7,8] were used to map probes from the Affymetrix HG-U133A and HuGene-1_0-st

GeneChip platforms to the Ensemble transcript database, combined in one probe set

per gene and normalized using the AROMA package with default parameters, resulting

in RMA normalized and log transformed gene expression values [9]. All RNA sequencing

data was processed by the PRADA pipeline [10]. Briefly, reads were aligned using BWA

against the genome and transcriptome. After initial mapping, the aligned reads were

filtered out if their best placements are only mapped to unique genomic coordinates.

Quality scores are recalibrated using the Genome Analysis Toolkit (GATK), and

duplicate reads are flagged using Picard. Mapped features were quantified and

normalized per kilo base of transcript per million reads (RPKM) and were converted to a

log2 scale to represent a gene expression level. RPKM values measuring the same

gene that mapped to the Ensemble transcript with longest size were selected to obtain

one expression value per gene and sample. RPKM values were converted to a log2

scale to represent gene expression level. The statistical environment R was used to

perform all the statistical analysis and graph plots.

Deriving new gene signatures

A pair-wise gene expression analysis identified 5,334 genes which are significant higher

expressed in glioma bulk samples compared to their derivative GSCs. These genes

were discarded from the gene list for developing tumor-specific molecular subtypes.

Consensus non-negative matrix factorization (CNMF) clustering method identified three

distinct subgroups among the 369 IDHwt primary GBMs. A set of 270 GBMs was

recognized as core samples based on a positive silhouette width. The gene expression

values of each subtype were compared with those from the other two subtypes

combined [6]. Signature genes per cluster were selected on the basis of differences in

gene expression level and were considered significant if they reached the cut-off value

with t-test p-value< 1E-3 for higher expressed in this class, while also showing a

significant lower expression with t-test p-value<1E-3 in the other two classes. In the

original gene signatures, genes could be either down-regulated or up-regulated, while

only up-regulated genes (n=70 per gene signature) were selected for revised gene

signatures. Only genes measured on both RNAseq and U133A platforms were

considered, and the U133A data from 162 GBM samples measured on both platforms

(which included the 144 cases used to compare U133A and RNAseq results) was used

in the final comparative analysis.

Molecular classifications based on ssGSEA enrichment scores

Single sample gene set enrichment analysis was performed as follows. For a given

GBM sample, gene expression values were rank-normalized and rank-ordered. The

Empirical Cumulative Distribution Functions (ECDF) of the signature genes and the

remaining genes were calculated. A statistic was calculated by integration of the

difference between the ECDFs, which is similar to the one used in GSEA but is based

on absolute expression rather than differential expression [11]. Since the ssGSEA test is

based on the ranking of genes by expression level, the uncentered and log-transformed

U133A and RPKM expression levels were used as input for ssGSEA. Since the scores

of the three signatures were not directly comparable, we performed a resampling

procedure to generate null distributions for each of the four subtypes. First we

generated a large number of virtual samples in which each gene obtains its expression

level by randomly selecting an expression value of the same gene in the remainder of

the samples. Then, the three ssGSEA scores for each signature were calculated.

Following this procedure we generated a large number (>1,000,000) of random ssGSEA

scores for each subtype, to build the null distribution and to give empirical p-values for

the raw ssGSEA scores of each sample. By testing on multiple datasets with different

sample sizes, we found the resampling generated distribution could be replaced with

Student-T distribution (sample size>30) or Normal distribution (sample size>50) for

getting very similar results. R-library with the code and expression matrices used is

provided upon request.

Evaluate the simplicity of subtype activation

For a single sample, we decreased rank the empirical p-values for each subtype to

generate order statistics as $R_{N-1}, R_{N-2} \dots R_1, R_0$. In particular, R_0 equals to the minimum

empirical p-value and points to the dominant subtype, i.e., the most significantly

activated subtype. The accumulative distance to the dominant subtype (ADDS) was

defined as:

$$ADDS = \sum_{i=1}^{N-1} (R_i - R_0)$$

642   Similarly, the accumulative distance between non-dominant subtypes (ADNS) as:

$$ADNS = \sum_{j>i>0} (R_j - R_i)$$

643   Obviously, the ADDS and ADNS are positive and negative correlated with single

644   activation, respectively. Hence, we defined the simplicity score by combing ADDS and

645   ADNS together and corrected with a constant $\frac{(R_{N-1}-R_0)}{N-1}$ as follows:

$$Simplicity\ score = [ADDS - ADNS] \times \frac{(R_{N-1} - R_0)}{N - 1}$$

646   In a three subtypes case, the simplicity score of subtype activation can be simplified as:

647 $$Simplicity\ score = (median\text{-}min) \times (max\text{-}min).$$

Tumor purity assessment

The ESTIMATE package was used to evaluate tumor purity on the basis of the expression level of marker genes in stromal and immune cells [12], where the fraction of stromal cells and immune cells in each sample were represented by stromal score and immune score respectively, and the mixed fraction of both stromal and immune cells was represented by estimate scores. The ABSOLUTE package was used to confirm the tumor purity on the basis of chromosome copy number and allele fraction ratios on samples for which single nucleotide polymorphism array data were available [13].

Sample collection and Neurosphere Cultures

After obtaining approval from the institutional review board of The University of Texas M.D. Anderson Cancer Center, glioblastoma tumor tissues were collected and named in the order that they were acquired. Each tissue was enzymatically and mechanically dissociated into single cells and grown in DMEM/F12 media supplemented with B27 (Invitrogen), EGF (20 ng/ml), and bFGF (20 ng/ml), resulting in neurosphere growth. All cell lines were tested to exclude the presence of Mycoplasma infection. To minimize any batch effect the downstream molecular analyses were performed on identical cell culture batches. Total RNA from formalin fixed, paraffin embedded tumor tissues and matching neurospheres was prepared using the Masterpure complete DNA and RNA isolation kit (Epicenter) after proteinase K digestion per to the instructions from the manufacturer. Paired-end Illumina HiSeq sequencing assays were performed resulting

in a medium number of 50 million 75bp paired end reads per sample. We employed the

PRADA pipeline to process the RNA sequencing data [10]. In short, Burroughs-Wheeler

alignment, Samtools, and Genome Analysis Toolkit were used to map short reads to the

human genome (hg19) and transcriptome (Ensembl 64) and RPKM gene expression

values were generated for each of the 135,994 transcripts of 21,165 protein coding

genes in Ensembl database.

Western blotting

Lysates were prepared from fresh frozen sections using RPPA lysis buffer (1% Triton X-

100 50mM HEPES pH 7.4, 150mM NaCl, 1.5mM MgCl2, 1mM EGTA, 100mM NaF,

10mM Na pyrophosphate, 1mM Na3VO4, 10% glycerol, plus protease and phosphatase

inhibitors cocktails from Roche Applied Science #05056489001 and 04906837001), with

sonication and clearing by centrifugation at 10,000g. Protein concentration was

measured using the BCA kit (Thermo Scientific - Pierce #23225). SDS-PAGE and

western blotting was performed using Midi gel system (Life Technologies - #WR0100)

and NuPage-Novex 4-12% Bis-Tris Midi (20-well) Protein Gels (Life Technologies -

#WG1402) using the following antibodies: ITGAM (CD11B) (Sigma Aldrich –

#HPA002274), IBA1 (AIF1) (Sigma Aldrich – #HPA049234), GFAP (Cell Signalling –

#3670), YKL40 (CHI3L1) YKL40 (CHI3L1, Santa Cruz Biotechnology - #sc-30465) a-

actinin (Sigma Aldrich A5044) and Tubulin (Sigma Aldrich T9026).

Immunohistochemistry

Formalin-fixed, paraffin-embedded tissue sections (4 µm thick) were collected on

Superfrost plus slides. Briefly, tissue sections were deparaffinized with xylene and

ethanol and re-hydrated with 95, 70 and 50% ethanol. Sections were antigen unmasked

using citrate buffer (Vector Labs #H-3300) and heating. Peroxidase block was

conducted with 3% H2O2 and blocking was with 5% goat serum (Vector Labs #S-1000).

Primary rabbit polyconal antibody against IBA1 (AIF1)(WAKO #016-20001) at 1:400

was used overnight. Secondary antibody was done using with the Rabbit-on-Rodent

HRP-Polymer (Biocare #RMR622L) for 1 hr at room temperature. The slides were

developed with Nova-red (Vector Labs #SK-4800) and counterstained with

haematoxylin, mounted and scanned with Pannoramic 250 slide scanner (Caliper Life

Sciences).

Analysis of the Human Protein Atlas database

We systematically queried the human protein atlas database (www.proteinatlas.org,

Version 12) for immunohistochemistry of specific markers of macrophage/microglial

lineage[14]. Typically, each antibody was used on a set of tissue microarrays covering

healthy controls and set of cancers of diverse sites, typically 10-15 individual cases for

each cancer. Only antibodies with high validation scores (concordant RNA seq/IHC

staining, western blot bands and antigen profile) were retained. ITGAM (Antibody HPA

002274), IBA1 (Antibody HPA 049234) and CD163 (Antibody HPA 002432) met these

criteria, as well as control protein vinculin (Antibody CAB002453).

Supplementary References

Phillips, H. S. *et al.* Molecular subclasses of high-grade glioma predict prognosis, delineate a pattern of disease progression, and resemble stages in neurogenesis. *Cancer cell* **9**, 157-173, doi:10.1016/j.ccr.2006.02.019 (2006).
Joo, K. M. *et al.* Patient-specific orthotopic glioblastoma xenograft models recapitulate the histopathology and biology of human glioblastomas in situ. *Cell Rep* **3**, 260-273, doi:10.1016/j.celrep.2012.12.013 (2013).
Kwon, S. M. *et al.* Recurrent Glioblastomas Reveal Molecular Subtypes Associated with Mechanistic Implications of Drug-Resistance. *PLoS One* **10**, e0140528, doi:10.1371/journal.pone.0140528 (2015).
Kim, H. *et al.* Whole-genome and multisector exome sequencing of primary and post-treatment glioblastoma reveals patterns of tumor evolution. *Genome research* **25**, 316-327, doi:10.1101/gr.180612.114 (2015).
Brennan, C. W. *et al.* The somatic genomic landscape of glioblastoma. *Cell* **155**, 462-477, doi:10.1016/j.cell.2013.09.034 (2013).
Verhaak, R. G. *et al.* Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1. *Cancer cell* **17**, 98-110, doi:10.1016/j.ccr.2009.12.020 (2010).
Dai, M. *et al.* Evolving gene/transcript definitions significantly alter the interpretation of GeneChip data. *Nucleic Acids Res* **33**, e175, doi:10.1093/nar/gni179 (2005).
Sandberg, R. & Larsson, O. Improved precision and accuracy for microarrays using updated probe set definitions. *BMC Bioinformatics* **8**, 48, doi:10.1186/1471-2105-8-48 (2007).
Bengtsson, H., Ray, A., Spellman, P. & Speed, T. P. A single-sample method for normalizing and combining full-resolution copy numbers from multiple platforms, labs and analysis methods. *Bioinformatics* **25**, 861-867, doi:10.1093/bioinformatics/btp074 (2009).
Torres-Garcia, W. *et al.* PRADA: pipeline for RNA sequencing data analysis. *Bioinformatics* **30**, 2224-2226, doi:10.1093/bioinformatics/btu169 (2014).
Barbie, D. A. *et al.* Systematic RNA interference reveals that oncogenic KRAS-driven cancers require TBK1. *Nature* **462**, 108-112, doi:10.1038/nature08460 (2009).
Yoshihara, K. *et al.* Inferring tumour purity and stromal and immune cell admixture from expression data. *Nature communications* **4**, 2612, doi:10.1038/ncomms3612 (2013).
Carter, S. L. *et al.* Absolute quantification of somatic DNA alterations in human cancer. *Nature biotechnology* **30**, 413-421, doi:10.1038/nbt.2203 (2012).
Murray, P. J. & Wynn, T. A. Protective and pathogenic functions of macrophage subsets. *Nat Rev Immunol* **11**, 723-737, doi:10.1038/nri3073 (2011).