Measuring Mechanodynamics using an Unsupported Epithelial Monolayer Grown at an Air-Water Interface

Corinne Gullekson[1], Matthew Walker[2], James L. Harden[1,3], Andrew E. Pelling[1,2,4,5,*]

[1]Centre for Interdisciplinary NanoPhysics, Department of Physics, MacDonald Hall, 150 Louis Pasteur, University of Ottawa, Ottawa, ON, K1N5N5 Canada
[2]Department of Biology, Gendron Hall, 30 Marie Curie, University of Ottawa, Ottawa, ON, K1N5N5 Canada
[3]Ottawa Institute of Systems Biology, Roger Guindon Hall, 451 Smyth Road, University of Ottawa, Ottawa, ON, K1N5N5 Canada
[4]Institute for Science Society and Policy, Simard Hall, 60 University, University of Ottawa, Ottawa, ON, K1N5N5 Canada
[5]SymbioticA, School of Anatomy, Physiology and Human Biology, University of Western Australia, Perth, WA, 6009

Short Title:
Mechanodynamics of an Unsupported Epithelial Monolayer

Keywords:
Actomyosin contractility, Cell mechanics, Tissue mechanics, Hanging drop

* Author for correspondence
Andrew E. Pelling
150 Louis Pasteur
MacDonald Hall
University of Ottawa
Ottawa, ON K1N 6N5
Canada
Tel. +1 613 562 5800 Ext 6965
Fax. +1 613 562 5190
Email: a@pellinglab.net
Web: http://www.pellinglab.net

ABSTRACT

Actomyosin contraction and relaxation in a monolayer is a fundamental

biophysical process in development and homeostasis. Current methods used to

characterize the mechanodynamics of monolayers often involve cells grown on solid

supports such as glass or gels. The results of these studies are fundamentally influenced

by these supporting structures. Here, we describe a new methodology for measuring the

mechanodynamics of epithelial monolayers by culturing cells at an air-liquid interface.

These model monolayers are grown in the absence of any supporting structures allowing

us to remove cell-substrate effects. This method's potential was evaluated by observing

and quantifying the generation and release of internal stresses upon actomyosin

contraction (320 ± 50Pa) and relaxation (190 ± 40Pa) in response to chemical treatments.

This is in contrast to the results observed in monolayers grown on solid substrates (glass

and gels) where movement was drastically muted. Tracking the displacement of cell

nuclei, cell edges and cluster perimeter allowed us to quantify the strain dynamics in the

monolayer indicating the reliability of this method. New insights were also revealed with

this approach. Although unsupported monolayers exhibited clear major and minor strain

axes, they were not correlated to the general alignment of cell nuclei. The situation was

dramatically different when the monolayers were grown on soft gels and hard glass

substrates. It was observed that both gels and glass substrates led to the promotion of

long-range alignment of cell nuclei. In addition, the strain orientation was correlated to

nuclear alignment on the soft deformable gels. This new approach provides us with a

picture of basal actomyosin mechanodynamics in a simplified system allowing us to infer

how the presence of a substrate impacts actomyosin contractility and long-range multi-

cellular organization and dynamics. This new methodology will also enable many new

questions to be asked about the molecular regulation of the mechanodynamics of

unsupported monolayers.

INTRODUCTION

Contractility is involved in the remodeling and organization of the cell interior

and plays a major role in multi-cellular morphogenesis over long distances and timescales

(1–3). There is increasing interest in understanding how contractility manifests in multi-

cellular systems (1,4,5). Contractile forces generated by the actomyosin cytoskeleton

within individual cells collectively generate tissue-level forces (6,7). A major challenge

in developmental biology is to understand how cytoskeletal activity is orchestrated to

produce higher order tissue organization (5). Therefore, understanding tissue

morphogenesis requires determining how cellular forces are integrated across cells and

tissues (7).

Actomyosin contraction is one of the major sources of internal force inside of the

cell. The actin cytoskeleton in epithelial monolayers is highly dependant on cell

attachment. In epithelia, the cell-cell attachment points, adherens junctions, link a thick

circumferential ring of actin and myosin around each cell, aligned with the cell borders

(8,9). The contractility of these marginal actin bundles is used for morphogenesis

facilitating epithelial sheet bending and invagination (10,11). The substrate attachment

points, focal adhesions, link actin stress fibers. Cell contractility is commonly described

and investigated in terms of changes in cortical cell elasticity as well as traction

dynamics, the resistance of the substrate to deformation, on flexible substrates (12,13).

Generally changes in cortical cell elasticity and traction dynamics are linked to sensing of

the mechanical microenvironment (14,15). Traction dynamics are linked to cortical

elasticity, focal adhesion organization and cell morphology (13,16,17). Often

fundamental studies of actomyosin contraction and relaxation are examined in situations

where cells are grown on, or embedded in, flexible substrates. (13,18–20). Currently

there is intense interest in understanding the interplay and feedback between the

mechanical properties of such substrates and contractility (14). Although such studies

typically employ substrates with systematically altered mechanical properties, the

substrate is always present. Therefore, it remains difficult to assess the intrinsic internal

mechanical dynamics and properties of cells in the absence of a mechanically supporting

surface.

There has been recent interest in intrinsic mechanical properties of substrate-free

epithelial sheets (3,21). Measurements of the elasticity and failure of monolayers

suspended between two flexible rods provided valuable insights into the understanding of

epithelial mechanics (3,21). Importantly, the measurements yielded mechanical

properties of the near substrate-free cell sheet. With tensile testing of the suspended

cultured monolayers, it was determined that rupture of intercellular junctions occurs after

doubling the monolayer in length with an average force approximately nine times larger

than measured in pairs of isolated cells (3,22), demonstrating that epithelial organization

leads to a strong enhancement in the integrity of the tissue. However this suspended

monolayer is still mechanically constrained at suspension points that provide an external

form of confinement and pre-stress.

This begs the question, how are cell contractility, the organization of the cell

interior and long range structural remodeling altered in the absence of mechanically

confining or supporting structures in a multi-cellular system? Therefore, the objective of

this study was to develop a new methodology to characterize actomyosin

mechanodynamics in a multi-cellular system when no resistance is offered by the

surroundings. Understanding these dynamics could provide us with a picture of basal

actomyosin mechanodynamics in a simplified model multi-cellular system. This method

will also allow us to indirectly infer how the presence of a substrate impacts actomyosin

contractility and long-range multi-cellular organization and dynamics.

Here, we employed an in-vitro hanging drop culture protocol. We created

Madine-Darby Canine Kidney (MDCK) epithelial monolayer clusters at an air-water

interface. Employing laser scanning confocal microscopy, cell nuclei within the

monolayer were used as fiduciary markers to determine the internal strain developed in

the clusters during basal and chemically-induced contractile changes. We validated our

method using two alternate methods for tracking cell deformation: 1) tracking the cell

membrane boundary of the cluster and 2) tracking cell boundaries marked with GFP-

actin. Our results indicate that these monolayers exhibited complex, often anisotropic,

contractile/relaxation dynamics. This is contrast to similar experiments performed in this

study on flexible and rigid substrates, where the presence of the substrate was found to

significantly impair contractile strain dynamics. Importantly, this new methodology for

studying the mechanodynamics of unsupported epithelial sheets opens the door for

several future research directions including investigations of the apico-basal polarization,

myosin distribution and actin ordering in the unsupported monolayer. This approach

allows one to observe the results of forces that are important for many processes that take

place in development (23) and may advance the understanding of how those forces are

being transmitted to neighbouring cells, and how they are integrated to trigger global

changes in tissue shape. The results of this study might also be relevant to 3D cell culture

where hard surfaces are not present.

MATERIALS AND METHODS

Cell culture

Madin-Darby Canine Kidney (MDCK) epithelial cells were cultured in DMEM

with 10% fetal bovine serum (FBS), 50mg/ml streptomycin and 50U/ml penicillin

antibiotics (all from Hyclone Laboratories Inc.). Cells were cultured at 37℃ in a 5% CO_2

incubator on 100mm tissue culture dishes (Corning). In some experiments, cells were

cultured on solid substrates. In these cases, $1x10^5$ cells were cultured in 34mm tissue

culture dishes (TPP). The dishes either contained a flexible (soft) polyacrylamide (PA)

hydrogel coating, previously described (24–26), or a bare glass coverslip bottom. To

produce a PA gel with a young's modulus of 4.8kPa (27), 7.5% acrylamide was

polymerized with 0.053% bisacrylamide on (3-Aminopropyl)trimethoxysilane coated

coverslips with 1mg/ml ammonium persulfate (Biorad) and 0.15%

tetramethylethylenediamine (Biorad). Prior to cell seeding, the hydrogel surface was

functionalized with 100μg/mL collagen (Gibco) in PBS using 0.5mg/mL sulfo-sanpah

(Pierce) as a crosslinker. In some experiments, cells were cultured in hanging drops.

MDCK cells were trypsinized and then resuspended in media at a low dilution of 4000

cells/ml. The cells were then placed in hanging drops of 40μl on the lower surface of the

lids of plastic Petri dishes containing Phosphate Buffered Saline (PBS). Hanging drops

were imaged after 24h of culture.

Drug treatments

Cells were treated with blebbistatin (Sigma) dissolved in DMSO at a final

concentration of $40\,\mu$M or calyculin A (CalA, Sigma) dissolved in water at a final

concentration of 300nM for 30 minutes in an incubator at 37℃ and 5% CO_2 between

before and after images.

Imaging

All images were acquired on a TiE A1-R laser scanning confocal microscope

(LSCM) (Nikon) with a 40X long working distance objective. Images were acquired with

a standard LSCM configuration with appropriate laser lines and filter blocks. Cell nuclei

were stained with Hoechst. In some cases, cells expressing GFP-actin were produced

with BacMam reagent (Invotrogen, CellLight® Reagent BacMam 2.0) using manufacture

protocols at 20 particles per cell. Cells were plated and incubated for at least 48h before

being placed into hanging drops using the protocol above. Cells were also stained with

wheat germ agglutinin (WGA) coupled to Texas Red-X (Invitrogen) to reveal the cell

membrane. Cells stained on glass were fixed with 3.5% paraformaldehyde and

permeabilized with Triton X-100 at 37°C. Actin was stained with Phalloidin Alexa Fluor

546 (Invitrogen). Following this vinculin was stained with monoclonal anti-vinculin

primary antibody (Sigma) and Alexa Fluor 488 rabbit anti-mouse immunoglobin

(Invitrogen) secondary antibody. DNA was labeled with DAPI (Invitrogen).

Strain Quantification

The strain in the monolayer was determined with the displacements of the cell

nuclei during a 30 minute treatment. This method was validated by tracking the cell

membrane boundary of the cluster and by tracking actin in the cell margins. Images were

processed and analyzed with ImageJ. Maximum intensity z-projections of 10, 5µm thick

slices were used for strain calculations with no enhancements applied (Fig 1A). The

positions of the nuclei in these images were determined using the analyze particles

function after thresholding. The positions of the nuclei in the before and after images

were manually linked and analyzed using Matlab. A Matlab program was used to account

for differences in dish placement in the before and after images, as well as to determine

nuclear displacements and calculate the strain tensor of the cluster.

FIGURE 1: The process used to analyze the images. A) Maximum intensity Z-

projections of the before and after confocal images of cells in a hanging drop used for a

control (scale bar, 10μm). B) The positions of the nuclei in the before (blue) and after

(red) images determined by threshholding. The raw nuclear positions (left) are shown as

well as the positions after the translation (middle) and after the rotation (right) of the after

positions. C) Positions of the before (blue) and after (red) nuclei in a mesh over a mean

strain map of the cluster. The major and minor principle axes (E_1 and E_2) of the cluster

are in the y and x directions respectively. Maps are shown for the control (left) case as

well as CalA (middle) and blebbistatin (right) treatments.

In this program, nuclei positions were translated in both images so that the center

of the nuclei was at the origin. These translated positions were then rotated so that the

sum of the nuclei displacements were minimized (Fig 1B). This process was needed so

that there were no artificially large displacements from the movement of the dish to and

from an incubator between images. The initial positions (x, y) and displacements (u, v) of

the nuclei were then used to fit the cluster with least squares to the displacement

equations: $u=c_1x+c_2y+c_3xy+c_4x^2+c_5y^2$ and $v=c_6x+c_7y+c_8xy+c_9x^2+c_{10}y^2$ (28). Second

order terms were included to account for non-constant strain throughout the cluster.

These equations were then used to determine the diagonal components of the Green-

Lagrangian strain tensor,

$E_{xx} = \frac{1}{2}\left(2\frac{\partial u}{\partial x} + \frac{\partial u}{\partial x}\frac{\partial u}{\partial x} + \frac{\partial v}{\partial x}\frac{\partial v}{\partial x}\right),$

$E_{yy} = \frac{1}{2}\left(2\frac{\partial v}{\partial y} + \frac{\partial u}{\partial y}\frac{\partial u}{\partial y} + \frac{\partial v}{\partial y}\frac{\partial v}{\partial y}\right)$ (29).

Substitution of our displacement equations resulted in

$E_{xx} = \frac{1}{2}\left(2(c_1 + c_3 y + 2c_4 x) + (c_1 + c_3 y + 2c_4 x)^2 + (c_6 + c_8 y + 2c_9 x)^2\right),$

$E_{yy} = \frac{1}{2}\left(2(c_7 + 2c_{10} y + c_8 x) + (c_2 + 2c_5 y + c_3 x)^2 + (c_7 + 2c_{10} y + c_8 x)^2\right).$

These were used along with the shape of the cluster to determine the average x-axis

extensional strain, E_{xx}, and average y-axis extensional strain, E_{yy}, in the cluster. The

cluster was then rotated to find the principle axes by maximizing the difference between

the y-axis strain and the x-axis strain. After this rotation, the y-axis strain was the major

principle strain (E_1) and the x-axis strain was the minor principle strain (E_2) (Fig 1C). The

mean of the principle strains, $E_M = \frac{E_1 + E_2}{2}$, is the deformation due to a change in area

(30).

For experiments measuring the deformation of the cluster outline, points along the

cluster outline labeled in the cell membrane stained images were tracked and used in the

code above in place of nuclei positions. In some experiments cellular-level strains were

determined in cells expressing actin-GFP. Cell deformation was determined by fitting cell

boundaries with polygons. The displacements of the vertices were least squares fit to the

displacement equations u=c_1x+c_2y and v=c_3x+c_4y to impose uniform strain inside the cell

(28). Only first order terms in the displacements u and v were required for these

calculations because the individual cells were much smaller than the cluster. The strain in

each cell was calculated using the principle strain axes of the cluster determined using the

nuclei displacements.

Order Parameter Calculations

Orientational order parameters were calculated using the angles between different

cluster features. The nematic order parameter was used, which in two dimensions is

$S = \langle 2\cos^2\theta - 1 \rangle$ (31). The order parameter will be 1 if the features are completely

aligned, -1 if they are completely antialigned, and 0 if there is no alignment. The nuclei in

the thresholded before images were fitted to ellipses in Matlab to determine the

orientations of the major length axes and the eccentricities of the nuclei. The orientation

of the major length axis of the cluster was determined using the positions of the nuclei

and finding the axis that minimized the square of the distances between the axis and the

nuclei. Four different types of order parameters were calculated. Nuclei-strain order

parameters (S_{ns}) were determined using the angles between the major length axes of the

nuclei and the major strain axis of the cluster. Nuclei-cluster orientation order parameters

(S_{nl}) were calculated using the angles between the major length axes of the nuclei and the

cluster major length axis. Cluster orientation-strain order parameters (S_{ls}) were also

calculated with the angles between the major length and major strain axes of clusters. In

addition, nuclei-nuclei order parameters (S_{nn}) were calculated using the angles formed

between the major axis of each nucleus and the major axis of its nearest neighbor. This

was also repeated for next nearest neighbors, third nearest neighbors and fourth nearest

neighbors.

RESULTS

Deformation of Monolayer Clusters on Solid Substrates

To examine the influence of a mechanically supporting substrate, monolayer clusters were created on hard glass substrates and soft polyacrylamide (PA) gels (4.8 kPa Young's modulus). These clusters naturally occur in MDCK cell culture at low confluence. The clusters on glass were fixed and stained for DNA, actin and vinculin (Fig 2 A). A strong actin signal was localized in the perimeter of the cluster. Similarly the vinculin also localized around the cluster perimeter. When epithelial cells form a cluster on solid substrates, more cellular adhesions and force generation appear at the cluster perimeter (32–34). In order to induce cluster relaxation and contraction in the clusters we employed the agents blebbistatin and CalA. Blebbistatin is a myosin II inhibitor that inhibits contraction and disrupts contractile filament organization (35). CalA is a phosphatase inhibitor that is well known to induce contraction (5,35). Clusters on glass were live stained with Hoechst and the nuclei were tracked with the addition of drugs in order to determine strain in the cluster, as described in the Materials and Methods. There was no significant difference in the mean strains of the clusters on glass with the different drug treatments (p>0.12) (Fig 2 B). To investigate the deformation on a softer substrate, clusters were cultured on PA gels. Consistent with the glass substrates, there was no significant difference in the mean strains of the clusters on the gels with the different drug treatments (p>0.07) (Fig 2 C). To more thoroughly examine the intrinsic deformation of a contracting or relaxing monolayer the solid substrate should be removed, for this reason the hanging drop method was introduced.

FIGURE 2: Deformation of the cluster on glass and PA. A) Confocal images of the actin (left), vinculin (middle) and merged (right, actin=red, vinculin=green, nucleus=blue) of a

cluster of MDCK cells on glass (scale bar, $10\mu m$). B) Major E_1 minor E_2 and mean E_M

strains of cell clusters on glass after the addition of blebbistatin (red), media (blue) or

calyliculin A (green). n=11, 9 and 12, no significance. C) Major E_1 minor E_2 and mean

E_M strains of cell clusters on a PA gel after the addition of blebbistatin (red), media (blue)

or calyliculin A (green). n=15, 10 and 15, no significance.

Formation of a cell monolayer cluster

As MDCK cells migrate and proliferate in 2D culture, the cells form islands that

eventually merge to form a continuous monolayer (36). In 3D culture, individual MDCK

cells plated within an extracellular matrix gel will assemble into a hollow sphere that is

lined by a monolayer of polarized epithelial cells (36). However, after incubation in a

hanging drop, MDCK cells will form monolayer clusters of cells (37–39). This is in

contrast to mesenchymal stem cells, which gradually coalesce into a single central

spheroid along the lower surface of the drop (40). It has been shown that MDCK cells in

a hanging drop will form a basement membrane-like sheet of cell-secreted proteins

providing the matrix for the proliferating cells (37,39). Previous immunohistochemical

studies demonstrated the presence of type IV collagen, laminin 1, and laminin 5 in the

membrane, and enzyme digestion experiments indicated that this membrane was sensitive

to collagenase (37). In previous studies, it has been shown that the tension carried by

basement membranes are modest compared with the tension carried by the cellular

portion of the epithelium (41) and that the basement membrane is thinner and softer than

the cellular component with a Young's modulus of 7.5kPa (42) compared with 20kPa (3).

Moreover, this basement membrane is relatively free to move along the air-water

interface. Therefore, this basement membrane-like sheet allows cells to form healthy

monolayers while still being relatively unrestrictive to collective cellular deformation.

Epithelial monolayer clusters were grown in these hanging drops and their nuclei

were used to track deformations during changes in contractility. The clusters were similar

in number of cells (average of 23±2 cells per cluster) to the ones used in the solid

substrate experiments (average of 21±2 cells per cluster). To label cell nuclei, we added

the common nuclear stain, Hoechst 33342, to the drop with a pipette in the lid's inverted

state (Fig 3 A and B). In all cases, 1μL of a stock solution of blebbistatin, CalA or media

(control) was added to the hanging drop for 30 minutes in a cell culture incubator. The

nuclei of suspended clusters were imaged before and after the addition of the drugs or

control. On average, only five monolayer clusters formed in each hanging drop and it was

straightforward to image the same cluster before and after the 1μl addition.

FIGURE 3: Experimental protocol used to create monolayer clusters in hanging drops.

A) The process used to grow, stain and image the monolayer clusters before and after the

addition of cytoskeletal drugs. 40μl droplets were incubated overnight with PBS in the

bottom of the dish. The dish lid was flipped and 1μl of Hoechst in media was added with

a pipette. The droplets were then imaged with a confocal microscope. After this, drugs

were added to the droplets, which were incubated for 30minutes and imaged again. B) A

typical set of clusters found in a droplet after the addition of Hoechst (scale bar, 20μm).

C) A diagram describing the shape of the droplet. The droplet shape can be approximated

as a spherical cap with a radius of curvature, R. The radius at the droplet's top, r, is

2.9±0.1mm. The approximate height of the droplet, h, is calculated with the known

volume of the droplet, $V = \frac{1}{6}\pi h(3r^2 + h^2)$.

Hanging drop shape

The shape of the droplet observed in this set up can be approximated as a

spherical cap (Fig 3 C). The radius, r, at the droplet's top, where it is attached to the dish

lid, was measured optically at 2.9±0.1mm (n=7). The droplet volumes were measured

using the mass of the droplets to be 39.49±0.04, 40.29±0.06 (after Hoechst addition) and

41.18±0.07μl (after drug addition, n=10). Using the spherical cap model, the height, h,

and radius of curvature, R, of the initial droplet were calculated to be 2.5±0.1mm and

2.9±0.1mm respectively. With the addition of 2μl, the height of the droplet increased by

74.4±0.5μm and the radius of curvature decreased by 14±8μm (0.5% decrease). The

average diameter of clusters was approximately 150μm. With the comparatively large

radius of curvature, the center and edge of the cluster had a height difference of only

0.96±0.04μm. For this reason it was reasonable to approximate the curved bottom of the

drop as a flat surface. Due to the small size of the droplet, scans were acquired quickly to

avoid evaporation effects. Under the imaging power employed in this study, some

evaporation of the droplet did occur, resulting in a height decrease of 0.24±0.02μm per

second, which corresponds to a change in volume of 0.32±0.02μl per minute (Fig S1).

For this reason, image collection was always maintained under 45 seconds. The radius of

curvature only decreased by a maximum of 1.5±0.1μm during imaging, having little

effect on the monolayer clusters' curvature (under 0.05%).

Deformation of Monolayer Clusters in Hanging Drops

The cytoskeletal drugs, blebbistatin and CalA, were used to relax and contract the

cell clusters. The displacements of the cell nuclei were fit to determine the major and

minor principle strains (E_1 and E_2) of the cluster. As a control, 1μl of media was added to

the drops followed by incubation for 30 minutes between imaging. We did observe slight

movement of nuclei within the cluster. However, the mean strain of the clusters in the

control study was not significantly different than zero ($p > 0.89$) (Fig 4). As expected there

was no significant dilation or contraction. The myosin II inhibitor, blebbistatin, caused

clusters to dilate appreciably. The mean strain was significantly higher ($p < 0.00001$) than

the control (Fig 4). Clusters also contracted with the addition of CalA resulting in a mean

strain that was significantly lower ($p < 0.0004$) than the control (Fig 4).

FIGURE 4: Deformation of the clusters in hanging drops with cytoskeletal drugs. Major

(E_1), minor (E_2) and mean (E_M) strains of cell clusters after the addition of blebbistatin

(Red), media (blue) or CalA (green). $n = 22$, $p < 0.05$.

Cluster Outline Deformation

To determine if the perimeter of the monolayer cluster itself deforms in a manner

consistent with the changes in nuclei position, the cluster outline was tracked by

fluorescently labeling cell membranes with WGA. Tracking points on the cluster

perimeter allowed us to determine monolayer strain (Fig S2 A). The major and minor

strains in the cluster were calculated from nuclei positions as well as from its outline. The

strains calculated using cell nuclei and cluster outlines were not significantly different for

the untreated control ($p > 0.90$), blebbistatin ($p > 0.85$) and CalA treatments ($p > 0.82$) (Fig

S2 B). This suggests the contraction and relaxation observed in the nuclear positions is a

good indicator of total cluster deformation.

Individual Cellular Strains

To determine individual cellular strains during cluster contraction and relaxation,

cells transiently expressing actin-GFP were used to form suspended clusters. It was found

that when a high percentage of cells were expressing actin-GFP, CalA treatments had no

effect on contraction. Therefore, we adjusted the transfection efficiency to ~25% of cells

in order to create suspended clusters that exhibited the same mechanical dynamics as

non-transfected cells. In this case, we felt confident that we would be able to examine

individual cellular strains, albeit only on a sub-population of cells within each cluster.

Distinct actin stress fibers were not evident but actin was found localized to the cell

margins allowing us to easily determine cell boundaries (Fig 5 A B). Experiments with

blebbistatin, CalA and control conditions were performed once again and nuclei positions

were tracked, as well as the deformation of single cells within the cluster.

FIGURE 5: Cellular deformations inside a deforming hanging drop cluster. A) The

before image of a MDCK cluster stained with Hoechst (blue) partially transduced with

GFP-actin (green) (scale bar, $10\mu m$). B) The mean strain of each cell overlaid on the

before image. C) Major, minor and mean strain of cell clusters after the addition of

blebbistatin, media, or CalA (blue) compared to the strain in the individual cells as

calculated with the actin cell boundaries (green).

In the control, the cluster underwent normal basal remodelling as seen by nuclear

displacements. With the addition of CalA and blebbistatin the clusters the nuclear

displacements showed contraction and relaxation of the cluster as expected (Fig 5 C). The

mean of the strains in the individual cells were not significantly different than the cluster

strains calculated by nuclear displacements in the control ($p>0.97$), blebbistatin ($p>0.96$)

and the CalA treatments ($p>0.98$). This demonstrates that the monolayer cluster strains

determined by nuclei correspond well to the deformation of the individual cells

themselves.

Substrate Effects on Strain

Comparing the blebbistatin or CalA induced strains on the glass or PA substrates to the hanging drop (Fig 6 A) reveals that they were significantly muted ($p<0.002$ in both cases on glass, and $p<0.04$ for blebbistatin on PA) by the substrate. Interestingly however, there was no significant difference in the controls ($p>0.87$).

FIGURE 6: Substrate dependence on nuclear deformation A) Mean E_M strains of cell clusters in the hanging drops (solid, n=22), on PA gels (stripes, n=15, 10 and 15) and glass (dots, n=11, 9 and 12) after the addition of blebbistatin (red), media (blue) or CalA (green). B) Order parameter, S_{ns}, for nuclei (defined by the angle between the major strain axis and the nuclear major axis) for blebbistatin (red), control (blue) and CalA (green) treatments for hanging drops, PA and glass. C) Order parameters, S_{nn} for nuclei (defined by the angle between the nuclear major axis and its neighbor's nuclear major axis) in hanging drops (circles), on PA (triangles) and glass (diamonds). The results are shown for nearest, next nearest, third nearest and fourth nearest neighbors. $p< 0.05$.

Feature Orientation

In order to determine if the deformation of the cluster exhibited a dependence on structural features of the cells, nematic order parameters were calculated as described in the Materials and Methods. The order parameter for nuclear alignment with strain (S_{ns}) was calculated for all the clusters (Fig 6 B). The order parameters in the hanging drops were not significantly different than 0 ($p>0.82$, 0.64 and 0.20 for blebbistatin, control and CalA, respectively), suggesting no preferential alignment. On glass, the order parameters were also not significantly different than 0 ($p>0.71$, 0.25 and 0.97 for blebbistatin, control and CalA), respectively, suggesting no preferential alignment. However, since the strains

were small on glass, the major strain axis was likely determined by basal remodeling of

the cells and not the contraction or relaxation of the actomyosin cytoskeleton. On PA, the

control order parameter was not significantly different than 0 (p>0.62), however the order

parameters of treated clusters' nuclei were significantly different than 0. In the

blebbistatin treated clusters, the nuclei preferentially aligned with the major strain axis

(p<0.04). In the CalA treated clusters, the nuclei preferentially aligned with the minor

strain axis (p<0.04). Therefore, the nuclei tended to be aligned with the relaxation and

contraction axes. This demonstrated that there was a correlation between nuclear

alignment and cluster strain on PA gels that was not observed in hanging drops or on

rigid glass substrates.

The order parameter S_{ls}, calculated with the angle between the major length axis

and the major strain axis of the cluster (Fig S3 A), was not significantly greater than 0 for

any treatment or substrate (p>0.09 in all cases). Another order parameter (S_{nl}) examines

if the nuclei were aligned with the major length axis on the cluster. Again, this parameter

was not significantly greater than 0 for any treatment or substrate (p>0.57 in all cases)

(Fig S3 B). The average eccentricities of the nuclei grown on the three different

substrates were not significantly different (p>0.15). The clusters also had varying

major/minor length aspect ratios, however the mean strains observed in the monolayer

clusters did not display any correlation with aspect ratio (Fig S4).

To examine the alignment of cells with their neighbors, the order parameters

between nuclei (S_{nn}) were calculated (Fig 5 D). For all substrates, the order parameters

for the nearest neighbors were significantly greater than 0 (p<0.002, 0.02 and 0.005 for

hanging drops, PA and glass, respectively) suggesting a local alignment of cells in direct

contact. For next nearest neighbors and third nearest neighbors, the order parameter was

not significantly greater than 0 for hanging drops ($p > 0.4$, 0.4), but was significantly

greater than 0 for PA ($p < 0.03$, 0.02) and glass ($p < 0.05$, 0.03). After this, the order

parameters were not significantly different than 0 in any case. This suggests that the cells

on substrates aligned with cells that were near to them even if they were not in direct

contact. Interestingly, this was not seen for cells in hanging drops, suggesting longer-

range cell alignment occurs only in monolayer clusters on solid substrates.

DISCUSSION AND CONCLUSION

Understanding actomyosin contractility is one key aspect of cell mechanics that is

required for explaining the dynamics of tissue remodeling. Direct experimental

measurements of force generation in monolayer clusters without a rigid surface can aid in

our understanding of the remodeling of epithelial monolayers. The objective of this study

was to develop a new experimental methodology and analysis framework in order to

provide an approach to understanding the mechanodynamics of unsupported epithelial

monolayers. The lack of a solid support provided a less complicated system that allowed

for remodeling not possible on a solid substrate. This method also provides a rational

approach to teasing apart substrate and intercellular connections.

As a validation of the practicality of the hanging drop model, we investigated the

effect of cytoskeletal drugs on the deformation of MDCK monolayer clusters in both the

absence and presence of mechanically supporting substrates. By comparing cell response

in these two support regimes, we were able to identify the intrinsic contractile dynamics

of an epithelial monolayer cluster. Since contraction/dilation was muted on solid

supports, removing the solid substrate allowed the monolayer to clusters contract/dilate.

The deformation we observed by tracking nuclei was echoed in the deformation of the

individual cell and the monolayer cluster outlines. This verified the use of nuclei as an

effective cluster deformation marker. The observed deformations established that the

contraction and dilation of monolayer clusters could be directly observed in the absence

of a substrate by tracking cell nuclei in hanging drop clusters.

To calculate stress in our droplets we used the mean strains obtained from our

data and the Young's modulus reported by Harris et al. for MDCK monolayers of

20 ± 2kPa (3). This results in an apparent decrease of stress of 320 ± 50Pa for the

blebbistatin treatment and an apparent increase of stress of 190 ± 40Pa for the CalA

treatment. There was a negligible change in stress for the control case. In other

experiments, MDCK cells grown on pillars have been found to have average traction

stresses of around 800Pa in a monolayer (43). Our stresses are smaller than, but of the

same order of magnitude as, stresses observed in cells grown on pillars. The calculated

stresses are reasonable considering that the hanging drop is expected to support less

traction force than a stiffer substrate. This demonstrates that the hanging drop system can

be used to calculate the intrinsic stress produced by actomyosin contraction and

relaxation in monolayer clusters.

Furthermore, varying the cluster substrate allowed us to relate contractile

dynamics to substrate rigidity. Our hanging drop method allowed us to measure the

intrinsic remodeling that can occur in response to actomyosin dynamics. This remodeling

was muted by the presence of a solid substrate, demonstrating that cluster contraction is

significantly restrained when bound to a rigid surface. In general, stiff substrates increase

cell-substratum adhesion (44–47) and increase traction forces (48). It has also been

shown that compliant substrates promote focal adhesions that are dynamic and irregular

in shape while stiff substrates promote the formation of stable arrays of elongated focal

adhesions (47). When the cells are grown on glass, a large portion of the work from the

actomyosin contraction was converted into substrate deformation instead of cellular

remodeling. On softer substrates (PA gels in particular) cells are still restrained, however

the smaller, more dynamic focal adhesions may allow for more movement. In this case, a

smaller portion of the work will go into deforming the substrate. The contraction energy

of the hanging drop cells is expected to be smaller than the contraction energy of cells on

glass or PA. However, the cells can move more freely in the cluster because they are not

attached by focal adhesions to a substrate that is difficult to deform. Only a negligible

amount of work should go into deforming the basement membrane-like sheet below the

monolayer cluster, allowing for the larger cellular remodeling events observed in this

paper.

Interestingly, when the clusters were grown on PA gels there was a correlation

between major strain direction and nuclear alignment. However, for cells grown on at the

air-water interface in a hanging drop, the direction of the major strain was independent of

nuclear alignment. The nuclei of single epithelial cells grown on flat rigid surfaces have

been previously found to align with the actin cytoskeleton and the major length axis of

the cell (49,50). This suggests that the actin fibers are aligned with the major strain

direction when on PA gels. This is congruous with the theory that individual cells are

contracting and relaxing preferentially in the direction of their actin cytoskeletons

resulting in a global deformation along the cluster axis that many cells were aligned with.

It has also been shown that matrix stiffness aids in regulating the polarization and

alignment of stress fibers within cells (51). The lack of correlation between nuclear

orientation and strain in the hanging drop may be due to reduced alignment of actin

within each cell and the reduction in longer-range correlated cell-cell alignment that was

observed on rigid substrates. In the case of the stiff glass substrate, although there should

be greater alignment due to the substrate, focal adhesions pinned cells to the substrate

leading to null cluster strains, making it arduous to correlate to strain. The softer PA

substrate allowed for substrate mediated alignment and limited cell movement permitting

a correlation between cluster strain and nuclear alignment to be observed.

Remarkably, a solid substrate appears to be required for longer-range cell-cell

alignment. By calculating nuclear order parameters, we found that cells in hanging drops

were only aligned with their nearest neighbors, while cells on substrates were aligned

with their next nearest and third nearest neighbors as well. This longer-range organization

may be a result of substrate cues not present in hanging drops. It suggests also that

signalling is necessary to develop long-range stresses. One possibility is that long-range

orientational correlations are the result of elastic coupling of cells mediated by

deformation of the compliant substrate. This is consistent with work showing that

intercellular forces develop within the cluster of cells grown on a gel (32). This

unexpected finding was made evident by comparison with the extreme condition of an

air-water interface substrate. It would not have been possible without the novel hanging

drop monolayer. Our new methodology has revealed some surprising and novel insights

into the mechanodynamics of epithelial monolayers. However, it also raises several new

questions that will warrant further study in the future.

For instance, do the cells in the monolayer establish apico-basal polarization? Is this polarization triggered without a solid substrate? This could be determined with more advanced staining techniques. Are the same structures formed on planar non-adhesive substrates or is the curved drop shape needed? Plating cells in dishes with non-adhesive coatings could be a method to investigate this. How do clusters grown on substrates recoil when detached? This could be investigated using trypsin to detach the clusters from the substrate or by using thermo-sensitive polyacrylamide derivatives to allow release of the sheets. Future work will include computational modeling of these scenarios to examine these effects in further detail. These hanging drop monolayers can also be made with epithelial cells stably expressing a fluorescently marked protein of interest and could also be utilized for a broad range of mechanical experiments investigating substrate effects on monolayers. Novel tools and methodologies for understanding the mechanics of monolayers will enable the advancement of the biophysics field.

ACKNOWLEDGEMENTS

This work was supported by a Natural Sciences and Engineering Research Council (NSERC) Discovery Grant. C.G. was supported by the uOttawa NSERC-CREATE program in Quantitative Biomedicine. A.E.P. gratefully acknowledges generous support from the Canada Research Chairs (CRC) program.

REFERENCES
1.   Rauzi M, Lenne P-F, Lecuit T. Planar polarized actomyosin contractile flows control epithelial junction remodelling. Nature [Internet]. Nature Publishing Group; 2010 Dec 23 [cited 2014 Jul 12];468(7327):1110–4. Available from: http://www.ncbi.nlm.nih.gov/pubmed/21068726
2.   Roh-Johnson M, Shemer G. Triggering a Cell Shape Change by Exploiting Pre-Existing Actomyosin Contractions. Science (80-) [Internet]. 2012 [cited 2014 Nov 14];335(6073):1232–5. Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3298882/
3.   Harris AR, Peter L, Bellis J, Baum B, Kabla AJ, Charras GT. Characterizing the

mechanics of cultured cell monolayers. Proc Natl Acad Sci USA [Internet]. 2012 Oct 9 [cited 2013 May 28];109(41):16449–54. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3478631&tool=pmcen trez&rendertype=abstract

4.   Nelson CM, Jean RP, Tan JL, Liu WF, Sniadecki NJ, Spector A a, et al. Emergent patterns of growth controlled by multicellular form and mechanics. Proc Natl Acad Sci USA [Internet]. 2005 Aug 16;102(33):11594–9. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1187971&tool=pmcen trez&rendertype=abstract

5.   Fernandez-Gonzalez R, Simoes SDM, Röper J-C, Eaton S, Zallen J a. Myosin II dynamics are regulated by tension in intercalating cells. Dev Cell [Internet]. 2009 Nov [cited 2013 Sep 18];17(5):736–43. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2854079&tool=pmcen trez&rendertype=abstract

6.   Vicente-Manzanares M, Ma X, Adelstein RS, Horwitz AR. Non-muscle myosin II takes centre stage in cell adhesion and migration. Nat Rev Mol Cell Biol [Internet]. Nature Publishing Group; 2009 Nov [cited 2014 Jul 10];10(11):778–90. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2834236&tool=pmcen trez&rendertype=abstract

7.   Martin AC, Gelbart M, Fernandez-Gonzalez R, Kaschube M, Wieschaus EF. Integration of contractile forces during tissue invagination. J Cell Biol [Internet]. 2010 Mar 8 [cited 2014 Jul 22];188(5):735–49. Available from: http://www.jcb.org/cgi/doi/10.1083/jcb.200910099

8.   Owaribe K, Kodama R, Eguchi G. Demonstration of contractility of circumferential actin bundles and its morphogenetic significance in pigmented epithelium in vitro and in vivo. J Cell Biol. 1981;90(2):507–14.

9.   Yonemura S, Itoh M, Nagafuchi a, Tsukita S. Cell-to-cell adherens junction formation and actin filament organization: similarities and differences between non-polarized fibroblasts and polarized epithelial cells. J Cell Sci. 1995;108 (Pt 1:127–42.

10. Leptin M. Gastrulation movements: The logic and the nuts and bolts. Dev Cell. 2005;8(3):305–20.

11. Lecuit T, Lenne P-F. Cell surface mechanics and the control of cell shape, tissue patterns and morphogenesis. Nat Rev Mol Cell Biol [Internet]. 2007 Aug [cited 2014 Jul 9];8(8):633–44. Available from: http://www.nature.com/doifinder/10.1038/nrm2222

12. Rauzi M, Verant P, Lecuit T, Lenne P-F. Nature and anisotropy of cortical forces orienting Drosophila tissue morphogenesis. Nat Cell Biol [Internet]. 2008 Dec [cited 2013 Sep 18];10(12):1401–10. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18978783

13. Discher DE, Janmey P, Wang Y-L. Tissue cells feel and respond to the stiffness of their substrate. Science [Internet]. 2005 Nov 18 [cited 2014 Jul 12];310(5751):1139–43. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16293750

14. Trichet L, Le Digabel J, Hawkins RJ, Vedula SRK, Gupta M, Ribrault C, et al.

Evidence of a large-scale mechanosensing mechanism for cellular adaptation to substrate stiffness. Proc Natl Acad Sci USA [Internet]. 2012 May 1 [cited 2014 Jul 10];109(18):6933–8. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3344951&tool=pmcentrez&rendertype=abstract

15. Vogel V, Sheetz M. Local force and geometry sensing regulate cell functions. Nat Rev Mol Cell Biol [Internet]. 2006 Apr [cited 2014 Jul 10];7(4):265–75. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16607289

16. Solon J, Levental I, Sengupta K, Georges PC, Janmey P a. Fibroblast adaptation and stiffness matching to soft elastic substrates. Biophys J [Internet]. 2007 Dec 15 [cited 2014 Jul 9];93(12):4453–61. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2098710&tool=pmcentrez&rendertype=abstract

17. Pelham R, Wang Y. Cell locomotion and focal adhesions are regulated by substrate flexibility. Proc Natl Acad Sci USA [Internet]. 1997 [cited 2014 Nov 14];95. Available from: http://www.pnas.org/content/94/25/13661.short

18. Ghibaudo M, Trichet L, Le Digabel J, Richert A, Hersen P, Ladoux B. Substrate topography induces a crossover from 2D to 3D behavior in fibroblast migration. Biophys J [Internet]. 2009 Jul 8 [cited 2014 Nov 21];97(1):357–68. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2711370&tool=pmcentrez&rendertype=abstract

19. Huh D, Matthews BD, Mammoto A, Montoya-Zavala M, Hsin HY, Ingber DE. Reconstituting organ-level lung functions on a chip. Science [Internet]. 2010 Jun 25 [cited 2013 Sep 16];328(5986):1662–8. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20576885

20. Tremblay D, Andrzejewski L, Leclerc A, Pelling AE. Actin and microtubules play distinct roles in governing the anisotropic deformation of cell nuclei in response to substrate strain. Cytoskeleton [Internet]. 2013 Dec [cited 2014 Apr 2];70(12):837–48. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24123894

21. Harris AR, Bellis J, Khalilgharibi N, Wyatt T, Baum B, Kabla AJ, et al. Generating suspended cell monolayers for mechanobiological studies. Nat Protoc [Internet]. Nature Publishing Group; 2013 Dec [cited 2014 Aug 13];8(12):2516–30. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24263091

22. Chu Y-S, Thomas W a, Eder O, Pincet F, Perez E, Thiery JP, et al. Force measurements in E-cadherin-mediated cell doublets reveal rapid adhesion strengthened by actin cytoskeleton remodeling through Rac and Cdc42. J Cell Biol [Internet]. 2004 Dec 20 [cited 2014 Mar 19];167(6):1183–94. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2172605&tool=pmcentrez&rendertype=abstract

23. Heisenberg C-P, Bellaïche Y. Forces in tissue morphogenesis and patterning. Cell [Internet]. 2013 May 23 [cited 2014 Jul 15];153(5):948–62. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23706734

24. Wang YN, Galiotis C, Bader DL. Determination of molecular changes in soft tissues under strain using laser Raman microscopy. J Biomech [Internet]. 2000 Apr;33(4):483–6. Available from: http://www.ncbi.nlm.nih.gov/pubmed/10768397

25. Tse JR, Engler AJ. Preparation of hydrogel substrates with tunable mechanical properties. Curr Protoc Cell Biol [Internet]. 2010 Jun [cited 2014 Jul 11];10(June):16. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20521229

26. Kandow CE, Georges PC, Janmey P a, Beningo K a. Polyacrylamide hydrogels for cell mechanics: steps toward optimization and alternative uses. Methods Cell Biol [Internet]. 2007 Jan [cited 2014 Nov 17];83(07):29–46. Available from: http://www.ncbi.nlm.nih.gov/pubmed/17613303

27. Quinlan AMT, Billiar KL. Investigating the role of substrate stiffness in the persistence of valvular interstitial cell activation. J Biomed Mater Res A [Internet]. 2012 Sep [cited 2014 Nov 21];100(9):2474–82. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3880130&tool=pmcentrez&rendertype=abstract

28. Bathe K-J. Finite Element Procedures. 2006. 195 p.

29. Rees D. Basic Engineering Plasticity: An Introduction with Engineering and Manufacturing Applications. 2012 p. 52.

30. Mase GTE, Smelser RE, Mase GTE. Continuum Mechanics for Engineers, Third Edition. 2009 p. 123.

31. Mercurieva A a., Birshtein TM. Liquid-crystalline ordering in two-dimensional systems with discrete symmetry. Die Makromol Chemie, Theory Simulations [Internet]. 1992;1(4):205–14. Available from: http://doi.wiley.com/10.1002/mats.1992.040010402

32. Mertz AF, Che Y, Banerjee S, Goldstein JM, Rosowski K a, Revilla SF. Cadherin-based intercellular adhesions organize epithelial cell–matrix traction forces. Proc Natl Acad Sci. 2013;110(3):842–7.

33. Notbohm J, Kim JH, Asthagiri a. R, Ravichandran G. Three-dimensional analysis of the effect of epidermal growth factor on cell-cell adhesion in epithelial cell clusters. Biophys J [Internet]. Biophysical Society; 2012;102(6):1323–30. Available from: http://dx.doi.org/10.1016/j.bpj.2012.02.016

34. Ng MR, Besser A, Brugge JS, Danuser G. Mapping the dynamics of force transduction at cell-cell junctions of epithelial clusters. Elife [Internet]. 2014;4:e03282. Available from: http://elifesciences.org/content/3/e03282.abstract

35. Lemmon C a, Chen CS, Romer LH. Cell traction forces direct fibronectin matrix assembly. Biophys J [Internet]. The American Society of Human Genetics; 2009 Jan [cited 2013 Sep 27];96(2):729–38. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2716473&tool=pmcentrez&rendertype=abstract

36. Zegers MMP, O'Brien LE, Yu W, Datta A, Mostov KE. Epithelial polarity and tubulogenesis in vitro. Trends Cell Biol [Internet]. 2003 Apr [cited 2014 Mar 12];13(4):169–76. Available from: http://linkinghub.elsevier.com/retrieve/pii/S0962892403000369

37. Eckhart L, Reinisch C, Inoue S, Messner P, Dockal M, Mayer C, et al. A basement membrane-like matrix formed by cell-released proteins at the medium/air interface supports growth of keratinocytes. Eur J Cell Biol [Internet]. 2003 Nov;82(11):549–55. Available from: http://www.ncbi.nlm.nih.gov/pubmed/14703011

38. Tchao R. Epithelial cell interaction in air-liquid interface culture. Vitr Cell Dev Biol [Internet]. 1989 [cited 2014 Apr 23];25(5). Available from:

http://link.springer.com/article/10.1007/BF02624633

39. Inoue S, Reinisch C, Tschachler E, Eckhart L. Ultrastructural characterization of an artificial basement membrane produced by cultured keratinocytes. J Biomed Mater Res A [Internet]. 2005 May 1 [cited 2014 Apr 23];73(2):158–64. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15754320

40. Bartosh T. Aggregation of human mesenchymal stromal cells (MSCs) into 3D spheroids enhances their antiinflammatory properties. Proc Natl Acad Sci USA [Internet]. 2010 [cited 2014 Apr 23];107:1–6. Available from: http://www.pnas.org/content/107/31/13724.short

41. Wiebe C, Wayne BG, Brodland GW. Tensile properties of embryonic epithelia measured using a novel instrument. J Biomech [Internet]. 2005 Oct [cited 2015 May 6];38(10):2087–94. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16084209

42. Last J, Liliensiek S, Nealey P, Murphy C. Determining the mechanical properties of human corneal basement membranes with atomic force microscopy. J Struct Biol [Internet]. 2009 [cited 2015 May 6];167(1):19–24. Available from: http://www.sciencedirect.com/science/article/pii/S104784770900080X

43. Roure O Du, Saez A, Buguin A, Austin RH, Chavrier P, Silberzan P, et al. Force mapping in epithelial cell migration. Proc Natl Acad Sci USA [Internet]. 2005 [cited 2014 Nov 21];102(32):2390–5. Available from: http://www.pnas.org/content/102/7/2390.short

44. Ghibaudo M, Saez A, Trichet L, Xayaphoummine A, Browaeys J, Silberzan P, et al. Traction forces and rigidity sensing regulate cell functions. Soft Matter. 2008;4(9):1836.

45. Yeung T, Georges PC, Flanagan L a., Marg B, Ortiz M, Funaki M, et al. Effects of substrate stiffness on cell morphology, cytoskeletal structure, and adhesion. Cell Motil Cytoskeleton. 2005;60(1):24–34.

46. Engler a., Sheehan M, Sweeney HL, Discher DE. Substrate compliance vs ligand density in cell on gel responses. Eur Cells Mater [Internet]. Elsevier; 2003;6(SUPPL. 1):7–8. Available from: http://dx.doi.org/10.1016/S0006-3495(04)74140-5

47. Pelham RJ. Cell Locomotion and Focal Adhesions Are Regulated by the Mechanical Properties of the Substrate. Biol Bull. 1998;(194):348–50.

48. Maruthamuthu V. Cell-ECM traction force modulates endogenous tension at cell–cell contacts. Proc Natl Acad Sci USA [Internet]. 2011 [cited 2015 Feb 19]; Available from: http://www.pnas.org/content/108/12/4708.short

49. McKee CT, Raghunathan VK, Nealey PF, Russell P, Murphy CJ. Topographic modulation of the orientation and shape of cell nuclei and their influence on the measured elastic modulus of epithelial cells. Biophys J [Internet]. Biophysical Society; 2011;101(9):2139–46. Available from: http://dx.doi.org/10.1016/j.bpj.2011.09.042

50. Raghunathan VK, McKee CT, Tocce EJ, Nealey PF, Russell P, Murphy CJ. Nuclear and cellular alignment of primary corneal epithelial cells on topography. J Biomed Mater Res - Part A. 2013;101 A(4):1069–79.

51. Zemel a., Rehfeldt F, Brown a. EX, Discher DE, Safran S a. Optimal matrix rigidity for stress-fibre polarization in stem cells. Nat Phys [Internet]. Nature

Publishing Group; 2010;6(6):468–73. Available from:
http://dx.doi.org/10.1038/nphys1613

Supporting Information Captions

FIGURE S1: Evaporation of the droplet under imaging power. Droplet height change

(blue squares) and volume change (red triangles) versus time while imaged under normal

imaging power. Imaging cut off time shown with grey line.

FIGURE S2: Deformation of the cluster outline with cytoskeletal drugs. A) Confocal

image of MDCK cells stained with Hoechst and WGA. Cluster outline positions

determined with WGA staining are shown with light blue xs (scale bar, 10m). B) Major,

minor and mean strains of cell clusters after the addition of blebbistatin, media (control)

or calyliculin A calculated from nuclear displacements (red, blue and green) and cluster

outline displacements determined with WGA (orange, light blue and light green). n=4, no

significance.

FIGURE S3: Cluster shape order parameters of clusters on glass and PA. A) Order

parameter, S_{ls}, for cluster shape and strain (defined by the angle between the major strain

axis and the cluster major axis) for blebbistatin (red), control (blue) and CalA (green)

treatments on glass and PA. B) Order parameter, S_{nl} for nuclear alignment and cluster

shape (defined by the angle between the major length axis of nuclei and the major length

axis of the cluster) for cell clusters in drops (solid), on PA (stripes) and on glass (dots).

FIGURE S4: Dependence of mean strain on cluster aspect ratio. Cluster mean strain of

droplets treated with blebbistatin (red diamonds), media (blue squares) and CalA (green

triangles) vs. the aspect ratio of the cluster. There is no significant correlation.

A Before | After

B Translation | Rotation

C E_1 E_2 E_M 0.3 0 −0.3

Control | Calyculin A | Blebbistatin

A Incubate   Inject   Image

B

C