Seed size drives species diversification across angiosperms

J. IGEA[1], E. F. MILLER[1,2], A. S. T. PAPADOPULOS[3], A. J. TANENTZAP[1]

[1] *Department of Plant Sciences, University of Cambridge, Downing St, Cambridge,*

CB2 3EA, UK

[2] *Department of Zoology, University of Cambridge, Downing St, Cambridge, CB2*

3EJ, UK

[3] *Royal Botanic Gardens, Kew, Richmond, TW9 3AB, UK.*

Species diversity varies greatly across the different taxonomic groups that

comprise the Tree of Life (ToL). This imbalance is particularly conspicuous

within angiosperms, but is largely unexplained [1]**. Seed mass is one factor that**

may help some lineages diversify more than others by influencing key life history

traits, such as dispersal, colonisation, environmental tolerance and reproductive

success [2]**. However, the extent and direction of these effects have not been**

assessed across the angiosperm ToL. Here, we show for the first time that

absolute seed size and the rate of change in seed size are both associated with

variation in diversification rates. Based on an unequalled phylogenetic tree that

included 4105 angiosperm genera, we found that smaller-seeded plants had

higher rates of diversification, possibly due to improved colonisation potential [3]**.**

The rate of phenotypic change in seed size was also strongly positively correlated

with speciation rates, supporting emerging evidence that rapid morphological

change is associated with species divergence [4]**. Our study now reveals that**

variation in morphological traits, as well as the rate at which traits evolve,

contribute significantly to the extremely uneven distribution of diversity across

the ToL.

Angiosperms are one of the most species-rich clades on Earth and have dominated

terrestrial plant communities since the Late Cretaceous [5]. The astounding diversity of

flowering plants is, however, extremely unevenly distributed across the ToL. Each of

the five most species-rich angiosperm families contains >10,000 species while more

than 200 families contain <100 species each. An enduring pursuit in evolutionary

biology is to explain this uneven distribution of biodiversity, not only in angiosperms,

but also across the whole ToL [1].

Seed mass influences individual life history in ways that can ultimately shape

angiosperm diversification [6]. Along with adult plant size, seed mass affects survival,

reproductive lifespan and dispersal [7]. These traits contribute to fitness and adaptation,

which are the ultimate determinants of whether lineages diversify or go extinct [8]. In

support of this idea, seed mass has been shown to correlate negatively with

diversification in the Polygoneaceae [9], but this has not been investigated across large

taxonomic scales. As seed mass varies over ten orders of magnitude in angiosperms,

from the minute 1 μg seeds of some orchids to the >18 kg seeds of the sea coconut

Lodoicea maldivica, this huge variation may coincide with variation in species

diversity. Generalising the direction and magnitude of an effect of seed mass on

diversification across taxonomic scales has however proved difficult. Some life

history traits encapsulated by seed mass are expected to promote speciation or

extinction, while others may simultaneously counteract such effects.

The rate of change in key life history traits such as seed size can be as important in

driving macroevolutionary dynamics as the absolute values of the traits themselves [4].

This is because phenotypic divergence may cause reproductive isolation that results in

speciation [10]. Nevertheless, few empirical studies have detected a correlation between

rates of phenotypic evolution and lineage diversification [11]. This correlation may be

expected where a trait can change more rapidly in some species than others in

response to selective pressures (i.e. high "evolvability" [12]); this enables greater access

to new ecological niches or quicker establishment of reproductive isolation, and

thereby increasing the rate of speciation (λ) [13]. In the case of seed mass, the ability to

switch rapidly, for example, from small seeds with high dispersal ability to larger

seeds with lower dispersal ability, might promote cycles of rapid colonisation and

isolation or permit adaptation to new dispersal vectors in novel environments. Rapid

evolution of new phenotypes may also allow individuals to escape harsh

environmental conditions and competitive interactions [14], thereby decreasing

extinction rates (μ). The net outcome of these processes on diversification ($r = \lambda-\mu$)

will ultimately depend upon which of these rates responds more strongly to

phenotypic change.

Here we show for the first time that both seed mass and its phenotypic rate of

evolution influence speciation and extinction across the angiosperm ToL. Our

approach combined the most comprehensive phylogenetic timetree available [15] with

an unparalleled dataset of seed mass measurements to obtain mean seed mass values

for 24% of all described angiosperm genera (n = 4105). We estimated rates of

speciation and extinction using Bayesian Analysis of Macroevolutionary Mixtures

(BAMM) [16], which models rate heterogeneity through time and lineages, and accounts

for incomplete taxon sampling. Similarly, we analysed the rates of change in seed size

across the phylogeny. We then estimated the correlations of the macroevolutionary

dynamics with absolute seed size and the rate of seed size evolution.

As expected, given the high degree of taxonomic imbalance observed in the

angiosperm phylogeny, we found strong support for multiple shifts in the rates of

diversification. The median number of rate shifts for the speciation/extinction analysis

was 123 (95% confidence interval (CI): 95-155) and 160 (95% CI: 62-110) for the

seed size evolution analysis (Extended Data Fig. 1). There was marked heterogeneity

in the rates of seed size evolution (Fig. 1), which varied over two orders of magnitude

(Extended Data Fig. 2). We then estimated whether shifts in macroevolutionary

dynamics (λ, μ and r) were significantly correlated with absolute mean genus seed

size and rates of seed size evolution by comparing the existing correlations to a null

distribution of correlations using STructured Rate Permutations on Phylogenies

(STRAPP), which is robust to phylogenetic pseudoreplication (see Methods for

details) [17].

For the first time, we were able to explain major differences in diversity across the

angiosperm ToL with a single trait that integrates multiple aspects of life history. We

specifically found evidence that increased speciation was associated with smaller seed

sizes (Spearman's ρ = -0.22, p-value = 0.016; Fig. 2a). Increased extinction rate was

similarly associated with smaller seeds (ρ = -0.20, p-value = 0.045), but given its

relatively weaker effect, the net outcome of λ-μ was that diversification rates

increased with decreasing seed size (ρ =-0.19, p-value = 0.049). We also identified a

stronger positive association between the rate of seed size evolution and both

speciation and extinction (ρ = 0.48, p-value < 0.0001 and ρ = 0.37, p-value = 0.003,

respectively; Fig. 2b). Again, as the effect of speciation was greater than that of

extinction, rates of diversification and phenotypic change were positively correlated

(ρ = 0.46, p-value = 0.0002; Fig. 2b). Generally, the observed correlations arose from

many phenotypically fast-evolving clades distributed across the phylogeny (Extended

Data Fig. 2).

Our results were unaffected by intra-generic variation in seed mass for two reasons.

First, there was no systematic bias in intra-generic variation across the phylogenetic

tree. We detected no correlation between the mean and the coefficient of variation

(CV) for seed mass of each genus (Extended Data Fig. 3, PGLS: $F_{1,131}$ = 0.67, p-value

= 0.416). Second, we could disregard intra-generic variation in rates of seed mass

evolution as influencing our results as we found that genera with larger variation in

seed size did not have different macroevolutionary dynamics (Extended Data Fig. 4,

Spearman's ρ_λ = -0.05, p-value = 0.114; ρ_μ = -0.03, p-value = 0.406; ρ_r = -0.067, p-

value = 0.058).

Our study supports the idea that variation in seed mass can explain disparity in

diversification across the angiosperm ToL by influencing dispersal and habitat

interactions. We specifically found that smaller-seeded genera had faster speciation

rates. This may be because they are capable of dispersing over larger distances [3] that

can result in isolated populations and eventually lead to speciation [18]. Large dispersal

distances may be especially important for isolation to occur in continuous habitats.

However, the positive effect on speciation from long-distance dispersal may be

dampened in highly fragmented landscapes where isolation is more common over

shorter distances [19]. Dispersal syndromes may also alter the effect of seed size on

speciation. Species with larger seeds are generally associated with biotic dispersal that

distributes seeds over greater distances than wind or gravity dispersal [7]. However,

broad scale predictions on the effects of dispersal syndromes on diversification may

be inaccurate, since the former depend on landscape connectivity [20] and can

sometimes be inconsistent, e.g. a wind-dispersed seed might be transported by an

animal. Detailed contextual data will be necessary to expand upon the mechanisms

underlying our findings. We also found that smaller-seeded genera had higher rates of

extinction, possibly due to smaller nutritional reserves that constrain establishment,

environmental tolerance and access to limiting resources [21].

Seed mass is associated with other traits that can affect diversification, but there is

little evidence that these better explain our observed correlations. For example,

genome size positively correlates with seed mass [22] and faster rates of genome size

evolution have been linked to increased speciation in angiosperms [23]. Shorter,

smaller-seeded plants also tend to have faster life cycles, which may accelerate

mutation rates [24] and promote diversification [25]. By comparing the effects of genome

size and life cycle across a subset of >900 genera in our dataset, we found that only

the distinction between strictly annual versus perennial genera influenced

macroevolutionary dynamics in a similar way as seed size (Supplementary

Information, Extended Data Fig. 5). Unlike other traits [23] both absolute seed size as

well as its rate of change were correlated with speciation and extinction. Thus, while

we cannot exclude unobserved traits as drivers of diversification [26], we argue that seed

151 mass plays a central role in plant life history both on its own and by integrating across

152 traits that should predictably shape macroevolutionary dynamics (Extended Data Fig.

153 6).

155 Our finding that high rates of phenotypic change correlate with diversification (Fig.

156 2b) has recently been observed in other taxonomic groups [4,11], but never across the

157 whole of the angiosperm ToL. Accelerated morphological evolution may allow

158 radiating lineages to occupy more complex adaptive landscapes [27]. Similarly, species

159 with greater rate of change in their seed mass (i.e., higher evolvability) could shift

160 between adaptive peaks or develop reproductive barriers more rapidly. However,

161 current methods do not allow us to distinguish whether speciation is responding to

162 morphological change or vice versa when reconstructing 250 million years of

163 evolutionary history [4].

165 The approach applied here can help to unravel the processes responsible for

166 generating large-scale asymmetries in biodiversity. It also offers the potential to test

167 how widely-varying traits influence other aspects of the evolution and adaptation of

168 flowering plants (e.g. [15]). Clade-specific exceptions arising from local interactions

169 with non-focal traits [29] and specific spatio-temporal contexts will undoubtedly interact

170 with broad-scale macroevolutionary patterns and may modulate the effects of seed

171 mass on diversification. Regardless, our results clearly demonstrate that seed size, and

172 its rate of change, drive speciation and extinction and help to explain why some

173 clades are much more species-rich than others.

175 **Methods**

Seed mass and phylogenetic dataset

Seed mass data for 31,932 species were obtained from the Royal Botanic Gardens Kew Seed Information Database [30]. Species names were standardised with The Plant List (TPL) nomenclature [31] and cleaned using the *Taxonstand* R package [32]. Further processing at the genus-level was carried out with the *taxonlookup* R package [33], which is a complete genus-family-order mapping for vascular plants that draws from TPL, the Angiosperm Phylogeny website [34] and a higher-level manually curated taxonomic lookup [15]. Seed mass mean values for each genus were calculated for a total of 4763 genera.

We used the most comprehensive phylogenetic tree for land plants [15,35] that comprises 31,389 species. Taxonomic information for our phylogenetic tree was run through *Taxonstand* and *taxonlookup* as described above to make it as comparable as possible to the seed mass dataset. Monophyly of the genera in the tree was assessed using the *Monophy* package [36] The initial estimate was 16%, but we removed taxa classified as outliers by *Monophy* (i.e., taxa that lay outside of an established "core clade" for each genus) and this resulted in only 8% of the genera not being monophyletic. The *Phyndr* [33] package was then used to generate a genus-level tree with as much overlap as possible between the phylogeny and the trait data. The final phylogenetic tree included representatives from 303 plant families (Extended Data Fig. 7).

Diversification and phenotypic evolution analysis

Speciation, extinction and net diversification rates and rates of seed size evolution were estimated separately on the phylogeny using BAMM version 2.5.0 [16]. BAMM

models shifts in macroevolutionary regimes across a phylogenetic tree using

reversible-jump Markov chain Monte Carlo (rjMCMC) sampling. Initial prior settings

were calculated with the *setBAMMpriors* function in *BAMMtools* [37], and the

expectedNumberOfShifts parameter was set at 25 and 10 for the speciation/extinction

and trait evolution analyses, respectively. We incorporated non-random incomplete

sampling information following BAMM protocols by calculating the proportion of

genera sampled inside each family and estimated the backbone sampling as the

overall proportion of sampled genera. *Taxonlookup* was used as a reference for these

calculations.

All analyses were run for 50 million generations. We verified convergence by plotting

chain traces and ensuring that the effective sample sizes of all relevant parameters

exceeded 200. The first 10 million generations were discarded as burn-in.

Correlation of diversification and trait evolution

We used STRAPP to test for multiple associations between macroevolutionary

dynamics and each of seed mass (using genus mean values as character states) and

seed mass rate of evolution (using seed mass evolutionary rates at the tips of the

phylogeny as character states). STRAPP compares the correlation between a focal

trait and a macroevolutionary parameter (λ, μ or r) to a null distribution of

correlations. The null correlations are generated by permuting the evolutionary rates

in the tips of the phylogenetic tree while maintaining the location of rate shift events

in the phylogeny. In each case, we calculated the absolute difference between the

observed correlation of the macroevolutionary rate and the trait state and the null

correlation obtained by the structured permutations across 5000 samples from the

BAMM posterior. The reported p-value was the proportion of replicates where the

null correlation coefficient was greater than the observed correlation. We found a low

type I error associated with our STRAPP correlation analysis (p-value = 0.11,

Extended Data Fig. 8).

Code availability

Scripts used to carry out the analysis described in the paper and generate the figures

will be deposited in Github.

<u>References</u>

1.   Davies, T. & Barraclough, T. in *Reconstructing the Tree of Life: Taxonomy and*
Systematics of Species Rich Taxa **72,** 149–163 (2007).
2.   Willson, M. & Traveset, A. in *Seeds: the ecology of regeneration in plant*
communities. 85–110 (2000). doi:10.1146/annurev.es.13.110182.001221
3.   Coomes, D. A. & Grubb, P. J. Colonization, tolerance, competition and seed-
size variation within functional groups. *Trends Ecol. Evol.* **18,** 283–291 (2003).
4.   Rabosky, D. L. *et al.* Rates of speciation and morphological evolution are
correlated across the largest vertebrate radiation. *Nat. Commun.* **4,** 1958 (2013).
5.   Crane, P. R., Friis, E. M. & Pedersen, K. R. The origin and early diversification
of angiosperms. *Nature* **374,** 27–33 (1995).
6.   Moles, A. T. *et al.* A Brief History of Seed Size. *Science (80-.).* **307,** 576–580
(2005).
7.   Moles, A. & Leishman, M. in *Seedling ecology and evolution* (eds. Leck, M.
A., Parker, V. T. & Simpson, R. L.) 215–235 (Cambridge University Press,
2008).
8.   Salguero-Gómez, R. *et al.* Fast-slow continuum and reproductive strategies
structure plant life-history variation worldwide. *Proc. Natl. Acad. Sci. U. S. A.*
113, 230–5 (2016).
9.   Kostikova, A., Salamin, N. & Pearman, P. B. The role of climatic tolerances
and seed traits in reduced extinction rates of temperate polygonaceae.
Evolution (N. Y). **68,** 1856–1870 (2014).
10.   Coyne, J. A. & Orr, H. A. *Speciation.* (Sinauer Associates, 2004).
11.   Price, S. L., Etienne, R. S. & Powell, S. Tightly congruent bursts of lineage and
phenotypic diversification identified in a continental ant radiation. *Evolution*
(2016). doi:10.1111/evo.12894
12.   Pigliucci, M. Is evolvability evolvable? *Nat. Rev. Genet.* **9,** 75–82 (2008).
13.   Ricklefs, R. E. & Renner, S. S. Species Richness Within Families of Flowering
Plants. *Evolution (N. Y).* **48,** 1619–1636 (1994).
14.   Rees, M. & Westoby, M. Game-Theoretical Evolution of Seed Mass in Multi-
Species Ecological Models. *Oikos* **78,** 116–126 (1997).

15.   Zanne, A. E. *et al.* Three keys to the radiation of angiosperms into freezing
environments. *Nature* **506,** 89–92 (2014).
16.   Rabosky, D. L. Automatic detection of key innovations, rate shifts, and
diversity-dependence on phylogenetic trees. *PLoS One* **9,** e89543 (2014).
17.   Rabosky, D. L. & Huang, H. A Robust Semi-Parametric Test for Detecting
Trait-Dependent Diversification. *Syst. Biol.* **65,** 181–93 (2016).
18.   Kisel, Y. & Barraclough, T. G. Speciation has a spatial scale that depends on
levels of gene flow. *Am. Nat.* **175,** 316–34 (2010).
19.   Claramunt, S., Derryberry, E. P., Remsen, J. V & Brumfield, R. T. High
dispersal ability inhibits speciation in a continental radiation of passerine birds.
Proc. Biol. Sci. **279,** 1567–74 (2012).
20.   Givnish, T. J. Ecology of plant speciation. *Taxon* **59,** 1326–1366 (2010).
21.   Leishman, M. R., Wright, I. J., Moles, A. T. & Westoby, M. in *The Ecology of*
Regeneration in Plant Communities 31–58 (2000).
22.   Beaulieu, J. M. *et al.* Correlated evolution of genome size and seed mass. *New*
Phytol. **173,** 422–37 (2007).
23.   Puttick, M. N., Clark, J. & Donoghue, P. C. J. Size is not everything: rates of
genome size evolution, not C-value, correlate with speciation in angiosperms.
Proc. Biol. Sci. **282,** 20152289– (2015).
24.   Lanfear, R., Ho, S. Y. W., Love, D. & Bromham, L. Mutation rate is linked to
diversification in birds. *Proc. Natl. Acad. Sci. U. S. A.* **107,** 20423–8 (2010).
25.   Lanfear, R. *et al.* Taller plants have lower rates of molecular evolution. *Nat.*
Commun. **4,** 1879 (2013).
26.   Beaulieu, J. M. & O'Meara, B. C. Detecting hidden diversification shifts in
models of trait-dependent speciation and extinction. *Syst. Biol.* syw022–
(2016). doi:10.1093/sysbio/syw022
27.   Lovette, I. J., Bermingham, E. & Ricklefs, R. E. Clade-specific morphological
diversification and adaptive radiation in Hawaiian songbirds. *Proc. Biol. Sci.*
269, 37–42 (2002).
28.   Eldredge, N. & Gould, S. J. in *Models in Paleobiology* (ed. Schopf, T.) 82–115
(Freeman Cooper, 1972).
29.   Donoghue, M. J. & Sanderson, M. J. Confluence, synnovation, and
depauperons in plant diversification. *New Phytol.* **207,** 260–274 (2015).
30.   Kew Seed Information Database. *Royal Botanic Gardens Kew* (2016). at
<http://data.kew.org/sid/>
31.   The Plant List. Version 1.1. (2013). at <http://www.theplantlist.org/>
32.   Cayuela, L., Granzow-de la Cerda, Í., Albuquerque, F. S. & Golicher, D. J.
taxonstand: An r package for species names standardisation in vegetation
databases. *Methods Ecol. Evol.* **3,** 1078–1083 (2012).
33.   Pennell, M. W., FitzJohn, R. G. & Cornwell, W. K. A simple approach for
maximizing the overlap of phylogenetic and comparative data. *Methods Ecol.*
Evol. n/a–n/a (2015). doi:10.1111/2041-210X.12517
34.   Stevens, P. . Angiosperm Phylogeny Website. *Angiosperm Phylogeny Website.*
Version 12 (2012). at <http://www.mobot.org/MOBOT/research/APweb/>
35.   Qian, H. & Jin, Y. An updated megaphylogeny of plants, a tool for generating
plant phylogenies and an analysis of phylogenetic community structure. *J.*
Plant Ecol. rtv047 (2015). doi:10.1093/jpe/rtv047
36.   Schwery, O. & O'Meara, B. C. MonoPhy: A simple R package to find and
visualize monophyly issues. (2015). doi:10.7287/peerj.preprints.1600v1
37.   Rabosky, D. L. *et al.* BAMMtools: an R package for the analysis of

evolutionary dynamics on phylogenetic trees. *Methods Ecol. Evol.* **5,** 701–707
(2014).
38. Bennett, M. & Leitch, I. J. Plant DNA C-values database (release 6.0, Dec
2012). (2012). at <http://www.kew.org/cvalues/>
320
321

Acknowledgments

We thank D. A. Coomes, A. J. Helmstetter, T. Jucker and W. G. Lee for useful

comments that helped improve the manuscript. J.I. and A.J.T. thank the Gatsby

Charitable Foundation, Wellcome Trust and Newton Trust for funding. E.F.M was

funded by the BBSRC DTP at the University of Cambridge.

Author contributions

J.I and A.S.T.P conceived the study. J.I. and E.F.M. performed the analysis. J.I. and

A.J.T interpreted the analysis and wrote the manuscript. All authors edited the

manuscript.

Author information

Scripts used to carry out the analyses described in the manuscript will be uploaded to

Github. Correspondence and requests for materials should be sent to J.I.

(ji247@cam.ac.uk)

Figure Legends

Figure 1. Seed mass and macroevolutionary rates across the angiosperm tree of life**.**

Phylogenetic tree of 4105 genera of plants with seed mass, rate of seed mass change,

and speciation (λ), extinction (μ) and net diversification (r) rates estimated by

BAMM. Seed mass and rate data were standardised to Z-scores so that variation could

be directly compared.

Figure 2. Correlation of seed mass and seed mass rate with macroevolutionary

dynamics. STRAPP correlations between speciation (λ), extinction (μ), and net

diversification (r) and each of a) seed mass and b) rate of seed mass change. Coloured

lines are Spearman correlations between λ (shown in red), μ (blue), r (purple) and

each trait for one sample of the BAMM posterior distribution, bold line is the median.

The insets show the density plots of the absolute difference between the observed and

null correlation calculated across 5000 structured permutations of the evolutionary

rates on the phylogenetic tree (myr = million years).

Extended Data Figure Legends

Extended Data Fig. 1. Posterior distribution of the number of rate shifts in BAMM

for a) the speciation/extinction and b) phenotypic evolution analyses.

Extended Data Fig. 2. Phylogenetic tree of 4105 angiosperm genera with branches

scaled by diversification rate as determined by BAMM. The colours of the branches

indicate the rate of seed mass evolution, also estimated with BAMM.

Extended Data Fig. 3. Mean genus seed mass and coefficient of variation for 2613

angiosperm genera. The two variables were not significantly correlated (PGLS test,

$F_{1,131} = 0.67$, p-value = 0.416).

Extended Data Fig. 4. Correlation of coefficient of variation in seed mass with

macroevolutionary dynamics. STRAPP correlations with speciation (λ), extinction (μ)

and net diversification (r) and the genus-level coefficient of variation in seed mass for

2613 angiosperm genera (i.e. each with seed mass data for >1 species in the Kew

SID). Correlations and insets calculated as in Fig. 2.

Extended Data Fig. 5. STRAPP correlations of diversification and phenotypic traits

for 938 angiosperm genera. The distribution of the absolute difference in the observed

correlation minus the null correlation is plotted for each trait. The coloured dotted

lines indicate the mean of that distribution, and the black dotted line indicates 0; a

distribution with mean = 0 would show no association between a focal trait and

speciation. STRAPP correlation of seed mass (shown in blue), C-value (shown in red)

and life cycle (shown in green) with a) speciation rate (λ), b) extinction rate (μ), and

c) net diversification rate (r).

Extended Data Fig. 6. Proposed effects of seed mass and other life history traits on

diversification (solid lines). Dashed lines indicate correlations between life history

traits. Numbers indicate reference where the link is proposed.

Extended Data Fig. 7. Phylogenetic tree of 303 angiosperm families with

representatives in this study. The red bars indicate the levels of sampling for each

family.

Extended Data Fig. 8. Type I error analysis. We estimated the type I error rate of our

analysis by simulating neutral traits on the angiosperm phylogenetic tree. We

performed 1000 simulations and then ran 1000 STRAPP tests with each simulated

dataset. We estimated the corresponding p-values for the association between traits

and diversification and calculated the type I error as the proportion of datasets where

a significant association (p-value < 0.05) was detected.

Extended Data Fig. 9. Mean genus seed mass of strict annual (n = 106) and perennial

(n = 832) genera. No significant difference between the means of the two groups was

found when phylogeny was accounted for (PGLS, $F_{1,937}$ < 0.01, p-value = 0.972).

Extended Data Table 1. STRAPP correlations for 938 genera of angiosperms with

seed mass, genome size (i.e., C-value) and life cycle data. Macroevolutionary

parameters were obtained from an analysis with 4105 genera (see Main text).

Supplementary Information

Seed mass is central to a network of inter-correlated life history traits that can impact

diversification. Two such traits are genome size or plant C-value (measured as

picograms of DNA per haploid nucleus) and life cycle. We compared the correlation

between macroevolutionary parameters (λ, μ, r) and each of seed mass, C-value and

life cycle across a dataset of 938 plant genera (106 strictly annuals and 832 strictly

perennials). Genome content data and life cycle data were downloaded from the Plant

DNA C-values database [38]. Surprisingly, mean seed mass did not differ between

strictly annual and perennial plants when accounting for phylogenetic relationships

(Extended Data Fig. 9, PGLS: $F_{1,937}$ <0.01, p-value = 0.972).

STRAPP requires large phylogenetic trees (n > 1000 species) to detect correlations [17].

In this reduced dataset, we ran STRAPP correlations for each focal trait with the

diversification parameters calculated from our 4105 genera BAMM analysis. We then

calculated the absolute differences in the observed and the null correlations between

the macroevolutionary parameters and seed mass, C-value and "annuality" (a binary

variable specifying whether the genus was annual or not). These differences were

similar and positive for seed mass and life cycle, while smaller for C-value (Extended

Data Fig. 5). However, with this reduced dataset, only life-cycle showed significant

correlations with macroevolutionary dynamics (Extended Data Table 1).   These

correlations were weaker than those observed with seed mass in the main text (Fig. 2),

and considerably weaker than correlations with the rate of seed size evolution.

Figure 1