3-Keto-Steroid transactivation of mineralocorticoid receptors from Amur sturgeon (Acipenser schrenckii) and tropical gar (Atractosteus tropicus)

Akira Sugimoto[1], Kaori Oka[1], Rui Sato[1], Shinji Adachi[2],

Michael E. Baker[3, *], Yoshinao Katsu[1, 4, *]

[1]Graduate School of Life Science, Hokkaido University, Sapporo, Japan; [2]Faculty of Fisheries

Sciences, Hokkaido University, Hakodate, Japan; [3]Department of Medicine, University of

California, San Diego, CA, USA; [4]Department of Biological Sciences, Hokkaido University,

Sapporo, Japan

*Corresponding author:

Michael E. Baker: mbaker@ucsd.edu

Yoshinao Katsu; ykatsu@sci.hokudai.ac.jp

Abstract

We investigated the response to a panel of steroids of the mineralocorticoid receptor (MR) in
Amur sturgeon and tropical gar, two basal ray-finned fish, whose ancestors evolved before
zebrafish and other teleosts.   Half-maximal responses (EC50s) for transcriptional activation of
sturgeon MR by 11-deoxycorticosterone, corticosterone, 11-deoxycortisol, cortisol and
aldosterone, and progesterone were between 13 pM and 150 pM.   Gar MR EC50s were between
8 pM and 55 pM.   Such low EC50s support regulation of MR activity by these steroids in
sturgeon and gar.   Companion studies with human MR and zebrafish MR found higher EC50s
compared to EC50s for sturgeon and gar MR, with EC50s for zebrafish MR closer to gar and
sturgeon MR than was human MR.   Zebrafish MR EC50s were between 75 pM and 740 pM; for
human MR, EC50s were between 65 pM and 2 nM.   Unexpectedly, progesterone was an agonist
for all ray-finned fish MRs, in contrast to its antagonist activity for human MR.   Sturgeon, gar and
zebrafish MRs contain a serine corresponding to serine-810 in human MR, hypothesized as critical
for progesterone antagonism.   Progesterone is an agonist forSer810Leu MR.   Our data suggests
an alternative mechanism for progesterone as an MR agonist in these three ray-finned fishes.

Keywords Amur sturgeon, tropical gar, mineralocorticoid receptor, mineralocorticoid evolution

Short title: ancient fish mineralocorticoid receptors

1. Introduction

The mineralocorticoid receptor (MR) is a transcription factor that belongs to the nuclear receptor family, a diverse group of transcription factors that also includes receptors for androgens (AR), estrogens (ER), glucocorticoids (GR) and progestins (PR), and other small lipophilic ligands, such as thyroid hormone and retinoids, as well as orphan receptors with no known ligand [1-7]. The MR and GR are descended from a common corticosteroid receptor (CR), which has descendants in jawless fish, such as lampreys and hagfish [7-9]. Several corticosteroids (Figure 1), including aldosterone (Aldo), cortisol (F), 11-deoxycortisol (S), corticosterone (B) and 11-deoxycorticosterone (DOC), as well as progesterone (Prog), are transcriptional activators of Atlantic sea lamprey CR and hagfish CR [8]. Among these steroids, Aldo, the main physiological activator of the MR in human and other terrestrial vertebrates [10-13], had the lowest half-maximal response (EC50) for transcriptional activation of the CR. This strong response to Aldo is surprising because Aldo is not found in either lamprey or hagfish serum [8]. S, which along with DOC is present in Atlantic sea lamprey serum, has been found to have mineralocorticoid activity in lamprey [14].

Distinct MR and GR genes first appear in cartilaginous fishes (Chondrichthyes), such as sharks, rays and skates [8, 15]. Carroll et al. [15] determined EC50s of several corticosteroids for skate MR; EC50s were 70 pM for Aldo, 30 pM for DOC, 90 pM for B, 1 nM for F and 22 nM for S. In teleosts, which comprise about 95% of known ray-fish species (*Actinopterygii*), corticosteroid activation of the MR has been investigated for cichlid [16], trout [17], carp [18], midshipman fish [19] and zebrafish [20], with Aldo, F and DOC being the principal steroids that were studied. Although Aldo has not been found in teleost fish [21], Aldo has a low EC50 for teleost MRs, similar to that for Aldo activation of lamprey CR and skate MR. DOC also has a low EC50 for teleost MRs, and DOC has been proposed as mineralocorticoid in fish [17, 22-25]. F also has been proposed to be ligand for teleost fish MR [23-25]. The response of the teleost MRs to B and S, which are found in fish [25, 26], has been studied only in trout, in which the EC50s are 10 nM for B and 3.7 nM for S [17]. These studies indicate that several corticosteroid(s) are potential transcriptional activators of teleost MRs [23, 25, 27, 28].

An important gap in our understanding of the evolution of selectivity of ray-finned fish MRs for steroids is the absence of data on the MR in Chondrostei (sturgeons, paddlefishes, reedfishes, bichirs) and Holostei (bowfins, gars), which evolved before a whole-genome duplication that occurred in Teleostei (trout, ,medaka, tilapia, carp, zebrafish) (Figure 2) [29, 30].

Figure 1. Structures of potential steroid regulators of fish MR.

Aldo, the physiological ligand for terrestrial vertebrate MRs, is not found in fish [21].   F and

DOC have been proposed to be mineralocorticoids in teleosts [23, 25].   S is a ligand for

corticosteroid receptor in lamprey [14].   Progesterone is an antagonist for human MR [31].

Figure 2. Phylogenetic relationship of sturgeon and gar MRs to other vertebrates.
To investigate the relationship of sturgeon and gar to other fish, we constructed a phylogenetic
tree of the steroid-binding domains on MRs in sturgeon, gar, selected teleosts, elasmobranchs
and tetrapods.   The phylogenetic tree was constructed using the maximum likelihood with
JTT+G model with 1000 bootstrap replications, which are shown as percentages at the nodes of
the tree.

        Our interest in the evolution of steroid hormone action [6, 7, 32] prompted us to
investigate transcriptional activation of the MR from Amur sturgeon, *Acipenser schrenckii*, and
tropical gar, *Atractosteus tropicus* by a broad panel of corticosteroids (Aldo, F, B, DOC, S) and
Prog.   To gain further insight into the evolution of steroid specificity teleost and in tetrapod
MRs, we compared our results with companion studies of zebrafish and human MRs.   In
agreement with studies of teleost MRs, we find that Aldo and DOC have the lowest EC50
(highest activity) for sturgeon and gar MRs.   However, we also find that S, B, F, and Prog have
low EC50s, consistent with these steroids also having a physiological role as ligands for these
MRs.   In comparison, zebrafish MR also has a strong response to Aldo and DOC and a good
response to B, F, S and Prog, while human MR has strong response to Aldo, DOC and B and a
good response to F and S, and a weak response to Prog.   The weak response to Prog of human
MR is in agreement with other studies [31, 33].   Thus, the strong response to Prog of sturgeon,
gar and zebrafish MR is unexpected because the basis for the low response of human MR is
thought to be due to the presence of Ser-810 on α-helix 5 [31, 34, 35].   Prog is an agonist for
human MR with Ser810Leu mutation [31, 34, 35].   Sturgeon, gar and zebrafish MRs contain a
serine corresponding to serine-810 in human MR, suggesting an alternative mechanism for Prog
acting as an MR agonist in these three ray-finned fishes.

The transcriptional response of ray-finned fish MR to five corticosteroids and progesterone also suggests one or more mechanisms for selectively regulating steroid activation of transcription of sturgeon, gar and zebrafish MRs. Potential mechanisms include tissue specific expression of enzymes that inactivate corticosteroids, such as found for selective inactivation of F and B by 11β-hydroxysteroid dehydrogenase-type2 (11β-HSD2) in mammalian kidney [10, 36-38] and enzymes that selectively synthesize a steroid [7, 39-42], selective sequestration of 3-keto-steroids by serum proteins [43-45], steroid-specific conformational changes in the MR that regulate binding of co-activators, [46-50], inter-domain interactions [20, 50, 51], as well as post-translational modifications [52-54] of the MR that alter steroid specificity.

2. Methods

2.1. Animals and chemical reagents

Amur sturgeon and tropical gar were obtained as described previously [55]. All experimental procedures involving live fish followed the policies and guidelines of the Hokkaido University Animal Care and Use Committee. Aldosterone (Aldo), corticosterone (B), cortisol (F), 11-deoxycortisol (S), 11-deoxycorticosterone (DOC), progesterone (Prog), 5α-dihydrotestosterone (DHT), and 17β-estradiol (E2) were purchased from Sigma-Aldrich. For the reporter gene assays, all hormones were dissolved in dimethyl-sulfoxide (DMSO) and the final concentration of DMSO in the culture medium did not exceed 0.1%.

2.2. Molecular cloning of mineralocorticoid receptors

Two conserved amino acid regions, GCHYGV and LYFAPD of vertebrate MRs were selected and degenerate oligonucleotides were used as primers for PCR. First-strand cDNA was synthesized from 2 µg of total RNA isolated from the liver after amplification, and an additional primer set (CKVFFK and LYFAPD) was used for the second PCR. The amplified DNA fragments were subcloned with TA-cloning plasmid pGEM-T Easy vector (Promega, Madison, WI), sequenced using a BigDye terminator Cycle Sequencing-kit (Applied Biosystems, Foster City, CA) with T7 and SP6 primers, and analyzed on the 3130 Genetic Analyzer (Applied Biosystems). The 5'- and 3'-ends of the mineralocorticoid receptor cDNAs were amplified by rapid amplification of the cDNA end (RACE) using a SMART RACE cDNA Amplification kit (BD Biosciences Clontech, Palo Alto, CA).

2.3. Database and sequence analysis

MRs for phylogenetic analysis were collected with Blast searches of GenBank. A phylogenetic tree for MRs was constructed by the Neighbor-Joining Method [56] after sequences

were aligned by MUSCLE [57] using several fish, frog, alligator, chicken, rat, mouse, human MRs.   Maximum likelihood (ML) analysis was conducted using the JTT+G model.   Statistical confidence for each branch in the tree was evaluated by the bootstrap method [58] with 1000 replications.   We used the MEGA5 program [59] for these analyses.

2.4. Construction of plasmid vectors

Full-coding regions of mineralocorticoid receptors were amplified by PCR with KOD DNA polymerase (TOYOBO Biochemicals, Osaka, Japan).   PCR products were gel-purified and ligated into pcDNA3.1 vector (Life Technologies, Carlsbad, CA).   Mouse mammary tumor virus-long terminal repeat (MMTV-LTR) was amplified from pMSG vector (Pharmacia) by PCR, and inserted into pGL3-basic vector containing the *Photinus pyralis* lucifease gene (Promega). All constructs were verified by sequencing.

2.5. Transactivation Assay

Human embryonic kidney 293 (HEK293) cells were used in the reporter gene assay. Transfection and reporter assays were carried out as described previously [32, 60], except that we used PEI-max as transfection reagent [61].   All transfections were performed at least three times, employing triplicate sample points in each experiment.   The values shown are mean ± SEM from three separate experiments, and dose-response data and EC50 were analyzed using GraphPad Prism.

2.6. Statistical methods

Results are presented as mean ± SE (SEM) from three separate experiments.   All multi-group comparisons were performed using one-way ANOVA followed by Bonferroni test. Dose-response data and EC50 were analyzed using GraphPad Prism. $P < 0.05$ was considered statistically significant.

3. Results

3.1. Isolation of mineralocorticoid receptors from sturgeon and gar

We cloned sturgeon MR cDNA containing an open reading frame encoding 953 amino acids (GenBank accession LC149818)], and gar MR cDNA containing an open reading frame encoding 987 amino acids (GenBank accession LC149819).   Sturgeon and gar MR sequences can be divided into four domains (Figure 3).   The overall amino acid identity between these two MRs was 72%, with particularly high sequence identities for the DBD (100%) and LBD (89%) (Figure 3).   Comparison of sturgeon MR with five other species (human, chicken, alligator,

Xenopus, and zebrafish) revealed that sturgeon MR had identities of 44-36% in A/B

domains,100-95% in DBDs, 67-47% in D domains, and 90-74% in LBDs   (Figure 3).

3.2. Phylogenetic analysis of ancient fish corticoid receptors

To investigate the evolutionary position of gar and sturgeon MR in relationship to other

fish MRs and tetrapods, we collected MR sequences from several teleosts, skates and elephant

shark and selected terrestrial vertebrates.   Consistent with the evolution of Acipenseriformes

and Holostei, phylogenetic analysis places sturgeon and gar MRs close to the base of ray-finned

fish (Figure 2).

Figure 3. Comparisons of functional domains in sturgeon, gar, zebrafish, *X. laevis*, alligator,

chicken and human MRs.

Comparison of the domains in sturgeon MR gar, zebrafish, *X. laevis*, alligator, and human MR

MR.   The functional A/B domain, C domain, D domain and E domain are schematically

represented with the numbers of amino acid residues at each domain boundary indicated.   The

percentage of amino acid identity between domains is depicted.   GenBank accession numbers

are: LC149818 for sturgeon MR; LC149819 for gar MR; NM_001100403 for zebrafish MR;

NM_001090605 for *Xenopus* MR; AB701406 for alligator MR; and NM_000901 for human MR.

3.3. Strong response to 3-keto-steroids by sturgeon and gar mineralocorticoid receptors

We examined the steroid-inducible transcriptional activation of gar and sturgeon MRs

using MMTV-driven reporter construct [32, 62].   For comparison, we also examined

transcriptional activation of human MR and zebrafish MR.   At 1 nM, Aldo, B, S, DOC, F and

Prog were strong inducers of luciferase activation by gar MR and sturgeon MR and by zebrafish

MR, with the exception of Prog which had a lower signal.   These MRs show little stimulation

by 1 nM DHT and E2 (Figure 4).   At 1 nM, Aldo, B, DOC were strong transcriptional activators

of human MR, which was activated to a lesser extent by S, and weakly activated by Prog (Figure

4).

Steroid Concentration = 1 nM

Figure 4. Ligand-specificities of fish and human MRs.

Full-length sturgeon MR (A), gar MR (B), zebrafish (C), and human MR (D) were expressed in

HEK293 cells with an MMTV-luciferase reporter.   Cells were treated with 10^{-8} M Aldo, F, B, S,

DOC, Prog, 5α-dihydrotestosterone (DHT), 17β-estradiol (E2) or vehicle alone (DMSO).

Results are expressed as means ± SEM, n=3.   Y-axis indicates fold-activation compared to the

activity of control vector with vehicle (DMSO) alone as 1.

1   We also examined concentration-dependent activation of gar, sturgeon, zebrafish, and

2   human MRs by Aldo, F, B, DOC, S and Prog (Figure 5, Table 1).   Both gar and sturgeon MRs

3   had similar low EC50s, which varied from 7.7 pM to 150 pM for these steroids.   For each

4   steroid, the EC50s for gar MR were a little lower than for sturgeon MR.

6   **Figure 5. Concentration-dependent transcriptional activities of fish and human MRs.**

Concentration-response profiles of full-length sturgeon MR (A and B), gar MR (C and D), zebrafish MR (E and F), and human MR (G and H) for various steroids.   HEK293 cells were transiently transfected with the MMTV-containing vector together with an MR expression vector. Cells were incubated with increasing concentrations of Aldo, B, and S (A, C, E, and G) or Prog, F, and DOC (B, D, F, and H) (10^{-13} to 10^{-6}M).   Data are expressed as a ration of steroid to vehicle (DMSO).   Each column represents the mean of triplicate determinations, and vertical bars represent the mean ± SEM.

Table 1. EC50 activities for 3-keto-steroid transcriptional activation of sturgeon, gar, zebrafish and human MRs

	Aldo	DOC	B	S	F	Prog
Sturgeon MR	2.7×10^{-11}	1.3×10^{-11}	4.8×10^{-11}	8.2×10^{-11}	1.5×10^{-10}	7.0×10^{-11}
Gar MR	1.7×10^{-11}	7.7×10^{-12}	3.1×10^{-11}	2.6×10^{-11}	5.3×10^{-11}	1.8×10^{-11}
Zebrafish MR	8.8×10^{-11}	7.4×10^{-11}	3.3×10^{-10}	5.0×10^{-10}	5.9×10^{-10}	7.4×10^{-10}
Human MR	6.5×10^{-11}	1.7×10^{-10}	2.2×10^{-10}	2.0×10^{-9}	6.5×10^{-10}	-

In comparison, EC50s of Aldo, B and F were similar for zebrafish and human MR and a little higher than their EC50s for sturgeon and gar MR.   EC50s of DOC, S and Prog for zebrafish MR were higher than their EC50s for sturgeon and gar MR, but lower than the EC50s for human MR.   Prog had a lower, but still significant, maximal activation for zebrafish MR and at 100 nM Prog had little activation of human MR.   Overall all corticosteroids and Prog had EC50s that would be consistent with a physiological role in transcription of the MR in sturgeon, gar and zebrafish (Table 1, Figure 5).

4. Discussion

The evolution of steroids that are transcriptional activators of the MR has been a puzzle because Aldo, the mineralocorticoid for terrestrial vertebrates first appears in lungfish [63]. Nevertheless, Aldo is a potent activator of the lamprey CR [8], which is ancestral to the MR [7-9].   Interestingly, F, DOC, B and S and Prog also are transcriptional activators of the CR in lamprey and hagfish [8], with only S, thus far, found to have mineralocorticoid activity in lamprey [14].   In skate, which has separate MR and GR genes, Aldo, F, DOC and B are strong transcriptional activators of the MR [15].   F, DOC, B and S are found in teleosts [26] and F and DOC have been proposed to be transcriptional activators of teleost MRs [6, 16-20, 22-25].

Absent, until now, was information about the response to corticosteroids of MRs in sturgeon and gar, two basal fish that fill in the gap between elasmobranchs and teleosts (Figure 2).   Here we report that sturgeon MR and gar MR have EC50s below 1 nM for F, DOC, B, S and Prog, as well as Aldo.   Interestingly, we find that zebrafish MR also has a similar strong

response to corticosteroids and Prog.   This low selectivity for 3-ketosteroids (Figure 1) that can activate these fish MRs resembles the response of lamprey and hagfish CR [8] and skate MR [15].   Thus, this strong response of the MR to a broad panel of 3-keto-steroids was conserved after the third whole-genome duplication at the base of the teleosts [29, 30]

In contrast, human MR is more selective for 3-keto-steroids with higher EC50s for S and Prog.   Our data showing weak activation by Prog of human MR is in agreement with other studies [33, 34].   The weak response of human MR to Prog combined with the high affinity of Prog explains the antagonist activity of Prog for human MR.   The strong response to Prog of ray-finned fish MR is interesting in the light of the finding of Geller et al. that human MR with a Ser810Leu mutation was activated by 1 nM Prog [31].   Mutagenesis studies and structural analyses of MR-Leu810 led to the hypothesis that Leu-810 on α-helix 5 has stabilizing contacts with Ala-773 on α-helix 3 [31, 34, 35], which promote strong transcriptional activation by Prog. This serine and alanine are conserved in zebrafish MR [6] and sturgeon and gar MRs (unpublished) indicating that other mechanism(s) can lead to a strong response of the MR to Prog.

4.1 Mechanisms for regulation of steroid activation of ray-finned fish MR

Although steroid activation of zebrafish MR was not the initial focus of the project, the strong response of zebrafish MR, as well as sturgeon and gar MRs, to five corticosteroids and Prog requires one or more mechanisms to provide steroid-specific regulation of transcriptional activation of these ray-finned MRs.   At this time, such mechanisms in gar, sturgeon and zebrafish MRs or other ray fined fish MRs are poorly understood.   Clues for possible mechanisms may be found from insights into regulation of mammalian MRs [6, 13, 36, 38, 43, 44, 51, 53].   One possibility is an important mechanism in epithelial cells for regulating access of F and B to mammalian MR by tissue specific expression of 11β-HSD2, which selectively converts F and B to cortisone (E) and 11-dehydrocortisone (A), two inactive steroids.   Aldo is inert to 11β-HSD2, allowing Aldo to occupy the MR in epithelial cells in which 11β-HSD2 inactivates F and B [13, 36, 38].   11β-HSD2 is found in ray fined fish [64, 65], including sturgeon and gar (unpublished).   Expression of 11β-HSD2 in MR-containing tissues provides a mechanism to exclude F and B from the MR.   DOC, S and Prog, which have low EC50s in gar, sturgeon and zebrafish, lack an 11β-hydroxyl group and are inert to 11β-HSD2.

Other potential regulatory mechanisms include tissue-selective synthesis of 3-keto-steroids, selective binding of 3-keto-steroids to plasma proteins [43-45], steroid-specific conformational changes that regulate MR binding of co-activators, [46-49], effects of inter-domain interactions between the NTD and the LBD [20, 50, 51, 66] and post-translational modifications, such as phosphorylation, and SUMOylation [52-54] of the MR of the response to

3-keto-steroids.

Acknowledgments

We thank colleagues in our laboratories.   We also thank Drs. Arlette Hernandez-Franyutti and

Mari Carmen for providing gar tissues.   K.O. was supported by the Japan Society for the

Promotion of Science (JSPS) Research Fellowships for Young Scientists.   This work was

supported in part by Grants-in-Aid for Scientific Research 23570067 and 26440159 (YK) from

the Ministry of Education, Culture, Sports, Science and Technology of Japan.

References

[1] Lu, N.Z., Wardell, S.E., Burnstein, K.L., Defranco, D., Fuller, P.J., Giguere, V.,

Hochberg, R.B., McKay, L., Renoir, J.M., Weigel, N.L., et al. 2006 International Union

of Pharmacology. LXV. The pharmacology and classification of the nuclear receptor

superfamily: glucocorticoid, mineralocorticoid, progesterone, and androgen receptors.

Pharmacological reviews **58**, 782-797. (doi:10.1124/pr.58.4.9).

[2] Markov, G.V., Tavares, R., Dauphin-Villemant, C., Demeneix, B.A., Baker, M.E. &

Laudet, V. 2009 Independent elaboration of steroid hormone signaling pathways in

metazoans. *Proc Natl Acad Sci U S A* **106**, 11913-11918.

(doi:10.1073/pnas.0812138106).

[3] Bridgham, J.T., Eick, G.N., Larroux, C., Deshpande, K., Harms, M.J., Gauthier,

M.E., Ortlund, E.A., Degnan, B.M. & Thornton, J.W. 2010 Protein evolution by

molecular tinkering: diversification of the nuclear receptor superfamily from a

ligand-dependent ancestor. *PLoS biology* **8**. (doi:10.1371/journal.pbio.1000497).

[4] Sladek, F.M. 2011 What are nuclear receptor ligands? *Molecular and cellular*

endocrinology **334**, 3-13. (doi:DOI 10.1016/j.mce.2010.06.018).

[5] Bertrand, S., Belgacem, M.R. & Escriva, H. 2011 Nuclear hormone receptors in

chordates. *Molecular and cellular endocrinology* **334**, 67-75.

[6] Baker, M.E., Funder, J.W. & Kattoula, S.R. 2013 Evolution of hormone selectivity

in glucocorticoid and mineralocorticoid receptors. *J Steroid Biochem Mol Biol* **137**,

57-70. (doi:10.1016/j.jsbmb.2013.07.009

S0960-0760(13)00138-6 [pii]).

[7] Baker, M.E., Nelson, D.R. & Studer, R.A. 2015 Origin of the response to adrenal

and sex steroids: Roles of promiscuity and co-evolution of enzymes and steroid

receptors. *J Steroid Biochem Mol Biol* **151**, 12-24. (doi:10.1016/j.jsbmb.2014.10.020).

[8] Bridgham, J.T., Carroll, S.M. & Thornton, J.W. 2006 Evolution of hormone-receptor

complexity by molecular exploitation. *Science* **312**, 97-101.

(doi:10.1126/science.1123348).

[9] Thornton, J.W. 2001 Evolution of vertebrate steroid receptors from an ancestral

estrogen receptor by ligand exploitation and serial genome expansions. *Proc Natl Acad*

Sci U S A **98**, 5671-5676. (doi:10.1073/pnas.091553298

091553298 [pii]).

[10] Funder, J.W. 2012 Aldosterone and mineralocorticoid receptors: a personal

reflection. *Molecular and cellular endocrinology* **350**, 146-150.

(doi:10.1016/j.mce.2011.11.026).

[11] Hawkins, U.A., Gomez-Sanchez, E.P., Gomez-Sanchez, C.M. & Gomez-Sanchez,

C.E. 2012 The ubiquitous mineralocorticoid receptor: clinical implications. *Curr*

Hypertens Rep **14**, 573-580. (doi:10.1007/s11906-012-0297-0).

[12] Martinerie, L., Munier, M., Le Menuet, D., Meduri, G., Viengchareun, S. &

Lombes, M. 2013 The mineralocorticoid signaling pathway throughout development:

expression, regulation and pathophysiological implications. *Biochimie* **95**, 148-157.

(doi:10.1016/j.biochi.2012.09.030

S0300-9084(12)00390-2 [pii]).

[13] Rossier, B.C., Baker, M.E. & Studer, R.A. 2015 Epithelial sodium transport and its

control by aldosterone: the story of our internal environment revisited. *Physiological*

reviews **95**, 297-340. (doi:10.1152/physrev.00011.2014).

[14] Close, D.A., Yun, S.S., McCormick, S.D., Wildbill, A.J. & Li, W. 2010

11-deoxycortisol is a corticosteroid hormone in the lamprey. *Proc Natl Acad Sci U S A*

107, 13942-13947. (doi:10.1073/pnas.0914026107).

[15] Carroll, S.M., Bridgham, J.T. & Thornton, J.W. 2008 Evolution of hormone

signaling in elasmobranchs by exploitation of promiscuous receptors. *Molecular*

biology and evolution **25**, 2643-2652. (doi:10.1093/molbev/msn204).

[16] Greenwood, A.K., Butler, P.C., White, R.B., DeMarco, U., Pearce, D. & Fernald,

R.D. 2003 Multiple corticosteroid receptors in a teleost fish: distinct sequences,

expression patterns, and transcriptional activities. *Endocrinology* **144**, 4226-4236.

(doi:10.1210/en.2003-0566

en.2003-0566 [pii]).

[17] Sturm, A., Bury, N., Dengreville, L., Fagart, J., Flouriot, G., Rafestin-Oblin, M.E.

& Prunet, P. 2005 11-deoxycorticosterone is a potent agonist of the rainbow trout

(Oncorhynchus mykiss) mineralocorticoid receptor. *Endocrinology* **146**, 47-55.

(doi:en.2004-0128 [pii]

10.1210/en.2004-0128).

[18] Stolte, E.H., de Mazon, A.F., Leon-Koosterziel, K.M., Jesiak, M., Bury, N.R.,

Sturm, A., Savelkoul, H.F., van Kemenade, B.M. & Flik, G. 2008 Corticosteroid receptors involved in stress regulation in common carp, Cyprinus carpio. *J Endocrinol* **198**, 403-417. (doi:10.1677/JOE-08-0100).

[19] Arterbery, A.S., Fergus, D.J., Fogarty, E.A., Mayberry, J., Deitcher, D.L., Lee Kraus, W. & Bass, A.H. 2011 Evolution of ligand specificity in vertebrate corticosteroid receptors. *BMC Evol Biol* **11**, 14. (doi:1471-2148-11-14 [pii] 10.1186/1471-2148-11-14).

[20] Pippal, J.B., Cheung, C.M., Yao, Y.Z., Brennan, F.E. & Fuller, P.J. 2011 Characterization of the zebrafish (Danio rerio) mineralocorticoid receptor. *Molecular and cellular endocrinology* **332**, 58-66. (doi:S0303-7207(10)00475-2 [pii] 10.1016/j.mce.2010.09.014).

[21] Jiang, J.Q., Young, G., Kobayashi, T. & Nagahama, Y. 1998 Eel (Anguilla japonica) testis 11beta-hydroxylase gene is expressed in interrenal tissue and its product lacks aldosterone synthesizing activity. *Molecular and cellular endocrinology* **146**, 207-211.

[22] Baker, M.E. 2003 Evolution of glucocorticoid and mineralocorticoid responses: go fish. *Endocrinology* **144**, 4223-4225. (doi:10.1210/en.2003-0843).

[23] Bury, N.R. & Sturm, A. 2007 Evolution of the corticosteroid receptor signalling pathway in fish. *Gen Comp Endocrinol* **153**, 47-56. (doi:S0016-6480(07)00117-7 [pii] 10.1016/j.ygcen.2007.03.009).

[24] Sakamoto, T., Mori, C., Minami, S., Takahashi, H., Abe, T., Ojima, D., Ogoshi, M. & Sakamoto, H. 2011 Corticosteroids stimulate the amphibious behavior in mudskipper: potential role of mineralocorticoid receptors in teleost fish. *Physiol Behav* **104**, 923-928. (doi:10.1016/j.physbeh.2011.06.002).

[25] Takahashi, H. & Sakamoto, T. 2013 The role of 'mineralocorticoids' in teleost fish: relative importance of glucocorticoid signaling in the osmoregulation and 'central' actions of mineralocorticoid receptor. *Gen Comp Endocrinol* **181**, 223-228. (doi:10.1016/j.ygcen.2012.11.016).

[26] Milla, S., Wang, N., Mandiki, S.N. & Kestemont, P. 2009 Corticosteroids: Friends or foes of teleost fish reproduction? *Comparative biochemistry and physiology. Part A, Molecular & integrative physiology* **153**, 242-251. (doi:10.1016/j.cbpa.2009.02.027).

[27] Gilmour, K.M. 2005 Mineralocorticoid receptors and hormones: fishing for answers. *Endocrinology* **146**, 44-46. (doi:10.1210/en.2004-1390).

[28] Prunet, P., Sturm, A. & Milla, S. 2006 Multiple corticosteroid receptors in fish: from old ideas to new concepts. *Gen Comp Endocrinol* **147**, 17-23. (doi:10.1016/j.ygcen.2006.01.015).

[29] Jaillon, O., Aury, J.M., Brunet, F., Petit, J.L., Stange-Thomann, N., Mauceli, E.,

Bouneau, L., Fischer, C., Ozouf-Costaz, C., Bernot, A., et al. 2004 Genome duplication in the teleost fish Tetraodon nigroviridis reveals the early vertebrate proto-karyotype. *Nature* **431**, 946-957. (doi:10.1038/nature03025).

[30] Kasahara, M., Naruse, K., Sasaki, S., Nakatani, Y., Qu, W., Ahsan, B., Yamada, T., Nagayasu, Y., Doi, K., Kasai, Y., et al. 2007 The medaka draft genome and insights into vertebrate genome evolution. *Nature* **447**, 714-719. (doi:10.1038/nature05846).

[31] Geller, D.S., Farhi, A., Pinkerton, N., Fradley, M., Moritz, M., Spitzer, A., Meinke, G., Tsai, F.T., Sigler, P.B. & Lifton, R.P. 2000 Activating mineralocorticoid receptor mutation in hypertension exacerbated by pregnancy. *Science* **289**, 119-123.

[32] Oka, K., Hoang, A., Okada, D., Iguchi, T., Baker, M.E. & Katsu, Y. 2015 Allosteric role of the amino-terminal A/B domain on corticosteroid transactivation of gar and human glucocorticoid receptors. *J Steroid Biochem Mol Biol* **154**, 112-119. (doi:10.1016/j.jsbmb.2015.07.025).

[33] Rupprecht, R., Reul, J.M., van Steensel, B., Spengler, D., Soder, M., Berning, B., Holsboer, F. & Damm, K. 1993 Pharmacological and functional characterization of human mineralocorticoid and glucocorticoid receptor ligands. *Eur J Pharmacol* **247**, 145-154.

[34] Bledsoe, R.K., Madauss, K.P., Holt, J.A., Apolito, C.J., Lambert, M.H., Pearce, K.H., Stanley, T.B., Stewart, E.L., Trump, R.P., Willson, T.M., et al. 2005 A ligand-mediated hydrogen bond network required for the activation of the mineralocorticoid receptor. *The Journal of biological chemistry* **280**, 31283-31293. (doi:M504098200 [pii] 10.1074/jbc.M504098200).

[35] Fagart, J., Huyet, J., Pinon, G.M., Rochel, M., Mayer, C. & Rafestin-Oblin, M.E. 2005 Crystal structure of a mutant mineralocorticoid receptor responsible for hypertension. *Nature structural & molecular biology* **12**, 554-555. (doi:10.1038/nsmb939).

[36] Chapman, K., Holmes, M. & Seckl, J. 2013 11beta-hydroxysteroid dehydrogenases: intracellular gate-keepers of tissue glucocorticoid action. *Physiological reviews* **93**, 1139-1206. (doi:10.1152/physrev.00020.2012).

[37] Odermatt, A. & Atanasov, A.G. 2009 Mineralocorticoid receptors: emerging complexity and functional diversity. *Steroids* **74**, 163-171. (doi:10.1016/j.steroids.2008.10.010).

[38] Odermatt, A. & Kratschmar, D.V. 2012 Tissue-specific modulation of mineralocorticoid receptor function by 11beta-hydroxysteroid dehydrogenases: an overview. *Molecular and cellular endocrinology* **350**, 168-186.

(doi:10.1016/j.mce.2011.07.020).

[39] Nebert, D.W. & Russell, D.W. 2002 Clinical importance of the cytochromes P450. *Lancet* **360**, 1155-1162. (doi:10.1016/S0140-6736(02)11203-7).

[40] Payne, A.H. & Hales, D.B. 2004 Overview of steroidogenic enzymes in the pathway from cholesterol to active steroid hormones. *Endocr Rev* **25**, 947-970. (doi:10.1210/er.2003-0030).

[41] Taves, M.D., Gomez-Sanchez, C.E. & Soma, K.K. 2011 Extra-adrenal glucocorticoids and mineralocorticoids: evidence for local synthesis, regulation, and function. *Am J Physiol Endocrinol Metab* **301**, E11-24. (doi:10.1152/ajpendo.00100.2011).

[42] Taves, M.D., Plumb, A.W., Sandkam, B.A., Ma, C., Van Der Gugten, J.G., Holmes, D.T., Close, D.A., Abraham, N. & Soma, K.K. 2015 Steroid profiling reveals widespread local regulation of glucocorticoid levels during mouse development. *Endocrinology* **156**, 511-522. (doi:10.1210/en.2013-1606).

[43] Funder, J. & Myles, K. 1996 Exclusion of corticosterone from epithelial mineralocorticoid receptors is insufficient for selectivity of aldosterone action: in vivo binding studies. *Endocrinology* **137**, 5264-5268. (doi:10.1210/endo.137.12.8940344).

[44] Sivukhina, E.V. & Jirikowski, G.F. 2014 Adrenal steroids in the brain: role of the intrinsic expression of corticosteroid-binding globulin (CBG) in the stress response. *Steroids* **81**, 70-73. (doi:10.1016/j.steroids.2013.11.001).

[45] Breuner, C.W. & Orchinik, M. 2002 Plasma binding proteins as mediators of corticosteroid action in vertebrates. *J Endocrinol* **175**, 99-112.

[46] Khan, S.H., Awasthi, S., Guo, C., Goswami, D., Ling, J., Griffin, P.R., Simons, S.S., Jr. & Kumar, R. 2012 Binding of the N-terminal region of coactivator TIF2 to the intrinsically disordered AF1 domain of the glucocorticoid receptor is accompanied by conformational reorganizations. *The Journal of biological chemistry* **287**, 44546-44560. (doi:10.1074/jbc.M112.411330).

[47] Li, Y., Suino, K., Daugherty, J. & Xu, H.E. 2005 Structural and biochemical mechanisms for the specificity of hormone binding and coactivator assembly by mineralocorticoid receptor. *Mol Cell* **19**, 367-380. (doi:S1097-2765(05)01429-2 [pii] 10.1016/j.molcel.2005.06.026).

[48] Bledsoe, R.K., Montana, V.G., Stanley, T.B., Delves, C.J., Apolito, C.J., McKee, D.D., Consler, T.G., Parks, D.J., Stewart, E.L., Willson, T.M., et al. 2002 Crystal structure of the glucocorticoid receptor ligand binding domain reveals a novel mode of receptor dimerization and coactivator recognition. *Cell* **110**, 93-105. (doi:S0092867402008176 [pii]).

[49] Hultman, M.L., Krasnoperova, N.V., Li, S., Du, S., Xia, C., Dietz, J.D., Lala, D.S., Welsch, D.J. & Hu, X. 2005 The ligand-dependent interaction of mineralocorticoid receptor with coactivator and corepressor peptides suggests multiple activation mechanisms. *Mol Endocrinol* **19**, 1460-1473. (doi:10.1210/me.2004-0537).

[50] Fuller, P.J. 2015 Novel interactions of the mineralocorticoid receptor. *Molecular and cellular endocrinology* **408**, 33-37. (doi:10.1016/j.mce.2015.01.027).

[51] Fuller, P.J., Yao, Y., Yang, J. & Young, M.J. 2012 Mechanisms of ligand specificity of the mineralocorticoid receptor. *J Endocrinol* **213**, 15-24. (doi:JOE-11-0372 [pii] 10.1530/JOE-11-0372).

[52] Bhargava, A. & Pearce, D. 2004 Mechanisms of mineralocorticoid action: determinants of receptor specificity and actions of regulated gene products. *Trends Endocrinol Metab* **15**, 147-153. (doi:10.1016/j.tem.2004.03.009).

[53] Pascual-Le Tallec, L. & Lombes, M. 2005 The mineralocorticoid receptor: a journey exploring its diversity and specificity of action. *Mol Endocrinol* **19**, 2211-2221. (doi:10.1210/me.2005-0089).

[54] Faresse, N. 2014 Post-translational modifications of the mineralocorticoid receptor: How to dress the receptor according to the circumstances? *J Steroid Biochem Mol Biol* **143**, 334-342. (doi:10.1016/j.jsbmb.2014.04.015).

[55] Katsu, Y., Kohno, S., Hyodo, S., Ijiri, S., Adachi, S., Hara, A., Guillette, L.J., Jr. & Iguchi, T. 2008 Molecular cloning, characterization, and evolutionary analysis of estrogen receptors from phylogenetically ancient fish. *Endocrinology* **149**, 6300-6310. (doi:10.1210/en.2008-0670).

[56] Saitou, N. & Nei, M. 1987 The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Molecular biology and evolution* **4**, 406-425.

[57] Edgar, R.C. 2004 MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research* **32**, 1792-1797. (doi:10.1093/nar/gkh340).

[58] Felsenstein, J. 1985 Confidence-Limits on Phylogenies - an Approach Using the Bootstrap. *Evolution* **39**, 783-791. (doi:Doi 10.2307/2408678).

[59] Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar, S. 2011 MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Molecular biology and evolution* **28**, 2731-2739. (doi:10.1093/molbev/msr121).

[60] Katsu, Y., Kubokawa, K., Urushitani, H. & Iguchi, T. 2010 Estrogen-dependent transactivation of amphioxus steroid hormone receptor via both estrogen and androgen response elements. *Endocrinology* **151**, 639-648.

[61] Kadlecova, Z., Nallet, S., Hacker, D.L., Baldi, L., Klok, H.A. & Wurm, F.M. 2012

Poly(ethyleneimine)-mediated large-scale transient gene expression: influence of

molecular weight, polydispersity and N-propionyl groups. *Macromol Biosci* **12**,

628-636. (doi:10.1002/mabi.201100404).

[62] Oka, K., Kohno, S., Urushitani, H., Guillette, L.J., Jr., Ohta, Y., Iguchi, T. & Katsu,

Y. 2013 Molecular cloning and characterization of the corticoid receptors from the

American alligator. *Molecular and cellular endocrinology* **365**, 153-161.

(doi:10.1016/j.mce.2012.10.014).

[63] Joss, J.M.P., Arnoldreed, D.E. & Balment, R.J. 1994 The Steroidogenic Response

to Angiotensin-Ii in the Australian Lungfish, Neoceratodus-Forsteri. *J Comp Physiol B*

164, 378-382. (doi:Doi 10.1007/Bf00302553).

[64] Kusakabe, M., Nakamura, I. & Young, G. 2003 11beta-hydroxysteroid

dehydrogenase complementary deoxyribonucleic acid in rainbow trout: cloning, sites of

expression, and seasonal changes in gonads. *Endocrinology* **144**, 2534-2545.

(doi:10.1210/en.2002-220446).

[65] Baker, M.E. 2010 11Beta-hydroxysteroid dehydrogenase-type 2 evolved from an

ancestral 17beta-hydroxysteroid dehydrogenase-type 2. *Biochemical and biophysical*

research communications **399**, 215-220. (doi:10.1016/j.bbrc.2010.07.057).

[66] Pippal, J.B., Yao, Y., Rogerson, F.M. & Fuller, P.J. 2009 Structural and functional

characterization of the interdomain interaction in the mineralocorticoid receptor. *Mol*

Endocrinol **23**, 1360-1370. (doi:me.2009-0032 [pii]

10.1210/me.2009-0032).