1   Fractional Dosing of Yellow Fever Vaccine to Extend Supply: A Modeling Study
2

3   **AUTHORS**
4   Joseph T. Wu[1], PhD, joewu@hku.hk*
5   Corey M. Peak[2], MS
6   Prof Gabriel M. Leung[1], MD
7   Prof Marc Lipsitch[2,3], DPhil mlipsitc@hsph.harvard.edu*
8
[1]WHO Collaborating Centre for Infectious Disease Epidemiology and Control, School of Public
Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong Special
Administrative Region, China
[2] Center for Communicable Disease Dynamics, Department of Epidemiology, Harvard T.H. Chan
School of Public Health, 677 Huntington Avenue, Boston MA 02115 USA
[3] Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public
Health, 665 Huntington Avenue, Boston MA 02115 USA. +1-617-432-4559.
* corresponding authors

17
18

Background
The ongoing yellow fever (YF) epidemic in Angola is placing strain on the global vaccine supply.
In order to extend vaccine supply and reduce the cost of mass-vaccination, dose sparing by
fractional-dose vaccination has received heightened consideration. Five-fold fractionation is
similar to the standard dose in safety and immunogenicity. However, no YF vaccine efficacy
trials have been performed in humans, so it is possible that fractional-dose vaccines may be less
efficacious even if equally immunogenic. There is an urgent need to study under what
conditions fractional dosing could provide epidemiologic benefits in reducing transmission.
27
Methods
We estimated the effective reproductive number for YF in Angola using disease natural history
and case report data. Using these results and simple mathematical models of YF transmission,
we calculated the expected final size of an epidemic under varying levels of vaccine coverage
with standard-dose vaccines and up to five-fold fractionation with varying efficacy. We consider
two allocation scenarios: random and whereby children receive standard-dose vaccines while
adults receive fractional-dose vaccines.
35
Findings
The effective reproductive number early in the outbreak ranged from approximately 5·2 to 7·1
transmission events per infectious individual. Intuition dictates, and we confirm with modeling
analysis, that five-fold fractional-doses can dramatically reduce the final epidemic size. If
vaccine efficacy is all-or-nothing, as we expect, the conclusion holds that n-fold fractionation is
beneficial as long as the efficacy is greater than $1/n$. We quantify how the threshold becomes
more stringent if fractional vaccines instead provide partial protection to every recipient (i.e.
"leaky" vaccine action).

45
46 **Interpretation**
47 We conclude that dose fractionation could be a very effective strategy for reducing infection
48 attack rate in populations with a large margin of error in case fractional-dose efficacy turns out
49 to be lower than expected.
50
51 **Funding**
52

INTRODUCTION

Yellow fever (YF) has resurged in Angola and threatens to spread to other countries with relatively low YF vaccine coverage. As of June 16, YF cases have been exported from Angola to Kenya (2 cases), China (11), and DRC (53), raising concern YF could resurge in other populations where competent vectors are present and vaccine coverage is low, especially during the rainy season which is beginning in West Africa.[1,2] A broad band of sub-Saharan Africa north of Namibia and Zambia is at risk (http://www.cdc.gov/yellowfever/maps/africa.html), as is much of the northern portion of South America (http://www.cdc.gov/yellowfever/maps/south_america.html). The global community is increasingly concerned for the risk of YF emergence in Asia, where the disease has been curiously absent despite seemingly amenable conditions.

There is a safe, highly effective live-attenuated vaccine against YF.[3] However, the global emergency stockpile of YF vaccines is low, with approximately 6.8 million doses currently available and 2-4 million more doses expected per month.[1] Given the large populations at risk for YF infection, the stockpile is expected to be inadequate to meet the need.[4] For this reason, the WHO is considering the possibility of dose-sparing by fractional-dose vaccination,[5] in which smaller volumes of vaccine would be used per dose in order to increase the number of persons who can be vaccinated with a given quantity of vaccine.[3] This strategy was previously proposed to extend pre-pandemic influenza vaccine supplies.[6] If dose-fractionation were consistently adopted, equity of YF vaccine access might also be enhanced both within and across countries at risk, as more people could benefit from vaccination without depriving others.[7]

A randomized, noninferiority trial has shown that 0·1 ml intradermal (ID) vaccination with the 17D YF vaccine was equally safe and immunogenic compared to the standard 0·5ml subcutaneous vaccination.[8] Another randomized trial of subcutaneous administration of the 17DD vaccine given in Brazil showed that there was no significant difference in immunogenicity and viremia kinetics when the currently administered vaccine (containing 27,476 IU of virus) was given at subdoses as low as 11% of the full dose (3,013 IU).[9] Even lower doses produced noninferior immune responses, but not equivalent viremia kinetics.[9] For comparison, the WHO minimum for YF vaccines is 1,000 IU per dose at the end of shelf life.[10]

No efficacy trial of YF vaccines has been performed in humans,[11] so the comparative efficacy of different doses and routes of administration remains uncertain. In particular, it is not known whether equal immunogenicity implies equal vaccine efficacy for YF vaccines. Moreover, the findings of equal immunogenicity of reduced doses are limited to healthy adults; no comparable data exist in children, elderly or immunocompromised individuals (e.g. HIV-infected people, pregnant women, etc). As such, while noninferior immunogenicity of fractional-dose vaccines provide a strong basis for an initial consideration of dose-sparing strategies for YF vaccines, it is unlikely that decision makers would change dosing recommendations without carefully evaluating the risk and implications of reduced vaccine efficacy in fractional-dose vaccines. Such an evaluation is nontrivial because even if dose fractionation reduces vaccine efficacy, higher vaccine coverage may confer higher population-level herd immunity in which

case the number of infections could be significantly reduced by the indirect effect of large-scale
vaccination.[12] The lower the transmissibility, the larger the number of infections that can be
averted by indirect protection, as has been illustrated by the previous study of dose-
fractionation for pre-pandemic influenza vaccines.[6] The importance of herd immunity for YF
vaccination is unknown because transmissibility of YF in urban settings has so far been poorly
characterized due to limited data.

To strengthen the evidence base for the public health impact of dose-fractionation of YF
vaccines, we use simple mathematical models to assess the potential reduction in infection
attack rate (IAR, defined as the proportion of population infected over the course of an
epidemic) conferred by five-fold dose-fractionation under different epidemic scenarios and
reductions in vaccine efficacy. We first estimate the transmissibility of YF during the recent
Angola outbreak in order to parameterize realistic epidemic scenarios. We then show that even
if vaccine efficacy for five-fold fractional-dose vaccination were considerably lower, higher
vaccine coverage could achieve significant reduction in IAR despite lower individual-level
efficacy, with the break-even point being 20% efficacy under the assumption that reduced
efficacy represents a mix of complete efficacy in some individuals and failure in others ("all-or-
nothing"), but higher if vaccines are partially effective in all individuals ("leaky"). Next, given the
lack of comparative immunogenicity data for fractional-dose YF vaccination in children, we
consider the results of a strategy that provides standard-dose vaccines to children and
fractional-dose vaccines to adults. We find that all dose-sparing strategies considered could
provide significant benefit epidemiologically, and that the best policy will be determined by
balancing logistical and regulatory considerations against the extent of epidemiologic benefit.

METHODS

Estimating the epidemiologic parameters for YF
To parameterize realistic epidemic scenarios for our analysis, we estimate the reproductive
number of YF over the course of the Angola outbreak and use the estimates during the early
epidemic stages (before large-scale vaccination affected transmission) as the range of basic
reproductive number for future outbreaks in other populations. To this end, we use the
Wallinga and Teunis method[13] to estimate the reproductive number of YF from the daily
number of confirmed YF cases recorded in the 17 April 2016 WHO Angola Situation Report,[14]
assuming that all cases were attributed to local transmission (i.e. no importation of cases).
When estimating the extrinsic incubation period, we assume that the average temperature in
Angola was 28 degrees Celsius during the outbreak. To estimate the serial interval distribution,
we make the following assumptions: (i) the extrinsic incubation period follows the Weibull
distribution estimated by ref. [15] which has mean 12·7 days at 28 degrees Celsius; (ii) the
intrinsic incubation period follows the lognormal distribution estimated by ref. [15] which has
mean 4·6 days; (iii) the infectious period in human is exponentially distributed with mean 4
days;[16] (iv) the mosquito lifespan is exponentially distributed with mean 7 to 14 days.[17] We
estimate the initial reproductive number of the YF outbreak in Angola as the average
reproductive number among all cases who developed symptoms one serial interval before
vaccination campaign began to affect disease transmission (see Figure 1).

Dose-response for fractional-dose vaccines
Let S_0 be the proportion of population susceptible just before the vaccination campaign begins
and V be the vaccine coverage achievable with standard-dose vaccines. Suppose each
standard-dose vaccine can be fractionated into *n*, *n*-fold fractional-dose vaccines (i.e. each of
which contains 1/n-th the amount of the antigen in a standard-dose vaccine) with vaccine
efficacy $VE(n)$. That is, the vaccine efficacy of standard-dose vaccines is $VE(1)$ which was
assumed to be 1. Given V, the highest fractionation sensible is $n_{max} = S_0/V$ if the susceptible
population can be identified for targeted vaccination and $n_{max} = 1/V$ otherwise, i.e. the
fractionation n must lie between 1 and n_{max}. To avoid overstating the benefit of dose-
fractionation, we assume that vaccine efficacy of n-fold fractional-dose vaccines for *n* between
1 and 5 increases linearly with the amount of antigen in the vaccines (see appendix for more
details). Potential increases in vaccine wastage during dose-sparing would be mostly due to
unused, reconstituted vaccines[18] or increased vaccine failure due to inexperience with
intradermal administration among vaccinators. In the setting of mass vaccination campaigns,
wastage due to unused vaccine doses will likely to be negligible because vaccination sessions
will be large. Increased vaccine failure is effectively the same as reduced vaccine efficacy if
vaccine action is all-or-nothing (as we have assumed in the main text). As such, for simplicity,
we do not explicitly model wastage.

Infection attack rate
We use infection attack rate or IAR (defined as the proportion of population infected over the
course of an epidemic) as the outcome measure for evaluating the impact of dose-

fractionation. We calculate IAR using the classical final size approach which is exact for directly
transmitted SIR-type diseases[19] but only an approximation for vector-borne diseases.[20]
Nonetheless, this approximation is excellent over realistic parameter ranges because only a
very small proportion of mosquitoes are infected with YF virus even during epidemics
(necessitating pooled testing).[21] See appendix for the mathematical details.

We denote the IAR under n-fold dose fractionation by $IAR(n)$. To evaluate the outcome of
fractional-dose vaccination against that of standard-dose vaccination, we calculate the absolute
and relative reductions in IAR as $IAR(1) - IAR(n)$ and $1 - IAR(n)/IAR(1)$, respectively. We
assume that the vaccination campaign is completed before the start of the epidemic.

Vaccine action
We assume that vaccine action is all-or-nothing, i.e. *n*-fold fractional-dose vaccines provide
100% protection against infection in a proportion $VE(n)$ of vaccinees and no protection in the
remainder. In this case, *n*-fold dose fractionation results in lower IAR if and only if the vaccine
efficacy of *n*-fold fractional-dose vaccines are at least 1/*n* times that of standard-dose vaccines,
i.e. $VE(n) > VE(1)/n$ (see appendix for details). We term this the benefit threshold for dose-
fractionation. We also consider the alternative case in which vaccine action is leaky, i.e. *n*-fold
fractional-dose vaccines reduce the hazard of infection (the probability of disease transmission
per mosquito bite) of each vaccinee by a proportion $VE(n)$.[22,23] Compared to all-or-nothing
vaccines, leaky vaccines have substantially higher benefit thresholds, especially when
transmissibility is high (see Results). However, we postulate that vaccine action is much more
likely to be all-or-nothing than leaky (see Discussion). As such, we present our main results in
the context of all-or-nothing vaccine action.

Dose-sparing strategies
We consider two dose-sparing strategies with at most five folds of dose fractionation:

1. *Random vaccination.* Each individual in the population has the same probability of
receiving vaccination regardless of their susceptibility. That is, the susceptible and
immune population are indiscernible, so targeted vaccination is not possible. If
susceptible individuals can be identified, then targeted vaccination has the same
epidemiologic outcome as random vaccination with vaccine coverage V/S_0.
2. *Standard-dose vaccination of children, fractional-dose vaccination of adults.* We assume
that children have vaccine priority. That is, all children receive standard-dose vaccines
before adults begin to receive fractional-dose vaccines. Let p be the proportion of
adults in the population. For a given standard-dose vaccine coverage V, the proportion
of children vaccinated is $V_{children} = \min(V, 1-p)$. If the stockpile is large enough to
vaccinate all children (i.e. $V > 1-p$), then adults receive n-fold fractional-dose
vaccination where $n = \min(5, p/(V-1+p))$. The proportion of adults vaccinated is
$V_{adults} = 1$ if $n < 5$ and $V_{adults} = 5(V-1+p)/p$ otherwise. Given the vector-borne nature of
yellow fever we assume that transmission between children and adults is well-mixed.

We assume that all individuals are susceptible before vaccination ($S_0 = 1$) unless specified
otherwise.

Role of the funding source
199 The sponsors of the study had no role in the study design, data collection, data analysis, writing
200 of the report, or the decision to publish. All authors had access to the data; the corresponding
201 authors had final responsibility to submit for publication.
202
203
RESULTS

Reproductive number of yellow fever in Angola. Figure 1 shows that the initial reproductive
number of YF in Angola was 5·2 (95% CI 4·3, 6·1) and 7·1 (5·5, 8·7) if the mean mosquito
lifespan was 7 and 14 days, respectively. While these estimates may reflect partial immunity
due to prior vaccination or exposure among some of the population, we assume that the
possible basic reproductive number (R_0) in a future outbreak in another population would range
between 4 and 12 due to varying vector ecology and levels of preexisting immunity in the
population. In principle, disease transmission can be halted if the effective vaccine coverage
(defined as vaccine efficacy times vaccine coverage) exceeds the herd immunity threshold 1-
$1/R_0$.

Random vaccination. Figure 2A shows the IAR under standard-dose and fractional-dose
vaccination as a function of standard-dose vaccine coverage *V* given varying levels of
transmission and five-fold fractionation vaccine efficacy. Figures 2B-C show the corresponding
absolute and relative reduction in IAR when vaccine action is all-or-nothing and confirm our
earlier claim that fractional-dose vaccination reduces IAR when $VE(5) > VE(1)/n = 0.2$.
Fractional-dose vaccination substantially reduces IAR if *V* > 10% and such reduction only
diminishes to insignificant levels when *V* is close to the herd immunity threshold $(1-1/R_0) \times 100\%$
(e.g. 75% and 88% for R_0 = 4 and 8, respectively). In short, dose-fractionation reduces IAR when
(i) the standard-dose vaccine supply is insufficient to halt disease transmission and (ii)
fractional-dose vaccine efficacy is above 20%.

If vaccine action is "leaky," then the benefit threshold (the efficacy of *n*-fold fractionated doses
necessary to reduce IAR) is higher than $1/n$ and increases with transmission intensity. This
occurs because under the leaky model each infectious bite is assumed to be less likely to cause
infection if the host is vaccinated, but the probability of infection grows as the person receives
more infectious bites. Figure 3 shows, under the leaky model of vaccine action, dose
fractionation is much less beneficial if vaccine action is leaky, efficacy is modest, and R_0 is high.

A recent study suggested that the mosquito biting rate for individuals aged 20 or above is 1.22
times higher than those age under 20.[24] We performed a sensitivity analysis to show that our
results are unaffected by such heterogeneity. See "Hetereogeneity in biting rates" in the
appendix for details.

Vaccination of adults with fractionated doses and children with standard doses. Given the
lack of immunogenicity data for fractionated doses in children and evidence of lower
seroconversion rates to standard doses,[25] a conservative strategy would be to fractionate doses
only for adults while providing full doses to children. Figure 4A shows the fold-increase in
vaccine coverage that could be achieved with five-fold fractionation in adults only, as a function
of proportion of adults in the population (p). Adult fractionation increases coverage more if a
larger fraction of the population is adults. In Angola in 2015 approximately 57% of the
population was adults (15 and older). If there were enough standard-dose vaccine supplies to
cover 70% of such a population, fractionation of only the adult doses would increase the
coverage by a factor of 1.43 to 100%. Figure 4B-C shows the same calculations as Figure 2A-B
where the five-fold fractional-dose vaccine efficacy is 60%. These calculations all assume that
there is no preexisting immunity in children or adults; if preexisting immunity existed mainly in
adults, then prioritizing children would have a greater benefit than projected here.

DISCUSSION

Our primary analysis shows that dose-fractionation of YF vaccine, if there were no loss of efficacy, could provide a substantial benefit to reducing the attack rate of YF in a population. We consider this assumption of full efficacy for five-fold fractionation to be the most likely scenario, despite the lack of efficacy data on any YF vaccine, for several reasons: 1) two studies of five- or greater-fold vaccination doses have shown indistinguishable immunogenicity in humans; 2) at least some preparations of YF vaccine substantially exceed the WHO minimum standard for potency of 1,000 IU/dose, so fractionation at some level could be performed without dropping below that threshold; 3) YF vaccine is live attenuated virus, so a biological rationale exists that if a productive vaccine-virus infection can be established by a fractionated dose, protection should be comparable to that with a higher dose. Nonetheless, to assess the the robustness of the conclusion that dose-fractionation is likely to beneficial, against the possibility that in fact efficacy of fractionated doses is lower than anticipated, we consider the possibility that five-fold fractionated dosing fails to immunize a proportion (1-VE(5)) of recipients. Consistent with intuition, we find that as long as at least 20% of recipients are fully immunized by the vaccine, more people would be immunized by vaccinating five times as many people with one-fifth the dose, and so the population-wide benefits of higher coverage would outweigh the lower efficacy of fractionated dosing for individual vaccinees. Even more unlikely, in our opinion, is that fractionated doses would be less efficacious according to a "leaky" model, in which all vaccinated individuals were imperfectly protected against infection from each infectious bite, with the same probability of infection from each bite, reduced by vaccine by a proportion VE. If this were the case, then especially in high-transmission areas, the fractionated-dose vaccine would need to be 80-90% efficacious to provide a benefit over standard dosing.

Based on the limited evidence on immunogenicity of fractional doses of YF vaccine to date, we consider it unlikely that reducing the dose five-fold or perhaps further from current preparations would result in dramatically lower efficacy of the leaky type. Visual inspection of the data from a dose-fractionation trial of the 17DD vaccine in Brazil shows that for doses down to 47x below the standard dose, the distribution of serologic responses was indistinguishable from those for the standard dose, suggesting that efficacy should be nearly equivalent to that for full doses. This was confirmed by the analysis of peak viremia, which was equivalent for standard dose and for doses down to 11% of the full dose (9-fold fractionation). It was further confirmed by peak cytokine responses, which were comparable to the standard dose for all cytokines tested, down to at least a 9-fold fractional dose. For even lower doses, the proportion seroconverting after vaccination was lower than the 97% observed for the full dose, but the antibody response among the seroconverters appears to be similar at all doses.[9] These data collectively suggest that down to approximately 9-fold fractional dosing of this vaccine the response should be equivalent, and that for further fractionation there may be a failure to induce any substantial response in a fraction of recipients, but the neutralizing antibody titres in those who do respond should be comparable. This pattern is consistent with an all-or-nothing model.

Our analysis is not intended to recommend extending coverage to the point of knowingly
compromising efficacy. Rather, our analysis indicates that a strategy of fractionation to a dose
that provides equivalent immunogenicity to standard dosing would be greatly beneficial if
efficacy is equivalent to standard dosing, and would still be beneficial if, unexpectedly, efficacy
was somewhat lower than standard dosing.

We have used five-fold fractionation as an example because it is the strategy with the best
evidence base of equal immunogenicity. However, some data suggest that more than five-fold
fractionation could be equally immunogenic, and of course the benefits of fractionation would
be greater if more than five-fold fractionation were logistically possible and comparably
efficacious.

On programmatic grounds a simpler strategy -- such as fractionated dosing for all -- may be
preferred to a more complex strategy that gives different doses to different groups, say age
groups. While either would provide epidemiologic benefit, the choice between such strategies
would be influenced by the number of available doses, logistical barriers, and location-specific
regulations regarding specific groups, such as children. Another group of interest are travelers,
for whom we must also consider longevity of response, lower levels of exposure, and more
detailed discussions on equity outside the scope of this modeling paper. The cost of fractional-
dose strategies will depend on the route of administration, but could potentially be
substantially less expensive per vaccine recipient.[18]

Our simple model has several limitations. We assume homogeneous mixing of the population
(reasonable at least locally for a vector-borne disease) and neglect preexisting immunity in our
main results, though the Methods show how our calculations could be modified to consider
preexisting immunity. The purpose is to provide basic calculations for the most at-risk
populations, those with little preexisting immunity. We also fix a particular value of R_0 for each
calculation, and assume this value is maintained until the epidemic has swept through a
population. In reality, R_0 will vary seasonally as vector abundance, extrinsic incubation period,
and other factors vary. The existence of a high-transmission season might enhance the benefits
of fractional-dose vaccination. Most importantly, there will be a premium on achieving high
vaccine coverage before the peak of transmission to maximally impact transmission, and this
will be limited by supply constraints that could be partially relieved by fractionation.

We conclude that dose fractionation could be a very effective strategy for improving coverage
of YF vaccines and reducing infection attack rate in populations -- possibly by a large absolute
and relative margin -- if high to moderate efficacy is maintained by reduced-dose formulations.
For vaccines whose standard formulations exceed WHO minimum concentration of viral
particles,[10] this dose-fractionation could be accomplished without changing the WHO
recommendations. Even if the efficacy of fractionated doses were substantially lower than
expected, increasing coverage by a factor greater than the reduction in efficacy would still be
predicted to reduce the population-wide infection attack rate. Substantial benefits could also
be achieved if fractional doses were given only to adults while providing standard-dose vaccines
to children. We urge consideration of means to implement dose-fractionation as a component

of a YF response strategy for the current situation. Rollout of fractionated dosing should

perhaps be preceded or accompanied by noninferiority studies of the intended vaccine's

immunogenicity at fractional doses in the intended populations. Ongoing programs should be

monitored by observational studies of safety, immunogenicity and, if possible, effectiveness[18]

to assure that the assumptions underlying the rationale for such programs continue to be met.

However, it is worth noting that if full-dose vaccine efficacy is indeed 100% or nearly so as

currently believed, estimating the relative efficacy of fractional vs. standard doses in a

comparative study would be challenging or impossible, as there might be few or no cases

accrued in the standard-dose arm.

REFERENCES

1   World Health Organization. Situation Report: Yellow Fever. Geneva, Switzerland, 2016
http://apps.who.int/iris/bitstream/10665/242438/1/yellowfeversitrep-16Jun2016-
eng.pdf?ua=1.

2   Woodall J, Yuill T. Why is the yellow fever outbreak in Angola a 'threat to the entire
world'? *Int J Infect Dis* 2016; : 1–4.

3   Monath TP, Woodall JP, Gubler DJ, *et al.* Yellow fever vaccine supply: a possible solution.
Lancet 2016; **387**: 1599–600.

4   Barrett AD. Yellow Fever in Angola and Beyond — The Problem of Vaccine Supply and
Demand. *N Engl J Med* 2016; : NEJMp1606997.

5   Schnirring L. WHO plans to stretch vaccine as yellow fever persists in Africa. CIDRAP
News. 2016. http://www.cidrap.umn.edu/news-perspective/2016/06/who-plans-stretch-
vaccine-yellow-fever-persists-africa.

6   Riley S, Wu JT, Leung GM. Optimizing the dose of pre-pandemic influenza vaccines to
reduce the infection attack rate. *PLoS Med* 2007; **4**: 1032–40.

7   Brearley L, Eggers R, Steinglass R, Vandelaer J. Applying an equity lens in the Decade of
Vaccines. *Vaccine* 2013; **31**: B103–7.

8   Roukens AH, Vossen AC, Bredenbeek PJ, van Dissel JT, Visser LG. Intradermally
Administered Yellow Fever Vaccine at Reduced Dose Induces a Protective Immune
Response: A Randomized Controlled Non-Inferiority Trial. *PLoS One* 2008; **3**: e1993.

9   Campi-Azevedo AC, de Almeida Estevam P, Coelho-Dos-Reis JG, *et al.* Subdoses of 17DD
yellow fever vaccine elicit equivalent virological/immunological kinetics timeline. *BMC
Infect Dis* 2014; **14**: 391.

10   World Health Organization. Expert Committee on Biological Standardization.
Requirements for yellow fever vaccine. Geneva, Switzerland, 2008.

11   Staples JE, Gershman M, Fischer M. Yellow Fever Vaccine: Recommendations of the
Advisory Committee. *Morb Mortal Wkly Rep* 2010; **59**: 1–27.

12   Fine P, Eames K, Heymann DL. 'Herd immunity': A rough guide. *Clin Infect Dis* 2011; **52**:
911–6.

13   Wallinga J, Teunis P. Different Epidemic Curves for Severe Acute Respiratory Syndrome
Reveal Similar Impacts of Control Measures. *Am J Epidemiol* 2004; **160**: 509–16.

14   World Health Organization. Yellow fever outbreak in Angola: Situation Report. 2016
http://www.afro.who.int/index.php?option=com_docman&task=doc_download&gid=10
121&Itemid=2593.

15   Johansson MA, Arana-Vizcarrondo N, Biggerstaff BJ, Staples JE. Incubation Periods of
Yellow Fever Virus. *Am J Trop Med Hyg* 2010; **83**: 183–8.

16   Hindle E. The Transmission of Yellow Fever. *Lancet* 1930; **219**: 835–42.

17   Johansson MA, Arana-Vizcarrondo N, Biggerstaff BJ, Gallagher N, Marano N, Staples JE.

Assessing the risk of international spread of yellow fever virus: a mathematical analysis
of an urban outbreak in Asuncion, 2008. *Am J Trop Med Hyg* 2012; **86**: 349–58.

18   Hickling J, Jones R, Tandem W. Yellow fever vaccination: The potential of dose-sparing to
increase vaccine supply and. *Path* 2013.
http://www.path.org/publications/detail.php?i=2285.

19   Diekmann O, Heesterbeek J. Mathematical Epidemiology of Infectious Diseases: Model
Building, Analysis and Interpretation. New York: Wiley, 2000.

20   Radcliffe J, Rass L. The Spatial Spread and Final Size of Models for the Deterministic Host-
Vector Epidemic. *Math Biosci* 1984; **70**: 123–46.

21   World Health Organization. Rapid field entomological assessment during yellow fever
outbreaks in Africa. Geneva, Switzerland, 2014.

22   Smith PG, Rodrigues LC, Fine PEM. Assessment of the Protective Efficacy of Vaccines
against Common Diseases Using Case-Control and Cohort Studies. *Int J Epidemiol* 1984;
13: 87–93.

23   Halloran ME, Longini IM, Struchiner CJ. Design and Analysis of Vaccine Studies:
Introduction. New York, NY: Springer, 2010.

24   Harrington LC, Fleisher A, Ruiz-Moreno D, *et al.* Heterogeneous Feeding Patterns of the
Dengue Vector, Aedes aegypti, on Individual Human Hosts in Rural Thailand. *PLoS Negl*
Trop Dis 2014; **8**.

25   Belmusto-Worn VE, Sanchez JL, McCarthy K, *et al.* Randomized, double-blind, phase III,
pivotal field trial of the comparative immunogenicity, safety, and tolerability of two
yellow fever 17D vaccines (Arilvax and YF-VAX) in healthy infants and children in Peru.
Am J Trop Med Hyg 2005; **72**: 189–97.

26   Martins RM, Maia M de LS, Farias RHG, *et al.* 17DD yellow fever vaccine: a double blind,
randomized clinical trial of immunogenicity and safety on a dose-response study. *Hum*
Vaccin Immunother 2013; **9**: 879.

27   World Health Organization. Situation Report: Yellow Fever. Geneva, Switzerland, 2016
http://apps.who.int/iris/bitstream/10665/208880/1/yellowfeversitrep_9Jun2016_eng.p
df?ua=1.

28   Pan American Health Organization. Control of Yellow Fever: FIeld Guide. Washington DC,
USA, 2005
https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=1&ved=0ahUKEw
jwl5GlganNAhWFLyYKHUILDvgQFggeMAA&url=http://www.paho.org/hq/index.php?opti
on=com_docman&task=doc_download&gid=14606&Itemid=&usg=AFQjCNF2BDqUicDur
0YzeGR-I7Lkd1wvbQ.

Contributors

415
Declaration of interests
416
417 ML reports consulting honoraria (which have been donated to charity) from Pfizer
418 and Affinivax, and research funding through his institution from Pfizer and PATH Vaccine
419 Solutions, all unrelated to yellow fever. JTW, CMP, and GML have no conflicts of interest.
420
Acknowledgements
We thank Jack Woodall and Rebecca Grais for comments on an earlier draft and Alejandro
Costa and the students of EPI 260 for helpful discussions. This study was funded by Cooperative
Agreement U54GM088558 from the National Institute Of General Medical Sciences, National
Research Service Award T32AI007535-16A1, and a commissioned grant from the Health and
Medical Research Fund from the Government of the Hong Kong Special Administrative Region.
The content is solely the responsibility of the authors and does not necessarily represent the
official views of the National Institute Of General Medical Sciences or the National Institutes of
Health.
430

431   **Panel: Research in context**
432
Systematic review
We searched PubMed and Google Scholar on June 10, 2016, with the terms "yellow fever" and
"vaccine" or "dose sparing". We did not find any reports of randomized trials of yellow fever
(YF) vaccine efficacy, at full or lower doses. Three relatively recent studies suggest similar
immunological responses at five-fold, or more, fractionation as compared to the current dose
antigen levels.[8,9,26] While several recent perspective articles propose the dose-sparing strategy
in response to the current shortage,[2–4] to our knowledge this is the first study to test the
intuition behind the strategy and assess the implications of uncertainties surrounding
fractional-dose YF vaccine efficacy and mode of action (e.g. "all-or-nothing" and "leaky").
442
443   **Added value of the study**
444   Our study provides a formal confirmation of intuition that dose-sparing can drastically reduce
445   the number of YF cases if high vaccine efficacy is retained. We show how the benefits of dose
446   fractionation are influenced by the transmission intensity of the setting, the target coverage,
447   and the fractional-dose vaccine efficacy and mode of action.
448
449   **Interpretation**
450   Our results support the growing evidence that dose-sparing strategies should be explored as an
451   option for extending the currently sparse YF vaccine supply.
452

Figure 1: **Estimates of reproductive number over the course of the Angola epidemic. A**
Epidemic curve of confirmed cases by dates of symptom onset in Angola and vaccine coverage
in Luanda province achieved by the reactive YF vaccination campaign that started on 2 February
2016.[27] The first cases of this YF outbreak were identified in Luanda province which accounted
for 90 of the 121 cases confirmed in Angola up to 26 February 2016. **B-C** Estimates of the daily
reproductive number (R_t) assuming that the mean mosquito lifespan was 7 and 14 days,
respectively. The red data points correspond to the cases that were used to estimate the initial
reproductive number. These cases had symptom onset one mean serial interval before the
vaccination campaign began to affect disease transmission (which was assumed to be 7 days
after the start of the campaign to account for the time it takes for adaptive immunity to
develop). The orange and purple horizontal bars indicate the length of the mean mosquito
lifespan and serial interval on the scale of the x-axis, respectively.

Standard-dose vaccine coverage, V

Figure 2: **The impact of five-fold fractional-dose vaccination with different vaccine efficacy and reproductive numbers.** Vaccine action is assumed to be all-or-nothing and standard-dose vaccine efficacy is assumed to be 1. If the standard-dose vaccine coverage V exceeds 20%, then everyone in the population can be vaccinated under five-fold fractionated-dose vaccination, in which case the fractionation would only be n = $1/V$. **A** Infection attack rate (IAR) as a function of standard-dose vaccine coverage, V. The solid and dashed curves correspond to standard-dose and five-fold fractional-dose vaccination, respectively. IAR is reduced to 0 when the effective vaccine coverage (V for solid curves, $VE(n) \times nV$ for dashed curves) reaches the herd immunity threshold $(1-1/R_0)$x100%. **B** Absolute reduction in IAR. IAR reduction is maximum when the five-fold fractional-dose effective vaccine coverage $VE(5) \times 5V$ reaches the herd immunity threshold $(1-1/R_0)$x100%. As V increases from 0, a kink appears when the herd-immunity threshold is attained or everyone is vaccinated under five-fold fractional-dose vaccination (i.e., V = 20%). If five-fold fractional-dose vaccination at 100% coverage cannot attain the herd immunity threshold (because of low fractional-dose vaccine efficacy), then a second kink appears when V is large enough such that fractional-dose vaccination attains herd-immunity

485   threshold due to the increase in $VE(n)$ resulting from lower fractionation factors (namely $n =$
486     $1/V$). **C** Relative reduction in IAR.

Figure 3: Benefit thresholds for leaky vaccines as a function of standard dose vaccine supply *V*
and basic reproductive number *R*₀. Five-fold fractionated dosing will reduce IAR compared to
standard dosing if the leaky vaccine efficacy of fractional-dose is above the line corresponding
to the basic reproductive number. This threshold becomes high for large values of R_0 because
under the "leaky" model of vaccine efficacy, multiple exposures eventually lead to infection of
vaccinated individuals, overcoming their protection from the vaccine.

497 **Figure 4: Vaccination of adults with fractionated doses and children with standard doses.** All
498 children are vaccinated with standard-dose vaccines before any adults receive vaccination. **A**
499 Fold-increase in the proportion of individuals vaccinated conferred by five-fold fractionated
500 dose vaccination. **B-C** Same as Figure 2A-B when 57% of the population are adults and five-fold
501 fractional-dose vaccine efficacy is 60%.

Appendix

Estimation of the effective reproductive number for YF in Angola

We use the Wallinga and Teunis method[13] to estimate the reproductive number over the course of the YF outbreak in Angola from the daily number of confirmed cases recorded in the 17 April 2016 WHO Angola Situation Report.[14] We assume that all cases were attributed by local transmission, i.e. no importation of cases. Let t_i be the date of symptom onset for case i. The relative likelihood that case i has been infected by case j is

$$p_{ij} = \frac{w(t_j - t_i)}{\sum_{k \neq j} w(t_j - t_i)}$$

where $w(\cdot)$ is the probability density function of the serial interval. Assuming that the probability of case j infecting case i is independent of the probability of case j infecting any other case, the reproductive number for case j is a Bernoulli random variable with mean $\sum_i p_{ij}$. The reproductive number on day t, namely R_t, is approximated as the average of the reproductive number of all cases who have symptom onset on day t, in which case the mean and standard deviation of R_t are

$$E[R_t] = \frac{1}{n_t} \sum_{j:t_j=t} \sum_i p_{ij}$$

$$s(R_t) = \frac{1}{n_t} \sqrt{\sum_i \left(\sum_{j:t_j=t} p_{ij}(1-p_{ij}) - \sum_{j,k:t_j=t_k=t, j \neq k} p_{ij}p_{ik} \right)}.$$

Assuming that R_t is normally distributed, the approximate $(1-\alpha)\times100\%$ confidence interval is

$$E[R_t] \pm z_{1-\alpha/2} s(R_t).$$

Estimation of the serial interval distribution for YF

We assume that the latent period is the same as the incubation period for all human infections of YF. Suppose an infected individual becomes infectious at time 0. Let t_1 be the time at which the infectious individual is bitten by a competent mosquito which becomes infected, t_2 be the time at which this mosquito becomes infectious, and t_3 be the time at which this mosquito bites and infects a human host. The probability distribution function for the serial interval is

$$f(a) = \frac{h(a)}{\int_0^\infty h(u)du}$$

where

$$h(a) = \int_0^a \int_0^{t_3} \int_0^{t_2} \underbrace{P(I > t_1)}_{\substack{\text{Probability that the}\\\text{human infectious period}\\\text{exceeds } t_1 \text{ days when}\\\text{the mean infectious duration}\\\text{is mean 4 days.}}} \cdot \underbrace{f_V(t_2 - t_1)}_{\substack{\text{Extrinsic incubation period}\\\text{at 28 degree Celsius; Weibull}\\\text{distributed with mean 12.7 days}\\\text{and CoV 0.61}}} \cdot \underbrace{e^{-d(t_3 - t_1)}}_{\substack{\text{Probability that the}\\\text{mosquito is still alive}\\t_3 - t_1 \text{ days after getting}\\\text{infected}}} \cdot \underbrace{f_H(a - t_3)}_{\substack{\text{Intrinsic incubation period;}\\\text{Lognormal distributed with}\\\text{mean 4.6 days and CoV 0.36}}} \quad dt_1 dt_2 dt_3$$

In this calculation, we assume that the infectious period in humans is exponentially distributed
with mean 4 days,[28] and mosquito lifespan is exponentially distributed with mean varying over
1-2 weeks (http://www.dengue.gov.sg/subject.asp?id=12;[17]). We assume that the extrinsic
incubation period follows the Weibull distribution with parameters $\nu = 1.7$ and
$\lambda_i = \exp(-7.6 + 0.11T)$ where T is the temperature (28 degrees Celsius) as estimated by ref. [15]
We assume that the intrinsic incubation period follows the lognormal distribution with
parameters $\mu = 1.46$ and $\tau = 8.1$ as estimated by ref. [15].

Dose-response relationship
We assume that vaccine efficacy of n-fold fractional-dose vaccines for n between 1 and 5
increases linearly with the amount of antigen in the vaccines which is proportional to $1/n$. In
general, if vaccine efficacy of n-fold fractional-dose vaccines for n between n_1 and n_2
increases linearly with the amount of antigen in the vaccines, then
$VE(n) = VE(n_2) + \dfrac{1/n - 1/n_2}{1/n_1 - 1/n_2}\left(VE(n_1) - VE(n_2)\right)$. We make this assumption to avoid
overestimating the benefit of dose-fractionation because:

1. If $VE(5)$ is at the all-or-nothing benefit threshold, namely $VE(1)/5$, then $VE(n)$ is also
at the benefit threshold (i.e. $VE(n) = VE(1)/n$) for all n between 1 and 5. That is, if five-
fold dose fractionation is not beneficial, then dose-fractionation is not beneficial for all
fractionation below five-fold.
2. The reduction in vaccine efficacy as fractionation increases from 1 is likely to be more
gradual than what we have assumed here given that standard dose vaccine efficacy
appears to be close to 100%.

Appendix Figure 1 illustrates this dose-response relationship for different values of $VE(5)$ with
$VE(1) = 1$.

Infection attack rate
We first provide mathematical details on IAR calculations for the case where the population is
not stratified into subgroups. If vaccine action is all-or-nothing, then IAR with fractionation n,
denoted by $IAR(n)$, is obtained by solving the equation

$$IAR(n) = S_0 \left(1 - VE(n)nV\right)\left[1 - \exp(-R_0 \cdot (I_0 + IAR(n)))\right]$$

where R_0 is the basic reproductive number, S_0 and I_0 are the initial proportion of population
that are susceptible and infectious. As such, dose-fractionation reduces IAR if and only if
$VE(n) > VE(1)/n$. If vaccine action is leaky, then $IAR(n)$ is obtained by solving the equation

562
$$IAR(n) = S_0 \left(1 - Vn\right)\left[1 - \exp(-R_0 \cdot (I_0 + IAR(n)))\right]$$
$$+ S_0 Vn\left[1 - \exp(-(1 - VE(n))R_0 \cdot (I_0 + IAR(n)))\right]$$

In this case, dose-fractionation reduces IAR if and only if

$$VE(n) > 1 + \frac{\ln(1 - Z)}{R_0(I_0 + IAR(1))} \quad \text{where} \quad Z = \frac{IAR(1)}{S_0 Vn} - \left(\frac{1}{Vn} - 1\right)\left[1 - \exp(-R_0 \cdot (I_0 + IAR(1)))\right]$$

In the special case where $VE(1) = 1$, the benefit threshold can be simplified as

$$VE(n) > 1 - \frac{\ln\left(1 - (1 - 1/n)\dfrac{IAR(1)}{S_0(1 - V)}\right)}{\ln\left(1 - \dfrac{IAR(1)}{S_0(1 - V)}\right)}.$$

Next, we provide mathematical details on IAR calculations for the general case where there are
m groups. Let $S_{0,i}$ and $I_{0,i}$ be the proportion of susceptible and infectious people in group i just
before the vaccination campaign begins. Let V_i be the vaccine coverage of standard-dose
vaccines for group i. If n_i is the fractionation for group i, then vaccine coverage of fractional-
dose vaccines for group i is $V_i n_i$. Let $R_0^{j,i}$ be the expected number of secondary infections in
group j caused by one infection in group i in a completely susceptible population. If vaccine
action is all-or-nothing, the group-specific IARs are obtained by solving the equations

$$IAR_i(n_i) = S_{0,i}\left(1 - V_i n_i VE(n_i)\right)\left[1 - \exp\left(-\sum_j R_0^{j,i}(I_{0,j} + IAR_j(n_i))\right)\right]$$

If vaccine action is leaky, then the group-specific IARs are obtained by solving the equations

576
$$IAR_i(n_i) = S_{0,i}\left(1 - n_i V_i\right)\left[1 - \exp\left(-\sum_j R_0^{j,i}(I_{0,j} + IAR_j(n_i))\right)\right]$$
$$+ S_{0,i} n_i V_i\left[1 - \exp\left(-(1 - VE(n_i))\sum_j R_0^{j,i}(I_{0,j} + IAR_j(n_i))\right)\right]$$

Heterogeneity in biting rates
A recent study suggested that the mosquito biting rate for individuals aged 20 or above is 1.22
times higher than those age under 20.[24] To test the robustness of our results against such
heterogeneity, we repeat the calculations in Figure 2 and 3 using a model in which the
population is stratified with age 20 as the cutoff. For illustration, we use the demographic
parameters of Angola where around 55% of the population are under 20. Appendix Figures 2-3
show that our results are unaffected by heterogeneity in biting rates.

Appendix Figure 1. The dose response relationship assumed in the model with *VE*(1) = 1.

590
591
592 **Appendix Figure 2. Repeating the calculations in Figure 2 using a 2-age-group model in which those 20**
593 **or older were 1.22 times more likely to be bitten by mosquitos compared to those under age 20.** The
594 results are essentially the same as that in Figure 2.
595

596
597
598 **Appendix Figure 3. Repeating the calculations in Figure 3 using a 2-age-group model in which those 20**
599 **or older were 1.22 times more likely to be bitten by mosquitos compared to those under age 20.** The
600 solid and dashed curves show the results without and with age stratification, respectively.
601
602