1 Fractional Dosing of Yellow Fever Vaccine to Extend Supply: A Modeling Study
2

3 **AUTHORS**
4 Joseph T. Wu[1], PhD, joewu@hku.hk*
5 Corey M. Peak[2], MS
6 Gabriel M. Leung[1], MD
7 Marc Lipsitch[2,3], DPhil mlipsitc@hsph.harvard.edu*
8

[1]WHO Collaborating Centre for Infectious Disease Epidemiology and Control, School of Public
Health, Li Ka Shing Faculty of Medicine, The University of Hong Kong, Hong Kong Special
Administrative Region, China
[2] Center for Communicable Disease Dynamics, Department of Epidemiology, Harvard T.H. Chan
School of Public Health, 677 Huntington Avenue, Boston MA 02115 USA
[3] Department of Immunology and Infectious Diseases, Harvard T.H. Chan School of Public
Health, 665 Huntington Avenue, Boston MA 02115 USA. +1-617-432-4559.
* corresponding authors

17
18

Background
The ongoing yellow fever (YF) epidemic in Angola is placing strain on the global vaccine supply.
In order to extend vaccine supply and reduce the cost of mass-vaccination, dose sparing by
fractional-dose vaccination has received heightened consideration. Five-fold fractionation is
similar to the standard dose in safety and immunogenicity. However, no YF vaccine efficacy
trials have been performed in humans, so it is possible that fractional-dose vaccines may be less
efficacious even if equally immunogenic. There is an urgent need to study under what
conditions fractional dosing could provide epidemiologic benefits in reducing transmission.
27

Methods
We estimated the effective reproductive number for YF in Angola using disease natural history
and case report data. Using these results and simple mathematical models of YF transmission,
we calculated the expected final size of an epidemic under varying levels of vaccine coverage
with standard-dose vaccine and up to five-fold fractionation with varying efficacy. We consider
three allocation scenarios: random; targeted at only susceptible individuals; and whereby
children receive standard-dose vaccines while adults receive fractional-dose vaccines.
35

Findings
The effective reproductive number early in the outbreak ranged from approximately 5 to 12
transmission events per infectious individual. As expected, if five-fold fractional-doses retain
100% efficacy, the final epidemic is dramatically reduced, especially if standard-dose vaccine
coverage is near 20%. If instead some fractional-dose recipients do not become immunized, the
five-fold fractional-dose strategy is always beneficial, as long as this dose produces immunity in
at least 20% of recipients. We quantify how the threshold becomes more stringent if vaccine
action is leaky.

Interpretation

We conclude that dose fractionation could be a very effective strategy for improving coverage

of YF vaccines and reducing infection attack rate in populations, possibly by a large absolute

and relative margin, if high to moderate efficacy is maintained by reduced-dose formulations.

Funding

INTRODUCTION

Yellow fever (YF) has resurged in Angola and threatens to spread to other countries with relatively low YF vaccine coverage. While the rainy season is ending in Angola as of April, it typically lasts through May in the Democratic Republic of Congo (DRC) and is just beginning in parts of West Africa. Reports of YF cases in Kinshasa, DRC, in mid-April 2016 and exportation of cases from Angola to other parts of the world raise concern that YF could resurge in other populations where competent vectors are present and vaccine coverage is low.[1] A broad band of sub-Saharan Africa north of Namibia and Zambia is at risk (http://www.cdc.gov/yellowfever/maps/africa.html), as is much of the northern portion of South America (http://www.cdc.gov/yellowfever/maps/south_america.html). The global community is increasingly concerned for the risk of YF emergence in Asia, where the disease has been curiously absent despite seemingly amenable conditions.

There is a safe, highly effective vaccine against YF.[2] However, the global stockpile of YF vaccines is low, with approximately 10 million doses currently available and a global manufacturing capacity of 25 million doses per month. Given the large populations at risk for YF infection, the stockpile might well be inadequate to meet the need; the entire current stockpile would be required just to vaccinate the population of Kinshasa. For this reason, the possibility of dose-sparing by fractional-dose vaccination is under consideration, in which smaller volumes of vaccine would be used per dose in order to increase the number of persons who can be vaccinated with a given quantity of vaccine.[2] Dose-fractionation has been proposed as a means of extending the supply and reducing the cost of mass-vaccination campaigns for YF; these benefits might also enhance equity of vaccine access.[3]

A randomized, noninferiority trial has shown that 0·1 ml intradermal (ID) vaccination with the 17D YF vaccine was equally safe compared to the standard 0·5ml subcutaneous dose.[4] Another randomized trial of the 17DD vaccine given in Brazil showed that there was no significant loss of immunogenicity when the currently administered vaccine (containing 27,476 IU of virus) was given at doses as low as 2·1% (i.e. a >40-fold reduction) of the full dose.[5] The lowest reduced dose in that study that was noninferior to the full dose was 587 IU. For comparison, the WHO minimum for YF vaccines is 1000 IU per dose.[6] No efficacy trial of YF vaccines has been performed in humans,[7] so the comparative efficacy of different doses and routes remains unknown.

While the equal immunogenicity of fractional-dose vaccines suggests that efficacy might also be equal to that of the standard dose, decision makers may be cautious about changing dosing recommendations. Challenges to such a change include obtaining regulatory approval, training vaccinators to deliver fractional doses, possibly by the intradermal route, and other logistical issues.[2]

Beyond these operational challenges, which we do not specifically consider in our modeling study, it is possible that the dose-sparing vaccines may be less efficacious even if equally immunogenic. Moreover, the findings of equal immunogenicity of reduced doses are limited to

adults; no comparable data exist in children. Here we use simple mathematical models to
assess the magnitude of benefit that could be achieved by dose-fractionation, by allowing
higher or more rapid vaccine coverage with a fixed supply of vaccine. Our initial analysis
assumes that the efficacy of five-fold fractional-dose vaccine[2,4] is equal to that of standard-dose
vaccine, and that both are 100% effective,[7] and estimates the benefit of higher coverage with
no loss of protection for individual vaccinees. We next show that even if vaccine efficacy were
considerably lower for fractional-dose vaccination, the population-level benefit of wider
coverage could still outweigh the lower individual-level efficacy, in terms of total cases
prevented. Finally, given the lack of comparative immunogenicity data for fractional-dose YF
vaccination in children, we consider the results of a strategy that provides standard-dose
vaccines to children and fractional-dose vaccines to adults. We find that each of these
strategies could provide significant benefit epidemiologically, and that the best policy will be
determined by balancing logistical and regulatory considerations against the extent of
epidemiologic benefit.

METHODS

Estimation of the effective reproductive number for YF in Angola

We used data on confirmed cases each day from the 18 April 2016 WHO Angola Situation
Report[8] and applied the method of Wallinga and Teunis[9]. We estimated the probability
distribution function for the generation time using the convolution of the intrinsic and extrinsic
incubation periods of yellow fever estimated by [10], with the assumption that the latent and
incubation period in the human host were the same. In the integral, t_1 gives the time post-
infection when the human host becomes infectious, t_2 gives the time at which he is bitten by a
competent mosquito, and t_3 is the time at which the mosquito becomes infectious. We assume
an exponentially distributed infectious period with mean 4 days in the human,[11] and an
exponentially distributed mosquito lifespan varying over a broad range around 2 weeks
(http://www.dengue.gov.sg/subject.asp?id=12). This expression corresponds to the
approximating assumption that once infected and infectious, vectors bite at a constant rate
until they die. Thus the probability distribution function f_{GT} for the generation time a is

121
$$f_{GT}(a) = \frac{h(a)}{\int_0^\infty h(u)\,du}$$

where

$$h(a) = \int_0^a \int_0^{t_3} \int_0^{t_2} \underbrace{f_H(t_1)}_{\substack{\text{Intrinsic incubation period;}\\ \text{Lognormal distributed with}\\ \text{mean 4.6 days and CoV 0.36}}} \cdot \underbrace{P(I > t_2 - t_1)}_{\substack{\text{Probability that the}\\ \text{human infectious period}\\ \text{exceeds } t_2-t_1 \text{ days when}\\ \text{the mean infectious duration}\\ \text{is mean 4 days.}}} \cdot \underbrace{f_V(t_3 - t_2)}_{\substack{\text{Extrinsic incubation period}\\ \text{at 28 degree Celsius; Weibull}\\ \text{distributed with mean 12.7 days}\\ \text{and CoV 0.61}}} \cdot \underbrace{e^{-d(a-t_2)}}_{\substack{\text{Probability that the}\\ \text{mosquito is still alive}\\ a-t_2 \text{ days after getting}\\ \text{infected}}} dt_1 dt_2 dt_3$$

We assumed that serial interval and generation time have the same probability distribution.

Infection attack rate calculations

Our basic approach was to calculate the expected final size of an epidemic (infection attack rate
or IAR) under the following homogeneous-mixing model. Let S_0 and I_0 be the proportion of
population susceptible and infectious just before vaccination. Let V be the vaccine coverage of
standard-dose vaccines. Suppose each standard-dose vaccine can be fractionated into n n-fold
fractional-dose vaccines (i.e. each n-fold fractional-dose vaccine contains 1/n-th amount of the
antigen in a standard-dose vaccine). We denote the vaccine efficacy of n-fold fractional-dose
vaccines by $VE(n)$, i.e. the vaccine efficacy of standard-dose vaccines is $VE(1)$. Given V, the
highest fractionation factor sensible is $n_{max} = S_0/V$ if the susceptible population can be
identified for targeted vaccination and $n_{max} = 1/V$ otherwise, i.e. the fractionation factor n
(hereafter, fractionation) must lie between 1 and n_{max}. Let $Q(V,n) = V n VE(n) w(n)$ be the

effective vaccine coverage assuming all-or-nothing vaccine action, where $w(n)$ is the

proportion of fractionated doses that do not get wasted. As such, the optimal fractionation that

gives the smallest IAR is the one that maximizes $nVE(n)w(n)$. Compared to standard-dose

vaccination, dose fractionation is better if and only if $nVE(n)w(n) > VE(1)$. That is, in the

absence of wastage, n-fold dose fractionation is better if and only if the vaccine efficacy of n-

fold fractional-dose vaccines are at least 1/n times that of standard-dose vaccines. The criterion

is substantially higher for leaky vaccines when transmissibility is high (see below). We assumed

that vaccine efficacy of n-fold fractional-dose vaccines is a linear function of the amount of

antigen in the vaccines (which is proportional to 1/n). That is, for n between 1 and 5,

$VE(n) = VE(5) + 1.25(1/n - 1/5)(VE(1) - VE(5))$.

We consider three scenarios:

1. *Random vaccination.* Vaccine coverage among susceptible and immune individuals are

148   the same, i.e. the susceptible and immune population are indiscernible, so targeted

149   vaccination is not possible. For any given fractionation n, the infection attack rate

150            $IAR(n)$ is obtained by solving the equation

151    $IAR(n) = S_0\left(1 - Q(V,n)\right)\left[1 - \exp(-R_0 \cdot (I_0 + IAR(n)))\right]$.

152   2. *Targeted vaccination.* Vaccinations are targeted only at susceptible individuals. For any

153   given fractionation n, the infection attack rate $IAR(n)$ is obtained by solving the

154   equation

$$\begin{aligned}IAR(n) &= \left(S_0 - Q(V,n)\right)\left[1 - \exp(-R_0 \cdot (I_0 + IAR(n)))\right]\\&= S_0\left(1 - Q(V,n)/S_0\right)\left[1 - \exp(-R_0 \cdot (I_0 + IAR(n)))\right]\end{aligned}$$

156   Hence, targeted vaccination has the same effects as random vaccination with effective

157   vaccine coverage $Q(V,n)/S_0$. We do not provide graphs of this scenario separately.

158   3. *Standard-dose vaccination of children, fractional-dose vaccination of adults*. We assume

159            $S_0 = 1$. Let p be the proportion of adults in the population. We assumed that children

160   have vaccine priority. That is, all children receive standard-dose vaccines before adults

161   begin to receive fractional-dose vaccines. For a given standard-dose vaccine coverage V

162   , the proportion of children vaccinated is $V_{children} = \min(V, 1-p)$. If the stockpile is large

163   enough to vaccinate all children (i.e. $V > 1-p$), then adults receive n-fold fractional-

164   dose vaccination where $n = \min(5, p/(V-1+p))$. The proportion of adults vaccinated is

165            $V_{adults} = 1$ if $n < 5$ and $V_{adults} = 5(V-1+p)/p$ otherwise. The effective vaccine coverage is

166            $Q(V,n) = V_{children}(1-p)VE(1) + V_{adults}\, p\, VE(n)w(n)$.

Leaky vaccine action

If vaccine efficacy is "leaky" (such that its efficacy is the independent probability of protecting
an immunized recipient on each exposure[12,13]) then the attack-rate calculations change slightly.
Suppose the expected number of secondary cases is the same for vaccinated and unvaccinated
individuals if they become infected. Given fractionation $n = \min(5, 1/V)$, the infection attack
rate $IAR(n)$ is obtained by solving the equation

174
$$IAR(n) = S_0\left(1 - Vn\right)\left[1 - \exp(-R_0 \cdot (I_0 + IAR(n)))\right]$$
$$+ S_0 Vn\left[1 - \exp(-(1 - VE(n))R_0 \cdot (I_0 + IAR(n)))\right]$$

175 if vaccination is random. Dose fractionation is better if and only if

176
$$VE(n) > \frac{1}{VE(1)}\left(1 + \frac{\ln(1 - Z)}{R_0(I_0 + IAR(1))}\right) \quad \text{where}$$
$$Z = \frac{IAR(1)}{Vn} - \left(\frac{1}{Vn} - 1\right)\left[1 - \exp(-R_0 \cdot (I_0 + IAR(1)))\right].$$

We define the right-hand side of this inequality as the "benefit threshold," the lowest leaky
efficacy that is needed for a fractional-dose strategy to produce a lower IAR than a standard-
dose strategy.

As in the case for all-or-nothing vaccines, the effects of targeted vaccination are the same as
that for random vaccination with effective vaccine coverage $Q(V, n)/S_0$ for leaky vaccines.

RESULTS

Reproductive number of Yellow Fever in Angola. Figure 1 shows the epidemic curve of yellow
fever cases in Angola[8] (A) and the estimated reproductive number R_t over the course of the
outbreak, estimated using natural history data from ref. [10] and the Wallinga-Teunis method.[9]
Early in the outbreak, estimates of R_t range from approximately 5 (lowest point estimate) to 12
(highest upper bound of 95% confidence interval). While these estimates may reflect partial
immunity due to vaccination or prior exposure among some of the population, we take this as
the center of a range for the possible basic reproductive number in a future outbreak in
another population, which we assume may be between 3 and 12, due to varying vector ecology
and levels of preexisting immunity in the population.

Reducing attack rate by increasing coverage. Figure 2A shows the attack rate expected for a
sustained epidemic in a simple mathematical model given varying levels of transmission (basic
reproductive number R_0, different curves) and standard-dose vaccine coverage (V, assumed to
be achieved prior to the epidemic's start) with and without five-fold dose-fractionation,
assuming 100% efficacy. Figures 2B-C show the corresponding absolute and relative reduction
in attack rate. Clearly a five-fold increase in coverage can dramatically reduce the attack rate,
by at least five-fold, and by more if the reproductive number is brought close to or below one
by herd immunity effects, achieved as vaccine coverage approaches the herd immunity
threshold $(1-1/R_0) \times 100\%$.

Lower-efficacy assumption. Figure 3 shows the robustness of the dose-fractionation strategy to
the possibility that fractionated doses have lower efficacy. In Figure 3, we repeat the
calculations of reduced attack rate from Figure 2, but assuming that fractional-dose vaccines
provide full immunity to 90%, 60%, 30% and 10% of those vaccinated (all-or-nothing
efficacy).[12,13] We find that the five-fold fractional-dose strategy is always beneficial, regardless
of the achievable coverage V with the standard-dose vaccine or the basic reproductive number
R_0, as long as five-fold fractional-dose vaccines are at least 20% efficacious. Mathematically, it
can be shown that for efficacy greater than 100%/n, an n-fold fractionation is always
advantageous in the sense of a lower attack rate expected with fractionated doses and higher
coverage than standard dosing at n-fold lower coverage. If we assume there is some wastage of
fractionated doses due to unfamiliar administration methods and lower seroconversion rates,
this threshold becomes slightly less favorable because x% wastage would cause a x% reduction
in effective vaccine coverage, which is a product of vaccine coverage, vaccine efficacy, and
wastage. For example, five-fold fractionation with no wastage and 20% vaccine efficacy has the
same effect as 20% wastage and 25% vaccine efficacy, which is still well above that which
seems plausible from the immunogenicity data.

Vaccination of adults with fractionated doses and children with standard doses. Given the
lack of immunogenicity data for fractionated doses in children and evidence of lower
seroconversion rates to standard doses,[14] a conservative strategy would be to fractionate doses
only for adults. Figure 4A shows the fold-increase in vaccine coverage that could be achieved
with five-fold fractionation in adults only, as a function of proportion of adults in the population

(P_{adults}). Adult fractionation increases coverage more if a larger fraction of the population is

adults. In Angola in 2015 approximately 70% of the population was adults (20 and older). If

there were enough standard-dose vaccine supplies to cover one half of such a population

(V=0.5), fractionation of only the adult doses would provide a 2-fold increase in coverage. Given

the vector-borne nature of yellow fever we assume that transmission between children and

adults is well-mixed. Figure 4B shows the same calculations as Figure 2B with the five-fold

fractional-dose vaccine efficacy is 60%. The benefits of n-fold fractionation for only adult doses

is maximized when standard-dose coverage $V = P_{children} + P_{adults} / n$. These calculations all assume

that there is no preexisting immunity in children or adults; if preexisting immunity existed

mainly in adults, then prioritizing children would have a greater benefit than projected here.

Leaky vaccine efficacy. Thus far, when considering the possibility that fractionated doses have

VE<100%, we have assumed that this follows the "all-or-nothing model" of 100% efficacy in a

proportion VE of vaccinees and 0% in the remainder. Alternatively, reduced efficacy could take

the form of less than 100% protection in all vaccinated persons. We consider this unlikely (see

Discussion) but note that in this situation, the efficacy of n-fold fractionated doses necessary to

provide a benefit in terms of IAR is higher than 1/n and is dependent on transmission intensity,

with higher efficacy required to provide an advantage in higher-transmission settings. This

occurs because under the "leaky" model each infectious bite is assumed to be less likely to

cause infection if the host is vaccinated, but the probability of infection grows as the person

receives more infectious bites. Figure 5 shows, under the "leaky" model of vaccine action, the

"benefit threshold," the minimum efficacy of fractionated vaccine dose for which greater

coverage outweighs lower efficacy. Dose fractionation is much less beneficial if vaccine action is

leaky, efficacy is modest, and R_0 is high.

DISCUSSION

Our primary analysis shows that dose-fractionation of YF vaccine, if there were no loss of
efficacy, could provide a substantial benefit to reducing the attack rate of YF in a population.
We consider this assumption of full efficacy for five-fold fractionation to be the most likely
scenario, despite the lack of efficacy data on any YF vaccine, for several reasons: 1) two studies
of five- or greater-fold vaccination doses have shown indistinguishable immunogenicity in
humans; 2) at least some preparations of YF vaccine substantially exceed the WHO minimum
standard for potency of 1000 IU/dose, so fractionation at some level could be performed
without dropping below that threshold; 3) YF vaccine is live attenuated virus, so a biological
rationale exists that if a productive vaccine-virus infection can be established by a fractionated
dose, protection should be comparable to that with a higher dose. Nonetheless, to test the
robustness of our analysis, we considered the possibility that five-fold fractionated dosing fails
to immunize a proportion (1-VE(5)) of recipients, and found that as long as at least 20% of
recipients are fully immunized by the vaccine, the population-wide benefits of higher coverage
would outweigh the lower efficacy of fractionated dosing for individual vaccinees. Even more
unlikely, in our opinion, is that fractionated doses would be less efficacious according to a
"leaky" model, in which all vaccinated individuals were imperfectly protected against infection
from each infectious bite, with the same probability of infection from each bite, reduced by
vaccine by a proportion VE. If this were the case, then especially in high-transmission areas, the
fractionated-dose vaccine would need to be 80-90% efficacious to provide a benefit over
standard dosing.

We have used five-fold fractionation as an example because it is the strategy with the best
evidence base of equal immunogenicity. However, some data suggest that more than five-fold
fractionation could be equally immunogenic, and of course the benefits of fractionation would
be greater if more than five-fold fractionation were logistically possible and comparably
efficacious.

On programmatic grounds a simpler strategy -- such as fractionated dosing for all -- may be
preferred to a more complex strategy that gives different doses to different groups, say age
groups. While either would provide epidemiologic benefit, the choice between such strategies
would be influenced by the number of available doses, logistical barriers, and location-specific
regulations regarding specific groups, such as children.

Our simple model has several limitations. We have assumed homogeneous mixing of the
population (reasonable at least locally for a vector-borne disease) and have neglected
preexisting immunity in our main results, though the Methods show how our calculations could
be modified to consider preexisting immunity. The purpose was to provide basic calculations for
the most at-risk populations, those with little preexisting immunity. We have also fixed a
particular value of R_0 for each calculation, and assumed this value is maintained until the
epidemic has swept through a population. In reality, R_0 will vary seasonally as vector
abundance, extrinsic incubation period, and other factors vary. The existence of a high-
transmission season might enhance the benefits of fractional-dose vaccination. Most

importantly, there will be a premium on achieving high vaccine coverage before the peak of
transmission to maximally impact transmission, and this will be limited by supply constraints
that could be partially relieved by fractionation.

We conclude that dose fractionation could be a very effective strategy for improving coverage
of YF vaccines and reducing infection attack rate in populations – possibly by a large absolute
and relative margin -- if high to moderate efficacy is maintained by reduced-dose formulations.
For vaccines whose standard formulations exceed WHO minimum concentration of viral
particles,[6] this dose-fractionation could be accomplished without changing the WHO
recommendations. Even if the efficacy of fractionated doses were substantially lower,
increasing coverage by a factor greater than the reduction in efficacy would still be predicted to
reduce the population-wide infection attack rate. Substantial benefits could also be achieved if
fractional doses were given only to adults while providing standard-dose vaccines to children.
We urge consideration of means to implement dose-fractionation as a component of a YF
response strategy for the current situation. Rollout of fractionated dosing should perhaps be
preceded or accompanied by noninferiority studies of the intended vaccine's immunogenicity in
the intended populations. Ongoing programs should be monitored by observational studies of
safety, immunogenicity and, if possible, effectiveness[15] to assure that the assumptions
underlying the rationale for such programs continue to be met.

REFERENCES

1   Woodall J, Yuill T. Why is the yellow fever outbreak in Angola a 'threat to the entire
305   world'? *Int J Infect Dis* 2016; : 1–4.

306   2   Monath TP, Woodall JP, Gubler DJ, *et al.* Yellow fever vaccine supply: a possible solution.
307   *Lancet* 2016; **387**: 1599–600.

308   3   Brearley L, Eggers R, Steinglass R, Vandelaer J. Applying an equity lens in the Decade of
309   Vaccines. *Vaccine* 2013; **31**: B103–7.

310   4   Roukens AH, Vossen AC, Bredenbeek PJ, van Dissel JT, Visser LG. Intradermally
311   Administered Yellow Fever Vaccine at Reduced Dose Induces a Protective Immune
312   Response: A Randomized Controlled Non-Inferiority Trial. *PLoS One* 2008; **3**.

313   5   Martins RM, Maia M de LS, Farias RHG, *et al.* 17DD yellow fever vaccine: a double blind,
314   randomized clinical trial of immunogenicity and safety on a dose-response study. *Hum*
315   *Vaccin Immunother* 2013; **9**: 879.

316   6   World Health Organization. Expert Committee on Biological Standardization.
317   Requirements for yellow fever vaccine. Geneva, Switzerland, 2008.

318   7   Staples JE, Gershman M, Fischer M. Yellow Fever Vaccine: Recommendations of the
319   Advisory Committee. *Morb Mortal Wkly Rep* 2010; **59**: 1–27.

320   8   World Health Organization. Yellow fever outbreak in Angola: Situation Report. 2016
321   http://www.afro.who.int/index.php?option=com_docman&task=doc_download&gid=10
322   121&Itemid=2593.

323   9   Wallinga J, Teunis P. Different Epidemic Curves for Severe Acute Respiratory Syndrome
324   Reveal Similar Impacts of Control Measures. *Am J Epidemiol* 2004; **160**: 509–16.

325   10   Johansson MA, Arana-Vizcarrondo N, Biggerstaff BJ, Staples JE. Incubation Periods of
326   Yellow Fever Virus. *Am J Trop Med Hyg* 2010; **83**: 183–8.

327   11   Pan American Health Organization. Control of Yellow Fever: FIeld Guide. Washington DC,
328   USA, 2005.

329   12   Smith PG, Rodrigues LC, Fine PEM. Assessment of the Protective Efficacy of Vaccines
330   against Common Diseases Using Case-Control and Cohort Studies. 1984; **13**.

331   13   Halloran ME, Longini IM, Struchiner CJ. Design and Analysis of Vaccine Studies:
332   Introduction. New York, NY: Springer, 2010.

333   14   Belmusto-worn VE, Sanchez JL, Carthy KMC, *et al.* Randomized, double–blind, phase III,
334   pivotal field trial of the comparative immunogenicity, safety and tolerability of the two
335   yellow fever 17D vacines (Arilvax and YF-Vax) in healthy infants and children. *Am J Trop*
336   *Med Hyg* 2005; **72**: 189–97.

337   15   Hickling J, Jones R, Tandem W. Yellow fever vaccination : The potential of dose- sparing
338   to increase vaccine supply and. *Path* 2013. http://www.path.org/news/press-room/427/.

339   16   Riley S, Wu JT, Leung GM. Optimizing the dose of pre-pandemic influenza vaccines to
340   reduce the infection attack rate. *PLoS Med* 2007; **4**: 1032–40.

Contributors

JTW, CMP, and ML reviewed the literature and designed the study. JTW and ML developed the mathematical model. JTW ran the mathematical model. JTW, CMP, GML, and ML interpreted the model results and approved the final version.

Declaration of interests

ML reports consulting honoraria (which have been donated to charity) from Pfizer and 323 Affinivax, and research funding through his institution from Pfizer and PATH Vaccine Solutions, 324 all unrelated to yellow fever. JTW, CMP, and GML have no conflicts of interest.

Acknowledgements

We thank Jack Woodall and Rebecca Grais for comments on an earlier draft and Alejandro Costa and the students of EPI 260 for helpful discussions. This study was funded by Cooperative Agreement U54GM088558 from the National Institute Of General Medical Sciences, National Research Service Award T32AI007535-16A1, and a commissioned grant from the Health and Medical Research Fund from the Government of the Hong Kong Special Administrative Region. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute Of General Medical Sciences or the National Institutes of Health.

Figure 1: **Estimates of reproductive number over the course of the Angola epidemic.** Panels show estimates for varying assumptions about the mean vector lifespan, which generate varying estimates of the mean serial interval.

Figure 2: **The impact of five-fold fractional-dose vaccination with no reduction in vaccine**
efficacy. A Infection attack rate (IAR) as a function of standard-dose vaccine coverage, V. The
solid and dashed curves correspond to standard-dose and five-fold fractional-dose vaccination,
respectively. IAR is reduced to 0 when coverage (V for solid curves, 5V for dashed curves)
reaches the herd immunity threshold $(1-1/R_0) \times 100\%$. **B-C** Absolute and relative reduction in IAR
conferred by five-fold fractional-dose vaccination. IAR reduction is maximum when the
fractional-dose vaccine coverage, namely 5V, reaches $(1-1/R0) \times 100\%$.

Figure **3: The impact of five-fold fractionated dose vaccination with reduced vaccine efficacy**
(all-or-nothing assumption) and different reproductive numbers. If the standard-dose vaccine
coverage V exceeds 20%, then everyone in the population can be vaccinated under five-fold
fractionated-dose vaccination, in which case the fractionation factor would only be n = 1/V.
Vaccine efficacy of n-fold fractional-dose vaccines, denoted by VE(n), is assumed to be a linear
function of vaccine dose (see Methods). **A** IAR. **B** Absolute reduction in IAR. As V increases, a
kink appears when herd-immunity threshold is attained or everyone is vaccinated under five-
fold fractional-dose vaccination (i.e., V = 0·2). If five-fold fractional-dose vaccination at 100%
coverage cannot attain the herd immunity threshold (because of low fractional-dose vaccine
efficacy), then a second kink appears when V is large enough such that fractional-dose
vaccination attains herd-immunity threshold due to the increase in VE resulting from lower
fractionation factors (namely 1/V). **C** Relative reduction in IAR.

389 *Figure* 4: **Vaccination of adults with fractionated doses and children with standard doses.** All
390 children are vaccinated with standard-dose vaccines before any adults receive vaccination. **A**
391 Fold-increase in the proportion of individuals vaccinated conferred by five-fold fractionated
392 dose vaccination. **B** Same as Figure 2B when 70% of the population are adults and five-fold
393 fractional-dose vaccine efficacy is 60%.

395   *Figure* **5: Benefit thresholds for leaky vaccines as a function of standard dose vaccine supply V**
396   **and basic reproductive number R_0.** Five-fold fractionated dosing will produce an infection
397   attack rate lower than standard dosing if the "leaky" vaccine efficacy of a fractionated dose is
398   above the line corresponding to the basic reproductive number. This threshold becomes high
399   for large values of R0 because under the "leaky" model of vaccine efficacy, multiple exposures
400   eventually lead to infection of vaccinated individuals, overcoming their protection from the
401   vaccine. Although we consider this model unlikely to apply to YF vaccines (and best evidence
402   suggests that five-fold fractionated doses are likely to be nearly 100% efficacious), we include
403   this as a formal possibility.