1
2
3
4 ***Cassava brown streak virus* has a rapidly evolving genome: implications for virus**
5 **speciation, variability, diagnosis and host resistance**
6
7
8

9 Titus Alicai[1], Joseph Ndunguru[2], Peter Sseruwagi[2], Fred Tairo[2], Geoffrey Okao-Okuja[1], Resty

10 Nanvubya[1], Lilliane Kiiza[1], Laura Kubatko[3], Monica A. Kehoe[4] and Laura M. Boykin[5,*]

11 [1]National Crops Resources Research Institute, P.O. Box 7084, Kampala, Uganda
12
13 [2]Mikocheni Agricultural Research Institute, Coca cola Road, Box 6226, Dar es Salaam, Tanzania
14
15 [3]The Ohio State University, 154W 12[th] Avenue, Columbus, Ohio 43210, USA

16 [4]Crop Protection Branch, Department of Agriculture and Food, Western Australia, Bentley Delivery Centre, Perth,
17 6983, Western Australia, Australia.

18 [5] The University of Western Australia, ARC Centre of Excellence in Plant Energy Biology and School of Chemistry
19 and Biochemistry, Crawley, Perth 6009, Western Australia, Australia.
20
21 * Corresponding author: laura.boykin@uwa.edu.au
22
23

24 **Keywords:** cassava, Uganda, *Manihot esculenta,* smallholder farmer, *Cassava brown streak virus*,

25 species tree estimation, SVD Quartets, nonsynonymous mutations

26
27
28
29

Abstract

Cassava is a major staple food for about 800 million people in the tropics and sub-tropical regions

of the world. Production of cassava is significantly hampered by cassava brown streak disease

(CBSD), which is caused by *Cassava brown streak virus* (CBSV) and *Ugandan cassava brown streak*

virus (UCBSV). The disease is suppressing cassava yields in eastern Africa at an alarming rate.

Previous studies have documented that CBSV is more devastating than UCBSV because it more

readily infects both susceptible and tolerant cassava cultivars, resulting in greater yield losses.

Using whole genome sequences from NGS data, we produced the first coalescent-based species

tree estimate for CBSV and UCBSV. This species framework led to the finding that CBSV has a

faster rate of evolution when compared with UCBSV. Furthermore, we have discovered that in

CBSV, nonsynonymous substitutions are more predominant than synonymous substitution and

occur across the entire genome. All comparative analyses between CBSV and UCBSV presented

here suggest that CBSV may be outsmarting the cassava immune system, thus making it more

devastating and harder to control.

Introduction

Cassava (*Manihot esculenta* Crantz) is a major staple food crop for over 800 million people in over 100 tropical and sub-tropical countries[1]. In sub-Saharan Africa, it is the main source of dietary calories for approximately 300 million people[2]. The tuberous storage roots of cassava are rich in carbohydrates and can be cooked or processed for human food, animal feeds and a wide range of industrial products. The crop is relatively drought tolerant and can yield well even in less fertile soils, hence, its importance to poor families farming marginal lands[3]. Cultivation of cassava is most adversely affected by two viral diseases: cassava mosaic disease (CMD) and cassava brown streak disease (CBSD)[4], which together were reported to cause production losses of more than US$1 billion every year[5] in Africa.

Serious yield losses due to CMD were first observed on mainland East Africa in the 1920s[6]. Recorded epidemics of CMD later occurred in the 1930s, 1940s and from 1990s to date[7,8]. By contrast, for about 70 years since it was first described[9], CBSD was confined to low altitudes (below 1000 meters above sea level) along coastal eastern Africa in Kenya, Tanzania, Mozambique and Malawi. However, in the early 2000s, outbreaks of CBSD were reported over 1000 km inland at mid-altitude locations (above 1000m) in multiple countries all around Lake Victoria in Uganda[10], western Kenya[11] and northern Tanzania[4]. Where it is already established in eastern Africa, the current CBSD epidemic prevails as the main cause of losses in cassava production. Over the last 10 years, the CBSD epidemic has expanded to other countries in East and Central Africa such as Rwanda, Burundi, Congo, DR Congo and South Sudan[12-14]. This has significantly increased the risk to countries in central and west Africa which are among the world's leading cassava producers, and where CBSD does not occur.

CBSD is caused by *Cassava brown streak virus* (CBSV) and *Ugandan cassava brown streak virus*

(UCBSV). Both viruses are (+) ssRNA viruses in the genus *Ipomovirus* and family *Potyviridae*[15-18],

and are often together referred to as cassava brown streak viruses (CBSVs). The CBSVs have

genomic organization of 10 segments, total size approximately 8.9 to 10.8 kb, and coding for a

polypeptide with about 2,900 amino acid residues[15,17,18]. The complete genome of a CBSD causal

virus was first sequenced in 2009[18], and to date there are only 26 publicly available[19]. Currently

there are two species recognized by the ICTV, but Ndunguru et al.[19] have suggested further

speciation in the UCBSV clade. Both viruses are transmitted in a semi-persistent manner by the

whitefly *Bemisia tabaci*[20] and mechanically[21]. Symptoms of CBSD on cassava vary with cultivar,

virus or plant age, but typically include leaf veinal choloris, brown stem lesions, as well as

constrictions, fissures and necrosis of the tuberous storage roots[22,23]. Overall, both CBSV and

UCBSV cause similar symptom types, however, infection with CBSV tend to result in more severe

sumptoms.

Although CBSD has become established in eastern Africa, there is limited knowledge on the

diversity of causal viruses, their distribution and evolutionary potential. Therefore, it is necessary

to obtain several full genome sequences of CBSD viral isolates, better understand the causal

viruses and design long-term control approaches for the disease.

In contrast to the growing knowledge on the causal agents of CBSD, host-pathogen interactions

are less clear. As such, little is known about specific responses of different cassava varieties to

prevailing species or strains of CBSD viral pathogens. Development and dissemination of CBSD-

tolerant varieties has been the main means adopted for CBSD control in eastern Africa. With

significant efforts geared at breeding for CBSD-resistant varieties, it is of great interest to know

if such resistance protects cassava against one or both CBSVs. The resistance may be expressed

as several related features including restricted infection, systemic spread or recovery of infected

plants from disease and the possibility that stem cuttings taken from these may give rise to

progeny that are virus-free (reversion). Recent studies have shown CBSV to be the more

aggressive virus, infecting both tolerant and susceptible cultivars as single or mixed infections

with UCBSV[15,24,25]. In contrast, tolerant varieties were infected with only CBSV, but free of UCBSV,

suggesting their resistance to the latter. Compared with UCBSV, CBSV isolates have been

reported to be more detectable, having higher infection rates by graft inoculation and inducing

more severe symptoms[26]. It has also been shown that plants of CBSD tolerant or resistant

cultivars graft-inoculated with UCBSV developed milder symptoms and a significantly higher

proportion of the progenies were virus-free (reverted) compared to those infected with CBSV[27].

To date, the underlying reasons for this more aggressive nature of CBSV compared with UCBSV

are not known.

In this study, CBSV and UCBSV molecular diversity was investigated by using next generation

sequencing to understand new complete genomes of three isolates from Uganda. The sequences

obtained were analyzed to determine species composition, CBSV and UCBSV evolutionary rates,

potential role of such changes in virus-host interactions, resulting into cassava cultivar

susceptibility or resistance. We set out to answer the following questions:

1) How do the three new complete genomes from Uganda compare to those already

published[19]?

2) Are CBSV and UCBSV distinct species and is there further speciation?

3) Why is CBSV more aggressive and harder to breed resistance for than UCBSV?

Results

CBSD Field Symptoms Associated with CBSV and UCBSV Isolates

Categorisation of CBSD foliar symptom distribution on symptomatic plants assessed revealed

that the most frequently encountered type was LL - symptoms only on lower leaves (68.4%),

followed by SW - systemic and on the whole plant (26.3%), and SL – systemic but localized (5.3%)

(table 1). Based on CBSVs detected and CBSD leaf symptom severity scores for 57 sampled plants,

whereas the majority of plants infected by UCBSV alone as determined by RT-PCR had mild

chlorosis (severity score 2), CBSV infections (single or mixture with UCBSV) tended to have

moderate to severe symptoms (scores 3-4) in same proportion to those exhibiting score 2 (Fig. 1,

Table 1). Regarding the three isolates used here for whole genome sequencing, U8 (UCBSV) was

from a plant with CBSD score 3 and LL symptom type. Both CBSV isolates (U1 and U4) were from

plants with severity scores 2 and 3, symptom types LL and SL, respectively.

Next Generation Sequencing

The three samples from Uganda produced raw reads ranging from 21,844,716 to 23,648,990.

After trimming for quality using CLCGW, these numbers were reduced to 21,582,374 to

23,373,606 (table 2). Following *de novo* assembly of the trimmed reads using CLCGW, the

numbers of contigs produced were 621-1,008. The contigs of interest from *de novo* assembly

were of lengths 2,214 to 8,954nt, with average coverage 24 to 366. After mapping to a reference

genome in Geneious, the lengths of the consensus sequences were 8,893 to 9,563 with average

coverages of 25 to 393. The final sequences consisted of a consensus between the *de novo* and

the mapped consensus with lengths of 8,700 to 8,748.

Genomic Variability and Positive Selection

The CBSV genomes included in this study were more variable when compared with those of

UCBSV (supplementary figs. S1 and S2). Characterizing amino acid usage at each position in the

whole genome revealed that CBSV genomes have non-synonymous substitutions present across

their entire genome (Fig. 2), and predominating when compared to synonymous substitutions.

In contrast, UCBSV had near equal non-synonymous and synonymous substitution rates across

the entire genome. Genes in the UCBSV genomes with non-synonymous substitutions at a higher

frequency were: P1, NIb and HAM1 (Fig. 2).

CBSV had 68 positively selected sites and 66 negatively selected sites, while UCBSV had zero

positively selected sites and 558 negatively selected sites (Table 3). Analyzed together there are

3 positively selected sites and 1383 negatively selected sites. The coat protein (CP) of CBSV had

the highest number of positively selected sites (16) while 6K2 had zero.

Rates of Evolution

CBSV and UCBSV have different rates of evolution (Table 4). We tested two hypothesis using

CODEML. The null hypothesis tested was that CBSV and UCBSV have equal rates of evolution

while the alternative hypothesis was that CBSV and UCBSV have different rates of evolution (two

omegas; model = 2). The Likelihood Ratio Test was used to test for significance. If the difference

in likelihood was greater than 3.84 (based on the Chi-squared distribution and one degree of

freedom) we rejected the null hypothesis that the rates between CBSV and UCBSV are equal.

CBSV whole genome sequences showed it is evolving 5 times faster than UCBSV overall. The

genes contributing to this accelerated rate of evolution for CBSV are Nla (D=29.95), followed by

6K2 (D=6.74), Nlb (D=5.18) and P1 (4.61) (Table 4). The transition/transversion ratios were also

estimated using CODEML, and show that the 6K1 (19.6) and CP (13.2) genes have the highest

estimates while the remaining 8 genes ranged from 5.05 – 9.93.

Species Tree Estimation - SVDQ

The species phylogeny (Fig. 3) shows strong support for a split into two primary viral clades, one

consisting of CBSV (Fig. 3 clades A and B) and the other consisting of UCBSV (Fig. 3 clades E-G),

with 100% bootstrap support separating the two clades. Figure 3 shows clades labeled A–G which

correspond to; 1) labels A-F from Ndunguru et al [19], and 2) a new clade G defined in this study.

Within the CBSV clade, there are several additional clades with 100% bootstrap support,

including the two new CBSV whole genomes from Uganda (U1 and U4). These are the first CBSV

whole genomes sequences from Uganda. The other CBSV grouping with 100% bootstrap support

labeled B in Figure 3 contains 4 Tanzania samples KoR6, Tan 79, Tan 19 1 and Nal 07. In the

UCBSV clade there are 6 nodes supported with a 100% bootstrap, including the new UCBSV whole

genome added from this study (U8) which is sister to Kab 07 from Uganda. In addition, the CBSV

clade had all samples from a given country grouping together while the UCBSV clade had

monophyletic clades from different countries (the multi-colored lines in Fig. 3).

Comparison of Gene Trees to Species Tree

Clades A and B, which partition the CBSV isolates into two groups, are consistently present with

high support in all genes except HAM1 and CP (Table 5). Clades D and G, which each consist of a

pair of UCBSV isolates, have high support across all genes, while clades C and E have relatively

high support across a majority of genes. Clade F is strongly supported by the CI gene, which is

relatively long, but is not found in the phylogenetic tree estimated for any of the other genes.

The whole genome concatenated analysis using MrBayes shows strong support (posterior

probability 1.0) for all clades (Table 5). However, this analysis does not take into account the

possibility of variation in the evolutionary processes across the individual genes. The SVDQ

analysis, on the other hand, uses a coalescent-based method to estimate the overall species tree,

and properly accounts for variation in the evolutionary history for each gene. In viewing the

bootstrap support values for each of the clades from the SVDQ analysis, we see that the level of

support for each clade across the genome is more accurately represented by the corresponding

bootstrap proportion. For example, clade F, which was found only in the phylogeny of the CI

gene, shows a bootstrap proportion of 0.44 for the SVDQ analysis (as compared to 1.0 for the

MrBayes concatenated analysis) (Table 5). Similarly, the SVDQ analysis gives a bootstrap

proportion of 0.87 for clade E, which showed posterior probabilities below 0.8 for 3 of the 10

genes, as compared to a posterior probability of 1.0 for the concatenated analysis with MrBayes.

All other clades are supported with bootstrap values of 1.0, consistent with the MrBayes analysis.

Sliding Window SVD Score

The SVD Score Sliding Window analysis (Fig. 4) shows several interesting patterns. First, note that

the gene boundaries track well with shifts in the magnitude of the SVD Score, indicating that

individual genes are subject to specific evolutionary processes that vary from gene to gene. In

particular, several genes show strong support for the primary CBSV/UCBSV split, as indicated by

their low scores, while other genes show variation from this basic process, as indicated by

increases in the scores. In addition, Fig. 4 shows the test statistic associated with the hypothesis

test for a shift in the rate of evolution between the two groups, with `*' indicating that the rate

difference between the two groups is statistically significant. It is readily apparent from the graph

that genes that show strong support (low SVD Score) for the primary CBSV/UCBSV split also show

strong evidence for statistically significant differences in evolutionary rate. These results support

the overall hypothesis that certain genes in CBSV have accelerated rates of evolution that may

be contributing to the higher aggressiveness of the virus.

220
221 **Discussion**
222

In this study, we analyzed new and all publicly available CBSVs whole genomes to elucidate

molecular mechanisms underlying the field and laboratory observations that CBSV more readily

infects cassava plants and tends to display severe symptoms when compared with UCBSV. Our

analyses included characterizing three new complete CBSV (2) and UCBSV (1) genomes, which

were combined with the 26 previously published. Our major findings show further speciation of

CBSV and UCBSV, a larger genetic landscape for CBSV, including many nonsynonymous sites, and

reveal that CBSV has a faster rate of evolution compared with UCBSV (Table 4 and Fig. 4). These

observations and their biological significance are discussed.

<u>**Genes with Accelerated Rates of Evolution in CBSV**</u>

We have identified P1, 6K2, NIb and NIa as the genes with accelerated rates of evolution in CBSV.

The function of P1 is as an RNA silencing suppressor (RSS), and there is also the suggestion that

it may be involved in virion binding to the whitefly stylet via a "bridge" formation by a virus-

encoded P1 protein for both CBSV and UCBSV. 6K2 is associated with cellular membrane and

responsible for systemic infection and viral long distance movement[28]. The NIb encodes for a

nuclear inclusion polymerase and the NIA for a nuclear inclusion protease[18,29].

In Potyviruses generally, when NIa and VPg are associated together they are located in the

cytoplasm and nucleus of infected cells. When 6K2-VPg-NIa forms a larger product, the VPg plays

a role in viral RNA replication[30]. Even though VPg is not one of the genes with a higher evolution

rate, both 6K2 and NIa are a part of the complex which affects replication, and this may go some

way to explaining their apparent accelerated evolution rate. Is it possible that the accelerated

rates of evolution for genes involved in replication could even be a response to the relatively

recent interaction of the viruses and cassava? The CBSVs are not present in South America where

cassava originates, so the viruses must be native to Africa. It would appear that the adaptation is

still occurring and the cassava immune system does not know how to effectively fight these

infections. Cassava was introduced to East Africa in the 18[th] century through oceanic movement.

The first report of cassava brown streak disease was in 1936[9,13]. There has been little opportunity

for the co-evolution of the viruses and the host, therefore natural resistance would be a hard

prospect. This raises the possibility of the original host of these viruses, a non-cassava host which

may be harboring the CBSVs or the most recent common ancestor of these viruses. This in turn

leads us to wonder just how old these viruses and their ancestors are, if any of the CBSVs is the

ancestral species, and the best way to answer such questions is to sequence more virus genomes

from both cassava and non-cassava hosts wherever they are found.

How Can CBSV Still Function with Such a Large Genetic Landscape?

CBSV and UCBSV have different evolutionary patterns as observed by characterizing the whole

genome sequences of CBSV and UCBSV separately. CBSV is genetically more diverse when

compared with UCBSV, as evident by the greater amino acid usage (supplementary Fig. 1), the

faster rates of evolution across the entire genome (table 4), and greater number of

nonsynonymous sites across the entire genome (Fig. 2). How can CBSV still function with such a

large genetic landscape? RNA viruses walk a very fine line of having the genetic arsenal to

overcome the host immune system and diverging to a point that key functions of genes are lost

[31]. Recent studies[32,33] have shown that viruses with a large genetic landscape adapt to host

changes much quicker and can overcome the host immune system faster. Viruses that occupy a

large portion of the possible sequence space might be less fit but they outcompete the fitter

strain when the host immune system shifts and hence these viruses have been described as

adapted to "survival of the flattest"[34,35]. This means that a virus that covers the most sequence

space will be able to adapt to host immune system faster than those with smaller spaces. Viruses

that are adapted in this category ("survival of the flattest") are going to be harder to breed

resistance for because the virus has a larger ability to adapt to changes. It is clear that in our

case, CBSV is the virus that has a larger sequence space (Supplemental Fig. 1) when compared to

that of UCBSV, which is clearly smaller (Supplemental Fig. 2). CBSV is one of the RNA viruses that

can be described as adapted to "survival of the flattest", while UCBSV is not. Therefore, CBSV is

more devastating because it has a larger genetic arsenal which it uses overcome the changes

breeders are introducing into cassava.

Not only are the CBSV genomes more genetically diverse, but are also characterized by a large

number of nonsynonymous changes in the genome (Fig. 2). An excess of nonsynonymous over

synonymous substitutions at individual amino acid sites signifies that positive selection has

affected the evolution of a protein between the extant sequences under study and their most

recent common ancestor[36]. Positive selection is the process by which new advantageous genetic

variants sweep a population and is the mechanism Darwin described to drive evolution. This is

further evidence that might suggest that CBSV has a greater capacity to evade the cassava

immune system as compared with UCBSV. CBSV had 66 sites under positive selection (Table 4)

while UCBSV had none. The CBSV sites under positive selection are found not only in the regions

that have gained the most attention, CP and HAM1-like[13], but are also found in all other genes

except 6K2. This is further support for CBSV's potential ability to outsmart the cassava immune

system. Every gene in the CBSV genome (except 6K2) has sites under positive selection indicating

effective RNA silencing of the virus will need to encompass many loci.

Using computational methods combined with field observations we have concluded that CBSV is

more devastating than UCBSV. This assertion is also supported by two recent biological studies.

The first was a test of reversion in three different cassava varieties (Albert, Kaleso and Kiroba)

infected with CBSV and UCBSV. Reversion is a type of resistance mechanism whereby virus-

infected plants will naturally recover from infection over time, and a proportion of their progeny

from stem cuttings are virus-free. A reversion event infers the host immune system was able to

clear or restrict the virus from systemic movement. It was shown that UCBSV-infected cassava

had a higher rate of reversion when compared to plants infected with CBSV[27] indicating that

plants infected with UCBSV recovered more often than those infected with CBSV. This is another

line of evidence supporting the more devastating nature of CBSV, and the possibility that cassava

immune systems of the three varieties tested are struggling to resist the virus.

The second study supporting the hypothesis that CBSV is more aggressive than UCBSV analyzed

virus-derived small RNAs within three cassava varieties (NASE 3, TME204 and 60444). Plants

infected with viruses are known to trigger RNAi antiviral defense that can be measured by

quantifying the abundance of 21-24 nucleotide (nt) segments produced by the dicer enzyme[37].

Cassava varieties were infected with either CBSV or UCBSV, NGS was used to detect virus-derived

small RNAs[24], and the 21-24 nt dicer fragments were mapped to either CBSV or UCBSV depending

on which virus was used to infect the plant. The results showed that CBSV infection triggered a

stronger immune response as measured by greater abundance of virus derived small RNA

fragments across the entire CBSV genome compared with UCBSV. In addition, across all three

genotypes they observed that cassava grafted with CBSV-infected buds showed more severe

symptoms compared to UCBSV-infected plants[24]. This is further evidence that CBSV is a more

aggressive virus and breeding for resistance to CBSV and UCBSV will require different

experimental approaches.

Implications of the Species Tree for CBSV and UCBSV

We have produced the first species tree estimate of the CBSD causal virus species using whole

genome sequences and the coalescent-based SVD Quartets species tree estimation algorithm.

Differences in the evolutionary history of the two viruses are seen in the branching patterns in

Figure 3. CBSV has diverged into two main clades A and B, while UCBSV has several well-

supported clades but the backbone is still unresolved, indicating more sampling is needed to fully

understand the diversity and evolutionary history of UCBSV. The species tree (Fig. 3) is similar to

the concatenated whole gene tree reported in Ndunguru et al.[19], except addition of the clade

labeled "G", and lack of support for clades E and F in the UCBSV species. It is well-documented

that concatenating genes without using the coalescent-based models can produce misleading

results [38,39]. In our case, only CI supports clade F, and it is also the longest gene (1,883 bp), and

therefore may swamp the signal of the other genes. The whole genome concatenation analysis

recovers clade F with a posterior probability of 1.00 (Table 4). With regard to clade E, the SVDQ

tree was more reflective of the individual gene tree signal by producing a bootstrap value of 0.87

versus 1.00 for the whole genome concatenated tree (Table 4). These results suggest that the

estimated topology in the UCBSV species may be further refined as more samples are added.

Our integrative approach of species tree estimation coupled with analyzing rates of evolution has

lead to a new framework for CBSV and UCBSV, which includes analyzing and treating these two

groups of viruses as separate species. Multiple putative species of both CBSV and UCBV have

been identified which means cassava needs to be resistant to the virus species that are prevalent

in farmers' fields. We argue that this genomic diversity and faster rate of evolution for CBSV is

what is causing the breeders to struggle with breeding resistant varieties and also why the

diagnostic primers are not working consistently. CBSV also has more positively selected sites than

UCBSV. It was first thought that CBSD was restricted to the coastal areas and below 1000 m [23]

but as more genetic data is gathered CBSV and UCBSV are found at all elevations in many

ecozones throughout eastern Africa[4,10,13,15,19,40]. We are still in the discovery phase with CBSV and

UCBSV species as there are only 29 (now with the three new included here) whole genome

sequences and other new species of both viruses are likely to be discovered. As we move forward

it is important to include all known samples and use appropriate species tree estimation methods

such as SVDQ.

Finally, the traditional gene regions (CP and HAM1-like) that are used to delimit species and serve

as the targets for diagnostic primers do not recover the species tree (Table 4). We recommend

designing new diagnostic regions for other genes that recover the species tree and also do not

have an accelerated rate of molecular evolution (Figure 4), such as CI or P3 for species-level

diagnoses. It is possible that the spread of CBSV and UCBSV could have been exacerbated

through dissemination of infected cuttings, as virus indexing with primers targeting CP may have

misleadingly returned negative results.

Implications of the Results for Cassava Breeding

During the last three decades worldwide, agricultural production has been compromised by a

series of epidemics caused by new variants of classic viruses that show new pathogenic and

epidemiological properties. An important determinant of the fitness of a virus in a given host is

its ability to overcome the defenses of the host. Overcoming plant resistance by changes in the

pathogenicity of viral populations represents a specific and important case of emergence, with

tremendous economic consequences since it jeopardizes the success and durability of resistance

factors in crops as an anti-viral control strategy. In this study, we found CBSV to be more variable,

to have more positively selected sites, and to evolve five times faster than UCBSV. These findings

have huge implications for cassava improvement efforts in Africa where CBSV is widely present.

Field and laboratory results have proven CBSV to be more virulent and more devastating than

UCBSV. Knowledge of specific virus species an improved cassava variety is resistant to will

determine where to screen, multiply and deploy such varieties. Cassava breeders have to take

into consideration the evolutionary and biological differences between CBSV and UCBSV in the

breeding programs. For example, cassava breeders can breed varieties that are resistant to CBSV

that can be strategically deployed in areas where CBSV is more prevalent, and similarly for

UCBSV. Furthermore, it becomes more appropriate to always screen cassava materials against

CBSV as a minimum, even if UCBSV is the more prevalent virus. Such a strategy will in effect

ensure durable resistance as opposed to the indiscriminate screening and distribution of the

improved CBSD resistant cassava varieties, without knowledge of the virus species in the area.

380
Methods

Field Plant Sample Collection

Farmers' fields in Uganda with cassava plants 3-6 months old were surveyed for CBSD in 20

districts. In each field, cassava plants were visually assessed to confirm typical CBSD symptoms

on leaves and stems. CBSD leaf symptom severity was scored on a 1-5 scale[41,42]; 1 = no visible

symptoms, 2 = mild vein yellowing or chlorotic blotches on some leaves, 3 =

pronounced/extensive vein yellowing or chlorotic blotches on leaves, but no lesions or streaks

on stems, 4 = pronounced/extensive vein yellowing or chlorotic blotches on leaves and mild

lesions or streaks on stems, 5 = pronounced/extensive vein yellowing or chlorotic blotches on

leaves and severe lesions or streaks on stems, defoliation and dieback. CBSD symptoms were also

categorized based on distribution of leaf chlorosis and stem lesions on the plant; systemic and

on the whole plant (SW), systemic on leaf or stem parts but localized (SL), only on lower leaves

(LL). On selected symptomatic plants, portions of the third fully expanded leaf on a shoot were

picked as samples, air-dried by pressing between sheets of newsprint and stored pending RNA

extraction.

RNA Extraction

About 0.25 g cassava leaf samples were frozen in liquid nitrogen, then ground using a mortar and

pestle. 2 ml CTAB lysis buffer (2% CTAB; 100 mM Tris–HCl, pH 8.0; 20 mM EDTA; 1.4 M 134 NaCl;

1% sodium sulphite; 2% PVP) was added and samples homogenized. The 1 ml of the homogenate

was incubated at 65°C for 15 min, an equal volume of chloroform: isoamyl alcohol (24:1) was

added, and the sample was centrifuged for 10 min at approximately 14,500rpm. 800µl of the

aqueous layer was transferred to a new tube with an equal volume of 4 M LiCl and incubated at

-20°C for 2 hrs. The samples were centrifuged for 25 min at 14,500 rpm and the supernatant was

poured off. The pelleted RNA was re-suspended in 200 µl TE buffer containing 1% SDS, 100 µl of

5M NaCl. 300 µl of ice-cold isopropanol were added and incubated at -20°C for 30 min. The

sample was centrifuged at 13,000 rpm for 10 min and the aqueous layer was decanted and RNA

pellets washed in 500 µl of 70% ethanol by centrifuging at 13,000 rpm for 5 min. The ethanol was

decanted off and RNA pellet dried to remove residual ethanol. The RNA was re-suspended in 50

µl nuclease-free water and stored at −80°C prior to testing.

412   CBSV and UCBSV Detection by RT-PCR

All samples were tested for presence of CBSV and UCBSV by a two-step RT-PCR assay [43]. The PCR

mixture consisted of 16.0 µl nuclease free water, 2.5 µl PCR buffer, 2.5 µl $MgCl_2$ (2.5 mM), 0.5µl

dNTPs (10 mM), 1.0 µl of each primer (10mM) [forward CBSDDF2 5'-

GCTMGAAATGCYGGRTAYACAA-3' and reverse CBSDDR 5'-GGATATGGAGAAAGRKCTCC-3'], 0.5 µl

Taq DNA polymerase and 1.0 µl of cDNA. The PCR thermo profile consisted of: $94^{o}C$ for 2 min

followed by 35cycles of $94^{o}C$ (30 s), $51^{o}C$ (30 s) and $72^{o}C$ (30 s) for denaturation, annealing and

extension, respectively. PCR products were analysed by electrophoresis in a x1 TAE buffer on a

1.2% agarose gel, stained with ethidium bromide, visualized under UV light and photographed

using a digital camera.

423   Sample Selection for Sequencing

From the data obtained in the diagnostic tests, samples for sequencing were selected to

represent different geographical regions, symptom types and severities. Three samples that

tested positive for either CBSV (2) or UCBSV (1) were selected for this study. The two samples for

which presence of CBSV was confirmed (U1 and U4) had been collected from different farmers'

fields in Mukono district, central Uganda. The sample with UCBSV (U8) selected for further

analysis originated from a field in Mayuge district, eastern Uganda.

431   Generation of the Transcriptomes

The three samples were transported to the laboratory and extracted as detailed above. Total

RNA was blotted on to FTA cards and later extracted using methods previously described[44]. Total

RNA from each sample was sent to the Australian Genome Research Facility (AGRF) for library

preparation and barcoding before 100 bp paired-end sequencing on an Illumina HiSeq2000.

De novo Sequence Assembly and Mapping

For each sample, reads were first trimmed using CLC Genomics Workbench 6.5 (CLCGW) with the

quality scores limit set to 0.01, maximum number of ambiguities to two and removing any reads

with <30 nucleotides (nt). Contigs were assembled using the *de novo* assembly function of

CLCGW with automatic word size, automatic bubble size, minimum contig length 500, mismatch

cost two, insertion cost three, deletion cost three, length fraction 0.5 and similarity fraction 0.9.

Contigs were sorted by length and the longest subjected to a BLAST search (blastn and blastx)[45].

In addition, reads were also imported into Geneious 6.1.6[46] and provided with reference

sequences obtained from Genbank (KR108828 for CBSV and KR108836 for UCBSV). Mapping was

performed with minimum overlap 10%, minimum overlap identity 80%, allow gaps 10% and fine

tuning set to iterate up to 10 times. A consensus between the contig of interest from CLCGW and

the consensus from mapping in Geneious was created in Geneious by alignment with MAFFT[47].

Open reading frames (ORFs) were predicted and annotations made using Geneious. Finalized

sequences were designated as "complete" based on comparison with the reference sequences

used in the mapping process, and "coding complete" if some of the 5' or 3' UTR was missing but

the coding region was intact[48,49], and entered into the European Nucleotide Archive (WEBIN ID

number Hx2000053576).

Genome Alignment and Annotation

Twenty-six whole genomes (12 CBSV and 14 UCBSV) were downloaded from GenBank and

imported into Geneious[46], and the MAFFT plugin[47] was used to align them with the 3 new whole

genome sequences obtained in this study. Nucleotide alignments were translated into protein

using the translate align option in Geneious and then visually inspected for quality. Annotations

were transferred to the 3 new genomes from the 26 previously published genomes using the live

annotation option in Geneious.

Characterizing the Genetic Diversity in CBSV and UCBSV Genomes

CBSV and UCBSV are distinct species (Fig. 2) therefore the genomes were treated separately in

the analyses in characterizing the genomes. Characterizing the genetic diversity of CBSV and

UCBSV was done using the Synonymous Non-synonymous Analysis Program (SNAP v2.1.1)

implemented in the Los Alamos National Laboratory HIV-sequence database

(http://www.hiv.lanl.gov)[50]. SNAP calculates synonymous and non-synonymous substitution

rates based on a set of codon-aligned nucleotide sequences. This program is based on the simple

methods for estimating the numbers of synonymous and nonsynonymous nucleotide

substitutions of[51], and incorporating a statistic developed for computing variances and

covariances of dS's and dN's[52]. An application of the SNAP package in HIV-1 research has also

been developed[53].

Estimating Rates of Evolution

To further characterize the CBSV and UCBSV genomes, we estimated the rates of molecular

evolution using CODEML implemented in PAML (Phylogenetic Analysis by Maximum

Likelihood)[54]. PAML is a package of programs for analysis of DNA or protein sequences by using

maximum likelihood methods in a phylogenetic framework. The null hypothesis tested was that

CBSV and UCBSV have equal rates of evolution (one omega; model = 0) while the alternative

hypothesis was that CBSV and UCBSV have different rates of evolution (two omegas; model = 2).

The Likelihood Ratio Test was used to test for significance. If the test statistic was greater than

3.84 (based on the Chi-squared distribution and one degree of freedom) we then rejected the

null hypothesis that the rates between CBSV and UCBSV are equal. Initial analyses were carried

out for the entire genome and showed that CBSV has a higher rate of evolution (Table 4). To

identify which gene or genes were contributing to the faster rate of evolution we repeated the

analysis separately for each individual gene.

Testing for Positive Selection

Sites under positive selection were identified using SLAC[55] implemented on the

http://www.datamonkey.org web server[56]. The settings used to run SLAC were as follows: the

best fitting model (GTR) was specified global dN/dS value was estimated and the significance

level was set to 0.01.

Gene Tree Estimation

Individual gene trees were estimated using MrBayes 3.2.1[57] run in parallel on Magnus (Pawsey

Supercomputing Centre, Perth, Western Australia) utilizing the BEAGLE library [58]. MrBayes 3.2.1

was run utilizing 4 chains for 30 million generations and trees were sampled every 1000

generations. All runs reached a plateau in likelihood score, which was indicated by the standard

deviation of split frequencies (0.0015), and the potential scale reduction factor (PSRF) was close

to one, indicating that the MCMC had converged.

Species Tree Estimation

The SVDQ method[59] implemented in PAUP*[60] was used to analyze the whole-genome data. This

method enables analysis of multi-locus data in a coalescent framework that allows for variation

in the phylogenetic histories of individual genes. The method was run with all possible quartets

(23,751) sampled in each of 100 bootstrap replicates, and the consensus across all bootstrap

replicates was used as the estimate of the species tree.   Bootstrap support values for each node

were used to quantify uncertainty in the species tree estimate. The entire analysis took

approximately 2.5 minutes on a MacBook Pro running OSX 10.11.2 with a 2.2 GHz Intel Core i7

processor.

Comparison of Gene Trees to Species Tree

We compared the single-gene phylogenies constructed using MrBayes with the overall species

tree phylogeny estimated using SVDQ and the concatenated phylogeny estimated by MrBayes.

For each tree, we evaluated presence or absence of the clades identified by Ndunguru et al.[19]

labeled A-F in Figure 3. We identified an additional clade (clade G, Fig. 3) that we noticed to be

consistently present across genes and methods. For each of these clades present in a particular

tree, we recorded the posterior probability (for trees constructed by MrBayes) or the bootstrap

proportion (for the tree estimated by SVDQ) in Table 5.

Sliding Window SVD Score

The SVD Score[61] was used to quantify support for two viral clades for portions of the genome in

a sliding window analysis. Briefly, the SVD Score measures the extent to which the data support

a phylogenetic "split" – a division of the taxa into two groups with specified group membership.

Low values of the SVD Score indicate strong support for the split of interest, while larger values

indicate *either* a lack of support for the split or a shift in the underlying evolutionary process (see

Allman et al. (2016) for details and examples). We computed the SVD Score with the split defined

by CBSV vs. UCBSV across the genome in windows of 500 bp, sliding in increments of 100 bp, and

plotted the resulting SVD Scores across the genome, with boundaries between genes marked

with vertical lines. The computations took less than one minute on a MacBook Pro running OSX

10.11.2 with a 2.2 GHz Intel Core i7 processor.

Acknowledgments

This work was supported by the Bill and Melinda Gates Foundation Grant no. 51466 "Regional

Cassava Virus Diseases Diagnostic Project" awarded to Mikocheni Agricultural Research Institute,

Tanzania, and a sub-grant to National Agricultural Research Organisation (Uganda).

Computational resources provided by the Pawsey Supercomputing Centre with funding from the

Australian Government and the Government of Western Australia supported this work.

Author contributions

Sample collection was carried out by T.A., G.O., R.N., L.Kiiza. Laboratory work was carried out

by G.O, R.N., L.Kiiza. J.N. P.S. M.K. Computational analyses were conducted by T.A., L.B.,

L.Kubatko., M.K, and all authors contributed to the writing of the manuscript.

Conflict of Interest

The authors declare that they have no conflicts of interest with the contents of this article.

References
551

552   1   Thresh, J. M. Control of tropical plant virus diseases. *Adv Virus Res* **67**, 245-295,
553   doi:10.1016/S0065-3527(06)67007-3 (2006).

554   2   Nweke, F. I. A Cash Crop in Africa. COSCA Working Paper No. 14. Collaborative Study of
555   Cassava in Africa, International Institute of Tropical Agriculture, Ibadan, Nigeria. (1996).

556   3   Robertson, A. I. & Ruhode, T. The potential advantages of cassava over hybrid maize as
557   a food security crop and a cash crop in the Southern Africa semi-arid zone. *African Crop*
558   *Science Crop Science Conference Proceedings* **5**, 539-542 (2001).

559   4   Legg, J. P. *et al.* Comparing the regional epidemiology of the cassava mosaic and cassava
560   brown streak virus pandemics in Africa. *Virus Res* **159**, 161-170,
561   doi:10.1016/j.virusres.2011.04.018 (2011).

562   5   Legg, J. P., Owor, B., Sseruwagi, P. & Ndunguru, J. Cassava mosaic virus disease in East
563   and Central Africa: epidemiology and management of a regional pandemic. *Adv Virus*
564   *Res* **67**, 355-418, doi:10.1016/S0065-3527(06)67010-3 (2006).

565   6   Hall, F. W. Annual Report of the Department of Agriculture, Uganda. *Government*
566   *Printer, Entebbe*, 35 (1928).

567   7   Jameson, J. D. Cassava mosaic disease in Uganda. *East African Agricultural and Forestry*
568   *Journal* **29**, 208-213 (1964).

569   8   Otim-Nape, G. W. *et al.* The current pandemic of cassava mosaic virus disease in East
570   Africa and its control. *NARO/NRI/DFID Publication. Chatham, UK. 100pp* (2000).

571   9   Storey, H. H. Virus diseases of East African plants: VI, A progress report of studies of the
572   diseases of cassava. *East African Agricultural Journal* **2**, 34-39 (1936).

573   10   Alicai, T. *et al.* Re-emergence of cassava brown streak disease in Uganda. *Plant Disease*
574   **91**, 24-29 (2007).

575   11   Ntawuruhunga, P. & Legg, J. P. New Spread of Cassava Brown Streak Virus Disease and
576   Its Implications for The Movement of Cassava Germplasm in The East and Central African
577   Region. International Institute of Tropical Agriculture-Uganda & Eastern Africa Root
578   Crops Research Network Report. (2007).

579   12   Bigirimana, S., Barumbanze, P., Ndayihanzamaso, P., Shirima, R. & Legg, J. P. First report
580   of cassava brown streak disease and associated Ugandan cassava brown streak virus in
581   Burundi. *New Disease Reports* **24**, 26 (2011).

582   13   Mbanzibwa, D. R. *et al.* Evolution of cassava brown streak disease-associated viruses. *J*
583   *Gen Virol* **92**, 974-987, doi:10.1099/vir.0.026922-0 (2011).

584   14   Mulimbi, W. *et al.* First report of Ugandan cassava brown streak virus on cassava in
585   Democratic Republic of Congo. *New Disease Reports* **26** (2012).

586   15   Winter, S. *et al.* Analysis of cassava brown streak viruses reveals the presence of distinct
587   virus species causing cassava brown streak disease in East Africa. *J Gen Virol* **91**, 1365-
588   1372, doi:10.1099/vir.0.014688-0 (2010).

589   16   Monger, W. A., Seal, S., Isaac, A. M. & Foster, G. D. Molecular characterization of cassava
590   brown streak virus coat protein. *Plant Pathol* **50**, 527-534 (2001).

591   17   Monger, W. A. *et al.* The complete genome sequence of the Tanzanian strain of Cassava
592   brown streak virus and comparison with the Ugandan strain sequence. *Arch Virol* **155**,
593   429-433, doi:10.1007/s00705-009-0581-8 (2010).

594   18   Mbanzibwa, D. R. *et al.* Genetically distinct strains of Cassava brown streak virus in the
595   Lake Victoria basin and the Indian Ocean coastal area of East Africa. *Arch Virol* **154**, 353-
596   359, doi:10.1007/s00705-008-0301-9 (2009).

597   19   Ndunguru, J. *et al.* Analyses of Twelve New Whole Genome Sequences of Cassava Brown
598   Streak Viruses and Ugandan Cassava Brown Streak Viruses from East Africa: Diversity,
599   Supercomputing and Evidence for Further Speciation. *PLoS One* **10**, e0139321,
600   doi:10.1371/journal.pone.0139321 (2015).

601   20   Maruthi, M. N. *et al.* Transmission of Cassava brown streak virus by Bemisia tabaci
602   (Gennadius). *J Phytopathol* **153**, 307-312 (2005).

603   21   Lister, R. M. Mechanical transmission of cassava brown streak virus. *Nature* **183**, 1588-
604   1589 (1959).

605   22   Jennings, D. L. Observations on virus diseases of cassava in resistant and susceptible
606   varieties. II. Brown streak disease. *Empire Journal of Experimental Agriculture* **28**, 261-
607   269 (1960).

608   23   Nichols, R. F. W. The brown streak disease of cassava: Distribution, climatic effects and
609   diagnostic symptoms. *East African Agricultural Journal* **15**, 154-160 (1950).

24   Ogwok, E., Ilyas, M., Alicai, T., Rey, M. E. & Taylor, N. J. Comparative analysis of virus-
derived small RNAs within cassava (Manihot esculenta Crantz) infected with cassava
brown streak viruses. *Virus Res* **215**, 1-11, doi:10.1016/j.virusres.2016.01.015 (2016).

25   Patil, B. L. *et al.* RNAi-mediated resistance to diverse isolates belonging to two virus
species involved in Cassava brown streak disease. *Mol Plant Pathol* **12**, 31-41,
doi:10.1111/j.1364-3703.2010.00650.x (2011).

26   Mohammed, I. U., Abarshi, M. M., Muli, B., Hillocks, R. J. & Maruthi, M. N. The symptom
and genetic diversity of cassava brown streak viruses infecting cassava in East Africa.
Adv Virol **2012**, 795697, doi:10.1155/2012/795697 (2012).

27   Mohammad, I. U., Ghosh, S. & Maruthi, M. N. Host and virus effects on reversion in
cassava affected by cassava brown streak disease. *Plant Pathology* **65**, 593-600 (2016).

28   Jiang, J., Patarroyo, C., Garcia Cabanillas, D., Zheng, H. & Laliberte, J. F. The Vesicle-
Forming 6K2 Protein of Turnip Mosaic Virus Interacts with the COPII Coatomer Sec24a
for Viral Systemic Infection. *J Virol* **89**, 6695-6710, doi:10.1128/JVI.00503-15 (2015).

29   Dombrovsky, A., Reingold, V. & Antignus, Y. Ipomovirus--an atypical genus in the family
Potyviridae transmitted by whiteflies. *Pest Manag Sci* **70**, 1553-1567,
doi:10.1002/ps.3735 (2014).

30   Revers, F. & Garcia, J. A. Molecular biology of potyviruses. *Adv Virus Res* **92**, 101-199,
doi:10.1016/bs.aivir.2014.11.006 (2015).

31   Sanjuan, R., Moya, A. & Elena, S. F. The distribution of fitness effects caused by single-
nucleotide substitutions in an RNA virus. *Proc Natl Acad Sci U S A* **101**, 8396-8401,
doi:10.1073/pnas.0400146101 (2004).

32   Lauring, A. S., Frydman, J. & Andino, R. The role of mutational robustness in RNA virus
evolution. *Nat Rev Microbiol* **11**, 327-336, doi:10.1038/nrmicro3003 (2013).

33   Acevedo, A., Brodsky, L. & Andino, R. Mutational and fitness landscapes of an RNA virus
revealed through population sequencing. *Nature* **505**, 686-690,
doi:10.1038/nature12861 (2014).

34   Wilke, C. O., Wang, J. L., Ofria, C., Lenski, R. E. & Adami, C. Evolution of digital organisms
at high mutation rates leads to survival of the flattest. *Nature* **412**, 331-333,
doi:10.1038/35085569 (2001).

35   Sanjuan, R., Cuevas, J. M., Furio, V., Holmes, E. C. & Moya, A. Selection for robustness in
mutagenized RNA viruses. *PLoS Genet* **3**, e93, doi:10.1371/journal.pgen.0030093 (2007).

36   Massingham, T. & Goldman, N. Detecting amino acid sites under positive selection and
purifying selection. *Genetics* **169**, 1753-1762, doi:10.1534/genetics.104.032144 (2005).

37   Blevins, T. *et al.* Four plant Dicers mediate viral small RNA biogenesis and DNA virus
induced silencing. *Nucleic Acids Res* **34**, 6233-6246, doi:10.1093/nar/gkl886 (2006).

38   Kubatko, L. S. & Degnan, J. H. Inconsistency of phylogenetic estimates from
concatenated data under coalescence. *Syst Biol* **56**, 17-24 (2007).

39   Degnan, J. H. & Salter, L. A. Gene tree distributions under the coalescent process.
Evolution **59**, 24-37 (2005).

40   Monger, W. A. *et al.* The complete genome sequence of the Tanzanian strain of Cassava
brown streak virus and comparison with the Ugandan strain sequence. *Arch Virol* **155**,
429-433 (2010).

41   Hahn, S. K., Isoba, J. C. G. & Ikotun, T. Resistance breeding in root and tuber crops at the
International Institute of Tropical Agriculture (IITA), Ibadan, Nigeria. *Crop Protection* **8**,
147-168 (1989).

42   Hillocks, R. J., Raya, M. D. & Thresh, J. M. The association between root necrosis and
above ground symptoms of brown streak virus infection of cassava in southern Tanzania.
International Journal of Pest Management **42**, 285-289 (1996).

43   Mbanzibwa, D. R. *et al.* Simultaneous virus-specific detection of the two cassava brown
streak-associated viruses by RT-PCR reveals wide distribution in East Africa, mixed
infections, and infections in Manihot glaziovii. *J Virol Methods* **171**, 394-400,
doi:10.1016/j.jviromet.2010.09.024 (2011).

663 44 Ndunguru, J., Legg, J. P., Aveling, T. A., Thompson, G. & Fauquet, C. M. Molecular
664   biodiversity of cassava begomoviruses in Tanzania: evolution of cassava geminiviruses
665   in Africa and evidence for East Africa being a center of diversity of cassava geminiviruses.
666   *Virol J* **2**, 21, doi:10.1186/1743-422X-2-21 (2005).

667 45 Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
668   search tool. *J Mol Biol* **215**, 403-410, doi:10.1016/S0022-2836(05)80360-2 (1990).

669 46 Geneious v5.1. Available from http://www.geneious.com/ (2010).

670 47 Katoh, K., Misawa, K., Kuma, K. & Miyata, T. MAFFT: a novel method for rapid multiple
671   sequence alignment based on fast Fourier transform. *Nucleic Acids Res* **30**, 3059-3066
672   (2002).

673 48 Kehoe, M. A., Coutts, B. A., Buirchell, B. J. & Jones, R. A. Split personality of a Potyvirus:
674   to specialize or not to specialize? *PLoS One* **9**, e105770,
675   doi:10.1371/journal.pone.0105770 (2014).

676 49 Kehoe, M. A., Coutts, B. A., Buirchell, B. J. & Jones, R. A. Plant virology and next
677   generation sequencing: experiences with a Potyvirus. *PLoS One* **9**, e104580,
678   doi:10.1371/journal.pone.0104580 (2014).

679 50 Korber, B. in *Computational Analysis of HIV Molecular Sequences* (eds A.G. Rodrigo &
680   G. H. Learn) Ch. 4, 55-72 (Kluwer Academic Publishers, 2000).

681 51 Nei, M. & Gojobori, T. Simple methods for estimating the numbers of synonymous and
682   nonsynonymous nucleotide substitutions. *Mol Biol Evol* **3**, 418-426 (1986).

683 52 Ota, T. & Nei, M. Variance and covariances of the numbers of synonymous and
684   nonsynonymous substitutions per site. *Mol Biol Evol* **11**, 613-619 (1994).

685 53 Ganeshan, S., Dickover, R. E., Korber, B. T., Bryson, Y. J. & Wolinsky, S. M. Human
686   immunodeficiency virus type 1 genetic evolution in children with different rates of
687   development of disease. *J Virol* **71**, 663-677 (1997).

54   Yang, Z. PAML: a program package for phylogenetic analysis by maximum likelihood. *Comput Appl Biosci* **13**, 555-556 (1997).

55   Kosakovsky Pond, S. L. & Frost, S. D. Not so different after all: a comparison of methods for detecting amino acid sites under selection. *Mol Biol Evol* **22**, 1208-1222, doi:10.1093/molbev/msi105 (2005).

56   Delport, W., Poon, A. F., Frost, S. D. & Kosakovsky Pond, S. L. Datamonkey 2010: a suite of phylogenetic analysis tools for evolutionary biology. *Bioinformatics* **26**, 2455-2457, doi:10.1093/bioinformatics/btq429 (2010).

57   Ronquist, F. *et al.* MrBayes 3.2: Efficient Bayesian Phylogenetic Inference and Model Choice Across a Large Model Space. *Systematic Biology* **61**, 539-542, doi:10.1093/sysbio/sys029 (2012).

58   Ayres, D. L. *et al.* BEAGLE: an application programming interface and high-performance computing library for statistical phylogenetics. *Systematic Biology* **61**, 170-173, doi:10.1093/sysbio/syr100 (2012).

59   Chifman, J. & Kubatko, L. Quartet inference from SNP data under the coalescent model. *Bioinformatics* **30**, 3317-3324, doi:10.1093/bioinformatics/btu530 (2014).

60   Phylogenetic Analysis using Parsimony (* and other methods) v. (open source version 4.0a147 downloaded from http://people.sc.fsu.edu/~dswofford/paup_test/ on March 25, 2016). (Sinauer Associates, Sunderland, MA 2002).

61   Allman, E. S., Kubatko, L. & Rhodes, J. A. Split scores: a tool to quantify phylogenetic signal in genome-scale data**, submitted, available at http://arxiv.org/abs/1608.00942** (2016).

710
711

712 Table 1: CBSD leaf symptom severities and types on plants infected by *Cassava brown streak virus*
713   and *Ugandan cassava brown streak virus*
714

Virus species	Number of plants with CBSD foliar symptom severity score[1]				Number of plants with CBSD foliar symptom type[2]		
	2	3	4	5	SW	LL	SL
CBSV	5	7	1	0	3	8	2
UCBSV	24	5	0	0	7	21	1
CBSV + UCBSV	9	6	0	0	5	10	0
Total	38	18	1	0	15	39	3
Percentage	66.7	31.6	1.7	0.0	26.3	68.4	5.3

715
716

[1]Foliar CBSD symptom severity score based on 1-5 scale; 1 = no visible symptoms, 2 = mild vein
yellowing or chlorotic blotches on some leaves, 3 = pronounced/extensive vein yellowing
or chlorotic blotches on leaves, but no lesions or streaks on stems, 4 = pronounced/extensive
vein yellowing or chlorotic blotches on leaves and mild lesions or streaks on stems, 5 =
pronounced/extensive vein yellowing or chlorotic blotches on leaves and severe lesions or
streaks on stems, defoliation and dieback.
[2]Types of foliar CBSD symptoms based on distribution of leaf chlorosis and stem lesions on the
plant; systemic and on the whole plant (SW), systemic on leaf or stem parts but localized (SL),
only on lower leaves (LL).
726

Table 2. Next generation sequencing data for samples from cassava brown streak disease symptomatic plants collected in Uganda.

Sample ID	Accession number	Virus	No. of reads obtained	No. of reads after trimming	Number of contigs produced (CLC)	Contig length (CLCGW, nt)	Average coverage (CLCGW)	Number of reads mapped to contig of interest	Ref seq. used for mapping	Length of consensus sequence from mapping (Geneious)	No. reads mapped to ref. sequence	Average coverage (Geneious)	Final sequence length (Coding region only)
U1		CBSV	23,335,344	23,053,082	726	3919, 2214	31, 24	1264, 549	KR108828	8,893	2,233	25	8,748
U4		CBSV	21,844,716	21,582,374	621	8,949	255	23,658	KR108828	8,949	22,987	256	8,748
U8		UCBSV	23,648,990	23,373,606	1,008	8,954	366	33,778	KR108836	9,563	178,117	393	8,700

731 Table 3. *Cassava brown streak virus* (CBSV) amino acid (AA) sites under positive selection
732   (analyses method: SLAC Hy-Phy). There were no sites under positive selection for
733   *Ugandan cassava brown streak virus* (UCBSV).
734
735

Gene	CBSV AA site under positive selection
P1	44, 46, 50, 174, 224, 230, 250, 283, 288, 349, 358
P3	415, 455, 467, 472, 499, 525, 618
6K1	658, 678
CI	735, 761, 820, 827, 848, 852, 894, 935, 1218
VPg	1465
NIa	1620, 1645, 1704, 1754, 1785
NIb	1879, 1880, 1890, 1907, 1929, 2109, 2145, 2156, 2161, 2285
HAM1	2320, 2345, 2404, 2432, 2453, 2475, 2519
CP	2550, 2555, 2588, 2611, 2631, 2635, 2640, 2659, 2728, 2745, 2783, 2818, 2843, 2860, 2877, 2884

751 Table 4. Rates of evolution tested using CODEML implemented in PAML. H_O was CBSV and UCBSV
752   have equal rates of evolution (one omega; model = 0), while H_1 was that CBSV and
753   UCBSV have different rates of evolution (two omegas; model = 2).
754

Gene	Assumptions	K (ts/tv rate ratio)	w (omega Dn/Ds) 0	w (omega Dn/Ds)1	Likelihood Ratio Test Statistic (if greater than 3.84 reject H_0) H_0=equal rates
WGS	UCBSV and CBSV equal rates	5.90944	0.06358		
	UCBSV and CBSV different rates	5.9598	0.05518	**0.07622**	26.29*
P1	UCBSV and CBSV equal rates	5.07336	0.10394		
	UCBSV and CBSV different rates	5.05456	0.09047	**0.12203**	4.61*
P3	UCBSV and CBSV equal rates	5.17197	0.08635		
	UCBSV and CBSV different rates	5.20559	0.0764	**0.10198**	2.22
6K1	UCBSV and CBSV equal rates	19.54143	0.00969		
	UCBSV and CBSV different rates	19.69676	**0.01527**	0.00316	3.01
CI	UCBSV and CBSV equal rates	9.9388	0.01722		
	UCBSV and CBSV different rates	8.06276	0.0155	**0.01977**	0.73
6K2	UCBSV and CBSV equal rates	8.40649	0.04684		
	UCBSV and CBSV different rates	8.82738	0.02354	**0.11057**	6.74*
VPg	UCBSV and CBSV equal rates	5.852	0.05759		
	UCBSV and CBSV different rates	5.87009	0.054	**0.06323**	0.29
NIa	UCBSV and CBSV equal rates	8.01105	0.02932		
	UCBSV and CBSV different rates	8.68283	0.01408	**0.06719**	29.95*
NIb	UCBSV and CBSV equal rates	6.07872	0.05329		
	UCBSV and CBSV different rates	6.14047	0.0452	**0.06508**	5.18*
HAM1	UCBSV and CBSV equal rates	7.25177	0.16144		
	UCBSV and CBSV different rates	7.2343	**0.17929**	0.14007	2.23
CP	UCBSV and CBSV equal rates	13.09297	0.06075		
	UCBSV and CBSV different rates	13.26752	0.05606	**0.07155**	1.29
			faster rate UCBSV	**faster rate CBSV**	* Rates are different

757 Table 5. Support for Clades A – G (Figure 3) in individual gene trees and whole genome analyses. Table entries represent
758 posterior probabilities from analysis with MrBayes, except values reported for SVDQ, which are bootstrap
759 proportions. Support values below 95% are indicated in bold, and '--' indicates that the clade was not present.

Genomic region	Clade A	Clade B	Clade C	Clade D	Clade E	Clade F	Clade G
P1	99.98	99.98	96.98	99.72	99.54	--	99.98
P3	99.98	99.98	98.57	99.98	99.98	--	99.98
6K1	**94.66**	99.85	98.87	99.95	--	--	98.42
CI	99.98	99.96	99.76	99.98	99.98	99.99	99.99
6K2	97.65	99.98	**71.52**	**91.70**	**66.46**	--	98.75
Vpg	99.98	99.99	--	99.18	**64.55**	--	99.99
NIa	99.96	99.99	99.98	99.41	**76.31**	--	**89.34**
NIb	99.99	99.98	99.98	99.99	99.98	--	99.98
HAM	--	--	--	99.89	99.83	--	98.55
CP	--	--	--	99.59	99.50	--	99.89
Whole genome	1.00	1.00	1.00	1.00	1.00	1.00	1.00
SVDQ	1.00	1.00	1.00	1.00	**87.00**	**44.00**	1.00

764   Figure Legend

766   Figure 1: Cassava brown streak disease symptoms on leaves and stems of sampled plants; **(a)**
767   Chlorosis along secondary and tertiary leaf veins of CBSV-infected plant of cultivar TME
768   204 (severity score 3), **(b)** Cultivar TME 14 plant with dual CBSV+UCBSV infection showing
769   chlorosis on secondary or tertiary veins, reverse chlorosis (general chlorosis and green
770   area along veins) (severity score 3), **(c)** UCBSV-infected plant of cultivar TME 204
771   exhibiting chlorosis on secondary veins, reverse chlorosis, chlorotic spots and mild stem
772   lesions (severity score 3), **(d)** Very severely diseased plant (severity score 5) of cultivar
773   TME 14 infected with both CBSV and UCBSV, and having chlorosis on leaves, severe stem
774   lesions/brown streaks, defoliation, stem dieback.

776   Figure 2. Genetic diversity of CBSV and UCBSV using the <u>S</u>ynonymous <u>N</u>on-synonymous <u>A</u>nalysis
777   Program (SNAP v2.1.1) implemented in the Los Alamos National Laboratory HIV-
778   sequence database ([http://www.hiv.lanl.gov)](http://www.hiv.lanl.gov) [50]. UCBSV is on the top panel, CBSV at
779   the bottom. The 10 gene segments are labeled from P1-CP.

781   Figure 3. Species tree generated from SVD Quartets using the whole genome sequences. Colors
782   at the tips are based on country of origin. Branches with mixed colors indicate a clade
783   that contains samples with mixed country of origin. For example, the ancestral branch
784   of UCBSV TZ Tan 23 KR108839 and UCBSV UG MI B3 FJ039520 is colored red and orange
785   to indicate a clade with sampled with mixed country of origin.

787   Figure 4. Computed SVD Score with the split defined by CBSV vs. UCBSV across the genome in
788   windows of 500 bp, sliding in increments of 100 bp, and resulting SVD Scores plotted
789   across the genome. Boundaries between genes are marked with vertical lines to further
790   characterize the CBSV and UCBSV genomes. Rates of molecular evolution were
791   estimated using CODEML implemented in PAML (Phylogenetic Analysis by Maximum
792   Likelihood)[54]. The results are shown for each gene and D represents the difference in
793   likelihoods from the null hypothesis (CBSV and UCBSV have equal rates) and the
794   alternative hypothesis (CBSV and UCBSV have different rates).

Figure 1: Cassava brown streak disease symptoms on leaves and stems of sampled plants; **(a)**
Chlorosis along secondary and tertiary leaf veins of CBSV-infected plant of cultivar TME 204
(severity score 3), **(b)** Cultivar TME 14 plant with dual CBSV+UCBSV infection showing chlorosis
on secondary or tertiary veins, reverse chlorosis (general chlorosis and green area along veins)
(severity score 3), **(c)** UCBSV-infected plant of cultivar TME 204 exhibiting chlorosis on secondary
veins, reverse chlorosis, chlorotic spots and mild stem lesions (severity score 3), **(d)** Very severely
diseased plant (severity score 5) of cultivar TME 14 infected with both CBSV and UCBSV, and
having chlorosis on leaves, severe stem lesions/brown streaks, defoliation, stem dieback.
804

806 Figure 2. Genetic diversity of CBSV and UCBSV using the Synonymous Non-synonymous Analysis Program (SNAP v2.1.1) implemented in the
807   Los Alamos National Laboratory HIV-sequence database (http://www.hiv.lanl.gov) [50]. UCBSV is on the top panel, CBSV at the
808   bottom. The 10 gene segments are labeled from P1-CP.
809

Figure 3. Species tree generated from SVD Quartets using the whole genome sequences. Colors at the tips are based on country of origin. Branches with mixed colors indicate a clade that contains samples with mixed country of origin. For example, the ancestral branch of UCBSV TZ Tan 23 KR108839 and UCBSV UG MI B3 FJ039520 is colored red and orange to indicate a clade with sampled with mixed country of origin.

Figure 4. Computed SVD Score with the split defined by CBSV vs. UCBSV across the genome in windows of 500 bp, sliding in increments of
100 bp, and resulting SVD Scores plotted across the genome. Boundaries between genes are marked with vertical lines to further
characterize the CBSV and UCBSV genomes. Rates of molecular evolution were estimated using CODEML implemented in PAML
(Phylogenetic Analysis by Maximum Likelihood)[54]. The results are shown for each gene and D represents the difference in
likelihoods from the null hypothesis (CBSV and UCBSV have equal rates) and the alternative hypothesis (CBSV and UCBSV have
different rates).

823
824
825