1   **Landscape connectivity of a noxious invasive weed:**

2   **human-aided or natural dispersal?**

4   Diego F. Alvarado-Serrano (corresponding author)
5   Department of Ecology and Evolutionary Biology
6   University of Michigan
7   E-mail: dalvarad@umich.edu
8
9   Megan Van Etten
10   Department of Ecology and Evolutionary Biology
11   University of Michigan
12   E-mail: mvanette@umich.edu
13
14   Shu-Mei Chang
15   Department of Plant Biology
16   University of Georgia
17   E-mail: smchang@uga.edu
18
19   Regina S. Baucom
20   Department of Ecology and Evolutionary Biology
21   University of Michigan
22   E-mail: rsbaucom@umich.edu
23

Landscape connectivity of a noxious invasive weed: human-aided or natural dispersal?

ABSTRACT

Examining how the landscape may influence gene flow is at the forefront of

understanding population differentiation and adaptation. Such understanding is crucial in light of

ongoing environmental changes and the elevated risk of ecosystems alteration. In particular,

knowledge of how humans may influence the structure of populations is imperative to elucidate

their role in shaping the evolution of other species, and specifically how humans may alter the

balance between genetic drift and selection. Here we characterize the population genetic structure

of *Ipomoea purpurea*, a noxious invasive weed, and assess the relative roles of natural and

human-driven landscapes on genetic differentiation. By combining rigorous statistical analyses

and a combination of different molecular markers, we detect both common and marker-specific

patterns of genetic connectivity and identify human-aided migration as an important component

shaping the evolutionary history of this species. In particular, we identified human population

density as an important predictor of pairwise population differentiation, suggesting that the

agricultural and/or horticultural trade may be involved in maintaining some level of connectivity

across agricultural fields. Climatic variation, primarily temperature, appears as an additional

predictor when considering agricultural fields in the northern United States. We discuss the

implications of these results and the approach we followed in the context of understanding

agricultural weed and invasive species' expansions, as well as the challenges and promises of

current landscape genetics research.

Keywords: human-aided migration, landscape genetics, morning glory, population structure

INTRODUCTION

Elucidating routes and levels of migration between subpopulations of a species is essential to understand the interplay between gene flow, adaptation, genetic drift, and selection, and hence the forces that shape its evolutionary trajectory (Barrowclough 1980; Slatkin 1985). Landscape features—such as rivers, mountain ranges, crop fields, and urban areas—can impact levels of gene flow between populations by determining dispersal rates and routes (McRae 2006; Cushman *et al.* 2006) as well as by influencing the likelihood of successful establishment of immigrants (Nosil *et al.* 2008; Sexton *et al.* 2014; Wang & Bradburd 2014). Landscape features can also indirectly condition the effect of gene flow through its effect on local effective population sizes since the actual role that migration plays in the evolution of a species is driven by the fraction of the local population size that correspond to immigrants (Wright 1949; Slatkin 1985). Consequently, the landscape, loosely defined as an area with spatially variable biotic and abiotic factors (Holderegger *et al.* 2010), creates the stage for spatially heterogeneous functional population connectivity, conditioned by species' specific physiological tolerances and behavioral preferences (Clobert *et al.* 2012). In this way, the landscape plays a pivotal role in the evolution of species.

Previous studies have identified common responses of species to their surrounding landscape across a range of very different systems, including evidence of frequent long distance dispersal (Berthouly-Salazar *et al.* 2013), dispersal over large unsuitable areas (Manel & Holderegger 2013), and a prevalent mixed effect of local and intervening landscapes on structuring populations (Sexton *et al.* 2014). Such findings have led to novel emergent evolutionary hypotheses that better represent the complex interaction of processes determining

the evolutionary fate of natural populations (Dyer 2015a). However, our current knowledge of the

interplay between landscape features and the genetic structure of populations comes mostly from

human-avoider species (Blair 2001) facing indirect human impact (e.g., Culley *et al.* 2007;

McRae & Beier 2007; Diniz-Filho *et al.* 2009; but see Harris *et al.* 2016). In contrast to species

that almost exclusively depend on natural dispersal agents, species in heavily human-dominated

ecosystems may exploit human activities to maintain gene flow among populations and expand

their ranges (Everman & Klawinski 2013; Fountain *et al.* 2014). Such species may be capable of

maintaining population connectivity over vast geographic ranges (Trakhtenbrot *et al.* 2005) by

overcoming landscape features that would otherwise represent natural barriers and reach dispersal

distances that could be orders of magnitude greater than those attain under natural agents or do it

under much smaller time frames (Mack & Lonsdale 2001; Ricciardi 2007). In this way, by

facilitating dispersal humans have the potential to: i) condition the balance between drift and

selection (Slatkin 1985; Lenormand 2002), ii) introduce relevant genetic variation to local

populations (Kolbe *et al.* 2004), iii) prevent local extinction or favor recolonization (Fountain *et*

al. 2014), and alter the overall genetic constitution of populations (Bataille *et al.* 2011).

Human-aided migration—intentional or unintentional— is particularly prevalent in plants

(Hodkinson *et al.* 2007; Wichmann *et al.* 2009; Auffret & Cousins 2013), where it has had major

impacts on the distribution of species and stability of communities (Simberloff 2013 and

references therein). Yet, the open question remains: how important is human-aided migration

relative to migration through other more natural means in maintaining population connectivity in

plant populations? This question is further complicated in plants because in this group dispersal is

a two-step process that involves the movement of both pollen and/or seeds (Holderegger *et al.*

2010), and these separate vectors of gene flow entail the movement of different genetic states

(haploid vs. diploid) as well as different dispersal agents and mechanisms (i.e., wind, water,

animal-mediated) (Levin & Kerster 1974).

A particularly amenable system to study the interaction between natural and human-aided

dispersal comes from agricultural weed populations. Agricultural weeds face a highly dynamic

landscape characterized by frequent spatial rearrangements (expansion of agricultural front,

increased fragmentation) and a constantly changing environment (crop rotation, agricultural

chemical use, climatic abnormalities) (Menchari *et al.* 2007; Meehan *et al.* 2011) that certainly

impact their opportunities for survival and local adaptation through its effect on population

connectivity (Margosian *et al.* 2009). Under these conditions, human-aided migration is expected

to be critical for weeds' success, but knowledge on how or if weedy plant populations are able to

maintain connectivity through such complex landscape matrix of croplands, grasslands, natural

and urban areas is limited. This is a striking omission given that agricultural weeds impose severe

economic costs (on the order of 33B USD per year in US agriculture alone, Pimentel *et al.* 2005).

As a first step into investigating the impact of human activities on structuring genetic

diversity, we estimate the intensity and extent of migration from genetic data and evaluate how

multiple landscape features influence genetic connectivity of a noxious agricultural weed, *I.*

purpurea, using two different sets of molecular markers (nuclear microsatellites and a genome-

wide panel of SNPs). Specifically, we ask the following questions: 1) Which natural or human-

influenced landscape features—soils, elevation, climate, landcover, crop types, human population

density—promote or constrain genetic connectivity between populations of this agricultural

weed? 2) Do human-associated landscapes disproportionally influence genetic connectivity in

this species, suggesting that this weed's evolutionary fate is strongly linked to human-aided

dispersal?, and 3) what additional insights can we gain from a broader representation of the

genome than traditionally used in landscape genetics studies (typically microsatellites and

organelle DNA)? Taken together, the answers to these questions offer deeper knowledge of the

interaction between human activities, landscape features, and population structure of noxious

weeds and hence contribute to improve effective management and control of these damaging

plants. More generally, these answers contribute to deepen our understanding of the interaction

between geographic setting and population differentiation, adaptation, and persistence (Taylor *et*

al. 1993).

MATERIALS AND METHODS

Study system

Ipomoea purpurea, the common morning glory, is an agricultural weed well suited for the

study of human-aided migration on population connectivity because of its relatively well-

characterized biology, its recent introduction into the United States, and its close association with

agricultural crops. Its tight association with humans since its introduction allows for assessing the

impact of recent landscapes on structuring genetic variance under the initial simplifying

assumption of evolutionary equilibrium (Marko & Hart 2011). *Ipomoea purpurea* is a noxious

weed with a widespread distribution that includes highly heterogeneous landscapes in the

Eastern, South- and Mid-western regions of the United States (Culpepper 2006; Webster &

Nichols 2012). It is a self-compatible annual bumblebee-pollinated vine, with heavy seeds, and is

found primarily in agricultural fields and disturbed areas (Tiffin & Rausher 1999; Baucom &

Mauricio 2008). It is currently one of the most problematic weeds of agriculture (Webster &

Nichols 2012) and is capable of infestations leading to substantial decline in crop (although not

quantified for *I. purpurea*, related *Ipomoea* species may cause declines up to 80% of crop yield,

Rogers *et al.* 1996). This species exhibits in general low-level resistance to the commonly used

herbicide glyphosate (Baucom & Mauricio 2004, 2008), although the exact resistance level varies

widely among populations of this species (Kuester *et al.* 2015). This species is also a major

concern for conservation given its naturalization in multiple regions throughout the world and its

aggressiveness as an invasive (Chaney & Baucom 2012; Fang *et al.* 2013). Previous analyses of

its genetic diversity and differentiation using a panel of 15 nuclear microsatellite markers identify

limited population structure and a less-than definitive isolation-by-distance pattern (Kuester *et al.*

2015). It remains to be seen, however, the role that landscape features, both natural and human-

driven, play in structuring genetic variation within this species.

Data compilation

To capture the plausible effect of both natural and disturbed landscapes on structuring genetic

diversity in *I. purpurea*, we compiled a diverse set of GIS data for the continental US from a

variety of sources (Table S1). These data encapsulate human activities (human population

density, landcover, and planted crops) as well as the geographical setting of *I. purpurea*

(elevation, climate—19 variables summarizing central tendencies and variability patterns in

temperature and precipitation, soil—8 variables summarizing the texture, pH, and organic and

inorganic content of the top 20cm of soil). We first processed all these data into landscape layers

at a common spatial resolution of 10km^2 and a common spatial extent around the US states with

available samples (Fig. 1). These spatial resolution and extent were chosen to maintain a practical

balance between scale and analytical manageability given available computational resources. To

reduce dimensionality, we opted to perform two separate Principal Component Analyses (PCAs)

on the 19 climatic and 8 soil layers, respectively. For all subsequent analyses we kept the

resulting first principal component of each of these analyses, which accounted for over 60% of

the variance and primarily summarized temperature temporal gradients and soils' pH and

sandiness, respectively (Table S2).

With the objective of estimating the genetic connectivity of populations of *I. purpurea*, we

compiled genetic data on an extensive panel of 15 previously optimized microsatellite loci

(Molecular Ecology Resources Primer Development Consortium; *et al.* 2013). These data

encompass a total of 597 individuals from 31 localities (with a minimum of 8 individuals per

locality) (Fig. 1; Table S3). All individuals were collected in 2012 from farms across the range of

I. purpurea in the United Sates (for further details and basic genetic variability analyses see

Kuester *et al.* 2015). In addition, to obtain a more comprehensive representation of the genome of

I. purpurea and assess the robustness of results in light of coalescent and mutational variance

(Nielsen & Slatkin 2013), we generated a Next Generation Sequencing (NGS) dataset from a

subsample of individuals. In conjunction with the microsatellite data, this complementary dataset,

which offers a more widespread representation of this species' genome, is expected to lead to

more comprehensive population genetic inferences (Bohonak & Vandergast 2011; Epps &

Keyghobadi 2015).

To generate the NGS dataset, we constructed genome-wide Genotype By Sequencing (GBS)

libraries for 80 individuals sampled across 8 of localities (10 individuals per locality; Fig 1). The

GBS library was developed using 7ng of genomic DNA extracted from leaf or cotyledon tissue

using SNPsaurus' (Oregon, USA) nextRAD technology, which uses a selective PCR primer to

amplify consistent genomic loci among individuals. Similarly to RAD-Seq sequences (Rowe *et*

al. 2011) in which the DNA flanking a restriction enzyme cut site is selected for amplification,

nextRAD amplifies sequences that correspond to the DNA downstream of a short selective

priming site. Samples were first fragmented and then ligated to short adapter and barcode

sequences using a partial Nextera reaction (Illumina; California, USA) before being amplified

using Phusion® Hot Start Flex DNA Polymerase (New England Biolabs; Massachusetts, USA).

The 80 dual-barcoded PCR-amplified samples were pooled and the resulting libraries were

purified using AMPure XP beads (Agencourt Bioscience Corporation; Massachusetts, USA) at

0.7x. The purified library was then size selected to 350-800 base pairs and sequenced using two

runs of an Illumina NextSeq500 sequencer (Genomics Core Facility, University of Oregon).

The resulting sequences were analytically processed by combining the reads of 16 randomly

selected individuals (of the 80 sequenced) to create a pseudo-reference genome. This was done

after removing loci with read counts above 20,000, which presumably corresponded to repetitive

genomic material, and loci with read counts below 100, which presumably corresponded to off-

target or read errors. The filtered reads were aligned to each other using BBMap

(http://sourceforge.net/projects/bbmap/). All parameters were set to default values with the

exception of minimum sequence identity, which was set to 0.93 to identify alleles. A single read

instance was chosen to represent the locus in the pseudo-reference. This resulted in a total of

263,658 loci. All reads from each of the 80 individuals were then aligned to the pseudo-reference

using BBMap (http://sourceforge.net/projects/bbmap/) and converted to a vcf genotype table,

using Samtools (Li *et al.* 2009) and bcftools (Li 2011), after filtering out nucleotides with a

quality score of 10 or worst. The resulting vcf table was filtered using vcftools (Danecek *et al.*

2011) for SNPs with a minimum allele frequency of 0.02, a minimum read depth of 5, and a

maximum 15% of missing data. This resulted in 9774 variable regions. Loci were further filtered

using vcftools to exclude loci with less than 5 high quality base-calls and with more than 20%

missing data or an average of less than 20 high quality base calls. This resulted in a final panel of

8210 Single Nucleotide Polymorphisms (SNPs) that we used in all subsequent analyses.

Population structure analyses

We first conducted a series of preliminary analyses to characterize the overall genetic

structure of *I. purpurea*. All analyses were run separately for the microsatellite (SSR, hereafter)

and SNP datasets given their different sampling and geographic coverage (Fig. 1). First, we

examined population differentiation by estimating F_{ST} using GenAlex v6.5 (Peakall & Smouse

2012) (because similar global F_{ST} and R_{ST} estimates were obtained for the SSR dataset, we opted

to report F_{ST} values only to allow direct comparisons with the SNP dataset). We then estimated

contemporary effective population size for each sampled locality in NeEstimator v2 using the

excess heterozygous method (Do *et al.* 2014). We performed this latter analysis to assess the

possibility of whether effective rate of migration (*Nm*) may be asymmetric in response to

differences in population size (e.g., greater migration into satellite populations).

In addition, we assessed population admixture and spatial genetic clustering using 2 spatially

explicit Bayesian clustering algorithms: BAPS (Corander *et al.* 2004) and TESS (Chen *et al.*

2007). BAPS was run using its spatial clustering algorithm (Cheng *et al.* 2013) with 10 runs for

each K value tested, whereas TESS was run using the admixture algorithm and a BYM model

(i.e., using a spatial prior of admixture proportions; Durand *et al.* 2009b) with 10 runs per K

value, and without using geographic weights. The TESS model, with the lowest DIC was chosen

as the optimal model (Durand *et al.* 2009a). For both analyses K values ranging from 2 to the

maximum number of sampled locations (31 and 10 for the SSR and SNP datasets, respectively)

were tested. Results were generally consistent between BAPS and TESS analyses; thus, here we

report exclusively the TESS results. Additionally, following Wang et al. (2009), we

complemented these analyses with an Analysis of Molecular Variance (AMOVA; Excoffier *et al.*

1992) run in GenAlex (Peakall & Smouse 2012) using 9999 permutation replicates. This analysis

partitioned the variance into regions based on the spatial genetic clusters previously identified to

quantify the fraction of the genetic variance explained by spatial genetic clusters, and hence the

relative importance of spatial genetic structure in *I. purpurea*.

Finally, we investigated population connectivity by estimating levels of recent migration

between sampled localities through the identification of individuals of mixed ancestry using

BayesAss (Wilson & Rannala 2003). BayesAss is a program that uses individual multilocus

genotypes and a Markov Chain Monte Carlo (MCMC) algorithm to probabilistically distinguish

between immigrants and long-term native individuals (Wilson & Rannala 2003). It provides

posterior probability distributions of individual immigrant ancestries (i.e., the probability that an

individual is a first or second generation migrant from each of the populations in the sample). We

ran BayesAss for 6 million generations using default parameter settings, and discarded the first 2

million generations as burn-in (Dyer 2009). For each marker dataset, we repeated this analysis 3

times (for a total of 18 million generations) and combined the results from the 3 replicates for our

final inference. Then, using a posterior probability cut-off of 0.75 we assign individuals'

ancestry. It is important to note that because of computational limits we had to randomly

subsample our set of SNPs to 400 SNPs for this analysis. We complemented this analysis with a

spatial autocorrelation analysis run in GenAlex (Peakall & Smouse 2012) to evaluate the overall

spatial scale of genetic turnover.

Landscape genetics analyses

After assessing overall population structure of *I. purpurea*, we evaluated the association

between landscape features and genetic differentiation. To do this, we first converted our

pairwise F_{ST} estimates into conditional genetic distances (Dyer *et al.* 2010) using GeneticStudio

(Dyer 2009). Briefly, conditional distances are measures of pairwise genetic distance derived

from population networks, constructed based on the degree of genetic similarity between sampled

localities (Dyer & Nason 2004). Because these networks are pruned based on the principle of

conditional independence of the total among population genetic covariance (the specific method

of pruning used is edge deviance; Magwene 2001), conditional distances reflect genetic similarity

between localities that better capture direct gene flow as opposed to connectivity driven by

intervening localities (Dyer 2015b).

The climate, crops, elevation, landcover, population density, and soils landscape layers (Table

S1) were converted into landscape resistance layers by assigning a resistance value to each

landscape feature in these layers to reflect the difficulty that each feature offers to the movement

of gametes or individuals (i.e., pollen or seeds). It is important to note that in contrast to previous

studies that typically rely on expert opinion for resistance assignment, we utilized an unbiased

statistical optimization to avoid the sensitivity of results to subjective resistance assignment

(Spear *et al.* 2010). Specifically, resistance values were optimized through a genetic algorithm

approach, which is a heuristic stochastic optimization algorithm that emulates the process of

inheritance, mutation, and natural selection (Mitchell 1996). Briefly, in this search algorithm a

population of individuals with traits encoded by unique combinations of model parameters

(resistance assignment proposals in our case) is allowed to compete with each other based on the

fitness associated with the traits it carries (Peterman *et al.* 2014). Specifically, in Peterman's

(2014) implementation of this algorithm, which we followed here, individuals' fitness is

estimated by the relative quality of a MLPE.lmm model (Maximum Likelihood Population

Effects – Linear Mixture Model) that evaluates the association between pairwise genetic distance

and landscape cumulative resistance between localities, estimated in Circuitscape (Shah &

McRae 2008). Individuals with parameter settings (resistance assignments in our case) that result

in better models, as measured by a Deviance Information Criterion (DIC) score, are preferentially

represented in the following generation. Offspring modifications introduced by mutations (i.e.,

small resistance assignment perturbations) allow for exploration of the parameter space. The

algorithm is stopped once a large number of generations have passed without significant

improvement in fitness.

We implemented Peterman's (2014) algorithm in R (package ResistanceGA; Peterman 2014)

allowing for the independent optimization of each of our landscape layers. The optimal resistance

landscapes identified in this way were then used to run a final univariate MLPE.lmm model to

identify the association between landscape features and conditional genetic distances between

localities. In addition, we run separate simple and partial db-RDA (distance-based Redundancy

Analyses; Legendre & Anderson 1999)—the latter uses the geographic distance between

populations as a covariate to account for the concurrent increase in cumulative resistance with

geographic distance. We opted for db-RDA instead of the most commonly used partial Mantel

test given the statistical issues of the latter (Raufaste & Rousset 2001; Guillot & Rousset 2013).

We run this latter analysis using the package vegan (Oksanen *et al.* 2015) in R and assessed

significance with 9999 permutations. Finally, to identify the relative contribution of natural and

human-driven landscape features we ran a Multiple Regression on Distance Matrices (MRDM;

Legendre *et al.* 1994), which has been identified as one of the best performing methods for

evaluating the interplay between landscape features and genetic connectivity (Balkenhol *et al.*

2009). Before running these regressions, we standardized all optimized resistance layers to mean

of zero and variance of 1 (Dyer *et al.* 2010). These final regressions included geographic distance

as a predictor and were run in R (using package ecodist; Goslee & Urban 2007) using 10,000

permutations to assess significance. In none of our analyses did we implement a Bonferroni

correction for multiple testing because of the overly conservative nature of this correction

(Nakagawa 2004; Glickman *et al.* 2014). Instead we applied a false recovery rate correction

(Benjamini & Hochberg 1995) using the function *p.adjust* in R (R Core Development Team

2016).

RESULTS

Population structure

The set of preliminary genetic analyses indicated that *I. purpurea* sampled localities were

in no major violation of Hardy-Weinberg equilibrium, as judged by the small difference between

expected and observed heterozygosity (mean He = 0.294±0.014 and 0.250±0.001; mean Ho =

0.291±0.009 and 0.260±0.001, respectively for SSR and SNP datasets). Levels of expected and

observed heterozygosity for the SSR dataset were only slightly greater than those estimated for

the SNP dataset (Table S3). Likewise, the estimated mean effective population size per sampled

locality was only slightly greater and more variable for the SSR dataset than for the SNP dataset

(13.71±5.59, 9.49±0.13, respectively; Table S3), but in neither case was there salient evidence of

a plausible source-sink dynamic, as judged by the similar effective sizes among populations

(Table S3; Diffendorfer 1998). Accordingly, spatial structure in this species was limited, as

evidenced by the modest F_{ST} estimates (0.151 and 0.140, respectively for SSR and SNP datasets)

and the admixed genetic composition of individuals (rather more structured in the SNP dataset;

Fig. 2). Furthermore, although TESS results indicated the existence of partially discernable

spatial genetic clusters (4 and 5 clusters for the SSR and SNP datasets, respectively; Fig. 2),

variation among the inferred spatial genetic clusters explained less than 10% of the variance

(Table 1). It is especially noteworthy that the inferred spatial genetic clusters were constrained to

geographically contiguous areas in the SNP dataset, but not in the SSR dataset (Fig. 2).

Despite these overall similarities between datasets, estimates of recent ancestry differed

between them. The analysis on the SSR dataset indicated that migration among localities is

widespread and hardly geographically constrained, with only four localities being primarily

constituted of native individuals (Fig. 3a). Across localities, on average 73.65% of individuals

were inferred to be 1st or 2nd generation immigrants. On the other hand, the analysis on the SNP

dataset showed that most populations have a limited number of recent immigrants, but that the

relatively few inferred immigrants (on average 27.42% of individuals across localities) did not

come exclusively from geographically proximate localities (Fig. 3c). Nevertheless, long distance

migration events were estimated to be infrequent in this dataset relative to the SSR dataset, in

which migration across geographically spread states was common (e.g., between North Carolina

and Tennessee, Fig. 3a,c). Furthermore, of the 6 localities shared between datasets only 1 was

similarly inferred to comprise mostly native individuals by both sets of markers; inferences for

the other five localities differed between datasets. Accordingly, the SSR-based pruned

conditional genetic network (Dyer *et al.* 2010) was significantly more interconnected (vertex

connectivity = 5 ; White & Harary 2001) than the SNP-based network (vertex connectivity 0)

(Fig. 3b,d). This relatively low complexity of the SNP-based network further confirmed the

inference of locally constrained migration in this dataset compared to more widespread migration

inferred with the SSR dataset.

In line with these findings, the spatial autocorrelation analyses indicated that genetic

clustering extended over wider spatial scales for the SSR dataset than for the SNP dataset (Fig.

S1). Specifically, nearby localities showed a significant tendency to be genetically similar to each

other up to a considerably greater distance in the case of the SSR dataset relative to the SNP

dataset (Fig. 3). Furthermore, at greater distances no significant association was recovered for the

SNP dataset, further confirming their genetic isolation from each other, whereas a significant

negative association (i.e., statistical tendency to be more different from each other than expected

by chance) was recovered in the SSR dataset.

Landscape genetics

Unsurprisingly given the distinct geographical and environmental ranges covered by each

dataset (Fig. 1), the optimization of landscape resistance layers resulted in alternative optimal

solutions for the SSR and SNP datasets (Fig. S2). It is important to note, however, that a formal

comparison is not possible as the associations recovered are statistical associations driven by the

fit of the resistance parameterization to the data under the statistical model implemented

(Martínez-Abraín 2008). While these associations are expected to recapitulate real biological

properties of the study system, they are constrained to the data at hand. Nevertheless, association

patterns that are robust to the data used are expected to better reflect the actual impact that

landscape features have on gene flow, independently of possible biases introduced by expert

opinion. Therefore, we focus below on the common biological findings between marker types,

while also denoting the most relevant differences. Such differences likely reflect not only the

different environmental ranges covered by each dataset, but most importantly, the particular

genetic connectivity pattern captured by each one of them (Fig. 3).

In spite of the underlying differences in inferred population connectivity (Fig. 3), some of

the resulting landscape resistance layers roughly resemble their counterpart in the other dataset in

terms of their relative resistance allocation (Fig. S2). There were some landscape features that

showed consistent resistance patterns between datasets. For example, cotton-dominated areas and

woody savannas were recovered as highly permeable landscape features, whereas areas

dominated with soybean fields, evergreen forests, open shrublands, and grasslands were

recovered as highly resistant landscape features. This difference between the permeability of

different crop and landcover features matches the prevalence and distribution of these features

across the study area. Soybean dominated areas occupied over 8.7% of the study area and were

most prevalent in its northwestern portion, where *I. purpurea,* being primarily a subtropical vine

(Fang *et al.* 2013)*,* is less prevalent. On the other hand, cotton-dominated areas occupied 0.8% of

the study area and were concentrated in its southern portion, including areas of high *I. purpurea*

concentration. Likewise, woody savannas represented 10.8% of the study system and were

constrained to the southern portion of it, whereas evergreen forests, open shrublands, and

grasslands represent together 1.1% of the study system and are prevalent in areas where *I.*

purpurea, being mainly found on agricultural fields, gardens, and roadsides (Defelice 2001), is

absent. Hence, these landscape associations most likely reflected the environmental preferences

of *I. purpurea*, as environments where this species thrives coincided with environments with a

low resistance assigned, whereas environments unoccupied by this species were assigned greater

resistances.

Given the differences in sampling between datasets (Fig. 1), the most revealing patterns

were similar landscape variables identified as significant (or marginally significant) predictors of

genetic differentiation in *I. purpurea* across datasets (Table 2). Both datasets pointed towards

human-impacted landscapes playing an important direct role in shaping genetic connectivity in

this species. While in both sets of MLPE.lmm models, natural (climate, elevation, and soils) and

human-related landscapes (landcover and human population density) were identified as

significant or marginally significant predictors of genetic similarity between localities, the

variable with the greatest association coefficient and lowest AICc value in these models was in

both cases a variable closely linked to human presence (landcover in the SSR dataset, and human

population density in the SNP dataset; Table 2). In contrast, the dbRDA analyses showed that

none of these features were consistently recovered as a significant predictor of genetic similarity

in the SNP dataset, especially after accounting for geographic distance between populations—

even though geographic distance by itself was not a significant predictor of genetic structure in

this species (i.e., no isolation by distance pattern was recovered). Yet, when considering all

variables together in a multivariate manner while accounting for geographic distance, human

population density was the only variable that remained a significant or marginally significant

predictor of population structure across both SSR and SNP datasets (Table 2). Specifically, these

results indicated that, after considering other landscape variables, human-population-density

resistance was associated with population differentiation across datasets, with high and low

populated areas identified as less favorable areas for migration (although conductance optimum

for the SNP dataset was notably lower than for the SSR datasets; Fig. 4). Accordingly, the

corresponding resistance-based conductance map (Shah & McRae 2008), showed local clusters

of high permeability for both datasets (although slightly less constrained in the SSR dataset),

surrounded by vast areas of high resistance (Fig. S3). It is important to note, however, that these

multivariate regressions explained a small, non-significant proportion of the variance (MRDM R^2

for SSR and SNP dataset were 0.055 (F = 3.776, p-val. = 0.110) and 0.309 (F = 1.275, p-val. =

0.261), respectively).

In addition to population density, climate was also recovered as a significant predictor of

genetic dissimilarity across analyses on the SSR dataset but not for the SNP dataset (Table 3) —

most likely because of the lack of northern samples, from colder areas, for the latter dataset (Fig.

1). Specifically, the first component of the climate PCA (see methods above), which primarily

summarized temperature variation, was consistently associated with genetic differentiation (Table

3), with intermediates values identified as a barrier to dispersal. Warmer areas were identified as

the most conducive to dispersal, followed by colder areas—the latter pattern exclusively driven

by connectivity between the northernmost localities (Fig. 4). The corresponding resistance-based

conductance map showed a less spatially constrained distribution of permeable areas with

connections between southern and between northern localities, and a strong barrier between them

(Fig. S3).

DISCUSSION

Taken together, the population structure analyses indicated limited to moderately high

(depending on the molecular marker) global levels of genetic differentiation in *I. purpurea* and

non-geographically structured migration. This non-geographically structured migration was

inferred to be rather rare when using the SNP dataset (Fig. 3). These results suggest that broadly

distributed populations of this agricultural weed are generally genetically distinct, but there is

some indication of long-distance and putatively human-mediated migration between localities, as

suggested by the recovered association between human population density and genetic similarity.

Such levels of differentiation and long-distance migration strongly contrast with this species'

patchy distribution, which is tightly linked to isolated agricultural patches that are surrounded by

a complex matrix of natural and urbanized areas. As suggested by the landscape genetics

analyses, this matrix does seem to impact connectivity in this species. Specifically, climate and

human population density were robustly recovered as predictors of genetic connectivity in this

species. Of these landscape variables, climate has a stronger effect, as judged by its greater

MRDM coefficient, but only when considering the SSR dataset, which covers the northern

portion of the range (Table 3). Otherwise, population density was the only variable across

datasets with a marginally significant effect—even after accounting for multiple tests. Along with

the recovered pattern of landscape features of high and low permeability, in particular in regards

to crop types (Fig. S2), this finding highlights the important role that humans play in structuring

populations of this species. In addition, the results highlight how inferences about population

structure and patterns of connectivity are dataset-dependent, with marked differences becoming

apparent only after careful dissection of roughly similar F_{ST} and heterozygosity estimates across

molecular markers. Below, we detail each of these novel findings and place them in the context

of agricultural weed movement across the landscape, invasive species, and landscape genetics

practice.

Human impact

Given that *I. pupurea* is a naturalized species in the United States that is found primarily

associated with cultivated crops and horticultural gardens (Defelice 2001; Baucom & Mauricio

2004; Fang *et al.* 2013), the finding that human population density is a predictor of genetic

similarity in this species is at first glance intuitive. Yet, because habitat requirements for

establishment and migration are not always the same, especially for organisms with distinct

migration stages (e.g., pollen or seeds in plants) and dormant stages (Murphy & Lovett-Doust

2004), this finding is not as straightforward as it seems. In particular, the fact that human

population density is an informative predictor throughout the entire sampled distribution, whereas

climate seems to mainly represent a gene flow barrier between southern and northern *I. purpurea*

sites (Fig. S1) highlights the influence that humans have on structuring the populations of this

species and helps to discern the factors involved in the spread of this noxious weed. In this sense,

the results point towards humans not only as likely responsible for the introduction of this weed

into the United States (Fang *et al.* 2013), but also as responsible for facilitating its current spatial

connectivity, and hence its opportunities for thriving in the complex landscapes it inhabits. As

evidenced by the inferred landscape conductivity, human population density seems to primarily

facilitate connectivity at local to intermediate scales (note the clusters of high conductivity

around sampled localities; Figs. 4 and S2), which encompass proximate agricultural fields,

suggesting that factors such as sharing of contaminated agricultural machines, trade between

nearby farmers, or local distribution of contaminated crop seeds are at play (Dastgheib 1989;

Thill & Mallory-Smith 1997; Benvenuti 2007; Boyd & White 2009). At the same time,

considering that i) the horticultural trade has been recognized as the main source of invasive

introductions and spread in the United States (Lehan *et al.* 2013), ii) that *I. purpurea* is an

appreciated horticultural species (Fang *et al.* 2013), and that, given current agricultural practices,

crop seed contamination is unlikely to be a major factor (Economic Research Service 1998), it is

probable that ornamentals' trade between population centers may help explain both the long

distance dispersal events recovered in both datasets (Fig. 3) and the local clustering (Fig. 4).

Alternatively, the impact of human populations on the distribution and abundance of bumblebees

(Martins *et al.* 2013; Jha 2015), which are *I. purpurea*'s predominant pollinators (Ennos 1981;

Baucom & Mauricio 2008), could also be partially responsible for the connectivity patterns

recovered as changes in the pollinators community would have strong effects on gene flow (Jha

& Kremen 2013). In reality a combination of all these factors may be involved.

While further analyses are needed to elucidate the ultimate causes behind the recovered

association between human population density and genetic dissimilarity in *I. purpurea*, our

findings bring much needed information to limit the spread of this noxious weed. Our findings

are not only relevant to *I. purpurea* and to the evolution of herbicide resistance in this species

(i.e., is herbicide resistance evolving independently across populations or is it being disseminated

through human-aided migration?), but also has important implications for other weeds of

agricultural concern as well as other human-exploiter species (Blair 2001), such as other

invasives. Specifically, in line with previous work (Bataille *et al.* 2011; Auffret *et al.* 2014;

Banks *et al.* 2015), the results here point towards the need of better strategies to minimize the

impact that humans have on the spread of species. In particular, our results further support that

humans may not only facilitate the introduction of invasive species into non-colonized areas, but

also contribute to the maintenance of gene flow among naturalized populations (Medley *et al.*

2015), which may be critical in providing relevant genetic variants for increased fitness as well as

prevent inbreeding depression in these newly colonized areas (Kolbe *et al.* 2004; Edelaar &

Bolnick 2012).

Landscape genetics practice

Under the common landscape genetics' assumption of an equilibrium between migration

and genetic drift (Marko & Hart 2011; Dyer 2015b), our results offer a valuable window into the

role that environmental setting plays in structuring genetic diversity. Our analyses take advantage

of recent methodological developments i) that surpass the need of arbitrary landscape resistance

assignment that make inferences sensitive to subjectivity of expert opinion (Dyer 2015a), ii) that

account for the indirect genetic similarity of populations (Dyer & Nason 2004), and iii) that use

rigorous statistical inferences (Balkenhol *et al.* 2009; Peterman *et al.* 2014). Furthermore, in

contrast to the common practice in the field of using one or a few loci that are either uni-

parentally inherited (although a few exceptions exist; e.g. Perry *et al.* 2013), which prevents an

assessment of common patterns across the genome (Bohonak & Vandergast 2011), our inferences

are derived from common findings among two rather different sets of molecular markers. In

doing so, we provide not only a better representation of the genome, and hence, less sensitive

inferences to i) ascertainment bias (Brandström & Ellegren 2008), ii) molecular markers'

idiosyncrasies (Buschiazzo & Gemmell 2006), and iii) coalescent and mutational variance

(Nielsen & Slatkin 2013; Steiner *et al.* 2013), but also the ability to distinguish differences in the

underlying population dynamics. Such differences have strikingly important implications. For

example, when evaluating plausible approaches to the threat of an invasive species such as *I.*

purpurea, recommendations would be quite different depending on whether gene flow is believed

to be relatively widespread (as inferred by the SSR dataset) or whether it is believed to be

minimal (as inferred by the SNP dataset). In this example, it is clear that evidence-based

conservation would clearly benefit from recognizing the current uncertainty in regards to the

exact population connectivity as opposed to automatically relying on a single-marker story.

Given recent advances in next generation sequencing, it seems straightforward to focus on

landscape genomics instead of few loci. Hence, development of methods for explicitly integrating

inferences from multiple genome regions, as it is customary in population genetics, would be of

great value.

Even more important, the differences identified between markers offer the opportunity to

explore the underlying causes for such differences and hence a more in-depth understanding of

the landscape influences on species' genetic structure. While it is theoretically possible that the

differences in inference between the two markers are exclusively driven by the particular

geographic sampling of each dataset, the robust differences that we report between the localities

and regions common to both datasets renders this possibility unlikely. A second alternative is that

the differences are driven by the different mutation rates underlying the two type of markers

(Wang 2010, but see Bohonak and Vandergast 2011). This time-differential hypothesis is based

on the argument that SSR mutation rates are typically estimated to be in the $10e^{-3}$-$10e^{-4}$ range,

whereas SNP mutation rates, although harder to estimate given their occurrence in multiple

heterogeneous genomic regions (Lercher & Hurst 2002), are typically presumed to be on average

much lower. Nonetheless, given mutational rates are still on the scale of thousand of years and

up, most genetic variation seen in current populations would precede the temporal window of the

majority of landscape genetics studies. It is then the sorting of standing genetic variation, rather

than the mutation rate, what would primarily drive genetic dissimilarity between current

subpopulations of a species. Such sorting is thus expected to be specific to different genomic

regions. That is, because of the genome-wide coverage of SNPs and the independent evolutionary

trajectories of SSR and SNP loci, recombination, effective population sizes, and other relevant

evolutionary parameters are expected to be highly heterogeneous across loci, which can play an

important role in the pace at which each loci segregates, making it unclear which temporal scale

would be reflected by each dataset (Ennos 1994; Bohonak & Vandergast 2011). Of all the factors

involved, effective population sizes of each locus, which directly affect the rate of genetic drift of

each loci (Storz *et al.* 2001; Piganeau & Eyre-Walker 2009), is probably the most impacted by

landscape conditions, and hence it is probably a major determinant of patterns of shared variants

across the landscape. Yet, traditionally estimates of effective population size are rarely

incorporated into landscape genetics studies.

In reality, both datasets likely encompass a range of temporal scales, and presumably reflect

both historical and contemporary processes. In this regard, working with an agricultural weed

with a relatively well-known history offers the advantage of better accounting for the historical

component of current genetic variation. Specifically, under the time-differential hypothesis

(Wang 2010; but see Bohonak & Vandergast 2011), it is plausible that the spatial structure

recovered by our SNP dataset is strongly influenced by ancestral subdivision and that the relative

lack of structure recovered by our SSR dataset reflect more recent gene flow. However,

considering the current understanding of the introduction of *I. purpurea* into the United Sates,

this possibility seems unlikely. If this weed was indeed introduced through horticulture from a

European bottlenecked population during the European colonization of North America (Fang *et*

al. 2013), ancestral structure would be unlikely. Likewise, if the species expanded from the point

of introduction, a nested pattern of genetic similarity driven by serial founder events would be

expected (Ramachandran *et al.* 2005; Slatkin & Excoffier 2012)—not a rather distinct clustering

of individual subpopulations as we recovered (Fig. 3). Instead, it seems more likely that the

obvious structure recovered in the SNP dataset reflects the isolation driven by the complex

agricultural matrix *I. purpurea* inhabits and the role that human intervention has had on its

evolutionary trajectory (e.g., by facilitating long distance dispersal events; Fig. 3). Yet, further

analyses (see below) are needed to test this hypothesis.

594  More importantly, the incorporation of both datasets into our analyses highlights a commonly

595 overlooked issue in landscape genetics. That is, any pattern of genetic variation cannot be

596 understood without explicit consideration of species' demographic history as current genetic

597 variation is the result of multiple processes thorough the history of a species (Marko & Hart

598 2011), and different genomic regions reflect different coalescent histories (Nielsen & Slatkin

599 2013). Hence, the importance of incorporating coalescent-based simulations that explicitly model

600 the presumed landscape effects on genetic population structure and that take into account its

601 demographic history (Balkenhol & Landguth 2011; Hoban *et al.* 2012). Advances in this area are

602 already being developed with promising perspectives (He *et al.* 2013; Alvarado-Serrano &

603 Knowles 2014). In this regard, traditional landscape genetics results can be interpreted as a

604 necessary first step towards a more comprehensive understanding of the interaction between

605 landscapes and species evolutionary trajectories. By offering a working hypothesis of the effect

606 of current landscapes on genetic differentiation, traditional landscape genetics results serve the

607 purpose of identifying relevant hypotheses for further testing (Dyer 2015a) and pave the way for

608 rigorous simulation-based assessments of the role of landscape features in promoting or deterring

609 population differentiation.

611 **ACKNOWLEDGEMENTS**

612 We thank Adam Kuester for seed collecting and for contributing valuable data for this study. We

613 also thank Ariana Wilson, Eva Fall, and Dan York for tissue collection. This research was funded

614 by USDA NIFA grants 04180 and 07191to R.S.B.

616 **DATA ACCESSIBILITY**

All data generated is in the process of being archived in Dryad. The corresponding doi would be

made available upon acceptance.

AUTHOR CONTRIBUTIONS

D.F.A.-S. and R.S.B conceived the study. D.F.A.-S. and M.V.E. generated and compiled the

molecular and GIS data. D.F.A.-S. analyzed the data. D.F.A.-S., M.V.E., S.M.C., and R.S.B.

wrote the manuscript. All authors read and approved the final submission.

LITERATURE CITED
625
Alvarado-Serrano DF, Knowles LL (2014) Ecological niche models in phylogeographic studies:

Applications, advances and precautions. *Molecular Ecology Resources*, **14**, 233–248.

Auffret AG, Berg J, Cousins SAO (2014) The geography of human-mediated dispersal. *Diversity*

and Distributions, **20**, 1450–1456.

Auffret AG, Cousins SAO (2013) Humans as Long-Distance Dispersers of Rural Plant

Communities. *PLoS ONE*, **8**.

Balkenhol N, Landguth EL (2011) Simulation modelling in landscape genetics: on the need to go

further. *Molecular ecology*, **20**, 667–70.

Balkenhol N, Waits LP, Dezzani RJ (2009) Statistical approaches in landscape genetics: an

evaluation of methods for linking landscape and genetic data. *Ecography*, **32**, 818–830.

Banks NC, Paini DR, Bayliss KL, Hodda M (2015) The role of global trade and transport

network topology in the human-mediated dispersal of alien species. *Ecology Letters*, **18**,

188–199.

Barrowclough GF (1980) Gene flow, effective population sizes and genetic variance components

in birds. *Evolution*, **34**, 789–798.

Bataille A, Cunningham AA, Cruz M, Cedeño V, Goodman SJ (2011) Adaptation, isolation by

distance and human-mediated transport determine patterns of gene flow among populations

of the disease vector Aedes taeniorhynchus in the Galapagos Islands. *Infection, Genetics and*

Evolution, **11**, 1996–2003.

Baucom RS, Mauricio R (2004) Fitness costs and benefits of novel herbicide tolerance in a

noxious weed. *Proceedings of the National Academy of Sciences*, **101**, 13386–13390.

Baucom RS, Mauricio R (2008) The evolution of novel herbicide tolerance in a noxious weed:

The geographic mosaic of selection. *Evolutionary Ecology*, **22**, 85–101.

Benjamini Y, Hochberg Y (1995) Controlling the false discovery rate [mdash] a practical and

powerful approach to multiple testing. *J. R. Stat. Soc. B*, **57**, 289–300.

Benvenuti S (2007) Weed seed movement and dispersal strategies in the agricultural

environment. *Weed Biology and Management*, **7**, 141–157.

Berthouly-Salazar C, Hui C, Blackburn TM *et al.* (2013) Long-distance dispersal maximizes

evolutionary potential during rapid geographic range expansion. *Molecular Ecology*, **22**,

5793–5804.

Blair RB (2001) Birds and butterflies along urban gradients in two ecoregions of the US. In:

Biotic homogenization (eds Lockwood JL, McKinney ML), pp. 33–56.

Bohonak AJ, Vandergast AG (2011) The value of DNA sequence data for studying landscape

genetics. *Molecular ecology*, **20**, 2477–9; authors reply 2480–2.

Boyd NS, White S (2009) Impact of Wild Blueberry Harvesters on Weed Seed Dispersal within

and between Fields. *Weed Science*, **57**, 541–546.

Brandström M, Ellegren H (2008) Genome-wide analysis of microsatellite polymorphism in

chicken circumventing the ascertainment bias Genome-wide analysis of microsatellite

polymorphism in chicken circumventing the ascertainment bias. *Genome Research*, 881–

887.

Buschiazzo E, Gemmell NJ (2006) The rise, fall and renaissance of microsatellites in eukaryotic

genomes. *BioEssays*, **28**, 1040–1050.

Chaney L, Baucom RS (2012) The evolutionary potential of Baker's weediness traits in the

common morning glory, Ipomoea purpurea (Convolvulaceae). *American Journal of Botany*,

99, 1524–1530.

Chen C, Durand E, Forbes F, François O (2007) Bayesian clustering algorithms ascertaining

spatial population structure: A new computer program and a comparison study. *Molecular*

Ecology Notes, **7**, 747–756.

Cheng L, Connor TR, Sir??n J, Aanensen DM, Corander J (2013) Hierarchical and spatially

explicit clustering of DNA sequences with BAPS software. *Molecular Biology and*

Evolution, **30**, 1224–1228.

Clobert J, Baguette M, Benton TG, Bullock JM, Ducatez S (2012) *Dispersal Ecology and*

Evolution. Oxford University Press, Oxford.

Consortium; MERPD, Aksoy S, Almeida-Val VMF *et al.* (2013) Permanent Genetic Resources

added to Molecular Ecology Resources Database 1 October 2012–30 November 2012.

Molecular Ecology Resources, **13**, 341–343.

Corander J, Waldmann P, Marttinen P, Sillanpää MJ (2004) BAPS 2: enhanced possibilities for

the analysis of genetic population structure. *Bioinformatics (Oxford, England)*, **20**, 2363–9.

Culley TM, Sbita SJ, Wick A (2007) Population genetic effects of urban habitat fragmentation in

the perennial herb Viola pubescens (Violaceae) using ISSR markers. *Annals of Botany*, **100**,

91–100.

Culpepper AS (2006) Glyphosate-Induced Weed Shifts. *Weed Technology*, **20**, 277–281.

Cushman S a, McKelvey KS, Hayden J, Schwartz MK (2006) Gene flow in complex landscapes:

testing multiple hypotheses with causal modeling. *The American naturalist*, **168**, 486–99.

Danecek P, Auton A, Abecasis G *et al.* (2011) The variant call format and VCFtools.

Bioinformatics, **27**, 2156–2158.

Dastgheib F (1989) Relative importance of crop seed, manure and irrigation water as sources of

weed infestation. *Weed Research*, **29**, 113–116.

Defelice MS (2001) Intriguing World of Weeds Tall Morningglory , Ipomoea purpurea (L .)

Roth-Flower or Foe ? *Weed Technology*, **15**, 601–606.

Diffendorfer JE (1998) Testing Models of Source-Sink Dynamics and Balanced Dispersal. *Oikos*,

81, 417–433.

Diniz-Filho JAF, Nabout JC, Bini LM *et al.* (2009) Niche modelling and landscape genetics of

Caryocar brasiliense ("Pequi" tree: Caryocaraceae) in Brazilian Cerrado: an integrative

approach for evaluating central–peripheral population patterns. *Tree Genetics & Genomes*,

5, 617–627.

Do C, Waples RS, Peel D *et al.* (2014) NeEstimator v2: Re-implementation of software for the

estimation of contemporary effective population size (Ne) from genetic data. *Molecular*

Ecology Resources, **14**, 209–214.

Durand E, Chen C, Francois O (2009a) Tess version 2.3 - Reference Manual August 2009 ∗.

Available at: memberstimc. imag. fr/Olivier. Francois/tess. html, 1–30.

Durand E, Jay F, Gaggiotti OE, François O (2009b) Spatial inference of admixture proportions

and secondary contact zones. *Molecular biology and evolution*, **26**, 1963–73.

Dyer RJ (2009) GeneticStudio: a suite of programs for spatial analysis of genetic-marker data.

Molecular ecology resources, **9**, 110–3.

Dyer RJ (2015a) Is there such a thing as landscape genetics? *Molecular Ecology*, **10**, 139–143.

Dyer RJ (2015b) Population Graphs and Landscape Genetics. *Annual Review of Ecology,*

Evolution, and Systematics, **46**, annurev–ecolsys–112414–054150.

Dyer RJ, Nason JD (2004) Population Graphs: the graph theoretic shape of genetic structure.

Molecular ecology, **13**, 1713–27.

Dyer RJ, Nason JD, Garrick RC (2010) Landscape modelling of gene flow: improved power

using conditional genetic distance derived from the topology of population networks.

Molecular ecology, **19**, 3746–59.

Economic Research Service U (1998) Seed Industry Structure Is Characterized by Growth and

Consolidation. *AIB*, **786**, 25–29.

Edelaar P, Bolnick DI (2012) Non-random gene flow: An underappreciated force in evolution

and ecology. *Trends in Ecology and Evolution*, **27**, 659–665.

Ennos RA (1981) Quantitative studies of the mating system in two sympatric species

ofIpomoea(Convolvulaceae). *Genetica*, **57**, 93–98.

Ennos R a (1994) Estimating the relative rates of pollen and seed migration among plant

populations. *Heredity*, **72**, 250–259.

Epps CW, Keyghobadi N (2015) Landscape genetics in a changing world: disentangling

historical and contemporary influences and inferring change. *Molecular Ecology*, **24**, 6021–

6040.

Everman E, Klawinski P (2013) Human-facilitated jump dispersal of a non-native frog species on

Hawai'i Island. *Journal of Biogeography*, **40**, 1961–1970.

Excoffier L, Smouse PE, Quattro JM (1992) Analysis of Molecular Variance Inferred From

Metric Distances Among DNA Haplotypes: Application. , **491**, 479–491.

Fang Z, Gonzales AM, Durbin ML *et al.* (2013) Tracing the geographic origins of weedy

Ipomoea purpurea in the Southeastern United States. *Journal of Heredity*, **104**, 666–677.

Fountain T, Duvaux L, Horsburgh G, Reinhardt K, Butlin RK (2014) Human-facilitated

metapopulation dynamics in an emerging pest species, Cimex lectularius. *Molecular*

Ecology, **23**, 1071–1084.

Glickman ME, Rao SR, Schultz MR (2014) False discovery rate control is a recommended

alternative to Bonferroni-type adjustments in health studies. *Journal of Clinical*

Epidemiology, **67**, 850–857.

Goslee SC, Urban DL (2007) The ecodist Package for Dissimilarity-based Analysis of Ecological

Data. *Journal Of Statistical Software*, **22**, 1–19.

Guillot G, Rousset F (2013) Dismantling the Mantel tests (L Harmon, Ed,). *Methods in Ecology*

and Evolution, **4**, 336–344.

Harris SE, Xue AT, Alvarado-serrano D *et al.* (2016) Urbanization shapes the demographic

history of a native rodent (the white- footed mouse , Peromyscus leucopus) in New York

City.

He Q, Edwards DL, Knowles LL (2013) Integrative testing of how environments from the past to

the present shape genetic structure across landscapes. *Evolution*, **67**, 3386–3402.

Hoban S, Bertorelle G, Gaggiotti OE (2012) Computer simulations: tools for population and

evolutionary genetics. *Nature Reviews Genetics*.

Hodkinson DJ, Thompson K, Journal T, Dec N (2007) Plant Dispersal : The Role of Man. , **34**,

1484–1496.

Holderegger R, Buehler D, Gugerli F, Manel S (2010) Landscape genetics of plants. *Trends in*

Plant Science, **15**, 675–683.

Jha S (2015) Contemporary human-altered landscapes and oceanic barriers reduce bumble bee

gene flow. *Molecular Ecology*, **24**, 993–1006.

Jha S, Kremen C (2013) Urban land use limits regional bumble bee gene flow. *Molecular*

Ecology, **22**, 2483–2495.

Kolbe JJ, Glor RE, Rodríguez Schettino L *et al.* (2004) Genetic variation increases during

biological invasion by a Cuban lizard. *Nature*, **431**, 177–81.

Kuester A, Chang S-M, Baucom RS (2015) The geographic mosaic of herbicide resistance

evolution in the common morning glory, *Ipomoea purpurea* : Evidence for resistance

hotspots and low genetic differentiation across the landscape. *Evolutionary Applications*,

n/a–n/a.

Legendre P, Anderson MJ (1999) Distance-based redundancy analysis: Testing multispecies

responses in multifactorial ecological experiments. *Ecological Monographs*, **69**, 1–24.

Legendre P, Lapointe FJ, Casgrain P (1994) Modeling Brain Evolution from Behavior - a

Permutational Regression Approach. *Evolution*, **48**, 1487–1499.

Lehan NE, Murphy JR, Thorburn LP, Bradley BA (2013) Accidental introductions are an

important source of invasive plants in the continental United States. *American Journal of*

Botany, **100**, 1287–1293.

Lenormand T (2002) Gene flow and the limits to natural selection. *Trends in Ecology and*

Evolution, **17**, 183–189.

Lercher MJ, Hurst LD (2002) Human SNP variability and mutation rate are higher in regions of

high recombination. *Trends in Genetics*, **18**, 337–340.

Levin DA, Kerster HW (1974) Gene flow in seed plants. *Evolutionary Biology*, **7**, 139–220.

Li H (2011) A statistical framework for SNP calling, mutation discovery, association mapping

and population genetical parameter estimation from sequencing data. *Bioinformatics*, **27**,

2987–2993.

Li H, Handsaker B, Wysoker A *et al.* (2009) The Sequence Alignment/Map format and

SAMtools. *Bioinformatics*, **25**, 2078–2079.

Mack RN, Lonsdale WM (2001) Humans as global plant dispersers: Getting more than we

bargained for. *BioScience*, **51**, 95.

Magwene PM (2001) New Tools for Studying Integration and Modularity. *Evolution*, **55**, 1734–

1745.

Manel S, Holderegger R (2013) Ten years of landscape genetics. *Trends in Ecology & Evolution*,

28, 614–621.

Margosian ML, Garrett K a., Hutchinson JMS, With K a. (2009) Connectivity of the American

Agricultural Landscape: Assessing the National Risk of Crop Pest and Disease Spread.

BioScience, **59**, 141–151.

Marko PB, Hart MW (2011) The complex analytical landscape of gene flow inference. *Trends in*

Ecology and Evolution, **26**, 448–456.

Martínez-Abraín A (2008) Statistical significance and biological relevance: A call for a more

cautious interpretation of results in ecology. *Acta Oecologica*, **34**, 9–11.

Martins AC, Goncalves RB, Melo G a R (2013) Changes in wild bee fauna of a grassland in

Brazil reveal negative effects associated with growing urbanization during the last 40 years.

Zoologia, **30**, 157–176.

McRae B (2006) Isolation by resistance. *Evolution*, **60**, 1551–1561.

McRae BH, Beier P (2007) Circuit theory predicts gene flow in plant and animal populations.

Proceedings of the National Academy of Sciences of the United States of America, **104**,

19885–90.

Medley KA, Jenkins DG, Hoffman EA (2015) Human-aided and natural dispersal drive gene

flow across the range of an invasive mosquito. *Molecular Ecology*, **24**, 284–295.

Meehan TD, Werling BP, Landis D a, Gratton C (2011) Agricultural landscape simplification and

insecticide use in the Midwestern United States. *Proceedings of the National Academy of*

Sciences of the United States of America, **108**, 11500–11505.

Menchari Y, Délye C, Le Corre V (2007) Genetic variation and population structure in black-

grass (Alopecurus myosuroides Huds.), a successful, herbicide-resistant, annual grass weed

of winter cereal fields. *Molecular Ecology*, **16**, 3161–3172.

Mitchell M (1996) *An introduction to genetic algorithms*. MIT Press, Cambridge Mass.

Murphy HT, Lovett-Doust J (2004) Context and connectivity in plant metapopulations and

landscape mosaics: Does the matrix matter? *Oikos*, **105**, 3–14.

Nakagawa S (2004) A farewell to Bonferroni: The problems of low statistical power and

publication bias. *Behavioral Ecology*, **15**, 1044–1045.

Nielsen R, Slatkin M (2013) *An introduction to population genetics : theory and applications*.

Sinauer Associates, Sunderland Mass.

Nosil P, Egan SP, Funk DJ (2008) Heterogeneous genomic differentiation between walking-stick

ecotypes: "isolation by adaptation" and multiple roles for divergent selection. *Evolution*, **62**,

316–36.

Oksanen J, Blanchet FG, Kindt R *et al.* (2015) vegan: community ecology package. ,

http://CRAN.R–project.org/package=vegan.

Peakall R, Smouse PE (2012) GenAlEx 6.5: genetic analysis in Excel. Population genetic

software for teaching and research--an update. *Bioinformatics (Oxford, England)*, **28**, 2537–

9.

Perry GH, Louis EE, Ratan A *et al.* (2013) Aye-aye population genomic analyses highlight an

important center of endemism in northern Madagascar. *Proceedings of the National*

Academy of Sciences of the United States of America, **110**, 5823–5828.

Peterman WE (2014) ResistanceGA: An R package for the optimization of resistance surfaces

using genetic algortihms. *bioRxiv*.

Peterman WE, Connette GM, Semlitsch RD, Eggert LS (2014) Ecological resistance surfaces

predict fine-scale genetic differentiation in a terrestrial woodland salamander. *Molecular*

Ecology, **23**, 2402–2413.

Piganeau G, Eyre-Walker A (2009) Evidence for variation in the effective population size of

animal mitochondrial DNA. *PLoS ONE*, **4**, 2–9.

Pimentel D, Zuniga R, Morrison D (2005) Update on the environmental and economic costs

associated with alien-invasive species in the United States. *Ecological Economics*, **52**, 273–

288.

R Core Development Team (2016) R: a language and environment for statistical computing,

3.2.4. , **0**, [http://www.r–project.org].

Ramachandran S, Deshpande O, Roseman CC *et al.* (2005) Support from the relationship of

genetic and geographic distance in human populations for a serial founder effect originating

in Africa. *Proceedings of the National Academy of Sciences of the United States of America*,

102, 15942–7.

Raufaste N, Rousset F (2001) Are Partial Mantel Tests Adequate? *Evolution*, **55**, 1703–1705.

Ricciardi A (2007) Are modern biological invasions an unprecedented form of global change?

Conservation Biology, **21**, 329–336.

Rogers JOYB, Murray DONS, Verhalen LM, Claypool PL (1996) Ivyleaf Morningglory (

Ipomoea hederacea) Interference with Cotton (Gossypium hirsutum)'. , **10**, 107–114.

Rowe HC, Renaut S, Guggisberg a (2011) RAD in the realm of next-generation sequencing

technologies. *Molecular ecology*, **20**, 3499–502.

Sexton JP, Hangartner SB, Hoffmann A a (2014) Genetic isolation by environment or distance:

854  which pattern of gene flow is most common? *Evolution; international journal of organic*

855  *evolution*, **68**, 1–15.

856 Shah VB, McRae BH (2008) Circuitscape : A Tool for Landscape Ecology. *Proceedings of the*

857  *7th Python in Science Conference*, 62–65.

858 Simberloff D (2013) *Invasive species : what everyone needs to know*.

859 Slatkin M (1985) Gene flow in natural populations. *Annual Review of Ecology and*

860  *Systematicsystematics*, **16**, 393–430.

861 Slatkin M, Excoffier L (2012) Serial founder effects during range expansion: a spatial analog of

862  genetic drift. *Genetics*, **191**, 171–181.

863 Spear SF, Balkenhol N, Fortin MJ, McRae BH, Scribner K (2010) Use of resistance surfaces for

864  landscape genetic studies: Considerations for parameterization and analysis. *Molecular*

865  *Ecology*, **19**, 3576–3591.

866 Steiner CC, Putnam AS, Hoeck PEA, Ryder OA (2013) Conservation genomics of threatened

867  animal species. *Annual Review of Animal Biosciences*, **1**, 261–281.

868 Storz JF, Ramakrishnan U, Alberts SC (2001) Determinants of effective population size for loci

869  with different modes of inheritance. *The Journal of heredity*, **92**, 497–502.

870 Taylor PD, Fahrig L, Henein K, Merriam G (1993) Connectivity Is a Vital Element of Landscape

871  Structure. *Nordic Society Oikos* , **68**, 571–573.

872 Thill DC, Mallory-Smith CA (1997) The Nature and Consequence of Weed Spread in Cropping

873  Systems. , **45**, 337–342.

874 Tiffin P, Rausher M (1999) Genetic Constraints and Selection Acting on Tolerance to Herbivory

875  in the Common Morning Glory Ipomoea purpurea. *The American naturalist*, **154**, 700–716.

876 Trakhtenbrot A, Nathan R, Perry G, Richardson DM (2005) The importance of long-distance

dispersal in biodiversity conservation. *Diversity and Distributions*, **11**, 173–181.

Wang IJ (2010) Recognizing the temporal distinctions between landscape genetics and

phylogeography. *Molecular ecology*, **19**, 2605–8.

Wang IJ, Bradburd GS (2014) Isolation by Environment. *Molecular Ecology*, n/a–n/a.

Wang IJ, Savage WK, Bradley Shaffer H (2009) Landscape genetics and least-cost path analysis

reveal unexpected dispersal routes in the California tiger salamander (Ambystoma

californiense). *Molecular Ecology*, **18**, 1365–1374.

Webster TM, Nichols RL (2012) Changes in the Prevalence of Weed Species in the Major

Agronomic Crops of the Southern United States: 1994/1995 to 2008/2009. *Weed Science*,

60, 145–157.

White DR, Harary F (2001) The cohesiveness of blocks in social networks: node connectivity

and conditional density. *Sociological Methodology*, **31**, 305–359.

Wichmann MC, Alexander MJ, Soons MB *et al.* (2009) Human-mediated dispersal of seeds over

long distances. *Proceedings. Biological sciences / The Royal Society*, **276**, 523–532.

Wilson GA, Rannala B (2003) Bayesian inference of recent migration rates using multilocus

genotypes. *Genetics*, **163**, 1177–1191.

Wright S (1949) The genetical structure of populations. *Annals of Eugenics*, **15**, 323–354.

 FIGURES AND TABLES - Landscape connectivity of a noxious weed

898 Table 1. Analysis of Molecular Variance (AMOVA) of SSR and SNP data. The contribution of
899 spatial clusters (region), individuals.

Effect	F-statistic	Variance explained		F-value		P-value	
		SSR	SNP	SSR	SNP	SSR	SNP
Regions	F_{RT}	3.94 %	8.51%	0.039	0.085	0.001	0.001
Localities (Regions)	F_{SR}	9.05 %	6.10 %	0.094	0.067	0.001	0.001
Individuals (Localities)	F_{ST}	0.67 %	24.85 %	0.130	0.146	0.001	0.001
Individuals	F_{IS}	86.33%	60.54 %	0.008	0.291	0.252	0.001
Total	F_{IT}	100%	100 %	0.137	0.395	0.001	0.001

901
902
903

Table 2. Summary of landscape genetics models. Model coefficients are reported followed
by associated p-value (in parenthesis) and, for MLPE.lmm models, followed by AIC
difference and ranking (in square brackets). Because of multiple testing , p-values reported
for MLPE.lmm and dbRDA analyses are adjusted using a false recovery rate correction.
Significant coefficients after a false-recovery-rate correction are in bold; marginally
significant ones are denoted with an asterisk.
910

Variable	MLPE.lmm		dbRDA		partial-dbRDA		MRDM	
	SSR	SNP	SSR	SNP	SSR	SNP	SSR	SNP
Geographic distance	-	-	1.234 (0.173)	2.503 (0.202)	-	-	0.006 (0.969)	0.321 (0.542)
Environmental layers								
Elevation	**0.244** (0.028) +10.742 [3]	0.840* (0.054) +1.423 [4]	1.236 (0.104)	1.369 (0.348)	1.195 (0.206)	1.256 (0.438)	-0.629 (0.303)	-1.778 (0.685)
PC1 – climate	**0.242** (0.034) +10.356 [2]	**0.967** (0.050) +0.032 [2]	**1.454** (0.009)	1.513 (0.302)	**1.402** (0.099)	1.031 (0.438)	**1.645** (0.002)	-0.288 (0.847)
PC1 – soil	**0.208** (0.031) +11.378 [5]	**1.044** (0.050) +0.471 [3]	**1.599** (0.009)	3.847 (0.194)	1.315 (0.146)	2.644 (0.340)	-0.191 (0.651)	1.383 (0.670)
Human-impact layers								
Crops	-0.226 (0.134) +14.491 [6]	0.858* (0.054) +15.371 [5]	0.819 (0.947)	2.484 (0.202)	0.836 (0.880)	1.270 (0.438)	-0.240 (0.279)	1.414 (0.297)
Landcover	**0.582** (<0.001) 0.000 [1]	**1.358** (0.006) +39.775 [6]	1.272* (0.054)	3.392 (0.194)	1.249 (0.201)	4.423 (0.318)	-0.222 (0.407)	-0.182 (0.816)
Population density	**0.227** (0.028) +10.821 [4]	0.912* (0.053) -0000 [1]	**1.656** (0.007)	3.808 (0.194)	1.401* (0.099)	2.873 (0.318)	-0.717* (0.091)	-2.071* (0.092)

911
912

913
914 Fig. 1. Distribution of *Ipomoea purpurea*'s sampled localities. Sample sizes for both SSR
915 (black bars) and SNP (white bars) datasets are indicated (numbers on top indicate the
916 actual number of individuals used in our analyses). Elevation is provided as background.

917
918
919 Fig. 2. Admixture and spatial clusters identified in *I. purpurea*. Results for the SSR (a) and
920 SNP (b) datasets are shown together with an inset map (c) denoting the location of those
921 clusters. Individuals in the admixture plot are sorted by sample locality.

Fig. 3. Inferred population connectivity. (a,c) The estimated origin of individuals for each sampled locality (i.e., sink) is depicted according to the locality they were inferred to have originated from (source). The color of each cell in these plots depicts the proportion of individuals in the sink population that were estimated to be recent immigrants from each locality along the x-axis. Cells on the minor diagonal correspond to the proportion of native individuals. Localities are sorted by State. (b,d) Pruned conditional genetic networks. Note the great difference in complexity and interconnectedness between networks. The top row shows SSR-based results, the bottom, the SNP-based results. Locality numbers follow Fig. 2.

Fig. 4. Landscape conductivity patterns recovered. Conductance maps for variables consistently recovered as significant predictors of genetic dissimilarity are overlaid on maps of those variables to indicate the characteristics of the landscape that facilitate connectivity between sampled localities (left column). Boxplots showing the association between each variable values (binned into categories for simplicity) and conductance (center column), and regression between landscape resistance corresponding to each variable and genetic dissimilarity (right column) are also shown. The 3 associations portrayed are: (a) human population density – SNP differentiation association, (b) human population density – SSR differentiation association, and (c) PC1 climate and SSR differentiation.