Landscape connectivity of a noxious invasive weed: promises and challenges of landscape

genomics for knowledge-based weed management?

Diego F. Alvarado-Serrano (corresponding author)
Department of Ecology and Evolutionary Biology
University of Michigan
E-mail: dalvarad@umich.edu
8
9 Megan Van Etten
10 Department of Ecology and Evolutionary Biology
11 University of Michigan
12 E-mail: mvanette@umich.edu
13
14 Shu-Mei Chang
15 Department of Ecology and Evolutionary Biology
16 University of Michigan
17 E-mail: smchang@uga.edu
18
Regina S. Baucom
Department of Ecology and Evolutionary Biology
University of Michigan
E-mail: rsbaucom@umich.edu

Landscape connectivity of a noxious invasive weed: promises and challenges of landscape

genomics for knowledge-based weed management?

ABSTRACT

Examining how the landscape may influence gene flow is at the forefront of

understanding population differentiation and adaptation. Such understanding is crucial in light of

ongoing environmental changes and the elevated risk of ecosystems alteration. In particular,

knowledge of how humans may influence the structure of populations is imperative to allow for

informed decisions in management and conservation. Here we characterize the population genetic

structure of *Ipomoea purpurea*, a noxious invasive weed, and assess the interaction between

natural and human-driven landscapes on genetic differentiation. By combining rigorous statistical

analyses and different molecular markers (nuclear microsatellites and a genome-wide panel of

SNPs), we detect both common and marker-specific patterns of genetic connectivity and identify

human population density as an important predictor of pairwise population differentiation,

suggesting that the agricultural and/or horticultural trade may be involved in maintaining some

level of connectivity across distant agricultural fields. Climatic variation appears as an additional

predictor. We discuss the implications of these results and the approach we followed in the

context of understanding agricultural weed and invasive species' connectivity, as well as the

challenges and promises of current landscape genomics research for knowledge-based weed

management.

Keywords: agricultural weeds, human-aided migration, landscape genetics, morning glory,

population structure

INTRODUCTION

Elucidating routes and levels of migration between subpopulations of a species is essential to understand the interplay between gene flow, adaptation, genetic drift, and selection, and hence the forces that shape its evolutionary trajectory (Barrowclough, 1980; Slatkin, 1985). Landscape features—such as rivers, mountain ranges, crop fields, and urban areas—can impact levels of gene flow between populations by determining dispersal rates and routes (Cushman, McKelvey, Hayden, & Schwartz, 2006; McRae, 2006) as well as by influencing the likelihood of successful establishment of immigrants (Nosil, Egan, & Funk, 2008; Sexton, Hangartner, & Hoffmann, 2014; Wang & Bradburd, 2014). Landscape features can also indirectly condition the effect of gene flow through its effect on local effective population sizes since the actual role that migration plays in the evolution of a species is driven by the fraction of the local population size that correspond to immigrants (Slatkin, 1985; Wright, 1949). Consequently, the landscape, loosely defined as an area with spatially variable biotic and abiotic factors (Holderegger, Buehler, Gugerli, & Manel, 2010), creates the stage for spatially variable levels of effective gene flow among populations, conditioned by species' specific physiological tolerances and behavioral preferences (Clobert, Baguette, Benton, Bullock, & Ducatez, 2012). In this way, the landscape plays a pivotal role in the evolution of species.

In contrast to species that depend almost exclusively on natural dispersal agents, species in heavily human-dominated ecosystems may exploit human activities to maintain gene flow among populations and expand their ranges (Everman & Klawinski, 2013; Fountain, Duvaux, Horsburgh, Reinhardt, & Butlin, 2014). Such species may be capable of maintaining population connectivity over vast geographic ranges (Trakhtenbrot, Nathan, Perry, & Richardson, 2005) by overcoming landscape features that would otherwise represent natural barriers and reach dispersal

distances that could be orders of magnitude greater than those attained under natural agents or do

it under much smaller time frames (Mack & Lonsdale, 2001; Ricciardi, 2007). In this way, by

facilitating dispersal humans have the potential to: i) condition the balance between drift and

selection (Lenormand, 2002; Slatkin, 1985), ii) introduce relevant genetic variation to local

populations (Kolbe et al., 2004), iii) prevent local extinction or favor recolonization (Fountain et

al., 2014), and alter the overall genetic constitution of populations (Bataille, Cunningham, Cruz,

Cedeño, & Goodman, 2011). Human-aided migration—intentional or unintentional—is

particularly prevalent in plants (Auffret & Cousins, 2013; Hodkinson, Thompson, Journal, &

Dec, 2007; Wichmann et al., 2009), where it has had major impacts on the distribution of species

and stability of communities (Simberloff, 2013 and references therein). Yet, the open question

remains: how might the interaction between human-dominated and more natural landscapes

affect population connectivity in plant populations, especially in human-exploiter species?

A particularly amenable system to study the interaction between natural and human-aided

dispersal comes from agricultural weed populations. Agricultural weeds face a highly dynamic

landscape characterized by frequent spatial rearrangements (expansion of agricultural front,

increased fragmentation) and a constantly changing environment (crop rotation, agricultural

chemical use, climatic abnormalities) (Meehan, Werling, Landis, & Gratton, 2011; Menchari,

Délye, & Le Corre, 2007) that certainly impact their opportunities for survival and local

adaptation through its effect on population connectivity (Margosian, Garrett, Hutchinson, &

With, 2009). At the same time, natural features such as climate, soil type, and topography are

expected to play a significant role in structuring populations provided intrinsic physiological

requirements and species-specific traits (Cimalová & Lososová, 2009; Navas, 2012). Under these

conditions, human-aided migration is expected to be critical for weeds' success (Epperson &

Clegg, 1986), but knowledge on how or if weedy plant populations are able to maintain

connectivity through a complex landscape matrix of croplands, grasslands, natural and urban

areas is limited. Addressing this limitation should not only improve our understanding of the

underlying processes governing weeds' population structure, but should also offer practical tools

to deal with this ever-growing agricultural problem that impose severe economic costs (on the

order of 33B USD per year in US agriculture alone; Pimentel, Zuniga, & Morrison, 2005).

As a first step into investigating the interplay between natural factors and human activities

on structuring genetic diversity in weed populations, we estimate the intensity and extent of

migration from genetic data and, under the preliminary simplifying assumption of evolutionary

equilibrium (Marko & Hart, 2011), evaluate how multiple landscape features influence genetic

connectivity of a noxious agricultural weed, *Ipomoea purpurea*, using two different sets of

molecular markers (nuclear microsatellites and a genome-wide panel of SNPs). Specifically, we

ask the following questions: 1) Which natural and/or human-influenced landscape features—

soils, elevation, climate, landcover, crop types, human population density—promote or constrain

genetic connectivity between populations of this agricultural weed? and 2) what additional

insights can we gain from a broader representation of the genome than traditionally used in

landscape genetics studies (typically microsatellites and organelle DNA)? By considering the

possible interactions between natural and human effects on migration, the answers to these

questions offer deeper knowledge of the interaction between human activities, landscape features,

and population structure of noxious weeds and hence contribute to improve effective

management and control of these damaging plants. More generally, these answers contribute to

deepen our understanding of the interaction between environmental setting and population

differentiation, adaptation, and persistence (Taylor, Fahrig, Henein, & Merriam, 1993).

MATERIALS AND METHODS

Study system

Ipomoea purpurea, the common morning glory, is an agricultural weed evolving under

the influence of human-driven and natural landscape factors. This species is a noxious weed, of

horticultural value (Defelice, 2001; Fang et al., 2013), with a widespread distribution that

includes highly heterogeneous landscapes in the Eastern, South- and Mid-western regions of the

United States (Culpepper, 2006; Webster & Nichols, 2012). It is a self-compatible annual

bumblebee-pollinated vine, with heavy seeds, and is found primarily in agricultural fields and

disturbed areas (Baucom & Mauricio, 2008; Tiffin & Rausher, 1999), as well as cultivated flower

gardens and yards (Defelice, 2001). While most details on the history of I. purpurea remain

unknown, it is hypothesized that I. purpurea originated in Central America, from where it was

taken to Spain to be grown in monasteries as an ornamental during the XVI century (Defelice,

2001; Fang et al., 2013). From there, it is hypothesized that its cultivation expanded to other

European countries, including England, and later to North America (Defelice, 2001; Fang et al.,

2013). By the early XVIII century it became a popular plant in gardens in the United States, but

also a known weed (Defelice, 2001). Since then, little is know about the demographic history of

this species in the United States, other than the fact that gene flow has probably been maintained

over time at least among populations in relatively close geographic proximity (Kuester, Chang, &

Baucom, 2015). What is clear is that since its introduction its history has been tightly linked to

human activities, making it suitable to assess the impact of natural and anthropogenic landscapes

on structuring genetic variance.

Ipomoea purpurea is currently one of the most problematic agricultural weeds (Webster

& Nichols, 2012) and is capable of infestations leading to substantial decline in crop (closely

related *Ipomoea* species cause declines of up to 80% of crop yield; Rogers, Murray, Verhalen, &

Claypool, 1996). This species exhibits resistance to the commonly used herbicide glyphosate

(Baucom & Mauricio, 2004, 2008), although the exact resistance level varies widely among

populations of this species (Kuester et al., 2015). This species is also a major concern for

conservation given its naturalization in multiple regions throughout the world and its

aggressiveness as an invasive (Chaney & Baucom, 2012; Fang et al., 2013). Hence, unraveling

the population structure of this species and how it is affected by the landscape should not only

improve our understanding of basic evolutionary processes, but should also inform practical

decisions for its management and control (e.g., Is herbicide resistance better controlled by

avoiding the spread of resistance genotypes or by local management of moderately isolated

populations?).

Data compilation

To capture the plausible effect of both natural and disturbed landscapes on structuring genetic

diversity in *I. purpurea*, we compiled a diverse set of GIS data for the continental US from a

variety of sources (Table S1). These data encapsulate human activities (human population

density, landcover, planted crops, and roads) as well as the geographical setting of *I. purpurea*

(elevation, climate—19 variables summarizing central tendencies and variability patterns in

temperature and precipitation, soil—8 variables summarizing the texture, pH, and organic and

inorganic content of the top 20cm of soil). We focused on both sets of data because of the

possible interaction between natural and human effects, which may lead to incorrect inferences if

not accounted for (e.g., spurious associations due to spatial correlation between crops distribution

and climate; Eberhardt & Teal, 2013). We first processed all these data into landscape layers at a

common spatial resolution of $10km^2$ and a common spatial extent around the US states with

available samples (Fig. 1). These spatial resolution and extent were chosen to maintain a practical

balance between scale and analytical manageability given available computational resources. To

reduce dimensionality, we opted to perform two separate Principal Component Analyses (PCAs)

on the 19 climatic and 8 soil layers, respectively. For all subsequent analyses we kept the

resulting first two principal components of each of these analyses, which accounted for over 78%

of the variance in each case, and primarily summarized temperature temporal gradients and

precipitation seasonality, and soils' pH, sandiness, and grain size, respectively (Table S2).

With the objective of estimating the genetic connectivity of populations of *I. purpurea*, we

compiled genetic data on an extensive panel of 15 previously optimized microsatellite loci

(Molecular Ecology Resources Primer Development Consortium 2013), which quality has been

verified by looking for scoring errors (Kuester et al., 2015) using Micro-Checker (Van

Oosterhout, Hutchinson, Wills, & Shipley, 2004). These data encompass a total of 597

individuals from 31 localities (with a minimum of 8 individuals per locality) (Fig. 1; Table S3).

All individuals were collected in 2012 from farms across the range of *I. purpurea* in the United

Sates (Kuester et al., 2015). In addition, to obtain a more comprehensive representation of the

genome of *I. purpurea* and assess the robustness of results in light of coalescent and mutational

variance (Nielsen & Slatkin, 2013), we generated a Next Generation Sequencing (NGS) dataset

from an additional set of individuals (from 6 localities represented in the SSR dataset, plus 2

additional localities in close geographic proximity to localities in the SSR dataset; Fig. 1).

To generate the NGS dataset, we constructed genome-wide Genotype By Sequencing (GBS)

libraries for 80 individuals sampled across the 8 localities. The GBS library was developed using

7ng of genomic DNA, extracted from leaf or cotyledon tissue, using SNPsaurus' (Oregon, USA)

nextRAD technology. This technology uses a selective PCR primer to amplify consistent

genomic loci among individuals. Similarly to RAD-Seq sequences (Rowe, Renaut, &

Guggisberg, 2011) in which the DNA flanking a restriction enzyme cut site is selected for

amplification, nextRAD amplifies sequences that correspond to the DNA downstream of a short

selective priming site. Samples were first fragmented and then ligated to short adapter and

barcode sequences using a partial Nextera reaction (Illumina; California, USA) before being

amplified using Phusion® Hot Start Flex DNA Polymerase (New England Biolabs;

Massachusetts, USA). The 80 dual-barcoded PCR-amplified samples were pooled and the

resulting libraries were purified using AMPure XP beads (Agencourt Bioscience Corporation;

Massachusetts, USA) at 0.7x. The purified library was then size selected to 350-800 base pairs

and sequenced using two runs of an Illumina NextSeq500 sequencer (Genomics Core Facility,

University of Oregon).

The resulting sequences were analytically processed using the SNPsaurus nextRAD pipeline

(SNPsaurus, Oregon, USA; Siliceo-Cantero, García, Reynolds, Pacheco, & Lister, 2016).

Specifically, reads of 16 randomly selected individuals (of the 80 sequenced) were combined to

create a pseudo-reference genome. This was done after removing loci with read counts above

20,000, which presumably corresponded to repetitive genomic material, and loci with read counts

below 100, which presumably corresponded to off-target or read errors. The filtered reads were

aligned to each other using BBMap (Bushnell, 2016). All parameters were set to default values

with the exception of minimum alignment identity, which was set to 0.93 to identify alleles, as it

is a threshold found to work well for non-reference species (SNPsaurus, Oregon, USA). A single

read instance was chosen to represent the locus in the pseudo-reference. This resulted in a total of

263,658 loci. All reads from each of the 80 individuals were then aligned to the pseudo-reference

using BBMap (Bushnell, 2016) and converted to a vcf genotype table, using Samtools (Li et al.,

2009) and bcftools (Li, 2011), after filtering out nucleotides with a quality score of 10 or worse

(an empirically informed threshold; SNPsaurus, Oregon, USA). The resulting vcf table was

filtered using vcftools (Danecek et al., 2011) for SNPs with a minimum allele frequency of 0.02,

a minimum read depth of 5, and a maximum 15% of missing data. We chose this filtering scheme

as a balance between accuracy and efficiency and to avoid inadvertent errors associated with our

use of a pseudo-reference genome. This resulted in 9774 variable regions. Loci were further

filtered using vcftools to exclude loci with less than 5 high quality base-calls and with more than

20% missing data or an average of less than 20 high quality base calls. This resulted in a final

panel of 8210 Single Nucleotide Polymorphisms (SNPs) that we used in all subsequent analyses.

Population structure analyses

We first conducted a series of preliminary analyses to characterize the overall genetic

structure of *I. purpurea*. All analyses were run separately for the microsatellite (SSR, hereafter)

and SNP datasets given their intrinsic differences and distinct geographic coverage (Fig. 1; Table

S3). In addition, we repeated all population structure analyses, separate for the SSR and SNP

datasets, for the subset of 6 localities with coincident data for both markers (referred as SSRc and

SNPc, hereafter) to assess the robustness of results to the difference in geographic coverage.

First, we examined population differentiation by estimating F_{ST} using GenAlex v6.5 (Peakall

& Smouse, 2012) (because similar global F_{ST} and R_{ST} estimates were obtained for the SSR

dataset, we opted to report F_{ST} values only to allow direct comparisons with the SNP dataset). We

then estimated contemporary effective population size for each sampled locality in NeEstimator

v2 using the excess heterozygous method (Do et al., 2014). We performed this latter analysis to

assess the possibility of whether differences in local population size underlie differences in

genetic variation (Weckworth et al., 2013) and/or promote asymmetric effective migration rate

(*Nm*).

In addition, we assessed population admixture and spatial genetic clustering using TESS

(Chen, Durand, Forbes, & François, 2007). TESS was run using the admixture algorithm and a

BYM model (Durand, Jay, Gaggiotti, & François, 2009) with 10 runs per K value, and without

using geographic weights. The TESS model, with the lowest DIC was chosen as the optimal

model (Durand, Chen, & Francois, 2009). K values tested ranged from two to the maximum

number of sampled localities. Additionally, following Wang et al. (2009), we complemented

these analyses with Analyses of Molecular Variance (AMOVA; Excoffier, Smouse, & Quattro,

1992) run in GenAlex (Peakall & Smouse, 2012) using 9999 permutation replicates. We run

these AMOVAs either partitioning the variance into regions based on the spatial genetic clusters

previously identified by TESS—to quantify the fraction of the genetic variance explained by

these clusters, or leaving it ungrouped (i.e., no regions), for comparison.

Additionally, we investigated population connectivity by estimating levels of recent

migration between sampled localities through the identification of individuals of mixed ancestry

using BayesAss (Wilson & Rannala, 2003). BayesAss is a program that uses individual

multilocus genotypes and a Markov Chain Monte Carlo (MCMC) algorithm to probabilistically

distinguish between immigrants and long-term native individuals (Wilson & Rannala, 2003). We

ran BayesAss for 6 million generations using default parameter settings, and discarded the first

two million generations as burn-in (Dyer, 2009). For each marker dataset, we repeated this

analysis three times (for a total of 18 million generations) and combined the results from the three

replicates for our final inference. Then, using a posterior probability cut-off of 0.75 we assign

individuals' ancestry. We chose this cut-off value as a minimum credibility score to

simultaneously maximize sample size and reliability (stringer thresholds show similar differences

between marker sets; results not shown). It is important to note that because of computational

limits we had to randomly subsample our set of SNPs to 400 SNPs for this analysis. The same

subsampled set was used for the full and reduced (i.e., on the SSRc and SNPc datasets) analyses.

Landscape genetics analyses

After assessing overall population structure of *I. purpurea*, we evaluated the association

between landscape features and genetic differentiation based on the full datasets. We limited our

analyses to the full datasets because of the robust genetic structure recovered between the full and

reduced datasets (see below) and the smaller sample size of the latter datasets, which limits

statistical inference power. First, we estimated conditional genetic distances (Dyer, Nason, &

Garrick, 2010) using GeneticStudio (Dyer, 2009). Briefly, conditional distances are measures of

pairwise genetic distance derived from population networks, constructed based on the degree of

genetic similarity between sampled localities (Dyer & Nason, 2004). Because these networks are

pruned based on the principle of conditional independence of the total among population genetic

covariance (using an edge deviance principle; Magwene, 2001), conditional distances reflect

genetic similarity between localities that better capture direct gene flow as opposed to

connectivity driven by intervening localities (Dyer, 2015b). The complexity of the associated

conditional genetic network was summarized by their vertex connectivity (White & Harary,

2001), whereas the congruence between networks derived from different marker sets was

measured by their structural congruence (a measure of wether the number of congruent edges

between networks is greater than expected by chance) (Dyer, 2009).

Climate, crops, elevation, landcover, population density, roads, and soils landscape layers

(Table S1) were converted into landscape resistance layers by assigning a resistance value to each

landscape feature in these layers to reflect the difficulty that each feature offers to the movement

of gametes or individuals. It is important to note that in contrast to previous studies that typically

rely on expert opinion for resistance assignment, we utilized an unbiased statistical optimization

to avoid the sensitivity of results to subjective resistance assignment (Spear, Balkenhol, Fortin,

McRae, & Scribner, 2010). Specifically, resistance values were optimized through a genetic

algorithm approach (Mitchell, 1996). Briefly, in this search algorithm a population of individuals

with traits encoded by unique combinations of model parameters (resistance assignment

proposals in our case) is allowed to compete with each other based on the fitness associated with

the traits it carries (Peterman, Connette, Semlitsch, & Eggert, 2014). Specifically, in Peterman's

(2014) implementation of this algorithm, which we followed here, individuals' fitness is

estimated by the relative quality of a MLPE.lmm model (Maximum Likelihood Population

Effects – Linear Mixture Model) that evaluates the association between pairwise genetic distance

and landscape cumulative resistance between localities, estimated in Circuitscape (Shah &

McRae, 2008). Individuals with parameter settings (i.e., resistance assignments) that result in

better models, as measured by a Deviance Information Criterion (DIC) score, are preferentially

represented in the following generation. Offspring modifications introduced by mutations (i.e.,

small resistance assignment perturbations) allow for exploration of the parameter space. The

algorithm was stopped once 25 generations have passed without significant improvement in

fitness.

We implemented Peterman's (2014) algorithm in R (package ResistanceGA; Peterman, 2014)

allowing for the independent optimization of each of our landscape layers. The optimal resistance

landscapes identified in this way were then used to run a final univariate MLPE.lmm model to

characterize the association between landscape features and conditional genetic distances

between localities. Because the roads-association resistance was not recovered as significant for

either marker dataset, we dropped this layer for all subsequent analyses. Finally, to identify the

simultaneous contribution of natural and human-driven landscape features we ran Multiple

Regression on Distance Matrices (MRDM; Legendre, Lapointe, & Casgrain, 1994), which has

been identified as one of the best performing methods for evaluating the interplay between

landscape features and genetic connectivity (Balkenhol, Waits, & Dezzani, 2009). Before running

these MRDM analyses, we standardized all optimized resistance layers to mean of zero and

variance of one (Dyer et al., 2010). These final regressions included geographic distance as a null

model predictor as well as effective population size and were run in R (package ecodist; Goslee

& Urban, 2007) using 10,000 permutations to assess significance. In none of our analyses did we

implement a Bonferroni correction for multiple testing because of the overly conservative nature

of this correction (Glickman, Rao, & Schultz, 2014; Nakagawa, 2004). Instead we applied a false

recovery rate correction (Benjamini & Hochberg, 1995) using the function *p.adjust* in R (R Core

Development Team, 2016).

RESULTS

Population structure

The set of preliminary genetic analyses indicated that *I. purpurea* sampled localities were

in no major violation of Hardy-Weinberg equilibrium, as judged by the small difference between

expected and observed heterozygosity (mean He = 0.294±0.014 and 0.250±0.001; mean Ho =

0.291±0.009 and 0.260±0.001, respectively for SSR and SNP datasets). Levels of expected and

observed heterozygosity for the SSR dataset were only slightly greater than those estimated for

the SNP dataset. Likewise, the estimated mean effective population size per sampled locality was

only slightly greater and more variable for the SSR dataset than for the SNP dataset (13.71±5.59,

9.49±0.13, respectively), but in neither case was there salient evidence of a plausible source-sink

dynamic, as judged by the similar effective sizes among populations. Neither were there salient

differences in F_{ST} estimates between datasets (0.151 and 0.140, respectively for SSR and SNP

datasets; Fig. S1), with F_{ST} estimates being within the range of F_{ST} values of other broadly

distributed agricultural weeds [F_{ST}: 0.14−0.38] (Bussell, 1999; Eschmann-Grupe, Neuffer, &

Hurka, 2004; Müller-Schärer & Fischer, 2001). Congruently, no major differences in genetic

estimates between the SSRc and SNPc estimates were found (Table S4). Further confirming the

limited spatial structure in this species, spatial genetic clusters identified by the best TESS model

(Fig. S2) explained less than 13% of the variance across datasets, and barely reduced the variance

explained solely by geographic location when compared to a null model with no regions assigned

(Tables 1, S5).

Despite these similarities between the SSR and SNP datasets, the underlying genetic

structure was markedly different. Estimates of recent ancestry differed between SSR and SNP

datasets. The analysis on the SSR dataset indicated that migration among localities is more

widespread and hardly geographically constrained, with only four localities being primarily

constituted of native individuals (Fig. 2a). Across localities, on average 73.65% of individuals

were inferred to be 1st or 2nd generation immigrants (it is important to note, however, that such

high migration rate surpasses the assumptions of the method, and hence they should be taken

cautiously). On the other hand, the analysis of the SNP dataset showed that most populations

have a much more limited number of recent immigrants, and that the relatively few inferred

immigrants (on average 27.42% of individuals) did not come exclusively from geographically

proximate localities (Fig. 2d). Accordingly, SSR and SNP pruned conditional genetic networks

(Dyer & Nason, 2004) indicated remarkably different underlying patterns of genetic connectivity

(structural congruence = 0.108; Fig. 2b,e). The SSR-based network was more interconnected

(vertex connectivity: 5) than the SNP-based network (vertex connectivity: 0). Furthermore,

strong admixture was recovered in the SSR dataset, whereas minimal admixture was identified in

the SNP dataset (Fig. 2c,f). As before, these differences were consistent when analyzing the

SSRc and SNPc datasets. The SNPc dataset was characterized by a smaller percentage of recent

immigrants (28.25%) than the SSRc dataset (44.93%) (Fig. S3a,d), and the corresponding genetic

networks were also clearly different from each other (structural congruence = 0.002)—with the

SSR-based network being more connected (and vertex connectivity = 2) than the SNPc-base

network (i.e., vertex connectivity = 0)—(Fig. S3b,e). Finally, as for the full data, a more admixed

genetic composition of individuals was recovered in the SSRc dataset than in the SNPc dataset

(Fig. S3c,f).

Landscape genetics

Unsurprisingly, given the distinct underlying genetic patterns between SSR and SNP

datasets (see above), the optimization of landscape resistance layers resulted in different

resistance optimization solutions for each dataset (Fig. S4). It is important to note, however, that

a formal comparison is unwarranted as the associations recovered are statistical associations

driven by the fit of the resistance parameterization to the data under the statistical model

implemented (Martínez-Abraín, 2008). While these associations are expected to recapitulate real

biological properties of the study system, they are constrained to the data at hand. Nonetheless,

association patterns that are robust to the data used are expected to better reflect the actual impact

that landscape features have on gene flow, independently of possible biases introduced by expert

opinion. Therefore, we focus below on the common biological findings between marker types,

while also denoting the most relevant differences. Such differences likely reflect not only the

different environmental ranges covered by each dataset (Fig. 1), but most importantly, the

particular population genetic structure underlying each dataset (Fig. 2).

In spite of the distinct underlying genetic structure, there were some landscape features

that showed consistent inferred conductance to migration between datasets (Fig. S5). For

example, a tendency towards greater landscape conductivity in relatively warm and precipitation-

seasonal areas was observed in both datasets along with a steep decline in connectivity towards

areas with the greatest temperatures and precipitation-seasonality values in the study area (Fig.

S5). Likewise, cotton-dominated areas were recovered as substantially more permeable landscape

features than areas dominated by soybean fields, evergreen forests, open shrublands, and

grasslands (Fig. S5). Furthermore, both datasets pointed towards human-impacted landscapes

playing an important role in shaping genetic connectivity in this species. While in both sets of

MLPE.lmm models, null (geographic distance), natural (climate, elevation, and soils), and

human-related landscapes (landcover and human population density) were identified as

significant (hereafter, $0.01 < p \leq 0.05$ after correction for multiple testing) or marginally

significant predictors (hereafter, $p \leq 0.01$ after correction for multiple testing) of genetic

similarity between localities, the variable with the greatest association coefficient and lowest

AICc value in these models was in both cases a variable closely linked to human presence

(landcover in the SSR dataset, and human population density in the SNP dataset; Table 2) (for

comparison, Table S6 shows comparable analyses based on distance-based Redundancy

Analysis—another commonly used algorithm in landscape genetics). However, when considering

all variables together in a multivariate manner while accounting for geographic distance, human

population density, local effective population size, and different aspects of climate were the only

variables that remained as significant or marginally significant predictors of genetic

differentiation across both SSR and SNP datasets (Table 2). Elevation and soil were identified as

significant or marginally significant predictors only in the SNP dataset.

In summary, across datasets results indicated that human-population-density resistance

was robustly associated with population differentiation, with highly populated areas identified as

less conducive areas for migration (although the exact association varied between datasets; Fig.

S5). Local effective population size was also a significant predictor when considering all other

variables. It is important to note, however, that these multivariate regressions explained a variable

proportion of the variance (MRDM R^2 for SSR and SNP dataset were 0.109 ($F_{1,29} = 3.654$, p-val.

= 0.063) and 0.532 ($F_{1,6} = 1.932$, p-val. = 0.113), respectively). In addition to population density

and effective population size, temperature was also recovered as a significant predictor of genetic

dissimilarity across most analyses for the SSR dataset but not for the SNP dataset (Table 2).

DISCUSSION

The results suggest that broadly distributed populations of *I. purpurea* are partially

genetically distinct (more so when analyzing the SNP dataset), although there is some indication

of long-distance and putatively human-mediated migration between localities—as suggested by

the recovered association between human population density and genetic similarity. The levels of

differentiation observed and inferred long-distance migration strongly contrast with this species'

patchy distribution, which is tightly linked to isolated agricultural patches that are surrounded by

a complex matrix of natural and urbanized areas. Contrary to what has been seen in other

agricultural weeds (Menchari et al., 2007; Ye, Mu, Cao, & Ge, 2004), populations of *I. purpurea*

in relatively close geographic proximity do not form clusters of genetically similar individuals.

This finding, supported by our spatial population structure and landscape genetics analyses,

suggests that the local agricultural matrix does not seem to have an overarching impact on the

connectivity in this species at the landscape level—albeit it likely influences connectivity at the

small spatial scales. Instead, climate and human population density were robustly recovered as

predictors of genetic connectivity in this species across datasets and analyses. Of these landscape

variables, climate has a stronger effect, as judged by its greater MRDM coefficient. Of note,

temperature (summarized by climate PC1) was recovered as marginally significant only when

considering the SSR dataset, which is the only dataset that covers the northern portion of the

range, whereas precipitation seasonality (summarized by climate PC2) was recovered as

marginally significant only in the SNP dataset. Otherwise, population density was the only

variable across datasets with a marginally significant effect—even after accounting for multiple

tests. In addition, local effective population size was found to be a significant predictor only after

accounting for all landscape variables, suggesting a plausible superseding effect of genetic drift

driving differentiation across localities (Weckworth et al., 2013). Taken together, these results

highlight the significant interplay between human-driven and natural landscapes in structuring

populations of this species. The role that humans play in this system is likely mediated by their

impact on migration patterns themselves as well as by the reduction of population size of this

weed through pest control (Barker, Thompson, & Godley, 1984; Baucom & Mauricio, 2008).

The results also highlight how inferences about population structure and patterns of

connectivity may be dataset-dependent, with marked differences becoming apparent only after

careful dissection of roughly similar F_{ST} and heterozygosity estimates across molecular markers.

Such differences cannot be attributed to the more widespread samples of our SSR dataset, as our

findings were robust to subsampling this dataset to match the available SNP samples (see

Supporting Information). This represents a rather unexpected finding as both the overall

population structure and the influence of landscape features on population connectivity should be

inherent species properties and no marker specific realizations of common underlying biological

processes (but see below). Next, we detail each of these novel findings and place them in the

context of agricultural weed movement across the landscape, invasive species, and landscape

genetics practice.

Human impact

Given that *I. purpurea* is a naturalized species in the United States that is found primarily

associated with cultivated crops and horticultural gardens (Baucom & Mauricio, 2004; Defelice,

2001; Fang et al., 2013), the finding that human population density is a predictor of genetic

similarity in this species is at first glance unsurprising. Yet, because habitat requirements for

establishment and migration are not always the same, especially for organisms with distinct

migration stages (e.g., pollen or seeds in plants) and dormant stages (Murphy & Lovett-Doust,

2004), this finding is not as straightforward as it seems. In particular, the fact that human

population density is recovered as an informative predictor throughout the entire sampled

distribution—even after accounting for climate and landcover variation, highlights the direct

influence that humans most likely have on structuring the populations of this species and helps to

discern the factors involved in the spread of this noxious weed. In this sense, the results point

towards humans not only as likely responsible for the introduction of this weed into the United

States (Fang et al., 2013), but also as likely responsible for facilitating its current spatial

connectivity and genetic structure, and hence its opportunities for thriving in the complex

landscapes it inhabits. While further testing is required to formally test this hypothesis, especially

considering the limitations of current landscape genetic approaches (see below), our findings

suggest a multifaceted effect of human activities.

For one, it is theoretically possible that human population density primarily facilitates

connectivity at local to intermediate scales, which encompasses agricultural fields in relatively

close geographic proximity, suggesting that factors such as regional sharing of contaminated

agricultural machines, regional trade between farmers, or regional distribution of contaminated

crop seeds were at play (Benvenuti, 2007; Boyd & White, 2009; Dastgheib, 1989; Thill &

Mallory-Smith, 1997). Yet, the limited spatial clustering at the regional scale and the lack of a

significant effect of roads (results not shown) make this possibility unlikely. Instead, considering

that i) the horticultural trade has been recognized as the main source of invasive introductions and

spread in the United States (Lehan, Murphy, Thorburn, & Bradley, 2013), ii) that *I. purpurea* is

an appreciated horticultural species (Fang et al., 2013), and that, given current agricultural

practices, crop seed contamination is unlikely to be a major factor (Economic Research Service,

1998), it is probable that ornamentals' trade between population centers may help explain both

the long distance dispersal events recovered in both datasets and the overall population structure.

Alternatively, the impact of human populations on the distribution and abundance of bumblebees

(Jha, 2015; Martins, Goncalves, & Melo, 2013), which are *I. purpurea*'s predominant pollinators

(Baucom & Mauricio, 2008; Ennos, 1981), could also be partially responsible for the landscape

genetic patterns recovered as changes in the pollinators community would have strong effects on

gene flow (Jha & Kremen, 2013).

In reality a combination of all these factors may be involved. While further analyses are

needed to elucidate the ultimate causes behind the recovered association between human

population density and genetic dissimilarity in *I. purpurea*, our findings bring much needed

insight to limit the spread of this noxious weed. Our findings are not only relevant to *I. purpurea*

and to the evolution of herbicide resistance in this species (i.e., is herbicide resistance evolving

independently across populations or is it being disseminated through human-aided migration?),

but also have important implications for other weeds of agricultural concern as well as other

human-exploiter species (Blair, 2001), such as other invasives. Specifically, in line with previous

work (Auffret, Berg, & Cousins, 2014; Banks, Paini, Bayliss, & Hodda, 2015; Bataille et al.,

2011), the results here point towards the need of better strategies to minimize the impact that

humans have on the spread of species. In particular, our results further support that humans may

not only facilitate the introduction of invasive species into non-colonized areas, but also

contribute to the maintenance of gene flow among naturalized populations (Medley, Jenkins, &

Hoffman, 2015), which may be critical in providing relevant genetic variants to respond to novel

selective regimes as well as prevent inbreeding depression in these newly colonized areas

(Edelaar & Bolnick, 2012; Kolbe et al., 2004).

SSR- vs. SNP-based inferences

The unique patterns observed for each marker offer the opportunity to explore the underlying

causes for such differences and hence a more in-depth understanding of the plausible landscape

influences on species' genetic structure. For instance, an important consideration in any

landscape genetics study, including this one, is the spatial distribution of samples and spatial

scale of environmental data (Wang & Bradburd, 2014), as it can strongly impact the associations

recovered. It is thus theoretically possible that the particular geographic sampling of each dataset

exclusively drives the differences in genetic structure recovered by the two markers. Yet, the

robust differences that we report between the localities and regions common to both datasets

(SSRc and SNPc subsampled datasets; see Supporting Information) renders this possibility highly

unlikely and suggest that our results are at least moderately robust to sample distribution. Still, it

is important to recognize that all our datasets contain sets of spatially clustered samples, partially

in reflection of the also spatially cluster distribution of agricultural fields (Ramankutty, Evan,

Monfreda, & Foley, 2008). Hence, it is in principle possible that our inferences on all 4 datasets

might be strongly impacted by the lack of samples from intervening areas. However, our analyses

do not show the pattern of genetic separation between geographic sample clusters that is expected

under clustered sampling (Schwartz & McKelvey, 2009), suggesting that the patterns recovered

are not simply a sampling artifact.

Instead, differences between SSR- and SNP-based patterns might be related to the different

mutation rates underlying the two type of markers (Wang, 2010, but see Bohonak and Vandergast

2011). SSR mutation rates per generation per site (μ) are typically estimated to be between 10^{-3}

and 10^{-4} mutations per genome site per generation (Garza & Freimer, 1996), whereas SNP

mutation rates are typically estimated to be on average orders of magnitudes slower, around 10^{-8}-

10^{-10} (Morin, Luikart, & Wayne, 2004). All else being equal, the inferential power of population

structure is tightly linked to the number of mutations (Hubisz, Falush, Stephens, & Pritchard,

2009; Turakulov & Easteal, 2003). As a consequence, unless there is widespread homoplasy it is

expected to be more likely to recover signatures of population differentiation using the faster

evolving SSR loci. This is true even considering the total number of loci on each dataset (15 SSR

loci vs. 8210 SNP loci) (Selkoe & Toonen, 2006). Yet, our results are in contrast with this

theoretical expectation as we recovered weaker population structure using the faster evolving

SSR loci than using SNP loci. It is still possible that the greater number of expected mutations for

SSR loci, which increase the opportunities for homoplasy (Garza & Freimer, 1996), may explain

the lower degree of population differentiation in this dataset. Nonetheless, the likelihood that

widespread loci homoplasy in SSR allele size across populations has been maintained over the

temporal window since the introduction of *I. purpurea* to the US seems small. Hence, this

mutation-differential hypothesis is unlikely to be solely responsible for the differences observed.

In fact, a large proportion of genetic variation in current populations might, depending on

effective population size and generation time, precede the temporal window of many landscape

genetics studies. In this regard, the greater number of SNP loci translates into a better genomic

representation. Since different genomic regions reflect different coalescent histories (Nielsen &

Slatkin, 2013), increased genomic coverage should better capture the range of processes

conditioning genetic patterns of populations and thus the combined effect of historical

demographic processes and current landscapes. It is then important to consider the relative

contribution of both processes: i) input from new mutations and ii) sorting of standing genetic

variation, rather than exclusively focus on mutation rates differences. Such sorting is expected to

be specific to different genomic regions, which most likely contribute to the differences observed

between our SSR and SNP datasets.

Advancing landscape genetics practice

Signals of population structure may arise from a wide range of evolutionary processes,

including historical demographic events (He, Edwards, & Knowles, 2013), local adaptation

(Orsini, Vanoverbeke, Swillen, Mergeay, & De Meester, 2013), and reproductive strategies (e.g.,

selling in mixed mating species such as *I. purpurea* can lead to a spurious identification of

structure; Gao, Williamson, & Bustamante, 2007). Yet, landscape genetics approaches

traditionally overlook these plausible confounding processes by working under the assumptions

of an equilibrium between migration and genetic drift and an implicit predominance of recent

landscape configurations over alternative explanations for the observed population structure

(Dyer, 2015b; He et al., 2013; Marko & Hart, 2011). Thus, traditional landscape genetics

analyses presumably present an incomplete picture of the evolutionary processes driving current

patterns of genetic diversity (Wang & Bradburd, 2014). Nonetheless, these approaches

undoubtedly offer a valuable hypothesis-generation framework about the possible role that

environmental setting plays in structuring genetic diversity against which the effect of other

demographic processes can be evaluated. Arguably, the integration of landscape genetics with

historical demographic reconstruction is key for robust population genetics inference since

disregarding the plausible effects of either current landscape processes or historical demographic

changes would impair the ability to understand species' complex responses to spatio-temporal

environmental variation.

In this context, considering the likely complex demographic dynamics of this introduced

agricultural weed, our results should be taken as a working hypothesis of the possible role of the

interaction between natural and anthropogenic landscapes in structuring *I. purpurea* populations.

Nonetheless, our analyses represent a step towards integrating traditional landscape genetics with

modern population genetics inference by taking advantage of recent analytical developments and

richer molecular datasets. On one hand, our analyses use novel methodological tools that i)

surpass the need of arbitrary landscape resistance assignment that make inferences sensitive to

subjectivity of expert opinion (Dyer, 2015a), ii) account for the indirect genetic similarity of

populations (Dyer & Nason, 2004), iii) use rigorous statistical inferences (Balkenhol et al., 2009;

Peterman et al., 2014), and account for plausible confounding processes (i.e., local effective

population size; Weckworth et al., 2013)—although more work on accounting for additional

processes such as historical demographic changes is needed. On the other hand, in contrast to the

common practice in the field of using a single analysis (commonly Mantel test; Guillot &

Rousset, 2013) and one or a few loci (although a few notable exceptions exist; e.g. Perry et al.,

2013), which prevents an assessment of common patterns across the genome (Bohonak &

Vandergast, 2011), our inferences are derived from common findings among two rather different

sets of molecular markers. In doing so, we provide not only statistically robust inferences, but

also a better representation of the genome. Hence, our inferences are not only less sensitive to

ascertainment bias (Brandström & Ellegren, 2008) and coalescent and mutational variance

(Buschiazzo & Gemmell, 2006; Nielsen & Slatkin, 2013; Steiner, Putnam, Hoeck, & Ryder,

2013), but have also the ability to uncover differences in the underlying population dynamics.

Such differences have strikingly important implications. For example, when evaluating plausible

approaches to the threat of an invasive species such as *I. purpurea*, recommendations would be

quite different depending on whether gene flow is believed to be relatively widespread (as

inferred by the SSR dataset) or whether it is believed to be minimal (as inferred by the SNP

dataset). In this example, it is clear that knowledge-based management would clearly benefit

from recognizing the current uncertainty in regards to the exact population connectivity as

opposed to automatically relying on a single-marker story.

Given recent advances in next generation sequencing, it seems straightforward to focus on

landscape genomics instead of few loci. Hence, development of methods for explicitly integrating

inferences from multiple genome regions and marker types, as it is customary in population

genetics, would be of great value. By incorporating multiple loci and coupling traditional

landscape genetic tools with coalescent-based simulations to explicitly model landscape effects

on genetic population structure, a robust hypothesis framework could be develop to

simultaneously account for both current landscape processes and demographic history of species

(Balkenhol & Landguth, 2011; Hoban, Bertorelle, & Gaggiotti, 2012). Advances in this area are

already being developed with promising perspectives (Alvarado-Serrano & Knowles, 2014;

Harris et al., 2016; He et al., 2013).

Final remarks

By offering a working hypothesis of the effect of current landscapes on genetic

differentiation, traditional landscape genetics results serve the purpose of identifying relevant

models for further testing (Baguette, Blanchet, Legrand, Stevens, & Turlure, 2013; Dyer, 2015a).

Under this framework, our results pave the way for rigorous simulation-based assessments of the

role of landscape features in promoting or deterring population differentiation in a noxious

agricultural weed, and hence for successful knowledge-based invasive management. Specifically,

we identify a probably major role of human-driven gene flow and long distance dispersal events

in the demographic history of this species. If this weed was, as hypothesized, singly introduced

through horticulture from a European bottlenecked population during the European colonization

of North America (Fang et al., 2013), distinct ancestral structure (pre-dating US introduction)

would be unlikely. Under this scenario, the rather distinct clustering of individual subpopulations

recovered would likely reflect the connectivity driven by agricultural and horticultural activities

and the complex natural/human landscape *I. purpurea* has experienced post-introduction to the

US. Alternatively, human trade might have allowed for recurrent introduction events, which

effects would have probably been amplified by local agricultural and horticultural activities.

Regardless, these findings call for future model-based inference that explicitly considers the

impact of human population density in conjunction with climate to further investigate the

evolutionary drivers of population structure in this noxious weed.

ACKNOWLEDGEMENTS

We thank Adam Kuester for seed collecting and for contributing valuable data for this study. We

also thank Ariana Wilson, Eva Fall, and Dan York for tissue collection. This research was funded

by USDA NIFA grants 04180 and 07191to R.S.B.

DATA ARCHIVING

All data generated is in the process of being archived in Dryad.

AUTHOR CONTRIBUTIONS

D.F.A.-S. and R.S.B conceived the study. D.F.A.-S. and M.V.E. generated and compiled the

molecular and GIS data. D.F.A.-S. analyzed the data. D.F.A.-S., M.V.E., S.M.C., and R.S.B.

wrote the manuscript. All authors read and approved the final submission.

LITERATURE CITED

Alvarado-Serrano, D. F., & Knowles, L. L. (2014). Ecological niche models in phylogeographic studies: Applications, advances and precautions. *Molecular Ecology Resources*, *14*, 233–248.

Auffret, A. G., Berg, J., & Cousins, S. A. O. (2014). The geography of human-mediated dispersal. *Diversity and Distributions*, *20*, 1450–1456.

Auffret, A. G., & Cousins, S. A. O. (2013). Humans as long-distance dispersers of rural plant communities. *PLoS ONE*, *8*, e62763.

Baguette, M., Blanchet, S., Legrand, D., Stevens, V. M., & Turlure, C. (2013). Individual dispersal, landscape connectivity and ecological networks. *Biological Reviews*, *88*, 310–326.

Balkenhol, N., & Landguth, E. L. (2011). Simulation modelling in landscape genetics: on the need to go further. *Molecular Ecology*, *20*, 667–670.

Balkenhol, N., Waits, L. P., & Dezzani, R. J. (2009). Statistical approaches in landscape genetics: an evaluation of methods for linking landscape and genetic data. *Ecography*, *32*, 818–830.

Banks, N. C., Paini, D. R., Bayliss, K. L., & Hodda, M. (2015). The role of global trade and transport network topology in the human-mediated dispersal of alien species. *Ecology Letters*, *18*, 188–199.

Barker, M. A., Thompson, L., & Godley, F. M. (1984). Control of annual morningglories (*Ipomoea* spp.) in soybeans (*Glycine max*). *Weed Science*, *32*, 813–818.

Barrowclough, G. F. (1980). Gene flow, effective population sizes and genetic variance components in birds. *Evolution*, *34*, 789–798.

Bataille, A., Cunningham, A. A., Cruz, M., Cedeño, V., & Goodman, S. J. (2011). Adaptation, isolation by distance and human-mediated transport determine patterns of gene flow among populations of the disease vector *Aedes taeniorhynchus* in the Galapagos Islands. *Infection,*

683  *Genetics and Evolution*, *11*, 1996–2003.

684 Baucom, R. S., & Mauricio, R. (2004). Fitness costs and benefits of novel herbicide tolerance in

685  a noxious weed. *Proceedings of the National Academy of Sciences*, *101*, 13386–13390.

686 Baucom, R. S., & Mauricio, R. (2008). The evolution of novel herbicide tolerance in a noxious

687  weed: the geographic mosaic of selection. *Evolutionary Ecology*, *22*, 85–101.

688 Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and

689  powerful approach to multiple testing. *Journal of the Royal Statistical Society B*, *57*, 289–

690  300.

691 Benvenuti, S. (2007). Weed seed movement and dispersal strategies in the agricultural

692  environment. *Weed Biology and Management*, *7*, 141–157.

693 Blair, R. B. (2001). Birds and butterflies along urban gradients in two ecoregions of the US. In J.

694  L. Lockwood & M. L. McKinney (Eds.), *Biotic homogenization* (pp. 33–56). New York:

695  Kluwer Academic.

696 Bohonak, A. J., & Vandergast, A. G. (2011). The value of DNA sequence data for studying

697  landscape genetics. *Molecular Ecology*, *20*, 2477–2479.

698 Boyd, N. S., & White, S. (2009). Impact of wild blueberry harvesters on weed seed dispersal

699  within and between fields. *Weed Science*, *57*, 541–546.

700 Brandström, M., & Ellegren, H. (2008). Genome-wide analysis of microsatellite polymorphism

701  in chicken circumventing the ascertainment bias. *Genome Research*, *18*, 881–887.

702 Buschiazzo, E., & Gemmell, N. J. (2006). The rise, fall and renaissance of microsatellites in

703  eukaryotic genomes. *BioEssays*, *28*, 1040–1050.

704 Bushnell, B. (2016). BBMap short read aligner. Retrieved from

705  http://sourceforge.net/projects/bbmap/

706 Bussell, J. D. (1999). The distribution of random amplified polymorphic DNA (RAPD) diversity

amongst populations of *Isotoma petraea* (Lobeliaceae). *Molecular Ecology*, *8*, 775–789.

Chaney, L., & Baucom, R. S. (2012). The evolutionary potential of Baker's weediness traits in

the common morning glory, *Ipomoea purpurea* (Convolvulaceae). *American Journal of*

Botany, *99*, 1524–1530.

Chen, C., Durand, E., Forbes, F., & François, O. (2007). Bayesian clustering algorithms

ascertaining spatial population structure: a new computer program and a comparison study.

Molecular Ecology Notes, *7*, 747–756.

Cimalová, Š.ˇ, & Lososová, Z. (2009). Arable weed vegetation of the northeastern part of the

Czech Republic: effects of environmental factors on species composition. *Plant Ecology*,

203, 45–57.

Clobert, J., Baguette, M., Benton, T. G., Bullock, J. M., & Ducatez, S. (2012). *Dispersal Ecology*

and Evolution (1st ed.). Oxford: Oxford University Press.

Culpepper, A. S. (2006). Glyphosate-induced weed shifts. *Weed Technology*, *20*, 277–281.

Cushman, S. A., McKelvey, K. S., Hayden, J., & Schwartz, M. K. (2006). Gene flow in complex

landscapes: testing multiple hypotheses with causal modeling. *The American Naturalist*,

168, 486–499.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin, R.

(2011). The variant call format and VCFtools. *Bioinformatics*, *27*, 2156–2158.

Dastgheib, F. (1989). Relative importance of crop seed, manure and irrigation water as sources of

weed infestation. *Weed Research*, *29*, 113–116.

Defelice, M. S. (2001). Tall morningglory, *Ipomoea purpurea* (L .) roth-flower or foe? *Weed*

Technology, *15*, 601–606.

Do, C., Waples, R. S., Peel, D., Macbeth, G. M., Tillett, B. J., & Ovenden, J. R. (2014).

NeEstimator v2: re-implementation of software for the estimation of contemporary effective

population size (Ne) from genetic data. *Molecular Ecology Resources*, *14*, 209–214.

Durand, E., Chen, C., & Francois, O. (2009). TESS version 2.3 - reference manual August 2009.

Available at: memberstimc.imag.fr/Olivier.Francois/tess.html, 1–30.

Durand, E., Jay, F., Gaggiotti, O. E., & François, O. (2009). Spatial inference of admixture

proportions and secondary contact zones. *Molecular Biology and Evolution*, *26*, 1963–1973.

Dyer, R. J. (2009). GeneticStudio: a suite of programs for spatial analysis of genetic-marker data.

Molecular Ecology Resources, *9*, 110–3.

Dyer, R. J. (2015a). Is there such a thing as landscape genetics? *Molecular Ecology*, *10*, 139–143.

Dyer, R. J. (2015b). Population graphs and landscape genetics. *Annual Review of Ecology,*

Evolution, and Systematics, *46*, 327–342.

Dyer, R. J., & Nason, J. D. (2004). Population graphs: the graph theoretic shape of genetic

structure. *Molecular Ecology*, *13*, 1713–1727.

Dyer, R. J., Nason, J. D., & Garrick, R. C. (2010). Landscape modelling of gene flow: improved

power using conditional genetic distance derived from the topology of population networks.

Molecular Ecology, *19*, 3746–3759.

Eberhardt, M., & Teal, F. (2013). No mangoes in the tundra: spatial heterogeneity in agricultural

productivity analysis. *Oxford Bulletin of Economics and Statistics*, *75*, 914–939.

Economic Research Service, U. S. D. A. (1998). Seed industry structure is characterized by

growth and consolidation. *AIB*, *786*, 25–29.

Edelaar, P., & Bolnick, D. I. (2012). Non-random gene flow: an underappreciated force in

evolution and ecology. *Trends in Ecology and Evolution*, *27*, 659–665.

Ennos, R. A. (1981). Quantitative studies of the mating system in two sympatric species of

Ipomoea (Convolvulaceae). *Genetica*, *57*, 93–98.

Epperson, B. K., & Clegg, M. T. (1986). Spatial-autocorrelation analysis of flower color

polymorphisms within substructured populations of morning glory (*Ipomoea purpurea*). *The*

American Naturalist, *128*, 840–858.

Eschmann-Grupe, G., Neuffer, B., & Hurka, H. (2004). Extent and structure of genetic variation

in two colonising *Diplotaxis* species (Brassicaceae) with contrasting breeding systems. *Plant*

Systematics and Evolution, *244*, 31–43.

Everman, E., & Klawinski, P. (2013). Human-facilitated jump dispersal of a non-native frog

species on Hawai'i Island. *Journal of Biogeography*, *40*, 1961–1970.

Excoffier, L., Smouse, P. E., & Quattro, J. M. (1992). Analysis of molecular variance inferred

from metric distances among DNA haplotypes: application to human mitochondrial DNA

restriction data. *Genetics*, *131*, 479–491.

Fang, Z., Gonzales, A. M., Durbin, M. L., Meyer, K. K. T., Miller, B. H., Volz, K. M., …

Morrell, P. L. (2013). Tracing the geographic origins of weedy *Ipomoea purpurea* in the

Southeastern United States. *Journal of Heredity*, *104*, 666–677.

Fountain, T., Duvaux, L., Horsburgh, G., Reinhardt, K., & Butlin, R. K. (2014). Human-

facilitated metapopulation dynamics in an emerging pest species, *Cimex lectularius*.

Molecular Ecology, *23*, 1071–1084.

Gao, H., Williamson, S., & Bustamante, C. D. (2007). A markov chain monte carlo approach for

joint inference of population structure and inbreeding rates from multilocus genotype data.

Genetics, *176*, 1635–1651.

Garza, J. C., & Freimer, N. B. (1996). Homoplasy for size at microsatellite loci in humans and

chimpanzees. *Genome Research*, *6*, 211–217.

Glickman, M. E., Rao, S. R., & Schultz, M. R. (2014). False discovery rate control is a

recommended alternative to Bonferroni-type adjustments in health studies. *Journal of*

Clinical Epidemiology, *67*, 850–857.

Goslee, S. C., & Urban, D. L. (2007). The ecodist package for dissimilarity-based analysis of

ecological data. *Journal of Statistical Software*, *22*, 1–19.

Guillot, G., & Rousset, F. (2013). Dismantling the Mantel tests. *Methods in Ecology and*

Evolution, *4*, 336–344.

Harris, S. E., Xue, A. T., Alvarado-Serrano, D. F., Boehm, T., Joseph, T., Hickerson, M. J., &

Munshi-South, J. (2016). Urbanization shapes the demographic history of a native rodent

(the white- footed mouse, *Peromyscus leucopus*) in New York City. *Biology Letters*, *12*, 1–

6.

He, Q., Edwards, D. L., & Knowles, L. L. (2013). Integrative testing of how environments from

the past to the present shape genetic structure across landscapes. *Evolution*, *67*, 3386–3402.

Hoban, S., Bertorelle, G., & Gaggiotti, O. E. (2012). Computer simulations: tools for population

and evolutionary genetics. *Nature Reviews Genetics*, *13*, 110–122.

Hodkinson, D. J., Thompson, K., Journal, T., & Dec, N. (2007). Plant dispersal: the role of man.

Journal of Applied Ecology, *34*, 1484–1496.

Holderegger, R., Buehler, D., Gugerli, F., & Manel, S. (2010). Landscape genetics of plants.

Trends in Plant Science, *15*, 675–683.

Hubisz, M. J., Falush, D., Stephens, M., & Pritchard, J. K. (2009). Inferring weak population

structure with the assistance of sample group information. *Molecular Ecology Resources*, *9*,

1322–1332.

Jha, S. (2015). Contemporary human-altered landscapes and oceanic barriers reduce bumble bee

gene flow. *Molecular Ecology*, *24*, 993–1006.

Jha, S., & Kremen, C. (2013). Urban land use limits regional bumble bee gene flow. *Molecular*

Ecology, *22*, 2483–2495.

Kolbe, J. J., Glor, R. E., Rodríguez Schettino, L., Lara, A. C., Larson, A., & Losos, J. B. (2004).

803  Genetic variation increases during biological invasion by a Cuban lizard. *Nature*, *431*, 177–

804  81.

805 Kuester, A., Chang, S.-M., & Baucom, R. S. (2015). The geographic mosaic of herbicide

806  resistance evolution in the common morning glory, *Ipomoea purpurea*: evidence for

807  resistance hotspots and low genetic differentiation across the landscape. *Evolutionary*

808  *Applications*, *8*, 821–833.

809 Legendre, P., Lapointe, F. J., & Casgrain, P. (1994). Modeling brain evolution from behavior: a

810  permutational regression approach. *Evolution*, *48*, 1487–1499.

811 Lehan, N. E., Murphy, J. R., Thorburn, L. P., & Bradley, B. A. (2013). Accidental introductions

812  are an important source of invasive plants in the continental United States. *American*

813  *Journal of Botany*, *100*, 1287–1293.

814 Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology and*

815  *Evolution*, *17*, 183–189.

816 Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association mapping

817  and population genetical parameter estimation from sequencing data. *Bioinformatics*, *27*,

818  2987–2993.

819 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., … Durbin, R. (2009). The

820  Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*, 2078–2079.

821 Mack, R. N., & Lonsdale, W. M. (2001). Humans as global plant dispersers: getting more than

822  we bargained for. *BioScience*, *51*, 95–102.

823 Magwene, P. M. (2001). New tools for studying integration and modularity. *Evolution*, *55*, 1734–

824  1745.

825 Margosian, M. L., Garrett, K. A., Hutchinson, J. M. S., & With, K. A. (2009). Connectivity of the

826  American agricultural landscape: assessing the national risk of crop pest and disease spread.

BioScience, *59*, 141–151.

Marko, P. B., & Hart, M. W. (2011). The complex analytical landscape of gene flow inference.

Trends in Ecology and Evolution, *26*, 448–456.

Martínez-Abraín, A. (2008). Statistical significance and biological relevance: a call for a more

cautious interpretation of results in ecology. *Acta Oecologica*, *34*, 9–11.

Martins, A. C., Goncalves, R. B., & Melo, G. A. R. (2013). Changes in wild bee fauna of a

grassland in Brazil reveal negative effects associated with growing urbanization during the

last 40 years. *Zoologia*, *30*, 157–176.

McRae, B. (2006). Isolation by resistance. *Evolution*, *60*, 1551–1561.

Medley, K. A., Jenkins, D. G., & Hoffman, E. A. (2015). Human-aided and natural dispersal

drive gene flow across the range of an invasive mosquito. *Molecular Ecology*, *24*, 284–295.

Meehan, T. D., Werling, B. P., Landis, D. A., & Gratton, C. (2011). Agricultural landscape

simplification and insecticide use in the Midwestern United States. *Proceedings of the*

National Academy of Sciences of the United States of America, *108*, 11500–11505.

Menchari, Y., Délye, C., & Le Corre, V. (2007). Genetic variation and population structure in

black-grass (*Alopecurus myosuroides* Huds.), a successful, herbicide-resistant, annual grass

weed of winter cereal fields. *Molecular Ecology*, *16*, 3161–3172.

Mitchell, M. (1996). *An introduction to genetic algorithms*. Cambridge Mass.: MIT Press.

Molecular Ecology Resources Primer Development Consortium, -, Aksoy, S., Almeida-Val, V.

M. F., Azevedo, V. C. R., Baucom, R., Bazaga, P., … Cian, M. J. (2013). Permanent genetic

resources added to molecular ecology resources database 1 October 2012–30 November

2012. *Molecular Ecology Resources*, *13*, 341–343.

Morin, P. A., Luikart, G., & Wayne, R. K. (2004). SNPs in ecology, evolution and conservation.

Trends in Ecology and Evolution, *19*, 208–216.

Müller-Schärer, H., & Fischer, M. (2001). Genetic structure of the annual weed *Senecio vulgaris*

in relation to habitat type and population size. *Molecular Ecology*, *10*, 17–28.

Murphy, H. T., & Lovett-Doust, J. (2004). Context and connectivity in plant metapopulations and

landscape mosaics: does the matrix matter? *Oikos*, *105*, 3–14.

Nakagawa, S. (2004). A farewell to Bonferroni: the problems of low statistical power and

publication bias. *Behavioral Ecology*, *15*, 1044–1045.

Navas, M. L. (2012). Trait-based approaches to unravelling the assembly of weed communities

and their impact on agro-ecosystem functioning. *Weed Research*, *52*, 479–488.

Nielsen, R., & Slatkin, M. (2013). *An introduction to population genetics: theory and*

applications. Sunderland Mass.: Sinauer Associates.

Nosil, P., Egan, S. P., & Funk, D. J. (2008). Heterogeneous genomic differentiation between

walking-stick ecotypes: "isolation by adaptation" and multiple roles for divergent selection.

Evolution, *62*, 316–36.

Oksanen, J., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., … Helene,

W. (2015). vegan: community ecology package. Available at: http://cran.r-

project.org/package=vegan.

Orsini, L., Vanoverbeke, J., Swillen, I., Mergeay, J., & De Meester, L. (2013). Drivers of

population genetic differentiation in the wild: isolation by dispersal limitation, isolation by

adaptation and isolation by colonization. *Molecular Ecology*, *22*, 5983–5999.

Peakall, R., & Smouse, P. E. (2012). GenAlEx 6.5: Genetic analysis in Excel. Population genetic

software for teaching and research—an update. *Bioinformatics*, *28*, 2537–2539.

Perry, G. H., Louis, E. E., Ratan, A., Bedoya-Reina, O. C., Burhans, R. C., Lei, R., … Miller, W.

(2013). Aye-aye population genomic analyses highlight an important center of endemism in

northern Madagascar. *Proceedings of the National Academy of Sciences of the United States*

of America, *110*, 5823–5828.

Peterman, W. E. (2014). ResistanceGA: an R package for the optimization of resistance surfaces

using genetic algortihms. *bioRxiv*, *7575*.

Peterman, W. E., Connette, G. M., Semlitsch, R. D., & Eggert, L. S. (2014). Ecological resistance

surfaces predict fine-scale genetic differentiation in a terrestrial woodland salamander.

Molecular Ecology, *23*, 2402–2413.

Pimentel, D., Zuniga, R., & Morrison, D. (2005). Update on the environmental and economic

costs associated with alien-invasive species in the United States. *Ecological Economics*, *52*,

273–288.

R Core Development Team. (2016). R: a language and environment for statistical computing,

3.2.4.

Ramankutty, N., Evan, A. T., Monfreda, C., & Foley, J. A. (2008). Farming the planet: 1.

Geographic distribution of global agricultural lands in the year 2000. *Global*

Biogeochemical Cycles, *22*, 1–19.

Ricciardi, A. (2007). Are modern biological invasions an unprecedented form of global change?

Conservation Biology, *21*, 329–336.

Rogers, J. B., Murray, D. S., Verhalen, L. M., & Claypool, P. L. (1996). Ivyleaf morningglory

(*Ipomoea hederacea*) interference with cotton (*Gossypium hirsutum*). *Weed Technology*, *10*,

107–114.

Rowe, H. C., Renaut, S., & Guggisberg, A. (2011). RAD in the realm of next-generation

sequencing technologies. *Molecular Ecology*, *20*, 3499–34502.

Schwartz, M. K., & McKelvey, K. S. (2009). Why sampling scheme matters: the effect of

sampling scheme on landscape genetic results. *Conservation Genetics*, *10*, 441–452.

Selkoe, K. A., & Toonen, R. J. (2006). Microsatellites for ecologists: a practical guide to using

and evaluating microsatellite markers. *Ecology Letters*, *9*, 615–629.

Sexton, J. P., Hangartner, S. B., & Hoffmann, A. A. (2014). Genetic isolation by environment or

distance: which pattern of gene flow is most common? *Evolution*, *68*, 1–15.

Shah, V. B., & McRae, B. H. (2008). Circuitscape: a tool for landscape ecology. *Proceedings of*

the 7th Python in Science Conference, 62–65.

Siliceo-Cantero, H. H., García, A., Reynolds, R. G., Pacheco, G., & Lister, B. C. (2016).

Dimorphism and divergence in island and mainland Anoles. *Biological Journal of the*

Linnean Society, *118*, 852–872.

Simberloff, D. (2013). *Invasive species: what everyone needs to know*. Oxford: Oxford

University Press.

Slatkin, M. (1985). Gene flow in natural populations. *Annual Review of Ecology and Systematics*,

16, 393–430.

Spear, S. F., Balkenhol, N., Fortin, M. J., McRae, B. H., & Scribner, K. (2010). Use of resistance

surfaces for landscape genetic studies: considerations for parameterization and analysis.

Molecular Ecology, *19*, 3576–3591.

Steiner, C. C., Putnam, A. S., Hoeck, P. E. A., & Ryder, O. A. (2013). Conservation genomics of

threatened animal species. *Annual Review of Animal Biosciences*, *1*, 261–281.

Taylor, P. D., Fahrig, L., Henein, K., & Merriam, G. (1993). Connectivity is a vital element of

landscape structure. *Oikos*, *68*, 571–573.

Thill, D. C., & Mallory-Smith, C. A. (1997). The nature and consequence of weed spread in

cropping systems. *Weed Science*, *45*, 337–342.

Tiffin, P., & Rausher, M. (1999). Genetic constraints and selection acting on tolerance to

herbivory in the common morning glory *Ipomoea purpurea*. *The American Naturalist*, *154*,

700–716.

Trakhtenbrot, A., Nathan, R., Perry, G., & Richardson, D. M. (2005). The importance of long-

distance dispersal in biodiversity conservation. *Diversity and Distributions*, *11*, 173–181.

Turakulov, R., & Easteal, S. (2003). Number of SNPs loci needed to detect population structure.

Human Heredity, *55*, 37–45.

Van Oosterhout, C., Hutchinson, W. F., Wills, D. P. M., & Shipley, P. (2004). MICRO-

CHECKER: software for identifying and correcting genotyping errors in microsatellite data.

Molecular Ecology Notes, *4*, 535–538.

Wang, I. J. (2010). Recognizing the temporal distinctions between landscape genetics and

phylogeography. *Molecular Ecology*, *19*, 2605–2608.

Wang, I. J., & Bradburd, G. S. (2014). Isolation by environment. *Molecular Ecology*, *23*, 5649–

5662.

Wang, I. J., Savage, W. K., & Bradley Shaffer, H. (2009). Landscape genetics and least-cost path

analysis reveal unexpected dispersal routes in the California tiger salamander (*Ambystoma*

californiense). *Molecular Ecology*, *18*, 1365–1374.

Webster, T. M., & Nichols, R. L. (2012). Changes in the prevalence of weed species in the major

agronomic crops of the southern United States: 1994/1995 to 2008/2009. *Weed Science*, *60*,

145–157.

Weckworth, B. V, Musiani, M., Decesare, N. J., Mcdevitt, A. D., Hebblewhite, M., & Mariani, S.

(2013). Preferred habitat and effective population size drive landscape genetic patterns in an

endangered species. *Proceedings of the Royal Society B*, *280*, 1–9.

White, D. R., & Harary, F. (2001). The cohesiveness of blocks in social networks: node

connectivity and conditional density. *Sociological Methodology*, *31*, 305–359.

Wichmann, M. C., Alexander, M. J., Soons, M. B., Galsworthy, S., Dunne, L., Gould, R., …

Bullock, J. M. (2009). Human-mediated dispersal of seeds over long distances. *Proceedings*

of the Royal Society B, *276*, 523–532.

Wilson, G. A., & Rannala, B. (2003). Bayesian inference of recent migration rates using

multilocus genotypes. *Genetics*, *163*, 1177–1191.

Wright, S. (1949). The genetical structure of populations. *Annals of Eugenics*, *15*, 323–354.

Ye, W. H., Mu, H. P., Cao, H. L., & Ge, X. J. (2004). Genetic structure of the invasive

Chromolaena odorata in China. *Weed Research*, *44*, 129–135.

TABLES

Table 1. Analysis of Molecular Variance (AMOVA) of SSR and SNP data. The contribution of

spatial clusters (regions), localities, and individuals is shown. For comparison, results from

an AMOVA analysis with no region category defined are presented in parentheses

underneath.

Effect	F-statistic	Variance explained		F-value		P-value	
		SSR	SNP	SSR	SNP	SSR	SNP
Regions	F_{RT}	3.94%	8.51%	0.006	0.085	0.001	0.001
Localities	F_{SR}	9.05% (11.06%)	6.10% (13.02%)	0.106	0.067	0.001	0.001
Individuals (among)	F_{ST}	0.67% (38.33%)	24.85% (25.31%)	0.112 (0.111)	0.146 (0.130)	0.001 (0.001)	0.001 (0.001)
Individuals (within)	F_{IS}	86.33% (50.61%)	60.54% (61.67%)	0.431 (0.431)	0.291 (0.291)	0.001 (0.001)	0.001 (0.001)
Total	F_{IT}	100% (100%)	100% (100%)	0.494 (0.494)	0.395 (0.383)	0.001 (0.001)	0.001 (0.001)

   Table 2. Summary of landscape genetics models. Model coefficients are reported followed

962   by associated p-value (in parenthesis) and, for MLPE.lmm models, followed by AICc

963   difference and ranking (in square brackets). Significant coefficients are in bold, marginally

964   significant coefficients are marked with an asterisk.

Feature	MLPE.lmm		MRDM	
	SSR	SNP	SSR	SNP
Intrinsic variables				
Geographic distance	0.187* (0.052) +12.303 [8]	0.780* (0.055) +0.396 [4]	0.051 (0.121)	0.255* (0.056)
Population size (Ne)	—	—	**-0.550** (0.001)	-2.650* (0.051)
Natural environment variables				
Climate PC1	**0.243** (0.034) +10.356 [3]	**0.967** (0.045) +0.032 [2]	0.533* (0.064)	1.782 (0.517)
Climate PC2	**0.205** (0.048) +12.030 [7]	0.814* (0.055) +1.560 [7]	-0.029 (0.978)	21.443* (0.075)
Elevation	**0.244** (0.034) +10.742 [4]	0.840* (0.054) +1.423 [6]	-0.607 (0.480)	-31.789* (0.095)
Soil PC1	**0.208** (0.039) +11.378 [6]	**1.044** (0.045) +0.471 [5]	0.305 (0.510)	**15.789** (0.038)
Soil PC2	**0.320** (0.018) +8.139 [2]	**0.941** (0.045) + 0.284 [3]	0.187 (0.703)	-6.838 (0.476)
Human-impact variables				
Crops	-0.226 (0.134) +14.491 [9]	0.858* (0.054) +15.371 [8]	0.154 (0.526)	0.360 (0.840)
Landcover	**0.582**	**1.358**	0.340	3.887

	(<0.001)	(0.003)	(0.218)	(0.184)
	–	+39.775		
	[1]	[9]		
Population density	**0.227**	**0.912**	-0.519*	**-3.271**
	(0.034)	(0.045)	(0.095)	(0.037)
	+10.821	–		
	[5]	[1]		

967 **FIGURES**

969 **Figure 1**. Distribution of *Ipomoea purpurea*'s sampled localities. Sample sizes for both SSR

970 (black bars) and SNP (white bars) datasets are indicated (locality numbers are given in

971 squares). Elevation is provided as background.

Figure 2. Inferred population connectivity. The estimated origin of individuals for each

sampled locality (i.e., sink) is depicted according to the locality they were inferred to have

originated from (source) (a, d). The color of each cell in these plots depicts the proportion

of individuals in the sink population that were estimated to be recent immigrants from each

locality along the x-axis. Cells on the minor diagonal correspond to the proportion of native

individuals. Pruned conditional genetic networks (b, e) and posterior estimates of

admixture proportion identified by TESS analysis (c, f) are also displayed. The top row

shows SSR-based results, the bottom shows the SNP-based results. Locality numbers follow

Fig. 1. Localities shared between SSR and SNP datasets are denoted by unique colors in all

panels (for a corresponding figure based exclusively on these shared localities, see Fig. S3).