Top-down beta enhances bottom-up gamma.

Abbreviated title: Top-down beta enhances bottom-up gamma.

Craig G. Richter[1,*], William H. Thompson[1,2,*], Conrado A. Bosman[3,4] Pascal Fries[1,3]

[1] Ernst Strüngmann Institute (ESI) for Neuroscience in Cooperation with Max Planck Society, Deutschordenstraße 46, 60528 Frankfurt, Germany

[2] Department of Clinical Neuroscience, Karolinska Institute, Nobels Väg 9, 171 77 Solna, Sweden

[3] Donders Institute for Brain, Cognition and Behaviour, Kapittelweg 29, Radboud University, 6525 EN Nijmegen, Netherlands

[4] Swammerdam Institute for Life Sciences, Center for Neuroscience, Faculty of Science, University of Amsterdam, Sciencepark 904, 1098 XH Amsterdam, Netherlands

*C.R. and W.T. contributed equally to this study

Correspondence should be addressed to Dr. Craig Richter or Dr. Pascal Fries, Ernst Strüngmann Institute (ESI) for Neuroscience in Cooperation with Max Planck Society, Deutschordenstraße 46, 60528 Frankfurt, Germany. E-mails: craig.richter@esi-frankfurt.de, pascal.fries@esi-frankfurt.de.

Pages: 31

Figures: 8

Number of words: Abstract: 126; Introduction: 406; Discussion: 1334

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgements: This work was supported by DFG (SPP 1665, FOR 1847, FR2557/5-1-CORNET), EU (HEALTH-F2-2008-200728-BrainSynch, FP7-604102-HBP), a European Young Investigator Award, National Institutes of Health (1U54MH091657-WU-Minn-Consortium-HCP), the LOEWE program (NeFF) and the FLAG-ERA JTC 2015 project CANON (co-funded by NWO to CAB). We would like to thank Julien Vezoli for his assistance in ROI definition, co-registration and construction of anatomical renderings.

1 Abstract

Several recent studies have demonstrated that the bottom-up signaling of a visual stimulus is
subserved by interareal gamma-band synchronization, whereas top-down influences are mediated by
alpha-beta band synchronization. These processes may implement top-down control of stimulus
processing if top-down and bottom-up mediating rhythms are coupled via cross-frequency interaction.
To test this possibility, we investigated Granger-causal influences among awake macaque primary
visual area V1, higher visual area V4 and parietal control area 7a during attentional task performance.
Top-down 7a-to-V1 beta-band influences enhanced visually driven V1-to-V4 gamma-band influences.
This enhancement was spatially specific and largest when beta-band activity preceded gamma-band
activity by ~0.1 s, suggesting a causal effect of top-down processes on bottom-up processes. We
propose that this cross-frequency interaction mechanistically subserves the attentional control of
stimulus selection.

13 Significance Statement

Contemporary research indicates that the alpha-beta frequency band underlies top-down control,
while the gamma-band mediates bottom-up stimulus processing. This arrangement inspires an
attractive hypothesis, which posits that top-down beta-band influences directly modulate bottom-up
gamma band influences via cross-frequency interaction. We evaluate this hypothesis determining that
beta-band top-down influences from parietal area 7a to visual area V1 are correlated with bottom-up
gamma frequency oscillations from V1 to area V4, in a spatially specific manner, and that this
correlation is maximal when top-down activity precedes bottom-up activity. These results show that for
top-down processes such as spatial attention, elevated top-down beta-band influences directly
enhance feedforward stimulus induced gamma-band processing, leading to enhancement of the
selected stimulus.

Richter et al., *Top-down beta enhances bottom-up gamma.*

Introduction

Many cognitive effects in vision can only be explained by invoking the concept of top-down influences (Gilbert and Sigman, 2007). For example, when top-down influences pre-allocate attention to specific spatial locations, stimulus processing is more accurate and/or faster. Correspondingly, neurons in higher visual areas show enhanced firing when processing attended stimuli. These neurophysiological consequences of top-down control must be mediated by corresponding anatomical projections. Indeed, anatomical studies have documented projections in the top-down direction that are as numerous as bottom-up projections. Bottom-up and top-down projections show different characteristic laminar patterns of origin and termination, and the pattern of interareal pairwise projections abides by a global hierarchy (Felleman and Van Essen, 1991; Hilgetag et al., 1996; Markov et al., 2014). Recently, it has been shown in both macaque and human visual cortex, that the pattern of anatomical projections is closely correlated to a pattern of frequency-specific directed interareal influences. Influences mediated by bottom-up projections are primarily carried by gamma-band synchronization; influences mediated by top-down projections are primarily carried by alpha-beta-band synchronization (Bastos et al., 2012; Bosman et al., 2012; Grothe et al., 2012; Jia et al., 2013; van Kerkoerle et al., 2014; Bastos et al., 2015b; 2015c; Michalareas et al., 2016). A similar association of higher frequency bands with bottom-up and lower frequency bands with top-down signaling has also been found in other systems (Colgin et al., 2009; Bieri et al., 2014; Fontolan et al., 2014) and related to respective task demands (von Stein and Sarnthein, 2000; Buschman and Miller, 2007).

Here, we investigate directly whether top-down influences actually modulate bottom-up influences. We simultaneously assess influences in both directions through multi-area electrophysiology and frequency-resolved Granger causality analysis. We find that on an epoch-by-epoch basis, top-down beta-band influences enhance bottom-up gamma-band influences. This effect is spatially specific, i.e. bottom-up gamma-band influences depend most strongly on the top-down beta-band influences that are directed to the origin of the bottom-up influence. Spontaneous enhancements in top-down beta-band influences are followed ~0.1 s later by enhancements in bottom-up gamma-band influences, suggestive of a causal relation.

We suggest that this is a mechanism for the implementation of spatially selective attention. Attentional control areas in posterior parietal cortex contain maps of visual space, with neuronal ensemble activity

Richter et al.*, Top-down beta enhances bottom-up gamma.*

53 describing the spatial location of a subject's attention (Bisley and Goldberg, 2003). These neurons

54 likely exert top-down beta-band influences on retinotopically corresponding parts of early visual cortex,

55 which in turn enhance the bottom-up forwarding of visual stimuli.

Materials and Methods

Visual stimulation and behavioral task. The experiment was approved by the ethics committee of the

Radboud University Nijmegen (Nijmegen, The Netherlands). Two adult male macaque monkeys

(monkey K and monkey P, both macaca mulatta) were used in this study. During experiments,

monkeys were placed in a dimly lit booth facing a CRT monitor (120 Hz non-interlaced). When they

touched a bar, a fixation point was presented, and gaze had to remain within the fixation window

throughout the trial (monkey K: 0.85 deg radius, monkey P: 1 deg radius), otherwise the trial would be

terminated and a new trial would commence. Once central fixation had been achieved and a

subsequent 0.8 s pre-stimulus interval had elapsed, two isoluminant and isoeccentric drifting

sinusoidal gratings were presented, one in each visual hemifield (diameter: 3 deg, spatial frequency:

≈1 cycle/deg, drift velocity: ≈1 deg/s, resulting temporal frequency: ≈1 cycle/s, contrast: 100%). Blue

and yellow tints were randomly assigned to each of the gratings on each trial (Fig. 1A). Following a

random delay interval (monkey K : 1 - 1.5 s; monkey P : 0.8 - 1.3 s), the central fixation point changed

color to match one of the drifting gratings, indicating that this grating was the target stimulus, i.e. the

fixation point color was the attentional cue. When the target stimulus was positioned in the visual

hemifield contralateral to the recorded hemisphere, we refer to this condition as attend-contra,

whereas when the target was in the ipsilateral hemifield with respect to the ECoG grid, this condition is

labeled attend-ipsi. Either the target or distracter stimulus could undergo a subtle change in shape

consisting of a transient bending of the bars of the grating (0.15 s duration of the full bending cycle).

This change could occur at any monitor refresh from 0.75 s to 5 s (monkey K), and 4 s (monkey P)

after stimulus onset. Bar releases within 0.15 - 0.5 s after target changes were rewarded. If stimulus

changes occurred before the cue indicated which stimulus was the target, reports were rewarded in a

random half of trials. Bar releases after distracter changes terminated the trial without reward. Trials

were pooled from both contra and ipsi conditions, except where explicit comparisons of these

conditions were made.

Richter et al.*, Top-down beta enhances bottom-up gamma.*

Neurophysiological recordings and signal preprocessing. LFP recordings were made via a 252

channel electrocorticographic grid (ECoG) implanted subdurally over the left hemisphere (Rubehn et

al., 2009). Data from the same animals, partly overlapping with the data used here, have been used in

several previous studies (Bosman et al., 2012; Pinotsis et al., 2014; Brunet et al., 2014a; 2014b; 2015;

Richter et al., 2015; Vinck et al., 2015; Bastos et al., 2015b; 2015c; Lewis et al., 2016). Recordings

were sampled at approximately 32 kHz with a passband of 0.159 – 8000 Hz using a Neuralynx Digital

Lynx system. The raw recordings were low-pass filtered at 250 Hz, and downsampled to 1 kHz. The

electrodes were distributed over eight 32-channel headstages, and referenced against a silver wire

implanted onto the dura overlying the opposite hemisphere. The electrodes were re-referenced via a

bipolar scheme to achieve 1) greater signal localization 2) cancellation of the common reference,

which could corrupt the validity of connectivity metrics, 3) to reject headstage specific noise. The

bipolar derivation scheme subtracted the recordings from neighboring electrodes (spaced 2.5 mm)

that shared a headstage, resulting in 218 bipolar derivations, henceforth referred to as "sites" (see

Bastos et al. (2015c) for a detailed description of the re-referencing procedure). The site locations are

shown as spheres in Figure 1b (monkey K: light gray, monkey P: black).

All signal processing was conducted in MATLAB (MathWorks, USA) and using the FieldTrip toolbox

(http://www.fieldtriptoolbox.org/) (Oostenveld et al., 2011). Raw data were cleaned of line noise via the

subtraction of 50, 100, and 150 Hz components fit to the data using a discrete Fourier transform.

Following trial epoching (see below for details), epochs for each site were de-meaned by subtracting

the mean over all time points in the epoch. Epochs exceeding 5 standard deviations of all data from

the same site in the same session were rejected. In addition, epochs were manually inspected and

epochs with artifacts were rejected. The remaining epochs were normalized by the standard deviation

across all data in all epochs from the same site in the same recording session. Subsequently, all

epochs of a given site were combined across sessions.

Region of interest definition. Three regions-of-interest (ROIs) were selected for the current study: V1,

V4, and area 7A (referred to simply as "7A"). ROIs were defined based on comparison of the electrode

locations (co-registered to each monkey's anatomical MRI and warped to the F99 template brain in

CARET (Van Essen, 2012), with multiple cortical atlases of the macaque (see Bastos et al. (2015c) for

a detailed description). Recording sites composing each ROI were co-registered to a common

Richter et al.*, Top-down beta enhances bottom-up gamma.*

template (INIA19, (Rohlfing et al., 2012)), as were the Paxinos ROI definitions (Paxinos et al., 1999).

The V1/V2 combined definition of Paxinos et al. (1999), is shown in Figures 1B, 3G (red) for simplicity

due to uncertainty across atlases of the V1/V2 border. Recording site selection was based on multiple

atlases with no recording sites selected that were believed to belong to area V2. Based on these ROI

definitions, 77 recording sites were selected from area V1 (monkey K: 29, monkey P: 48), 31 from

area V4 (monkey K: 17, monkey P: 14), and 18 from area 7A (monkey K: 8, monkey P: 10).

Segmenting data into periods and epochs. Each successfully completed trial contained three periods:

The fixation, the stimulation and the attention period. The fixation period was the time between fixation

onset and stimulus onset. During the fixation period, monkeys fixated on a fixation point on a gray

screen, and there was no stimulus presented and no cue had been nor was presented during that

time. The stimulation period was the time between onset of the stimuli and either a change in one of

the stimuli or cue onset. During the stimulation period, monkeys kept fixation, the stimuli were

continuously present, one tinted yellow the other blue, chosen randomly, and the fixation point had not

yet assumed a color, and thereby the attentional cue had not been given. The attention period was the

time between cue onset and a target or distracter change, whichever occurred first. During the

attention period, monkeys kept fixation, the stimuli were continuously present with their tints, and the

fixation point was tinted in one of these colors, thereby providing the attentional cue.

The fixation, stimulation and attention periods all were of variable lengths across trials. The spectral

analysis was based on epochs of fixed lengths. Therefore, the task periods were cut into epochs.

Longer trials contributed correspondingly more epochs. We aimed at excluding data soon after events,

like stimulus onset and cue onset, to minimize effects of post-event transients and non-stationarities

on the metrics of rhythmicity and synchronization. Therefore, from the fixation period, we used the last

0.5 s before stimulus onset; from the stimulation period, the first 0.3 s after stimulus onset were

discarded; for the attention period, the first 0.3 s after cue onset were discarded.

For the analyses of Figure 2 and Figure 3, the remaining parts of the described periods were

segmented into 0.5 s epochs with 60% overlap. This overlap allows for the application of Welch's

method (Welch, 1967) and was selected as an optimal overlap for the multitaper method, while

maintaining a reasonable computational load (Thomson, 1977; Percival and Walden, 1993). This

resulted in 6822 fixation epochs (Monkey K: 3384, Monkey P: 3438), 13675 stimulation epochs

(Monkey K: 8109, Monkey P: 5566), and a total of 16212 attend epochs (Monkey K: 7275, Monkey P:

8937), of which 8313 were attend-contra epochs (Monkey K: 3819, Monkey P: 4494) and 7899 were

attend-ipsi epochs (Monkey K: 3456, Monkey P: 4443).

For the analyses of Figures 4-7, the same procedure was followed, but using only the attention period

and segmenting it into epochs without overlap, because the correlation analyses shown in these

figures require independent epochs. This resulted in fewer epochs, totaling 6414 attend epochs

(Monkey K: 2655, Monkey P: 3759).

For the analyses of Figure 8, the attention period was segmented into many overlapping 0.5 s epochs,

with epochs centers stepped by 0.005 s and epochs fully contained within 0.3-2 s after cue onset. This

was done, because the analysis required GC influence time series. It resulted in a total of 878 time

series (Monkey K: 398, Monkey P: 480).

Spectral Analysis of power, phase locking and directed influences. The 0.5 s epochs were tapered and

Fourier transformed using the Fast Fourier Transform (FFT). For frequencies from 0-50 Hz, a Hann

taper was utilized, whereas for frequencies above 50 Hz, the multitaper method (MTM) was used to

improve the spectral concentration of the gamma rhythm (Thomson, 1982; Percival and Walden,

1993). We applied 5 tapers, resulting in a spectral smoothing of +/- 6 Hz. All epochs were zero-padded

to 1 s, resulting in a spectral resolution of 1 Hz. The coefficients resulting from the FFT were used to

determine power-spectral densities (PSDs) and cross-spectral densities (CSDs), which are the basis

for the two employed connectivity metrics: pairwise phase consistency (PPC) (Vinck et al., 2010), and

Granger causality (GC) (Granger, 1969; Bressler and Seth, 2011). For the display of power spectra,

the power dropoff with frequency, often referred to as 1/f, was partly compensated by multiplying

power values at each frequency by the respective frequency values; for example, the power at 40 Hz

was multiplied by 40 (Sirota et al., 2008) (Fig. 2A-F). The PPC metric (Fig. 2G-J, Fig. 3A, B), in

contrast to the classic coherence metric, does not contain a sample-size bias, that is, it can be directly

compared between different numbers of epochs. The GC metric (Fig. 3C-F) was computed directly

from the CSDs, using non-parametric spectral matrix factorization, in contrast to the traditional

parametric method based on autoregressive modeling (Dhamala et al., 2008). Connectivity metrics

were computed for all interareal site pairs of the ROI pairs V1-V4 and V1-7A.

Richter et al.*, Top-down beta enhances bottom-up gamma.*

Statistical comparison of spectral metrics. Statistical comparison of interaction metrics between
conditions used non-parametric randomization, entailing correction for multiple comparisons across
frequencies (Maris and Oostenveld, 2007). We illustrate this for PPC. The observed PPC spectra
(without randomization) are derived by 1) Calculating PPC spectra across all epochs of a given
condition in a given animal, separately for all site pairs of the relevant ROI pair, 2) Averaging PPC
spectra across those site pairs, 3) Averaging across the two animals after aligning the peaks of the
respective frequency bands. We performed 1000 randomizations, each constituting the null
hypothesis. For each randomization, the following steps were performed: 1) Epochs were randomly
distributed between conditions. 2) The same steps as for the observed PPC spectra were followed.
3) The maximal absolute difference value across all frequencies was retained and placed into the
randomization distribution. The observed differences were compared against the distribution of
maximal absolute differences. Observed absolute difference values greater than the 97.5[th] percentile
of the randomization distribution were considered significant at $p<0.05$. The selection of the maximal
absolute difference value per randomization implements the correction for multiple comparisons
across frequencies (Nichols and Holmes, 2002). This procedure was used for both PPC and
GC influences spectra, separately for the respective frequency bands, either comparing stimulation to
fixation or attend-contra to attend-ipsi.

The GC influence metric is known to be biased by sample size (Bastos and Schoffelen, 2015a), thus
the number of epochs per attention condition needed to be balanced for each monkey. This was
accomplished by finding the condition with the fewest epochs, and randomly selecting this number of
epochs from the other condition.

The comparison of top-down versus bottom-up GC was performed on the pooled data from the attend-
contra and attend-ipsi conditions. Since this was a within-condition comparison, no balancing of epoch
numbers was needed, and all epochs from both attention conditions were used. The statistical
analysis of the difference between top-down and bottom-up GC could not be obtained using a non-
parametric randomization framework, because top-down and bottom-up GC are not properties of
specific sets of epochs, but rather are expressed by all trials simultaneously. Therefore, an alternative
statistical approach was used, namely the bootstrap (Efron and Tibshirani, 1994). Like with the
randomization approach, the statistic of interest (in this case the top-down/bottom-up GC difference) is

Richter et al._, Top-down beta enhances bottom-up gamma._

recomputed on each bootstrap resample, giving rise to a distribution of surrogate values. Following

Efron and Tibshirani (1994), a confidence interval was constructed from the surrogate distribution. To

assess the statistical significance at p<0.05 (two-tailed), we find the 2.5^{th} and 97.5^{th} percentile values

from the surrogate distribution of differences between top-down and bottom-up GC. This naturally

forms the 95% confidence interval such that if zero lies outside of this interval, we may conclude that

the result is significant at a level of p<0.05. This method does not control for multiple comparisons, but

it can be easily modified to do so using the same logic as employed by Maris and Oostenveld (2007).

We performed 1000 bootstrap resamples. For each resample we determined the absolute difference

across frequencies between the bootstrap resample spectrum and the average of all bootstrap

resamples, and retained the maximum of this value across frequencies. Thus, we are guaranteed to

form the largest confidence interval possible across frequencies and in so doing construct an omnibus

confidence interval that controls for the multiple comparisons. This confidence interval is applied to

each frequency, and where it does not contain zero, the result is significant at p<0.05. To conduct

group level statistics, the omnibus statistic is derived from the mean of each bootstrap resample of the

difference between top-down and bottom-up spectra across both monkeys (first averaged within-

subject across pairs), such that the mean of the empirical difference across the monkeys can be

assessed for significance.

Median split analysis. For the median split analysis, we used a jackknife approach, which leaves out

one epoch at a time. The remaining epochs form a jackknife replication (JR). There are as many JRs

as there are epochs. 7A-to-V1 beta-band GC (beta peak ±1 Hz) was computed for all JRs. The

resulting values were median split, and V1-to-V4 GC computed for the corresponding sets of epochs.

The jackknife procedure inverts the distribution of single-epoch GC values (Richter et al., 2015), such

that the upper half of JR values corresponds to the lower half of the true single epoch values. The

observed differences in V1-to-V4 GC were tested for significance as described for the spectral

metrics, using 200 randomizations. The analysis always included a 7A-to-V1 site pair and a V1-to-V4

site pair, sharing the same V1 site. We refer to this configuration as a triplet, or sometimes as

7A-to-V1-to-V4 triplet. There was a total of 10664 triplets (Monkey K: 3944, Monkey P=6720). The

median split analysis is presented in three forms: 1) for an example triplet, 2) for all triplets, followed

by averaging of V1-to-V4 GC, 3) for median splits based on the averaged 7A-to-V1 GC, followed by

averaging of V1-to-V4 GC. For approaches 2) and 3), individual peak frequencies were aligned across

monkeys. The median split analysis was repeated for the inverse direction, splitting epochs based on

V1-to-V4 gamma-band GC jackknife replications (average of the peak ± 5 Hz), followed by averaging

of 7A-to-V1 GC.

Binning of sorted trials. To determine correlation values after sorting and binning the data, JR values

of 7A-to-V1 beta-band GC were sorted and binned into either 5, 10, 50, or 100 bins of equal size. Per

bin, JR values of 7A-to-V1 beta-band GC and of V1-to-V4 gamma GC were averaged. The Spearman

rank correlation was determined across bins. When data were combined across monkeys, correlation

coefficients were averaged across triplets for each monkey, and then averaged across monkeys.

Statistical testing was based on permutations of bin order. Per permutation, bin order was

randomized, and correlation coefficients were computed and averaged as for the observed data. This

was applied 10000 times, resulting in minimal p-values of 0.0001.

Jackknife Correlation. We aimed at quantifying the correlation between epoch-by-epoch fluctuations in

two GC influences. This is normally precluded by the fact that GC influences are not defined per single

data epoch (without substantially sacrificing spectral resolution and/or signal-to-noise ratio). Therefore,

we used the Jackknife Correlation (JC), which quantifies the correlation by first calculating GC

influences for all leave-one-out subsamples (i.e. the jackknife replications of all epochs) and then

correlating these values (Richter et al., 2015). For each leave-one-out subsample, the GC or any other

smooth function F of the data can be defined as follows:

$$F_{x_j} = F\left(x_1, x_2, \ldots, x_{j-1}, x_{j+1}, \ldots, x_n\right) \tag{1}$$

246 , where x specifies the pair of recording sites and j specifies the index of the left-out observation, here

247 the epoch. Attend contra and attend ipsi conditions were combined for the JC analysis. The JC is

248 defined using the following formula:

Richter et al.*, Top-down beta enhances bottom-up gamma.*

$$JC_{F_x F_y} = \frac{1}{n-1} \sum_{j=1}^{n} \left(\frac{F_{x_j} - \overline{F}_x}{s_{F_x}} \right) \left(\frac{F_{y_j} - \overline{F}_y}{s_{F_y}} \right) \tag{2}$$

, where n is defined as the number of jackknife replications and is equal to the total number of epochs,

F_{x_j} and F_{y_j} are the jackknife replications, $\overline{F}_x$ and $\overline{F}_y$ are the means of the jackknife replications, and s_{F_x}

and s_{F_y} are the standard deviations of the jackknife replications. To use the JC with the Spearman

correlation metric, we applied the above formula on the ranks of F_{x_j} and F_{y_j}.

For the statistical analysis of the observed JC values, we created a distribution of 1000 JC values

under a realization of the null hypothesis of independence between the 7A-to-V1 and V1-to-V4 GC

components of each triplet. This was realized by calculating JC between randomly permuted orderings

of jackknife replications of 7A-to-V1 and V1-to-V4 GC influences, which is equivalent to calculating the

JC between GC influences after leaving out a random epoch for the 7A-to-V1 GC and a random epoch

for the V1-to-V4 GC influence without replacement. To control for multiple comparisons across the

frequency-frequency combinations, the max-based approach (see above) was again employed. For

each permutation, the maximum absolute Spearman's rho value was selected, giving rise to an

omnibus distribution of surrogate correlation coefficients. For the example triplet, the observed values

were compared to the distribution of maximum absolute surrogate correlation values. For the average

across triplets, the average was first calculated over triplets, then over monkeys, after aligning to their

respective peak frequencies. For statistical testing, the same randomization was applied to all triplets

of a given monkey, subsequent averaging was performed as for the observed data, and the maximum-

based multiple comparison correction was applied as for the example triplet. An additional JC analysis

first averaged all 7A-to-V1 site pairs and all V1-to-V4 site pairs to form one triplet per monkey. JC

values were then averaged over monkeys. For statistical testing, the randomization was applied to this

triplet per monkey, followed by averaging over monkeys, and the max-based multiple comparison

correction.

For testing spatial specificity, we analyzed recording site triplets, which did not share the same V1 site

(see Results for additional details): 7A-to-V1$_a$V1$_b$-to-V4. Since a vast number of such triplets exist, yet

we wished to select a number equal to the original number of triplets to control potential statistical

bias, we randomly selected a number of 7A-to-V1$_a$V1$_b$-to-V4 triplets that matched the original number

of 7A-to-V1-to-V4 triplets evaluated for each monkey. We repeated this procedure 100 times and

averaged the outcomes. Results were plotted against the average distance between the two V1 sites,

V1$_a$ and V1$_b$, obtained for each distance interval.

Weighted Jackknife Correlation. JCs might be particularly strong for triplets with strong 7A-to-V1 beta-

band GC and/or strong V1-to-V4 gamma-band GC. To test for this, we weighted JC by multiplying it

with the product of the 7A-to-V1 beta and the V1-to-V4 gamma GC magnitudes. Weights were

normalized such that the average weight across triplets was one. The weighted and the unweighted

JC values were separately averaged per monkey and then across monkeys. Statistics were based on

random exchanges between weighted and unweighted values, within each monkey, and subsequent

averaging as for the observed data. Randomization was applied 1000 times.

Lagged Jackknife Correlation (LJC). We were interested in whether the correlation between top-down

beta GC and bottom-up gamma GC depended on their time lag. We started out by using the JC as

described above. To smooth the results against shifts in peak frequencies over time, we used the

average of the range of the top-down beta GC peak ±5 Hz, and the bottom-up gamma GC peak ±10

Hz. Since a given jackknife replication eliminated the same epoch for the calculation of both GC

influences, this established the correlation at zero time-lag. To investigate the dependence of JC on

time lag between GCs, we computed the JC between GC influences calculated from epochs that were

offset by a variable lag. The epochs were stepped at intervals of t=0.005 s. The offsets were stepped

at τ=0.005 s. Note that stepping of intervals and offsets was in principle independent and could have

been different, but it was chosen to be identical to speed up computation. We refer to this as lagged

JC (LJC):

$$LJC_{F_xF_y}(\tau,t) = \frac{1}{n-1}\sum_{j=1}^{n}\left(\frac{F_{x_{j,t}} - \overline{F}_x}{s_{F_{x_t}}}\right)\left(\frac{F_{y_{j,t+\tau}} - \overline{F}_y}{s_{F_{y_{t+\tau}}}}\right) \tag{3}$$

300 τ was chosen to cover a range of lags from -0.5 s to 0.5 s. The GC calculation itself was as in the

301 previous zero-lag JC, using 0.5 s and the tapering specified above. LJC was calculated across trials,

Richter et al*., Top-down beta enhances bottom-up gamma.*

i.e. leaving out an entire trial at a time (this is different from the previous zero-lag JC, which used

multiple non-overlapping epochs per trial if available). The data that was available per trial allowed for

multiple realizations of the two epochs with a particular lag. For each lag, LJC was calculated

separately for all possible realizations and averaged. The number of possible realizations decreases

as the lag between top-down beta GC and bottom-up gamma GC increases, resulting in fewer LJC

computations that are averaged. This results in a noisier estimate at larger lags, but no systematic

bias in the average JC value. The number of epochs that each LJC is computed upon always equals

the number of trials. Formally, this implementation of the LJC is defined as:

$$LJC_{F_xF_y}(\tau) = \frac{1}{m-\tau}\sum_{t=1}^{m-\tau}\left[\frac{1}{n-1}\sum_{j=1}^{n}\left(\frac{F_{x_{j,t}} - \bar{F}_{x_t}}{s_{F_{x_t}}}\right)\left(\frac{F_{y_{j,t+\tau}} - \bar{F}_{y_{t+\tau}}}{s_{F_{y_{t+\tau}}}}\right)\right] \tag{4}$$

, where m is the number of 0.5 s windows, stepped at 0.005 s, that fit into the trial length of 1.7 s.

Statistical significance was assessed using the same logic as used for the JC, where the epoch order

of one member of the JC was permuted with respect to the other. For the LJC, the permutation was

identical for each time step and lag, to be conservative. Multiple comparison correction must take

place over the multiple lags, which was achieved by taking the maximum absolute Spearman's rho

value across lags for each permutation. The resulting distribution was used to assess the probability

that the observed result at each lag occurred by chance. The observed and the permutation metrics

were first averaged over all triplets per monkey and then averaged over the two monkeys, to give

equal weight to both subjects.

We wished to assess whether the LJC peak lag of -0.105 s was significantly different from a lag of

zero. We did so using a jackknife method to determine the standard error of the peak position in

milliseconds (Efron, 1981). In this case, we leave out a specific triplet to assess the variability of the

peak. The jackknife procedure causes a compression of the variance (Richter et al., 2015), thus the

0.005 s sampling grid would not be sufficient to represent the peak positions of the jackknife

replications. To account for this, we cubic-spline interpolated each replication to a resolution of 1e-6 s,

which proved adequate to represent the variance of the peak. The peak of each jackknife replication

Richter et al., *Top-down beta enhances bottom-up gamma.*

was found using a Gaussian fit of the smoothed correlation as a function of lag (findpeaksG.m by T.C

O'Haver). We then derived the standard error of the estimator, and converted this to a t-score by

dividing the mean peak lag value of the jackknife replications by the estimated standard error. The

significance of this t-value was then assessed against Student's t-distribution. At the group level, this

procedure entails concatenating the data from both monkeys, and leaving out each triplet once. Based

on this group estimate of the standard error, a t-value is derived, as above, and assessed for statistical

significance.

Results

Top-Down Versus Bottom-Up Spectral Asymmetries and their Stimulus and Task Dependence

We performed electrocorticographic (ECoG) recordings from large parts of the left hemisphere in two

macaque monkeys performing a selective attention task (Fig. 1). From the ECoG recording sites, we

selected three ROIs (Fig. 1B), according to the following criteria: 1) One ROI should be a high-level

control area, one ROI a low-level visual area, and one ROI a higher-level visual area; 2) ECoG

coverage should be as good as possible; 3) ROIs should not be directly abutting, to avoid ambiguity in

boundary definition and to minimize volume conduction. These criteria led to the selection of areas 7A,

V1 and V4. The ROI pair 7A-V1 constitutes a clear top-down pathway with documented projections

from a very high-level control area to primary visual cortex (Markov et al., 2014; Bastos et al., 2015c;

Michalareas et al., 2016). The ROI pair V1-V4 constitutes a clear bottom-up pathway emerging from

V1, i.e. the area targeted by the top-down 7A-to-V1 influence. For both ROI pairs, the ECoG provided

good coverage. The central aim was to determine whether the top-down influence was modulating the

bottom-up influence.

To assess the individual frequency bands for each monkey (monkey K and monkey P), we first

computed the power spectra during visual stimulation. Area 7A shows strong beta-band peaks in both

monkeys (Fig. 2E, F, monkey K: ≈17 Hz; monkey P: ≈13 Hz). Areas V1 and V4 show gamma

frequency peaks (Fig. 2A-D, monkey K: ≈76 Hz; monkey P: ≈60 Hz). Beta activity is visible in V4 and

V1 of both monkeys at their matching peak frequencies found in area 7A. In area V4 of both monkeys,

there are distinct beta peaks. In area V1, monkey K shows a distinct beta peak, and monkey P shows

a shoulder in the power spectrum, at the respective beta frequency.

We determined the dominant interareal communication frequencies for each monkey by calculating

the pair-wise phase consistency (PPC), a frequency-resolved measure of synchronization (Vinck et

al., 2010), between the V1-V4 and V1-7A ROI pairs (Fig. 2G-J). Gamma band synchronization was

present for both ROI pairs in both monkeys with peaks at ≈76 Hz in monkey K and in a range of 58–65

Hz in monkey P. Beta peaks were present between both ROI pairs: at ≈17 Hz in monkey K and at ≈12

Hz in monkey P. For the further analyses, data from both monkeys were combined, by aligning their

individual beta and gamma peaks ±10 Hz and averaging across monkeys.

When determining individual beta and gamma frequencies, we selected the dominant peaks in the

respective frequency ranges. It has been shown that both beta and gamma frequencies show

substantial inter-individual variability, which can largely be explained by genetic factors (Vogel, 1970;

van Pelt et al., 2012). Note that while the beta peak in monkey P has its maximum at 12-13 Hz, at the

border between the alpha and beta band, the peak is strongly asymmetric with a sharp rise to the

maximal value and a slower falloff, such that most of the peak falls into the classical beta-frequency

band. This observation, together with the fact that it corresponded phenomenologically to the clear

beta peak in monkey K, led us to refer to it as beta, rather than alpha. The further analyses confirmed

that the beta rhythms in both monkeys exerted qualitatively the same effects. Yet whether they reflect

the same underlying physiological process cannot be determined on the basis of the ECoG recordings

alone. Some of the power and PPC spectra showed also a theta-band peak, which is not further

investigated, because the focus of this study is on the interaction between beta and gamma rhythms.

To demonstrate that interareal gamma-band synchronization is stimulus driven (Bosman et al., 2012;

Grothe et al., 2012), we contrasted PPC between the fixation and stimulation conditions. Figure 3A, B

shows significantly enhanced gamma-band synchronization between ROI pairs V1-V4 and 7A-V1

once the stimulus has appeared, in contrast to an almost flat spectrum when no stimulus is present.

This finding is consistent with gamma-band oscillations occurring as a result of stimulus drive. In

contrast, beta-band synchronization for both ROI pairs is present already during the pre-stimulus

fixation period, suggesting an endogenous origin (Fig. 3A, B). Beta synchronization is maintained

during the stimulation period, consistent with an ongoing top-down influence.

Richter et al.*, Top-down beta enhances bottom-up gamma.*

We next assessed the dominant directionality of interareal synchronization and its attentional

modulation. We quantified directionality of synchronization by means of Granger causality (GC)

(Granger, 1969; Ding et al., 2006; Bressler and Seth, 2011). As shown by Bastos et al. (2015c), and

extended to humans by Michalareas et al. (2016), the top-down beta-band influence of area 7A to V1

is significantly greater than the bottom-up beta-band influence of V1 to 7A (Fig. 3E). This top-down

beta-band influence is significantly increased when attention is directed to the visual hemifield

contralateral to the recording grid (Fig. 3C), consistent with an earlier report (Bastos et al., 2015c).

Between V1 and V4, the gamma-band influence is stronger in the bottom-up direction from area V1 to

V4 (Fig. 3F). The bottom-up gamma-band influence of V1 to V4 was significantly increased with

attention (Fig. 3D).

Note that the beta-band GC influence between V1 and V4 was stronger in the bottom-up than top-

down direction (Fig. 3F), in contrast to what has been reported for the same dataset (Bastos et al.,

2015c). This is due to differences in preprocessing between the present and the previous study. It

shows that for this particular pair of areas, the beta-band directionality is not robustly determined by

their hierarchical relationship. However, the difference in preprocessing did not affect other area pairs,

leaving the overall pattern intact, in which gamma-band influences are stronger in the bottom-up

direction and beta-band influences are stronger in the top-down direction. This overall pattern has also

been confirmed in an independent dataset from 43 human subjects (Michalareas et al., 2016). Note

that also in this large human dataset, the beta-band influence between V1 and V4 does not show an

unequivocal directionality, despite the overall pattern across all area pairs.

Top-Down Beta-Band Influences Correlate with Bottom-Up Gamma-Band Influences

Spontaneous endogenous increases in 7A-to-V1 beta GC may lead to increases in stimulus-driven

V1-to-V4 gamma GC. A resulting correlation between epoch-by-epoch fluctuations in the two GC

influences is difficult to quantify, because GC influences are not defined per single data epoch (without

substantially sacrificing spectral resolution and/or signal-to-noise ratio). To surmount this problem, we

used a jackknife approach. A jackknife replication (JR) of all epochs consists of all epochs except one,

that is, an all-but-one subsample of the epochs. There are as many JRs as there are epochs, because

each epoch can be left out once. If the GC in a given JR is high, this reflects a low GC in the left-out

epoch, and vice versa. This approach allows to median split epochs according to GC, and to sort and

Richter et al., *Top-down beta enhances bottom-up gamma.*

bin epochs according to GC. Ultimately, it also allows to calculate the correlation between epoch-wise

top-down and bottom-up GC, a technique that we recently described and named Jackknife Correlation

(JC) (Richter et al., 2015).

We first investigate this for one example triplet of recording sites from monkey K. Data epochs were

median split based on the GC JRs. When we select the epochs in which V1-to-V4 gamma GC is

weak, 7A-to-V1 beta GC is almost absent; by contrast, when we select the epochs in which V1-to-V4

gamma GC is strong, 7A-to-V1 beta GC shows a pronounced peak (Fig.4A, left panel). A similar

dependency exists in the other direction. When we select epochs with weak 7A-to-V1 beta GC, V1-to-

V4 gamma GC is small; when we select epochs with strong 7A-to-V1 beta GC, V1-to-V4 gamma GC

is much larger (Fig. 4A, right panel). These results indicate a relationship between 7A-to-V1 beta GC

and V1-to-V4 gamma GC.

We next asked whether this relationship holds when we move from the coarse median split to finer

and finer bins, and calculate the Spearman rank correlation across bins. We sorted epochs according

to 7A-to-V1 beta GC JRs and binned them into 5, 10, 50 or 100 bins. Per bin, the 7A-to-V1 beta GC

JRs and the V1-to-V4 gamma GC JRs were averaged, and the correlation between them was

calculated across bins. This analysis revealed that the relationship found for the median split indeed

held for finer bins (Fig. 4B). Finally, we used JC to base the correlation analysis on single epochs,

which confirmed the relationship even at this most fine-grained level (Fig. 4B, rightmost bars). The

finer the binning, the lower the correlation value, and the lower the p-value, that is, the more significant

the correlation. This dependence of correlation and p-value on bin size has been previously described

as a general consequence of binning (Richter et al., 2015). Essentially, the widely-used sorting-and-

binning approach, through averaging observations per bin and thereby removing noise, leads to

dramatic increases in correlation values; yet this comes at the expense of statistical power and it

precludes an inference on the epoch-by-epoch correlation, which requires the JC approach.

To investigate the frequency-specificity of this correlation, JC was calculated between all possible

combinations of top-down and bottom-up frequencies, both ranging from 1-100 Hz. This analysis

revealed a correlation between 7A-to-V1 GC and V1-to-V4 GC in the beta band and in the gamma

band (Fig. 4C, lower left and upper right quadrants). Critically, and further confirming the median-split

and the sorting-and-binning analysis, 7A-to-V1 beta GC shows a significant positive correlation with

Richter et al.*, Top-down beta enhances bottom-up gamma.*

V1-to-V4 gamma GC (Fig. 4C, lower right quadrant). The peak of this cross-frequency interaction is

well aligned with the 7A-to-V1 beta and V1-to-V4 gamma GC peak frequencies (Figure 4C, dashed

lines). There is no significant JC between 7A-to-V1 gamma and V1-to-V4 beta GC, even though

7A-to-V1 gamma GC is significantly correlated to V1-to-V4 gamma GC, and 7A-to-V1 beta GC is

significantly correlated to V1-to-V4 beta GC.

We then repeated these analyses for all possible triplets (Fig. 5). Figure 5A, B, C uses the same

format as Figure 4A, B, C, but shows the grand average over all triplets and the two monkeys after

alignment by their respective top-down beta, and bottom-up gamma peak frequencies. The pattern of

results found for the example triplet held in the grand average, even though average effect size was

smaller. Figure 5D shows the probability distribution across triplets of JC values between 7A-to-V1

beta GC and V1-to-V4 gamma GC, averaged over monkeys. The distribution shows a positive skew

and greater mass above zero, indicating a greater number and magnitude of positive correlations,

though with a substantial number of negative correlations, which partially accounts for the low average

JC value.

To determine whether greater GC values resulted in greater JC values, we weighted the JC value of

each triplet by the product of the respective triplet's top-down beta and bottom-up gamma GC values,

followed by averaging over triplets. This resulted in an increase of the mean JC value from 0.006 to

0.014, that is, a ≈230% increase (p<0.001, two-tailed non-parametric randomization test). This

indicates that site pairs with larger GC magnitudes, which are less susceptible to the influence of

noise, give rise to higher correlation coefficients.

Next, we tested whether the correlation was affected by attentional state. We calculated the JC

between 7A-to-V1 beta GC and V1-to-V4 gamma GC, separately for the attend-contra and attend-ipsi

conditions. The JC was significant for each attention condition separately (p<0.001). Selective

attention to the right hemifield stimulus, activating the recorded left hemisphere, enhanced JC values

by 17% (attend-contra: 0.007; attend-ipsi: 0.006, p=0.006, two-tailed paired non-parametric

randomization test). This effect is likely related to the observed increase in JC, when triplets are

weighted by GC values, because attention increased both 7A-to-V1 beta GC and V1-to-V4 gamma

GC (Fig. 3 C, D).

Richter et al.*, Top-down beta enhances bottom-up gamma.*

The sorting-and-binning analysis showed that the JC increases substantially when 7A-to-V1 GC and

V1-to-V4 GC are averaged per bin, thereby removing noise across individual triplets within bins. The

bin-wise averaging precludes an inference on the epoch-by-epoch correlation. Therefore, we explored

an alternative approach to reduce noise, while retaining an epoch-by-epoch correlation. Per epoch, we

averaged GC JRs over all 7A-to-V1 site pairs, and we averaged GC JRs over all V1-to-V4 site pairs.

Thus, per monkey, we constructed one triplet comprising the average 7A-to-V1 GC and the average

V1-to-V4 GC. Figure 6A, B, C shows the resulting analyses in the same format as the respective

panels in Figures 4 and 5. After reduction of noise across site pairs, the pattern of results was

confirmed, yet with substantially increased correlation values. Importantly, this also holds for the

epoch-wise JC values shown in the rightmost bars of Figure 6B and in Figure 6C. The JC value

between 7A-to-V1 beta GC and V1-to-V4 gamma GC increased from 0.006 in the average across

individual triplets to 0.12, that is, it showed a 20-fold increase. This is a striking demonstration of the

utility of spatial averaging for exposing the epoch-wise correlation. When we additionally give up the

epoch-wise calculation and smooth the GC values further through binning of epochs, correlations

increase even more and exceed values of 0.6 for a typical 10-bin scheme (Fig. 6 B).

To investigate whether V1-to-V4 gamma GC is truly dependent on 7A-to-V1 beta GC influences,

rather than merely on 7A beta power, we stratified for beta power. We applied this to the median split

analysis, because it lends itself best to stratification. As before, we median split epochs based on 7A-

to-V1 beta GC JRs. The 7A power spectra aligned to the monkeys' individual beta peak frequencies

showed that stronger 7A-to-V1 beta GC was in fact related to higher 7A beta power (Fig. 6D, left

panel, solid lines). This power difference is fully in line with the hypothesis that the top-down beta-

band influence is generated in 7A. Power differences were sufficiently small, such that stratification

rendered the power spectra almost identical (Fig. 6D, left panel, dashed lines). After stratification, V1-

to-V4 gamma-band GC values remained almost unchanged compared to the values before

stratification (compare right panels of Fig. 6A and D). Thus, V1-to-V4 gamma GC depended not on 7A

power differences, but rather on actual 7A-to-V1 GC influences through synchronization.

Spatial Resolution of the Correlation between Top-Down and Bottom-Up Influences

We next investigated whether the JC between 7A-to-V1 beta GC and V1-to-V4 gamma GC depended

on involving the same V1 site, which would demonstrate spatial specificity at the level of recording

sites. We tested this spatial specificity by pairing 7A-to-V1 beta GC to a specific V1 site, with V1-to-V4

gamma GC from a different V1 site. The distance that separated the two V1 sites was parametrically

varied. For each V1 site, 5 sets of other V1 sites were defined that fell into pre-specified distance

intervals (1 cm per interval, stepped by 2.5 mm, between 0 and 2 cm). Figure 7A shows one example

V1 site (arrow) and illustrates with 5 colored lines the five distance intervals (colored lines were slightly

displaced for illustration purposes). The average distance for each distance interval is marked with a

filled circle. Figure 7B shows the resulting JCs, averaged over triplets and monkeys, as a function of

distance. It can be seen that as the distance between the two V1 sites increases, there is a monotonic

falloff of the correlation coefficient between 7A-to-V1 beta and V1-to-V4 gamma GC. This indicates

that the physiological process linking 7A-to-V1 beta GC to V1-to-V4 gamma GC is not global, but

rather spatially specific. As a consequence, any spatially non-specific fluctuations, for example of

neuromodulators reflecting arousal fluctuations, are unlikely to cause the observed correlation.

Top-down Beta Leads Bottom-up Gamma in Time

We have established that spontaneous fluctuations in endogenous top-down beta GC are correlated

with fluctuations in stimulus-driven bottom-up gamma GC. To investigate whether the data contain

evidence in support of a causal relation, we assessed whether top-down beta GC is predictive of

subsequent bottom-up gamma GC. To accomplish this, we extended the JC by adding a temporal

dimension, similar to time-lagged cross-correlation. We compute the JC on time-frequency data,

where we systematically offset 7A-to-V1 beta GC JRs from V1-to-V4 gamma GC JRs by positive or

negative lags. We call this procedure lagged jackknife correlation (LJC). This quantifies at what time

delay between the top-down beta-band influence and the bottom-up gamma-band influence the JC

between them is largest. We computed LJC for each triplet. The red curve in Figure 8 shows the LJC

averaged over triplets and monkeys and exhibits a peak at -0.105 s, indicating that 7A-to-V1 beta GC

led V1-to-V4 gamma GC by 0.105 s ($t(10663) = -7.576$, $p \ll 0.001$, two-tailed jackknife-based t-test).

The LJC dropped off slowly, which likely reflects slow dynamics of the underlying GC influences

together with the fact that the windows used for GC influence estimation result in low-pass filtering of

GC influence time courses. For comparison, we performed the same LJC analysis between 7A-to-V1

gamma GC and V1-to-V4 beta GC. This combination had not shown a significant correlation in the

Richter et al.*, Top-down beta enhances bottom-up gamma.*

527   non-lagged case (Fig. 5C, upper left quadrant), which was confirmed for the lagged case (Fig. 8, gray

528   line).

529   Discussion

We used LFP recordings from 252-channel ECoG arrays covering large parts of the left hemispheres

of two macaques to analyze the interaction between top-down and bottom-up influences, both

quantified by Granger-causality (GC). Top-down influences were assessed between area 7a at the top

of the visual hierarchy and V1 at the bottom. Bottom-up influences were assessed between V1 and

V4, a known feedforward pathway carrying stimulus driven input. 7A-to-V1 GC showed a beta-band

peak, which did not require visual stimulation and thus was endogenously generated, which was

significantly larger in the 7A-to-V1 than the V1-to-7A direction, and which increased with selective

attention. V1-to-V4 GC showed a gamma-band peak, which was stimulus driven, which was

significantly larger in the bottom-up than the top-down direction, and which also increased with

selective attention. Jackknife Correlation between top-down beta-band influences and bottom-up

gamma-band influences revealed a positive cross-frequency interaction. This interaction was spatially

specific, as it was maximal between top-down and bottom-up interareal influences that shared the

same V1 site. Finally, top-down beta-band influences best predicted bottom-up gamma-band

influences ~0.1 s later, suggesting that the cross-frequency interaction is causal. Therefore, we

conclude that 7A-to-V1 beta-band influences enhance V1-to-V4 gamma-band influences.

Noise can affect GC estimates (Nalatore et al., 2007; Vinck et al., 2015). Fluctuating shared noise

could in principle generate correlation between GC fluctuations. Yet, a predominant source of shared

noise, volume conduction, is strongly attenuated in our signals due to bipolar derivation

(Trongnetrpunya et al., 2015). Furthermore, influences of shared noise on the two GC metrics should

typically occur nearly simultaneously and therefore, the lagged JC would peak at zero-lag, whereas

we found a significant lag of ~0.1 s. Also, local noise can affect GC estimates, yet these effects do not

necessarily explain a positive correlation between GC influences. For example, enhanced noise in a

recording site can lead to an overestimation of the GC influence onto that site, and at the same time to

an underestimation of the GC influence of that site onto other sites (Bastos and Schoffelen, 2015a). If

this applied to our V1 recording sites, fluctuations in their noise level would artefactually reduce the

Richter et al.*, Top-down beta enhances bottom-up gamma.*

observed correlation and thereby lead to an underestimation of the true correlation value. Finally, the

observed increase in average JC after weighting site triplets by the GC values of their constituent site

pairs argues against a confounding role of noise. The effect of noise on GC is larger for weaker GC.

Thus, if noise had generated the observed JC, more weight for strong-GC triplets would have reduced

the average JC. By contrast, we observed an increased average JC, which is consistent with our

interpretation of actual top-down influences affecting actual bottom-up influences.

Additional alternative scenarios concern physiological modulatory effects that act independently on

both, 7A-to-V1 beta GC and V1-to-V4 gamma GC, without a direct causal link between the two GC

influences. Those interareal GC influences can probably only be affected at the circuits containing the

respective projection neurons and at the circuits containing their postsynaptic target neurons, that is,

not at the corresponding interareal axonal projections. Thus, to explain our observations, such

modulatory effects would need to modify 7A circuits to enhance their beta outflow, and 0.1 s thereafter

modify V1 circuits to enhance their gamma outflow. Alternatively, they would need to modify V1

circuits to enhance their beta susceptibility, and 0.1 s thereafter modify V4 circuits to enhance their

gamma susceptibility. Additional mechanisms would be necessary to explain the observed spatial

specificity. Thus, these alternative interpretations require complex sets of assumptions. Certainly,

widespread or even global neuromodulatory fluctuations, as potentially associated with arousal

fluctuations, cannot explain the observed pattern of results.

The most parsimonious interpretation seems to be the following: The 7A-to-V1 beta-band influence

onto a given V1 site enhances with 0.1 s delay that site's V1-to-V4 gamma influence. We propose that

this cross-frequency interaction constitutes a mechanism for spatially selective attention. If correct, this

entails that the top-down control of selective attention corresponds fully or partly to top-down beta-

band influences, and the ensuing preferential bottom-up routing of the attentionally selected stimulus

corresponds fully or partly to bottom-up gamma-band influences. Thus, epoch-by-epoch fluctuations in

spatially selective attention could well generate the observed correlations. In fact, we would like to

identify the epoch-by-epoch GC fluctuations that are the basis for the observed correlations with

epoch-by-epoch fluctuations of spatially selective attention. Several recent studies have revealed that

attention samples stimuli rhyhthmically, with a predominant sampling rhythm in the theta and/or alpha

range (7-13 Hz), which is multiplexed across the sampled stimuli (Landau and Fries, 2012; Fiebelkorn

Richter et al*., Top-down beta enhances bottom-up gamma.*

et al., 2013; Holcombe and Chen, 2013; Fries, 2015; Landau et al., 2015; VanRullen, 2016). Future

studies will need to investigate whether the correlations described here are specifically driven by those

attentional sampling rhythms.

Numerous studies in visual cortex have reported gamma-band synchronization within and between

visual areas (Gray and Singer, 1989; Engel et al., 1991; Kreiter and Singer, 1996; Tallon-Baudry et al.,

1996; Fries et al., 1997; Fries, 2001; Bichot et al., 2005; Taylor et al., 2005; Hoogenboom et al., 2006;

Womelsdorf et al., 2006; Wyart and Tallon-Baudry, 2008), and numerous studies in parietal cortex

have reported beta-band synchronization within parietal areas and between parietal and frontal areas

(Buschman and Miller, 2007; Salazar et al., 2012; Dotson et al., 2014; Stetson and Andersen, 2014).

Recent ECoG recordings covering both visual and parietal areas revealed that interareal beta-band

influences predominate in the top-down and interareal gamma-band influences predominate in the

bottom-up direction (Bastos et al., 2015c). These findings link parietal beta-band activity with visual

gamma-band activity and suggest a concrete case of cross-frequency interaction (Bressler and

Richter, 2015). In the present paper, we have tested some of the resulting predictions and found direct

experimental support for such a cross-frequency interaction that allows top-down beta-band influences

to enhance bottom-up gamma-band influences.

Cortical anatomy has revealed a distinct laminar pattern of top-down and bottom-up projections

(Felleman and Van Essen, 1991; Markov et al., 2014). Bottom-up projections originate predominantly

in superficial layers, and this predominance increases with the number of hierarchical levels bridged

by the bottom-up projection. Note that an additional bottom-up pathway via the pulvinar originates in

layer 5, and it might mediate the observed V1-to-V4 GC influences in the beta band (Shipp, 2003;

Sherman, 2007; Schmid et al., 2012). Furthermore, bottom-up projections terminate predominantly in

layer 4. Top-down projections originate predominantly in deep layers, and this predominance

increases with the number of hierarchical levels bridged by the top-down projection. Furthermore, top-

down projections terminate predominantly outside layer 4, primarily in layers 1 and 6. Determining how

the respective top-down influences interact with local processing, and thereby ultimately with bottom-

up influences, remains a central neuroscientific quest. One potential mechanism has been proposed in

a model that entails details of both layer-specific anatomy and cellular biophysics (Lee et al., 2013),

and that replicates the effect of top-down selective attention on bottom-up gamma-band coherence.

Richter et al., *Top-down beta enhances bottom-up gamma.*

The model implicates a subclass of inhibitory interneurons, the slow-inhibitory (SI) interneurons, as

targets of top-down modulation. These cells may span multiple cortical laminae and thus are suitably

situated for integration of neuronal activity across layers. A subpopulation of these cells, low-threshold

spiking (LTS) cells, are found in deep layers of the cortex. In the model, LTS cells: 1) are hypothesized

to receive top-down input, 2) are implicated in the generation of beta oscillations and in a resonant

response to beta-rhythmic top-down input, 3) selectively modulate gamma-band activation in layer 2/3,

leading to an enhanced gamma band output. Our present analysis confirms the central prediction of

the Lee et al. (2013) paper, namely that specifically top-down beta-band influences enhance stimulus-

driven gamma-band processes. Lee et al. show how this mechanism can support the implementation

of attentional stimulus selection. The current results, which mechanistically link the previously reported

attentional enhancements of top-down beta and bottom-up gamma influences, provide the hitherto

missing experimental bridge. Together, experiments, modeling and model-testing data analysis have

led to an intriguingly coherent understanding of the neuronal processes behind the implementation of

attentional stimulus selection.

Richter et al*., Top-down beta enhances bottom-up gamma.*

References

Bastos AM, Schoffelen J-M (2015a) A Tutorial Review of Functional Connectivity Analysis Methods and Their Interpretational Pitfalls. Front Syst Neurosci 9:175.

Bastos AM, Usrey WM, Adams RA, Mangun GR, Fries P, Friston KJ (2012) Canonical microcircuits for predictive coding. Neuron 76:695–711.

Bastos AM, Litvak V, Moran R, Bosman CA, Fries P, Friston KJ (2015b) A DCM study of spectral asymmetries in feedforward and feedback connections between visual areas V1 and V4 in the monkey. Neuroimage 108:460–475.

Bastos AM, Vezoli J, Bosman CA, Schoffelen J-M, Oostenveld R, Dowdall JR, De Weerd P, Kennedy H, Fries P (2015c) Visual areas exert feedforward and feedback influences through distinct frequency channels. Neuron 85:390–401.

Bichot NP, Rossi AF, Desimone R (2005) Parallel and serial neural mechanisms for visual search in macaque area V4. Science 308:529–534.

Bieri KW, Bobbitt KN, Colgin LL (2014) Slow and fast γ rhythms coordinate different spatial coding modes in hippocampal place cells. Neuron 82:670–681.

Bisley JW, Goldberg ME (2003) Neuronal Activity in the Lateral Intraparietal Area and Spatial Attention. Science 299:81–86.

Bosman CA, Schoffelen J-M, Brunet N, Oostenveld R, Bastos AM, Womelsdorf T, Rubehn B, Stieglitz T, De Weerd P, Fries P (2012) Attentional Stimulus Selection through Selective Synchronization between Monkey Visual Areas. Neuron 75:875–888.

Bressler SL, Richter CG (2015) Interareal oscillatory synchronization in top-down neocortical processing. Curr Opin Neurobiol 31:62–66.

Bressler SL, Seth AK (2011) Wiener-Granger causality: a well established methodology. Neuroimage 58:323–329.

Brunet N, Bosman CA, Vinck M, Roberts M, Oostenveld R, Desimone R, De Weerd P, Fries P (2014a) Stimulus repetition modulates gamma-band synchronization in primate visual cortex. Proc Natl Acad Sci USA 111:3626–3631.

Brunet N, Vinck M, Bosman CA, Singer W, Fries P (2014b) Gamma or no gamma, that is the question. Trends Cogn Sci (Regul Ed) 18:507–509.

Brunet N, Bosman CA, Roberts M, Oostenveld R, Womelsdorf T, De Weerd P, Fries P (2015) Visual Cortical Gamma-Band Activity During Free Viewing of Natural Images. Cereb Cortex 25:918–926.

Buschman TJ, Miller EK (2007) Top-down versus bottom-up control of attention in the prefrontal and posterior parietal cortices. Science 315:1860–1862.

Colgin LL, Denninger T, Fyhn M, Hafting T, Bonnevie T, Jensen O, Moser M-B, Moser EI (2009) Frequency of gamma oscillations routes flow of information in the hippocampus. Nature 462:353–357.

Richter et al., *Top-down beta enhances bottom-up gamma.*

Dhamala M, Rangarajan G, Ding M (2008) Analyzing information flow in brain networks with
nonparametric Granger causality. Neuroimage 41:354–362.

Ding M, Chen Y, Bressler SL (2006) Granger Causality: Basic Theory and Application to
Neuroscience Schelter B, Winterhalder M, Timmer J, eds. Handbook of Time Series
Analysis: Recent Theoretical Developments and Applications.

Dotson NM, Salazar RF, Gray CM (2014) Frontoparietal correlation dynamics reveal
interplay between integration and segregation during visual working memory. 34:13600–
13613.

Efron B (1981) Nonparametric estimates of standard error: The jackknife, the bootstrap and
other methods. Biometrika.

Efron B, Tibshirani RJ (1994) An Introduction to the Bootstrap. Boca Raton: CRC Press.

Engel AK, Kreiter AK, König P, Singer W (1991) Synchronization of oscillatory neuronal
responses between striate and extrastriate visual cortical areas of the cat. Proc Natl Acad Sci
USA 88:6048–6052.

Felleman D, Van Essen DC (1991) Distributed hierarchical processing in the primate cerebral
cortex. Cereb Cortex 1:1–47.

Fiebelkorn IC, Saalmann YB, Kastner S (2013) Rhythmic Sampling within and between
Objects despite Sustained Attention at a Cued Location. Curr Biol 23:2553–2558.

Fontolan L, Morillon B, Liegeois-Chauvel C, Giraud A-L (2014) The contribution of frequency-
specific activity to hierarchical information processing in the human auditory cortex. Nat
Commun 5:4694.

Fries P (2001) Modulation of Oscillatory Neuronal Synchronization by Selective Visual
Attention. Science 291:1560–1563.

Fries P (2015) Rhythms for Cognition: Communication through Coherence. Neuron 88:220–
235.

Fries P, Roelfsema PR, Engel AK, König P, Singer W (1997) Synchronization of oscillatory
responses in visual cortex correlates with perception in interocular rivalry. Proc Natl Acad Sci
USA 94:12699–12704.

Gilbert CD, Sigman M (2007) Brain states: top-down influences in sensory processing.
Neuron 54:677–696.

Granger C (1969) Investigating causal relations by econometric models and cross-spectral
methods. Econometrica 37:424–438.

Gray CM, Singer W (1989) Stimulus-specific neuronal oscillations in orientation columns of
cat visual cortex. Proc Natl Acad Sci USA 86:1698–1702.

Grothe I, Neitzel SD, Mandon S, Kreiter AK (2012) Switching neuronal inputs by differential
modulations of gamma-band phase-coherence. 32:16172–16180.

Hilgetag CC, O'Neill MA, Young MP (1996) Indeterminate organization of the visual system.
Science 271:776–777.

Richter et al., *Top-down beta enhances bottom-up gamma.*

Holcombe AO, Chen W-Y (2013) Splitting attention reduces temporal resolution from 7 Hz for
tracking one object to <3 Hz when tracking three. J Vis 13:12–12.

Hoogenboom N, Schoffelen J-M, Oostenveld R, Parkes LM, Fries P (2006) Localizing human
visual gamma-band activity in frequency, time and space. Neuroimage 29:764–773.

Jia X, Tanabe S, Kohn A (2013) Gamma and the Coordination of Spiking Activity in Early
Visual Cortex. Neuron 77:762–774.

Kreiter AK, Singer W (1996) Stimulus-dependent synchronization of neuronal responses in
the visual cortex of the awake macaque monkey. 16:2381–2396.

Landau AN, Fries P (2012) Attention Samples Stimuli Rhythmically. Curr Biol 22:1000–1004.

Landau AN, Schreyer HM, van Pelt S, Fries P (2015) Distributed Attention Is Implemented
through Theta-Rhythmic Gamma Modulation. Curr Biol 25:2332–2337.

Lee JH, Whittington MA, Kopell NJ (2013) Top-down beta rhythms support selective attention
via interlaminar interaction: a model. PLoS Comput Biol 9:e1003164.

Lewis CM, Bosman CA, Womelsdorf T, Fries P (2016) Stimulus-induced visual cortical
networks are recapitulated by spontaneous local and interareal synchronization. Proc Natl
Acad Sci USA 113:E606–E615.

Maris E, Oostenveld R (2007) Nonparametric statistical testing of EEG- and MEG-data. J
Neurosci Methods 164:177–190.

Markov NT, Vezoli J, Chameau P, Falchier A, Quilodran R, Huissoud C, Lamy C, Misery P,
Giroud P, Ullman S, Barone P, Dehay C, Knoblauch K, Kennedy H (2014) Anatomy of
hierarchy: feedforward and feedback pathways in macaque visual cortex. J Comp Neurol
522:225–259.

Michalareas G, Vezoli J, van Pelt S, Schoffelen J-M, Kennedy H, Fries P (2016) Alpha-Beta
and Gamma Rhythms Subserve Feedback and Feedforward Influences among Human
Visual Cortical Areas. Neuron 89:384–397.

Nichols TE, Holmes AP (2002) Nonparametric permutation tests for functional neuroimaging:
a primer with examples. Hum Brain Mapp 15:1–25.

Oostenveld R, Fries P, Maris E, Schoffelen J-M (2011) FieldTrip: Open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. Comput Intell
Neurosci 2011:156869.

Paxinos G, Huang X, Toga AW (1999) The Rhesus Monkey Brain in Stereotaxic
Coordinates. San Diego: Academic Press.

Percival DB, Walden AT (1993) Spectral Analysis for Physical Applications. Cambridge
University Press.

Pinotsis DA, Brunet N, Bastos A, Bosman CA, Litvak V, Fries P, Friston KJ (2014) Contrast
gain control and horizontal interactions in V1: A DCM study. Neuroimage 92:143–155.

Richter CG, Thompson WH, Bosman CA, Fries P (2015) A jackknife approach to quantifying
single-trial correlation between covariance-based metrics undefined on a single-trial basis.

Richter et al., *Top-down beta enhances bottom-up gamma.*

Neuroimage 114:57–70.

Rohlfing T, Kroenke CD, Sullivan EV, Dubach MF, Bowden DM, Grant KA, Pfefferbaum A
(2012) The INIA19 Template and NeuroMaps Atlas for Primate Brain Image Parcellation and
Spatial Normalization. Front Neuroinform 6:27.

Rubehn B, Bosman C, Oostenveld R, Fries P, Stieglitz T (2009) A MEMS-based flexible
multichannel ECoG-electrode array. J Neural Eng 6:036003.

Salazar RF, Dotson NM, Bressler SL, Gray CM (2012) Content-specific fronto-parietal
synchronization during visual working memory. Science 338:1097–1100.

Schmid MC, Singer W, Fries P (2012) Thalamic Coordination of Cortical Communication.
Neuron 75:551–552.

Sherman SM (2007) The thalamus is more than just a relay. Current Opinion in Neurobiology
17:417–422.

Shipp S (2003) The functional logic of cortico–pulvinar connections. Philos Trans R Soc Lond
B Biol Sci 358:1605–1624.

Sirota, A., Montgomery, S., Fujisawa, S., Isomura, Y., Zugaro, M., and Buzsaki, G.
(2008) Entrainment of neocortical neurons and gamma oscillations by the hippocampal theta
rhythm. Neuron 60:683-697.

757
Stetson C, Andersen RA (2014) The parietal reach region selectively anti-synchronizes with
dorsal premotor cortex during planning. 34:11948–11958.

Tallon-Baudry C, Bertrand O, Delpuech C, Pernier J (1996) Stimulus specificity of phase-
locked and non-phase-locked 40 Hz visual responses in human. 16:4240–4249.

Taylor K, Mandon S, Freiwald WA, Kreiter AK (2005) Coherent oscillatory activity in monkey
area v4 predicts successful allocation of attention. Cereb Cortex 15:1424–1437.

Thomson DJ (1977) Spectrum Estimation Techniques for Characterization and Development
of WT4 Waveguide-I. Bell System Technical Journal 56:1769–1815.

Thomson DJ (1982) Spectrum estimation and harmonic analysis. Proceedings of the IEEE
70:1055–1096.

Trongnetrpunya A, Nandi B, Kang D, Kocsis B, Schroeder CE, Ding M (2015) Assessing
Granger Causality in Electrophysiological Data: Removing the Adverse Effects of Common
Signals via Bipolar Derivations. Front Syst Neurosci 9:189.

Van Essen DC (2012) Cortical cartography and Caret software. Neuroimage 62:757–764.

van Kerkoerle T, Self MW, Dagnino B, Gariel-Mathis M-A, Poort J, van der Togt C,
Roelfsema PR (2014) Alpha and gamma oscillations characterize feedback and feedforward
processing in monkey visual cortex. Proc Natl Acad Sci USA 111:14332–14341.

van Pelt S, Boomsma DI, Fries P (2012) Magnetoencephalography in twins reveals a strong
genetic determination of the peak frequency of visually induced γ-band synchronization. J
Neurosci 32:3388–3392.

Richter et al., *Top-down beta enhances bottom-up gamma.*

VanRullen R (2016) Perceptual Cycles. Trends Cogn Sci (Regul Ed) 20:723–735.

Vinck M, Huurdeman L, Bosman CA, Fries P, Battaglia FP, Pennartz CMA, Tiesinga PH
(2015) How to detect the Granger-causal flow direction in the presence of additive noise?
Neuroimage 108:301–318.

Vinck M, van Wingerden M, Womelsdorf T, Fries P, Pennartz CMA (2010) The pairwise
phase consistency: a bias-free measure of rhythmic neuronal synchronization. Neuroimage
51:112–122.

Vogel F (1970) The genetic basis of the normal human electroencephalogram (EEG).
Humangenetik 10:91–114.

von Stein A, Sarnthein J (2000) Different frequencies for different scales of cortical
integration: from local gamma to long range alpha/theta synchronization. Int J Psychophysiol
38:301–313.

Welch P (1967) The use of fast Fourier transform for the estimation of power spectra: A
method based on time averaging over short, modified periodograms. IEEE Trans Audio
Electroacoust 15:70–73.

Womelsdorf T, Fries P, Mitra PP, Desimone R (2006) Gamma-band synchronization in visual
cortex predicts speed of change detection. Nature 439:733–736.

Wyart V, Tallon-Baudry C (2008) Neural dissociation between visual awareness and spatial
attention. 28:2667–2679.

Richter et al., *Top-down beta enhances bottom-up gamma.*

 ## Figure Legends

Figure 1. Behavioral task and recording locations. ***A***, The task commenced with a fixation period followed by presentation of two differently colored stimuli. The fixation point color then indicated the visual stimulus to covertly attend in either the visual hemifield ipsilateral (attend-ipsi) or contralateral (attend-contra) to the recording grid. The presentation timings for each monkey are shown as a timeline. ***B***, Recording sites for areas V1 and V2 (red), V4 (blue), and 7A (yellow) from monkey K (light gray spheres) and monkey P (black spheres), co-registered to a common macaque template.

Figure 2. Power spectra and phase locking (PPC) spectra during visual stimulation. V4 (***A, B)***, V1 (***C, D***), and 7A (***E, F***) power spectra averaged over all respective site pairs of monkey K (***A, C, E***) and P (***B, D, F***). Power values at each frequency were multiplied by that frequency value to reduce the 1/f component. (***G-J***) LFP-LFP PPC for V1-V4 (***G, H***) and V1-7A (***I, J***), for monkey K (***G, I***) and P (***H, J***), respectively. Error regions show ±1 standard error of the mean (SEM) over sites or site pairs. Frequencies from 45-55 Hz were omitted due to line-noise pollution.

Figure 3. Average phase locking (PPC) and Granger causality (GC) spectra. All spectra were first averaged over site pairs, then over monkeys after alignment to their beta and gamma peak frequencies. ***A,*** LFP-LFP PPC for 7A-V1 and ***B,*** V1-V4 during fixation (dark shading) and visual stimulation (light shading). Error regions show ±1 standard error of the mean (SEM) over site pairs, merely for illustration. Gray background shading indicates frequency regions with significant condition differences ($p < 0.05$, two-tailed non-parametric randomization test, corrected for comparisons across multiple frequencies). Inset brackets denote the minimum separation required for significance. ***C,*** 7A-to-V1 GC influence spectra and ***D***, V1-to-V4 GC influence spectra, for attend-contra (light shading) and attend-ipsi (dark shading). Same statistics conventions as ***A*** and ***B***. ***E,*** Bidirectional GC influences for 7A-V1 and ***F***, V1-V4. For 7A-V1, the 7A-to-V1 influences are shown in blue shading, and the V1-to-7A influences in dark shading. For V1-V4, V1-to-V4 influences are shown in green shading, and V4-to-V1 influences in light shading. Same statistics conventions as ***A*** and ***B***, but with statistical tests based on a non-parametric bootstrap procedure. ***G***, Top-down modulatory stream (blue arrow) from 7A-to-V1, and bottom-up feedforward path from V1-to-V4 (green arrow).

Richter et al.*, Top-down beta enhances bottom-up gamma.*

Figure 4. Example triplet: Median split, correlation across binned epochs, and JC. *A*, left panel:

7A-to-V1 GC for epochs median split by the V1-to-V4 gamma GC jackknife replications; right panel:

V1-to-V4 GC for epochs median split by the 7A-to-V1 beta GC jackknife replications. Gray background

shading indicates significant differences (p<0.05, two-tailed non-parametric randomization test,

corrected for multiple comparisons across frequencies). Inset brackets denote the minimum

separation required for significance. *B*, The correlation between 7A-to-V1 beta GC and V1-to-V4

gamma GC for the sorted data divided into 5, 10, 50, or 100 bins, and without binning. Corresponding

p-values for each correlation coefficient are shown below with a dashed line marking p=0.05. *C*, The 4

colored panels show JC for the selected 7A-to-V1-to-V4 triplet. The frequencies of 7A-to-V1 GC are

shown on the vertical axis, the frequencies of V1-to-V4 GC are shown on the horizontal axis. The

frequency ranges 1-50 Hz and 51-100 Hz are shown separately, because they required slightly

different spectral analyses (see Materials and Methods). Non-significant regions are partially masked

by white (p<0.001, two-tailed non-parametric randomization test, corrected for multiple comparisons

across both frequency axes). The line plots at the bottom show the GC spectrum for the

corresponding V1-to-V4 site pair. The line plots to the left show the GC spectrum for the

corresponding 7A-to-V1 site pair. Dashed lines mark the top-down beta GC spectral peak and the

bottom-up gamma GC spectral peak.

Figure 5. Average over all triplets and both monkeys: Median split, correlation across binned epochs,

and JC. (*A, B, C*) Same format as Figure 4A, B, C, but averaging over all triplets and both monkeys,

after aligning to their individual beta and gamma peak frequencies. *D*, Probability distribution across

triplets of JC values between 7A-to-V1 beta GC and V1-to-V4 gamma GC, averaged over monkeys.

Figure 6. Triplet based on average 7A-to-V1 GC and average V1-to-V4 GC: Median split, correlation

across binned epochs, and JC. (*A, B, C*) Same format as Figure 5A, B, C, but for a triplet formed by

the average 7A-to-V1 GC jackknife replications and the average V1-to-V4 GC jackknife replications,

per monkey, and then averaged over monkeys. *D,* Left panel: 7A beta-band power spectra after

median split by 7A-to-V1 beta GC jackknife replications. Solid lines: Before stratification; Dashed lines:

After stratification. Right panel: Same as right panel of *A*, but after stratification for 7A beta power.

Richter et al.*, Top-down beta enhances bottom-up gamma.*

Figure 7. Spatial specificity of correlation between top-down beta and bottom-up gamma GC. *A*, Inset: MRI of monkey K with the surgical trepanation and prominent sulci overlaid. The blow-up shows recording sites in 7A (yellow), V1 (red), and V4 (blue) that were analyzed. The V1 site marked by the arrow serves as an example to demonstrate the distance intervals from which another V1 site may have been chosen, which were 0 to 1 cm (magenta), 0.25 to 1.25 cm (orange), 0.5 to 1.5 cm (yellow), 0.75 to 1.75 cm (green), and 1 to 2 cm (cyan). Solid circles mark the average distance between V1 sites that were chosen for each distance interval. The particular arrangement of the distance intervals was chosen solely for the purpose of illustration. *B*, Average jackknife correlation between 7A-to-V1 beta GC and V1-to-V4 gamma GC, averaged over triplets, then over monkeys. The solid black circle shows the result for standard triplets, that is, triplets where the 7A-to-V1 beta GC and the V1-to-V4 gamma GC are connected through the same V1 site. As the distance between V1 sites increases, there is a monotonic falloff in the correlation coefficient. The mean distance between V1 sites is color-coded as solid circles matching the mean distances of the five distance intervals shown in *A*. The error region shows ±1 SEM across triplets per monkey, then averaged over monkeys.

Figure 8. Lagged jackknife correlation (LJC) analysis. *A*, The solid red curve shows the LJC between 7A-to-V1 beta GC and V1-to-V4 gamma GC, averaged over triplets, then over monkeys. The blue dashed line denotes the significance threshold (p=0.05, two-tailed non-parametric randomization test, corrected for multiple comparisons across lags). The green vertical line indicates the lag of the maximum LJC value, with ±1 SEM indicated in red. The lag of the peak LJC value was significantly different from zero (t(10663) = -7.576, p<<0.001, two-tailed). Solid gray curve: LJC between 7A-to-V1 gamma GC and V1-to-V4 beta GC, averaged over triplets, then over monkeys, which showed no significant peaks.

Richter et al., *Top-down beta enhances bottom-up gamma.*

Richter et al., Fig. 1

Richter et al., Fig. 2

Richter et al., Fig. 3

Richter et al., Fig. 4

Richter et al., Fig. 5

Richter et al., Fig. 6

A

1 cm

B

x 10^{-3}

Mean rho (peak +/− 5 Hz)

Mean distance (mm)

Richter et al., Fig. 7

Richter et al., Fig. 8