NCS1 regulates Ca^{2+}-Dependent Focal Exocytosis of Golgi-derived Vesicles to Help Phagocytic uptake in Macrophages

Nimi Vashi[1], Syed Bilal Ahmad Andrabi[$,1], Swapnil Ghanwat[$,1], Mrutyunjay Suar[2] and Dhiraj Kumar*[,1]

[1]Cellular Immunology Group, International Centre for Genetic Engineering and Biotechnology, Aruna Asaf Ali Marg, New Delhi-110067, India

and

[2]School of Biotechnology, KIIT University, Bhubaneswar-751024, India

Running title: *Role of Golgi-derived vesicles in phagocytosis*

[$] Equal contribution

*Correspondence: dhiraj@icgeb.res.in
Phone: +91-11-26742357
Fax: +91-11-26742316

Abstract

During phagocytic uptake by macrophages, role of Golgi apparatus or vesicles

derived from it has never been established. Using fluorescently tagged Mannosidase-

II, a marker for Golgi-derived vesicles, we show these vesicles are recruited during

uptake of diverse targets including latex beads, *E. coli*, *Salmonella* Typhimurium and

Mycobacterium tuberculosis in both human and mouse macrophages. The recruitment

of Mannosidase-II vesicles occurred very early during phagocytosis, which was

mediated by focal exocytosis. The focal movement of Mannosidase-II vesicles

required Ca^{2+} from both extra- and intra-cellular sources apart from PI3Kinase,

microtubules and dynamin-2. At the site of uptake voltage-gated Ca^{2+} channels help

establish a Ca^{2+}-dependent local PIP3 gradient, which guide the focal movement.

Mannosidase-II vesicles also contained Neuronal Calcium Sensor-1 (NCS1) and

resembled secretory vesicles. Depleting NCS1 blocked the recruitment of

Mannosidase-II vesicles and inhibited phagocytic uptake of diverse targets. We

propose Golgi-derived vesicles provide membrane for phagosome biogenesis and are

universally required for phagocytosis, the key innate defense function.

Introduction

Phagocytosis lies at the core of innate defense mechanisms in higher eukaryotes. The process of phagocytosis involves internalization of external particles including pathogens, cellular debris etc. into a specialized membrane bound organelle called phagosomes. The phagosome thus formed undergoes a series of fusion and fission events leading to acquisition of hydrolytic enzymes and microbicidal properties to mature into phago-lysosomes (Aderem and Underhill, 1999; Flannagan et al., 2012; Jutras and Desjardins, 2005).

It was perceived initially that the membrane for phagosomes could be solely derived from the plasma membrane (Cannon and Swanson, 1992). However in macrophages, the professional phagocytes, which can engulf objects bigger than their own size without significantly altering their function, there is no apparent decline in the membrane surface area following phagocytosis(Holevinsky and Nelson, 1998). Rather capacitance measurement experiments showed that a concomitant increase in the membrane surface area precedes resealing of the phagosome, suggesting supply of phagosome membrane from intracellular sources(Holevinsky and Nelson, 1998). Professional phagocytes are therefore expected to be under tremendous pressure to sustain the supply of the membrane so as to effectively phagocytize their targets. Consequently a variety of membrane bound organelles including the vesicles originating from recycling endosomes and lysosomes as well as ER were later shown to provide membrane for the nascent phagosome(Bajno et al., 2000; Gagnon et al., 2002; Tardieux et al., 1992).

Vesicle recruitment during phagocytosis involves exocytosis of vesicles in the vicinity of nascent phagosomes (Braun and Niedergang, 2006). Studies in the past have shown secretion of vesicular contents like lysosomal hydrolases and azurophil

granules during engulfment by macrophages and neutrophils respectively, thereby

coupling the process of phagocytosis with targeted exocytosis, a process also termed

as focal exocytosis (Bajno et al., 2000; Braun and Niedergang, 2006). Recycling

endosome marker VAMP3, a soluble N-ethylmaleimide-sensitive factor attachment

protein receptor proteins on vesicles (v-SNARE), was shown to accumulate at the

early phagosomes through focal exocytosis (Bajno et al., 2000). Moreover

Synaptotagmin V (SytV), a Ca^{2+} sensor on the recycling endosomes was shown to

regulate phagocytosis but not phagosome maturation, suggesting a key role of Ca^{2+} in

the regulation of vesicle exocytosis (Vinet et al., 2008). Similarly SytVII, a lysosome

resident Ca^{2+} sensor was reported to regulate delivery of lysosome membrane to the

nascent phagosomes (Czibener et al., 2006). Interestingly, SytVII was also shown to

regulate Ca^{2+} dependent exocytosis of lysosomes during plasma membrane repair

(Reddy et al., 2001). Recently TRPML1, a key Ca^{2+} channel in the lysosomes was

shown to regulate focal exocytosis during phagocytosis of large particles (Samie et

al., 2013). Phosphoinositide3kinase (PI3K) is yet another key regulator of

phagocytosis, and is believed to help pseudopod extension around the particles during

phagocytosis (Cox et al., 1999). However, inhibition of PI3K by wortmannin may not

limit membrane availability for phagocytosis (Cox et al., 1999). There are some

reports however suggesting the role of PI3K in recruiting the membranes from

intracellular sources (Underhill and Ozinsky, 2002). Yet another study revealed

recruitment of endoplasmic reticulum (ER) at the site of phagocytosis to provide

membrane for the newly formed phagosomes, a process that was regulated by the ER

resident SNARE protein ERS24 (Gagnon et al., 2002; Jutras and Desjardins, 2005).

Intriguingly, despite being intricately involved with the recycling endocytic network

and as one of the major sources of vesicles destined to the plasma membrane like

secretory vesicles, involvement of Golgi Apparatus (GA) or vesicles derived from GA

in the process of phagocytosis has been ruled out (Becker et al., 2005). However in an

interesting observation, fragmentation and reorganization of GA was reported during

a process of frustrated phagocytosis (Bainton et al., 1989). Interestingly majority of

the reports that conclude no direct involvement of GA during phagocytic uptake

relied on the experiments that were limited to Fcγ-Receptor mediated phagocytosis

(Beemiller et al., 2006; Braun et al., 2007; Zhang et al., 1998). Here we studied the

role of vesicles derived from the GA during phagocytic uptake. Mannosidase-II, a

medial and trans-Golgi marker was also detected within secretory granules and at the

surface of certain cell types like enterocytes, pancreatic acinar cells and goblet cells

(Velasco et al., 1993). Later, using Mannosidase-II as the marker of Golgi-derived

vesicles its recruitment to cell surface was used to track the role of these vesicles

during plasma membrane repair in mouse bone marrow derived primary macrophages

(Divangahi et al., 2009; Velasco et al., 1993). Here we show, Mannosidase-II

containing Golgi-derived vesicles are recruited during phagocytosis of a variety of

targets including inert particles (latex beads), non-pathogenic bacteria (*E. coli*) and

pathogenic bacteria (*Salmonella* Typhimurium and *Mycobacterium tuberculosis*) in

mouse (RAW264.7 and BMDMs) and human (THP-1 and U937) macrophages. The

recruitment of Mannosidase-II vesicles at the site of phagocytosis occurred through

the process of focal exocytosis. This process was dependent on voltage-gated Ca^{2+}

channels that helped establish a Ca^{2+} dependent PIP3 gradient for uptake.

Mechanistically, TGN and secretory vesicle resident protein NCS1 sensed and

triggered the movement of vesicles to help phagosome biogenesis.

Results

α-Mannosidase-II (MAN2A) localizes to Golgi Apparatus (GA) and Golgi derived vesicles

To confirm the localization of Mannosidase-II to GA in our experimental set-up, we nucleofected THP-1 human macrophages and RAW264.7 mouse macrophages with a mCherry-tagged Mannosidase-II construct [mCherry-MannII-N-10 (mCherry-MAN2A), Addgene plasmid # 55074] and visualized the cells under Nikon A1R confocal microscope (see methods). Expression of Mannosidase-II in these cells distinctly labeled the GA as shown in figure 1A. We also immuno-stained Mannosidase-II using specific antibody, which labeled the GA (Fig. 1B). Treatment of mCherry-MAN2A expressing cells with Brefeldin A, the classical inhibitor of Golgi function, resulted in the disruption and fragmentation of the GA structure (Fig 1C). In both transfected as well as antibody stained cells, in addition to the main GA we could also observe several smaller punctate structures, representing the Golgi-derived vesicles (Fig. 1A and B). In U937 cells stably expressing mRuby-tagged Beta-1,4-Galactosyltransferase (B4GALT1-mRuby), a mid-Golgi resident enzyme, no such vesicle like structures could be seen (Fig. 1D), Mannosidase-II however co-localized with B4GALT1-mRuby (Fig. 1D). These results confirmed that the Mannosidase-II construct used in this study labeled GA and Golgi-derived vesicles.

Mannosidase-II is recruited at the site of phagocytic uptake

RAW264.7 cells transfected with mCherry tagged Mannosidase-II were incubated with mouse-IgG coated latex beads (see methods) that were either unstained or stained with anti-mouse IgG-Alexa488. At 30 minutes and one-hour post-addition of beads to the macrophages, cells were fixed and visualized under the microscope. We observed the recruitment of Mannosidase-II to the sites where latex beads were

phagocytosed, forming ring like structure around the beads at many instances (Fig.

2A and Fig. S1A). Overall intensity of Mannosidase-II at the bead surface had a

median of ~900 at 30 minutes and ~1500 arbitrary units (A.U.) at 1 hour as

determined using the 3-D module tool in Imaris 7.2 (see methods, Fig 2A). To check

whether Mannosidase-II recruitment during phagocytosis was a general phenomenon

we also monitored uptake of non-pathogenic bacterium *E. coli* and pathogenic

bacteria *Salmonella* Typhimurium. In both the cases, we could see Mannosidase-II

recruited at very early stages of phagocytosis. In the case of *E. coli,* nearly 60% of the

bacteria at the site of entry showed Mannosidase-II recruitment at 15 and 30 minutes

post-infection while nearly 30% of *Salmonella* containing phagosomes showed

Mannosidase-II recruitment at 5 and 10 minutes (Fig 2B). In the case of yet another

pathogenic bacteria *Mycobacterium tuberculosis* strain H37Rv, Mannosidase-II was

recruited at 30-40% phagosomes at the time of entry (Fig. 2C). Recruitment of

Mannosidase-II at the early phagosomes in case of H37Rv was also verified by

immuno-staining against Mannosidase-II (Fig. S1B). To confirm that the

Mannosidase-II positive phagosomes indeed represented early stages of phagosomes,

we immuno-stained the mCherry-MAN2A expressing cells, that were either incubated

with mouse-IgG coated latex beads or infected with *Mtb*, with anti-transferrin

receptor (TfR) antibody. We observed significant overlap between recruited

Mannosidase-II and TfR around the internalized beads and *Mtb* (Fig. 2D and 2E). TfR

recruitment around phagocytized beads showed saturating levels with median of

around 4095 A.U., much higher than the intensity distribution of Mannosidase-II

(~1000 A.U., Fig. 2D). In case of *Mtb* too, TfR showed relatively higher co-

localization with the phagosomes (~40-60%) compared to Mannosidase-II (30-50%,

Fig. 2E). Expectedly, TfR on *Mtb* phagosomes declined rapidly from one-hour post

infection to two hours post infection (Fig. 2E). To further establish that the membrane

recruitment of Mannosidase-II during phagocytosis was an early event, we took U937

cells that stably expressed EGFP fused with a plasma membrane targeting sequence

from Neuromodulin (Liu et al., 1991). We nucleofected these cells with mCherry-

MAN2A followed by incubation with the latex beads. As shown in figure 2F,

Mannosidase-II was recruited at the site of phagocytosis where it mostly localized to

the cytosolic face of the newly forming phagosome while the latex beads were still

getting internalized. We also verified the recruitment of Mannosidase-II in mouse

bone marrow derived macrophages (BMDMs) during phagocytic uptake of *E. coli*,

thus ensuring that it was not a phenomenon restricted to only cell-lines (Fig. S1C).

Characterization of early phagosomes reveals direct involvement of secretory

vesicles derived from the GA in the phagosome biogenesis

As shown above and by others (Divangahi et al., 2009; Velasco et al., 1993),

Mannosidase-II marks both GA and Golgi-derived vesicles. In most of the fields

observed in the previous sections, we could see Mannosidase-II organized into a

separate Golgi structure, underscoring the fact that the phagosome associated

Mannosidase-II were derived from the recruitment of Golgi-derived vesicles at the

site of infection. To further confirm the selective enrichment of Mannosidase-II at the

nascent phagosomes, we purified latex beads phagosomes using sucrose density

gradient ultra-centrifugation from macrophages within 1 hour of uptake (Fig. 3A).

The phagosome preparation showed presence of expected early phagosome markers

like TfR and RAB5 (Fig. 3B). They also showed presence of previously reported

markers like VAMP3 from recycling endosomes and Calnexin from ER (Fig. 3B). At

the same time, the phagosome preparation was devoid of any mitochondrial, nuclear

or cytosolic contamination (Fig. 3B). In agreement with the microscopy data,

phagosomes also showed presence of Mannosidase-II, the marker for Golgi-derived

vesicles (Fig. 3B). In addition, we could also score the presence of another Golgi-

derived secretory vesicles marker NCS1 in the phagosomes (Fig. 3B). We verified

that there was no GA contamination in the preparation as the phagosomes were

devoid of B4GALT1 (Fig. 3B).

Trafficking of vesicles in the cells is regulated by a large number of proteins

including small GTPases like RABs, ADP-Ribosylation factors (ARFs) and SNAREs

(Chen and Scheller, 2001; Gillingham and Munro, 2007; Stenmark, 2009). In order to

understand which RABs, SNAREs or ARFs could be involved in the recruitment of

Mannosidase-II vesicles at the phagosomes we performed mass-spectrometry of the

purified latex beads phagosome preparations, specifically to identify molecules below

25kDa molecular weight (Fig. S2, see methods). We were able to identify about 290

proteins from this preparation, all of them below 25kDa molecular weight

(Supplementary table S1). We performed a gene ontology analysis on these proteins

using Amigo2.0 database to specifically see enrichment of eleven classes including

ER, GA, secretory vesicles, recycling endosomes, endocytic vesicles,

neurotransmitter release, TGN, vesicle mediated transport, phagocytosis/engulfment,

exocytosis and endosomal transport (Fig. 3C, Supplementary table S2). A large

number of proteins belonging to ER were found in the phagosomes (Fig. 3C). The

phagosome mass spectrometry data revealed presence of 17 different RABs, 5 out of

6 known ARFs and four VAMPs. Many of these proteins are involved with Golgi

apparatus, exocytosis, secretory vesicles and neurotransmitter release (Fig. 3C).

VAMP2 gets enriched in the secretory vesicles that are released from the GA and

targeted to the membranes (Faurschou and Borregaard, 2003). It also helps in the

release of neurotransmitters by regulating the exocytosis of secretory vesicles (Shen et

al., 2015). Presence of a large variety of proteins that could regulate vesicle

trafficking on the early phagosomes could potentially mean a complex interplay of

different vesicle trafficking pathways in the process of phagosome biogenesis. Since

the role of vesicles derived from recycling endosomes and involvement of ER was

already established during phagosome biogenesis, presence of markers associated

with vesicles derived from the GA reconfirmed the utility of this organelle as an

additional source of membranes for phagosome biogenesis.

Mannosidase-II positive Golgi-derived vesicles are recruited at the phagosomes

through focal exocytosis

We next wanted to understand how Mannosidase-II vesicle recruitment was regulated

during phagocytosis. It was previously reported that Mannosidase-II positive Golgi-

derived vesicles get recruited at the site of membrane damage during repair

(Divangahi et al., 2009). It is also well established that vesicular movement during

membrane repair occurs through focal exocytosis (Reddy et al., 2001). To verify the

recruitment of Mannosidase-II during membrane repair, we treated RAW264.7

macrophages expressing mCherry-MAN2A with detergent for 15 minutes to cause

membrane damage. The cells were then followed live for next 2-3 hours to assess

recruitment of Mannosidase-II at the damaged site. We could observe formation of

membrane lesion upon detergent treatment and subsequent recruitment of

Mannosidase-II that eventually sealed the lesion (Fig. S3) thereby highlighting the

targeted exocytic movement of Mannosidase-II vesicles.

One of the key regulators of focal exocytosis, specifically for post-Golgi transport

vesicles towards the plasma membrane, is dynamin (Kreitzer et al., 2000). Dynamins

are the member of large GTPase family and dynamin-2 is the ubiquitously expressed

isoform while dynamin-1 is mostly neuronal (Henley and McNiven, 1996; Warnock

et al., 1997). Their involvement in budding of transport vesicles from the Golgi is also well documented along with their role in regulating focal exocytosis (Praefcke and McMahon, 2004; Samie et al., 2013). In mCherry-MAN2A expressing RAW264.7 cells, we observed significant overlap of Mannosidase-II with dynamin-2 on the phagosomes containing either latex beads or *Mtb* (Fig. 4A and 4B). Again as in the case of TfR, dynamin-2 recruitment on the nascent phagosomes was relatively higher than Mannosidase-II recruitment for both latex beads and *Mtb* (Fig. 4A and 4B). Presence of dynamin-2 at the phagosomes was consistent with previous reports, which showed their recruitment at early phagosomes during phagocytosis (Gold et al., 1999).

We next inhibited dynamin-2 by dynasore treatment for varying period of time and observed its effect on Mannosidase-II recruitment and phagocytosis. In the case of latex beads, four hours of pre-treatment with dynasore (at 40 and 80μM) resulted in a significant inhibition in the recruitment of Mannosidase-II at the phagosomes (Fig. 4C). It also markedly reduced the phagocytosis of latex beads (Fig. 4D). Dynasore treatment also inhibited uptake of *Mtb* in THP-1 macrophages (Fig. S4A). There was noticeable decline in the recruitment of Mannosidase-II at the *Mtb* phagosomes (Fig. S4B). Dynamin assisted movement of vesicles also require microtubules (Kreitzer et al., 2000). Inhibition of microtubules by nocodazole treatment resulted in a transient decline in the uptake of *Mtb* and latex beads (Fig. S4C and D). Further, nocodazole treatment resulted in a marked decline in the recruitment of Mannosidase-II to the phagosomes containing latex beads and *Mtb* (Fig. S4E and F). These results established that the Mannosidase-II positive Golgi-derived vesicles were recruited at the phagosomes through focal exocytosis and were assisted by dynamins.

Golgi-derived vesicles co-operate with the vesicles derived from the recycling endosomes during phagocytosis

Recruitment of vesicles from endocytic origin (VAMP3 positive) and lysosomal origin (LAMP1 positive) during phagocytosis has been reported earlier (Bajno et al., 2000; Czibener et al., 2006). We also observed presence of VAMP3 on the early latex beads phagosomes as shown above (Fig. 3B). This finding was reconfirmed in mCherry-MAN2A expressing RAW264.7 cells, where both latex beads and *Mtb* very early during phagocytosis (30 minutes and 1 hour for beads; 1 hour and 2 hours for *Mtb*) showed very high recruitment of and co-localization with VAMP-3 (Fig. 4E and F). In case of *Mtb*, more than 90% of the Mannosidase-II positive phagosomes were also positive for VAMP-3 (Fig. 4F). Overall, VAMP-3 recruitment was much higher compared to Mannosidase-II recruitment at the site of phagocytosis (Fig. 4E and F). Thus spatially and temporally, Mannosidase-II recruitment at the phagosomes was strikingly similar to VAMP3 recruitment. Moreover similar to the case of Golgi-derived vesicles here, the recruitment of VAMP-3 positive vesicles from the recycling endosomes is also dependent on focal exocytosis (Bajno et al., 2000). Thus it is very likely that vesicles from different origins co-operate in aiding the biogenesis of phagosomes.

Phosohatidylinositol3Kinase (PI3K) is required for Mannosidase-II recruitment at the nascent phagosomes and phagocytosis

The role of PI3K in regulating phagocytosis is well known where its function is believed to be mostly required for pseudopod extension around the cargo during phagocytosis (Cox et al., 1999). We wanted to test whether some of the effects of PI3K on phagocytosis were due to its involvement in the recruitment of Mannosidase-II vesicles for phagocytosis. Inhibition of PI3K by wortmannin led to nearly 80%

decline in the uptake of latex beads at 30 and 60 minutes (Fig. 5A). In case of *Mtb*,

the decline in uptake was ~85% at 1 and 2 hours post-infection and 70% at 4 hours

post-infection (Fig. 5B). In the case of *Mtb* we assayed multiple doses of wortmannin

and three different MOIs (Fig. S5A). The effect of wortmannin on the uptake of *Mtb*

was dependent on the inhibitor concentration however independent of the MOIs used

(Fig. S5A). Wortmannin was equally effective in inhibiting the phagocytic uptake of

Salmonella Typhimurium in THP-1 macrophages (Fig. S5B). Curiously, PI3K

inhibition also resulted in a marked decrease in the recruitment of Mannosidase-II at

the nascent phagosomes (Fig. 5C and D). Thus at least to some extent, it seems the

effect of inhibition of PI3K on phagocytosis may be due to a reduced recruitment of

Mannosidase-II positive Golgi-derived vesicles at the phagosomes. Similar effect of

PI3K inhibition on the membrane recruitment for the newly forming phagosomes has

been discussed previously (Underhill and Ozinsky, 2002).

Extracellular Calcium (Ca^{2+}) is needed for phagocytosis and focal exocytosis of

Golgi-derived vesicles at the site of phagocytosis

We next wanted to understand the immediate early mediators of focal exocytosis

during phagocytosis. Focal exocytosis of vesicles derived from the recycling

endosomes as well as lysosomes depends on the functioning of key Ca^{2+} sensors in

these compartments (Czibener et al., 2006; Samie et al., 2013; Vinet et al., 2008). It

was therefore imperative to test the role of Ca^{2+} in the focal movement of Golgi-

derived vesicles. Release of Ca^{2+} from intracellular stores is one of the key signaling

events during the course of phagocytosis and downstream maturation of the

phagosomes (Koul et al., 2004). Expectedly, phagocytosis of both latex beads and

H37Rv in THP-1 macrophages was severely hampered in the presence of TMB-8

(Fig. 5E and F), an inhibitor of the IP3 receptor that serves as the Ca^{2+} release channel

from intracellular stores like ER upon binding with IP3 (Singh et al., 2005).

Intriguingly, presence of Ca^{2+} chelator EGTA in the extracellular media also inhibited

phagocytic uptake of both latex beads and H37Rv in THP-1 macrophages (Fig. 5E

and F) thereby suggesting the involvement of extracellular Ca^{2+} in the phagocytic

uptake. However it could simply imply the effect of capacitative Ca^{2+} influx, which

typically requires the CRAC channels and happens immediately after the intracellular

stores are exhausted of Ca^{2+} (Singh et al., 2005). Treatment with both TMB-8 and

EGTA had additive effect on the uptake of latex beads but not on the uptake of *Mtb*

by THP-1 macrophages (Fig. 5E and F). We also noted a much more pronounced and

sustained effect of TMB-8 and EGTA+TMB8 (~70% and ~85% inhibition

respectively, Fig. 5E) on the uptake of latex beads as against that of *Mtb* (~25% and

~45% inhibition respectively, Fig. 5F). As observed in the case of wortmannin

treatment, inhibition of Ca^{2+} signaling also resulted in a marked decline in the

recruitment of Mannosidase-II at the nascent phagosomes (Fig. 5C and D).

Entry of extracellular Ca^{2+} through voltage-gated Ca^{2+} channels helps create the

foci for the recruitment of vesicles during phagocytosis

The role and mechanism of recruitment of extracellular Ca^{2+} during phagocytosis is

limited to the capacitative influx (Lee et al., 2003) or through passive accumulation of

Ca^{2+} in the phagosomes during phagocytosis (Lundqvist-Gustafsson et al., 2000). An

increase in the local Ca^{2+} concentration around phagosomes has been reported to

facilitate phagocytosis (Stendahl et al., 1994). Interestingly, the role of extracellular

Ca^{2+} during membrane repair process and focal exocytosis of vesicles to the damaged

site has been extensively reported (Miyake and McNeil, 1995; Steinhardt et al., 1994;

Togo et al., 1999). Similarly during neurotransmitter release in the neuronal cells,

activation of voltage-gated Ca^{2+} channels during action potential helps focal

exocytosis of secretory vesicles (Sudhof, 2012). We hypothesized that one of the

earliest signals during phagocytic uptake could consist of Ca^{2+} entry into the cells

through the voltage-gated Ca^{2+} channel. In THP-1 macrophages treated with

loperamide hydrochloride or amlodipine besylate (inhibitors of L/P-type and L –type

voltage gated Ca^{2+} channels respectively) (Church et al., 1994; Kochegarov, 2003),

phagocytosis of *E. coli* declined considerably (Fig. 6A). These treatments were also

effective against phagocytosis of *Mtb* and latex beads in THP-1 macrophages (Fig.

S6A and B). Treatment with any of these two inhibitors also blocked Mannosidase-II

recruitment at the nascent phagosomes during uptake of latex beads in THP-1

macrophages (Fig. 6B). Thus voltage-gated channel seems to play an important role

during phagocytosis of diverse targets.

We next wanted to understand whether extracellular Ca^{2+} helped decide the foci for

the recruitment of Golgi-derived vesicles during phagosome formation (Sudhof,

2012). We investigated the accumulation of PIP3 at the site of engulfment. Inhibition

of PIP3 by wortmannin treatment abrogates the uptake as observed in this study (Fig.

5) as well as reported earlier (Cox et al., 1999). PIP3 is known to interact with the

Pleckstrin Homology (PH) domain of proteins. In cells expressing AKT-PH domain

fused with mCherry (AKT-PH-mCherry), at 5 minutes post-addition of *E. coli*, we

observed, expectedly, a significant accumulation of PIP3 at the site of bacterial entry

(Fig. 6C). Also there was a gradual decline in the PIP3 levels as we move further into

the cells from the site of phagocytosis (Fig. 6C). However in cells pre-treated with the

Ca^{2+} chelator (EGTA, 3mM), the selective accumulation of PIP3 at the site of

phagocytosis was lost (Fig. 6C) and PIP3 was almost uniformly distributed across the

cell irrespective of the site of engagement with the bacteria (Fig. 6C). Surprisingly,

similar effect on PIP3 distribution was observed when these cells were pre-treated

with either loperamide or amlodipine (Fig. 6C). In both these cases, the distribution of

PIP3 in the cells was not influenced by the site of engagement with the bacteria (Fig.

6C). In cells treated with the PI3K inhibitor wortmannin, AKT-PH-mCherry was as

expected, more uniformly distributed (Fig. 6C). Quantitatively, it was evident from

the signal intensity plots (see methods) in figure 6D, that a gradient of PIP3 is

established during phagocytosis, with the highest concenration at the site of entry.

Moreover since blocking the entry of extracellular Ca^{2+} either through chelators or

inhibitors of voltage gated channels resulted in a loss of PIP3 gradient in a similar

fashion as in the case of wortmannin treatment, it was evident that the formation of

PIP3 gradient was dependent on Ca^{2+}, more liekly on the entry of Ca^{2+} from

extracellular milieu through the voltage-gated Ca^{2+} channels (Fig. 6D). Together these

results imply that the extracellular Ca^{2+}, entering through a voltage-gated Ca^{2+}

channel helps forming the foci for the recruitment of PI3K and initiate the signaling

cascade for phagocytic uptake in the macrophages.

Mannosidase-II positive early phagosomes are also NCS1 positive

Having established the role of Ca^{2+} in regulating the movement of Mannosidase-II

vesicles during phagocytosis, we next wanted to understand how the focal movement

of Golgi-derived vesicles is triggered in phagocytosing macrophages. NCS1 is an EF-

hand motif containing protein, originally believed to exclusively express in the

neuronal cells (Haynes et al., 2005). Subsequently it was shown to get expressed in a

variety of cell lines as TGN resident Ca^{2+} interacting protein (Haynes et al., 2005).

NCS1 was reported to be involved in the recruitment of Golgi-derived vesicles during

plasma membrane repair in macrophages and was integral to these vesicles (Behar et

al., 2010; Divangahi et al., 2009). In cells expressing mCherry-MAN2A we stained

for NCS1 during uptake of latex beads or Mtb. In both the cases, NCS1 co-localized

with the cargo (Fig. 7A, Fig. S7A). In the case of Mtb, nearly all Mtb containing

phagosomes were also positive for NCS1 whereas some NCS1 positive phagosomes

did not contain Mannosidase-II (Fig. 7A). Similarly for beads, NCS1 intensity was

much higher compared to Mannosidase-II intensity (Fig. 7B). Moreover, we observed

NCS1 to co-localize with Mannosidase-II in vesicles that were not part of the

phagosome (Fig. S7B), corroborating with the fact that both Mannosidase-II and

NCS1 are components of the GA secretory vesicles derived from the GA (Behar et

al., 2010; Divangahi et al., 2009).

Depleting Neuronal Calcium Sensor 1 (NCS1) inhibits focal exocytosis of

MAN2A vesicle during phagocytosis

To test whether Ca^{2+} dependent trigger of vesicular trafficking relied on the ability of

NCS1 activation upon Ca^{2+} binding, we compared the recruitment of Mannosidase-II

at the nascent phagosome during phagocytosis of *E. coli* in THP-1 macrophages that

were either treated with scrambled siRNA control or NCS1 specific siRNA (Fig.

S7C). Knocking down NCS1 resulted in more than 50% decline in the recruitment of

Mannosidase-II at the early *E. coli* phagosomes (Fig. 7C). Expectedly, knockdown of

NCS1 in THP-1 macrophages resulted in a marked decline in the uptake of latex

beads, *E. coli*, *Salmonella* and *Mtb* (Fig. 7D). True to all other treatments, which

inhibited the phagocytic uptake, NCS1 siRNA knockdown also had both quantitative

and kinetic effects on the uptake of all of the targets except in *Salmonella* where the

effects were persistent (Fig. 7D). Thus uptake in case of latex beads was ~35, 60 and

85% of the control in the siRNA treated sets at 30 minutes, 1 hours and 2 hours

respectively (Fig. 7D). For *E. coli* these numbers were 8, 10 and 22% of control at 5,

15 and 30 minutes (Fig. 7D). In case of *Salmonella*, uptake was about 65% of the

control set at 15 and 30 minutes while in case of *Mtb*, the relative uptake in the NCS

knockdown cells was 40, 45 and 55% of control at 1 hour, 2 hours and 4 hours

respectively (Fig. 7D).

Discussions

Professional phagocytes like macrophages require continuous supply of membrane in order to form phagosomes around the phagocytosed particles (Vicker, 1977). It is now understood that cellular compartments like recycling endosomes, lysosomes and ER could supply membrane for the nascent phagosomes (Bajno et al., 2000; Tardieux et al., 1992)ʼ (Gagnon et al., 2002). Adding to the existing pool of membrane sources available for phagosome formation, in this study we show recruitment of Golgi-derived vesicles at the site of phagocytosis in macrophages. However unlike previous reports, where the studies were mostly restricted to either latex beads or select organism, we show here a more universal requirement of the Golgi-derived vesicles during phagocytosis by macrophages using inert particles, non-pathogenic bacteria and two different pathogenic bacterial species as the cargo. It is important to note here that GA serves as the origin for most vesicles in the cell, including recycling endosomes, endocytic machinery and vesicles destined to the plasma membrane for exocytosis (Bonifacino and Glick, 2004). Bacterial pathogens once inside the host cells further subvert the membrane trafficking pathways to ensure prolonged survival and escape from innate defense mechanisms(Alix et al., 2011). Interestingly GA serves as the hub for trafficking inside the cells (Gillingham and Munro, 2016). It therefore may make strong sense that GA gets involved and alerted of incoming pathogen while the cell has just started to engulf it. This line of investigation seems extremely fascinating at present as it may open a new understanding in the functioning of innate immune system.

Role of Golgi-derived vesicles in supplying the membrane at plasma membrane is well known at least in one context - the membrane repair pathway (Divangahi et al., 2009). Using Mannosidase-II as specific marker for Golgi-derived vesicles they

showed participation of GA and lysosomes in the repair process through targeted

exocytosis (Divangahi et al., 2009). The movement of vesicles during phagocytosis

from the recycling endosomes and lysosomes was also reported to follow a similar

targeted exocytosis where vesicles are directed towards the phagocytic cup (Bajno et

al., 2000; Niedergang and Chavrier, 2004; Samie et al., 2013). Interestingly more than

90% of Mannosidase-II positive phagosomes at the site of entry were also found to be

positive for VAMP-3 and dynamin-2. While VAMP-3 gets recruited through focal

exocytosis, dynamin-2 is known to regulate post-Golgi transport of vesicles and

facilitate focal exocytosis (Samie et al., 2013). Dynamins have earlier been implicated

in regulating the focal exocytosis in phagocytosing macrophages (Di et al., 2003;

Gold et al., 1999; Samie et al., 2013). Inhibition of dynamins by dynasore treatment

resulted in a reduced phagocytic uptake, which may simply occur due to inhibition of

dynamin-mediated scission of phagosomes from the membrane (Jones et al., 1998).

However dynasore treatment also resulted in reduced Mannosidase-II recruitment at

the nascent phagosomes, implying the involvement of roles dynamin plays in vesicle

budding and their focal movement towards the site of uptake. High co-localization of

Mannosidase-II with TfR further emphasized very early recruitment of Golgi-derived

vesicles. Yet another instance where vesicle exocytosis from Golgi has been

extensively studied is in the context of neurotransmitter release, which shows

remarkable similarity with the focal exocytosis during membrane repair. Thus it is

intriguing to witness the brilliance of cellular economy, where a common mechanism

could be utilized to address three entirely independent cellular requirements.

Next to understand how target recognition at the cell surface for phagocytic uptake

could trigger GA to elicit the movement of the vesicles, we took cues from membrane

repair pathway where the damage is typically sensed via entry of extracellular Ca^{2+}

into the cells (Miyake and McNeil, 1995; Steinhardt et al., 1994). We indeed observed

inhibiting either extracellular Ca^{2+} or the release of Ca^{2+} from intracellular stores

resulted in a loss of phagocytic function in the macrophages. The possibility of a

membrane breach during phagocytic uptake in macrophages has never been

discussed. Thus most plausible source for the entry of extracellular Ca^{2+} was some

membrane channels, which are classically involved at a similar step during

neurotransmitter release(Augustine, 2001; Sudhof, 2012). Experiments with

loperamide and amlodipine strongly support a critical involvement of voltage-gated

Ca^{2+} channel in the extracellular Ca^{2+} entry thereby facilitating phagocytosis.

Incidentally the role of voltage-gated channel in regulating podosome formation in the

macrophages was recently shown (Carrithers et al., 2009). Yet another channel

TRPV2 was recently shown to be important for phagocytosis in macrophages and its

absence resulted in loss of Ca^{2+} influx from the extracellular milieu and abrogated

phagocytic uptake (Link et al., 2010). Curiously macrophages lacking TRPV2 were

also defective in chemoattractant-evoked motility (Link et al., 2010). It may not be

unusual to assume that some of the inhibitory effects shown in this study were most

likely due to inhibition of TRPV channels at the plasma membrane. Therefore it

seems macrophage membrane depolarization could be a more general mechanism of

cellular functioning including phagocytosis, adherence and motility.

Phagocytosis was also dependent on the release of Ca^{2+} from intracellular stores, as

TMB8 treatment inhibited phagocytosis. A combination of EGTA and TMB8 had

more dramatic effects on phagocytosis, which also showed some sort of selectivity in

terms of the cargo. Thus, for latex beads, blocking both intra- and extra-cellular cargo

abolished their uptake by the macrophages, however in case of bacteria, the block in

uptake was more kinetic in nature. It strongly supports the possibility that Ca^{2+} may

be extremely critical for the uptake of cargos where cognate receptors are not

known/available thereby rely a lot on the membrane depolarization and associated

Ca^{2+} entry. The graded importance of Ca^{2+} during phagocytosis of diverse targets

needs further exploration for better understanding. Inhibition of Ca^{2+} also resulted in

reduced Mannosidase-II recruitment at the phagosomes.

The role of PI3Kinase in phagocytosis is well known, which supposedly helps

pseudopod extension while engulfing the target (Cox et al., 1999). However, it has

also been shown that inhibition of PI3Kinase may limit the membrane availability

(Underhill and Ozinsky, 2002). We indeed observed uptake of *Mtb*, *Salmonella* or

latex beads by macrophages were severely compromised in the presence of

wortmannin. *Salmonella* entry is known to occur via either macropinocytosis or

phagocytosis, however the former is insensitive to PI3Kinase inhibition(Drecktrah et

al., 2006). Thus the strains used in this study taken up by the macrophages thorugh

phagocytosis. Interestingly vesicle recruitment has also been reported to be important

for membrane ruffle formation, which helps in Salmonella macropinocytosis (Dai et

al., 2007). We also noted significantly reduced recruitment of Mannosidase-II vesicles

at the nascent phagosomes in the presence of wortmannin. Thus recruitment of Golgi-

derived vesicles at the nascent phagosomes indeed required PIP3. Experiments with

AKT-PH-mCherry clearly support the establishment of PIP3 gradient during

phagocytosis. Selective PIP3 enrichment has previously been shown regulating

cellular polarity, chemotaxis and pseudopod extension (Cox et al., 1999; Funamoto et

al., 2002; Ridley et al., 2003). The most dramatic observation was however the use of

voltage-gated Ca^{2+} channels by macrophages to set the foci for PIP3 accumulation,

which eventually results in all the downstream signaling and recruitments.

Interestingly, dynamins have conserved PH-domain at their C-terminal, making them

responsive to the PIP3 levels. It has been shown that mutations in the PH domain of

dynamin could result in severe defects in its key functioning like post-Golgi transport

and endocytosis (Achiriloaie et al., 1999; Zuchner et al., 2005). Thus PI3K and

dynamins seem to work together for the focal recruitment of Golgi-derived vesicles

during phagosome biogenesis.

The Ca^{2+} sensing function in the GA is attributed to the resident molecule NCS1,

which was initially discovered in the neuronal cells regulating the synaptic

transmission (Haynes et al., 2005). NCS1 was also shown as part of the secretory

vesicles (Scalettar et al., 2002). In macrophages, ablation of NCS1 resulted in a block

in the membrane-resealing pathway (Divangahi et al., 2009). We therefore

hypothesized; NCS1 could also serve as the Ca^{2+} sensor in Golgi apparatus for

triggering the exodus of the vesicles during phagocytosis. Interestingly, NCS1

knockdown not only reduced the phagocytic function of macrophages, there was also

significantly lower recruitment of Mannosidase-II at the phagosomes.

The results form purified latex beads phagosomes supported the overall observation in

this study that phagocytic uptake is a complex process and involves several players.

We deliberately focused on the molecules below 25kDa cut-off to make sure

enrichment of three key classes of molecules including RABs (~25 kDa), ARFs (~20

kDa) and VAMPs (~14 kDa). This was to ensure that the presence of other high

abundance and high molecular weight proteins did not mask these low abundant low

molecular weight proteins. We identified 17 different RABs on the phagosomes out of

which nine were associated with Golgi function. We also identified five out of six

known ARFs, where ARF1 and ARF3 were particularly important for their known

association with Golgi apparatus, phagosome/engulfment and exocytosis(Beck et al.,

2008; Faundez et al., 1997; Krauss et al., 2008). We could identify four different

VAMPs in the phagosomes including VAMP2, 3, 4 and 8. While role of VAMP3 in

phagosome biogenesis is well described, VAMP2 is particularly known for its

involvement in exocytosis, neurotransmitter release and secretory vesicles (Faurschou

and Borregaard, 2003; Shen et al., 2015). Previously in a proteomic study of the latex-

beads containing phagosome, VAMP4 was identified as phagosome membrane

associated protein (Shui et al., 2008). VAMP4 are trans-Golgi resident proteins and

are also involved in the immature secretory granules (Steegmaier et al., 1999). We

also found SEC22B in the phagosome MS data, an important SNARE known to be

important for the recruitment of ER during phagocytic uptake (Becker et al., 2005).

Considering the number of RABs, ARFs and VAMPs associated with the exocytosis

and secretory vesicles, it seems very likely that the true identity of the Golgi-derived

vesicles that are recruited during phagosome biogenesis could be secretory vesicles.

However the role of GA during phagocytosis was ruled out in the past since

phagocytic uptake was found insensitive to Brefeldin A (Becker et al., 2005), a fungal

metabolite, which disrupts the GA (Lippincott-Schwartz et al., 1989). Interestingly

during membrane repair process, which is also dependent on secretory vesicles from

GA in addition to other membrane sources, while the repair of first wound is typically

independent of Brefeldin A, it has been reported that repair of any subsequent wounds

are sensitive to Brefeldin A (Andrews, 2002; Togo et al., 2000). Similarly during

secretion, Brefeldin A while inhibits the formation of new secretory vesicles, does not

interfere with the membrane fusion of pre-formed vesicles (Miller et al., 1992; Rosa

et al., 1992). Thus it is possible that following exhaustion of pre-formed vesicles

during phagocytosis, uptake of any subsequent cargo encountered by the cell may

show sensitivity to BFA. This is an interesting proposition and certainly warrants

further investigations.

At the molecular level it seems SNARE mediated fusion may be involved in the

phagosome biogenesis. We were able to track the presence of syntaxin 1, the t-

SNARE at the plasma membrane involved in SNARE mediated vesicle fusion, along

with Mannosidase-II at the early phagosomes (Fig S7D). Syntaxin-1 is known to

facilitate fusion of secretory vesicles(Sollner et al., 1993). Moreover, its presence will

also be required for the fusion of VAMP3 positive vesicles, a vSNARE typically

located in the recycling endosomes. Given the multiplicity of v-SNAREs and t-

SNAREs with possibly overlapping functions makes it challenging to clearly establish

the fusion machinery involved. The identity of SNAREs involved in this process

therefore constitutes the next set of challenge following this study. Taken together the

results in this study allowed us to reconstruct the process of phagocytosis (Fig. 7E).

We note that Golgi-derived vesicles are one of the several sub-cellular vesicular pools

available for phagosome biogenesis including recycling endosomes and lysosomes. It

actually explains why in many cases except for PI3K inhibition, the effects were more

kinetic in nature. We did observe in our results cases where some phagosomes were

only positive for VAMP3 and not Mannosidase-II or vice-versa, leaving us with an

intriguing question that whether the outcome of phagocytosis could vary depending

on which organelle contributed the most for the formation of phagosome.

In conclusion, we show here an as yet unknown function of Golgi-derived vesicles

during phagocytic uptake in macrophages. The targeted exocytosis coupled with

phagocytosis has been studied in the past, however involvement of vesicles from the

GA in macrophages is unprecedented. Similarly regulation of phagocytic uptake by

NCS1 comprehensively extends the known functions of this molecule in

macrophages. Finally our finding that the voltage-gated Ca^{2+} channel could play a

role in the process of phagocytosis may be potentially harnessed in future for

developing better therapeutic interventions for various infectious diseases including
tuberculosis.

Methods

Ethics statement

The animal experiments were performed upon prior approval from the institutional

ethics committee (IAEC) of International Center for Genetic Engineering and

Biotechnology (Approval no.: ICGEB/AH/2013/03/IMM-38).

Reagents And Antibodies: The following reagents were used in this study:

Wortmannin (Sigma Aldrich,W1628), Dynasore Hydrate (Sigma Aldrich, D7693),

Nocodazole (Sigma Aldrich, M1404), TMB8 (Sigma Aldrich, T111), EGTA

(Amresco, 0732), Loperamide (Sigma Aldrich, L4762), Amlodipine besylate (Sigma

Aldrich, A5605), PKH 67 (Sigma Aldrich, MINI 26), RIPA buffer (Amresco,N653),

BSA (Sigma Aldrich, A2153), Saponin (Sigma Aldrich, 47036), Puromycin

(Invivogen, ant-pr-1), 4 μm aldehydate-sulphatelatex beads (Life Technologies,

A37304) and 1μm yellow green aldehyde latex beads (Life Technologies, F8823).

All the siRNA used in this study were siGenome siRNA SmartPool (Dharmacon Inc).

The transfection reagent used for siRNA transfections was Dharmafect-2 (Dharmacon

Inc). The primary antibodies used in this study are: Mannosidase II (abcam, ab12277),

Dynamin (Santacruz, sc-6401), VAMP3 (abcam, ab43080), Transferrin (abcam,

ab84036). The secondary antibodies used in this study are: Alexa fluor 405 and Alexa

fluor 568 conjugates from (Life Technologies). **Plasmid Constructs:** The plasmid

constructs used in this study are mCherry-ManII-N-10 (Addgene plasmid #55073),

pcDNA3.1_AktPH-mCherry (Addgene plasmid #67301), pCT- Golgi-GFP

(CYTO104-VA-1) and pCT-Mem-GFP (System Biosciences, CYTO100-PA-1).

Bacterial Culture Maintainence: Mycobacterial cultures were maintained in 7H9

media (Difco) supplemented with 10% OADC. Single cell suspension of

Mycobacterial strains were prepared by aspiration of the culture eight times with 26

gauge needle and six times 30 gauge needles. Quantification of this prepared culture

was done by taking absorbance at 600 nm wavelength (0.6 O.D. corresponds to ~100

x 10^6 bacteria). The bacteria thus appropriately calculated were added to the cells at

the mentioned MOI (Multiplicity of Infection). For microscopy experiments, the

desired number of bacteria was stained with PKH67 (Sigma Aldrich), a lipophilic

green fluorescent dye, as per the manufacturer's protocol. The stained bacteria were

then passed thrice through a 26-gauge needle and used for infection. *E. coli* and

Salmonella Typhimurium strains were maintained in LB (Difco). For infection culture

density was determined by OD at 600 for both *E. coli* and *S.*Typhimurium containing

GFP, an OD of 1= 1 x 10^8 bacteria for both these strains.

Tissue Culture: THP-1 cells (a kind gift from Dr. Dong An, UCLA) and U937 cells

(ATCC) were cultured in RPMI 1640 medium (Life Technologies) and RAW264.7

(ATCC) murine macrophages were grown in Dulbecco's modified Eagle's media

(DMEM, Life Technologies) supplemented with 10% Fetal Bovine Serum (FBS,

GIBCO) and maintained $37^{o}C$ in a humidified, 5% CO_2 atmosphere. THP1 cells were

differentiated with 20ng/µl PMA for 24 hours, washed with plain RPMI and

maintained in 10% FCS supplemented RPMI for another 24 hours. The cells were

then infected with respective bacteria/beads. RAW264.7 macrophages were seeded in

respective plates and differentiated with 200ng/ml LPS for 12 hours. The cells were

washed once with plain DMEM and infected with respective bacteria/beads.

Animals and isolation of BMDMs

Bone marrow derived macrophages (BMDMs) were isolated from femurs of BALB/C

mice (4-6 weeks old, female) obtained from institutional animal house. BMDMs were

obtained by culturing the marrow cells in the presence of macrophage-colony

stimulating factor (M-CSF, eBioscience, 14-8983-80) for 7 days. Fully differentiated

macrophages were harvested and seeded for infection with H37Rv. The infection

protocol was same as described above for THP-1 macrophages.

Experiments with Latex beads: For microscopy experiments 4μm aldehyde sulfate

latex beads (Life Technologies) were incubated with human/mouse IgG on an agitator

overnight. The beads were washed twice with plain DMEM by centrifugation at 2000

rpm for 10 minutes and added to cells at respective MOI. For flow cytometry

experiments, 1μm FITC labeled aldehyde beads were added at respective MOI to the

differentiated THP1 cells.

Transfection & Nucleofection Assays: Transfection & Transduction was carried out

using JetPrime Reagent (Himedia) as per the manufacturer's protocol. Nucleofection

was carried out with the 4D-Nucleofector™ System Lonza in 20 μl Nucleocuvette™

Strips with program DS-136, as per the manufacturer's protocol. At 24 hours post

Nucleofection these cells expressed the inserted vector as determined by visualization

with a fluorescence microscope.

siRNA Transfection & Inhibitor Assays: Post 24 hours of PMA treatment, the cells

were washed once with plain RPMI. The siRNA's were added as per the

manufacturer's protocol. Post 48 hours of incubation the respective infection/assays

were performed. For the inhibitor assays such as, Dynasore(40μM, 80μM),

Wortmannin (as specified) were added 4 hours before infection. While

Nocodazole(25 μM) was added 2.5 hours before infection and EGTA(3mM),

TMB8(100 μM), Loperamide hydrochloride(100 μM) and Amlodipine besylate(100

μM) were added for 30 minutes before infection.

Creation of stable cell lines: The stable cell lines were created from lentivirus

cytotracer plasmids (B4GALT1-mRuby and NEUM-EGFP, System Biosciences).

$1X10^6$ U937 cells were added to a 24 well plate. The harvested media containing

lentivirus (refer to transfection section) was added to them. At 72 hours post infection

the cells were selected on Puromycin at 350ng/ml for 21 days. The population of

positive cells was routinely checked by microscopy and it was found that post

selection for 21 days the population of cells remained stable (in our case 30-40%).

Staining for Confocal Microscopy: At specific time points cells were fixed with 4%

Paraformaldehyde (PFA) for 15 minutes. This was followed by 2 washes with 1x

PBS. The cells were permeabilized with 0.4% Triton X- 100 for 30 minutes and

washed once with 1X PBS. Blocking was performed using 3 %(w/v) BSA and

0.5%Tween-20 in 1X PBS for one hour, followed by one wash with 1X PBS. The

cells were now stained with respective primary antibody made in the blocking

solution at specific dilution for 90 minutes. The coverslips were then washed once

with 1X PBST and twice with 1X PBS. This was followed by staining with respective

secondary antibody tagged with fluorophore of choice for 90 minutes. The primary

and secondary antibodies were diluted in blocking solutions for use. After three

washes with 1X PBS, the coverslips were then mounted on glass slides with anti-fade

reagent(Life Technologies). Images were acquired with Nikon A1R Laser Scanning

Confocal Microscope with a 100X/1.4NA Plan Apochromat VC, DIC N2 objective

lens. Image processing viz. 3D reconstruction, co-localization and intensity

measurements were done via Imaris 7.2 (Bitplane).

Live Cell Imaging: The live cell imaging dish pre-seeded was RAW264.7 macrophages expressing mCherry-MAN2A. The live cell system was set at 5% CO_2 and 37^0C. The field was set and imaged for 10 minutes. This was followed by an on-stage addition of 0.0001% saponin for 5 minutes. The cells were washed once, supplemented with complete media and imagining continued. The entire procedure was performed on stage to analyze the pre- and post treatment effects on the membrane under our experimental set up.

Flow Cytometry experiments: At respective time points, the infected cells were washed thrice with sterile 1X PBS to remove extracellular bacteria/beads and fixed with 2% PFA. The cells were scraped and run on BD FACS Canto or BD FACS Influx cytometer's.

Western Blot: Post SDS PAGE, the proteins were blotted onto a nitrocellulose membrane using a semidry-transfer system. Following incubation with primary and secondary antibodies, the blots were scanned using Odyssey InfraRed Imaging System (LI-COR BioSciences, Lincoln, NE, USA) at various intensities in order to obtain a blot scan with minimum background. All settings were rigorously maintained for all experiments. The scans were quantitatively analyzed using Odyssey InfraRed Imaging System Application Software Version.3.O (commercially available from (LI-COR BioSciences, Lincoln, NE, USA).

Phagosome Isolation: Latex bead phagosome for 1μm latex beads were isolated as per protocol described earlier (Desjardins et al., 1994). At 1 hour post addition, the plates were washed 4 times with cold 1XPBS, ensuring all the extracellular beads were completely washed off. The cells were scraped off with a rubber policeman and centrifuged twice with cold 1XPBS at 1200 RPM for 5 minutes. The cells were pooled and washed once with homogenization buffer at 1200 RPM for 5 Minutes

(3mM imidazole, 250 mM sucrose, pH 7.4). The cells were then incubated in

appropriate volume of homogenization buffer (with protease inhibitor cocktail –

Amresco) for 30 minutes. The cells were lysed by dounce homogenizer till 90% of the

cells were disrupted as observed under a light microscope. This lysate was

centrifuged at 1200 RPM for 5 minutes at 4^0C to remove the unbroken cells. This was

followed by preparation of sucrose gradient for ultracentrifugation. The homogenate

was made to be consisting of 40% sucrose by mixing with equal volume of 62%

sucrose and 2.25 ml of the same was layered upon 2.0ml of 62% sucrose solution. We

then added 2.25 ml each of 35%, 25% and 10% sucrose solutions. The tubes were

centrifuged at 100,000g for 1 hour at 4^0C in an SW28 rotor (Beckman Coulter). The

LBC band obtained between 10 and 25% sucrose layers was collected and washed

once with cold 1XPBS at 40,000g at 4^0C in an SW28 rotor (Beckman Coulter).

Mass Spectrometry & Data Analysis: For LC-LTQ Orbitrap MS analysis, samples

were re-solubilized in 2% [v/v] acetonitrile, 0.1% [v/v] formic acid in water and

injected onto the trap column at a flow rate of 20 µl/min subsequently peptides were

separated on Zorbax 300SB-C18 (Agilent, Santa Clara, CA, USA) by a gradient

developed from 2% [v/v] acetonitrile, 0.1% [v/v] formic acid to 80% [v/v]

acetonitrile, 0.1% [v/v] formic acid in water over 180 min at a flow rate of 300 nl/min

onto an Agilent 1200 (Agilent, Santa Clara, CA, USA) nano-flow LC-System that

was in-line coupled to the nano-electrospray source of a LTQ-Orbitrap discovery

hybrid mass spectrometer (Thermo Scientific, San Jose, CA, USA). Full MS in a mass

range between m/z 300-2,000 was performed in the Orbitrap mass analyzer with a

resolution of 30,000 at m/z 400 and an AGC target of 2x 105. The strongest five

signals were selected for CID (collision induced dissociation)-MS/MS in the LTQ ion

trap at a normalized collision energy of 35% using an AGC target of 1x 105 and two

microscans. Dynamic exclusion was enabled with one repeat counts during 45 s and

an exclusion period of 180 s. Peptide identification was performed by CID-based

MS/MS of the selected precursors. For protein/peptide identification, MS/MS data

were searched against the *Mus musculus* amino acid sequence database (downloaded

in August 2015) using an in-house Mascot server (version 2.4) through the Proteome

Discoverer 1.4 software. The search was set up for full tryptic peptides with a

maximum of three missed cleavage sites- carbamidomethyl on cysteine, and oxidized

methionine were included as variable modifications. The precursor mass tolerance

threshold was 10 ppm, and the maximum fragment mass error was 0.8 Da. The

significance threshold of the ion score was calculated based on a false discovery rate

of <1%, estimated by the peptide validator node of the Proteome Discoverer software.

Analysis for functional classes:

The non-redundant list of proteins identified was matched against a series of gene

ontology classes from AMIGO2 database. The selection of gene ontology was

empirically made based on known and perceived classes, which together could

represent the set of trafficking proteins identified. The obtained matches were then

represented as a network using Cytoscape 3.2.0.

Statistical Analysis:

Comparative groups were analyzed using paired two-tailed t-test using inbuilt

function in MS Excel.

3-D recreation and analysis: The intensity of a particular fluorophore on the bead

was estimated by first creating a three dimensional bead surface in the captured image

(z-stack) using "spot module" in Imaris Version 7.2 (Bitplane) which automatically

detects spheres in an image depending upon the dimensions fed. The software then

allows one to determine fluorescence intensity on the surface individually for each

fluorophore. For each case maximum fluorescence intensity was determined and

plotted.

PIP3 gradient analysis: The gradient of fluorescence in the AKT-pH mCherry

experiments was determined using "intensity profile line tool" in NIS-Elements

(NIKON).

ACKNOWLEDGEMENTS

The work was supported by a grant from the Department of Biotechnology (DBT),

Govt. of India (BT/PR14730/BRB/10/874/2010, DK) and partly by International

AIDS Society and CFAR funded CNIHR grant from NIH, USA (DK). The work

related to *Mycobacterium tuberculosis* infections were carried out at DBT funded

TACF facility at ICGEB. We thank Prof. Sarman Singh for access to the flow-

cytometer at AIIMS and Purnima for her help in the confocal microscopy. We thank

C-CAMP, NCBS, Bangalore for the mass spectrometry facility. NV is a recipient of a

senior research fellowship from the University Grants Commission, India.

References

786
Achiriloaie, M., B. Barylko, and J.P. Albanesi. 1999. Essential role of the dynamin
pleckstrin homology domain in receptor-mediated endocytosis. *Molecular*
and cellular biology. 19:1410-1415.
Aderem, A., and D.M. Underhill. 1999. Mechanisms of phagocytosis in
macrophages. *Annual review of immunology*. 17:593-623.
Alix, E., S. Mukherjee, and C.R. Roy. 2011. Subversion of membrane transport
pathways by vacuolar pathogens. *The Journal of cell biology*. 195:943-952.
Andrews, N.W. 2002. Lysosomes and the plasma membrane: trypanosomes
reveal a secret relationship. *The Journal of cell biology*. 158:389-394.
Augustine, G.J. 2001. How does calcium trigger neurotransmitter release?
Current opinion in neurobiology. 11:320-326.
Bainton, D.F., R. Takemura, P.E. Stenberg, and Z. Werb. 1989. Rapid
fragmentation and reorganization of Golgi membranes during frustrated
phagocytosis of immobile immune complexes by macrophages. *The*
American journal of pathology. 134:15-26.
Bajno, L., X.R. Peng, A.D. Schreiber, H.P. Moore, W.S. Trimble, and S. Grinstein.
2000. Focal exocytosis of VAMP3-containing vesicles at sites of
phagosome formation. *The Journal of cell biology*. 149:697-706.
Beck, R., Z. Sun, F. Adolf, C. Rutz, J. Bassler, K. Wild, I. Sinning, E. Hurt, B. Brugger,
J. Bethune, and F. Wieland. 2008. Membrane curvature induced by Arf1-

GTP is essential for vesicle formation. *Proceedings of the National*
Academy of Sciences of the United States of America. 105:11731-11736.
Becker, T., A. Volchuk, and J.E. Rothman. 2005. Differential use of endoplasmic
reticulum membrane for phagocytosis in J774 macrophages. *Proceedings*
of the National Academy of Sciences of the United States of America.
102:4022-4026.
Beemiller, P., A.D. Hoppe, and J.A. Swanson. 2006. A phosphatidylinositol-3-
kinase-dependent signal transition regulates ARF1 and ARF6 during
Fcgamma receptor-mediated phagocytosis. *PLoS biology*. 4:e162.
Behar, S.M., M. Divangahi, and H.G. Remold. 2010. Evasion of innate immunity by
Mycobacterium tuberculosis: is death an exit strategy? *Nature reviews.*
Microbiology. 8:668-674.
Bonifacino, J.S., and B.S. Glick. 2004. The mechanisms of vesicle budding and
fusion. *Cell*. 116:153-166.
Braun, V., C. Deschamps, G. Raposo, P. Benaroch, A. Benmerah, P. Chavrier, and F.
Niedergang. 2007. AP-1 and ARF1 control endosomal dynamics at sites of
FcR mediated phagocytosis. *Molecular biology of the cell*. 18:4921-4931.
Braun, V., and F. Niedergang. 2006. Linking exocytosis and endocytosis during
phagocytosis. *Biology of the cell / under the auspices of the European Cell*
Biology Organization. 98:195-201.
Cannon, G.J., and J.A. Swanson. 1992. The macrophage capacity for phagocytosis.
Journal of cell science. 101 (Pt 4):907-913.
Carrithers, M.D., G. Chatterjee, L.M. Carrithers, R. Offoha, U. Iheagwara, C. Rahner,
M. Graham, and S.G. Waxman. 2009. Regulation of podosome formation in
macrophages by a splice variant of the sodium channel SCN8A. *The*
Journal of biological chemistry. 284:8114-8126.
Chen, Y.A., and R.H. Scheller. 2001. SNARE-mediated membrane fusion. *Nature*
reviews. Molecular cell biology. 2:98-106.
Church, J., E.J. Fletcher, K. Abdel-Hamid, and J.F. MacDonald. 1994. Loperamide
blocks high-voltage-activated calcium channels and N-methyl-D-
aspartate-evoked responses in rat and mouse cultured hippocampal
pyramidal neurons. *Molecular pharmacology*. 45:747-757.
Cox, D., C.C. Tseng, G. Bjekic, and S. Greenberg. 1999. A requirement for
phosphatidylinositol 3-kinase in pseudopod extension. *The Journal of*
biological chemistry. 274:1240-1247.
Czibener, C., N.M. Sherer, S.M. Becker, M. Pypaert, E. Hui, E.R. Chapman, W.
Mothes, and N.W. Andrews. 2006. Ca2+ and synaptotagmin VII-dependent
delivery of lysosomal membrane to nascent phagosomes. *The Journal of*
cell biology. 174:997-1007.
Dai, S., Y. Zhang, T. Weimbs, M.B. Yaffe, and D. Zhou. 2007. Bacteria-generated
PtdIns(3)P recruits VAMP8 to facilitate phagocytosis. *Traffic*. 8:1365-
1374.
Desjardins, M., L.A. Huber, R.G. Parton, and G. Griffiths. 1994. Biogenesis of
phagolysosomes proceeds through a sequential series of interactions with
the endocytic apparatus. *The Journal of cell biology*. 124:677-688.
Di, A., D.J. Nelson, V. Bindokas, M.E. Brown, F. Libunao, and H.C. Palfrey. 2003.
Dynamin regulates focal exocytosis in phagocytosing macrophages.
Molecular biology of the cell. 14:2016-2028.

Divangahi, M., M. Chen, H. Gan, D. Desjardins, T.T. Hickman, D.M. Lee, S. Fortune,
S.M. Behar, and H.G. Remold. 2009. Mycobacterium tuberculosis evades
macrophage defenses by inhibiting plasma membrane repair. *Nature*
immunology. 10:899-906.
Drecktrah, D., L.A. Knodler, R. Ireland, and O. Steele-Mortimer. 2006. The
mechanism of Salmonella entry determines the vacuolar environment and
intracellular gene expression. *Traffic*. 7:39-51.
Faundez, V., J.T. Horng, and R.B. Kelly. 1997. ADP ribosylation factor 1 is required
for synaptic vesicle budding in PC12 cells. *The Journal of cell biology*.
138:505-515.
Faurschou, M., and N. Borregaard. 2003. Neutrophil granules and secretory
vesicles in inflammation. *Microbes and infection / Institut Pasteur*. 5:1317-
1327.
Flannagan, R.S., V. Jaumouille, and S. Grinstein. 2012. The cell biology of
phagocytosis. *Annual review of pathology*. 7:61-98.
Funamoto, S., R. Meili, S. Lee, L. Parry, and R.A. Firtel. 2002. Spatial and temporal
regulation of 3-phosphoinositides by PI 3-kinase and PTEN mediates
chemotaxis. *Cell*. 109:611-623.
Gagnon, E., S. Duclos, C. Rondeau, E. Chevet, P.H. Cameron, O. Steele-Mortimer, J.
Paiement, J.J. Bergeron, and M. Desjardins. 2002. Endoplasmic reticulum-
mediated phagocytosis is a mechanism of entry into macrophages. *Cell*.
110:119-131.
Gillingham, A.K., and S. Munro. 2007. The small G proteins of the Arf family and
their regulators. *Annual review of cell and developmental biology*. 23:579-
611.
Gillingham, A.K., and S. Munro. 2016. Finding the Golgi: Golgin Coiled-Coil
Proteins Show the Way. *Trends in cell biology*.
Gold, E.S., D.M. Underhill, N.S. Morrissette, J. Guo, M.A. McNiven, and A. Aderem.
1999. Dynamin 2 is required for phagocytosis in macrophages. *The*
Journal of experimental medicine. 190:1849-1856.
Haynes, L.P., G.M. Thomas, and R.D. Burgoyne. 2005. Interaction of neuronal
calcium sensor-1 and ADP-ribosylation factor 1 allows bidirectional
control of phosphatidylinositol 4-kinase beta and trans-Golgi network-
plasma membrane traffic. *The Journal of biological chemistry*. 280:6047-
6054.
Henley, J.R., and M.A. McNiven. 1996. Association of a dynamin-like protein with
the Golgi apparatus in mammalian cells. *The Journal of cell biology*.
133:761-775.
Holevinsky, K.O., and D.J. Nelson. 1998. Membrane capacitance changes
associated with particle uptake during phagocytosis in macrophages.
Biophysical journal. 75:2577-2586.
Jones, S.M., K.E. Howell, J.R. Henley, H. Cao, and M.A. McNiven. 1998. Role of
dynamin in the formation of transport vesicles from the trans-Golgi
network. *Science*. 279:573-577.
Jutras, I., and M. Desjardins. 2005. Phagocytosis: at the crossroads of innate and
adaptive immunity. *Annual review of cell and developmental biology*.
21:511-527.
Kochegarov, A.A. 2003. Pharmacological modulators of voltage-gated calcium
channels and their therapeutical application. *Cell calcium*. 33:145-162.

Koul, A., T. Herget, B. Klebl, and A. Ullrich. 2004. Interplay between mycobacteria
and host signalling pathways. *Nature reviews. Microbiology*. 2:189-202.
Krauss, M., J.Y. Jia, A. Roux, R. Beck, F.T. Wieland, P. De Camilli, and V. Haucke.
2008. Arf1-GTP-induced tubule formation suggests a function of Arf
family proteins in curvature acquisition at sites of vesicle budding. *The*
Journal of biological chemistry. 283:27717-27723.
Kreitzer, G., A. Marmorstein, P. Okamoto, R. Vallee, and E. Rodriguez-Boulan.
2000. Kinesin and dynamin are required for post-Golgi transport of a
plasma-membrane protein. *Nature cell biology*. 2:125-127.
Lee, W.L., R.E. Harrison, and S. Grinstein. 2003. Phagocytosis by neutrophils.
Microbes and infection / Institut Pasteur. 5:1299-1306.
Link, T.M., U. Park, B.M. Vonakis, D.M. Raben, M.J. Soloski, and M.J. Caterina. 2010.
TRPV2 has a pivotal role in macrophage particle binding and
phagocytosis. *Nature immunology*. 11:232-239.
Lippincott-Schwartz, J., L.C. Yuan, J.S. Bonifacino, and R.D. Klausner. 1989. Rapid
redistribution of Golgi proteins into the ER in cells treated with brefeldin
A: evidence for membrane cycling from Golgi to ER. *Cell*. 56:801-813.
Liu, Y.C., E.R. Chapman, and D.R. Storm. 1991. Targeting of neuromodulin (GAP-
43) fusion proteins to growth cones in cultured rat embryonic neurons.
Neuron. 6:411-420.
Lundqvist-Gustafsson, H., M. Gustafsson, and C. Dahlgren. 2000. Dynamic
ca(2+)changes in neutrophil phagosomes A source for intracellular
ca(2+)during phagolysosome formation? *Cell calcium*. 27:353-362.
Miller, S.G., L. Carnell, and H.H. Moore. 1992. Post-Golgi membrane traffic:
brefeldin A inhibits export from distal Golgi compartments to the cell
surface but not recycling. *The Journal of cell biology*. 118:267-283.
Miyake, K., and P.L. McNeil. 1995. Vesicle accumulation and exocytosis at sites of
plasma membrane disruption. *The Journal of cell biology*. 131:1737-1745.
Niedergang, F., and P. Chavrier. 2004. Signaling and membrane dynamics during
phagocytosis: many roads lead to the phagos(R)ome. *Current opinion in*
cell biology. 16:422-428.
Praefcke, G.J., and H.T. McMahon. 2004. The dynamin superfamily: universal
membrane tubulation and fission molecules? *Nature reviews. Molecular*
cell biology. 5:133-147.
Reddy, A., E.V. Caler, and N.W. Andrews. 2001. Plasma membrane repair is
mediated by Ca(2+)-regulated exocytosis of lysosomes. *Cell*. 106:157-169.
Ridley, A.J., M.A. Schwartz, K. Burridge, R.A. Firtel, M.H. Ginsberg, G. Borisy, J.T.
Parsons, and A.R. Horwitz. 2003. Cell migration: integrating signals from
front to back. *Science*. 302:1704-1709.
Rosa, P., F.A. Barr, J.C. Stinchcombe, C. Binacchi, and W.B. Huttner. 1992.
Brefeldin A inhibits the formation of constitutive secretory vesicles and
immature secretory granules from the trans-Golgi network. *European*
journal of cell biology. 59:265-274.
Samie, M., X. Wang, X. Zhang, A. Goschka, X. Li, X. Cheng, E. Gregg, M. Azar, Y.
Zhuo, A.G. Garrity, Q. Gao, S. Slaugenhaupt, J. Pickel, S.N. Zolov, L.S.
Weisman, G.M. Lenk, S. Titus, M. Bryant-Genevier, N. Southall, M. Juan, M.
Ferrer, and H. Xu. 2013. A TRP channel in the lysosome regulates large
particle phagocytosis via focal exocytosis. *Developmental cell*. 26:511-524.

Scalettar, B.A., P. Rosa, E. Taverna, M. Francolini, T. Tsuboi, S. Terakawa, S.
Koizumi, J. Roder, and A. Jeromin. 2002. Neuronal calcium sensor-1 binds
to regulated secretory organelles and functions in basal and stimulated
exocytosis in PC12 cells. *Journal of cell science*. 115:2399-2412.
Shen, C., S.S. Rathore, H. Yu, D.R. Gulbranson, R. Hua, C. Zhang, N.E. Schoppa, and
J. Shen. 2015. The trans-SNARE-regulating function of Munc18-1 is
essential to synaptic exocytosis. *Nature communications*. 6:8852.
Shui, W., L. Sheu, J. Liu, B. Smart, C.J. Petzold, T.Y. Hsieh, A. Pitcher, J.D. Keasling,
and C.R. Bertozzi. 2008. Membrane proteomics of phagosomes suggests a
connection to autophagy. *Proceedings of the National Academy of Sciences
of the United States of America*. 105:16952-16957.
Singh, D.K., D. Kumar, Z. Siddiqui, S.K. Basu, V. Kumar, and K.V. Rao. 2005. The
strength of receptor signaling is centrally controlled through a
cooperative loop between Ca2+ and an oxidant signal. *Cell*. 121:281-293.
Sollner, T., S.W. Whiteheart, M. Brunner, H. Erdjument-Bromage, S. Geromanos, P.
Tempst, and J.E. Rothman. 1993. SNAP receptors implicated in vesicle
targeting and fusion. *Nature*. 362:318-324.
Steegmaier, M., J. Klumperman, D.L. Foletti, J.S. Yoo, and R.H. Scheller. 1999.
Vesicle-associated membrane protein 4 is implicated in trans-Golgi
network vesicle trafficking. *Molecular biology of the cell*. 10:1957-1972.
Steinhardt, R.A., G. Bi, and J.M. Alderton. 1994. Cell membrane resealing by a
vesicular mechanism similar to neurotransmitter release. *Science*.
263:390-393.
Stendahl, O., K.H. Krause, J. Krischer, P. Jerstrom, J.M. Theler, R.A. Clark, J.L.
Carpentier, and D.P. Lew. 1994. Redistribution of intracellular Ca2+ stores
during phagocytosis in human neutrophils. *Science*. 265:1439-1441.
Stenmark, H. 2009. Rab GTPases as coordinators of vesicle traffic. *Nature reviews.
Molecular cell biology*. 10:513-525.
Sudhof, T.C. 2012. Calcium control of neurotransmitter release. *Cold Spring
Harbor perspectives in biology*. 4:a011353.
Tardieux, I., P. Webster, J. Ravesloot, W. Boron, J.A. Lunn, J.E. Heuser, and N.W.
Andrews. 1992. Lysosome recruitment and fusion are early events
required for trypanosome invasion of mammalian cells. *Cell*. 71:1117-
1130.
Togo, T., J.M. Alderton, G.Q. Bi, and R.A. Steinhardt. 1999. The mechanism of
facilitated cell membrane resealing. *Journal of cell science*. 112 (Pt 5):719-
731.
Togo, T., T.B. Krasieva, and R.A. Steinhardt. 2000. A decrease in membrane
tension precedes successful cell-membrane repair. *Molecular biology of
the cell*. 11:4339-4346.
Underhill, D.M., and A. Ozinsky. 2002. Phagocytosis of microbes: complexity in
action. *Annual review of immunology*. 20:825-852.
Velasco, A., L. Hendricks, K.W. Moremen, D.R. Tulsiani, O. Touster, and M.G.
Farquhar. 1993. Cell type-dependent variations in the subcellular
distribution of alpha-mannosidase I and II. *The Journal of cell biology*.
122:39-51.
Vicker, M.G. 1977. On the origin of the phagocytic membrane. *Experimental cell
research*. 109:127-138.

1000 Vinet, A.F., M. Fukuda, and A. Descoteaux. 2008. The exocytosis regulator
1001   synaptotagmin V controls phagocytosis in macrophages. *Journal of*
1002   *immunology*. 181:5289-5295.
1003 Warnock, D.E., T. Baba, and S.L. Schmid. 1997. Ubiquitously expressed dynamin-
1004   II has a higher intrinsic GTPase activity and a greater propensity for self-
1005   assembly than neuronal dynamin-I. *Molecular biology of the cell*. 8:2553-
1006   2562.
1007 Zhang, Q., D. Cox, C.C. Tseng, J.G. Donaldson, and S. Greenberg. 1998. A
1008   requirement for ARF6 in Fcgamma receptor-mediated phagocytosis in
1009   macrophages. *The Journal of biological chemistry*. 273:19977-19981.
1010 Zuchner, S., M. Noureddine, M. Kennerson, K. Verhoeven, K. Claeys, P. De Jonghe,
1011   J. Merory, S.A. Oliveira, M.C. Speer, J.E. Stenger, G. Walizada, D. Zhu, M.A.
1012   Pericak-Vance, G. Nicholson, V. Timmerman, and J.M. Vance. 2005.
1013   Mutations in the pleckstrin homology domain of dynamin 2 cause
1014   dominant intermediate Charcot-Marie-Tooth disease. *Nature genetics*.
1015   37:289-294.
1016

Figure 1

 FIGURE LEGENDS

 Figure 1: Mannosidase-II is a marker of Golgi apparatus and vesicles derived

 from the Golgi

A) RAW264.7 cells and THP-1 derived macrophages were nucleofected with

mCherry-MAN2A. At 24 hours post nucleofection, cells were visualized under the

confocal microscope (see methods for detail). The 3-D plots at the right were created

using Imaris 7.2 software tool (scale bar: 5μm).

B) THP-1 derived macrophages were permeabilized using 0.5% TritonX-100 and

stained with anti-Mannosidase-II antibody followed by the secondary antibody tagged

with Alexa-560, fixed and visualized under the confocal microscope (scale bar: 5μm).

C) THP-1 derived macrophages were nucleofected with mCherry-MAN2A. At 24

hours post nucleofection, the cells were treated with Brefeldin A (20μM) for 4 hours

and visualized under the microscope (scale bar: 5μm).

D) U937 human macrophages, stably expressing the Golgi marker mRuby-

B4GALT1 (red, β-galactosyl transferase). For the lower panel, B4GALT1 (red)

expressing U937 macrophages were stained with anti-Mannosidase-II antibody

followed by secondary antibody (Alexa-405, blue, scale bar: 5μm).

1034

 Figure 2: Mannosidase-II is recruited at the site of phagocytosis

A)   RAW264.7 macrophages expressing mCherry-MAN2A (red) were incubated

with mouse-IgG coated latex beads for 30 minutes and 1 hour. The images shown are

representative from the 30 minutes samples. At the right, the total intensity of

mCherry-MAN2A puncta on the bead surface was estimated using 3D spot creation

module in Imaris 7.2 and the intensity distribution of the population has been plotted

(see methods for detail). The box plot represents data from more than 200 beads

analyzed from two different experiments (values ± S.D.; scale bar: 4μm).

B)   THP-1 derived macrophages were infected with PKH67 labeled *E. coli* (green,

top panel) or GFP expressing *Salmonella* Typhimurium (green, lower panel) for 15

and 30 minutes. Cells were then stained with anti-Mannosidase-II antibody followed

by Alexa 568 tagged secondary antibody (red). Images shown are representative from

the 15 minutes samples from both the experiments. The bar plots at the right show %

co-localization of *E. coli* or *Salmonella* with Mannosidase-II at the surface at 5 and 10

or 15 and 30 minutes respectively post-infection. Data represents average of more

than 200 bacteria from two different experiments (values ± S.D.; scale bar: 3μm).

C)   mCherry-MAN2A (red) expressing THP-1 derived macrophages were infected

with PKH67 labeled H37Rv (green) for 1 and 2 hours. Samples were fixed and

visualized under the microscope. The bar-plot at the right shows % co-localization of

H37Rv with Mannosidase-II at both these time points. Data represents average of

more than 200 bacteria from two different experiments (values ± S.D.; scale bar:

4μm).

D)   mCherry-MAN2A (red) expressing RAW264.7 macrophages were incubated

with mouse-IgG coated latex beads for 30 minutes or 1 hour. At the respective time

points, cells were immune-stained with anti-TfR antibody followed by a secondary

antibody tagged with Alexa 405 (blue). Presence of TfR or Mannosidase-II at the

bead surface was calculated in terms of fluorescence intensity using the 3D spot

creation module in Imaris 7.2software. The box-plot at the right shows data from

more than 100 beads from two independent experiments (scale bar: 5μm).

E)   mCherry-MAN2A (red) expressing RAW264.7 macrophages were infected

with PKH67 labeled H37Rv (green) for 1 and 2 hours. At the respective time points,

samples were fixed and stained with anti-Transferrin receptor antibody followed by

Alexa-405 tagged secondary antibody (blue). The images are representative from the

1 hour time point. For the plots at the right, % co-localization of H37Rv with

Mannosidase-II, TfR or both Mannosidase-II and TfR was calculated using Imaris

7.2. The data represents average of more than 150 bacteria from three different

experiments (values ± S.D.; scale bar: 5μm).

F)   U937 cells stably expressing EGFP (green) at the plasma-membrane using a

membrane targeting tag from Neuromodulin (NEUM-EGFP) were nucleofected with

mCherry-MAN2A (red) and incubated with human-IgG coated latex beads for 30

minutes. The 3-D constructions at the right hand side were performed using 3D spot

creation module in Imaris 7.2 to visualize the juxtaposition of Mannosidase-II and

latex beads against the plasma membrane (scale bar: 5μm).

1078

Figure 3

Figure 3: Proteomic analysis reveals presence of several VAMPs, ARFs and

RABs in the early phagosomes

A)   Preparation of latex bead phagosomes from THP-1 derived macrophages on a

sucrose density gradient (see methods)

B)   THP-1 derived macrophages were incubated with latex beads (1μm size) for 1

hour. Phagosomes were isolated using differential density ultracentrifugation and

samples were probed for indicated markers using Western blots.

C)   Latex beads phagosomes isolated from RAW264.7 macrophages were lysed

and resolved on a 10% SDS-PAGE (Fig. S2). The lane below 25kDa molecular

weight was analyzed using mass spectrometry to identify enrichment of low

molecular weight proteins (see methods). The list of genes identified was then

searched in the AMIGO2.0 database to establish functional association. Finally the

representative network was constructed using Cytoscape 2.6.1. The pink nodes and

edges in the network denote association with the Golgi apparatus.

1093

Figure 4

Figure 4: Golgi-derived vesicles are recruited through focal exocytosis

A)   mCherry-MAN2A (red) expressing RAW264.7 macrophages were incubated

with mouse-IgG coated latex beads for 30 minutes or 1 hour. At the respective time

points, cells were immune-stained with anti-dynamin2 antibody followed by a

secondary antibody tagged with Alexa 405 (blue). Presence of dynamin2 (Dyn-2) or

Mannosidase-II at the bead surface was calculated using the Imaris 7.2 software. The

box-plot at the right shows data from more than 100 beads from two independent

experiments (values ± S.D.; scale bar: 4μm).

B)   mCherry-MAN2A (red) expressing THP-1 macrophages were infected with

PKH67 labeled H37Rv (green) for 1 and 2 hours. At the respective time points,

samples were fixed and stained with anti-dynamin2 antibody followed by Alexa-405

tagged secondary antibody (blue). The images are representative from the 1hour time

point. For the plots at the right, % co-localization of H37Rv with Mannosidase-II,

dynamin2 or both Mannosidase-II and dynamin2 was calculated using Imaris 7.2. The

data represents average of more than 150 bacteria from three different experiments

(values ± S.D.; scale bar: 4μm).

C) mCherry-MAN2A (red) expressing RAW264.7 macrophages were pretreated with

80μM Dynasore and incubated with mouse-IgG coated latex beads for 30 minutes or

1 hour. At 30 minutes, samples were fixed and stained with anti-dynamin2 antibody

followed by Alexa-405 tagged secondary antibody (blue). Intensity of Mannosidase-II

and dynamin at the bead surface was calculated using the 3D spot creation module in

Imaris 7.2 software. The box-plot shows data from more than 100 beads from two

independent experiments (*p-value<0.05).

D) THP 1 derived macrophages were treated with the respective dynasore

concentrations for 4 hours and incubated with FITC labeled latex beads (1μm). At the

respective time points cells were fixed and analyzed by flow cytometry. To calculate

% uptake, data for each cargo for a given time point, the dynasore treated set was

normalized against the respective untreated control set (values ± S.D.).

E) mCherry-MAN2A (red) expressing RAW264.7 macrophages were incubated with

mouse-IgG coated latex beads for 30 minutes or 1 hour. At the respective time points,

cells were immune-stained with anti-VAMP-3 antibody followed by a secondary

antibody tagged with Alexa 405 (blue). Presence of VAMP-3 or Mannosidase-II at

the bead surface was calculated using the 3D spot creation module in Imaris 7.2

software. The box-plot at the right shows data from more than 100 beads from two

independent experiments (scale bar: 5μm).

F) mCherry-MAN2A (red) expressing THP-1 derived macrophages were infected

with PKH67 labeled H37Rv (green) for 1 and 2 hours. At the respective time points,

samples were fixed and stained with anti-VAMP-3 antibody followed by Alexa-405

tagged secondary antibody (blue). The images are representative from the 1hour time

point. For the plots at the right, % co-localization of H37Rv with Mannosidase-II,

VAMP-3 or both Mannosidase-II and VAMP-3 was calculated using Imaris 7.2. The

data represents average of more than 150 bacteria from three different experiments

(values ± S.D.; scale bar: 4μm).

1137

Figure 5

Figure 5: Role of PIP3 and Calcium from intra and extra-cellular sources in recruitment of Mannosidase II vesicles:

A) THP-1 derived macrophages were pre-treated with 5μM wortmannin for 4 hours. FITC latex beads (1μm) were added to these. The cells were fixed at respective time points and analyzed by flow cytometry. Data shown are average from three different experiments and represented as % uptake in treated cells with respect to the untreated cells (values ± S.D.).

B) THP-1 derived macrophages were pre-treated with 5μM wortmannin for 4 hours. They were infected with PKH labeled H37Rv. At respective time points the cells were fixed and analyzed by flow cytomtery. Data shown are average from three different experiments and represented as % uptake in treated cells with respect to the untreated cells (values ± S.D.).

C) mCherry-MAN2A (red) expressing RAW264.7 macrophages were pre-treated with 5μM Wortmannin or EGTA (3mM)+TMB8 (100 μM) for 30 minutes respectively. The cells were incubated with mouse-IgG coated latex beads for 30 minutes or 1 hour. At the respective time points, cells were fixed and analyzed by confocal microscopy. The images shown are representative from the 30 minutes samples. The 3D construction of latex bead was done using spot creation module of Imaris 7.2.

D) RAW264.7 macrophages expressing mCherry-MAN2A (red) were incubated with mouse-IgG coated latex beads for 30 minutes and 1 hour. The total intensity of mCherry-MAN2A puncta on the bead surface was determined using 3D spot creation module in Imaris 7.2 and the intensity distribution of the population has been plotted. The box plot represents data from more than 200 beads analyzed from two different experiments (*p-value<0.05).

E) THP-1 derived macrophages were pre-treated with EGTA (3mM), TMB8

(100µM) and EGTA+TMB8 for 30 minutes before addition of FITC latex beads

(1µm). At respective time points the cells were fixed and analyzed by flow cytometry.

Data shown are average from three different experiments and represented as % uptake

in treated cells with respect to the untreated cells (values ± S.D.).

F) THP-1 derived macrophages were pre-treated with EGTA (3mM), TMB8

(100µM) and EGTA+TMB8 for 30 minutes before addition of PKH labeled H37Rv.

At respective time points the cells were fixed and analyzed by flow cytometry. Data

shown are average from three different experiments and represented as % uptake in

treated cells with respect to the untreated cells (values ± S.D.).

1173

Figure 6

1174 **Figure 6: Ca^{2+} entry through voltage-gated Ca^{2+} channel helps establish PIP3**

1175 **gradient to aid phagocytosis**

1176 **A)** THP-1 derived macrophages were pre-treated with Amlodipine (100μM) or

1177 Loperamide (100μM) for 30 minutes before addition of GFP expressing *E. coli*. At

1178 respective time points the cells were fixed and analyzed by flow cytometry**.** Data

1179 shown are average from three different experiments and represented as % uptake in

1180 the treated cells with respect to the untreated cells (values ± S.D.).

1181 **B)** RAW264.7 macrophages expressing mCherry-MAN2A (red) were pre-treated

1182 with amlodipine (100μM) or loperamide (100μM) for 30 minutes and subsequently

1183 incubated with mouse-IgG coated latex beads for 30 minutes and 1 hour. The total

1184 intensity of mCherry-MAN2A puncta on the bead surface was determined using 3D

1185 spot creation module in Imaris 7.2 and the intensity distribution of the population has

1186 been plotted. The box-plot represents data from more than 200 beads analyzed from

1187 two different experiments (*p-value<0.05, **p-value<0.01 and ***p-value<0.005).

1188 **C)** RAW264.7 macrophages were transfected with AKT-PH-mCherry. At 24

1189 hours of cells were incubated with GFP expressing E coli for 5 minutes. In parallel we

1190 also had AKT-PH-mCherry expressing cells that were pre-treated with EGTA (3mM),

1191 loperamide (100μM), amlodipine (100μM) or wortmannin (5uM) followed by

1192 incubation with GFP expressing *E. coli* for 5 minutes. Samples were fixed at 5

1193 minutes and analyzed by confocal microscopy. The arrows in the extreme right image

1194 in each of the panel highlight the fluorescence intensity measurements for the analysis

1195 presented in figure 6D (scale bar: 4μm).

1196 **D)** Images in figure 6C (arrows) were analyzed using intensity profile line tool in

1197 the NIS-elements software (see methods). The data represent median from more than

1198 20 fields for each case.

Figure 7

Figure 7: The neuronal calcium sensor (NCS1) in the Golgi apparatus recognizes

Ca2+ signal for focal release of Mannosidase-II vesicles

A)   mCherry-MAN2A (red) expressing U937 derived macrophages were infected

with PKH67 labeled H37Rv (green) for 1 and 2 hours. At the respective time points,

samples were fixed and stained with anti-NCS1 antibody followed by Alexa-405

tagged secondary antibody. The images are representative from the 1hour time point.

Percent co-localization of H37Rv with Mannosidase-II, NCS1 or both Mannosidase-II

and NCS1 was calculated using Imaris 7.2. The data represents average of more than

150 bacteria from three different experiments (values ± S.D.).

B)   In U937 macrophages, incubated with beads for 30 minutes or 1 hour, samples

were stained with anti-NCS1 antibody. Presence of NCS1 at the bead surface was

calculated using the 3D spot creation module in Imaris 7.2 software. The box-plot at

the right shows data from more than 100 beads from two independent experiments.

C)   THP-1 derived macrophages treated with NCS1 siRNA were incubated with

GFP expressing *E. coli* for 5 and 15 minutes. Cells were stained with anti-

Mannosidase-II antibody to assess the recruitment of Mannosidase-II at the

phagosomes in NCS1 depleted cells. The total intensity of mCherry-MAN2A puncta

on the E coli surface was determined using 3D spot creation module in Imaris 7.2. For

lower panel, percent co-localization of *E. coli* with Mannosidase-II was calculated

using Imaris 7.2. The data represents average of more than 150 bacteria from three

different experiments (values ± S.E.M, *p-value<0.05; scale bar: 2μm).

D)   THP-1 macrophages were treated with siRNA against NCS1 or scrambled

control. At 48 hours post siRNA treatment, cells were monitored to uptake latex beads

(1μm), *E coli*, *Salmonella* Typhimurium. or H37Rv for indicated time points. Data are

shown as % uptake in the siRNA treated cells with respect to the scrambled siRNA

control treated cells. Data are representative of three independent experiments (values

± S.D.).

E) A model to explain the Ca2+ dependent activation of NCS1 to trigger the

focal movement of Golgi derived vesicles. i) Initial recognition of an object (bacteria,

beads, cell debris etc) for phagocytosis results in membrane depolarization possibly

due to torsional stress and resulting in the activation of voltage gated Ca^{2+} channels,

leading to the entry of extracellular Ca^{2+} into the cells. The entry of $Ca2^+$ through

voltage-gated Ca^{2+} channels sets the focus for quick and efficient recruitment of PI3K

and results in the accumulation of PIP3 at the site of phagocytosis. These early events

are further aided by signaling through specific pattern recognition receptors and

release of Ca^{2+} from intracellular stores. ii) Increased cytosolic Ca^{2+} is sensed by

Golgi-resident NCS1, which triggers the movement of Mannosidase-II vesicles

towards the site of phagocytosis, guided by a gradient of PIP3, along with dynamin

and microtubule. iii) Mannosidase-II vesicles fuse with the membrane at the site of

phagocytosis and help grow the phagosome around the cargo before final scission and

internalization. iv) The membrane of the nascent phagosome is contributed in part by

the plasma membrane and rest from the Golgi-derived vesicles (in this model for

clarity, we have excluded lysosome and recycling endosomes as additional sources,

see text).