

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22

**A Re-examination of the Role of *AUXIN RESPONSE FACTOR 8* in  
Developmental Abnormalities Caused by the P1/HC-Pro Viral  
Suppressor of RNA Silencing**

Sizolwenkosi Mlotshwa<sup>1,#a</sup>, Gail J. Pruss<sup>1</sup>, John L. MacArthur<sup>1</sup>, Jason W. Reed<sup>2</sup> and  
Vicki Vance<sup>1\*</sup>

<sup>1</sup>Department of Biological Sciences, University of South Carolina, Columbia, South  
Carolina, United States of America

<sup>2</sup>Department of Biology, University of North Carolina, Chapel Hill, North Carolina, United  
States of America

<sup>#a</sup>Current Address: Department of Molecular Genetics, Ohio State University,  
Columbus, Ohio, United States of America

\* Corresponding author

E-mail: [vance@biol.sc.edu](mailto:vance@biol.sc.edu) (VV)

## 1 **Abstract**

2

3 Plant viral suppressors of RNA silencing induce developmental defects similar to those  
4 caused by mutations in genes involved in the microRNA (miRNA) pathway. These  
5 abnormalities were originally thought to reflect a pleiotropic impact of silencing  
6 suppressors on miRNA control of plant development. However, subsequent work with  
7 the P1/HC-Pro potyviral suppressor of silencing showed that global impairment of the  
8 miRNA pathway was not responsible for the phenotypical anomalies. More recently,  
9 developmental defects caused by a *P1/HC-Pro* transgene under control of the 35S  
10 promoter were attributed to moderate upregulation of *AUXIN RESPONSE FACTOR 8*  
11 (*ARF8*), a target of miR167. The key piece of evidence in that work was that the  
12 developmental defects in the 35S-pro:*P1/HC-Pro* transgenic *Arabidopsis* were greatly  
13 alleviated in the F1 progeny of a cross with plants carrying the *arf8-6* mutation. *Arf8-6* is  
14 a SALK line T-DNA insertion mutant, a class of mutations prone to inducing  
15 transcriptional silencing of transgenes expressed from the 35S promoter. Here we  
16 report a re-investigation of the role of *ARF8* in P1/HC-Pro-mediated developmental  
17 defects. We characterized the progeny of a cross between our 35S-pro:*P1/HC-Pro*  
18 transgenic *Arabidopsis* line and the same *arf8-6* T-DNA insertion mutant used in the  
19 earlier study. The T-DNA mutation had little effect in the F1 generation, but almost all  
20 *arf8-6/P1/HC-Pro* progeny had lost the P1/HC-Pro phenotype in the F2 generation.  
21 However, this loss of phenotype was not correlated with the number of functional copies  
22 of the *ARF8* gene. Instead, it reflected transcriptional silencing of the 35S-pro:*P1/HC-*  
23 *Pro* transgene, as evidenced by a pronounced decrease in P1/HC-Pro mRNA

1 accompanied by the appearance of 35S promoter siRNAs. Furthermore, *arf8-8*, an  
2 independent loss-of-function point mutation, had no detectable effects on P1/HC-Pro  
3 phenotype in either the F1 or F2 generations. Together these data argue against the  
4 reported role of increased *ARF8* expression in mediating developmental defects in  
5 *P1/HC-Pro* transgenic plants.

6

## 7 **Author Summary**

8

9 RNA silencing is an important antiviral defense in plants that uses small RNA molecules  
10 to target the invading RNA. Plant viruses, however, have countered with proteins that  
11 suppress RNA silencing, and one of the best-studied plant viral suppressors of silencing  
12 is P1/HC-Pro. When the genetic model plant *Arabidopsis thaliana* is bioengineered to  
13 express P1/HC-Pro, the resulting plants display distinct developmental abnormalities.  
14 These abnormalities are thought to arise because P1/HC-Pro also interferes with the  
15 arm of RNA silencing that uses small RNAs called microRNAs (miRNAs) to regulate  
16 expression of the plant's own genes. Earlier work, however, showed that interference  
17 with all miRNAs in general could not be responsible for these developmental defects.  
18 More recently, it was reported that enhanced expression of a single miRNA-controlled  
19 gene, *AUXIN RESPONSE FACTOR 8 (ARF8)*, underlies the developmental defects  
20 caused by P1/HC-Pro. However, using the same *ARF8* mutation as that report, as well  
21 as a second, independent *ARF8* loss-of function mutation, we now show that mis-  
22 regulation of *ARF8* is not responsible for those defects. One or a few key miRNA-

1 controlled factors might, in fact, underlie the developmental defects caused by P1/HC-  
2 Pro; however, our results show that *ARF8* is not one of the key factors.

3

## 4 **Introduction**

5

6 Eukaryotes have evolved an elaborate network of RNA silencing pathways  
7 mediated by small regulatory RNAs (for reviews, see [1, 2]). These pathways play two  
8 major roles in plants. One is to regulate expression of endogenous genes, and the other  
9 is to defend against invading nucleic acids, such as viruses, transposons, and - more  
10 recently – transgenes. MicroRNAs (miRNAs) are the small RNAs involved in regulation  
11 of endogenous genes and play an important role in development, while small interfering  
12 RNAs (siRNAs) function in defense against invading nucleic acids. Both of these  
13 classes of small RNA are produced by RNase III-like ribonucleases called DICER-LIKE  
14 (DCL) in plants; however, miRNAs and siRNAs differ not only in function, but also in  
15 biogenesis. In *Arabidopsis*, DCL1 processes miRNAs from endogenous, imperfectly  
16 double-stranded transcripts to produce a duplex containing the miRNA and the opposite  
17 strand, the miRNA\*. In contrast, siRNAs are processed by DCL2, DCL3, or DCL4 from  
18 the perfectly double-stranded RNA that triggers the defensive arm of RNA silencing.  
19 MiRNA and siRNA duplexes are methylated by HUA ENHANCER 1 (HEN1), and one  
20 strand incorporates into an RNA-induced silencing complex to guide the Argonaute  
21 (AGO) silencing effector proteins to complementary target sequences. Thus, partially  
22 overlapping biosynthetic pathways lead to two classes of small RNAs with distinct  
23 functions.

1           To counteract antiviral silencing, many plant viruses have evolved proteins that  
2 suppress RNA silencing. It was noted early on that transgenic plants expressing P1/HC-  
3 Pro, the first discovered plant viral suppressor of silencing [3, 4], exhibited pronounced  
4 developmental abnormalities [5], in addition to being suppressed for silencing. These  
5 developmental defects were first observed in tobacco [5, 6], but were subsequently  
6 characterized in greater detail in *Arabidopsis* [7-9]. In *Arabidopsis*, P1/HC-Pro-mediated  
7 developmental defects include reduced size, serrated leaves, late flowering, and  
8 abnormalities in floral morphology resulting in reduced fertility. Some of these  
9 developmental abnormalities resemble ones caused by mutations in genes involved in  
10 plant miRNA biogenesis and function. In addition, *P1/HC-Pro* transgenic lines display  
11 clear defects in both the biogenesis and function of miRNAs. MiRNAs are present at  
12 higher levels in the *P1/HC-Pro* lines, but the usually labile miRNA\* also accumulates [7,  
13 10]. In addition, although miRNA levels are generally increased, so are their mRNA  
14 targets, suggesting a reduction in miRNA function [7, 8]. This correlation of  
15 developmental defects with defects in the miRNA pathway led to the logical idea that  
16 the abnormalities in morphology are caused by interference with miRNA control of  
17 developmental pathways [7, 8].

18           Following up on this idea, a subsequent study investigated the possibility that  
19 ectopic over-expression of *DCL1*, which encodes the Dicer that produces miRNAs in  
20 plants, might rescue the defects in the miRNA pathway and thereby alleviate the  
21 developmental abnormalities in *P1/HC-Pro* transgenic *Arabidopsis*. Surprisingly,  
22 although over-expression of *DCL1* did largely alleviate the developmental defects, it did  
23 not correct the P1/HC-Pro-associated defects in the miRNA pathway: levels of miRNAs

1 and their targets were unchanged from those seen in the parental *P1/HC-Pro* line [9].  
2 These data indicate that P1/HC-Pro-mediated defects in development do not result from  
3 general impairments in miRNA biogenesis or function. However, because only a few  
4 miRNAs and their targets were examined, the study could not rule out the possibility  
5 that the developmental defects were due to interference with one or a small set of  
6 miRNA-controlled genes. Consistent with that idea, it was later reported that the  
7 developmental defects in *P1/HC-Pro* transgenic plants are due to upregulation of  
8 *AUXIN RESPONSE FACTOR 8 (ARF8)*, a target of miR167 that plays a key role in  
9 auxin signaling and developmental patterning [11].

10 We were interested in further investigating the role of *ARF8* in P1/HC-Pro-  
11 mediated developmental defects because the conclusion that upregulation of *ARF8* is  
12 responsible for the defects is in conflict with the earlier work on the effect of over-  
13 expression of *DCL1* on P1/HC-Pro phenotype [9]. Importantly, one of the miRNA and  
14 target pairs looked at in that work was miR167 and its target, *ARF8*. Increased  
15 accumulation of *ARF8* mRNA was unaffected by over-expression of *DCL1* in *P1/HC-Pro*  
16 plants, even though the P1/HC-Pro developmental phenotype was largely eliminated.  
17 Moreover, the developmental phenotypes of *P1/HC-Pro*-expressing plants differ from  
18 those caused by mutation of the *miR167* target site in *ARF8* or the redundantly acting  
19 *ARF6*. *P1/HC-Pro*-expressing plants have narrow serrated leaves, narrow sepals and  
20 petals, and short stamen filaments [7-9]. In contrast, *mARF8* or *mARF6* plants with  
21 mutated *miR167* target sites do not have serrated leaves or narrow flower organs, but  
22 do have elongated stamen filaments and swollen (indehiscent) anthers, and are also  
23 female sterile due to arrested ovule integument growth [12] Based on those results,

1 upregulation of *ARF8* would not appear to be correlated with or responsible for P1/HC-  
2 Pro-mediated developmental defects. To resolve these apparent discrepancies,  
3 therefore, we further characterized the effects of *arf8* mutations on the P1/HC-Pro  
4 phenotype using the same mutation as in the published work [11] as well as a second,  
5 independent mutation.

6

## 7 **Results**

8

### 9 **The P1/HC-Pro developmental phenotype is largely retained in the F1 generation** 10 **of a cross with the *arf8-6* T-DNA insertion mutant**

11       Developmental defects in a *P1/HC-Pro* transgenic *Arabidopsis* line were reported  
12 to be greatly alleviated in the F1 progeny of a cross with plants carrying the *arf8-6*  
13 mutation [11]. To try to reproduce that result, we used the same *arf8-6*  
14 SALK line as in that study and crossed it to our established *P1/HC-Pro* transgenic  
15 *Arabidopsis* line [9, 13, 14]. The *P1/HC-Pro* line is hemizygous for the Turnip Mosaic  
16 Virus (TuMV) coding region, expresses high levels of the *P1/HC-Pro* mRNA, and has a  
17 severe developmental phenotype ([9] and Fig 1). In the F1 generation, about half  
18 (124/238) of the progeny of the cross had the P1/HC-Pro phenotype, while about half  
19 (114/238) did not. Genotyping 69 of the non-phenotypic plants revealed that none had  
20 the *P1/HC-Pro* transgene, and all but three had the *arf8-6* mutation, indicating that  
21 absence of P1/HC-Pro phenotype in the F1 progeny population did not reflect any effect  
22 of the *arf8-6* mutation but simply reflected absence of the *P1/HC-Pro* transgene.  
23 Among 71 F1 plants confirmed to be both hemizygous for *P1/HC-Pro* and heterozygous

1 for *arf8-6*, all had the P1/HC-Pro phenotype. Thus, in our system, hemizygous *arf8-6*  
2 does not cause wide-spread, strong reduction in P1/HC-Pro phenotype in the F1  
3 generation. A minority of our P1/HC-Pro phenotypic F1 progeny, however, showed a  
4 subtle reduction in developmental abnormalities (Fig 1, F1), suggesting that effects of  
5 the *arf8-6* mutation might be better detected in the F2 generation of the cross.

6

7 **Fig 1. The P1/HC-Pro phenotype is largely retained in the F1 but lost in the F2**  
8 **generation of a cross with the *arf8-6* mutant.**

9 Representative parental plants plus F1 and F2 progeny of the cross between *arf8-6* and  
10 P1/HC-Pro lines are shown. **(A)** The *arf8-6/arf8-6* parental line has a wild-type  
11 phenotype, whereas the *P1/HC-Pro/GUS/hpGUS* parent exhibits the characteristic  
12 phenotype of our *P1/HC-Pro* lines. **(B)** P1/HC-Pro phenotypic F1 progeny of the cross,  
13 showing that most have the full P1/HC-Pro phenotype and that only a minority have a  
14 reduced phenotype (arrow). **(C)** Full and reduced P1/HC-Pro phenotype F1 plants from  
15 which RNA was isolated were allowed to self. Seeds from each of the F1 phenotypic  
16 groups were pooled separately before planting. No P1/HC-Pro phenotypic F2 progeny  
17 carrying *arf8-6* were obtained from the reduced phenotype F1 plants (right pot), while a  
18 minority of the F2 progeny from the full phenotype F1 plants exhibited the P1/HC-Pro  
19 phenotype (left pot, arrow).

20

21



1 **The P1/HC-Pro phenotype is largely lost in the F2 generation of the *arf8-6* cross,**  
2 **but the loss is independent of whether the *arf8-6* mutation is hemizygous or**  
3 **homozygous**

4       Because an effect of the *arf8-6* mutation on the phenotype of our *P1/HC-Pro* line  
5 might be more evident when plants are homozygous rather than hemizygous for the  
6 mutation, we examined the phenotypes of the F2 progeny of the cross (after self-  
7 pollination of the F1 plants). Progeny of the slightly reduced P1/HC-Pro phenotypic F1  
8 plants had completely lost the phenotype in the F2 generation (Fig 1, F2) whether plants  
9 were hemizygous or homozygous for *arf8-6*. P1/HC-Pro phenotypic progeny obtained  
10 from the slightly reduced P1/HC-Pro phenotypic F1 plants all lacked the *arf8-6* T-DNA  
11 insertion. Thus, although complete loss of P1/HC-Pro phenotype occurred in this  
12 population of the F2 progeny of the *arf8-6* cross, the loss did not require that *arf8-6* be  
13 homozygous. The P1/HC-Pro phenotype was also lost in the F2 progeny of the fully  
14 P1/HC-Pro phenotypic F1 plants, but was not total in this case (Fig 1, F2). Only about  
15 25% (67/271) of this F2 population had a P1/HC-Pro phenotype, and most (50/67) had  
16 a reduced phenotype. All of these reduced P1/HC-Pro phenotype plants had *arf8-6*, but  
17 only 11 out of the 50 were homozygous for the mutation. Thus, progressive loss of  
18 P1/HC-Pro phenotype occurred in successive generations of the *arf8-6* cross, but was  
19 independent of whether the *arf8-6* mutation was hemizygous or homozygous. Of the  
20 17 fully P1/HC-Pro phenotypic plants obtained in this F2 population, 10 had the *arf8-6*  
21 mutation. Two of the 10 were homozygous for *arf8-6*, showing that the full P1/HC-Pro  
22 phenotype can occur even in the complete absence of *ARF8* expression.

1           Altogether, our results suggest that some mechanism other than reduced *ARF8*  
2 expression is responsible for alleviation and loss of P1/HC-Pro phenotype in plants  
3 carrying the *arf8-6* mutation. Because the *arf8-6* mutation is a SALK line T-DNA  
4 insertion mutation, one likely candidate is transcriptional silencing induced by the T-  
5 DNA. The T-DNAs used to generate SALK lines and several other T-DNA insertion lines  
6 carry an extraneous CaMV 35S promoter that is known to transcriptionally silence 35S  
7 promoter-driven transgenes elsewhere in the plant genome [15].

8           Such unexpected silencing effects can generate misleading results, especially in  
9 studies involving viral suppressors of silencing because such suppressors are not  
10 effective against transcriptional silencing. The *P1/HC-Pro* transgene in our lines and  
11 the one in the work of Jay et al. (2011) are both expressed from the 35S promoter.  
12 Therefore, one likely possibility is that transcriptional silencing is the mechanism  
13 responsible for alleviation and loss of P1/HC-Pro phenotype in plants also carrying the  
14 *arf8-6* mutation.

15  
16 **Indications of 35S promoter-induced transcriptional silencing are evident in the**  
17 **reduced P1/HC-Pro phenotype F1 progeny of the *arf8-6* cross**

18           To see whether transcriptional silencing of 35S promoter-driven transgenes was  
19 occurring in our *arf8-6* mutant plants, we performed RNA gel blot analysis of high and  
20 low molecular weight RNA isolated from F1 and F2 progeny of the *arf8-6* cross. For the  
21 purpose of the molecular analysis, the *P1/HC-Pro* line we used for the cross contained  
22 two additional transgenes that provide an assay for silencing suppression [14]. These  
23 additional loci are a sense *uidA* transgene (*GUS*) encoding  $\beta$ -glucuronidase and a

1 hairpin transgene (*hpGUS*) that post-transcriptionally silences the *GUS* locus [16].  
2 Expression of each of the three transgenes in this *P1/HC-Pro* line is under the control of  
3 the Cauliflower Mosaic Virus (CaMV) 35S-promoter. The *hpGUS* and *GUS* transgenes  
4 function not only as reporters for suppression of post-transcriptional silencing by *P1/HC-*  
5 *Pro*, but also as additional indicators of transcriptional silencing because each is  
6 expressed from a 35S promoter. On their own, however, these two transgenes do not  
7 induce transcriptional silencing [14, 16].

8 . For the RNA gel blot analysis, we examined RNA from plants that had *P1/HC-*  
9 *Pro*, the *arf8-6* mutation, as well as both the *hpGUS* and *GUS* transgenes. In the F1  
10 progeny of the cross, 50 plants were heterozygous for the *arf8-6* mutation and  
11 contained all three of the other transgenes. Of these, 38 plants retained the full *P1/HC-*  
12 *Pro* phenotype, while 12 showed a slight reduction in the phenotype. RNA gel blot  
13 analysis shows that fully *P1/HC-Pro* phenotypic *arf8-6* plants have the same levels as  
14 the *P1/HC-Pro* parent for all high and low molecular weight RNA species examined (Fig  
15 2, lanes 1-6): *P1/HC-Pro* is expressed at about the same level and suppresses post-  
16 transcriptional silencing equally well, as indicated by the highly increased accumulation  
17 of *GUS* and *hpGUS* mRNA compared to the non-*P1/HC-Pro* controls (Fig 2, compare  
18 lanes 1-6 with lanes 15-19). The *GUS* locus gives rise to a transcript that is slightly  
19 longer than the full-length *hpGUS* transcript; however, a prominent high molecular  
20 weight *hpGUS* RNA species that also accumulates corresponds to the loop of the  
21 hairpin [14]. With respect to low molecular weight RNA species, the fully *P1/HC-Pro*  
22 phenotypic *arf8-6* plants and the *P1/HC-Pro* parent show the same increased

1 accumulation of miRNAs and of primary siRNAs, which are the siRNAs from the stem of  
2 the *hpGUS* transcript (Fig 2, compare lanes 1-6 with lanes 15-19).

3

4 **Fig 2. Transcriptional silencing is evident in F1 progeny of the *arf8-6* cross that**  
5 **have a reduced P1/HC-Pro phenotype.**

6 Accumulation of high and low molecular weight RNA from parental and F1 progeny  
7 plants having the indicated genotypes and phenotypes was determined using RNA gel  
8 blot analysis with RNA probes. RNA from the *GUS* line was included as a control to  
9 show the position of the *GUS* transcript. Grouped lanes are all from the same gel. The  
10 high molecular weight blot was hybridized with a probe for *GUS* and *hpGUS*, then  
11 stripped and probed for *P1/HC-Pro*. Two separate low molecular weight (LMW) RNA  
12 gels were run. One LMW blot was probed for *hpGUS* primary siRNAs, while the other  
13 was probed for 35S siRNAs, then successively stripped and probed for the indicated  
14 miRNAs. Ethidium bromide (EtBr) stained rRNA and the major RNA species in LMW  
15 RNA are shown as loading controls.

16

17 Although the difference in phenotype between F1 progeny having the full P1/HC-  
18 Pro phenotype and those having a slightly reduced phenotype is quite subtle, the  
19 difference in the RNA data of these two groups is very striking. The RNA data for the  
20 slightly reduced phenotype plants present a very complicated picture because  
21 transcriptional silencing and P1/HC-Pro suppression of post-transcriptional silencing  
22 both appear to be occurring to some degree. 35S promoter siRNAs are associated with  
23 transcriptional silencing induced by T-DNA insertion mutants [17], providing a good

1 indicator for 35S promoter homology-dependent transcriptional silencing. 35S siRNAs  
2 were not evident in any of the fully P1/HC-Pro phenotypic *arf8-6* F1 plants (Fig 2, lanes  
3 3-6) or in plants that lacked the *arf8-6* mutation (Fig 2, lanes 1-2 and 18-19). In  
4 contrast, we detected 35S promoter siRNAs in RNA from all of the slightly reduced  
5 P1/HC-Pro phenotype *arf8-6* F1 plants assayed (Fig 2, lanes 7-14), indicating that  
6 transcriptional silencing had begun in this population. Consistent with the very subtle  
7 reduction in phenotype, however, accumulation of P1/HC-Pro mRNA appears  
8 unaffected (Fig 2, compare lanes 7-14 with lanes 1-6). In contrast, accumulation of  
9 *GUS* and *hpGUS* mRNA as well as primary siRNAs from *hpGUS* is much lower than in  
10 the fully P1/HC-Pro phenotypic plants (Fig 2, compare lanes 7-14 to lanes 3-6),  
11 although still higher in some cases than in plants lacking *P1/HC-Pro* (Fig 2 compare  
12 lanes 7-14 to 15-19). Thus, despite the onset of transcriptional silencing, P1/HC-Pro  
13 suppression of post-transcriptional silencing appears to be occurring in at least some of  
14 the mildly reduced phenotype F1 progeny. P1/HC-Pro activity in the mildly reduced  
15 phenotype F1 progeny is more clearly demonstrated by an increase in miRNA  
16 accumulation comparable to that in fully P1/HC-Pro phenotypic plants (Fig 2 compare  
17 lanes 7-14 to 1-6 and 15-19).

18 Thus, the *P1/HC-Pro/arf8-6* F1 plants are a mixed population. Most are  
19 completely normal for transgene expression and P1/HC-Pro activity, but some have  
20 started to transcriptionally silence 35S promoter-driven transgenes. It is interesting to  
21 note that 35S promoter-induced transcriptional silencing appears to have started even  
22 in *arf8-6* progeny that do not have the *P1/HC-Pro* transgene: 35S promoter siRNAs are

1 very faintly detectable in RNA from two of the three *arf8-6/GUS/hpGUS* plants we  
2 analyzed (Fig 2, compare lanes 15-16 with lanes 18-19).

3

4 **35S promoter-induced transcriptional silencing is widespread and well developed**  
5 **in the F2 progeny of the *arf8-6* cross**

6 RNA gel blot analysis of the F2 progeny of the *arf8-6* cross confirms that  
7 transcriptional silencing of 35S promoter-driven transgenes is occurring, even in the  
8 absence of the *P1/HC-Pro* transgene. All three of the *arf8-6<sup>-1</sup>/GUS/hpGUS* plants  
9 analyzed are transcriptionally silenced for the *GUS* and *hpGUS* transgenes. These  
10 plants show high levels of 35S siRNAs and have no detectable *GUS* and *hpGUS* mRNA  
11 or primary siRNAs from *hpGUS*, in contrast to *GUS/hpGUS* controls lacking *arf8-6* (Fig  
12 3, compare lanes 15-17 with lanes 18-19). Similarly, *P1/HC-Pro/arf8-6<sup>-1</sup>/GUS/hpGUS*  
13 F2 plants that have completely lost the *P1/HC-Pro* phenotype are clearly  
14 transcriptionally silenced for *P1/HC-Pro* as well as for the *GUS* and *hpGUS* transgenes.  
15 These plants have easily detectable levels of 35S siRNAs and accumulate little or no  
16 *P1/HC-Pro* mRNA and no *GUS* and *hpGUS* mRNA or primary siRNAs from *hpGUS*,  
17 compared to *P1/HC-Pro* controls lacking the *arf8-6* mutation (Fig 3, compare lanes 7-14  
18 with lanes 1-2). Surprisingly, however, miR168 accumulation is slightly elevated  
19 compared to non-*P1/HC-Pro* controls (Fig 3, compare lanes 7-14 with lanes 15-19).

20

21

22

1 **Fig 3. Transcriptional silencing is widespread in the F2 progeny of the *arf8-6***  
2 **cross.**

3 Accumulation of high and low molecular weight RNA from the original parental lines and  
4 from F2 progeny having the indicated genotypes and F1 parental phenotypes was  
5 determined using RNA gel blot analysis with RNA probes. RNA from the *GUS* line was  
6 included as a control to show the position of the *GUS* transcript. Grouped lanes are all  
7 from the same gel. Two high molecular weight (HMW) RNA gels were run. One HMW  
8 blot was hybridized with a probe for *GUS* and *hpGUS*, while the other was probed for  
9 *P1/HC-Pro*. Two low molecular weight (LMW) RNA gels were run. One LMW blot was  
10 probed for *hpGUS* primary siRNAs, while the other was probed for 35S siRNAs, then  
11 successively stripped and probed for the indicated miRNAs. Ethidium bromide (EtBr)  
12 stained rRNA and the major RNA species in LMW RNA are shown as loading controls.

13  
14 As discussed above, some *P1/HC-Pro/arf8-6<sup>-/-</sup>*F2 plants had a *P1/HC-Pro*  
15 phenotype (Fig 1, F2). Most of these had a reduced phenotype, but a few had the full  
16 *P1/HC-Pro* phenotype. RNA gel blot analysis shows that 35S promoter-induced  
17 transcriptional silencing had started in the reduced phenotype plants, as evidenced by  
18 the presence of 35S siRNAs as well as reduced levels of *P1/HC-Pro*, *GUS* and *hpGUS*  
19 mRNA and *hpGUS* primary siRNAs (Fig 3, compare lanes 3-4 with lanes 1-2). In  
20 contrast, 35S promoter-induced transcriptional silencing does not appear to have  
21 started in the few *P1/HC-Pro/arf8-6<sup>-/-</sup>*F2 plants that still have the full *P1/HC-Pro*  
22 phenotype. These plants look identical to the *P1/HC-Pro* parent for all high and low  
23 molecular weight RNA species examined (Fig 3, compare lanes 5-6 with lanes 1-2).

1 Thus, 35S promoter-induced transcriptional silencing appears well established in the  
2 vast majority of F2 progeny of the *arf8-6* cross, supporting our hypothesis that  
3 transcriptional silencing is the mechanism responsible for loss of P1/HC-Pro phenotype  
4 in *P1/HC-Pro/arf8-6* plants. The few *P1/HC-Pro/arf8-6* plants that retain some degree  
5 of P1/HC-Pro phenotype are ones in which transcriptional silencing has not yet been  
6 completely established.

7

### 8 **The *arf8-8* loss of function point mutation does not alleviate P1/HC-Pro-mediated** 9 **developmental defects**

10 To test further whether loss of *ARF8* function can alleviate the P1/HC-Pro  
11 phenotype, we used the *arf8-8* mutant line, which has a point mutation at the junction of  
12 the 5<sup>th</sup> intron and 6<sup>th</sup> exon of *ARF8* [18]. The *arf8-8* mutation causes vegetative and  
13 flower growth phenotypes indistinguishable from those caused by the loss-of-function T-  
14 DNA insertion mutation *arf8-3*, both as a single mutant and in combination with the  
15 *ARF6* loss-of-function mutation, *arf6-2* (Fig 4). Because the *ARF8* and *ARF6* genes act  
16 partially redundantly [19], the loss of function of either gene alone has only a subtle  
17 effect, whereas the double mutant is profoundly affected (Fig 4 and [19]). Thus, *arf8-8*  
18 appears phenotypically as a null allele. We crossed our *P1/HC-Pro/GUS/hpGUS* line to  
19 the *arf8-8* mutant line and examined the phenotypes of the F1 progeny. There were two  
20 classes of phenotype: P1/HC-Pro and wild type. All of the F1 progeny we obtained that  
21 had *P1/HC-Pro* in the heterozygous *arf8-8* background retained the distinctive P1/HC-  
22 Pro phenotype (Fig 5, F1). None of the non-phenotypic plants had the *P1/HC-Pro*  
23 transgene, and none of the *P1/HC-Pro* heterozygous *arf8-8* plants showed any



1 reduction in P1/HC-Pro phenotype. The P1/HC-Pro phenotype was also retained in the  
2 F2 generation of the cross, whether plants were homozygous or heterozygous for the  
3 *arf8-8* mutation (Fig 5, F2).

4

5 **Fig 4. Comparison of *arf8-3* and *arf8-8* developmental phenotypes.**

6 **(A-C)** 5-week-old wild-type, *arf6-2 arf8-3*, and *arf6-2 arf8-8* plants. **(D-H)** Inflorescence  
7 apices of 5-week-old plants of indicated genotypes. Panels D, E, and F were each  
8 assembled from multiple overlapping photographs. **(I)** Measurements of floral organ  
9 lengths and petal widths of mature flowers of each genotype. Data are means  $\pm$  sd  
10 (n=13 to 18 flowers). In I, letters above each bar indicate values that were not  
11 statistically distinguishable by ANOVA at  $P < 0.05$  based on Tukey's HSD statistic,  
12 calculated within each measurement class. As ARF6 and ARF8 act redundantly, the  
13 *arf6-2* background exposes the full phenotypic effects of the *arf8* mutations.

14

15 **Fig 5. The *arf8-8* mutation has no effect on the P1/HC-Pro phenotype in either the**  
16 **F1 or F2 generation of a cross to our *P1/HC-Pro* line.**

17 Representative parental plants plus F1 and F2 progeny of the cross between *arf8-8* and  
18 *P1/HC-Pro* lines are shown. **(A)** The *arf8-8/arf8-8* parental line has a wild-type  
19 phenotype, whereas the *P1/HC-Pro/GUS/hpGUS* parent exhibits the characteristic  
20 phenotype of our *P1/HC-Pro* lines. **(B)** P1/HC-Pro phenotypic F1 progeny of the cross,  
21 showing that all have the full P1/HC-Pro phenotype. **(C)** P1/HC-Pro phenotypic F1  
22 plants from which RNA was isolated were allowed to self, and seeds were pooled

1 before planting. All *P1/HC-Pro* F2 progeny carrying *arf8-8* exhibited the full *P1/HC-Pro*  
2 phenotype.

3

#### 4 **The *arf8-8* loss of function point mutation does not affect *P1/HC-Pro* function**

5 Because *P1/HC-Pro* phenotype was unaffected by the *arf8-8* mutation (Fig 5),  
6 we expected that molecular measures of *P1/HC-Pro* function would also be  
7 unaffected. RNA gel blot analysis of high and low molecular weight RNA isolated from  
8 F1 and F2 progeny of the *arf8-8* cross support that expectation. *P1/HC-Pro* plants  
9 hemizygous (Fig 6) or homozygous (Fig 7) for *arf8-8* look nearly identical to the *P1/HC-Pro*  
10 *Pro* parent for all high and low molecular weight RNA species examined (Fig 6 compare  
11 lanes 3-7 with lanes 1-2; Fig 7 compare lanes 3-6 with lanes 1-2). *P1/HC-Pro*  
12 suppresses the *hpGUS* transgene-induced silencing in these plants, as indicated by the  
13 highly increased accumulation of *GUS* and *hpGUS* mRNA, and mediates increased  
14 accumulation of miRNAs and *hpGUS* primary siRNAs compared to the non-*P1/HC-Pro*  
15 controls (Fig 6, compare lanes 3-7 and 1-2 with lanes 8-14; Fig 7 compare lanes 3-6  
16 and 1-2 with lanes 7-12). Thus, reduced levels of functional *ARF8* in the absence of  
17 35S promoter-induced transcriptional silencing affect neither *P1/HC-Pro* phenotype nor  
18 molecular measures of *P1/HC-Pro* function.

19

20 **Fig 6. The *arf8-8* mutation has no effect on molecular measures of *P1/HC-Pro***  
21 **function in the F1 generation of a cross to our *P1/HC-Pro* line.**

22 Accumulation of high and low molecular weight RNA from parental and F1 progeny  
23 plants having the indicated genotypes was determined using RNA gel blot analysis with

1 RNA probes. RNA from the *GUS* line was included as a control to show the position of  
2 the *GUS* transcript. Grouped lanes are all from the same gel. The high molecular weight  
3 (HMW) blot was hybridized with a probe for *GUS* and *hpGUS*, then stripped and probed  
4 for *P1/HC-Pro*. Two separate low molecular weight (LMW) RNA gels were run. One  
5 LMW blot was probed for *hpGUS* primary siRNAs, while the other was probed for 35S  
6 siRNAs, then successively stripped and probed for the indicated miRNAs. Ethidium  
7 bromide (EtBr) stained rRNA and the major RNA species in LMW RNA are shown as  
8 loading controls.

9

10 **Fig 7. The *arf8-8* mutation has no effect on molecular measures of *P1/HC-Pro***  
11 **function in the F2 generation of a cross to our *P1/HC-Pro* line.**

12 Accumulation of high and low molecular weight RNA from parental and F2 progeny  
13 plants having the indicated genotypes was determined using RNA gel blot analysis with  
14 RNA probes. RNA from the *GUS* line was included as a control to show the position of  
15 the *GUS* transcript. Grouped lanes are all from the same gel. Two high molecular  
16 weight (HMW) RNA gels were run. One HMW blot was hybridized with a probe for *GUS*  
17 and *hpGUS*, while the other was probed for *P1/HC-Pro*. Two low molecular weight  
18 (LMW) RNA gels were run. One LMW blot was probed for *hpGUS* primary siRNAs,  
19 while the other was probed for 35S siRNAs, then successively stripped and probed for  
20 the indicated miRNAs. Ethidium bromide (EtBr) stained rRNA and the major RNA  
21 species in LMW RNA are shown as loading controls.

22

23

24

## 1 Discussion

2

3 Our results clearly show that *ARF8* is not the key factor responsible for  
4 developmental defects in *Arabidopsis* expressing a TuMV *P1/HC-Pro* transgene.  
5 Moderate upregulation of *ARF8* was originally reported to underlie developmental  
6 defects in *Arabidopsis* induced by P1/HC-Pro and two other viral suppressors of  
7 silencing [11]. The identification of *ARF8* as a candidate for this role came from  
8 determining what miRNA targets are upregulated in common in *Arabidopsis* expressing  
9 any one of the viral suppressors, P1/HC-Pro, P19, and P15, as well as in *Arabidopsis*  
10 *hen1-1* and *dcl1-9* mutants. The upregulation in all cases was about 1.5- to 2-fold  
11 suggesting that loss of just one copy of *ARF8* would greatly reduce or eliminate the  
12 developmental defects caused by the viral suppressors. Alleviation of the  
13 developmental defects by one copy of the *arf8-6* SALK insertion mutation was then  
14 taken as evidence that misregulation of *ARF8* was the key factor underlying viral  
15 suppressor-mediated developmental anomalies.

16 We were led to question the conclusions of the Voinnet group [11] for three  
17 reasons. The first was our prior work, which showed no correlation between P1/HC-Pro  
18 phenotype and *ARF8* expression [9]. The second reason was a paradoxical result  
19 presented in the gel blot analysis of RNA isolated from *P1/HC-Pro* plants that carried  
20 the *arf8-6* mutation and a hairpin transgene that targets the gene encoding chalcone  
21 synthase (*CHS*) [[11]: Fig 5E]. The paradoxical result was that the gel blot analysis  
22 showed a total absence of *CHS* siRNAs in those plants, although that same panel of the  
23 figure presented evidence that P1/HC-Pro was suppressing the hairpin transgene-

1 induced post-transcriptional silencing. It has been well established, however, that  
2 primary siRNAs, such as those that would derive from the stem of a hairpin transgene,  
3 are not eliminated in P1/HC-Pro suppression of silencing [13, 14, 20, 21]. Therefore,  
4 *CHS* siRNAs should have been present if P1/HC-Pro suppression of silencing was truly  
5 occurring, and their absence strongly suggested that some confounding factor had  
6 affected the Jay et al. P1/HC-Pro studies [11]. Lastly, the P1/HC-Pro phenotypes differ  
7 in several aspects from those caused by loss of miR167 regulation of *ARF6* or *ARF8*  
8 [12].

9 Our work clearly shows that all the effects of the *arf8-6* T-DNA insertion mutation  
10 on P1/HC-Pro phenotype and on molecular indicators of P1/HC-Pro function are due to  
11 35S promoter homology-dependent transcriptional silencing induced by the *arf8-6* T-  
12 DNA insertion. In addition, the *arf8-8* loss of function point mutation has no effect on  
13 P1/HC-Pro phenotypes or on molecular measures of P1/HC-Pro function, further  
14 arguing against *ARF8* as the key factor responsible for developmental defects caused  
15 by *P1/HC-Pro* expression. Thus, the exact mechanism responsible for P1/HC-Pro-  
16 mediated developmental defects remains an open question. Interference with miRNA-  
17 controlled developmental pathways is a very attractive hypothesis, but ectopic over-  
18 expression of *DCL1* has been shown to separate P1/HC-Pro phenotype from effects on  
19 general miRNA biogenesis and function [9]. Therefore, one or a few key miRNA-  
20 controlled factors might, in fact, underlie the developmental defects caused by P1/HC-  
21 Pro; however, our work shows that *ARF8* is not one of these key factors.

22

23

## 1 **Materials and Methods**

2

### 3 **Arabidopsis lines**

4 The mutant and transgenic lines used are all in the Columbia (Col-0) ecotype and have  
5 been described in previous studies. The *P1/HC-Pro/GUS/hpGUS* line is hemizygous for  
6 the *P1/HC-Pro* transgene and was previously published as *P1/HC-Pro/6b4/306* [14].  
7 The *arf8-6* line is the University of Wisconsin T-DNA line WiscDsLox324F09 [22]. The  
8 *arf8-8* line has an ethyl methane sulfonate (EMS)-generated G to A mutation in a splice  
9 acceptor site in *ARF8* [18].

10

### 11 **Genotyping**

12 Hemizygous and homozygous *arf8-8* plants were genotyped using the PCR primers and  
13 EcoNI digestion of the PCR product as described previously [18]. For hemizygous and  
14 homozygous *arf8-6* genotyping, the T-DNA primer LB (5'-  
15 AACGTCCGCAATGTGTTATTAAGTTGTC-3') was used with primers ARF8-6LP (5'-  
16 CGAGGAAAGGTGAAACCTAC-3') and ARF8-6RP(5'-AGCTGTCAACATCTGGATTGG-  
17 3'). Primers ARF8-6LP and ARF8-6RP amplify a 1014 bp product from the wild-type  
18 locus, and primers ARF8-6RP and LB amplify a 500 bp product from the *arf8-6* locus.  
19 Genotyping for the *hpGUS* (= 306) and *GUS* (= 6b4) transgenes was performed as  
20 described previously [14]. *P1/HC-Pro* primers HC-F (GTGCCCAGAAGTTCAAGAGC)  
21 and HC-R (GTCAACGACTATGCCACTCCAACC) were used to confirm the presence of  
22 the *P1/HC-Pro* transgene in combination with phenotypic selection. *P1/HC-Pro*

1 phenotype was evaluated based on the degree of dwarfing and leaf serration compared  
2 to wild-type *Arabidopsis*.

3

#### 4 **RNA isolation and gel-blot analysis**

5 High and low molecular weight total RNA was isolated from aerial tissues of individual  
6 flowering plants and gel blot analysis performed as described previously [9, 14] [ $\alpha$ -  
7  $^{32}\text{P}$ ]UTP-labeled antisense RNA probes to detect *GUS*, *hpGus*, and P1/HCPPro mRNAs  
8 or the sense RNA probes to detect *hpGUS* primary siRNAs and 35S-promoter siRNAs  
9 were prepared using an Ambion MAXIscript *in vitro* transcription kit (Ambion,  
10 <http://www.ambion.com>) as described previously [14, 17]. The probes were hybridized  
11 to mRNA blots at 68°C in Ambion ULTRAhyb buffer, or to siRNA blots at 42°C in  
12 Ambion ULTRAhyb-oligo buffer. Antisense oligonucleotide probes for detection of  
13 miRNAs were prepared by end-labeling with  $^{32}\text{P}$  using T4 polynucleotide kinase (New  
14 England Biolabs) and hybridized to miRNA blots at 42°C in Ambion ULTRAhyb-oligo  
15 buffer.

16

#### 17 ***arf* mutant phenotypic analyses**

18 Flower organs were dissected from mature open flowers of wild-type and *arf8* single  
19 mutants (2-3 flowers per apex), or from arrested flower buds at the equivalent position  
20 on the inflorescence of *arf6 arf8* double mutants (5-7 flowers per apex). Organs were  
21 placed flat on agar plates, and measured using a camera lucida attached to a dissecting  
22 microscope [18]. Mean lengths or widths measured from at least two sepals, petals,  
23 and long stamens for each flower were used in the summary graph in Figure 4.

## 1 **References**


2


- 3 1. Borges F, Martienssen RA. The expanding world of small RNAs in plants. *Nat*  
4 *Rev Mol Cell Biol.* 2015;16(12):727-41.
- 5 2. Axtell MJ, Westholm JO, Lai EC. Vive la difference: biogenesis and evolution of  
6 microRNAs in plants and animals. *Genome Biol.* 2011;12(4):221.
- 7 3. Anandalakshmi R, Pruss GJ, Ge X, Marathe R, Mallory AC, Smith TH, et al. A  
8 viral suppressor of gene silencing in plants. *Proc Natl Acad Sci U S A.*  
9 1998;95(22):13079-84.
- 10 4. Kasschau KD, Carrington JC. A counterdefensive strategy of plant viruses:  
11 suppression of posttranscriptional gene silencing. *Cell.* 1998;95(4):461-70.
- 12 5. Anandalakshmi R, Marathe R, Ge X, Herr JM, Jr., Mau C, Mallory A, et al. A  
13 calmodulin-related protein that suppresses posttranscriptional gene silencing in  
14 plants. *Science.* 2000;290(5489):142-4.
- 15 6. Mallory AC, Parks G, Endres MW, Baulcombe D, Bowman LH, Pruss GJ, et al.  
16 The amplicon-plus system for high-level expression of transgenes in plants. *Nat*  
17 *Biotechnol.* 2002;20:622-5.
- 18 7. Chapman EJ, Prokhnovsky AI, Gopinath K, Dolja VV, Carrington JC. Viral RNA  
19 silencing suppressors inhibit the microRNA pathway at an intermediate step.  
20 *Genes Dev.* 2004;18(10):1179-86.
- 21 8. Kasschau KD, Xie Z, Allen E, Llave C, Chapman EJ, Krizan KA, et al. P1/HC-  
22 Pro, a viral suppressor of RNA silencing, interferes with Arabidopsis development  
23 and miRNA function. *Dev Cell.* 2003;4(2):205-17.




- 1 9. Mlotshwa S, Schauer SE, Smith TH, Mallory AC, Herr JM, Jr., Roth B, et al.  
2 Ectopic DICER-LIKE1 expression in P1/HC-Pro Arabidopsis rescues phenotypic  
3 anomalies but not defects in microRNA and silencing pathways. *Plant Cell*.  
4 2005;17(11):2873-85.
- 5 10. Mallory AC, Reinhart BJ, Bartel D, Vance VB, Bowman LH. From the Cover: A  
6 viral suppressor of RNA silencing differentially regulates the accumulation of  
7 short interfering RNAs and micro-RNAs in tobacco. *Proc Natl Acad Sci U S A*.  
8 2002;99(23):15228-33.
- 9 11. Jay F, Wang Y, Yu A, Taconnat L, Pelletier S, Colot V, et al. Misregulation of  
10 AUXIN RESPONSE FACTOR 8 underlies the developmental abnormalities  
11 caused by three distinct viral silencing suppressors in Arabidopsis. *PLoS Pathog*.  
12 2011;7(5):e1002035.
- 13 12. Wu MF, Tian Q, Reed JW. Arabidopsis microRNA167 controls patterns of ARF6  
14 and ARF8 expression, and regulates both female and male reproduction.  
15 *Development*. 2006;133(21):4211-8.
- 16 13. Endres MW, Gregory BD, Gao Z, Foreman AW, Mlotshwa S, Ge X, et al. Two  
17 plant viral suppressors of silencing require the ethylene-inducible host  
18 transcription factor RAV2 to block RNA silencing. *PLoS Pathog*.  
19 2010;6(1):e1000729.
- 20 14. Mlotshwa S, Pruss GJ, Peragine A, Endres MW, Li J, Chen X, et al. *DICER-*  
21 *LIKE2* plays a primary role in transitive silencing of transgenes in *Arabidopsis*.  
22 *PLoS ONE*. 2008;3(3):e1755.


- 1 15. Daxinger L, Hunter B, Sheikh M, Jauvion V, Gasciolli V, Vaucheret H, et al.  
2 Unexpected silencing effects from T-DNA tags in Arabidopsis. *Trends Plant Sci.*  
3 2008;13(1):4-6.
- 4 16. Beclin C, Boutet S, Waterhouse P, Vaucheret H. A branched pathway for  
5 transgene-induced RNA silencing in plants. *Curr Biol.* 2002;12(8):684-8.
- 6 17. Mlotshwa S, Pruss GJ, Gao Z, Mgutshini NL, Li J, Chen X, et al. Transcriptional  
7 silencing induced by Arabidopsis T-DNA mutants is associated with 35S  
8 promoter siRNAs and requires genes involved in siRNA-mediated chromatin  
9 silencing. *Plant J.* 2010;64(4):699-704.
- 10 18. Reeves PH, Ellis CM, Ploense SE, Wu MF, Yadav V, Tholl D, et al. A regulatory  
11 network for coordinated flower maturation. *PLoS Genet.* 2012;8(2):e1002506.
- 12 19. Nagpal P, Ellis CM, Weber H, Ploense SE, Barkawi LS, Guilfoyle TJ, et al. Auxin  
13 response factors ARF6 and ARF8 promote jasmonic acid production and flower  
14 maturation. *Development.* 2005;132(18):4107-18.
- 15 20. Johansen LK, Carrington JC. Silencing on the spot. Induction and suppression of  
16 RNA silencing in the Agrobacterium-mediated transient expression system. *Plant*  
17 *Physiol.* 2001;126(3):930-8.
- 18 21. Zhang X, Du P, Lu L, Xiao Q, Wang W, Cao X, et al. Contrasting effects of HC-  
19 Pro and 2b viral suppressors from Sugarcane mosaic virus and Tomato aspermy  
20 cucumovirus on the accumulation of siRNAs. *Virology.* 2008;374(2):351-60.
- 21 22. Goetz M, Vivian-Smith A, Johnson SD, Koltunow AM. AUXIN RESPONSE  
22 FACTOR8 is a negative regulator of fruit initiation in Arabidopsis. *Plant Cell.*  
23 2006;18(8):1873-86.













**A** parents  
*arf8-8/arf8-8*



*HC-Pro/GUS/hpGUS*



**B** F1



**C** F2



