1 **FGIN-1-27, an agonist at translocator protein 18 kDa (TSPO), produces anti-anxiety and anti-**
2 **panic effects in non-mammalian models**
3

4 Jonathan Cueto-Escobedo[1], Juan Francisco Rodríguez Landa[1], Caio Maximino[2]
5
6 [1] Instituto de Neuroetología, Universidad Veracruzana, Xalapa/Mexico
7 [2] Instituto de Estudos em Saúde e Biológicas, Universidade Federal do Sul e Sudeste do Pará,
8 Marabá/Brazil

Corresponding author
Caio Maximino
Instituto de Estudos em Saúde e Biológicas – Universidade Federal do Sul e Sudeste do Pará
Campus III – Av. dos Ipês, s/n
Marabá/PA, Brazil
cmaximino@unifesspa.edu.br

FGIN-1-27 is an agonist at the translocator protein 18 kDa (TSPO), a cholesterol transporter that is associated with neurosteroidogenesis. This protein has been identified as a peripheral binding site for benzodiazepines; in anamniotes, however, a second TSPO isoform that is absent in amniotes has been implicated in erythropoiesis. Functional conservation of the central benzodiazepine binding site located in GABA$_A$ receptors has been demonstrated in anamniotes and amniotes alike; however, the same was not previously demonstrated for TSPO. The present investigation explored the behavioral effects of FGIN-1-27 on an anxiety test in zebrafish and on a mixed anxiety/panic test on wall lizaeds. Results showed that FGIN-1-27 reduced anxiety-like behavior in the zebrafish light/dark preference test which are comparable to diazepam with fewer sedative effects. Similarly, FGIN-1-27 also reduced anxiety- and fear-like behaviors in the defense test battery in wall lizards, again producing fewer sedative-like effects than diazepam; the benzodiazepine was also unable to reduce fear-like behaviors in this species. These results A) underline the functional conservation of TSPO in defensive behavior in anamniotes; B) strengthen the proposal of using anamniote behavior as models in behavioral pharmacology; and C) suggest TSPO/neurosteroidogenesis as a target in treating anxiety disorders.

Keywords: translocator protein 18 kDa; fear; anxiety; zebrafish; wall lizard; benzodiazepines

1. Introduction

The translocator protein 18 kDa (TSPO, mitochondrial benzodiazepine receptor, peripheral benzodiazepine receptor) was first identified as a peripheral binding site for diazepam, but later identified as part of the mitochondrial cholesterol transport pathway that is associated with the regulation of cellular proliferation, immunomodulation, porphyrin transport and heme biosynthesis, anion transport, regulation of steroidogenesis, and apoptosis (Casellas et al., 2002). This transporter is highly expressed in steroidogenic tissues. In the central nervous system, its expression is mainly restricted to ependymal cells and glia, in which it is responsible for the local synthesis of neuroactive steroids such as allopregnanolone (Papadopoulos et al., 2006). This latter neurosteroid, in its turn, positively modulates GABA$_A$ receptors, especially those involved in tonic inhibition (Smith et al., 2009; Maguire et al., 2012). As such, TSPO has been proposed as a pharmacological target for the treatment of neurological and psychiatric disorders associated with decreased

GABAergic tone, such as anxiety disorders (Romeo et al., 1993; de Mateos-Verchere et al., 1998; Kita et al., 2004; Costa et al., 2011; Matsuda et al., 2011; Nin et al., 2011; Pinna and Rasmusson, 2012; Pinna and Rasmusson, 2012; Perna et al., 2014) and epilepsy (Ugale et al., 2004), as well as for the fine control of stress responses (Gunn et al., 2011; Maguire et al., 2012; Maguire, 2014). TSPO agonists have been demonstrated to produce anti-anxiety and anti-conflict effects in rodents with both systemic (Kita et al., 2004; Costa et al., 2011) and intra-hippocampal (Bitran et al. 2000) injections; these effects are blocked by $GABA_A$ receptor antagonists (i.e. picrotoxin) and/or 5α-reductase blockers (i.e. 4-MA) and TSPO antagonists (PK 11195), implicating neurosteroidogenesis and chloride ion channel at the $GABA_A$ receptors in these responses (Bitran et al. 2000). These effects are spared in adrenalectomized and castrated animals, suggesting that they are not mediated by peripheral steroidogenesis, but rather by the production of neurosteroids in the brain (Romeo et al., 1993). Nonetheless, octadecaneuropeptide, a diazepam-binding inhibitor peptide which acts through both the central benzodiazepine receptor (CBR) and TSPO, produces anxiety-like behavior in mice and rats (de Mateos-Verchere et al., 1998), as well as in goldfish (Matsuda et al., 2011). TSPO is highly conserved, being present in Bacteria, Archaea and Eukarya domains (Fan and Papadopoulos, 2013). Anamniotes and invertebrates possess a single isoform, while amniotes possess two TSPO isoforms (Fan et al., 2009). Interestingly, while no functional divergence is predicted to appear between *tspo* (found in invertebrates and basal vertebrates) and *tspo1* (found in amniotes), a functional divergence was detected in TSPO2 (Fan and Papadopoulos, 2013). Some of the neurobehavioral functions of this protein, on the other hand, seen to be conserved. In zebrafish, for example, benzodiazepines have been shown to affect a plethora of anxiety-like behaviors, from bottom-dwelling (Bencan et al., 2009; Egan et al., 2009) and dark preference (Maximino et al., 2010, 2011) to shoal cohesion (Gebauer et al., 2011) and cocaine withdrawal-induced anxiety (López-Patiño et al., 2008). Likewise, benzodiazepines decrease tonic immobility duration and the following freezing and explosive behavior in a defensive behavior battery in the wall lizard

(*Tropidurus oreadicus*), and also increase exploratory behavior in the same test (Maximino et al.,

2014). In the separation stress paradigm, benzodiazepines attenuate separation stress-induced

distress vocalizations in chicks in the anxiety phase, but not in the depression phase (Warnick et al.,

2009). Thus, agonists at the CBR decrease fear- and anxiety-like behavior in both amniotes and

anamniotes. Moreover, some evidence regarding the neurosteroidogenesis pathway in behavioral

control has been suggested by the observation that allopregnanolone has an anticonvulsant effect in

zebrafish (Baxendale et al. 2012), and that chronic fluoxetine treatment upregulates the expression

of genes from the neurosteroidogenesis pathway in this species (Wong et al. 2013). These results

suggest that some downstream effectors of neurosteroidogenesis are conserved, although it is not

known whether the role of TSPO in behavioral control *per se* is conserved. A comparative approach

could untangle this question, especially if species at the base of the amniote and anamniote clades

are used. In this paper, we describe the behavioral effects of FGIN-1-27, a TSPO agonist, in

zebrafish (*Danio rerio*) and wall lizards (*Tropidurus oreadicus*) and compare these responses with

the effects of diazepam, an agonist at the CBR with tested anxiolytic and anticonvulsant actions at

preclinical and clinical research.

2. Experimental procedures

2.1. Experiment 1: Effects of FGIN-1-27 and diazepam on dark preference in zebrafish

2.1.1. Animals and husbandry

120 adult zebrafish from the *longfin* phenotype were acquired in a local aquarium shop and kept in

collective tanks at the laboratory for at least 2 weeks before experiments started. Conditions in the

maintenance tank were kept stable, as per recommendations in Lawrence (2007). Recommendations

in the Brazilian legislation (Conselho Nacional de Controle de Experimentação Animal - CONCEA,

2017) were followed to ensure ethical principles in animal care and throughout experiments. This

manuscript is a complete report of all the studies performed to test the effect of diazepam or FGIN-

1-27 on anxiety-like behavior in zebrafish. We report how we determined our sample size, all data

exclusions (if any), all manipulations, and all measures in the study

2.1.2. Drug administration

Diazepam was dissolved in 40% propylene glycol, 10% ethyl alcohol, 5% sodium benzoate, and

1.5% benzyl alcohol (Maximino et al. 2010). FGIN-1-27 was dissolved in 1% DMSO to which one

or two drops of Tween 80 was added before sonication into a fine suspension (Auta et al. 1993).

Drugs were diluted to their final concentrations and injected i.p. in a volume of 1 µL/0.1 g b.w.

(Kinkel et al. 2010). For both diazepam and FGIN-1-27, the doses used were 0.14, 0.28, 0.57, 1.1,

and 2.3 mg/kg . The three highest doses were chosen based on reported effects of diazepam on

zebrafish (Maximino et al. 2010) or wall lizard (Maximino et al. 2014) defensive behavior, and

were identical for both drugs to facilitate comparison; the lower doses were chosen in order to

define a minimal effect. All drugs were acquired from Tocris Biosciene (UK).

2.1.3. Experimental groups and dosage

120 adult zebrafish were assigned to 12 independent groups (n= 10 fish per group): a vehicle group

(propylene glycol:ethyl alcohol:sodium benzoate:benzyl alcohol for the diazepam controls, DSMO

for the FGIN-1-27 controls) that received 1 µL/0.1 g b.w.; and 10 additional groups that received

the doses described above for diazepam (5 groups) or FGIN-1-27 (5 groups). Sample sizes were

based on Maximino et al. (2010). All drugs were i.p. injected in a volume of 1 µL/0.1 g b.w., 30

min before of the behavioral test. Animals were randomly drawn from the holding tank immediately

before injection, and the order with which doses were tested was randomized *via* generation of ran-

dom numbers using the randomization tool in http://www.randomization.com/. Experimenters were

blinded to treatment by using coded vials for drugs. The data analyst was blinded to phenotype by

using coding to reflect treatments in the resulting datasets; after analysis, data was unblinded.

2.1.4. Scototaxis assay

The light/dark preference (scototaxis) assay was performed as described by Maximino et al., (2013). Briefly, 30 min after injection animals were transferred individually to the central compartment of a black and white tank (15 cm height X 10 cm width X 45 cm length) for a 3-min acclimation period, after which the doors which delimit this compartment were removed and the animal was allowed to freely explore the apparatus for 15 min. While the whole experimental tank was illuminated from above by an homogeneous light source, due to the reflectivity of the apparatus walls and floor average illumination (measured just above the water line) above the black compartment was 225 ± 64.2 (mean $\pm$ S.D.) lux, while in the white compartment it was 307 ± 96.7 lux. The following variables were recorded:

time spent on the white compartment: the time spent in the white half of the tank (percentage of the trial);

squares crossed: the number of 10 cm² squares crossed by the animal in the white compartment;

latency to white compartment: the time in seconds (s) to first entry in the white compartment;

erratic swimming: defined as the number of zig-zag, fast, and unpredictable swimming behavior of short duration;

freezing: the proportional duration of freezing events (in % of time in the white compartment), defined as complete cessation of movements with the exception of eye and operculae movements;

thigmotaxis: the proportional duration of thigmotaxis events (in % of time in the white compartment), defined as swimming in a distance of 2 cm or less from the white compartment's walls;

risk assessment: the number of "risk assessment" events, defined as a fast (<1 s) entry in the white compartment followed by re-entry in the black compartment, or as a partial entry in the

white compartment (i.e., the pectoral fin does not cross the midline);

A digital video camera (Samsung ES68, Carl Zeiss lens) was installed above the apparatus to record

the behavioral activity of the zebrafish. Two independent observers, blinded to treatment, manually

measured the behavioral variables using X-Plo-Rat 2005 (https://github.com/lanec-unifesspa/x-plo-

rat). Inter-observer reliability was at least > 0.85.

2.2. Experiment 2: Effects of FGIN-1-27 and diazepam on the defense test battery in *Tropidu-*

rus oreadicus

2.2.1. Animals and husbandry

120 adult wall lizards (*Tropidurus oreadicus*) of either sex, ranging from 61-96 mm in rostro-cloa-

cal length, were captured in Marabá, PA, Brazil, between February and March 2015. The animals

were inspected for mites, which were removed with forceps before treatment with de-miting solu-

tion as described by the manufacturer (Reptile Relief, Natural Chemistry, Norwalk, USA). All of

the lizards were treated with 50 mg/kg fenbendazole (Vetnil, Brazil), p.o., and then housed accord-

ing to recommendations for anoline lizards (Sanger et al. 2008) for at least 2 weeks before the ex-

periments began. Animals were housed in groups of four in Plexiglas standard laboratory cages (42

cm length x 27.5 cm width x 21 cm height) with mango tree sticks collected from the outdoors to

provide perches. Before using the sticks, they were sterilized for 15 min in an autoclave. To prevent

escape, screen meshes were inserted in the cage tops. The bottoms of the cages were covered with

synthetic cage carpet (Repti Cage Carpet CC-10, Zoo Med, Costa Mesa, USA) placed above a

heater plate (Repti Therm U.T.H. Under Tank RH-6, Zoo Med, Costa Mesa, USA) that kept the

temperature above the carpet at an average of 28°C. The cages were misted with water twice daily,

thus raising the humidity within each cage to approximately 85% (Sanger et al. 2008). The animals

had *ad libitum* access to drinking water. The animals were fed three times weekly with commercial

ration (Shrimp mix, Nutral, Monte Mor, Brazil) and once per week with captured crickets. In the

absence of Brazilian guidelines for lizards, the guidelines for the husbandry of *Anolis* lizards by

Sanger et al. (2008) were used, in an attempt to ensure ethical principles in animal care throughout

experiments.

2.2.2. Drug administration

FGIN-1-27 and diazepam were prepared as in Experiment 1 and injected intraperitoneally in a vol-

ume of 0.5 ml of either vehicle or drug 30 min before behavioral tests.

2.2.3. Experimental groups and dosage

120 adult lizards were asigned to 12 independent groups (n= 10 lizards per group): a vehicle group

(propylene glycol:ethyl alcohol:sodium benzoate:benzyl alcohol for the diazepam controls, DSMO

for the FGIN-1-27 controls) that received 1 µL/0.1 g b.w.; and 10 additional groups that received

the doses described above for diazepam (5 groups) or FGIN-1-27 (5 groups). Sample sizes were

based on Maximino et al. (2014). All drugs were i.p. injected in a volume of 0.5 ml per animal, 30

min before of the behavioral test. Animals were randomly drawn from the cage immediately before

injection, and the order with which doses were tested was randomized *via* generation of random

numbers using the randomization tool in http://www.randomization.com/. Experimenters were

blinded to treatment by using coded vials for drugs. The data analyst was blinded to phenotype by

using coding to reflect treatments in the resulting datasets; after analysis, data was unblinded.

2.2.4. Defense test battery

The defense test battery was applied as described by Maximino et al. (2014). Briefly, tonic immo-

bility was induced by carefully placing the animal on its back in the center of a 10 cm diameter cir-

cular open field and applying mild pressure to the thorax and pelvis while restraining the limbs.

When the lizard ceased struggling, it was slowly released, and the time taken for it to resume an up-

right posture was recorded (Hennig 1979). After the animal spontaneously ceased tonic immobility (TI), the following behavioral endpoints were recorded after each of these manipulations:

> *tonic immobility (TI):* the total duration, in minutes (min), in a rigid supine posture after release;
>
> *freezing:* the lack of limb, neck, or tongue movements for more than 5 s in an upright position;
>
> *circling:* a high-velocity escape attempt with a latency of less than 10 s after release, usually leading to circling around the edges of the apparatus, and quantified as the number of complete circles made near the walls;
>
> *tongue-flicking:* repeatedly licking the air with the tongue;
>
> *ventilatory frequency:* the average number of inspiratory responses per minute;
>
> *total locomotion:* the number of 2 cm² squares crossed by normal locomotor responses (i.e., not concomitant to circling); can be superimposed to tongue-flicking.

A digital video camera (Samsung ES68, Carl Zeiss lens) was installed above the apparatus to record the behavioral activity of the lizards. Two independent observers, blinded to treatment, manually measured the behavioral variables using EthoLog 2.2 (Ottoni 2000). Inter-observer reliability was at least > 0.85.

2.3. Statistical analysis

The data were analyzed using two-way ANOVAs, with drug and dose as between-subjects factors. Values of $p \leq 0.05$ in the ANOVA were followed by Tukey's HSD whenever appropriate; planned comparisons were between different doses of a given drug and its vehicle and between same doses of both drugs. All statistical analyses were made using R 3.1.3. Data are presented graphically as means ± S.E.M.

3. Results

3.1. Experiment 1

Main effects of drug ($F_{1, 107} = 91.877$, $p < 0.0001$) and dose ($F_{5, 107} = 36.464$, $p < 0.0001$), as well as an interaction effect ($F_{5,107} = 4.924$, $p = 0.000428$) were observed in time spent into the white compartment (Figure 1A). Post-hoc tests uncovered differences between all doses except the highest in relation to controls in diazepam-treated animals ($p < 0.001$) and between all doses in relation to controls in FGIN-1-27-treated animals ($p < 0.001$). Finally, differences were also observed between FGIN-1-27- and diazepam-treated animals at doses of 0.56 mg/kg ($p = 0.0012$), 1.2 mg/kg ($p < 0.0001$), and 2.4 mg/kg ($p = 0.0002$).

Main effects of drug ($F_{1, 107} = 17.9$, $p < 0.0001$) and dose ($F_{5, 107} = 211.6$, $p < 0.0001$), as well as an interaction effect ($F_{5, 107} = 20.0$, $p < 0.0001$), were found for risk assessment (Figure 1B). Post-hoc tests uncovered differences between vehicle-treated and diazepam-treated animals at all doses ($p < 0.0001$), and between vehicle-treated and FGIN-1-27-treated animals at all doses ($p < 0.0001$). Moreover, differences between diazepam- and FGIN-1-27-treated animals were observed at 1.1 mg/kg ($p = 0.012$) and 2.3 mg/kg ($p < 0.0001$).

A main effect of dose ($F_{5, 107} = 111.547$, $p < 0.0001$), but not drug ($F_{1, 107} = 0.643$), was found for thigmotaxis (Figure 1C), and an interaction effect was also found ($F_{5, 107} = 10.536$, $p < 0.0001$). Post-hoc tests unveiled differences between vehicle-treated and diazepam-treated animals at doses above 0.28 mg/kg ($p < 0.01$), as well as between all FGIN-1-27 doses and vehicle-treated animals ($p < 0.0001$). Differences were also observed between diazepam- and FGIN-1-27-treated zebrafish at doses of 0.14 mg/kg ($p = 0.0065$) and 1.1 mg/kg ($p = 0.044$).

Main effects of drug ($F_{1, 107} = 89.134$, $p < 0.0001$) and dose ($F_{5, 107} = 47.418$, $p < 0.0001$), as well as an interaction effect ($F_{5. 107} = 6.921$, $p < 0.0001$), were found for freezing (Figure 1D). Post-hoc tests uncovered differences between vehicle-treated and diazepam-treated animals at all doses ($p < 0.001$), and between FGIN-1-27-treated and vehicle-treated animals at 0.28 mg/kg and higher doses

(p < 0.001). Differences were also found between diazepam- and FGIN-1-27-treated zebrafish at

doses of 0.14 mg/kg (p = 0.03), 1.1 mg/kg (p < 0.0001), and 2.3 mg/kg (p < 0.0001).

Main effects of drug ($F_{1, 107}$ = 14.46, p < 0.0001) and dose ($F_{5, 107}$ = 42.67, p < 0.0001), as well as an

interaction effect ($F_{5, 107}$ = 26.31, p < 0.0001), were found for erratic swimming (Figure 1E). Post-

tests uncovered no statistically significant differences between vehicle-treated and diazepam-treated

animals at any dose (p > 0.05), while FGIN-1-27-treated animals were significantly different be-

tween vehicle-treated animals at doses of 0.28 mg/kg (p = 0.0001), 1.1 mg/kg (p < 0.001) and 2.3

mg/kg (p < 0.0001). Also, differences were observed between FGIN-1-27- and diazepam-treated

animals at 0.28 mg/kg (p = 0.021), 1.1 mg/kg (p = 0.05), and 2.3 mg/kg (p < 0.0001).

Finally, main effects of drug ($F_{1, 107}$ = 9.297, p = 0.00289) and dose ($F_{5, 107}$ = 12.298, p < 0.0001), as

well as an interaction effect ($F_{5, 107}$ = 4.054, p = 0.00208) were found for number of entries into the

white compartment (Figure 1F). Post-hoc tests uncovered differences between vehicle-treated and

diazepam-treated zebrafish at doses of 0.28 mg/kg and 0.57 mg/kg (p < 0.01) and vehicle-treated

and FGIN-1-27-treated zebrafish at the highest dose (p = 0.003). No differences were found be-

tween diazepam- and FGIN-1-27-treated animals.

3.2. Experiment 2

Main effects of drug ($F_{1, 108}$ = 248.5, p < 0.0001) and dose ($F_{1, 108}$ = 93.39, p < 0.0001), as well as an

interaction effect ($F_{1, 108}$ = 60.09, p < 0.0001) were found for TI duration (Figure 2A). Post-hoc tests

found statistically significant differences between vehicle-treated and diazepam-treated lizards at

doses of 0.28-1.1 mg/kg (p < 0.05) and vehicle-treated and FGIN-1-27-treated lizards at all doses

except the higher (p < 0.0001). Differences between FGIN-1-27- and diazepam-treated lizards were

observed at all doses except the higher (p < 0.0001).

Main effects of drug ($F_{1, 108}$ = 771.29, p < 0.0001) and dose ($F_{5, 108}$ = 59.58, p < 0.0001), as well as an

interaction effect ($F_{5, 108}$ = 66.98, p < 0.0001) were found for circling behavior (Figure 2B). Vehicle-

treated animals differed from diazepam-treated animals at 0.14 mg/kg (p = 0.019), 0.57 mg/kg (p <

0.0001), and 1.1 mg/kg (p = 0.003); vehicle-treated lizards differed from FGIN-1-27-treated ani-

mals at all doses (p < 0.0001). Diazepam- and FGIN-1-27-treated animals differed at all doses (p <

0.0001).

Main effects of drug ($F_{1, 108}$ = 291.58, p < 0.0001) and dose ($F_{5, 108}$ = 45.09, p < 0.0001), as well as an

interaction effect ($F_{5, 108}$ = 14.63, p < 0.0001) were found for freezing (Figure 2C). Post-hoc tests

identified significant differences between vehicle- and diazepam-treated animals at 1.1 mg/kg (p <

0.0001) and between vehicle- and FGIN-1-27-treated animals at all doses (p < 0.0001). Differences

between diazepam- and FGIN-1-27-treated lizards were also found for all doses (p < 0.0001).

A drug main effect ($F_{1, 108}$ = 1482.4, p < 0.0001) and a dose main effect ($F_{5, 108}$ = 110.18, p <

0.0001), as well as an interaction effect ($F_{5, 108}$ = 86.33, p < 0.0001) were observed for ventilatory

frequency (Figure 2D). Post-hoc tests uncovered differences between vehicle-treated and diazepam-

treated animals at 1.1 mg/kg (p < 0.0001), and between vehicle-treated and FGIN-1-27-treated ani-

mals at all doses (p < 0.0001). Finally, statistically significant differences were found between di-

azepam- and FGIN-1-27-treated animals at all doses (p < 0.0001).

Main effects of drug ($F_{1, 108}$ = 38.4, p < 0.0001) and dose ($F_{5, 108}$ = 437.2, p < 0.0001), as well as in-

teraction effect ($F_{5, 108}$ = 3735, p < 0.0001) were found for tongue-flicking (Figure 2E). Post-hoc

tests revealed significant differences between vehicle- and diazepam-treated lizards at doses of 1.1

and 2.3 mg/kg (p < 0.0001), and between vehicle- and FGIN-1-27-treated lizards at all doses except

the highest (p < 0.0001). Moreover, statistically significant differences between diazepam- and

FGIN-1-27-treated lizards at all evaluated doses were found (p < 0.0001).

Main effects of both drug ($F_{1, 108}$ = 87.41, p < 0.0001) and dose ($F_{5, 108}$ = 36.46, p < 0.0001), as well

as an interaction effect ($F_{5, 108}$ = 56.92, p < 0.0001) were found for total locomotion (Figure 2F). Sig-

nificant differences were found between vehicle- and diazepam-treated lizards at 0.57 and 2.3

mg/kg (p < 0.0003) and vehicle- and FGIN-1-27-treated lizards at 1.1 and 2.3 mg/kg (p < 0.002).

Finally, statistically significant differences were also found between diazepam- and FGIN-1-27-

treated animals at 2.3 mg/kg (p < 0.0001).

4. Discussion

In the present work we demonstrated that the TSPO agonist FGIN-1-27 produced a dose-dependent

decrease in defensive behavior in both wall lizards and zebrafish. Specifically, in wall lizards

FGIN-1-27 decreased tonic immobility in the intermediate dose range (0.14-1.1 mg/kg); decreased

post-TI ventilatory frequency, freezing and circling at all doses; and increased exploratory behavior

(tongue-flicking) and decreased thigmotaxis at 0.14-1.1 mg/kg. Similarly, in zebrafish FGIN-1-27

decreased scototaxis, thigmotaxis, freezing and risk assessment at all doses, increasing erratic

swimming at 0.14 and 0.28 mg/kg. The highest dose (2.3 mg/kg) decreased locomotion in both

species, suggesting a sedative effect. Moreover, diazepam, a CBR and TSPO agonist, decreased TI

duration at the smaller doses and increased it at high doses, increased tongue-flicking and decreased

thigmotaxis at doses above 0.28 mg/kg; no effects were observed in post-TI behavior. In zebrafish,

diazepam produced an inverted-U effect on scototaxis and entries into the white compartment,

monotonically decreasing freezing, thigmotaxis and risk assessment.

In the lizard defense test battery, induction of TI produces a stereotypical behavioral pattern in

which is followed by either freezing or "explosive" circling behavior; after that, the animal emits

exploratory behavior, marked by thigmotaxis ("wall-hugging") and tongue-flicking (Maximino et

al., 2014). This sequence resembles defensive behavior at increasing predatory imminence continua

(Fanselow and Lester, 1988) – that is, defensive behavior in this test shifts from panic-like, circa-

strike behavior towards sustained risk assessment (i.e., tongue-flicking, thigmotaxis). Moreover, in

a previous experiment panicolytic drugs (i.e., alprazolam, imipramine) decreased TI duration, freez-

ing and circling behavior, while diazepam (0.5 mg/kg) increased tongue-flicking and decreased

thigmotaxis (Maximino et al., 2014). In the present experiment, FGIN-1-27 produced a more wide

range of effects than diazepam: the TSPO agonist affected both exploratory behavior (consistent

with an anxiolytic-like effect), post-TI behavior (consistent with a panicolytic effect) and TI dura-

tion, which could also represent a panicolytic-like effect. Diazepam, on the other hand, only af-

fected exploratory behavior and had a hormetic (inverted-U) profile on TI duration.

A complementary profile was observed in the zebrafish scototaxis assay. Benzodiazepines have

been shown to produce a hormetic profile in several behavioral tests in zebrafish (Bencan et al.,

2009; Cachat et al., 2010; Sackerman et al., 2010; Vada et al., 2015). Previous experiments also

demonstrated that diazepam was anxiolytic at 1.25 mg/kg, but not 2.5 mg/kg, in the scototaxis assay

(Maximino et al., 2011); in addition to this observation, the present work also demonstrated effects

of diazepam on freezing, thigmotaxis and risk assessment. Importantly, FGIN-1-27 treatment also

produced anxiolytic-like effects, with locomotor-impairing effects at the highest dose.

Little is known about the role of TSPO in behavioral control in anamniotes. In goldfish (*Carassius*

auratus), octadecaneuropeptide increased the latency to enter a white compartment in a light/dark

box, suggesting an anxiogenic-like effect (Matsuda et al., 2011). While octadecaneuropeptide is an

endozepine which acts as an agonist at TSPO (Papadopoulos et al., 1991), it also acts as an antago-

nist at the CBR (Ferrero et al., 1986); consistently with the hypothesis that the anxiogenic-like ef-

fect of octadecaneuropeptide is mediated by the CBR, flumazenil, but not a metabotropic en-

dozepine receptor antagonist, blocked the scototaxis effects in the goldfish (Matsuda et al. 2011).

Overall, the present results suggest that drugs targeting the CBR and MBR exert anti-anxiety effects

in anamniotes, while drugs acting at the MBR also exert anti-panic effects. It is plausible that some

of the effects of diazepam were mediated by the MBR; further experiments are needed to untangle

the precise mechanisms. Thus, while the present results suggest a functional conservation of TSPO

that is concomitant to gene duplication, it is not known whether FGIN-1-27 (or even diazepam) pro-

duced its behavioral effects in lizards and zebrafish by acting on the (conserved) *tspo1* or whether

some effects are also mediated by *tspo*. Further experiments will clarify the issue. The present re-

sults also add to the mounting evidence that TSPO/MBR ligands could be used to treat fear disorders, including panic disorder, in human populations. The degree of conservation of *tspo1* suggest that anamniotes could be used as experimental models to study potential anti-panic drugs to contribute to develop of pharmacological therapeutic strategies to ameliorate panic disorders symptoms in the human.

References

Auta J, Romeo E, Kozikowski AP, et al (1993) Participation of mitochondrial diazepam binding inhibitor receptors in the anticonflict, antineophobic and anticonvulsant action of 2-aryl-3-indoleacetamide and imidazopyridine derivatives. J Pharmacol Exp Ther 265:649–656.

Baxendale S, Holdsworth CJ, Santoscoy PLM, et al (2012) Identification of compounds with anticonvulsant properties in a zebrafish model of epileptic seizures. Dis Model Mech 5:773–784. doi: 10.1242/dmm.010090

Bencan Z, Sledge D, Levin ED (2009) Buspirone, chlordiazepoxide and diazepam effects in a zebrafish model of anxiety. Pharmacol Biochem Behav 94:75–80. doi: 10.1016/j.pbb.2009.07.009

Bitran D, Foley M, Audette D, et al (2000) Activation of peripheral mitochondrial benzodiazepine receptors in the hippocampus stimulates allopregnanolone synthesis and produces anxiolytic-like effects in the rat. Psychopharmacology (Berl) 151:64–71.

Cachat J, Canavello P, Elegante M, et al (2010) Modeling withdrawal syndrome in zebrafish. Behav Brain Res 208:371–376.

Casellas P, Galiegue S, Basile AS (2002) Peripheral benzodiazepine receptors and mitochondrial function. Neurochem Int 40:475–486.

Conselho Nacional de Controle de Experimentação Animal - CONCEA. Diretriz brasileira para o cuidado e a utilização de animais para fins científicos e didáticos - DBCA. Anexo I. Peixes mantidos em instalações de instituições de ensino ou pesquisa científica, Pub. L. No. Resolução Normativa 30/2016 (2017). Brasil.

Costa B, Da Pozzo E, Chelli B, et al (2011) Anxiolytic properties of a 2-phenylindolglyoxylamide TSPO ligand: Stimulation of in vitro neurosteroid production affecting GABAA receptor activity. Psychoneuroendocrinology 36:463–472. doi: 10.1016/j.psyneuen.2010.07.021

De Mateos-Verchere JG, Leprince JJ, Tonon M-C, et al (1998) The octadecaneuropeptide ODN induces anxiety in rodents: Possible involvement of a shorter biologically active fragment. Peptides 19:841–848.

Egan RJ, Bergner CL, Hart PC, et al (2009) Understanding behavioral and physiological phenotypes of stress and anxiety in zebrafish. Behav Brain Res 205:38–44. doi: 10.1016/j.bbr.2009.06.022

Fan J, Papadopoulos V (2013) Evolutionary origin of the mitochondrial cholesterol transport machinery reveals a universal mechanism of steroid hormone biosynthesis in animals. PLoS One 8:e76701. doi: 10.1371/journal.pone.0076701

Fan J, Rone MB, Papadopoulos V (2009) Translocator protein 2 is involved in cholesterol redistribution during erythropoiesis. J Biol Chem 284:30484–30497. doi: 10.1074/jbc.M109.029876

Fanselow MS, Lester L (1988) A functional behavioristic approach to aversively motivated behavior: Predatory imminence as a determinant of the topography of defensive behavior. In: Bolles RC, Beecher MD (eds) Evol. Learn. Erlbaum, Hillsdale, pp 185–211

Ferrero P, Santi MR, Conti-Tronconi B, et al (1986) Study of an octadecaneuropeptide derived from diazepam binding inhibitor (DBI): Biological activity and presence in rat brain. Proc Natl Acad Sci U S A 83:827–831. doi: 10.1073/pnas.83.3.827

Gebauer DL, Pagnussat N, Piato ÂL, et al (2011) Effects of anxiolytics in zebrafish: Similarities and differences between benzodiazepines, buspirone and ethanol. Pharmacol Biochem Behav 99:480–486. doi: 10.1016/j.pbb.2011.04.021

Gunn BG, Brown AR, Lambert JJ, Belelli D (2011) Neurosteroids and GABAA receptor interactions: A focus on stress. Front Neurosci 5:Article 131. doi: 10.3389/fnins.2011.00131

Hennig CW (1979) The effects of physical environment, time in captivity, and defensive distance on tonic immobility, freezing, and flight behaviors in Anolis carolinensis. Anim Learn Behav 7:106–110.

Kinkel MD, Eames SC, Philipson LH, Prince VE (2010) Intraperitoneal injection into adult zebrafish. J Vis Exp 42:2126. doi: 10.3791/2126

Kita A, Kohayakawa H, Kinoshita T, et al (2004) Antianxiety and antidepressant-like effects of AC-5216 , a novel mitochondrial benzodiazepine receptor ligand. Br J Pharmacol 142:1059–1072. doi: 10.1038/sj.bjp.0705681

Lawrence C (2007) The husbandry of zebrafish (Danio rerio): A review. Aquaculture 269:1–20. doi: 10.1016/j.aquaculture.2007.04.077

López-Patiño MA, Yu L, Cabral H, Zhdanova I V (2008) Anxiogenic effects of cocaine withdrawal in zebrafish. Physiol Behav 93:160–171. doi: 10.1016/j.physbeh.2007.08.013

Maguire J (2014) Stress-induced plasticity of GABAergic inhibition. Front Cell Neurosci 8:Article 157. doi: 10.3389/fncel.2014.00157

Maguire J, Mody I, Maguire J (2012) The reciprocal regulation of stress hormones and GABA(A) receptors. Front Cell Neurosci 6:Article 4. doi: 10.3389/fncel.2012.00004

Matsuda K, Wada K, Azuma M, et al (2011) The octadecaneuropeptide exerts an anxiogenic-like action in goldfish. Neuroscience 181:100–108. doi: 10.1016/j.neuroscience.2011.02.058

Maximino C, Carvalho CM, Morato S (2014) Discrimination of anxiety- versus panic-like behavior in the wall lizard Tropidurus oreadicus. Psychol Neurosci 7:227–231. doi: 10.3922/j.psns.2014.020

Maximino C, de Brito TM, Dias CA de M, et al (2010) Scototaxis as anxiety-like behavior in fish. Nat Protoc 5:209–216. doi: 10.1038/nprot.2009.225

Maximino C, Puty B, Benzecry R, et al (2013) Role of serotonin in zebrafish (Danio rerio) anxiety: Relationship with serotonin levels and effect of buspirone, WAY 100635, SB 224289, fluoxetine and para-chlorophenylalanine (pCPA) in two behavioral models. Neuropharmacology 71:83–97. doi: 10.1016/j.neuropharm.2013.03.006

Maximino C, Silva AWB da, Gouveia Jr. A, Herculano AM (2011) Pharmacological analysis of zebrafish (Danio rerio) scototaxis. Prog Neuropsychopharmacol Biol Psychiatry 35:624–631. doi: 10.1016/j.pnpbp.2011.01.006

Nin MS, Martinez LA, Pibiri F, et al (2011) Neurosteroids reduce social isolation-induced behavioral deficits: A proposed link with neurosteroid-mediated upregulation of BDNF expression. Front Endocrinol (Lausanne) 2:Article 73. doi: 10.3389/fendo.2011.00073

Ottoni EB (2000) EthoLog 2.2: A tool for the transcription and timing of behavior observation sessions. Behav Res Methods 32:446–449.

Papadopoulos V, Berkovich A, Krueger KE, et al (1991) Diazepam binding inhibitor and its processing products stimulate mitochondrial steroid biosynthesis via an interaction with mitochondrial benzodiazepine receptors. Endocrinology 129:1481–1488. doi: 10.1016/S0028-3908(11)80011-0

Papadopoulos V, Lecanu L, Brown RC, et al (2006) Peripheral-type benzodiazepine receptor in neurosteroid biosynthesis, neuropathology and neurological disorders. Neuroscience 138:749–756. doi: 10.1016/j.neuroscience.2005.05.063

Perna G, Schruers K, Alciati A, Caldirola D (2014) Novel investigational therapeutics for panic disorder. Expert Opin Investig Drugs. doi: 10.1517/13543784.2014.996286

Piato ÂL, Rosemberg DB (2014) Princípios éticos no uso do peixe-zebra como organismo-modelo na pesquisa científica. In: Pichler NA, Giacomini ACVV (eds) Ética em Pesqui. com animais e humanos. Bem-estar e Dign. UPF Editora, Passo Fundo, pp 63–73

Pinna G, Rasmusson AM (2012) Up-regulation of neurosteroid biosynthesis as a pharmacological strategy to improve behavioural deficits in a putative mouse model of post-traumatic stress disorder. J Neuroendocrinol 24:102–116. doi: 10.1111/j.1365-2826.2011.02234.x

Romeo E, Cavallaro S, Korneyev A, et al (1993) Stimulation of brain steroidogenesis by 2-aryl-indole-3-acetamide derivatives acting at the mitochondrial diazepam-biding inhibitor complex. J Pharmacol Exp Ther 267:462–470.

Sackerman J, Donegan JJ, Cunningham CS, et al (2010) Zebrafish behavior in novel environments: Effects of acute exposure to anxiolytic compounds and choice of Danio rerio line. Int J Comp Psychol 23:43–61.

Sanger TJ, Hime PM, Johnson MA, et al (2008) Laboratory protocols for husbandry and embryo collection of Anolis lizards. Herpetol Rev 39:58–63.

Smith SS, Shen H, Gong QH, Zhou X (2009) Neurosteroid regulation of GABAA receptors: Focus
on the α4 and δ subunits. Pharmacol Ther 115:1069–1083. doi: 10.1037/a0013572.An

Ugale RR, Mittal N, Hirani K, Chopde CT (2004) Essentiality of central GABAergic neuroactive
steroid allopregnanolone for anticonvulsant action of fluoxetine against pentylenetetrazole-
induced seizures in mice. Brain Res 1023:102–111. doi: 10.1016/j.brainres.2004.07.018

Vada S, Goli D, Bose A, Sharma UR (2015) Effects of anti anxiety drug midazolam as pretreatment
therapy for anxiety of zebra fishes induced by the exposure of unfamiliar environment of
aquatic white/black maze. Asian J Pharm Clin Res 8:3–6.

Warnick JE, Huang CJ, Acevedo EO, Sufka KJ (2009) Modelling the anxiety-depression continuum
in chicks. J Psychopharmacol 23:143–156. doi: 10.1177/0269881108089805

Wong RY, Oxendine SE, Godwin J (2013) Behavioral and neurogenomic transcriptome changes in
wild-derived zebrafish with fluoxetine treatment. BMC Genomics 14:348. doi: 10.1186/1471-
2164-14-348

Figure captions

Figure 1 – Effects of FGIN-1-27 (filled circles) and diazepam (empty circles) on (A) scototaxis, (B)

risk assessment, (C) thigmotaxis, (D) freezing, (E) erratic swimming, and (F) total locomotion in

the light/dark test in zebrafish (*Danio rerio*). Error bars represent the standard errors.

Figure 2 – Effects of FGIN-1-27 (filled circles) and diazepam (empty circles) on (A) tonic immo-

bility, (B) circling responses, (C) freezing, (D) ventilatory frequency, (E) tongue-flicking, and (F)

total locomotion on wall lizards (*Tropidurus oreadicus*). Error bars represent the standard errors.