Real-time Zika risk assessment in the United States

Lauren A Castro*,[1], Spencer J Fox*,[1][@], Xi Chen[2], Kai Liu[3], Steve Bellan[4], Nedialko B Dimitrov[2], Alison

P Galvani[5,6], Lauren Ancel Meyers[1,7]

Affiliations:

1 - Department of Integrative Biology, The University of Texas at Austin, Austin, TX, USA

2 - Graduate Program in Operations Research Industrial Engineering, The University of Texas at Austin,

Austin, TX, USA

3 - Institute for Cellular and Molecular Biology, The University of Texas at Austin, Austin, TX, USA

4 - Center for Computational Biology and Bioinformatics, The University of Texas at Austin, Austin, TX,

USA

5 - Center for Infectious Disease Modeling and Analysis, Yale School of Public Health, New Haven, CT,

USA

6 - Department of Ecology and Evolution, Yale University, New Haven, CT, USA

7 - The Santa Fe Institute, Santa Fe, NM, USA

* contributed equally to this manuscript

[@] corresponding author: spncrfx@gmail.com

Abstract

Recent Zika Virus (ZIKV) outbreaks in southern Florida have heightened public health concern

across the southern United States. As autochthonous (locally-acquired) cases accumulate within the US,

policymakers seek early and accurate indicators of self-sustaining transmission to inform intervention

efforts in high risk areas. However, given ZIKV's low reporting rates and the geographic variability in

both importations and transmission potential, a small cluster of reported cases may reflect diverse

scenarios, ranging from multiple self-limiting but independent introductions to a self-sustaining local

epidemic. We developed a stochastic model that captures variation and uncertainty in ZIKV case

reporting, importations, and transmission, and applied it to assess county-level epidemic risk throughout

the state of Texas. For each of the 254 counties, we estimated the future epidemic risk as a function of

reported autochthonous cases and evaluated a national recommendation to trigger interventions

immediately following the first two reported cases of locally-transmitted ZIKV. Our analysis suggests

that the regions of greatest risk for sustained ZIKV transmission include 21 Texas counties along the

Texas-Mexico border, in the Houston Metro Area, and throughout the I-35 Corridor from San Antonio to

Waco. Variation in vector habitat suitability and importation risk drives epidemic risk variation. Upon

detection of two locally transmitted cases in a spatiotemporal cluster, the threat of epidemic expansion

depends critically on local vulnerability. For high risk Texas counties, we estimate this likelihood to be

64%, assuming an August 2016 risk projection and a 20% reporting rate. With reliable estimates of key

epidemiological parameters, including reporting rates and vector abundance, this framework can help

optimize the timing and spatial allocation of public health resources to fight ZIKV in the US.

Author Summary

Given the growing threat of Zika Virus (ZIKV) in the southern US and the critical importance of

early intervention, public health decision makers seek early and accurate indicators of local transmission.

However, given ZIKV's low reporting rates and the geographic variability in both importations and

transmission potential, a small cluster of reported cases may reflect diverse scenarios, ranging from

multiple self-limiting but independent introductions to a self-sustaining local epidemic. To support real-

time risk assessment of emerging ZIKV outbreaks at a county level, we developed a quantitative

framework that estimates current ZIKV burden and future epidemic threat based on recent reported cases

and the underlying risks of ZIKV importation and transmission. We assessed ZIKV risk in Texas and

found that 21 of the 254 counties are potentially vulnerable to locally-transmitted ZIKV outbreaks. The

high risk region includes much of the Texas-Mexico border, the Houston Metro Area, and patches along

the I-35 Corridor from San Antonio to Waco. If two locally transmitted and epidemiologically linked

cases are detected in one of these counties, we estimate a 64% risk of imminent epidemic expansion.

Introduction

In February 2016, the World Health Organization (WHO) declared Zika virus (ZIKV) a Public

Health Emergency of International Concern [1]. As of 18 August of 2016, the WHO confirmed

mosquito-transmitted cases in 70 countries and territories, with over 500,000 suspected and confirmed

cases in the Americas alone [2,3]. In the US, the 46 reported autochthonous (local) ZIKV cases all

occurred in Southern Florida, but the potential range of a primary ZIKV vectors, *Aedes aegypti*, may

include over 30 states [4,5]. Texas is a particularly vulnerable state, given its suitable climate,

international airports, and geographical proximity to affected countries [4,6–10]. Of the 2,487 imported

ZIKV cases in the US that have been identified, 137 have occurred in Texas through the end of August.

While these importations have yet to spark autochthonous transmission within the state, Texas has

historically sustained several small, autochthonous outbreaks (ranging from 4 - 25 confirmed cases) of

another arbovirus vectored by *Ae. Aegypti*—dengue (DENV) [11–13].

As peak mosquito season continues in the US and more cases are introduced via international

travelers from the Americas, public health decision makers will face considerable uncertainty in gauging

the severity of the threat and in effectively initiating interventions, given the large fraction of undetected

ZIKV cases (asymptomatic and symptomatic) as well as the challenge in estimating the economic tradeoff

of disease prevention versus disease response [14–17]. Depending on the ZIKV symptomatic fraction,

reliability and rapidity of diagnostics, importation rate, and transmission rate, the detection of five

autochthonous cases in a single Texas county, for example, may indicate a small chain of cases from a

single importation, a self-limiting outbreak, or even a large, hidden epidemic underway (Fig 1). These

diverging possibilities have historical precedents. In French Polynesia, a handful of suspected ZIKV cases

were reported by October 2013; two months later an estimated 14,000-29,000 individuals had been

infected [14,15]. By contrast, Anguilla had 17 confirmed cases from late 2015 into 2016 without a

subsequent epidemic, while many surrounding countries experienced large ZIKV epidemics [18]. To

address the uncertainty, the CDC recently issued conservative guidelines for state and local agencies; they

recommend initiation of public health responses following local reporting of *two* non-familial

autochthonous ZIKV cases [19].

Fig 1. ZIKV emergence scenarios. A ZIKV infection could spark (A) a self-limiting outbreak or (B) a growing epidemic. Cases

are partitioned into symptomatic (grey) and asymptomatic (black). Arrows indicate new ZIKV importations by infected travelers

and vertical dashed lines indicate case reporting events. On the 75th day, these divergent scenarios are almost indistinguishable to

public health surveillance, as exactly three cases have been detected in both. By the 100th day, the outbreak (A) has died out with

21 total infections while the epidemic (B) continues to grow with already 67 total infections. Each scenario is a single stochastic

realization of the model with R_0=1.1, reporting rate of 10%, and introduction rate of 0.1 case/day.

Here, we develop a model to support real-time ZIKV risk assessment that accounts for

uncertainty regarding ZIKV epidemiology, including importation rates, reporting rates, and local vector

population density. This framework can be readily updated as our understanding of ZIKV evolves to

provide situational awareness and actionable guidance for public health officials. By simulating ZIKV

transmission using a stochastic branching process model [20] based on recent ZIKV data, we estimated

ZIKV epidemic risk for each of the 254 counties in Texas, both initially and as a function of the

cumulative number of reported autochthonous ZIKV cases. Further, we estimate county-level epidemic

risks corresponding to the recommended two-case trigger and demonstrate the design of county-specific

triggers indicative of imminent epidemic expansion. Our results suggest that counties along the Texas-

Mexico border, in the Houston Metro Area, and throughout the I-35 Corridor from San Antonio to Waco

are at highest risk for sustained epidemics.

Methods

Risk-assessment methodological overview

In our study, we first estimated county-level ZIKV importation rates and sustained transmission

risk (R_0) for Texas in August of 2016. For each county, we then used these estimates to simulate 10,000

stochastic ZIKV outbreaks. Finally, we analyzed the simulated outbreaks to determine spatiotemporal

variation in ZIKV risk and estimate current and future epidemic risk from the number of reported

autochthonous cases (Fig S1).

Estimating County Importation Rates

Our analysis assumes that any ZIKV outbreaks in Texas originate with infected travelers

returning from regions with ZIKV activity. To estimate the ZIKV importation rate for specific counties,

we (1) estimated the Texas statewide importation rate (expected number of imported cases per day), (2)

estimated the probability that the next Texas import will arrive in each county, and (3) took the product of

the state importation rate and each county importation probability.

1. During the first quarter of 2016, 27 travel-associated cases of ZIKV were reported in Texas, with

11 occurring in Houston's Harris County [13], yielding a baseline first quarter estimate of 0.3

imported cases per day throughout Texas. In 2014 and 2015, arbovirus introductions into Texas

were threefold higher during the third quarter than the first quarter of each year, perhaps driven

by seasonal increases in arbovirus activity in endemic regions and the approximately 40%

increase from quarter 1 to quarter 3 in total number of international travelers to the US [21].

Taking this as a *baseline* (lower bound) scenario, we projected a corresponding increase in ZIKV

importations to 0.9 cases per day (statewide) for the third quarter. We also considered an *elevated*

137   importation scenario, in which the first quarter cases (27) in Texas represent only the

138   symptomatic (20%) imported cases, corresponding to a projected third quarter statewide

139   importation rate of 4.5 cases per day.

140   2. We defined county-specific import risk as the probability that the next imported case in Texas

141   will occur in that county. To build a predictive model for import risk, we fit a probabilistic model

142   (maximum entropy) [22] of importation risk to 183 DENV, 38 CHIKV, and 31 ZIKV Texas

143   importations reported at the county level from 2002 to 2016 and informative socioeconomic,

144   environmental, and travel variables (Supplement §1.1). Given the geographic and biological

145   overlap between ZIKV, DENV and Chikungunya (CHIKV), we used historical DENV and

146   CHIKV importation data to supplement ZIKV importations in the importation risk model, while

147   recognizing that future ZIKV importations may be fueled by large epidemic waves in neighboring

148   regions and summer travel, and thus far exceed recent DENV and CHIKV importations [23].

149   Currently, DENV, CHIKV, and ZIKV importation patterns differ most noticeably along the

150   Texas-Mexico border. Endemic DENV transmission and sporadic CHIKV outbreaks in Mexico

151   historically have spilled over into neighboring Texas counties. In contrast, ZIKV is not yet as

152   widespread in Mexico as it is in Central and South America, with only one reported ZIKV

153   importation along the border to date (Val Verde County). We included DENV and CHIKV

154   importation data in the model fitting so as to consider potential future importation pressure from

155   Mexico, as ZIKV continues its increasing trend since March 2016 [24]. To find informative

156   predictors for ZIKV importation risk, we analyzed 72 socio-economic, environmental, and travel

157   variables, and removed near duplicate variables and those that contributed least to model

158   performance, based on out-of-sample cross validation [25,26], reducing the original set of 72

159   variables to 10 (Tables S3-S4).

<u>Estimating County Transmission Risk</u>

The risk of ZIKV emergence following an imported case will depend on the likelihood of

mosquito-borne local transmission. For each Texas county, we used the Ross-Macdonald formulation to

estimate the ZIKV reproduction number (R_0), which is the average number of secondary infections caused

by the introduction of a single infectious individual into a fully susceptible population (Supplement §1.2)

[27]. To parameterize the model (Table S5), we used mosquito life history estimates from a combination

of DENV and ZIKV studies and estimates of *Ae. aegypti* abundance for each county [6]. For parameters

that are sensitive to temperature (i.e., mosquito mortality and the extrinsic incubation period), we adjusted

the mosquito parameters using average reported Texas county temperatures for the month of August [25].

Given the uncertainty in using DENV and ZIKV parameters from experimental and field studies from

other localities, we have included a sensitivity analysis in Supplement §4 to explore upper and lower

bounds of absolute values with changes in the average parameter values.

<u>ZIKV Outbreak Simulation Model</u>

To transmit ZIKV, a mosquito must bite an infected human, the mosquito must get infected with

the virus, and then the infected mosquito must bite a susceptible human. We assumed that mosquito-borne

transmission would be the main driver of epidemic dynamics, so we did not include sexual transmission

in our model. Rather than explicitly model the full transmission cycle, we aggregated the two-part cycle

of ZIKV transmission (mosquito-to-human and human-to-mosquito) into a single exposure period where

the individual has been infected by ZIKV, but not yet infectious, and do not explicitly model mosquitos.

For the purposes of this study, we need only ensure that the model produces a realistic human-to-human

generation time of ZIKV transmission.

The resultant model thus follows a Susceptible-Exposed-Infectious-Recovered (SEIR)

transmission process stemming from a single ZIKV infection using a Markov branching process model

(Fig S2). The temporal evolution of the compartments is governed by daily probabilities of infected

individuals transitioning between S, E, I, and R states, and new ZIKV cases arising from importations or

autochthonous transmission (Table S6). We treat days as discrete time steps, and the next disease state

progression depends solely on the current state and the transition probabilities. We assume that infectious

cases cause a Poisson distributed number of secondary cases per day (via human to mosquito to human

transmission), but this assumption can be relaxed as more information regarding the distribution of

secondary cases becomes available. We also assume infectious individuals are introduced daily according

to a Poisson distributed number of cases around the importation rate. Furthermore, *Infectious* cases are

categorized into reported and unreported cases according to a reporting rate. We assume that reporting

rates approximately correspond to the percentage (~20%) of symptomatic ZIKV infections [16] and occur

at the same rate for imported and locally acquired cases. Additionally, we make the simplifying

assumption that reported cases transmit ZIKV at the same rate as unreported cases. We track imported

and autochthonous cases separately, and conduct risk analyses based on reported autochthonous cases

only, under the assumption that public health officials will have immediate and reliable travel histories for

all reported cases [28].

To accurately model the timing of ZIKV outbreaks, we fit the ZIKV generation time to recent

estimates (Supplement §2.4) [29]. The generation time measures the average duration from initial

symptom onset to the subsequent exposure of a secondary case, and is estimated to range from 10 to 23

days for ZIKV [29]. In our model, the generation time corresponds to the exposure period followed by

half of the infectious period. First, we fit the infectious period in our model to human ZIKV estimates for

duration of viral shedding, which we assumed to be the length of the infectious period, but may be an

underestimate of the total length. Specifically, we solved for transition rates of a Boxcar Model [30] that

produce an infectious period with mean duration of 9.88 days (Table S6) [14]. Then, we fit the exposure

period to achieve the empirical ZIKV generation time distribution [29], yielding a mean exposure period

of 10.4 days (95% CI 6-17) and a mean generation time of 15.3 days (95% CI 9.5-23.5). Given that the

exposure period includes human and mosquito incubation periods and mosquito biting rates, this range is

consistent with the estimated 5.9 day human ZIKV incubation period [14]. While there is much

uncertainty regarding ZIKV transmission, our model framework is flexible to extensions and updates as

we learn more about ZIKV epidemiology including reporting rates, transmission probabilities, clinical

features, and outside importation pressures.

Simulations

For each county risk scenario, defined by a specified importation rate, transmission rate, and

reporting rate, we ran 10,000 stochastic simulations. Each simulation began with a single imported

infectious case and terminated either when there were no individuals left in either the *Exposed* or

Infectious classes or the cumulative number of autochthonous infections reached 2,000. We classified

simulations as either epidemics or self-limiting outbreaks; epidemics were all simulations that fulfilled

two criteria: reached 2,000 cumulative autochthonous infections and had a maximum daily prevalence

(which we defined as the number of current infectious cases) exceeding 50 autochthonous cases (Fig S3).

The second criterion distinguishes simulations resulting in large self-sustaining outbreaks (that achieve

substantial peaks) from those that accumulate infections through a series of small, independently sparked

clusters (that fail to reach the daily prevalence threshold). The latter occurs occasionally under low R_0s

and high importation rates scenarios.

For simulations resulting in epidemics, the cumulative reported autochthonous infections may

include cases from several co-occurring or proximate transmission clusters, as might occur in actual

outbreaks. To verify that our simulations do not aggregate cases from clusters with clear temporal

separation, we calculated the distribution of times between sequential cases (Fig S4). In our simulated

epidemics, almost all sequentially occurring cases occur within 14 days of each other, consistent with the

CDC's threshold for identifying local transmission events (based on the estimated maximum duration of

the ZIKV incubation period) [19].

Outbreak Analysis

Policymakers must often make decisions in the face of uncertainty, such as when and where to

initiate ZIKV interventions. Our stochastic framework allows us to provide real-time county-level risk

assessments as reported cases accumulate. For each county, we found the probability that an outbreak will

progress into an epidemic (reach 2,000 cases with a maximum daily prevalence over 50), as a function of

the number of reported cases. We call this *epidemic risk*. To solve for epidemic risk in a county following

the xth reported autochthonous case, we first find all simulations (of 10,000 total) that experience at least

x reported autochthonous cases, and then calculate the proportion of those that are ultimately classified as

epidemics. For example, consider a county in which 1,000 of 10,000 simulated outbreaks reach at least

two reported autochthonous cases; the remaining 9,000 simulations dissipate with only one or zero case

reports. If only 50 of the 1,000 simulations ultimately fulfill the two epidemic criteria, then the estimated

epidemic risk following two reported cases in that county would be 5%. This simple classification scheme

performs quite well, only rarely misclassifying a string of small outbreaks as an epidemic, with the

probability of such an error increasing with the importation rate. For example, epidemics should not occur

when R_0=0.9. If the importation rate is high, however, overlapping series of moderate outbreaks may

occasionally meet the two epidemic criteria. Even under the highest importation rate we considered (0.3

cases/day), only 1% of outbreaks were misclassified.

This method can be applied to evaluate universal triggers (like the recently recommended two-

case trigger) or derive robust triggers based on local importation and transmission risks as well as the risk

tolerance of public health agencies. For example, if a policymaker would like to initiate interventions as

soon as the risk of an epidemic reaches 30%, we would simulate local ZIKV transmission and solve for

the number of reported cases at which the probability of an epidemic first exceeds 30%. Generally, the

recommended triggers decrease as the policymaker threshold for action decreases (for example,

policymakers would act sooner (fewer reported cases) for a 10% versus 30% threshold) and as the local

transmission potential increases (*e.g.* R_0 = 1.5 versus R_0 = 1.2). A policymaker wishing to trigger

interventions early, upon even a low probability of epidemic spread, has a low tolerance for failing to

intervene but may waste resources on unnecessary interventions; a policymaker willing to wait longer,

has a higher risk tolerance, but may implement interventions too late in the course of the outbreak.

County Uncertainty Analysis

We took two approaches to addressing uncertainty in the model parameters. First, we conducted a

sensitivity analysis to address the considerable uncertainty regarding several inputs into our estimation of

R_0, including mosquito biology, ZIKV epidemiology, and human-mosquito interactions (Supplement §4).

For most factors, the county estimates of R_0 simply scale linearly with changes in the factor. However,

county-specific human-mosquito contact rates can change relative county risks based on assumptions

regarding the socioeconomic effect on human-mosquito interactions (Fig S6-7), and county risk moves

southward as the summer heat subsides (Fig S8). Second, given the considerable uncertainty regarding

ZIKV epidemiology, we examined a scenario where the absolute values of both R_0 and importation rate

are unknown, but lie within plausible ranges for Texas. To do so, we randomly sampled 10,000

simulations from the high risk Texas county outbreaks (counties with estimates of R_0>1), creating an

amalgamous high risk county, and completed the outbreak analysis as we do with individual counties.

Results

To develop a ZIKV risk assessment framework for Texas counties, we first estimate county-level

ZIKV importation and transmission rates for August 2016. ZIKV importation risk within Texas is

predicted by variables reflecting urbanization, mobility patterns, and socioeconomic status (Table S3),

and is concentrated in metropolitan counties of Texas (Fig 2A). The two highest risk counties--Harris,

which includes Houston and has an estimated 27% chance of receiving the next imported Texas case, and

Travis, which includes Austin and has a 10% chance of the next importation--contain international

airports. Other high risk regions include Brazos County, the Dallas and San Antonio metropolitan areas,

and several counties along the Texas-Mexico border.

Fig 2. ZIKV importation and transmission risk estimates across Texas. (A) Color indicates the probability that the next ZIKV import will occur in a given county for each of the 254 Texas counties. Probability is colored on a log scale. The 10 most populous cities in Texas are labeled. Houston's Harris County has 2.7 times greater chance than Austin's Travis County of receiving the next imported case. **(B)** Estimated county-level R_0 for ZIKV. Counties with estimated R_0s ≤ 1 are shades of green, and yellow increasing to red counties indicate R_0s > 1 (See Fig S7-11 for sensitivity analysis).

Our county-level estimates of autochthonous ZIKV transmission risk (Fig 2B) suggest that the majority of Texas counties (87%) have an estimated R_0 below one, and thus are unlikely to sustain epidemics. The Southeast region of Texas has the highest estimated transmission risk, driven primarily by high mosquito habitat suitability. These estimates are sensitive to uncertainty in several parameters (Fig S7-11), and can be updated as we learn more about ZIKV. While the average transmission risk may be higher or lower than our baseline assumption, and certainly varies seasonally, the relative high and low risks areas of the state are robust (Fig S7-11), and allow us to conduct plausible case studies and identify at risk areas for enhanced surveillance and preparedness efforts. However given the uncertainty underlying specific county R_0 estimates, we also aggregate the 21 highest absolute estimates into a plausible distribution for any high risk Texas county, ranging from $R_0 = 1.0$ to $R_0 = 2.2$ with a median of 1.1.

Under a single set of epidemiological conditions, wide ranges of outbreaks are possible (Fig 3A).

The relationship between what policymakers can observe (cumulative reported cases) and what they wish

to know (current underlying disease prevalence) can be obscured by such uncertainty, and will depend

critically on both the transmission and reporting rates (Fig 3B). If key drivers, such as R_0, can be

estimated with confidence, then the breadth of possibilities narrows, enabling more precise surveillance.

For example, under a known high risk R_0 scenario and with a 20% reporting rate, ten linked cumulative

autochthonous reported cases corresponds to 6 currently circulating cases with a 95% CI of 1-16; under

an unknown but high risk R_0 scenario, the same number of cases corresponds to an expected daily

prevalence of 10 cases with a much wider 95% CI of 2-32 (Fig 3B).

Fig 3. Real-time risk-assessment for ZIKV situational awareness and forecasting. (A) Simulated outbreaks, assuming an

importation rate of 0.01 case per day, for a known high risk R_0 (blue) or an unknown high risk R_0 (red). Two thousand randomly

sampled simulations are shown for each scenario. (B) Total number of (current) autochthonous cases as a function of the

cumulative reported autochthonous cases, assuming an importation rate of 0.01 case per day, for a known R_0 (blue) or an

unknown high-risk R_0 (red), and a relatively high (dashed) or low (solid) reporting rate. Ribbons indicate 50% quantiles. (C) The

increasing probability of imminent epidemic expansion as reported autochthonous cases accumulate, compared across the high

risk (red) and known moderate risk (blue) for a low (solid) and high (dashed) reporting rate. Suppose cases arise in a high risk

county and a policymaker wishes to trigger a response as soon as two cases are reported (vertical line). With a 20% reporting rate

there is a 64% probability of an ensuing epidemic.

We apply our model to characterize time-varying epidemic risk as cases accumulate in a given

county. Under both a known and unknown high risk scenario, we track the probability of epidemic

expansion following each additional reported case (Fig 3C). Across the full range of reported cases, the

probability of epidemic spread is always higher in the unknown, compared with the known high risk

scenario showing more sensitivity to the reporting rate. These curves can support both real-time risk

assessment as cases accumulate and the identification of surveillance triggers indicating when risk

exceeds a specified threshold. For example, suppose a policymaker wanted to initiate an intervention

upon two reported cases and thought the reporting rate was 20%, this would correspond with a 64%

probability of an epidemic in the unknown high risk county but only 35% in the known high risk county.

Alternatively suppose a policy maker wishes to initiate an intervention when the chance of an epidemic

exceeds 50%. In the unknown high risk scenario, they should act immediately following the 1st

autochthonous reported case; in the known high risk scenario, the corresponding trigger ranges from two

to seven autochthonous reported cases, depending on the reporting rate. As the policymaker's threshold

(risk tolerance) increases, the recommended surveillance triggers can be adjusted accordingly.

To evaluate a universal intervention trigger of two reported autochthonous cases, we estimate

both the probability of a trigger event (two such cases) in each county and the level of epidemic risk at the

moment a trigger event occurs (second case reported) in each county. Assuming a baseline importation

rate extrapolated from recent importations to August 2016 and a 20% reporting rate, only a minority of

counties are likely to experience a trigger event (Fig 4A). While 231 of the 254 counties (91%) have non-

zero probabilities of experiencing two reported autochthonous cases, only 63 counties have at least a 10%

chance of such an event, with the remaining 168 counties having a median probability of 0.017 (range

0.0004 to 0.089). Next, assuming that a second autochthonous case has indeed been reported, we find that

the underlying epidemic risk varies widely among the 231 counties, with most counties having near zero

epidemic probabilities and a few counties far exceeding a 50% chance of epidemic expansion. For

example, two reported autochthonous cases in Starr County, along the Texas-Mexico border, correspond

to a 99% chance of ongoing transmission that would proceed to epidemic proportions without

intervention. The greater San Antonio metropolitan region appears to be the highest risk metropolitan

region with four of its eight counties having a higher than 25% probability of experiencing two reported

autochthonous cases; in those four counties, the epidemic risk upon detection of a second case ranges

from 19-90%. Houston metropolitan region is also a high risk region with its second (Fort Bend) and

fourth (Brazoria) largest counties having a 39% and 45% chance of sustaining two reported

autochthonous cases, respectively, with corresponding epidemic risks of 67% and 86% thereafter.

Fig 4. Texas county ZIKV risk assessment. (A) Probability of an outbreak with at least two reported autochthonous ZIKV

cases. **(B)** The probability of epidemic expansion at the moment the second autochthonous ZIKV case is reported in a county.

White counties never reach two reported cases across all 10,000 simulated outbreaks; light gray counties reach two cases, but

never experience epidemics. **(C)** Recommended county-level surveillance triggers (number of reported autochthonous cases)

indicating that the probability of epidemic expansion has exceeded 50%. White counties indicate that fewer than 1% of the

10,000 simulated outbreaks reached two reported cases. All three maps assume a 20% reporting rate and a baseline importation

scenario for August 2016 (81 cases statewide per 90 days) projected from historical arbovirus data. (Figure S5 provides

corresponding estimates under a worse case elevated importation scenario).

Given that a universal trigger may signal highly disparate levels of ZIKV risk, policy makers

might seek to adapt their triggers to local conditions. Suppose a policymaker wishes to design triggers

that indicate a 50% chance of an emerging epidemic (Fig 4C). Under the baseline importation and

reporting rates, only 21 of the 254 counties in Texas are expected to reach a 50% epidemic probability,

with triggers ranging from one (Starr County) to 21 (Dimmit County) reported autochthonous cases, with

a median of two cases. The remaining counties have less than a 1% chance of experiencing sustained

ZIKV transmission. Under an elevated importation scenario, assuming that only one fifth of ZIKV

importations (the symptomatic proportion) have been observed, we find that the recommended triggers

decrease by a mean of 1.5 reported cases (Fig S5) and the size of Texas' population at risk for sustained

ZIKV transmission is expected to increase from ~14% to ~30%, largely driven by increased risk in the

Houston metropolitan area.

Discussion

US public health authorities are responding to ZIKV outbreaks in Southern Florida, and preparing

for possible ZIKV outbreaks in other vulnerable regions. A key challenge is knowing when and where to

initiate interventions based on potentially sparse and biased ZIKV case reports. Our simple model is

designed to address this challenge by providing county-specific epidemic risk estimates as a function of

reported cases. We demonstrate its application across the 254 ecologically and demographically diverse

counties of Texas, a high risk state for ZIKV [7,8,10]. Based on county-level estimates for ZIKV

importation and transmission rates (Fig 2), we expect that most Texas counties are not at risk for a

sustained ZIKV epidemic (Fig 4). However, 30% of Texas' population may reside in vulnerable regions,

including the cities of Austin, San Antonio, and Waco along the I-35 corridor, Houston, and the Rio

Grande Valley. The higher the ZIKV importation rate in these locations, the higher the chance of an

epidemic (Fig S5). However, even in the most high risk regions of Texas, we expect far more limited

ZIKV transmission than observed in Central America, South America, and Puerto Rico, where R_0 has

been estimated to be as high as 11 [27,31,32]. Our analysis is consistent with recent introductions of

DENV and CHIKV into Texas that have caused small pockets of local transmission but have failed to

spark large epidemics.

Surveillance triggers--guidelines specifying situations that warrant intervention--are a key

component of many public health response plans. Given the urgency and uncertainty surrounding ZIKV,

universal recommendations can be both pragmatic and judicious. In choosing an appropriate trigger for an

environmentally and socioeconomically diverse region, policymakers must weigh the risks of acting

unnecessarily with those of responding too late. A risk averse policymaker will likely select a low trigger

(a small number of reported cases), ensuring early detection in the riskiest sites at the cost of false alarms

in low risk locations (Fig 4C). In that vein, the CDC recently issued conservative guidelines to state and

local public health agencies that suggest a two-case trigger for initiating interventions [28]. To assist

Texas policymakers in interpreting this recommendation, we assessed the likelihood and implication of a

two-case trigger for each of Texas' 254 counties, under a scenario projected from recent ZIKV data to

August 2016. Across counties, there is enormous variation in both the chance of a trigger and the

magnitude of the public health threat if and when two cases are reported. If and when two autochthonous

ZIKV cases are reported, only 18 of Texas' 254 counties have over a 20% chance of experiencing

sustained transmission (under our baseline scenario); in most of the remaining 236 counties, the threat is

much lower.

Rather than implement a universal *trigger,* which may indicate different threats in different

locations, one could design local surveillance triggers that correspond to a universal *risk threshold*. Our

modeling framework can readily identify triggers (numbers of reported cases) for indicating any specified

epidemic event (e.g., prevalence reaching a threshold or imminent epidemic expansion) with any

specified risk tolerance (e.g., 10% or 50% chance of that the event has occurred), given local

epidemiological conditions. As a case study, we identify *epidemic expansion* triggers in each of Texas'

254 counties, each designed to indicate when the probability of an epidemic exceeds 50%. Across the 21

counties with non-negligible probabilities of an epidemic, the recommended triggers range from one to 21

reported autochthonous cases, highlighting Texas' spatial risk heterogeneity. These findings apply only to

the early, pre-epidemic phase of ZIKV in Texas, when travel from affected regions outside the contiguous

US is the primary importation source, and as the epidemiological situation in those affected regions

remains constant. If self-sustaining outbreaks emerge within the United States, there may be county-to-

county or state-to-state importations, particularly in high risk regions, that are not yet included in the

model. For example, if the frequency of travel-related ZIKV importations from Florida increases, like the

imported case in El Paso, our model will need to be recalibrated.

This simple framework offers a flexible means for bringing current data and expert knowledge to

assist critical public health decision making. The design of a trigger--both the event to be detected and the

probability threshold upon which to take action--requires extensive public health expertise and

deliberation. Our case studies were motivated by formal recommendations and informal discussions with

state and national public health agencies. They demonstrate our data-driven approach for relating

observed disease activity to underlying risk in the face of great uncertainty, and address the pros, cons and

complexity of developing universal versus regional action plans. However, they are not meant to critique

or validate any specific policies, and highlight the prudence of recent ZIKV planning efforts.

Importantly, our analyses rest on the recent and limited scientific investigations of ZIKV's

biology and epidemiology, and should be continually updated as our understanding matures. Specifically,

our county-level estimates of R_0 are sensitive to many underlying assumptions, particularly regarding

temperature-dependent and socioeconomic factors (Supplement §4). Although the risk of ZIKV

transmission may be much higher or lower than assumed in our baseline scenario, the relative

vulnerabilities of counties are fairly robust to our assumptions. Thus, counties with the highest estimated

risks should likely be prioritized for surveillance and interventions resources. Given the minimal

incursions of DENV and CHIKV into Texas, we suspect that, if anything, we may be underestimating the

socioeconomic and behavioral impediments to ZIKV transmission in the contiguous US, and thus

overestimating the transmission risk of ZIKV across Texas. Given current understanding of ZIKV, we

believe that our county-level estimates of R_0 are reasonable, but should be interpreted and applied with

full consideration of their underlying uncertainty. As detailed ZIKV outbreak data become available, the

research community will continue to improve accuracy and reduce uncertainty in such estimates, which

can be readily incorporated into this risk assessment framework.

The reporting rate dictates the relationship between observed cases and the underlying outbreak,

and its magnitude impacts the timeliness and precision of detection. If only a small fraction of cases are

reported, the first few reported cases may correspond to a wide range of underlying epidemiological

conditions, from isolated introductions to a growing epidemic. In contrast, if most cases are reported,

policymakers can wait longer (in terms of the number of reported cases) to trigger interventions and have

more confidence in their epidemiological assessments. ZIKV reporting rates are expected to remain quite

low, because an estimated 80% of infections are asymptomatic, and DENV reporting rates have

historically matched its asymptomatic proportion [16,33]. Obtaining a realistic estimate of the ZIKV

reporting rate is arguably as important as increasing the rate itself, with respect to reliable situational

awareness and forecasting. An estimated 8-22% of ZIKV infections were reported during the 2013-2014

outbreak in French Polynesia [32]; however estimates ranging from 1 to 10% have been reported during

the ongoing epidemic in Columbia [31,34]. While these provide a baseline estimate for the US, there are

many factors that could increase (or decrease) the reporting rate, such as ZIKV awareness among both the

public and health-care practitioners, or active surveillance of regions with recent ZIKV cases (our

analysis assumes continuous passive surveillance). Thus, rapid estimation of the reporting rate should be a

high priority. While some methods require extensive epidemiological data not typically available early in

an outbreak [35], a new method exploiting early outbreak viral sequence data was introduced during the

recent West African Ebola epidemic [36]. However, as of July 2016, there are no US ZIKV sequences

available on GenBank and few available from other regions.

Unlike other risk estimation methods, this framework is designed to support risk assessments

both prior to ZIKV outbreaks and, in real-time, upon the early detection of cases through the lens of

potentially sparse and biased surveillance data [37,38]. It can facilitate situational awareness, allowing

analysts to translate case counts into estimates of overall prevalence and the potential for future spread,

albeit with large error bars. It can also support the development of response plans, by forcing

policymakers to be explicit about risk tolerance, that is, the certainty needed before sounding an alarm,

and quantifying the consequences of premature or delayed interventions. For example, should ZIKV-

related pregnancy advisories be issued when there is only 5% chance of an impending epidemic? 10%

chance? 80%? A policymaker has to weigh the costs of false positives--resulting in unnecessary fear

and/or intervention--and false negatives--resulting in suboptimal disease control and prevention--

complicated by the difficulty inherent in distinguishing a false positive from a successful intervention.

The more risk averse the policymaker (with respect to false negatives), the earlier the trigger should be,

which can be exacerbated by low reporting rates, high importation rate, and inherent ZIKV transmission

potential. In ZIKV prone regions with low reporting rates, even risk tolerant policymakers should act

quickly upon seeing initial cases; in lower risk regions, longer waiting periods may be prudent.

Acknowledgements

We acknowledge the Texas Department of State Health Services (DSHS) for providing historic

arbovirus county importation data, and MUG Kraemer and A Perkins for providing Texas mosquito

abundance data and technical guidance. We also thank M Johansson and M Meltzer of the Centers for

Disease Control and Prevention (CDC) and Dr. John Hellerstedt, commissioner of the Texas DSHS, for

critical conversations regarding the modeling framework and its application to Texas and national ZIKV

planning efforts. We finally acknowledge the Texas Advanced Computing Center (TACC) at The

University of Texas at Austin for providing High performance computing resources that have contributed

to the research results reported within this paper. URL: http://www.tacc.utexas.edu.

Code Availability

The R package, rtZIKVrisk, has been created to accompany the manuscript and can be found

here: https://github.com/sjfox/rtZIKVrisk. The package provides functionality to simulate and analyze

ZIKV outbreaks according to our framework, and recreate the figures presented in the manuscript.

References

1. Gulland A. Zika virus is a global public health emergency, declares WHO. BMJ. BMJ Publishing

Group Ltd; 2016;352. doi:10.1136/bmj.i657

2. PAHO, WHO. Suspected and confirmed Zika cases reported by countries and territories in the

Americas, 2015-2016 [Internet]. 2016 p. http://ais.paho.org/phip/viz/ed_zika_epicurve.asp.

3. World Health Organization. Zika situation report: 18 August 2016. 2016.

4.   About estimated range of Aedes aegypti and Aedes albopictus in the United States, 2016 Maps. In:

Atlanta: Centers for Disease Control and Prevention [Internet]. [cited 1 Jan 2016]. Available:

http://www.cdc.gov/zika/vector/range.html

5.   Florida Department of Health. Department of Health Daily Zika Update - August 30, 2016. 2016.

6.   Kraemer MUG, Sinka ME, Duda KA, Mylne A, Shearer FM, Barker CM, et al. The global

distribution of the arbovirus vectors Aedes aegypti and Ae. albopictus. Elife. eLife Sciences

Publications Limited; 2015;4: e08347. Available: http://elifesciences.org/content/4/e08347v3

7.   Messina JP, Kraemer MU, Brady OJ, Pigott DM, Shearer FM, Weiss DJ, et al. Mapping global

environmental suitability for Zika virus. Elife. eLife Sciences Publications Limited; 2016;5:

e15272. Available: https://elifesciences.org/content/5/e15272

8.   Gardner LM, Chen N, Sarkar S. Global risk of Zika virus depends critically on vector status of

Aedes albopictus. Lancet Infect Dis. Elsevier; 2016;16: 522–523. Available:

http://www.thelancet.com/article/S1473309916001766/fulltext

9.   Peterson AT, Osorio J, Qiao H, Escobar LE. Zika virus, elevation, and transmission risk. PLoS

Curr Outbreaks. 2016;9. doi:10.1371/currents.outbreaks.a832cf06c4bf89fb2e15cb29d374f9de

10.   Monaghan AJ, Morin CW, Steinhoff DF, Wilhelmi O, Hayden M, Quattrochi DA, et al. On the

Seasonal Occurrence and Abundance of the Zika Virus Vector Mosquito Aedes Aegypti in the

Contiguous United States. PLoS Curr. 2016; 1–31.

doi:10.1371/currents.outbreaks.50dfc7f46798675fc63e7d7da563da76

11.   Texas Department of State Health and Human Services. Arbovirus Activity in Texas 2014

Surveillance Report [Internet]. 2014. Available:

https://www.google.com/url?sa=t&rct=j&q=&esrc=s&source=web&cd=3&ved=0ahUKEwiE8Oz

U0I_OAhVG5SYKHTBTA9QQFggoMAI&url=https://www.dshs.texas.gov/WorkArea/linkit.asp

x?LinkIdentifier=id&ItemID=8589999503&usg=AFQjCNGO9cW43_oOrfmhjWIKseBh9Q1n9g

&s

12.   CDC, Texas Department of State Health and Human Services. Dengue Hemorrhagic Fever ---

U.S.-Mexico Border, 2005 [Internet]. 2007. Available:

http://www.cdc.gov/mmwr/preview/mmwrhtml/mm5631a1.htm

13. Zika in Texas. In: Texas Department of State Health Services [Internet]. 2016 [cited 5 Apr 2016].

Available: http://www.texaszika.org/

14. Lessler JT, Ott CT, Carcelen AC, Konikoff JM, Williamson J, Bi Q, et al. Times to key events in

the course of Zika infection and their implications: a systematic review and pooled analysis. Bull

World Health Organ. 2016; doi:10.2471/BLT.16.174540

15. Lloyd-Smith JO, Funk S, McLean AR, Riley S, Wood JLN. Nine challenges in modelling the

emergence of novel pathogens. Epidemics. Elsevier B.V.; 2014;10: 35–39.

doi:10.1016/j.epidem.2014.09.002

16. Duffy MR, Chen T-H, Hancock WT, Powers AM, Kool JL, Lanciotti RS, et al. Zika Virus

Outbreak on Yap Island, Federated States of Micronesia. N Engl J Med. 2009;360: 2536–2543.

doi:10.1056/NEJMoa0805715

17. Alfaro-Murillo JA, Parpia AS, Fitzpatrick MC, Tamagnan JA, Medlock J, Ndeffo-Mbah ML, et al.

A Cost-Effectiveness Tool for Informing Policies on Zika Virus Control. Carabin H, editor. PLoS

Negl Trop Dis. 2016;10: e0004743. doi:10.1371/journal.pntd.0004743

18. PAHO and WHO. Zika-Epidemiological Report - Anguilla. 2016; 1–2.

19. ZIKA CDC Interim Response Plan [Internet]. 2016. Available: http://www.cdc.gov/zika/pdfs/zika-

draft-interim-conus-plan.pdf

20. Getz WM, Lloyd-Smith JO. Basic methods for modeling the invasion and spread of contagious

diseases. Dis Evol Model concepts, data Anal AMS-DIMACS Ser. 2006;71: 87. Available:

http://books.google.com/books?hl=en&lr=&id=5uMf4qCmghEC&oi=fnd&pg

=PA87&dq=Basic+Methods+for+Modeling+the+Invasion+and+Spread+of+Contagious+Dis

eases&ots=M3miy-

szrR&sig=Re_k7i7YQoXGHkktCbLilsv41XQ\nhttp://books.google.com/books?h

21. Office of Travel & Tourism Industries. US Monthly Arrivals Trend Line [Internet]. 2014.

Available: http://travel.trade.gov/view/m-2014-I-001/index.html

22.   Jaynes ET. Information Theory and Statistical Mechanics. Phys Rev. American Physical Society;

1957;106: 620–630. doi:10.1103/PhysRev.106.620

23.   Musso D, Cao-Lormeau VM, Gubler DJ. Zika virus: following the path of dengue and

chikungunya? Lancet. Elsevier Ltd; 2015;386: 243–244. doi:10.1016/S0140-6736(15)61273-9

24.   WHO P and. Zika-Epidemiological Report - Mexico. 2016.

25.   Wolsey LA. Integer programming. Wiley; 1998.

26.   Merow C, Smith MJ, Silander JA. A practical guide to MaxEnt for modeling species'

distributions: what it does, and why inputs and settings matter. Ecography (Cop). 2013;36: 1058–

1069. doi:10.1111/j.1600-0587.2013.07872.x

27.   Alex Perkins T, Siraj AS, Ruktanonchai CW, Kraemer MUG, Tatem AJ. Model-based projections

of Zika virus infections in childbearing women in the Americas. Nat Microbiol. 2016;1: 16126.

doi:10.1038/nmicrobiol.2016.126

28.   Interim CDC recommendations for Zika vector control in the continental United States. In:

Centers for Disease Control and Prevention [Internet]. 2016 [cited 5 Jan 2016]. Available:

http://www.cdc.gov/zika/public-health-partners/vector-control-us.html

29.   Majumder MS, Cohn E, Fish D, Brownstein JS. Estimating a feasible serial interval range for Zika

fever. Bull World Health Organ. 2016; doi:10.2471/BLT.16.171009

30.   Lloyd AL. Realistic Distributions of Infectious Periods in Epidemic Models: Changing Patterns of

Persistence and Dynamics. Theor Popul Biol. 2001;60: 59–71. doi:10.1006/tpbi.2001.1525

31.   Rojas DP, Dean NE, Yang Y, Kenah E, Quintero J, Tomasi S, et al. The Epidemiology and

Transmissibility of Zika Virus in Girardot and San Andres Island, Colombia. bioRxiv. 2016;

Available: http://biorxiv.org/content/early/2016/04/24/049957.abstract

32.   Kucharski AJ, Funk S, Eggo RM, Mallet H, Edmunds WJ, Nilles EJ. Transmission dynamics of

Zika virus in island populations : a modelling analysis of the 2013 – 14 French Polynesia outbreak.

bioRxiv. 2016; 1–15.

33.   Dechant E, Rigau-Perez J. Hospitalizations for suspected dengue in Puerto Rico, 1991-1995:

583   estimation by capture-recapture methods. The Puerto Rico Association of Epidemiologists. Am J

584   Trop Med Hyg. 1999;61: 574–578. Available: http://www.ajtmh.org/content/61/4/574.short

585   34.   Zhang Q, Sun K, Chinazzi M, Pastore-Piontti A, Dean NE, Rojas DP, et al. Projected spread of

586   Zika virus in the Americas. bioRxiv. 2016; 66456. doi:10.1101/066456

587   35.   Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution

588   and burden of dengue. Nature. Nature Publishing Group; 2013;496: 504–507.

589   doi:10.1038/nature12060

590   36.   Scarpino S V, Iamarino A, Wells C, Yamin D, Ndeffo-Mbah M, Wenzel NS, et al.

591   Epidemiological and viral genomic sequence analysis of the 2014 ebola outbreak reveals clustered

592   transmission. Clin Infect Dis. Oxford University Press; 2015;60: 1079–82. Available:

593   http://cid.oxfordjournals.org/content/early/2015/01/05/cid.ciu1131.full

594   37.   Cauchemez S, Epperson S, Biggerstaff M, Swerdlow D, Finelli L, Ferguson NM. Using Routine

595   Surveillance Data to Estimate the Epidemic Potential of Emerging Zoonoses: Application to the

596   Emergence of US Swine Origin Influenza A H3N2v Virus. Peiris JSM, editor. PLoS Med.

597   2013;10: e1001399. doi:10.1371/journal.pmed.1001399

598   38.   Blumberg S, Lloyd-Smith JO. Comparing methods for estimating R0 from the size distribution of

599   subcritical transmission chains. Epidemics. 2013;5: 131–145. doi:10.1016/j.epidem.2013.05.002

S1 Supporting Information. Supporting information for "Real-time Zika risk assessment in the United

States."

S2 Supporting Data. Raw Texas county arbovirus importation data used for estimating county ZIKV

importation risk. Data provided by Texas DSHS.

S3 Supporting Data. Raw county socioeconomic data used for estimating county ZIKV importation risk.

Data accessed from: https://www.census.gov/programs-surveys/acs/data.html

S4 Supporting Data. Texas county ZIKV risk estimates used as inputs for the ZIKV simulation model.

Outputs of the importation and transmission risk analysis.

610
611