A Common Class of Transcripts with 5'-Intron Depletion, Distinct Early Coding

Sequence Features, and N[1]-Methyladenosine Modification

Can Cenik[1,2], Hon Nian Chua[3,4], Guramrit Singh[2,5,7,8], Abdalla Akef[6], Michael P Snyder[1],

Alexander F. Palazzo[6], Melissa J Moore[2,7,8#], and Frederick P Roth[3,9,10#]

Affiliations:

[1] Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305, USA

[2] Department of Biochemistry and Molecular Pharmacology, University of Massachusetts Medical

School, Worcester, MA 01605, USA

[3] Donnelly Centre and Departments of Molecular Genetics and Computer Science, University of Toronto

and Lunenfeld-Tanenbaum Research Institute, Mt Sinai Hospital, Toronto M5G 1X5, Ontario, Canada

[4] DataRobot, Inc., 61 Chatham St. Boston MA 02109, USA

[5] Department of Molecular Genetics, Center for RNA Biology, The Ohio State University, Columbus, OH

43210, USA

[6] Department of Biochemistry, University of Toronto, 1 King's College Circle, MSB Room 5336, Toronto,

ON M5S 1A8, Canada

[7] Howard Hughes Medical Institute, University of Massachusetts Medical School, Worcester, MA 01605,

USA

[8] RNA Therapeutics Institute, University of Massachusetts Medical School, Worcester, MA 01605, USA

[9] Center for Cancer Systems Biology (CCSB), Dana-Farber Cancer Institute, Boston 02215, MA, USA

[10] The Canadian Institute for Advanced Research, Toronto M5G 1Z8, ON, Canada

#Correspondence to: Melissa.Moore@umassmed.edu , Fritz.Roth@utoronto.ca

Short Title: First intron position defines a transcript class

Keywords: 5' UTR introns, random forest, N[1] methyladenosine, Exon Junction Complex

Can Cenik

Abstract:

Introns are found in 5' untranslated regions (5'UTRs) for 35% of all human transcripts.

These 5'UTR introns are not randomly distributed: genes that encode secreted, membrane-

bound and mitochondrial proteins are less likely to have them. Curiously, transcripts lacking

5'UTR introns tend to harbor specific RNA sequence elements in their early coding regions. To

model and understand the connection between coding-region sequence and 5'UTR intron

status, we developed a classifier that can predict 5'UTR intron status with >80% accuracy

using only sequence features in the early coding region. Thus, the classifier identifies

transcripts with <u>5</u>' proximal-<u>i</u>ntron-<u>m</u>inus-like-coding regions ("5IM" transcripts).

Unexpectedly, we found that the early coding sequence features defining 5IM transcripts are

widespread, appearing in 21% of all human RefSeq transcripts. The 5IM class of transcripts is

enriched for non-AUG start codons, more extensive secondary structure both preceding the

start codon and near the 5' cap, greater dependence on eIF4E for translation, and association

with ER-proximal ribosomes. 5IM transcripts are bound by the Exon Junction Complex (EJC) at

non-canonical 5' proximal positions. Finally, N^1-methyladenosines are specifically enriched in

the early coding regions of 5IM transcripts. Taken together, our analyses point to the existence

of a distinct 5IM class comprising ~20% of human transcripts. This class is defined by

depletion of 5' proximal introns, presence of specific RNA sequence features associated with

low translation efficiency, N^1-methyladenosines in the early coding region, and enrichment for

non-canonical binding by the Exon Junction Complex.

Introduction:

Approximately 35% of all human transcripts harbor introns in their 5' untranslated

regions (5'UTRs) (Cenik et al. 2010; Hong et al. 2006). Among genes with 5'UTR introns (5UIs),

those annotated as "regulatory" are significantly overrepresented, while there is an under-

representation of genes encoding proteins that are targeted to either the endoplasmic

reticulum (ER) or mitochondria (Cenik et al. 2011). For transcripts that encode ER- and

mitochondria-targeted proteins, 5UI depletion is associated with presence of specific RNA

sequences (Cenik et al. 2011; Palazzo et al. 2007, 2013). Specifically, nuclear export of an

otherwise inefficiently exported microinjected mRNA or cDNA transcript can be promoted by

an ER-targeting signal sequence-containing region (SSCRs) or mitochondrial signal sequence

coding region (MSCRs) from a gene lacking 5' UTR introns (Cenik et al. 2011; Lee et al. 2015).

However, more recent studies suggest that many SSCRs have little impact on nuclear export for

RNAs transcribed *in vivo* (Lee et al. 2015), but rather enhance translation in a RanBP2-

dependent manner (Mahadevan et al. 2013).

Among SSCR- and MSCR-containing transcripts (referred to hereafter as SSCR and

MSCR transcripts), ~75% lack 5' UTR introns ("5UI⁻" transcripts) and ~25% have them ("5UI⁺"

transcripts). These two groups have markedly different sequence compositions at the 5' ends

of their coding sequences. 5UI⁻ transcripts tend to have lower adenine content (Palazzo et al.

2007) and use codons with fewer uracils and adenines than 5UI⁺ transcripts (Cenik et al.

2011). Their signal sequences also contain leucine and arginine more often than the

biochemically similar amino acids isoleucine and lysine, respectively. Leucine and arginine

codons contain fewer adenine and thymine nucleotides, consistent with adenine and thymine

depletion. This depletion is also associated with the presence of a specific GC-rich RNA motif

in the early coding region of 5UI⁻ transcripts (Cenik et al. 2011).

Despite some knowledge as to their early coding region features, key questions about

this class of 5UI⁻ transcripts have remained unanswered: Do the above sequence features

extend beyond SSCR- and MSCR-containing transcripts to other 5UI⁻ genes? Do 5UI⁻

transcripts having these features share common functional or regulatory features? What

binding factor(s) recognize these RNA elements? A more complete model of the relationship of

early coding features and 5UI⁻ status would begin to address these questions.

Here, to better understand the relationship between early coding region features and

5UI status, we undertook an integrative machine learning approach. We reasoned that a

machine learning classifier which could identify 5UI⁻ transcripts solely from early coding

sequence would potentially provide two types of insight: First, it could systematically identify

predictive features. Second, the subset of 5UI⁻ transcripts which could be identified by the

classifier might then represent a functionally distinct transcript class. Having developed such

a classifier, we found that it identified ~21% of all human transcripts as harboring coding

regions characteristic of 5UI⁻ transcripts. While many of these transcripts encode ER- and

mitochondrial-targeted proteins, many others encode nuclear and cytoplasmic proteins. This

class of transcripts shares characteristics such as a tendency to lack 5' proximal introns, to

contain non-canonical Exon Junction Complex binding sites, to have multiple features

associated with lower intrinsic translation efficiency, and to have an increased incidence of N^1-

methyladenosine modification.

Results:

Can Cenik

A classifier that predicts 5UI status using only early coding sequence information.

To better understand the previously-reported enigmatic relationship between certain early

coding region sequences and the absence of a 5UI, we sought to model this relationship.

Specifically, we used a random forest classifier (Breiman 2001) to learn the relationship

between 5UI absence and a collection of 36 different nucleotide-level features extracted from

the first 99 nucleotides of all human coding regions (CDS) (**Figs 1A-C; Table S1**; Methods).

We then used all transcripts known to contain an SSCR (a total of 3743 transcripts clusters;

Methods), regardless of 5UI status, as our training set. This training constraint ensured that all

input nucleotide sequences were subject to similar functional constraints at the protein level.

Thus, we sought to identify sequence features that differ between 5UI$^-$ and 5UI$^+$ transcripts at

the RNA level.

Our classifier assigns to each transcript a "5'UTR-intron-minus-predictor" (5IMP) score

between 0 and 10, where higher scores correspond to a higher likelihood of being 5UI$^-$ (**Fig**

1C). Interestingly, preliminary ranking of the 5UI$^-$ transcripts by 5IMP score revealed a

relationship between the position of the first intron in the coding region and the 5IMP score.

5UI$^-$ transcripts for which the first intron was more than 85 nts downstream of the start codon

had the highest 5IMP scores. Furthermore, the closer the first intron was to the start codon,

the lower the 5IMP score (**Fig 1D**). We explored this relationship further by training classifiers

that increasingly excluded from the training set 5UI$^-$ transcripts according to the distance of

the first intron from the 5' end of the coding region. This revealed that classifier performance,

as measured by the area under the precision recall curve (AUPRC), increased as a function of

the distance from start codon to first intron distance (Methods, **Fig 1E**). Thus, the RNA

sequence features we identified as being predictive of 5UI$^-$ transcripts are more accurately

described as being predictors of transcripts without 5'-proximal introns.

To minimize the impact of transcripts that may 'behave' as though they were 5UI$^+$ due

to an intron early in the coding region, we eliminated 5UI$^-$ SSCR transcripts with a first intron

<90 nts downstream of the start codon (Methods) and generated a new classifier.

Discriminative motif features were learned independently (Methods), and performance of this

new classifier was gauged using 10-fold cross validation. We assessed cross-validation

performance in two ways: 1) in terms of the area under the receiver operating curve (AUC)—

which can be thought of as a measure of average recall across a range of false positive rates; 2)

in terms of area under the precision vs recall curve (AUPRC), which can be thought of as the

average precision (fraction of predictions which are correct) across a range of recall values.

Specifically, the classifier showed an AUC of 74% and AUPRC of 88% (**Fig 2A, yellow curves**;

exceeding AUC 50% and AUPRC 71%, the performance value expected of a naïve predictor).

We used this optimized classifier for all subsequent analyses.

SSCR transcripts exhibited markedly different 5IMP score distributions for the 5UI$^+$ and

5UI$^-$ subsets (**Fig 2B**). The 5UI$^+$ score distribution was unimodal with a peak at ~2.4. In

contrast, the 5UI$^-$ score distribution was bimodal with one peak at ~3.6 and another at ~9,

suggesting the existence of at least two underlying 5UI$^-$ transcript classes. The peak at score

3.6 resembled the 5UI$^+$ peak. Also contributing to the peak at 3.6 is the set of 5UI$^-$ transcripts

harboring an intron in the first 90 nts of the CDS (55% of all 5UI$^-$ transcripts). The other

distinct high-scoring 5UI$^-$ class (peak at score 9) is composed of transcripts that have specific

5UI$^-$-predictive RNA sequence elements within the early coding region.

We next wished to evaluate whether our classifier was discriminating 5UI$^+$ and 5UI$^-$

SSCR transcripts using signals that appear specifically in the early coding region as opposed to

signals that appear broadly across the coding region. To do so, for every transcript we

randomly chose 99 nts from the region downstream of the 3rd exon. The 5IMP score

distributions of these '3' proximal exon' sets were identical for 5UI+ and 5UI- transcripts (**Figs**

2A, and 2C), confirming that the sequence features that distinguish 5UI+ and 5UI- transcripts

are specific to the early coding region.

RNA elements associated with 5UI- transcripts are pervasive in the human genome

Having trained the classifier on SSCR transcripts, we wondered how well it would predict the

5UI status of other transcripts. Despite having been trained exclusively on SSCR transcripts,

the classifier performed remarkably well on MSCR transcripts, achieving an AUC of 86% and

AUPRC of 95% (**Fig 2A, purple line; Fig 2D**; as compared with 50% and 77%, respectively,

expected by chance). This result suggests that RNA elements within early coding regions of

5UI- MSCR-transcripts are similar to those in 5UI- SSCR-transcripts despite distinct functional

constraints at the protein level.

We next wondered whether the class of 5UI- transcripts that can be predicted on the

basis of early coding region features is restricted to transcripts encoding proteins trafficked to

the ER or mitochondria, or is instead a more general class of transcripts. We therefore asked

whether the classifier could predict 5UI- status in transcripts that contain neither an SSCR nor

an MSCR ("S-/M-" transcripts). Because unannotated SSCRs could confound this analysis, we

first used SignalP 3.0 to identify S-/M- transcripts most likely to contain an unannotated SSCR

(Bendtsen et al. 2004). These 'SignalP+' transcripts had a 5IMP score distribution comparable

to those of known SSCR and MSCR transcripts (**Fig 2E**), and the classifier worked well to

identify the 5UI- subset of these transcripts (AUC 82% and AUPRC 95%, **Fig 2A, light blue**

line). While 5UI+ SignalP+ transcripts had predominantly low 5IMP scores, 5UI- SignalP+ 5IMP

scores were strongly skewed towards high 5IMP scores (peak at ~9; **Fig 2E**). These results

were consistent with the idea that SignalP⁺ transcripts do contain many unannotated SSCRs.

Having considered SignalP⁺ transcripts as well as SSCR- and MSCR-containing

transcripts, we used the classifier to calculate 5IMP scores for all remaining "S⁻/M⁻/SignalP⁻"

transcripts. Although the performance was weaker on this gene set, it was still better than

expected of a naive predictor (**Fig 2A, green line**). 5UI⁺ S⁻/M⁻/SignalP⁻ transcripts were

strongly skewed toward low 5IMP scores (**Fig 2F**). Surprisingly, however, a significant

fraction of 5UI⁻ S⁻/M⁻/SignalP⁻ transcripts had high 5IMP scores (~18%). Thus, our results

suggest a broad class of transcripts with early coding regions carrying sequence signals that

predict the absence of a 5'proximal intron, or in other words, a class of transcripts with

<u>5</u>'proximal-<u>i</u>ntron-<u>m</u>inus-like coding regions. Hereafter we refer to transcripts in this class as

"5IM" transcripts.

We sought to identify what fraction of transcripts have 5IMP scores that exceed what

would be expected in the absence of 5UI⁻-predictive early coding region signals. To establish

this expectation, we used the above-described negative control set of equal-length coding

sequences from outside of the early coding region. By quantifying the excess of high-scoring

sequences in the real distribution relative to this control distribution, we estimate that 21% of

all human transcripts are 5IM transcripts (a 5IMP score of 7.41 corresponds to a 5% False

Discovery Rate; **Fig 2G**). The set of 5IM transcripts defined by our classifier (**Table S2**)

includes many that do not encode ER-targeted or mitochondrial proteins. The distribution of

various classes of mRNAs among the 5IM transcripts was: 38% ER-targeted (SSCR or SignalP⁺),

9% mitochondrial (MSCR) and 53% other classes (S⁻/M⁻/SignalP⁻) (Figure 2G). These results

suggest that RNA-level features prevalent in the early coding regions of 5UI⁻ SSCR and MSCR

transcripts are also found in other transcript types (**Figs 2F-G**), and that 5IM transcripts

represent a broad class.

Functional characterization of 5IM transcripts

5IM transcripts are defined by mRNA sequence features. Hence, we hypothesized that 5IM

transcripts may be functionally related through shared regulatory mechanisms mediated by

the presence of these common features. To this end, we collected large-scale datasets

representing diverse attributes covering six broad categories (see Table S3 for a complete list):

(1) Curated functional annotations -- e.g., Gene Ontology terms, annotation as a 'housekeeping'

gene, genes subject to RNA editing; (2) RNA localization -- e.g., to dendrites, to mitochondria;

(3) Protein and mRNA half-life, ribosome occupancy and features that decrease stability of one

or more mRNA isoforms -- e.g., AU-rich elements (4) Sequence features associated with

regulated translation -- e.g., codon optimality, secondary structure near the start codon; (5)

Known interactions with RNA-binding proteins or complexes such as Staufen-1, TDP-43, or the

Exon Junction Complex (EJC) (6) RNA modifications – i.e., N^1-methyladenosine (m^1A).

We adjusted for multiple hypotheses testing at two levels. First, we took a conservative

approach (Bonferroni correction) to correct for the number of tested functional

characteristics. Second, some of the functional categories were analyzed in more depth and

multiple sub-hypotheses were tested within the given category. In this in-depth analysis a false

discovery-based correction was adopted. Below, all reported p-values remain significant (p-

adjusted < 0.05) after multiple hypothesis test correction.

No associations between 5IM transcripts and features in categories (1), (2) and (3) were

found, other than the already-known enrichments for ER- and mitochondrial-targeted mRNAs.

However, analyses for the remaining categories yielded the significant results described below.

5IM transcripts have features suggesting lower translation efficiency

Translation regulation is a major determinant of protein levels (Vogel and Marcotte 2012). To

investigate potential connections between 5IM transcripts and translational regulation, we

examined features associated with translation. Features found to be significant were:

(I) Secondary structures near the start codon can affect initiation rate by modulating

start codon recognition (Parsyan et al. 2011). We observed a positive correlation between

5IMP score and the free energy of folding (-ΔG) of the 35 nucleotides immediately preceding

the start codon (**Fig 3A**; Spearman rho=0.39; $p < 2.2e-16$). This suggests that 5IM transcripts

have a greater tendency for secondary structure near the start codon, presumably making the

start codon less accessible.

(II) Similarly, secondary structures near the 5'cap can modulate translation by

hindering binding by the 43S-preinitiation complex to the mRNA (Babendure et al. 2006). We

observed a positive correlation between 5IMP score and the free energy of folding (-ΔG) of the

5'most 35 nucleotides (**Fig 3B**; Spearman rho=0.18; $p = 7.9e-130$). This suggests that 5IM

transcripts have a greater tendency for secondary structure near the 5'cap, presumably

hindering binding by the 43S-preinitiation complex.

(III) eIF4E overexpression. The heterotrimeric translation initiation complex eIF4F

(made up of eIF4A, eIF4E and eIF4G) is responsible for facilitating the translation of

transcripts with strong 5'UTR secondary structures (Parsyan et al. 2011). The eIF4E subunit

binds to the 7mGpppG 'methyl-G' cap and the ATP-dependent helicase eIF4A (scaffolded by

eIF4G) destabilizes 5'UTR secondary structure (Marintchev et al. 2009). A previous study

identified transcripts that were more actively translated under conditions that promote cap-

dependent translation (overexpression of eIF4E) (Larsson et al. 2007). In agreement with the

observation that 5IM transcripts have more secondary structure upstream of the start codon

and near the 5'cap, transcripts with high 5IMP scores were more likely to be translationally,

but not transcriptionally, upregulated upon eIF4E overexpression (**Fig 3C**; Wilcoxon Rank Sum

Test p = 2.05e-22, and p = 0.28, respectively).

(IV) <u>Non-AUG start codons</u>. Transcripts with non-AUG start codons also have

intrinsically low translation initiation efficiencies (Hinnebusch and Lorsch 2012). These

mRNAs were greatly enriched among transcripts with high 5IMP scores (Fisher's Exact Test p

= 0.0003; odds ratio = 3.9) and have a median 5IMP score that is 3.57 higher than those with

an AUG start (**Fig 3D**).

(V) <u>Codon optimality</u>. The efficiency of translation elongation is affected by codon

optimality (Hershberg and Petrov 2008). Although some aspects of this remain controversial

(Charneski and Hurst 2013; Shah et al. 2013; Zinshteyn and Gilbert 2013; Gerashchenko and

Gladyshev 2015), it is clear that decoding of codons by tRNAs with different abundances can

affect the translation rate under conditions of cellular stress (reviewed in Gingold and Pilpel

2011). We therefore examined the tRNA adaptation index (tAI), which correlates with copy

numbers of tRNA genes matching a given codon (dos Reis et al. 2004). Specifically, we

calculated the median tAI of the first 99 coding nucleotides of each transcript, and found that

5IMP score was negatively correlated with tAI (**Fig S1A**; Spearman Correlation rho= -0.23; p <

2e-16 median tAI and 5IMP score). This effect was restricted to the early coding regions as the

negative control set of randomly chosen sequences downstream of the 3rd exon from each

transcript did not exhibit a relationship between 5IMP score and codon optimality (**Fig S1B-C**).

Thus, 5IM transcripts show reduced codon optimality in early coding regions, suggesting that

5IM transcripts have decreased translation elongation efficiency.

To more precisely determine where the codon optimality phenomenon occurs within

the entire early coding region, we grouped transcripts by 5IMP score. For each group, we

calculated the mean tAI at codons 2-33 (i.e., nts 4-99). Across this entire region, 5IM

transcripts (5IMP >7.41; 5% FDR) had significantly lower tAI values at every codon except

codons 24 and 32 (**Fig 3E**; Wilcoxon Rank Sum test Holm-adjusted $p < 0.05$ for all

comparisons). To eliminate potential confounding variables, including nucleotide composition,

we performed several additional control analyses (Methods); none of these altered the basic

conclusion that 5IM transcripts have lower codon optimality than non-5IM transcripts across

the entire early coding region.

(VI) Ribosomes per mRNA. Finally, we examined the relationship between 5IMP score

and translation efficiency, as measured by the steady-state number of ribosomes per mRNA

molecule. To this end, we used a large dataset of ribosome profiling and RNA-Seq experiments

from human lymphoblastoid cell lines (Cenik et al. 2015). From this, we calculated the average

number of ribosomes on each transcript and identified transcripts with high or low ribosome

occupancy (respectively defined by occupancy at least one standard deviation above or below

the mean; see Methods). 5IM transcripts were slightly but significantly depleted in the high

ribosome-occupancy category (**Fig 3F**; Fisher's Exact Test $p = 0.0006$, odds ratio = 1.3).

Moreover, 5IMP scores exhibited a weak but significant negative correlation with the number

of ribosomes per mRNA molecule (Spearman rho=-0.11; p = 5.98e-23).

Taken together, all of the above results reveal that 5IM transcripts have sequence

features associated with lower translation efficiency, at the stages of both translation initiation

and elongation.

Non-ER trafficked 5IM transcripts are enriched in ER-proximal ribosome occupancy

We next investigated the relationship between 5IMP score and the localization of translation

within cells. Exploring the subcellular localization of translation at a transcriptome-scale

remains a significant challenge. Yet, a recent study described proximity-specific ribosome

profiling to identify mRNAs occupied by ER-proximal ribosomes in both yeast and human cells

(Jan et al. 2014). In this method, ribosomes are biotinylated based on their proximity to a

marker protein such as Sec61, which localizes to the ER membrane (Jan et al. 2014). For each

transcript, the enrichment for biotinylated ribosome occupancy yields a measure of ER-

proximity of translated mRNAs.

We reanalyzed this dataset to explore the relationship between 5IMP scores and ER-

proximal ribosome occupancy in HEK-293 cells. As expected, transcripts that exhibit the

highest enrichment for ER-proximal ribosomes were SSCR-containing transcripts and

transcripts with other ER-targeting signals. Yet, we noticed a surprising positive correlation

between ER-proximal ribosome occupancy and 5IM transcripts with no ER-targeting evidence

(**Fig 4**). This relationship was true for both mitochondrial genes (**Fig 4;** Spearman rho=0.43; p

< 2.2e-16), and genes with no evidence for either ER- or mitochondrial-targeting (**Fig 4**;

Spearman rho=-0.36; p < 2.2e-16). These results suggest that 5IM transcripts are more likely

than non-5IM transcripts to engage with ER-proximal ribosomes.

5IM transcripts are strongly enriched in non-canonical EJC occupancy sites

Shared sequence features and functional traits among 5IM transcripts causes one to wonder

what common mechanisms might link 5IM sequence features to 5IM traits. For example, 5IM

transcripts might share regulation by one or more RNA-binding proteins (RBPs). To

investigate this idea further, we tested for enrichment of 5IM transcripts among the

experimentally identified targets of 23 different RBPs (including CLIP-Seq, and variants; see

Methods). Only one dataset was substantially enriched for high 5IMP scores among targets: a

transcriptome-wide map of binding sites of the Exon Junction Complex (EJC) in human cells,

obtained via tandem-immunoprecipitation followed by deep sequencing (RIPiT) (Singh et al.

2014, 2012). The EJC is a multi-protein complex that is stably deposited upstream of exon-

exon junctions as a consequence of pre-mRNA splicing (Le Hir et al. 2000). RIPiT data

confirmed that canonical EJC sites (cEJC sites; sites bound by EJC core factors and appearing

~24 nts upstream of exon-exon junctions) occupy ~80% of all possible exon-exon junction

sites and are not associated with any sequence motif. Unexpectedly, many EJC-associated

footprints outside of the canonical -24 regions were observed (**Fig 5A**) (Singh et al. 2012).

These 'non-canonical' EJC occupancy sites (ncEJC sites) were associated with multiple

sequence motifs, three of which were similar to known recognition motifs for SR proteins that

co-purified with the EJC core subunits (Singh et al. 2012). Interestingly, another motif (**Fig 5B**;

top) that was specifically found in first exons is not known to be bound by any known RNA-

binding protein (Singh et al. 2012). This motif was CG-rich, a sequence feature that also

defines 5IM transcripts. This similarity presages the possibility of enrichment of first exon

ncEJC sites among 5IM transcripts.

Position analysis of called EJC peaks revealed that while only 9% of cEJCs reside in first

exons, 19% of all ncEJCs are found there. When we investigated the relationship between 5IMP

scores and ncEJCs in early coding regions, we found a striking correspondence—the median

5IMP score was highest for transcripts with the greatest number of ncEJCs (**Fig 5C**; Wilcoxon

Rank Sum Test; $p < 0.0001$). When we repeated this analysis by conditioning on 5UI status, we

similarly found that ncEJCs were enriched among transcripts with high 5IMP scores regardless

of 5UI status (Fisher's Exact Test, $p < 3.16e-14$, odds ratio > 2.3; **Fig 5D**). These results suggest

that the striking enrichment of ncEJC peaks in early coding regions was generally applicable to

all transcripts with high 5IMP scores regardless of 5UI presence.

Transcripts harboring N^1-methyladenosine (m^1A) have high 5IMP scores

It is increasingly clear that ribonucleotide base modifications in mRNAs are highly prevalent

and can be a mechanism for post-transcriptional regulation (Frye et al. 2016). One RNA

modification present towards the 5' ends of mRNA transcripts is N^1-methyladenosine (m^1A)

(Li et al. 2016; Dominissini et al. 2016), initially identified in total RNA and rRNAs (Dunn 1961;

Hall 1963; Klootwijk and Planta 1973). Intriguingly, the position of m^1A modifications has

been shown to be more correlated with the position of the first intron than with

transcriptional or translational start sites (Figure 2g from Dominissini et al. 2016). When the

distance of m^1As to each splice site in a given mRNA was calculated, the first splice site was

found to be the nearest for 85% of m^1As (Dominissini et al. 2016). When 5'UTR introns were

present, m^1A was found to be near the first splice site regardless of the position of the start

codon (Dominissini et al. 2016). Given that 5IM transcripts are also characterized by the

position of the first intron, we investigated the relationship between 5IMP score and m^{1}A RNA

modification marks.

We analyzed the union of previously identified m^{1}A modifications (Dominissini et al.

2016) across all cell types and conditions. Although there is some evidence that these marks

depend on cell type and growth condition, it is difficult to be confident of the cell type and

condition-dependence of any particular mark given experimental variation (see Methods).

Nevertheless, we found that mRNAs with m^{1}A modification early in the coding region (first 99

nucleotides) had substantially higher 5IMP scores than mRNAs lacking these marks (**Fig 6A**;

Wilcoxon Rank Sum Test p = 3.4e-265), and were greatly enriched among 5IM transcripts (**Fig**

6A; Fisher's Exact Test p = 1.6e-177; odds ratio = 3.8). In other words, the sequence features

within the early coding region that define 5IMP transcripts also associate with m^{1}A

modification in the early coding region.

We next wondered whether 5IMP score was related to m^{1}A modification generally, or

only associated with m^{1}A modification in the early coding region. Indeed, many of the

previously identified m^{1}A peaks were within the 5'UTRs of mRNAs (Li et al. 2016).

Interestingly, 5IMP scores were only associated with m^{1}A modification in the early coding

region, and not with m^{1}A modification in the 5'UTR (**Fig 6B**). This offers the intriguing

possibility that the sequence features that define 5IMP transcripts are co-localized with m^{1}A

modification.

Discussion:

Can Cenik

Coordinating the expression of functionally related transcripts can be achieved by post-

transcriptional processes such as splicing, RNA export, RNA localization or translation (Moore

and Proudfoot 2009). Sets of mRNAs subject to a common regulatory transcriptional process

can exhibit common sequence features that define them to be a class. For example, transcripts

subject to regulation by particular miRNAs tend to share certain sequences in their 3'UTRs that

are complementary to the regulatory miRNA (Ameres and Zamore 2013). Similarly, transcripts

that share a 5' terminal oligopyrimidine tract are coordinately regulated by mTOR and

ribosomal protein S6 kinase (Meyuhas 2000). Here we quantitatively define '5IM' transcripts

as a class that shares common sequence elements and functional properties. We estimate the

5IM class to comprise 21% of all human transcripts.

Whereas 35% of human transcripts have one or more 5'UTR introns, the majority of

5IM transcripts have neither a 5'UTR intron nor an intron in the first 90 nts of the ORF. Other

shared features of 5IM transcripts include sequence features associated with low translation

initiation rates. These are: (1) a tendency for RNA secondary structure in the region

immediately preceding the start codon (**Fig 3A**), and near the 5'cap (**Fig 3B**); (2) translational

upregulation upon overexpression of eIF4E (**Fig 3C**); and (3) more frequent use of non-AUG

start codons (**Fig 3D**). Also consistent with low intrinsic translation efficiencies, 5IM

transcripts additionally tend to depend on less abundant tRNAs to decode the beginning of the

open reading frame (**Fig 3E**).

We had previously reported that transcripts encoding proteins with ER- and

mitochondrial-targeting signal sequences (SSCRs and MSCRs, respectively) are over-

represented among the 65% of transcripts lacking 5'UTR introns (Cenik et al. 2011).

Transcripts in this set are enriched for the sequence features detected by our 5IM classifier. By

examining these enriched sequence features, we showed that the 5IM class extends beyond

mRNAs encoding membrane proteins.   Jan et al. recently developed a transcriptome-scale

method to identify mRNAs occupied by ER-proximal ribosomes in both yeast and human cells

(Jan et al. 2014). As expected, transcripts known to encode ER-trafficked proteins were highly

enriched for ER-proximal ribosome occupancy. However, their data also showed many

transcripts encoding non-ER trafficked proteins to also be engaged with ER-proximal

ribosomes (Reid and Nicchitta 2015b). Similarly, several other studies have suggested a critical

role of ER-proximal ribosomes in translating several cytoplasmic proteins (Reid and Nicchitta

2015a). Here, we found that 5IM transcripts including those that are not ER-trafficked or

mitochondrial were significantly more likely to exhibit binding to ER-proximal ribosomes

(**Figs 4A-B**).

In addition to ribosomes directly resident on the ER, an interesting possibility is the

presence of a pool of peri-ER ribosomes (Jan et al. 2015; Reid and Nicchitta 2015a).

Association of 5IM transcripts with such a peri-ER ribosome pool could potentially explain the

observed correlation of 5IM status with binding to ER-proximal ribosomes. The ER is

physically proximal to mitochondria (Rowland and Voeltz 2012), so peri-ER ribosomes may

include those translating mRNAs on mitochondria (i.e., mRNAs with MSCRs) (Sylvestre et al.

2003). However, even when transcripts corresponding to ER-trafficked and mitochondrial

proteins were excluded from consideration, ER-proximal ribosome enrichment and 5IMP

scores were highly correlated (**Fig 4A**). Thus another shared feature of 5IM transcripts is their

translation on or near the ER regardless of the ultimate destination of the encoded protein.

In an attempt to identify a common factor binding 5IM transcripts, we asked whether

5IM transcripts were enriched among the experimentally identified targets of 23 different

RBPs. Only one RBP emerged--the exon junction complex (EJC). Specifically, we observed a

dramatic enrichment of non-canonical EJC (ncEJC) binding sites within the early coding region

of 5IM transcripts. Further, the CG-rich motif identified for ncEJCs in first exons is strikingly

similar to the CG-rich motif enriched in the first exons of 5IM transcripts (**Fig 5B**). Previous

work implicated RanBP2, a protein associated with the cytoplasmic face of the nuclear pore, as

a binding factor for some SSCRs (Mahadevan et al. 2013). This finding suggests that nuclear

pore proteins may influence EJC occupancy on these transcripts.

EJC deposition during the process of pre-mRNA splicing enables the nuclear history of

an mRNA to influence post-transcriptional processes including mRNA localization, translation

efficiency, and nonsense mediated decay (Chang et al. 2007; Kervestin and Jacobson 2012;

Choe et al. 2014). While canonical EJC binding occurs at a fixed distance upstream of exon-exon

junctions and involves direct contact between the sugar-phosphate backbone and the EJC core

anchoring protein eIF4AIII, ncEJC binding sites likely reflect stable engagement between the

EJC core and other mRNP proteins (e.g., SR proteins) recognizing nearby sequence motifs.

Although some RBPs were identified for ncEJC motifs found in internal exons (Singh et al.

2014, 2012), to date no candidate RBP has been identified for the CG-rich ncEJC motif found in

the first exon. If this motif does result from an RBP interaction, it is likely to be one or more of

the ~70 proteins that stably and specifically bind to the EJC core (Singh et al. 2012).

Finally, we observed a dramatic enrichment for m^1A modifications among 5IM

transcripts, with specific enrichment for m^1A modifications in the early coding region. Given

this striking enrichment perhaps it is perhaps not surprising that m^1A containing mRNAs were

also shown to have more structured 5'UTRs that are GC-rich compared to m^1A lacking mRNAs

(Dominissini et al. 2016). Similar to 5IM transcripts, m^1A containing mRNAs were found to

decorate start codons that appear in a highly structured context. While ALKBH3 has been

identified as a protein that can demethylate m^{1}A, it is currently unknown whether there are

any proteins that can specifically act as "readers" of m^{1}A. Recent studies have begun to identify

such readers for other mRNA modifications such as YTHDF1, YTHDF2, WTAP and HNRNPA2B1

(Ping et al. 2014; Liu et al. 2014; Wang et al. 2014, 2015; Alarcón et al. 2015). Our study

highlights a possible link between non-canonical EJC binding and m^{1}A. Hence, our results yield

the intriguing hypothesis that one or more of the ~70 proteins that stably and specifically bind

to the EJC core can function as an m^{1}A reader. Future work involving directed experiments

would be needed to test this hypothesis.

Given that 5IM transcripts are enriched for ER-targeted and mitochondrial proteins, it

is plausible that the observed functional characteristics of 5IM transcripts are driven solely by

SSCR and MSCR-containing transcripts. Hence, we repeated all analyses for the subclasses of

5IM transcripts (MSCR-containing, SSCR-containing, S$^-$/M$^-$/SignalP$^+$, or S$^-$/M$^-$/SignalP$^-$). We

found the observed associations remained statistically significant and had the same direction

of effect, even after eliminating SSCR- and MSCR-containing transcripts, despite the fact that all

training of the 5IM classifier was performed only using SSCR transcripts. We also found that

5IMP score was equally or more strongly associated with each of the functional characteristics

compared to the 5UI status. In conclusion, the molecular associations we report apply to 5IM

transcripts as a whole, and are not driven solely by the subset of 5IM transcripts encoding ER-

or mitochondria-targeting signal peptides, and seem to indicate shared features beyond simple

lack of a 5'UTR intron.

An intriguing possibility is that 5IM transcript features associated with lower intrinsic

translation efficiency may together enable greater 'tunability' of 5IM transcripts at the

translation stage. Regulated enhancement or repression of translation, for 5IM transcripts,

could allow for rapid changes in protein levels. There are analogies to this scenario in

transcriptional regulation, wherein highly regulated genes often have promoters with low

baseline levels that can be rapidly modulated through the action of regulatory transcription

factors. As more ribosome profiling studies are published examining translational responses

transcriptome-wide under multiple perturbations, conditions under which 5IM transcripts are

translationally regulated may be revealed. Directed experiments will be needed to test

translational features of 5IM transcripts hypothesized via this computational analysis.

Taken together, our analyses reveal the existence of a distinct '5IM' class comprising

21% of human transcripts. This class is defined by depletion of 5' proximal introns, presence

of specific RNA sequence features associated with low translation efficiency, non-canonical

binding by the Exon Junction Complex and an enrichment for N^1-methyladenosine

modification.

Can Cenik

Materials and Methods:

Datasets and Annotations

Human transcript sequences were downloaded from the NCBI human Reference Gene

Collection (RefSeq) via the UCSC table browser (hg19) on Jun 25 2010 (Kent et al. 2002; Pruitt

et al. 2005). Transcripts with fewer than three coding exons, and transcripts where the first 99

coding nucleotides straddle more than two exons were removed from further consideration.

The criteria for exclusion of genes with fewer than three coding exons was to ensure that the

analysis of downstream regions was possible for all genes that were used in our analysis of

early coding regions. Specifically, the downstream regions were selected randomly from

downstream of the 3rd exons. Hence, genes with fewer exons would not be able to contribute a

downstream region potentially creating a skew in representation. In total there were ~3000

genes that were removed from consideration due to this filter. Therefore, our classifier is

limited in its ability to assess transcripts from these genes. However, the performance

measures reported in our manuscript are robust to exclusion of these genes, in the sense that

the same class of transcripts was used in both training and test datasets.

Transcripts were clustered based on sequence similarity in the first 99 coding

nucleotides. Specifically, each transcript pair was aligned using BLAST with the DUST filter

disabled (Altschul et al. 1990). Transcript pairs with BLAST E-values < 1e-25 were grouped

into transcript clusters. In total, there were 15576 transcript clusters that were considered

further. These clusters that were subsequently assigned to one of four categories: MSCR-

containing, SSCR-containing, S$^-$/MSCR$^-$ SignalP$^+$, or S$^-$/MSCR$^-$ SignalP$^-$ as follows:

MSCR-containing transcripts were annotated using MitoCarta and other sources as

described in (Cenik et al. 2011). SSCR-containing transcripts were the set of transcripts

annotated to contain signal peptides in the Ensembl Gene v.58 annotations, which were

downloaded through Biomart on Jun 25 2010. For transcripts without an annotated MSCR or

SSCR, the first 70 amino acids were analyzed using SignalP 3.0 (Bendtsen et al. 2004). Using

the eukaryotic prediction mode, transcripts were assigned to the S$^-$/MSCR$^-$ SignalP$^+$ category

if either the Hidden Markov Model or the Artificial Neural Network classified the sequence as

signal peptide containing. All remaining transcript clusters were assigned to the S$^-$/MSCR$^-$

SignalP$^-$ category. The number of transcript clusters in each of the four categories was: 3743

SSCR, 737 MSCR, 696 S$^-$/MSCR$^-$ SignalP$^+$, 10400 S$^-$/MSCR$^-$ SignalP$^-$.

For each transcript cluster, we also constructed matched control sequences. Control

sequences were derived from a single randomly chosen in-frame 'window' downstream of the

3rd exon from the evaluated transcripts. If an evaluated transcript had fewer than 99

nucleotides downstream of the 3rd exon, no control sequence was extracted. 5UI labels and

transcript clustering for the control sequences were inherited from the evaluated transcript.

The rationale for this decision is that our analysis depends on the position of the first intron;

hence genes with fewer than two exons need to be excluded, as these will not have introns. We

further required the matched control sequences to fall downstream of the early coding region.

In the vast majority of cases the third exon fell outside the first 99 nucleotides of the coding

region, making this a convenient criterion by which to choose control regions.

Sequence Features and Motif Discovery

36 sequence features were extracted from each transcript (**Table S1**). The sequence

features included the ratio of arginines to lysines, the ratio of leucines to isoleucines, adenine

content, length of the longest stretch without adenines, preference against codons that contain

adenines or thymines. These features were previously found to be enriched in SSCR-

containing and certain 5UI⁻ transcripts (Cenik et al. 2011; Palazzo et al. 2007). In addition, we

extracted ratios between several other amino acid pairs based on having

biochemical/evolutionary similarity, i.e. having positive scores, according to the BLOSUM62

matrix (Henikoff and Henikoff 1992). To avoid extreme ratios given the relatively short

sequence length, pseudo-counts were added to amino acid ratios using their respective

genome-wide prevalence.

In addition, we used three published motif finding algorithms (AlignACE, DEME, and

MoAN (Roth et al. 1998; Redhead and Bailey 2007; Valen et al. 2009)) to discover RNA

sequence motifs enriched among 5UI⁻ transcripts. AlignACE implements a Gibbs sampling

approach and is one of the pioneering efforts in motif discovery (Roth et al. 1998). We

modified the AlignACE source code to restrict motif searches to only the forward strand of the

input sequences to enable RNA motif discovery. DEME and MoAN adopt discriminative

approaches to motif finding by searching for motifs that are differentially enriched between

two sets of sequences (Redhead and Bailey 2007; Valen et al. 2009). MoAN has the additional

advantage of discovering variable length motifs, and can identify co-occurring motifs with the

highest discriminative power.

In total, six motifs were discovered using the three motif finding algorithms (**Table S1**).

Position specific scoring matrices for all motifs were used to score the first 99 - *l* positions in

each sequence, where *l* is the length of the motif. We assessed the significance of each motif

instance by calculating the p-value of enrichment (Fisher's Exact Test) among 5UI⁻ transcripts

considering all transcripts with a motif instance achieving a PSSM score greater than equal to

the instance under consideration. The significance score and position of the two best motif

instances were used as features for the classifier (**Table S1**).

5IM Classifier Training and Performance Evaluation

We modified an implementation of the Random Forest classifier (Breiman 2001) to

model the relationship between sequence features in the early coding region and the absence

of 5'UTR introns (5UIs). This classifier discriminates transcripts with 5'proximal-intron-

minus-like-coding regions and hence is named the '5IM' classifier. The training set for the

classifier was composed of SSCR transcripts exclusively. There were two reasons to restrict

model construction to SSCR transcripts: 1) we expected the presence of specific RNA elements

as a function of 5UI presence based on our previous work (Cenik et al. 2011); and 2) we

wanted to restrict model building to sequences that have similar functional constraints at the

protein level.

We observed that 5UI$^-$ transcripts with introns proximal to 5' end of the coding region

have sequence characteristics similar to 5UI$^+$ transcripts (**Fig 1D**). To systematically

characterize this relationship, we built different classifiers using training sets that excluded

5UI$^-$ transcripts with a coding region intron positioned at increasing distances from the start

codon. We evaluated the performance of each classifier using 10-fold cross validation.

Given that a large number of motif discovery iterations were needed, we sought to

reduce the computational burden. We isolated a subset of the training examples to be used

exclusively for motif finding. Motif discovery was performed once using this set of sequences,

and the same motifs were used in each fold of the cross validation for all the classifiers.

Imbalances between the sizes of positive and negative training examples can lead to

detrimental classification performance (Wang and Yao 2012). Hence, we balanced the training

set size of 5UI$^-$ and 5UI$^+$ transcripts by randomly sampling from the larger class. We

constructed 10 sub-classifiers to reduce sampling bias, and for each test example, the

prediction score from each subclassifier was summed to produce a combined score between 0

and 10. For the rest of the analyses, we used the classifier trained using 5UI⁻ transcripts where

first coding intron falls outside the first 90 coding nucleotides (**Fig 1E**).

We evaluated classifier performance using a 10-fold cross validation strategy for SSCR-

containing transcripts (i.e. the training set). In each fold of the cross-validation, the model was

trained without any information from the held-out examples, including motif discovery. For all

the other transcripts and the control sets (see above), the 5IMP scores were calculated using

the classifier trained using SSCR transcripts as described above. 5IMP score distribution for

the control set was used to calculate the empirical cumulative null distribution. Using this

function, we determined the p-value corresponding to the 5IMP score for all transcripts. We

corrected for multiple hypotheses testing using the qvalue R package (Storey 2003). Based on

this analysis, we estimate that a 5IMP score of 7.41 corresponds to a 5% False Discovery Rate

and suggest that 21% of all human transcripts can be considered as 5IM transcripts.

While the theoretical range of 5IMP scores is 0-10, the highest observed 5IMP is 9.855.

We note that for all figures that depict 5IMP score distributions, we displayed the entire

theoretical range of 5IMP scores (0-10).

Functional Characterization of 5IM Transcripts:

We collected genome-scale dataset from publically available databases and from

supplementary information provided in selected articles. For all analyzed datasets, we first

converted all gene/transcript identifiers (IDs) into RefSeq transcript IDs using the Synergizer

webserver (Berriz and Roth 2008). If a dataset was generated using a non-human species (ex.

Targets identified by TDP-43 RNA immunoprecipitation in rat neuronal cells), we used

Homologene release 64 (downloaded on Sep 28 2009) to identify the corresponding ortholog

in humans. If at least one member of a transcript cluster was associated with a functional

phenotype, we assigned the cluster to the positive set with respect to the functional phenotype.

If more than one member of a cluster had the functional phenotype, we only retained one copy

of the cluster unless they differed in a quantitative measurement. For example, consider two

hypothetical transcripts: NM_1 and NM_2 that were clustered together and have a 5IMP score

of 8.5. If NM_1 had an mRNA half-life of 2hr while NM_2's half-life was 1hr than we split the

cluster while preserving the 5IMP score for both NM_1 and NM_2.

Once the transcripts were partitioned based on the functional phenotype, we ran two

statistical tests: 1) Fisher's Exact Test for enrichment of 5IM transcripts within the functional

category; 2) Wilcoxon Rank Sum Test to compare 5IMP scores between transcripts partitioned

by the functional phenotype. Additionally, for datasets where a quantitative measurement was

available (ex. mRNA half-life), we calculated the Spearman rank correlation between 5IMP

scores and the quantitative variable. In these analyses, we assumed that the test space was the

entire set of RefSeq transcripts. For all phenotypes where we observed a preliminary

statistically significant result, we followed up with more detailed analyses described below.

Analysis of Features Associated with Translation:

For each transcript, we predicted the propensity for secondary structure preceding the

translation start site and the 5'cap. Specifically, we extracted 35 nucleotides preceding the

translation start site or the first 35 nucleotides of the 5'UTR. If a 5'UTR is shorter than 35

nucleotides, the transcript was removed from the analysis. hybrid-ss-min utility (UNAFold

package version 3.8) with default parameters was used to calculate the minimum folding

energy (Markham and Zuker 2008).

Can Cenik

Codon optimality was measured using the tRNA Adaptation Index (tAI), which is based

on the genomic copy number of each tRNA (dos Reis et al. 2004). tAI for all human codons

were downloaded from Tuller et al. 2010 Table S1. tAI profiles for the first 30 amino acids

were calculated for all transcripts. Codon optimality profiles were summarized for the first 30

amino acids for each transcript or by averaging tAI at each codon.

We carried out two control experiments to test whether the association between 5IMP

score and tAI could be explained by confounding variables. First, we permuted the nucleotides

in the first 90 nts and observed no relationship between 5IMP score and mean tAI when these

permuted sequences were used (**Fig S1**). Second, we selected random in-frame 99 nucleotides

from 3rd exon to the end of the coding region and found no significant differences in tAI (**Fig**

S1). These results suggest that the relationship between tAI and 5IMP score is confined to the

first 30 amino acids and is not explained by simple differences in nucleotide composition.

Ribosome profiling and RNA expression data for human lymphoblastoid cells (LCLs)

were downloaded from NCBI GEO database accession number: GSE65912. Translation

efficiency was calculated as previously described (Cenik et al. 2015). Median translation

efficiency across the different cell-types was used for each transcript.

Analysis of Proximity Specific Ribosome Profiling Data:

We downloaded proximity specific ribosome profiling data for HEK 293 cells from Jan

et al. 2014; Table S6. We converted UCSC gene identifiers to HGNC symbols using g:Profiler

(Reimand et al. 2011). We retained all genes with an RPKM >5 in either input or pulldown and

required that at least 30 reads were mapped in either of the two libraries. We used ER-

targeting evidence categories "secretome", "phobius", "TMHMM", "SignalP", "signalSequence",

"signalAnchor" from (Jan et al. 2014) to annotate genes as having ER-targeting evidence. The

genes that did not have any ER-targeting evidence or "mitoCarta" /"mito.GO" annotations were

deemed as the set of genes with no ER-targeting or mitochondrial evidence. We calculated the

$\log_2$ of the ratio between ER-proximal ribosome pulldown RPKM and control RPKM as the

measure of enrichment for ER-proximal ribosome occupancy (as in Jan et al. 2014). A moving

average of this ratio was calculated for genes grouped by their 5IMP score. For this calculation,

we used bins of 30 mitochondrial genes or 100 genes with no evidence of ER- or mitochondrial

targeting.

Analysis of Genome-wide Binding Sites of Exon Junction Complex:

Dr. Gene Yeo and Gabriel Pratt generously shared uniformly processed peak calls for

experiments identifying human RNA binding protein targets. These datasets include various

CLIP-Seq datasets and its variants such as iCLIP. A total 49 datasets from 22 factors were

analyzed. These factors were: hnRNPA1, hnRNPF, hnRNPM, hnRNPU, Ago2, hnRNPU, HuR,

IGF2BP1, IGF2BP2, IGF2BP3, FMR1, eIF4AIII, PTB, IGF2BP1, Ago3, Ago4, MOV10, Fip1, CF

Im68, CF Im59, CF Im25, and hnRNPA2B1. We extracted the 5IMP scores for all targets of each

RBP. We calculated the Wilcoxon Rank Sum test statistic comparing the 5IMP score

distribution of the targets of each RBP to all other transcripts with 5IMP scores. None of the

tested RBP target sets had an adjusted p-value < 0.05 and a median difference in 5IMP score >

1 when compared to non-target transcripts.

In addition, we used RNA:protein immunoprecipitation in tandem (RIPiT) data to

determine Exon Junction Complex (EJC) binding sites (Singh et al. 2014, 2012). We analyzed

the common peaks from the Y14-Magoh immunoprecipitation configuration (Singh et al. 2012;

Kucukural et al. 2013). Canonical EJC binding sites were defined as peaks whose weighted

center were 15 to 32 nucleotides upstream of an exon-intron boundary. All remaining peaks

were deemed as "non-canonical" EJC binding sites. We extracted all non-canonical peaks that

overlapped the first 99 nucleotides of the coding region and restricted our analysis to

transcripts that had an RPKM greater than one in the matched RNA-Seq data.

Analysis of m^1A *modified transcripts*

We downloaded the list of RNAs observed to contain m^1A from Li et al. (2016) and

Dominissini et al. (2016). RefSeq transcript identifiers were converted to HGNC symbols using

g:Profiler (Reimand et al. 2011). The overlap between the two datasets was determined using

HGNC symbols. m^1A modifications that overlap the first 99 nucleotides of the coding region

were determined using bedtools (Quinlan and Hall 2010).

Li et al. (2016) identified 600 transcripts with m^1A modification in normal HEK293

cells. Of these, 368 transcripts were not found to contain the m^1A modification in HEK293 cells

by Dominissini et al. (2016). Yet, 81% of these were found to be m^1A modified in other cell

types. Li et al. (2016) also analyzed m^1A upon H_2O_2 treatment and serum starvation in HEK293

cells and identified many m^1A modifications that are only found these stress conditions.

However, 20% of 371 transcripts harboring stress-induced m^1A modifications were found in

normal HEK293 cells by Dominissini et al. (2016). Taken together, these analyses suggest that

transcriptome-wide m^1A maps remain incomplete. Hence, we analyzed the 5IMP scores of all

mRNAs with m^1A across cell types and conditions. We reported results using the Dominissini

et al. dataset but the same conclusions were supported by m^1A modifications from Li et al.

Can Cenik

Acknowledgements:

We would like to thank Gene Yeo, and Gabriel Pratt for sharing peak calls for RNA-

binding protein target identification experiments (CLIP, PAR-CLIP, iCLIP). We thank Elif

Sarinay Cenik, Başar Cenik, and Alper Küçükural for helpful discussions. F.P.R. and H.N.C. were

supported by the Canada Excellence Research Chairs Program. This work was supported in

part by NIH grants GM50037 (M.J.M.), HG001715 and HG004233 (F.P.R.), the Krembil

Foundation, an Ontario Research Fund award, and the Avon Foundation (F.P.R.). A.F.P. was

supported by a grant from the Canadian Institutes of Health Research to A.F.P. (FRN 102725).

Author contributions: CC, MJM, and FPR designed the study. CC and HNC implemented the

random forest classifier and carried out all analyses. GS and AA performed initial experiments.

MPS and AFP provided critical feedback. MJM and FPR jointly supervised the project. CC, FPR

and MJM wrote the manuscript with input from all authors.

Can Cenik

1 References:

Alarcón CR, Goodarzi H, Lee H, Liu X, Tavazoie S, Tavazoie SF. 2015. HNRNPA2B1 Is a Mediator of
m(6)A-Dependent Nuclear RNA Processing Events. *Cell* **162**: 1299–1308.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol*
215: 403–410.

Ameres SL, Zamore PD. 2013. Diversifying microRNA sequence and function. *Nat Rev Mol Cell Biol* **14**:
475–488.

Babendure JR, Babendure JL, Ding J-H, Tsien RY. 2006. Control of mammalian translation by mRNA
structure near caps. *RNA* **12**: 851–861.

Bendtsen JD, Nielsen H, von Heijne G, Brunak S. 2004. Improved prediction of signal peptides: SignalP
3.0. *J Mol Biol* **340**: 783–795.

Berriz GF, Roth FP. 2008. The Synergizer service for translating gene, protein and other biological
identifiers. *Bioinformatics* **24**: 2272–2273.

Breiman L. 2001. Random Forests. *Mach Learn* **45**: 5–32.

Cenik C, Chua HN, Zhang H, Tarnawsky SP, Akef A, Derti A, Tasan M, Moore MJ, Palazzo AF, Roth FP.
2011. Genome analysis reveals interplay between 5'UTR introns and nuclear mRNA export for
secretory and mitochondrial genes. *PLoS Genet* **7**: e1001366.

Cenik C, Derti A, Mellor JC, Berriz GF, Roth FP. 2010. Genome-wide functional analysis of human 5'
untranslated region introns. *Genome Biol* **11**: R29.

Cenik C, Sarinay Cenik E, Byeon GW, Grubert F, Candille SI, Spacek D, Alsallakh B, Tilgner H, Araya CL,
Tang H, et al. 2015. Integrative analysis of RNA, translation and protein levels reveals distinct
regulatory variation across humans. *Genome Res*. http://dx.doi.org/10.1101/gr.193342.115.

Chang Y-F, Imam JS, Wilkinson MF. 2007. The nonsense-mediated decay RNA surveillance pathway.
Annu Rev Biochem **76**: 51–74.

Charneski CA, Hurst LD. 2013. Positively charged residues are the major determinants of ribosomal
velocity. *PLoS Biol* **11**: e1001508.

Choe J, Ryu I, Park OH, Park J, Cho H, Yoo JS, Chi SW, Kim MK, Song HK, Kim YK. 2014. eIF4AIII enhances
translation of nuclear cap-binding complex-bound mRNAs by promoting disruption of secondary
structures in 5'UTR. *Proc Natl Acad Sci U S A* **111**: E4577–86.

Dominissini D, Nachtergaele S, Moshitch-Moshkovitz S, Peer E, Kol N, Ben-Haim MS, Dai Q, Di Segni A,
Salmon-Divon M, Clark WC, et al. 2016. The dynamic N(1)-methyladenosine methylome in
eukaryotic messenger RNA. *Nature* **530**: 441–446.

dos Reis M, Savva R, Wernisch L. 2004. Solving the riddle of codon usage preferences: a test for
translational selection. *Nucleic Acids Res* **32**: 5036–5044.

Dunn DB. 1961. The occurrence of 1-methyladenine in ribonucleic acid. *Biochim Biophys Acta* **46**: 198–
200.

Frye M, Jaffrey SR, Pan T, Rechavi G, Suzuki T. 2016. RNA modifications: what have we learned and
where are we headed? *Nat Rev Genet* **17**: 365–372.

Gerashchenko MV, Gladyshev VN. 2015. Translation inhibitors cause abnormalities in ribosome
profiling experiments. *Nucleic Acids Res* **42**: e134.

Gingold H, Pilpel Y. 2011. Determinants of translation efficiency and accuracy. *Mol Syst Biol* **7**: 481.

Hall RH. 1963. Method for isolation of 2′-O-methylribonucleosides and N1-methyladenosine from
ribonucleic acid. *Biochimica et Biophysica Acta (BBA) - Specialized Section on Nucleic Acids and*
Related Subjects **68**: 278–283.

Henikoff S, Henikoff JG. 1992. Amino acid substitution matrices from protein blocks. *Proc Natl Acad Sci*
U S A **89**: 10915–10919.

Hershberg R, Petrov DA. 2008. Selection on codon bias. *Annu Rev Genet* **42**: 287–299.

Hinnebusch AG, Lorsch JR. 2012. The mechanism of eukaryotic translation initiation: new insights and
challenges. *Cold Spring Harb Perspect Biol* **4**. http://dx.doi.org/10.1101/cshperspect.a011544.

Hong X, Scofield DG, Lynch M. 2006. Intron size, abundance, and distribution within untranslated
regions of genes. *Mol Biol Evol* **23**: 2392–2404.

Jan CH, Williams CC, Weissman JS. 2015. LOCAL TRANSLATION. Response to Comment on "Principles of
ER cotranslational translocation revealed by proximity-specific ribosome profiling." *Science* **348**:
1217.

Jan CH, Williams CC, Weissman JS. 2014. Principles of ER cotranslational translocation revealed by
proximity-specific ribosome profiling. *Science* **346**: 1257521.

Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM, Haussler D. 2002. The human
genome browser at UCSC. *Genome Res* **12**: 996–1006.

Kervestin S, Jacobson A. 2012. NMD: a multifaceted response to premature translational termination.
Nat Rev Mol Cell Biol **13**: 700–712.

Klootwijk J, Planta RJ. 1973. Analysis of the methylation sites in yeast ribosomal RNA. *Eur J Biochem* **39**:
325–333.

Kucukural A, Özadam H, Singh G, Moore MJ, Cenik C. 2013. ASPeak: an abundance sensitive peak
detection algorithm for RIP-Seq. *Bioinformatics* **29**: 2485–2486.

Larsson O, Li S, Issaenko OA, Avdulov S, Peterson M, Smith K, Bitterman PB, Polunovsky VA. 2007.
Eukaryotic Translation Initiation Factor 4E–Induced Progression of Primary Human Mammary
Epithelial Cells along the Cancer Pathway Is Associated with Targeted Translational Deregulation
of Oncogenic Drivers and Inhibitors. *Cancer Res* **67**: 6814–6824.

Lee ES, Akef A, Mahadevan K, Palazzo AF. 2015. The consensus 5' splice site motif inhibits mRNA
nuclear export. *PLoS One* **10**: e0122743.

Le Hir H, Izaurralde E, Maquat LE, Moore MJ. 2000. The spliceosome deposits multiple proteins 20-24
nucleotides upstream of mRNA exon-exon junctions. *EMBO J* **19**: 6860–6869.

Liu J, Yue Y, Han D, Wang X, Fu Y, Zhang L, Jia G, Yu M, Lu Z, Deng X, et al. 2014. A METTL3-METTL14
complex mediates mammalian nuclear RNA N6-adenosine methylation. *Nat Chem Biol* **10**: 93–95.

Li X, Xiong X, Wang K, Wang L, Shu X, Ma S, Yi C. 2016. Transcriptome-wide mapping reveals reversible
and dynamic N1-methyladenosine methylome. *Nat Chem Biol*.
http://dx.doi.org/10.1038/nchembio.2040 (Accessed March 29, 2016).

Mahadevan K, Zhang H, Akef A, Cui XA, Gueroussov S, Cenik C, Roth FP, Palazzo AF. 2013.
RanBP2/Nup358 potentiates the translation of a subset of mRNAs encoding secretory proteins.
PLoS Biol **11**: e1001545.

Marintchev A, Edmonds KA, Marintcheva B, Hendrickson E, Oberer M, Suzuki C, Herdy B, Sonenberg N,
Wagner G. 2009. Topology and regulation of the human eIF4A/4G/4H helicase complex in
translation initiation. *Cell* **136**: 447–460.

Markham NR, Zuker M. 2008. UNAFold: software for nucleic acid folding and hybridization. *Methods*
Mol Biol **453**: 3–31.

Meyuhas O. 2000. Synthesis of the translational apparatus is regulated at the translational level. *Eur J*
Biochem **267**: 6321–6330.

Moore MJ, Proudfoot NJ. 2009. Pre-mRNA processing reaches back to transcription and ahead to
translation. *Cell* **136**: 688–700.

Palazzo AF, Mahadevan K, Tarnawsky SP. 2013. ALREX-elements and introns: two identity elements
that promote mRNA nuclear export. *Wiley Interdiscip Rev RNA* **4**: 523–533.

Palazzo AF, Springer M, Shibata Y, Lee C-S, Dias AP, Rapoport TA. 2007. The signal sequence coding
region promotes nuclear export of mRNA. *PLoS Biol* **5**: e322.

Parsyan A, Svitkin Y, Shahbazian D, Gkogkas C, Lasko P, Merrick WC, Sonenberg N. 2011. mRNA
helicases: the tacticians of translational control. *Nat Rev Mol Cell Biol* **12**: 235–245.

Ping X-L, Sun B-F, Wang L, Xiao W, Yang X, Wang W-J, Adhikari S, Shi Y, Lv Y, Chen Y-S, et al. 2014.
Mammalian WTAP is a regulatory subunit of the RNA N6-methyladenosine methyltransferase. *Cell*
Res **24**: 177–189.

Pruitt KD, Tatusova T, Maglott DR. 2005. NCBI Reference Sequence (RefSeq): a curated non-redundant
sequence database of genomes, transcripts and proteins. *Nucleic Acids Res* **33**: D501–4.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**: 841–842.

Redhead E, Bailey TL. 2007. Discriminative motif discovery in DNA and protein sequences using the
DEME algorithm. *BMC Bioinformatics* **8**: 385.

Reid DW, Nicchitta CV. 2015a. Diversity and selectivity in mRNA translation on the endoplasmic
reticulum. *Nat Rev Mol Cell Biol* **16**: 221–231.

Reid DW, Nicchitta CV. 2015b. LOCAL TRANSLATION. Comment on "Principles of ER cotranslational
translocation revealed by proximity-specific ribosome profiling." *Science* **348**: 1217.

Reimand J, Arak T, Vilo J. 2011. g: Profiler—a web server for functional interpretation of gene lists
(2011 update). *Nucleic Acids Res* gkr378.

Roth FP, Hughes JD, Estep PW, Church GM. 1998. Finding DNA regulatory motifs within unaligned
noncoding sequences clustered by whole-genome mRNA quantitation. *Nat Biotechnol* **16**: 939–

945.

Rowland AA, Voeltz GK. 2012. Endoplasmic reticulum–mitochondria contacts: function of the junction.
Nat Rev Mol Cell Biol **13**: 607–625.

Shah P, Ding Y, Niemczyk M, Kudla G, Plotkin JB. 2013. Rate-limiting steps in yeast protein translation.
Cell **153**: 1589–1601.

Singh G, Kucukural A, Cenik C, Leszyk JD, Shaffer SA, Weng Z, Moore MJ. 2012. The cellular EJC
interactome reveals higher-order mRNP structure and an EJC-SR protein nexus. *Cell* **151**: 750–764.

Singh G, Ricci EP, Moore MJ. 2014. RIPiT-Seq: a high-throughput approach for footprinting RNA:protein
complexes. *Methods* **65**: 320–332.

Storey JD. 2003. The positive false discovery rate: a Bayesian interpretation and the q-value. *Ann Stat*
31: 2013–2035.

Sylvestre J, Vialette S, Corral Debrinski M, Jacq C. 2003. Long mRNAs coding for yeast mitochondrial
proteins of prokaryotic origin preferentially localize to the vicinity of mitochondria. *Genome Biol* **4**:
R44.

Tuller T, Carmi A, Vestsigian K, Navon S, Dorfan Y, Zaborske J, Pan T, Dahan O, Furman I, Pilpel Y. 2010.
An evolutionarily conserved mechanism for controlling the efficiency of protein translation. *Cell*
141: 344–354.

Valen E, Sandelin A, Winther O, Krogh A. 2009. Discovery of regulatory elements is improved by a
discriminatory approach. *PLoS Comput Biol* **5**: e1000562.

Vogel C, Marcotte EM. 2012. Insights into the regulation of protein abundance from proteomic and
transcriptomic analyses. *Nat Rev Genet* **13**: 227–232.

Wang S, Yao X. 2012. Multiclass Imbalance Problems: Analysis and Potential Solutions. *IEEE Trans Syst*
Man Cybern B Cybern. http://dx.doi.org/10.1109/TSMCB.2012.2187280.

Wang X, Lu Z, Gomez A, Hon GC, Yue Y, Han D, Fu Y, Parisien M, Dai Q, Jia G, et al. 2014. N6-
methyladenosine-dependent regulation of messenger RNA stability. *Nature* **505**: 117–120.

Wang X, Zhao BS, Roundtree IA, Lu Z, Han D, Ma H, Weng X, Chen K, Shi H, He C. 2015. N(6)-
methyladenosine Modulates Messenger RNA Translation Efficiency. *Cell* **161**: 1388–1399.

Zinshteyn B, Gilbert WV. 2013. Loss of a conserved tRNA anticodon modification perturbs cellular
signaling. *PLoS Genet* **9**: e1003675.

Can Cenik

Figure Legends:

Fig 1 | Modeling the relationship between sequence features in the early coding

region and the absence of 5'UTR introns (5UIs). (a) For all human transcripts,

information about 36 nucleotide-level features of the early coding region (first 99

nucleotides) and 5UI presence was extracted. **(b)** Transcripts containing a signal sequence

coding region (SSCR) were used to train a Random Forest classifier that modeled the

relationship between 5UI absence and 36 sequence features. **(c)** With this classifier, all

human transcripts were assigned a score that quantifies the likelihood of 5UI absence

based on specific RNA sequence features in the early coding region. Transcripts with high

scores are thus considered to have 5'-proximal intron minus-like coding regions (5IMs).

(d) "5'UTR-intron-minus-predictor" (5IMP) score distributions for SSCR-containing

transcripts shift to higher scores with later-appearing first introns, suggesting that 5IM

coding region features not only predict lack of a 5UI, but also lack of early coding region

introns. **(e)** Classifier performance was optimized by excluding 5UI⁻ transcripts with

introns appearing early in the coding region. Cross-validation performance (area under the

precision recall curve, AUPRC) was examined for a series of alternative 5IM classifiers

using different first-intron-position criterion for excluding 5UI⁻ transcripts from the

training set (Methods).

Fig 2 | Predicting 5UI status accurately using only early coding sequences. **(a)** As

judged by area under the receiver operating characteristic curve (AUROC) and AUPRC, The

5IM classifier performed well for several different transcript classes. **(b)** The distribution

of 5IMP scores reveals clear separation of 5UI⁺ and 5UI⁻ transcripts for SSCR-containing

transcripts, where each SSCR-containing transcript was scored using a classifier that did

not use that transcript in training (Methods). **(c)** Coding sequence features that are

predictive of 5' proximal intron presence are restricted to the early coding region. This was

supported by identical 5IM classifier score distributions with respect to 5UI presence for

negative control sequences, each derived from a single randomly chosen 'window'

downstream of the 3rd exon from one of the evaluated transcripts. **(d)** MSCR transcripts

exhibited a major difference in 5IMP scores based on their 5UI status even though no MSCR

transcripts were used in training the classifier. **(e)** Transcripts predicted to contain signal

peptides (SignalP+) had a 5IMP score distribution similar to that of SSCR-containing

transcripts. **(f)** After eliminating SSCR, MSCR, and SignalP+ transcripts, the remaining S−

/MSCR− SignalP− transcripts were still significantly enriched for high 5IM classifier scores

among 5UI− transcripts. **(g)** The control set of randomly chosen sequences downstream of

the 3rd exon from each transcript was used to calculate an empirical cumulative null

distribution of 5IMP scores. Using this function, we determined the p-value corresponding

to the 5IMP score for all transcripts. The red dashed line indicates the p-value

corresponding to 5% False Discovery Rate. The inset depicts the distribution of various

classes of mRNAs among the input set and 5IM transcripts.

Fig 3 | 5IM transcripts are more likely to be differentially expressed and have

sequence features associated with lower translation efficiency (a) The 5IM classifier

score was positively correlated with the propensity for mRNA structure preceding the start

codon (-ΔG) (Spearman rho=0.39; p < 2.2e-16). For each transcript, 35 nucleotides

immediately upstream of the AUG were used to calculate -ΔG (Methods). **(b)** The 5IM

classifier score was positively correlated with the propensity for mRNA structure near the

5'cap (-ΔG) (Spearman rho=0.18; p = 7.9e-130; Methods). **(c)** Transcripts that are

translationally upregulated in response to eIF4E overexpression (Larsson et al. 2007)

(blue) were enriched for higher 5IMP scores. Light green shading indicates 5IMP scores

corresponding to 5% FDR. **(d)** Transcripts with non-AUG start codons (blue) exhibited

significantly higher 5IMP scores than transcripts with a canonical ATG start codon (yellow).

(e) Higher 5IMP scores were associated with less optimal codons (as measured by the

tRNA adaptation index, tAI) for the first 33 codons. For all transcripts within each 5IMP

score category (blue-high, orange-low), the mean tAI was calculated at each codon position.

Start codon was not shown. **(f)** Transcripts with lower translation efficiency were enriched

for higher 5IMP scores. Transcripts with translation efficiency one standard deviation

below the mean ("LOW" translation, yellow) and one standard deviation higher than the

mean ("HIGH" translation, blue) were identified using ribosome profiling and RNA-Seq data

from human lymphoblastoid cell lines (Methods).

Fig 4 | 5IM transcripts with no evidence of ER-targeting are more likely to exhibit ER-

proximal ribosome occupancy. A moving average of ER-proximal ribosome occupancy

was calculated by grouping genes by 5IMP score (see Methods). We plotted the moving

average of 5IMP scores for transcripts with no evidence of ER- or mitochondrial targeting

(green) or for transcripts predicted to be mitochondrial (purple). We plotted a random

subsample of transcripts on top of the moving average (circles).

Fig 5 | 5IM transcripts harbor non-canonical Exon Junction Complex (EJC) binding

sites (a) Observed EJC binding sites (Singh et al. 2012) are shown for an example 5IM

transcript (*LAMC1*). Canonical EJC binding sites (purple) are ~24nt upstream of an exon-

intron boundary. The remaining binding sites are considered to be non-canonical (green).

(b) A CG-rich sequence motif previously identified to be enriched among ncEJC binding

sites in first exons (Singh et al. 2012) is shown **(c)** 5IMP score for transcripts with zero,

one, two or more non-canonical EJC binding sites in the first 99 coding nucleotides reveals

that transcripts with high 5IMP scores frequently harbor non-canonical EJC binding sites.

(d) Transcripts with high 5IMP scores are enriched for non-canonical EJCs regardless of

5UI presence or absence.

Fig 6 | 5IM transcripts are enriched for mRNAs with early coding region m^{1}A

modifications (a) Transcripts with m^{1}A modifications (blue) in the first 99 coding nucleotides

exhibit significant enrichment for 5IM transcripts and have higher 5IMPS scores than

transcripts without m^{1}A modifications in the first 99 coding nucleotides (yellow). **(b)**

Transcripts with m^{1}A modifications (blue) in the 5'UTR do not display a similar enrichment.

Supporting Information

Figure S1 | Association between 5IMP scores and codon optimality is restricted to the

first 30 amino acids and is not explained by nucleotide content. (a) 5IM transcripts tend

to have less optimal codons in their first 30 amino acids as measured by tRNA adaptation index

(tAI). The median tAI for each transcript was calculated and transcripts were grouped by their

5IMP scores. The distribution of median tAI was plotted as a boxplot. **(b)** We permuted the

nucleotides of the first 99 nucleotides and found that the relationship between 5IMP and tAI

was lost. This result suggested that nucleotide composition alone doesn't explain the

relationship between 5IMP and tAI. **(c)** We used the previously described negative control

sequences, each derived from a single randomly chosen in-frame 'window' downstream of the

3rd exon from one of the evaluated transcripts. We found no relationship between 5IMP score

and tAI in these regions suggesting that the observed association is restricted to the early

coding region.

Table S1- | List of features used by the 5IM classifier.

Table S2- | 5IMP scores of all human transcripts.

Table S3- | List of functional features tested for association with 5IMP scores.

A

Extract CDS features

B

Model the relationship between early coding sequence and 5UI status

C

Assign '5IMP' score reflecting 5UI– propensity

D **E**

Figure 2

A

A

B

C

All Transcripts
First 99 Coding Nucleotides

ncEJC Peaks

2+ Peaks

1 Peak

No Peaks

Density

5IMP Score

D

First 99 Coding Nucleotides

5UI⁺ Transcripts

No ncEJC peak

1+ ncEJC peak

5UI⁻ Transcripts

1+ ncEJC peak

No ncEJC peak

Density

5IMP Score

5IMP Score

A

First 99
Coding Nucleotides

B

5'UTR