1 **<u>Transcriptome analysis in whole blood reveals increased microbial diversity in schizophrenia</u>**
2
3

Authors: Loes M Olde Loohuis[#1], Serghei Mangul[#2], Anil PS Ori[1], Guillaume Jospin[3], David
Koslicki[4], Harry Taegyun Yang[2], Timothy Wu[1], Marco P Boks[5], Catherine Lomen-Hoerth[6],
Martina Wiedau-Pazos[7], Rita M Cantor[8], Willem M de Vos[9,10], René S Kahn[5], Eleazar Eskin[2], Roel
A Ophoff[*,1,5,8]

Affiliations:

[1]Center for Neurobehavioral Genetics, Semel Institute for Neuroscience and Human Behavior,
University California Los Angeles, Los Angeles, USA

[2]Department of Computer Science, University of California Los Angeles, Los Angeles, USA

[3]Davis Genome Center, University of California, Davis, USA

[4]Mathematics Department, Oregon State University, Corvallis, USA

[5]Department of Psychiatry, Brain Center Rudolf Magnus, University Medical Center Utrecht,
Utrecht, The Netherlands

[6]Department of Neurology, University of California San Francisco, San Francisco, USA

[7]Department of Neurology, David Geffen School of Medicine, University of California Los
Angeles, Los Angeles, USA

[8]Department of Human Genetics, University of California Los Angeles, Los Angeles, USA

[9]Laboratory of Microbiology, Wageningen University, Wageningen, The Netherlands

[10]Department of Bacteriology and Immunology, Immunobiology Research Program, University
of Helsinki, Helsinki, Finland.

* Correspondence to: Roel A. Ophoff ophoff@ucla.edu

# Equal contribution

Key words: RNA sequencing, microbiome, schizophrenia, unmapped reads, blood

Abstract:

The role of the human microbiome in health and disease is increasingly appreciated. We

studied the composition of microbial communities present in blood across 192 individuals,

including healthy controls and patients with three disorders affecting the brain: schizophrenia,

amyotrophic lateral sclerosis and bipolar disorder. By using high quality unmapped RNA

sequencing reads as candidate microbial reads, we performed profiling of microbial transcripts

detected in whole blood. We were able to detect a wide range of bacterial and archaeal phyla

in blood. Interestingly, we observed an increased microbial diversity in schizophrenia patients

compared to the three other groups. We replicated this finding in an independent

schizophrenia case-control cohort. This increased diversity is inversely correlated with

estimated cell abundance of a subpopulation of CD8^{+} memory T cells in healthy controls,

supporting a link between microbial products found in blood, immunity and schizophrenia.

Introduction

Microbial communities in and on the human body represent a complex mixture of eukaryotes, bacteria, archaea and viruses. In recent years, mounting evidence has demonstrated the involvement of the microbiome in human health and disease. In particular, through the 'microbiota-gut-brain-axis' (1, 2), the microbiome has been implicated in complex psychiatric disorders, including schizophrenia and major depressive disorder(3-8), possibly via an impact on intestinal permeability(9).

High-throughput sequencing offers a powerful culture-independent approach to study the underlying diversity of microbial communities in their natural habitats across different human tissues (10) and diseases (3, 11-15). The majority of current microbiome studies use fecal samples and target 16S ribosomal RNA gene sequencing (16). With the availability of comprehensive compendia of reference microbial genomes and phylogenetic marker genes (17), it has become feasible to use non-targeted sequencing data to identify the microbial species across different human tissues and diseases in a relatively inexpensive and easy way.

Other than in cases of sepsis, we currently lack a comprehensive understanding of the human microbiome in blood, as blood has been generally considered a sterile environment lacking proliferating microbes (18). However, over the past few decades, this assumption has been challenged (19, 20), and the presence of a microbiome in the blood has received increasing attention (21-23).

To explore potential connections between the microbiome and diseases of the brain, we performed a comprehensive analysis of microbial products detected in blood in almost two

hundred individuals, including patients with schizophrenia, bipolar disorder and sporadic

amyotrophic lateral sclerosis. These three disease groups represent complex polygenic traits

that affect the central nervous system with largely unknown etiology. Moreover, roles for the

microbiome in all the diseases have been previously hypothesized (5, 24-26). We used available

high quality RNA sequencing (RNA-Seq) reads from whole blood that fail to map to the human

genome as candidate microbial reads for microbial classification. We observed an increased

diversity of microbial communities in schizophrenia patients, and we replicated this finding in

an independent dataset. Careful analyses, including the use of positive and negative control

datasets, suggest that these detected phyla represent true microbial communities in whole

blood and are not present in samples due to contaminants. With the increasing number of RNA-

Seq data sets, our approach may have great potential for application across different tissues

and disease types.

Materials and Methods

A brief description of Materials and Methods follows below; see Supplementary Methods for

the full details.

Sample Description

The discovery sample consists of unaffected controls (Controls, n=49) and patients with three

brain-related disorders: schizophrenia (SCZ, n=48), amyotrophic lateral sclerosis (ALS, n=47) and

bipolar disorder (BPD, n=48). The replication sample includes Controls (n=88) and SCZ samples

(n=91). Sample recruitment of the cohorts is described in the Supplementary Methods. All

study methods were approved by the institutional review board of the University of California

at Los Angeles, San Francisco or the Medical Research Ethics Committee of the University

Medical Center Utrecht at The Netherlands. All participants provided written informed consent.

Sample sequencing

For the discovery sample, RNA-Seq libraries were prepared using Illumina's TruSeq RNA v2

protocol, including ribo-depletion protocol (Ribo-Zero Gold). In total, we obtained 6.8 billion

2x100bp paired-end reads for the primary study (35.3M ± 6.0 paired-end reads per sample).

The replication sample was processed at the same core facility using the same standardized

procedures as the discovery sample. However, the RNA-Seq libraries were prepared with

poly(A) enrichment, a procedure more selective than the total RNA that was used for the

discovery sample. A total of 3.8 billion reads were obtained (26.3M ± 12.0).

Sequence Analysis

We separated human and non-human reads, and use the latter as candidate microbial reads for taxonomic profiling of microbial communities. To identify potentially microbial reads, we developed the following pipeline. First, we filtered read pairs and singleton reads mapped to the human genome or transcriptome. Because total number of reads may affect microbial profiling, we performed normalization by sub-sampling to 100,000 reads for each sample. Next, we filtered out low-quality and low-complexity reads using FASTX and SEQCLEAN (see urls). Finally, the remaining reads were realigned to the human references using the Megablast aligner (27) in order to exclude any potentially human reads. The remaining reads were used as candidate microbial reads in subsequent analyses. Figure 1 displays an overview of our pipeline.

Taxonomic profiling

To access the assembly and richness of the microbiomial RNA in blood, we used phylogenetic marker genes to assign the candidate microbial reads to the bacterial and archaeal taxa. We used PhyloSift (v 1.0.1 with default parameters) to perform taxonomic profiling of the whole blood samples (17). PhyloSift makes use of a set of protein coding genes found to be relatively universal (i.e., present in nearly all bacterial and archaeal taxa) and have low variation in copy number between taxa. Homologs of these genes in new sequence data (e.g., the transcriptomes used here) are identified and then placed into a phylogenetic and taxonomic context by comparison to references from sequenced genomes. For our replication study, we used MetaPhlAn for microbial profiling v.1.7.7(28). MetaPhlAn was run in 2 stages; the first stage identifies the candidate microbial reads (i.e., reads hitting a marker) and the second stage

profiles meta-genomes in terms of relative abundances. We used MetaPhlAn, rather than

PhyloSift, due to differences in library preparation (polyA enrichment versus Ribo-Zero); there

were an insufficient number of reads matching the database of the marker genes curated by

PhyloSift for adequate microbial profiling of the replication sample.

Estimating Microbial diversity

Microbial diversity, or alpha diversity, within each sample was determined using the inverse

Simpson index. This index simultaneously assesses both richness (corresponding to the number

of distinct taxa) and relative abundance of the microbial communities within each sample (29).

In particular, it enables effective differentiation between the microbial communities shaped by

the dominant taxa and the communities with many taxa with even abundances (30) (*asbio* R

package). To measure sample-to-sample dissimilarities between microbial communities, we use

Bray-Curtis beta diversity index, which accounts for both changes in the abundances of the

shared taxa and for taxa uniquely present in one of the samples (*vegan* R package). Higher beta

diversity indicates higher level of dissimilarity between microbial communities, providing a link

between diversity at local scales (alpha diversity) and the diversity corresponding to total

microbial richness of the subject group (gamma diversity (31)).

Statistical analysis of microbiome diversity

To test for differences in alpha diversity between disease groups, we fit an analysis of

covariance (ANCOVA) model using normalized values of alpha, including sex and age, and

technical covariates (RNA INtegrity value (RIN), batch, flow cell lane and RNA concentration)

into the model. Bonferroni correction for multiple testing was used. To determine the relative

effect size of alpha diversity on schizophrenia status, we fit a logistic regression model including

the same covariates and measure reduction in R^2 comparing the full logistic regression model

versus a reduced model with alpha removed. Analysis of beta diversity was performed

analogously (see Supplementary Methods).

Reference-free microbiome analysis

We complement the reference-based taxonomic analysis with a reference independent

analysis. We use EMDeBruijn (https://github.com/dkoslicki/EMDeBruijn), a reference-free

approach capable of quantifying differences in microbiome composition between the samples.

EMDeBruijn compresses the k-mer counts of two given samples onto de Bruijn graphs and then

measures the minimal cost of transforming one of these graphs into the other. To determine

overlap between the results from PhyloSift and EMdeBruin, we correlated principal

components of EMdeBruin and PhyloSift by Spearman rank correlation, including all samples.

Estimation of cell proportions in whole blood

We assessed DNA methylation data from 65 controls taken from our replication sample, and we

compared methylation-derived blood cell proportions estimated using Houseman's estimation

method (32, 33) to alpha diversity after adjusting for age, gender, RIN and all technical

parameters. We tested whether alpha diversity levels are associated with cell type abundance

estimates. More details on the method, quality control pipeline of the methylation data and

statistical analysis can be found in Supplementary Methods.

Results

Studying microbial RNA in blood

To study the composition of microbial RNA in blood, we determined the microbial meta-
transcriptome present in the blood of unaffected controls (Controls, n=49) and patients with
three brain-related disorders: schizophrenia (SCZ, n=48), amyotrophic lateral sclerosis (ALS,
n=47) and bipolar disorder (BPD, n=48) (Figure 1, Table 1).

[Figure 1 about here]
[Table 1 about here]
173
Using our filtering pipeline, an average of 33,546 of 100,000 unmapped reads are identified as
high quality, unique non-host reads and were used as candidate microbial reads in our analyses.
From these, PhyloSift was able to assign an average of 1,235 reads (1.24% ± 0.41%, mean ±
standard deviation) to the bacterial and archaeal gene families. A total of 1,880 taxa were
assigned, with 23 taxa at the phylum level (Figure 2). Most of the taxa we observed derived
from bacteria (relative genomic abundance 89.8% ± 7.4%), and a smaller portion derived from
archaea (relative genomic abundance 12.28% ±6.4%)**.**

[Figure 2 about here]

In total, we observed 23 distinct microbial phyla with on average 4.1 ± 2.0 phyla per
individual. The large majority of taxa observed in our sample is not universally present in all
individuals; the single exception is Proteobacteria, which dominates all samples with 73.4% ±
18.3% relative abundance (Figure 2 dark green color). Several bacterial phyla show a broad

prevalence across individuals and disorders (present in 1/4 of the samples of each subject

group). Those phyla include Proteobacteria, Firmicutes and Cyanobacteria, with relative

abundance 73.4% ± 18.3%, 14.9 ±10.9%, and 11.0% ± 8.9% (Table S2). This is in line with recent

published work on the blood microbiome using 16S targeted metagenomic sequencing

reporting relative abundance of 80.4-87.4% and 3.0-6.4% for Proteobacteria and Firmicutes,

respectively (23).   The other two phyla identified in this study (Actinobacteria and

Bacteroidetes) were also detected in our sample in more than 25 individuals.   Although

Proteobacteria and Firmicutes, are commonly associated with the human microbiome (34),

some members of these phyla might be associated with reagent and environmental

contaminants (35, 36).

To validate our pipeline and investigate the possibility of contamination introduced

during RNA isolation, library preparation and sequencing steps, we performed both negative

and positive control experiments (see Supplementary Results and Methods for details). In brief:

no microbiome sequences were detected in transcriptome data in lymphopblast cell lines

(negative control), and we only detected the Chlamydiae phylum in RNA-Seq from cells infected

with Chlamydiae (positive control). We examined experimental procedures and technical

parameters on microbial composition, and we observed no link between the presence of

microbial communities and possible confounders.

To compare the inferred microbial composition found in blood with that in other body

sites, we used taxonomic composition of 499 meta-genomic samples from Human Microbiome

Project (HMP) obtained by MetaPhlAn or five major body habitats (gut, oral, airways, and skin)

(10). Of the 23 phyla discovered in our sample, 15 were also found in HMP samples, of which 13

are confirmed by at least ten samples. Our data suggest that the predominant phyla detected in

blood are most closely related to the known oral and gut microbiome (Table S2). Comparing the

microbial composition of whole blood with the microbiome detected in atherosclerotic plaques

(37), we observe that the four phyla that together make up for >97% of the microbiome in

plaques are also identified in our sample (Firmicutes, Bacteroidetes, Proteobacteria,

Actinobacteria).

Finally, it should be noted that the sequencing technology does not allow for

identification of the origin of microbial RNA. That is, we can not distinguish whether the

observed microbial signatures in blood are originate from bacterial communities actually

present in the blood, or whether the RNA crossed into the blood stream from elsewhere.

Increased microbial diversity in schizophrenia samples

To evaluate potential differences in microbial profiles of individuals with the different disorders

(SCZ, BPD, ALS) and unaffected controls, we explored the composition and richness of the

microbial communities across the groups.

We observed increased alpha diversity in schizophrenia samples compared to all other

groups (*ANCOVA* P < 0.005 for all groups, Figure 3a, Table 2 and Table S5, *Bonferroni*

correction). These differences are corrected for covariates and are independent of potential

confounders, such as experimenter and RNA extraction run (Figure S1 and S2), and they are not

the consequence of a different number of reads being detected as microbial in schizophrenia

samples (see Supplementary Results). No significant differences were observed between the

three remaining groups (BPD, ALS, Controls). In our sample, alpha diversity was found to be a

significant predictor of schizophrenia status and explained 5.0% of the variation as measured by

reduction in Nagelkerke's R^2 from logistic regression. We observe no correlation between

polygenic risk scores (39) and alpha diversity in our schizophrenia sample (n= 32, Kendall's tau=

0.008, *P* = 0.96, Supplementary Methods). We also did not observe differences in alpha

diversity between sexes or across ages, nor are our results driven by the relatively younger

schizophrenia cohort (Supplementary Results). Alpha diversity at other main taxonomic ranks

yields a similar pattern of increased diversity in schizophrenia (Figure S3).

The increased diversity observed in schizophrenia patients may be due to specific phyla

characteristic to schizophrenia, or due to a more general increased microbial diversity in people

affected by the disease. To investigate this, we compared diversity across individuals within the

schizophrenia group to control samples. We compared beta diversity across pairs of samples

with schizophrenia and controls, resulting in three subject groups: SCZ_Controls, SCZ_SCZ and

Controls_Controls. The lowest diversity was observed in the Controls_Controls group (0.43 ±

0.21), followed by SCZ_SCZ (0.50 ± 0.14), and the highest beta diversity values were observed

for SCZ_Controls (0.51 ± 0.17) (P< 0.05 for each comparison, by ANCOVA after correcting for

three tests). This result was confirmed by permanova (P<0.001) based on 1,000 permutations.

Thus, the observed increased alpha diversity in schizophrenia is not caused by a particular

microbial profile, but most likely represents a non-specific overall increased microbial burden

(see also Figure S4 and Supplementary Results).

In addition to measuring individual microbial diversity (alpha), and diversity between

individuals (beta), we measured the total richness of the microbiome by the total number of

distinct taxa of the microbiome community observed within an entire subject group (gamma

diversity (40)). We observed that all 23 distinct phyla are observed in schizophrenia:

gamma(SCZ)=23 compared to gamma(Controls)=20, gamma(ALS)=16 and gamma(BPD)=18.

We complemented reference-based methods (PhyloSift and MetaPhlAn) with

EMDeBruijn, a reference-independent method. EMDeBruin distances measured between

samples correlated significantly with beta diversity (Spearman rank P < 2.2e-16, rho = 0.37,

including SCZ and Controls). Also, EMDeBruijn PCs correlated with principal components

obtained from edge PCA based on the PhyloSift taxonomic classification (correlation between

EMDeBruijn PC1, and PhyloSift PC1 is P = 1.824e-09; Spearman rank correlation is rho = -0.42;

see also Figure S5). After correcting covariates, the first three EMDeBruijn PCs are significant

predictors of schizophrenia status, and jointly explained 7.1% of the variance (P< 0.05 for each

PC).

[Figure 3 about here]

[Table 2 about here]

Group differences of individual phyla

In addition to a global difference between schizophrenia and the other groups, we also

investigated whether there are particular individual phyla contributing to the differences

between schizophrenia and other groups. There are two phyla detected more often in

schizophrenia cases versus all the other groups: Plactomycetes, observed in 20 SCZ cases

compared to 3(ALS) 2(BPD) 5(Controls) (P= 0.0002 Fisher's exact for four groups, Bonferroni

corrected for 23 tests P=0.0057) and Thermotogae, observed in 20 SCZ cases compared to 6

ALS, 3 BPD and 6 Controls (P= 0.0006 Fisher's exact, corrected P=0.014). No outliers were

observed for the other groups (see Table S7).

Replication

We performed a replication experiment in an independent case-control sample: schizophrenia

(SCZ n=91) and healthy controls (Controls n=88) (See Table S1.D). MetaPhlAn was able to assign

5,174 reads (0.089% ± 0.039%, mean ± standard deviation) on average to the bacterial gene

families.

Schizophrenia samples showed increased alpha diversity on genus level (2.73 ± 0.77 for

cases, versus 2.32 ± 0.57 for controls, corrected P = 0.003 Figure 3b) and explained 2.5% of

variance as measured by reduction in Nagelkerke R^2, thus replicating our main finding of

increased diversity in schizophrenia. While our original analysis was performed on the phylum

level, in our discovery sample we observe a similar increase of diversity at the genus level (see

Figure S3). Similar to our discovery cohort, we observed no significant correlation between

alpha diversity and age or differences across gender. Beta diversity and EMDeBruijn analyses

also show similar, though not identical, patterns of nonspecific increased diversity in

schizophrenia samples (Supplementary Results).

Cell type composition and diversity

We hypothesized that differences in microbial diversity may be linked to whole blood cell type

composition. Our analysis shows that the proportion of one cell type, $CD8^+ CD28^- CD45RA^-$ cells,

is significantly negatively correlated with alpha diversity after correction for all other cell-count

estimates as estimated from whole blood DNA methylation data (correlation = -0.41, P=7.3e-4,

n= 65 Controls from the Replication study, Figure S6, Table S6). These cells are T cells that lack

$CD8^+$ naïve cell markers CD28 and CD45RA and are thought to represent a subpopulation of

$CD8^+$ memory T cells (41, 42). We observed that low alpha diversity correlates with high levels

of cell abundance of this population of T cells.

Discussion

We used high throughput RNA sequencing from whole blood to perform microbiome profiling

and identified an increased diversity in schizophrenia patients.

While other studies of human microbiome using RNA-Seq have been conducted (43, 44),

this is the first assessing the microbiome from whole blood by using unmapped non-human

reads. Despite the fact that transcripts are present at much lower fractions than human reads,

we were able to detect microbial transcripts from bacteria and archaea in almost all samples.

The microbes found in blood are thought to be originating from the gut as well as oral cavities

(45, 46), which is in line with our finding that the microbial profiles found in our study most

closely resemble the gut and oral microbiome as profiled by the HMP (10). The taxonomic

profile of the cohort samples suggests the prevalence of the several phyla, Proteobacteria,

Firmicutes and Cyanobacteria, across individuals and different disorders included in our study.

This is in line with a recent study that used 16S targeted meta-genomic sequencing, which

reported Proteobacteria and Firmicutes among the most abundant phyla detected in blood

(23).

Rigorous quality control is critically important for any high-throughput sequencing

project, especially in the context of studying the microbiome (35). To this end, we performed

both negative and positive quality control experiments, and we carefully evaluated possible

contamination effects introduced during the experiments. Our results suggest that the detected

phyla represent true microbial communities in whole blood and are not due to contaminants.

However, it should be noted that whether only the microbial products crossed into the

bloodstream or whether the microbes themselves are present in blood cannot be answered

using sequencing techniques. Future experiments, for example, using microscopy, culturing or

direct measures of gut permeability, may be able to shed light on this question.

The most striking finding of our study that relates to diseases affecting the central

nervous system is the increased microbial alpha diversity in schizophrenia patients compared to

controls and the other two disease groups (ALS, BPD). We replicate this finding in an

independent cohort of schizophrenia cases and controls. The replication experiment, while

based on different library preparation (Ribo-Zero versus Poly(A)), provides strong evidence for

an increased alpha diversity of the microbiome detected in blood in schizophrenia and explains

roughly 5% of disease variation. We not only observe an increased individual microbial

diversity, but also an increased diversity between individuals (Beta diversity) with schizophrenia

compared to controls, rendering it unlikely that a single phylum or microbial profile is causing

the disease-specific increase in diversity. Nevertheless, in our study we observed that two phyla

in particular, Planctomycetes and Thermotogae, were present in significantly more

schizophrenia samples when compared to the other groups. Interestingly, Planctomycetes is

group of gram-negative bacteria closely related to Verrucomicrobia and Chlamydiae; together

these comprise the Planctomycetes-Verrucomicrobia-Chlamydiae superphylum (47). From

peripheral blood, infection with Chlamydiaceae species has been reported to be increased in

schizophrenia (40%) compared to controls (7%) (48). Since Chlamydiae is one of the taxa of the

superphylum, it is possible that the increase in Planctomycetes we observe is related to the

observed increase in Chlamydiaceae species. As the collection of available reference genomes

continues to grow and improve, future studies are needed to corroborate and refine these

findings.

For the study of microbiome diversity, we employed reference-based methods

(PhyloSift and MethPhlAn) and the EMDebruin method, a purely reference-agnostic approach.

The latter showed strong correspondence to both reference-based methods, highlighting the

value of this unbiased sequence-based analysis for investigating microbial differences across

groups. However, in addition to differences in distribution of microbial transcripts, EMDebruin

may capture variation of other, yet unknown, origin.

In addition to our observation that microbial diversity is more generally increased in

schizophrenia, our study demonstrates the value of analyzing non-human reads present in the

RNA-Seq data to study the microbial composition of a tissue of interest (49, 50). The RNA-Seq

approach avoids biases introduced by primers in targeted 16S ribosomal RNA gene profiling. In

addition, since *mRNA stability is low in prokaryotes,* RNA-Seq might offer a potential advantage

of avoiding contamination of genomic DNA by dead cells compared to genome sequencing (51).

Given the many large-scale RNA-Seq datasets that are becoming available, we anticipate that

high-throughput meta-transcriptome-based microbiome profiling will find broad applications as

a hypothesis-generating tool in studies across different tissues and disease types.

The increased microbial diversity observed in schizophrenia could be part of the disease

etiology (i.e., causing schizophrenia) or may be a secondary effect of disease status. In our

sample, we observed no correlation between increased microbial diversity and genetic risk for

schizophrenia as measured by polygenic risk scores (52). In addition, it is remarkable that

bipolar disorder, which is genetically and clinically correlated to schizophrenia (53), does not

show a similar increased diversity. We did observe a strong inverse correlation between

increased diversity and estimated cell abundance of a population of T-cells in healthy controls.

Even though this finding is based on indirect cell count measures using DNA methylation data

(41), the significant correlation highlights a likely close connection between the immune system

and the blood microbiome, a relationship that has been documented before (54). More

extensive cell count measures and/or better markers of immune sensing of microbial products

could be used to study this relationship more directly. In the absence of a direct link with

genetic susceptibility and the reported correlation with the immune system, we hypothesize

that the observed effect in schizophrenia may be mostly a consequence of disease. This may be

affected by lifestyle and/or health status differences of schizophrenia patients, including

smoking, treatment plans, (chronic) infection, GI status, the use of probiotics, antibiotics and

other drug use or other environmental exposures. Future targeted and/or longitudinal studies

with larger sample sizes, detailed clinical phenotypes and more in-depth sequencing are

needed to corroborate this hypothesis. Another interesting direction for future work is to study

gut permeability in the context of our findings more directly. For example, how does damage to

the gut (such as measured using I-FABP) affect observed microbial diversity in blood? These

studies would likely result in an expanded understanding of the functional mechanisms

underlying the connection between the human immune system, microbiome, and disease

etiology. In particular, we hope that these future efforts will provide a useful quantitative and

qualitative assessment of the microbiome and its role across the gut-blood barrier in the

context of psychiatric disorders.

Availability of Data and Materials:

The data discussed in this publication have been deposited in NCBI's Gene Expression Omnibus (55) and are accessible through GEO Series accession number GSE80974 (https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE80974).

Acknowledgments:

This work is supported by NIH/NIMH R01 5R01MH090553 (BP samples), 5R01NS058980 (ALS samples) and R01MH078075 (SCZ, Control samples), R21MH098035 (replication sample) awarded to R.A.O. L.M.O.L was financially supported by the National Institute of Neurological Disorders And Stroke of the National Institutes of Health under Award Number T32NS048004. The authors thank Dr. Jonathan Eisen for helpful discussions and insights throughout the course of this project. We thank Dr. Lana Martin for helpful edits to the manuscript. Finally, we thank the study subjects for their participation.

Financial Declarations:

The authors have no conflicts of interest to declare.

References and Notes

1. Clarke G, Grenham S, Scully P, Fitzgerald P, Moloney RD, Shanahan F, et al. The microbiome-gut-brain axis during early life regulates the hippocampal serotonergic system in a sex-dependent manner. Mol Psychiatry. 2013;18(6):666-73.

2. Cryan JF, Dinan TG. Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour. Nat Rev Neurosci. 2012;13(10):701-12.

3. Hsiao EY, McBride SW, Hsien S, Sharon G, Hyde ER, McCue T, et al. Microbiota modulate behavioral and physiological abnormalities associated with neurodevelopmental disorders. Cell. 2013;155(7):1451-63.

4. Erny D, Hrabe de Angelis AL, Jaitin D, Wieghofer P, Staszewski O, David E, et al. Host microbiota constantly control maturation and function of microglia in the CNS. Nat Neurosci. 2015;18(7):965-77.

5. Foster JA, McVey Neufeld KA. Gut-brain axis: how the microbiome influences anxiety and depression. Trends Neurosci. 2013;36(5):305-12.

6. Castro-Nallar E, Bendall ML, Perez-Losada M, Sabuncyan S, Severance EG, Dickerson FB, et al. Composition, taxonomy and functional diversity of the oropharynx microbiome in individuals with schizophrenia and controls. PeerJ. 2015;3:e1140.

7. Rogers GB, Keating DJ, Young RL, Wong ML, Licinio J, Wesselingh S. From gut dysbiosis to altered brain function and mental illness: mechanisms and pathways. Mol Psychiatry. 2016;21(6):738-48.

8. Zheng P, Zeng B, Zhou C, Liu M, Fang Z, Xu X, et al. Gut microbiome remodeling induces depressive-like behaviors through a pathway mediated by the host's metabolism. Mol Psychiatry. 2016;21(6):786-96.

9. Kelly JR, Kennedy PJ, Cryan JF, Dinan TG, Clarke G, Hyland NP. Breaking down the barriers: the gut microbiome, intestinal permeability and stress-related psychiatric disorders. Front Cell Neurosci. 2015;9:392.

10. Human Microbiome Project C. Structure, function and diversity of the healthy human microbiome. Nature. 2012;486(7402):207-14.

11. Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An obesity-associated gut microbiome with increased capacity for energy harvest. Nature. 2006;444(7122):1027-31.

12. Turnbaugh PJ, Hamady M, Yatsunenko T, Cantarel BL, Duncan A, Ley RE, et al. A core gut microbiome in obese and lean twins. Nature. 2009;457(7228):480-4.

13. Abu-Shanab A, Quigley EM. The role of the gut microbiota in nonalcoholic fatty liver disease. Nature reviews Gastroenterology & hepatology. 2010;7(12):691-701.

14. Cho I, Blaser MJ. The human microbiome: at the interface of health and disease. Nature reviews Genetics. 2012;13(4):260-70.

15. Greenblum S, Turnbaugh PJ, Borenstein E. Metagenomic systems biology of the human gut microbiome reveals topological shifts associated with obesity and inflammatory bowel disease. Proc Natl Acad Sci U S A. 2012;109(2):594-9.

16. de Vos WM, de Vos EA. Role of the intestinal microbiome in health and disease: from correlation to causation. Nutrition reviews. 2012;70 Suppl 1:S45-56.

17. Darling AE, Jospin G, Lowe E, Matsen FAt, Bik HM, Eisen JA. PhyloSift: phylogenetic analysis of genomes and metagenomes. PeerJ. 2014;2:e243.

18. Drennan MR. What is "Sterile Blood"? British Medical Journal. 1942;2(4269):526-.

19.   McLaughlin RW, Vali H, Lau PC, Palfree RG, De Ciccio A, Sirois M, et al. Are there
naturally occurring pleomorphic bacteria in the blood of healthy humans? J Clin Microbiol.
2002;40(12):4771-5.
20.   Nikkari S, McLaughlin IJ, Bi W, Dodge DE, Relman DA. Does blood of healthy subjects
contain bacterial ribosomal DNA? J Clin Microbiol. 2001;39(5):1956-9.
21.   Amar J, Serino M, Lange C, Chabo C, Iacovoni J, Mondot S, et al. Involvement of tissue
bacteria in the onset of diabetes in humans: evidence for a concept. Diabetologia.
2011;54(12):3055-61.
22.   Sato J, Kanazawa A, Ikeda F, Yoshihara T, Goto H, Abe H, et al. Gut dysbiosis and
detection of "live gut bacteria" in blood of Japanese patients with type 2 diabetes. Diabetes Care.
2014;37(8):2343-50.
23.   Paisse S, Valle C, Servant F, Courtney M, Burcelin R, Amar J, et al. Comprehensive
description of blood microbiome from healthy donors assessed by 16S targeted metagenomic
sequencing. Transfusion. 2016.
24.   Wu S, Yi J, Zhang YG, Zhou J, Sun J. Leaky intestine and impaired microbiome in an
amyotrophic lateral sclerosis mouse model. Physiol Rep. 2015;3(4).
25.   Dinan TG, Borre YE, Cryan JF. Genomics of schizophrenia: time to consider the gut
microbiome? Mol Psychiatry. 2014;19(12):1252-7.
26.   Dickerson F, Severance E, Yolken R. The microbiome, immunity, and schizophrenia and
bipolar disorder. Brain Behav Immun. 2016.
27.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
architecture and applications. BMC bioinformatics. 2009;10:421.
28.   Segata N, Waldron L, Ballarini A, Narasimhan V, Jousson O, Huttenhower C.
Metagenomic microbial community profiling using unique clade-specific marker genes. Nature
methods. 2012;9(8):811-4.
29.   Simpson EH. Measurement of diversity. Nature. 1949.
30.   Whittaker RH. Evolution and measurement of species diversity. Taxon. 1972:213-51.
31.   Koleff P, Gaston KJ, Lennon JJ. Measuring beta diversity for presence–absence data.
Journal of Animal Ecology. 2003;72(3):367-82.
32.   Houseman EA, Accomando WP, Koestler DC, Christensen BC, Marsit CJ, Nelson HH, et
al. DNA methylation arrays as surrogate measures of cell mixture distribution. BMC
bioinformatics. 2012;13:86.
33.   Aryee MJ, Jaffe AE, Corrada-Bravo H, Ladd-Acosta C, Feinberg AP, Hansen KD, et al.
Minfi: a flexible and comprehensive Bioconductor package for the analysis of Infinium DNA
methylation microarrays. Bioinformatics. 2014;30(10):1363-9.
34.   Human Microbiome Project C. A framework for human microbiome research. Nature.
2012;486(7402):215-21.
35.   Salter SJ, Cox MJ, Turek EM, Calus ST, Cookson WO, Moffatt MF, et al. Reagent and
laboratory contamination can critically impact sequence-based microbiome analyses. BMC Biol.
2014;12:87.
36.   Strong MJ, Xu G, Morici L, Splinter Bon-Durant S, Baddoo M, Lin Z, et al. Microbial
contamination in next generation sequencing: implications for sequence-based analysis of
clinical samples. PLoS Pathog. 2014;10(11):e1004437.
37.   Koren O, Spor A, Felin J, Fak F, Stombaugh J, Tremaroli V, et al. Human oral, gut, and
plaque microbiota in patients with atherosclerosis. Proc Natl Acad Sci U S A. 2011;108 Suppl
1:4592-8.

38.   Consortium GT. Human genomics. The Genotype-Tissue Expression (GTEx) pilot
analysis: multitissue gene regulation in humans. Science. 2015;348(6235):648-60.
39.   Schizophrenia Working Group of the Psychiatric Genomics Consortium. Biological
insights from 108 schizophrenia-associated genetic loci. Nature. 2014;511:421--7
.
40.   Jost L. Partitioning diversity into independent alpha and beta components. Ecology.
2007;88(10):2427-39.
41.   Horvath S, Levine AJ. HIV-1 Infection Accelerates Age According to the Epigenetic
Clock. J Infect Dis. 2015.
42.   Koch S, Larbi A, Derhovanessian E, Ozcelik D, Naumova E, Pawelec G. Multiparameter
flow cytometric analysis of CD4 and CD8 T cell subsets in young and old people. Immun
Ageing. 2008;5:6.
43.   Croucher NJ, Thomson NR. Studying bacterial transcriptomes using RNA-seq. Curr Opin
Microbiol. 2010;13(5):619-24.
44.   McClure R, Balasubramanian D, Sun Y, Bobrovskyy M, Sumby P, Genco CA, et al.
Computational analysis of bacterial RNA-Seq data. Nucleic Acids Res. 2013;41(14):e140.
45.   Potgieter M, Bester J, Kell DB, Pretorius E. The dormant blood microbiome in chronic,
inflammatory diseases. FEMS Microbiol Rev. 2015;39(4):567-91.
46.   Spadoni I, Zagato E, Bertocchi A, Paolinelli R, Hot E, Di Sabatino A, et al. A gut-
vascular barrier controls the systemic dissemination of bacteria. Science. 2015;350(6262):830-4.
47.   Hou S, Makarova KS, Saw JH, Senin P, Ly BV, Zhou Z, et al. Complete genome
sequence of the extremely acidophilic methanotroph isolate V4, Methylacidiphilum infernorum,
a representative of the bacterial phylum Verrucomicrobia. Biol Direct. 2008;3:26.
48.   Fellerhoff B, Laumbacher B, Mueller N, Gu S, Wank R. Associations between
Chlamydophila infections, schizophrenia and risk of HLA-A10. Mol Psychiatry. 2007;12(3):264-
72.
49Jorth P, Turner KH, Gumus P, Nizam N, Buduneli N, Whiteley M. Metatranscriptomics of the
human oral microbiome during health and disease. MBio. 2014;5(2):e01012-14.
50.   Kostic AD, Ojesina AI, Pedamallu CS, Jung J, Verhaak RG, Getz G, et al. PathSeq:
software to identify or discover microbes by deep sequencing of human tissue. Nat Biotechnol.
2011;29(5):393-6.
51.   Ben-Amor K, Heilig H, Smidt H, Vaughan EE, Abee T, de Vos WM. Genetic diversity of
viable, injured, and dead fecal bacteria assessed by fluorescence-activated cell sorting and 16S
rRNA gene analysis. Applied and environmental microbiology. 2005;71(8):4679-89.
52.   Ripke S, O'Dushlaine C, Chambert K, Moran JL, Kahler AK, Akterin S, et al. Genome-
wide association analysis identifies 13 new risk loci for schizophrenia. Nat Genet.
2013;45(10):1150-9.
53.   Bulik-Sullivan B, Finucane HK, Anttila V, Gusev A, Day FR, Loh PR, et al. An atlas of
genetic correlations across human diseases and traits. Nature genetics. 2015;47(11):1236-41.
54.   Belkaid Y, Hand TW. Role of the microbiota in immunity and inflammation. Cell.
2014;157(1):121-41.
55.   Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression
and hybridization array data repository. Nucleic Acids Res. 2002;30(1):207-10.
542
543

Figure Legends

Figure 1. Microbial profiling using RNA-Seq data from whole blood. (A) We analyzed a cohort of

192 individuals from four subject groups, i.e. Schizophrenia (SCZ, n=48), amyotrophic lateral

sclerosis (ALS n=47), bipolar disorder (BPD n=48), unaffected control subjects (Controls n=49).

(B) Peripheral blood was collected for RNA collection. (B) RNA-Seq libraries were prepared from

total RNA using ribo-depletion protocol. Reads that failed to map to the human reference

genome and transcriptome were sub-sampled and further filtered to exclude low-quality, low

complexity, and remaining potentially human reads. High quality, unique, non-host reads were

used to determine the taxonomic composition and diversity of the detected microbiome. See

also Table S1.

Figure 2. Relative abundances of microbial taxa at phylum level. Phylogenetic classification is

performed using PhyloSift, which is able to assign the filtered candidate microbial reads to the

microbial genes from 23 distinct taxa on the phylum level.

Figure 3. Increased diversity of microbiome detected in blood in schizophrenia samples. (A)

Alpha diversity per sample for four subject groups (Controls, ALS, BPD, SCZ), measured using

the inverse Simpson index on the phylum level of classification. Schizophrenia samples show

increased diversity compared to all three other groups (ANCOVA $P < 0.005$ for all groups, after

adjustment of covariates, see also Methods, Table S5 and Figure S3). (B) Alpha diversity per

sample of schizophrenia cases and controls, measured using the inverse Simpson index on the

567   genus level of classification. Schizophrenia samples show increased within-subject diversity

568   compared to Controls (P = 0.003 after adjustment of covariates).