Thanatotranscriptome: genes actively expressed after organismal death

Authors: Alex E. Pozhitkov[1,2], Rafik Neme[2], Tomislav Domazet-Lošo[3,4], Brian G. Leroux[1], Shivani Soni[5], Diethard Tautz[2] and Peter A. Noble[,5,6,7*].

Affiliations:

[1]Department of Oral Health Sciences, University of Washington, Box 357444, Seattle, WA USA 98195.

[2]Max-Planck-Institute for Evolutionary Biology, August-Thienemann-Strasse 2, 24306 Ploen, Germany.

[3]Laboratory of Evolutionary Genetics, Division of Molecular Biology, Ruđer Bošković Institute, 10002 Zagreb, Croatia.

[4]Catholic University of Croatia, Ilica 242, Zagreb, Croatia.

[5]Department of Periodontics, School of Dentistry, University of Washington, Box 357444, Seattle, WA USA 98195.

[6]Department of Biological Sciences, Alabama State University, Montgomery, AL, USA 36101-0271.

[7]PhD Program in Microbiology, Alabama State University, Montgomery, AL, USA 36101-0271.

*Correspondence to:

Peter A. Noble

Email: panoble@washington.edu

Phone: 206-409-6664.

AEP: pozhit@uw.edu

RN: rneme@evolbio.mpg.de

TDL: tdomazet@unicath.hr

BGL: leroux@uw.edu

SS: ssoni@alasu.edu

DT: tautz@evolbio.mpg.edu

Short Title: Transcriptional dynamics after organismal death

ABSTRACT

A continuing enigma in the study of biological systems is what happens to highly ordered structures, far from equilibrium, when their regulatory systems suddenly become disabled. In life, genetic and epigenetic networks precisely coordinate the expression of genes -- but in death, it is not known if gene expression diminishes gradually or abruptly stops or if specific genes are involved. We investigated the 'unwinding of the clock' by identifying upregulated genes, assessing their functions, and comparing their transcriptional profiles through postmortem time in two species, mouse and zebrafish. We found transcriptional abundance profiles of 1,063 genes were significantly changed after death of healthy adult animals in a time series spanning from life to 48 or 96 h postmortem. Ordination plots revealed non-random patterns in profiles by time. While most thanatotranscriptome (thanatos-, Greek *defn.* death) transcript levels increased within 0.5 h postmortem, some increased only at 24 and 48 h. Functional characterization of the most abundant transcripts revealed the following categories: stress, immunity, inflammation, apoptosis, transport, development, epigenetic regulation, and cancer. The increase of transcript abundance was presumably due to thermodynamic and kinetic controls encountered such as the activation of epigenetic modification genes responsible for unraveling the nucleosomes, which enabled transcription of previously silenced genes (e.g., development genes). The fact that new molecules were synthesized at 48 to 96 h postmortem suggests sufficient energy and resources to maintain self-organizing processes. A step-wise shutdown occurs in organismal death that is manifested by the apparent upregulation of genes with various abundance maxima and durations. The results are of significance to transplantology and molecular biology.

KEYWORDS

Postmortem transcriptome; postmortem gene expression; Gene meters; calibrated DNA microarrays, thanatotranscriptome; cancer; signaling, development, immunity, apoptosis, transport, inflammation; epigenetic regulation; thermodynamic sink; transplantology, forensic science.

INTRODUCTION

A healthy adult vertebrate is a complex biological system capable of highly elaborate functions such as the ability to move, communicate, and sense the environment -- all at the same time. These functions are tightly regulated by genetic and epigenetic networks through multiple feedback loops that precisely coordinate the expression of thousands of genes at the right time, in the right place, and in the right level [1]. Together, these networks maintain homeostasis and thus sustain 'life' of a biological system.

While much is known about gene expression circuits in life, there is a paucity of information about what happens to these circuits after organismal death. For example, it is not well known whether gene expression diminishes gradually or abruptly stops in death -- nor whether specific genes are newly expressed or upregulated. In organismal 'death', defined here as the cessation of the highly elaborate system functions in vertebrates, we conjecture that there is a gradual disengagement and loss of global regulatory networks, but this could result in a regulatory response of genes involved in survival and stress compensation. To test this, we examined postmortem gene expression in two model organisms: the zebrafish, *Danio rerio,* and the house mouse, *Mus musculus.* The purpose of the research was to investigate the "unwinding of the clock" by identifying genes whose expression increases (i.e., nominally upregulated) and assessing their functions based on the primary literature. The biological systems investigated in this study are different from those examined in other studies, such as individual dead and/or injured cells in live organisms, i.e., apoptosis and necrosis (reviewed in refs. [2-5]). In contrast to previous studies, gene expression from the entire *D. rerio* body, and the brains and livers of *M. musculus* were assessed through postmortem time. Gene expression was measured using the 'Gene Meter' approach that precisely reports gene transcript abundances based on a calibration curve for each microarray probe [6].

MATERIALS AND METHODS

Induced death and postmortem incubation. Zebrafish Forty-four female *Danio rerio* were transferred from several flow-through aquaria kept at 28°C to a glass beaker containing 1 L of aquarium water. Four individuals were immediately taken out, snap frozen in liquid nitrogen, and stored in Falcon tubes at -80°C (two zebrafish per tube). These samples were designated as the first set of live controls. A second set of live controls was immersed in an open cylinder (described below). Two sets of live controls were used to determine if putting the zebrafish back into their native environment had any effects on gene expression (we later discovered no significant effects).

The rest of the zebrafish were subjected to sudden death by immersion in a "kill" chamber. The chamber consisted of an 8 L styrofoam container filled with chilled ice water. To synchronize the death of the rest of the zebrafish, they were transferred to an open cylinder with a mesh-covered bottom and the cylinder was immersed into the kill chamber. After 20 to 30 s of immersion, four zebrafish were retrieved from the chamber, snap frozen in liquid nitrogen, and stored at -80°C (two zebrafish per Falcon tube). These samples were designated as the second set of live controls. The remaining zebrafish were

kept in the kill chamber for 5 min and then the cylinder was transferred to a flow-through aquarium kept at 28°C so that they were returned to their native environment.

Postmortem sampling of the zebrafish occurred at: time 0, 15 min, 30 min, 1 h, 4 h, 8 h, 12 h, 24 h, 48 h, and 96 h. For each sampling time, four expired zebrafish were retrieved from the cylinder, snap frozen in liquid nitrogen, and stored at -80°C in Falcon tubes (two zebrafish to a tube). One zebrafish sample was lost, but extraction volumes were adjusted to one organism.

Mouse The mouse strain C57BL/6JRj (Janvier SAS, France) was used for our experiments. The mice were 20-week old males of approximately the same weight. The mice were highly inbred and were expected to have a homogenous genetic background. Prior to euthanasia, the mice were kept at room temperature and were given *ad libitum* access to food and water. Each mouse was euthanized by cervical dislocation and placed in an individual plastic bag with holes to allow air / gas exchange. The bagged carcasses were kept at room temperature in a large, open polystyrene container. Sampling of the deceased mice began at 0 h (postmortem time zero) and continued at 30 min, 1 h, 6 h, 12 h, 24 h and 48 h postmortem. At each sample time, 3 mice were sampled (except for 48h where 2 mice were sampled) and the entire brain (plus stem) and two portions of the liver were extracted from each mouse. For liver samples, clippings were taken from the foremost and rightmost lobes of the liver. The brain and liver samples were snap frozen in liquid nitrogen and stored individually in Falcon tubes at -80°C.

The euthanasia methods outlined above are approved by the American Veterinary Medical Association (AVMA) Guidelines for Euthanasia (www.avma.org) and carried out by personnel of the Max-Planck-Institute for Evolutionary Biology (Ploen, Germany). All animal work: followed the legal requirements, was registered under number V312-72241.123-34 (97-8/07) and approved by the ethics commission of the Ministerium für Landwirtschaft, Umwelt und ländliche Räume, Kiel (Germany) on 27. 12. 2007.

RNA extraction, labeling, hybridization and DNA microarrays. The number of biologically distinct organisms was 43 for zebrafish and 20 for mice. Samples from two fish were pooled for analysis, resulting in two replicate measurements at each time point. The number of replicated measurements for mice was three at each of the first six time points and two at 48h. Thus, the total number of samples analyzed was 22 for zebrafish and 20 for mice. For the zebrafish, samples were mixed with 20 ml of Trizol and homogenized using a TissueLyzer (Qiagen). For the mice, 100 mg of brain or liver samples were mixed with 1 ml of Trizol and homogenized. One ml of the emulsion from each sample was put into a fresh 1.5 ml centrifuge tube for RNA extraction and the rest was frozen at -80°C.

RNA was extracted by adding 200 µl of chloroform, vortexing the sample, and incubating it at 25°C for 3 min. After centrifugation (15 min at 12000 x g at 4°C), the supernatant (approx. 350 µl) was transferred to a fresh 1.5 ml tube containing an equal volume of 70% ethanol. The tube was vortexed, centrifuged and purified following the procedures outlined in the PureLink RNA Mini Kit (Life Technologies, USA).

The isolated RNA, 400 ng per sample, was labeled, purified and hybridized according to the One-Color Microarray-based Gene Expression Analysis (Quick Amp Labeling) with Tecan HS Pro Hybridization kit (Agilent Technologies). For the zebrafish, the labeled

RNA was hybridized to the Zebrafish (v2) Gene Expression Microarray (Design ID
019161). For the mouse, the labeled RNA was hybridized to the SurePrint G3 Mouse GE
8x60K Microarray Design ID 028005 (Agilent Technologies). The microarrays were
loaded with 1.65 µg of labeled cRNA for each postmortem sample.

Microarray calibration. Oligonucleotide (60 nt) probes on the zebrafish and mouse
microarrays were calibrated using pooled labeled cRNA of all zebrafish and all mouse
postmortem samples, respectively. The dilution series for the Zebrafish array was created
using the following concentrations of labeled cRNA: 0.41, 0.83, 1.66, 1.66, 1.66, 3.29,
6.60, and 8.26 µg. The dilution series for the Mouse arrays was created using the
following concentrations of labeled cRNA: 0.17, 0.33, 0.66, 1.32, 2.64, 5.28, 7.92, and
10.40 µg. Calibration involved plotting the signal intensities of the probes against a
dilution factor and determining the isotherm model (e.g., Freundlich and/or Langmuir)
that best fit the relationship between signal intensities and gene abundances.

Consider zebrafish gene transcripts targeted by A_15_P110618 (which happens to be one
of the transcriptional profiles of gene *Hsp70.3* shown in Fig 2A). External file
FishProbesParameters.txt shows that a Freundlich model best fit the dilution curve with
R^2=0.99. The equation for this probe is the following:

$SI = \exp(7.1081)x^{0.67632}$

where *SI* is the observed average signal intensity for dilution *x*. The gene abundance *G*
was calculated by inverting this equation. For this probe, signal intensity at each
postmortem time, SI_t, is determined by the equation: $G = (SI_t/ \exp(7.1081))^{\wedge}(1/0.67632)$.
Specifically, consider two biological replicates of 15 min postmortem zebrafish, the
signal intensities of the probe A_15_P110618 are 770.5 and 576.0, which translates into
the abundances 0.50 and 0.33 arbitrary units (a.u.) respectively. The target abundances
were further converted to log10 and are shown in external file Fish_log10_AllProfiles.txt.

Details of the calibration protocols to calculate gene expression values, i.e., mRNA
relative abundances, are provided in our recent paper where we describe the "Gene
Meter" [6].

Statistical analysis. Abundance levels were log-transformed for analysis to stabilize the
variance. A one-sided Dunnett's T-statistic was applied to test for increase at one or more
postmortem times compared to live control (fish) or time 0 (mouse). A bootstrap
procedure with 10^9 simulations was used to determine the critical value for the Dunnett
statistics in order to accommodate departures from parametric assumptions and to
account for multiplicity of testing. The profiles for each gene were centered by
subtracting the mean values at each postmortem time point to create "null" profiles.
Bootstrap samples of the null profiles were generated to determine the 95th percentile of
the maximum (over all genes) of the Dunnett statistics. Significant postmortem
upregulated genes were selected as those having Dunnett T values larger than the 95th
percentile. Only significantly upregulated genes were retained for further analyses.

Orthogonal transformation of the abundances to their principal components (PC) was
conducted and the results were graphed on a 2 dimensional ordination plot. The *m* x *n*
matrix of abundances (sampling times by number of gene transcripts), which is 10 x 548
for zebrafish and 7 x 515 for mouse, was used to produce an *m* x *m* matrix D of Euclidean

distances between all pairs of sampling times. Principal component analysis (PCA) was
performed on the matrix of distances, D. To investigate and visualize differences
between the sampling times, a scatterplot of the first two principal components (PC1 and
PC2) was created. To establish relative contributions of the gene transcripts, the
projection of each sampling time onto the (PC1, PC2) plane was calculated and those
genes with high correlations (>=0.70) between abundances and either component (PC1 or
PC2) were displayed as a biplot.

Gene annotation and functional categorization. Microarray probe sequences were
individually annotated by performing a BLASTN search of the zebrafish and mouse
NCBI databases (February, 2015). The gene annotations were retained if the bit score
was greater than or equal to 100 and the annotations were in the correct 5' to 3'
orientation. Transcription factors, transcriptional regulators, and cell signaling
components (e.g., receptors, enzymes, and messengers) were identified as global
regulatory genes. The rest were considered response genes.

Functional categorizations were performed by querying the annotated gene transcripts in
the primary literature and using UniProt (www.uniprot.org). Genes not functionally
categorized to their native organism (zebrafish or mouse) were categorized to genes of
phylogenetically related organisms (e.g., human). Cancer-related genes were identified
using a previously constructed database (see Additional File 1: Table S1 in [7]).

RESULTS

Similar quantities of total mRNA were extracted from zebrafish samples for the first 12 h
postmortem (avg. 1551 ng per µl tissue extract) then the quantities abruptly decreased
with postmortem time (Table S1). The quantities of total mRNA extracted from the
mouse liver samples were about the same for the first 12 h postmortem (avg. of 553 ng
per µl tissue extract) then they increased with time (Table S2). The quantities of total
mRNA extracted from the mouse brain samples were similar (avg. of 287 ng per µl tissue
extract) for all postmortem times (Table S2). Hence, the amount of total mRNA
extracted depended of the organism/organ/tissue and time.

Calibration of the microarray probes and determination of the transcript abundances at
each postmortem sampling time produced a fine-grain series of data for the zebrafish and
the mouse. Approximately 84.3% (36,811 of 43,663) zebrafish probes and 67.1%
(37,368 of 55,681) mouse probes were found to provide a suitable dose-response curve
for calibration (data available upon request).

Figure 1 shows the sum of all gene abundances calculated from the calibrated probes with
postmortem time. In general, the sum of all abundances decreased with time, which
means that less targets hybridized to the microarray probes. In the zebrafish, mRNA
decreased abruptly at 12 h postmortem (Fig 1A), while for the mouse brain Fig 1B), total
mRNA increased in the first hour and then gradually decreased. For the mouse liver,
mRNA gradually decreased with postmortem time. The fact that total mRNA shown in
Figs 1A and 1B mirrors the electrophoresis banding patterns shown in Fig S1 and S2
(ignoring the 28S and 18S rRNA bands) indicates a general agreement of the Gene Meter

approach to another molecular approach (i.e., Agilent Bioanalyzer). Hence, total mRNA
abundances depended on the organism (zebrafish, mouse), organ (brain, liver), and
postmortem time, which are aligned with previous studies [8,9,10,11,12,13].

238
Fig 1. Total mRNA abundance (arbitrary units, a.u.) by postmortem time
determined using all calibrated microarray probes. A, extracted from whole
zebrafish; B, extracted from brain and liver tissues of whole mice. Each datum
point represents the mRNA from two organisms in the zebrafish and a single
organism in the mouse.

The abundance of a gene transcript is determined by its rate of synthesis and its rate of
degradation [14]. We focused on genes that show a significant increase in RNA
abundance -- relative to live controls -- because these genes are likely to be actively
transcribed in organismal death despite an overall decrease in total mRNA with time. An
upregulated transcription profile was defined as one having at least one time point where
the abundance was statistically higher than that of the control (Fig 2 A to 2C). It is
important to understand that the entire profiles, i.e., 22 data points for the zebrafish and
20 points for the mouse, were subjected to a statistical test to determine significance (see
Materials and Methods). We found 548 zebrafish transcriptional profiles and 515 mouse
profiles were significantly upregulated. The fact that there are upregulated genes is
consistent with the notion that there is still sufficient energy and functional cellular
machinery for transcription to occur -- long after organismal death.

Fig 2. Transcriptional profiles of representative genes (a.u.), ordination plots based on transcript abundances by postmortem time (h) with corresponding transcript contributions (biplots), and averaged transcript abundances by group. (A) Transcriptional profile of the *Hsp70.3* gene, (B) the *Tox2* gene, and (C) the NULL (i.e., no annotation, probe number shown) gene as a function of postmortem time. Each datum point was derived from the mRNA of two zebrafish or one mouse. (D) Ordination plots of the zebrafish and (E) mouse were based on all upregulated gene profiles by postmortem time (h). Gene transcripts in the biplots were arbitrarily assigned alphabetical groups based on their positions in the ordination. The average transcript abundances for each group are shown.

Based on GenBank gene annotations, we found that among the upregulated genes for the zebrafish, 291 were protein-coding genes (53%) and 257 non-coding mRNA (47%) and, for the mouse, 324 known protein-coding genes (63%), 190 non-coding mRNA (37%), and one control sequence of unknown composition. Hence, about 58% of the total upregulated genes in the zebrafish and mouse are known and the rest (42%) are non-coding RNA.

Examples of genes yielding transcripts that significantly increased in abundance with postmortem time are: the 'Heat shock protein' (*Hsp70.3)* gene, the 'Thymocyte selection-associated high mobility group box 2' (*Tox2*) gene, and an unknown (*NULL*) gene (Fig

2A to 2C). While the *Hsp70.3* transcript increased after 1 h postmortem to reach a
maximum at 12 h, the *Tox2* transcript increased after 12 h postmortem to reach a
maximum at 24 h, and the *NULL* transcript consistently increased with postmortem time.
These figures provide typical examples of transcript profiles and depict the high
reproducibility of the sample replicates as well as the quality of output obtained using the
Gene Meter approach.

Non-random patterns in transcript profiles

Ordination plots of the significantly upregulated transcript profiles revealed prominent
differences with postmortem time (Fig 2D and 2E), suggesting the expression of genes
followed a discernible (non-random) pattern in both organisms. The biplots showed that
203 zebrafish transcript profiles and 226 mouse profiles significantly contributed to the
ordinations. To identify patterns in the transcript profiles, we assigned them to groups
based on their position in the biplots. Six profile groups were assigned for the zebrafish
(A to F) and five groups (G to K) were assigned for the mouse. Determination of the
average gene transcript abundances by group revealed differences in the shapes of the
averaged profiles, particularly the timing and magnitude of peak transcript abundances,
which accounted for the positioning of data points in the ordinations.

Genes coding for global regulatory functions were examined separately from others (i.e.,
response genes). Combined results show that about 33% of the upregulated genes in the
ordination plots were involved in global regulation with 14% of these encoding
transcription factors/transcriptional regulators and 19% encoding cell signaling proteins
such as enzymes, messengers, and receptors (Table S3). The response genes accounted
for 67% of the upregulated transcripts.

The genes were assigned to 22 categories (File S8) with some genes having multiple
categorizations. For example, the Eukaryotic Translation Initiation Factor 3 Subunit J-B
(*Eif3j2*) gene was assigned to protein synthesis and cancer categories [15].

Genes in the following functional categories were investigated: stress, immunity,
inflammation, apoptosis, solute/ion/protein transport, embryonic development, epigenetic
regulation and cancer. We focused on these categories because they were common to
both organisms and contained multiple genes, and they might provide explanations for
postmortem upregulation of genes (e.g., epigenetic gene regulation, embryonic
development, cancer). The transcriptional profiles of the genes were plotted by category
and each profile was ordered by the timing of the upregulation and peak transcript
abundance. This allowed comparisons of gene expression dynamics as a function of
postmortem time for both organisms. For each category, we provided the gene name and
function and compared expression dynamics within and between the organisms.

Stress response

In organismal death, we anticipated the upregulation of stress response genes because
these genes are activated in life to cope with perturbations and to recover homeostasis
[16]. The stress response genes were assigned to three groups: heat shock protein (*Hsp*),
hypoxia-related, and 'other' responses such as oxidative stress.

Hsp In the zebrafish, upregulated *Hsp* genes included: 'Translocated promoter region'
(*Tpr*), *Hsp70.3,* and *Hsp90* (Fig 3). The *Tpr* gene encodes a protein that facilitates the
export of the *Hsp* mRNA across the nuclear membrane [17] and has been implicated in
chromatin organization, regulation of transcription, mitosis [18] and controlling cellular
senescence [19]. The *Hsp70.3* and *Hsp90* genes encode proteins that control the level of
intracellular calcium [20], assist with protein folding, and aid in protein degradation [21].

In the mouse, upregulated *Hsp* genes included: *Tpr*, *Hsp*-associated methyltransferase
(*Mettl21*) and Heat Shock Protein 1 (*Hspe1*) (Fig 3). The *Mettl21* gene encodes a protein
modulating *Hsp* functions [22]. The *Hspe1* gene encodes a chaperonin protein that
assists with protein folding in the mitochondria [23].

The timing and duration of *Hsp* upregulation varied by organism. In general, activation
of *Hsp* genes occurred much later in the zebrafish than the mouse (4 h vs. 0.5 h
postmortem, respectively). There were also differences in transcript abundance maxima
since, in the zebrafish, maxima were reached at 9 to 24 h, while in the mouse maxima
were reached at 12 to 24 h. Previous studies have examined the upregulation of *Hsp70.3*
with time in live serum-stimulated human cell lines [24]. In both the zebrafish and
human cell lines the *Hsp70.3* gene transcript reached maximum abundance at about 12 h
(Fig 2A), indicating the same reactions are occurring in life and death.

Fig 3. Upregulated stress response genes by postmortem time (h) and stress
category. A. zebrafish; B, mouse. Green, intermediate value; Red, maximum value.

Hypoxia
In the zebrafish, upregulated hypoxia-related genes included: Carbonic anhydrase 4
(*Ca4c*), Nuclear factor interleukin-3 (*Nfil3*), Hypoxia-inducible factor 1-alpha (*Hiflab*)
and Arginase-2 (*Arg2*) (Fig 3). The Carbonic anhydrase 4 (*Ca4c*) gene encodes an
enzyme that converts carbon dioxide into bicarbonate in response to anoxic conditions
[25]. The *Nfil3* gene encodes a protein that suppresses hypoxia-induced apoptosis [26]
and activates immune responses [27]. The *Hiflab* gene encodes a transcription factor that
prepares cells for decreased oxygen [28]. The *Arg2* gene encodes an enzyme that
catalyzes the conversion of arginine to urea under hypoxic conditions [29]. Of note, the
accumulation of urea presumably triggered the upregulation of the *Slc14a2* gene at 24 h,
reported in the Transport Section (below).

In the mouse, upregulated hypoxia genes included: Methyltransferase hypoxia inducible
domain (*Methig1*) and Sphingolipid delta-desaturase (*Degs2*) (Fig 3). The *Methig1* gene
encodes methyltransferase that presumably is involved in gene regulation [30]. The
Degs2 gene encodes a protein that acts as an oxygen sensor and regulates ceramide
metabolism [31]. Ceramides are waxy lipid molecules in cellular membranes that
regulate cell growth, death, senescence, adhesion, migration, inflammation, angiogenesis
and intracellular trafficking [32].

The activation of the *Ca4c* gene in the zebrafish indicates a build up of carbon dioxide at
0.1 to 1 h postmortem in the zebrafish presumably due lack of blood circulation. The
upregulation of the *Nfil3* gene in the zebrafish and *Methig1* gene in the mouse suggests
hypoxic conditions existed within 0.5 h postmortem in both organisms. The upregulation
of other hypoxia genes varied with postmortem time, with the upregulation of *Hif1ab,*
Arg2, and *Degs2* genes occurring at 4 h, 12 h and 24, respectively.

Other stress responses
In the zebrafish, upregulated response genes included: Alkaline ceramidase 3 (*Acer3*),
Peroxirodoxin 2 (*Prdx2*), Immediate early (*Ier2*), Growth arrest and DNA-damage-
inducible protein (*Gadd45a*), Zinc finger CCH domain containing 12 (*Zcchc12*),
Corticotropin releasing hormone receptor 1 (*Crhr1*), and Zinc finger AN1-type domain 4
(*Zfand4*) (Fig 3). The *Acer3* gene encodes a stress sensor protein that mediates cell-
growth arrest and apoptosis [33]. The *Prdx2* gene encodes an antioxidant enzyme that
controls peroxide levels in cells [34] and triggers production of *Tnfa* proteins that induce
inflammation [35]. The *Ier2* gene encodes a transcription factor involved in stress
response [36]. The *Gadd45a* gene encodes a stress protein sensor that stops the cell cycle
[37], modulates cell death and survival, and is part of the signaling networks in immune
cells [38]. The *Zcchc12* gene encodes a protein involved in stress response in the brain
[39]. The *Crhr1* and *Zfand4* genes encode stress proteins [40,41].

While the *Acer3, Prdx2,* and *Ier2* genes were activated within 0.3 h postmortem,
indicating a changed physiological state; the *Gadd45a* gene was activated at 9 h and the
other genes (*Zcchc12, Crhr1, Zfand4*) were activated at 24 h postmortem.

In the mouse, upregulated stress response genes included: Membrane-associated RING-
CH 4 (*March4*), Homocysteine-responsive endoplasmic reticulum-resident ubiquitin-like
domain member 2 (*Herpud2*), Prohibitin-2 (*Phb2*), *Gadd45a*, and Two-oxoglutarate and
iron-dependent oxygenase domain-containing 1 (*Ogfod1*) (Fig 3). The *March4* gene
encodes an immunologically-active stress response protein [42]. The *Herpud2* gene
encodes a protein that senses the accumulation of unfolded proteins in the endoplasmic
reticulum [43]. The *Phb2* gene encodes a cell surface receptor that responds to
mitochondrial stress [44]. The *Ogfod1* gene encodes a stress-sensing protein [45].

Note that the stress genes in the mouse were all activated within 1 h postmortem and
remained upregulated for 48 h.

Summary of stress response
In both organisms, organismal death activated heat shock, hypoxia, and other stress
genes, which varied in the timing and duration of upregulation within and between
organisms. Consider, for example, the *Tpr* and *Gadd45a* genes, which were common to
both organisms. While the *Tpr* genes were upregulated within 0.5 h postmortem in both

organisms, the *Gadd45a* gene was upregulated at 9 h in the zebrafish and 0.5 h in the
mouse. In addition, the transcription profile of the *Tpr* gene was more variable in the
zebrafish than the mouse since it was activated at 0.3 h, 9 h and 24 h postmortem, which
suggest the gene might be regulated through a feedback loop. In contrast, in the mouse,
the *Tpr* gene was upregulated at 0.5 h and the transcripts reached peak abundance at 12
and 24 h postmortem.

Taken together, the stress genes were activated in both organisms to compensate for a
loss in homeostasis.

Innate and adaptive immune responses

In organismal death, we anticipated the upregulation of immune response genes because
vertebrates have evolved ways to protect the host against infection in life, even under
absolutely sterile conditions [46]. Inflammation genes were excluded from this section
(even though they are innate immune genes) because we examined them in a separate
section (below).

In the zebrafish, upregulated immunity genes included: Early growth response-1 and -2
(*Egr1, Egr2*), Interleukin-1b (*Il1b*), L-amino acid oxidase (*Laao*), Interleukin-17c (*Il17c*),
Membrane-spanning 4-domains subfamily A member 17A.1 (*Ms4a17.a1*), Mucin-2
(*Muc2*), Immunoresponsive gene 1 (*Irg1*), Interleukin- 22 (*Il22*), Ubl carboxyl-terminal
hydrolase 18 (*Usp18*), ATF-like 3 (*Batf3*), Cytochrome b-245 light chain (*Cyba*), and
'Thymocyte selection-associated high mobility group" box protein family member 2
(*Tox2*) (Fig 4). The *Egr1* and *Egr2* genes encode proteins that regulate B and T cell
functions in adaptive immunity [47,48]. The *Il1b* gene encodes an interleukin that kills
bacterial cells through the recruitment of other antimicrobial molecules [49]. The *Laao*
gene encodes an oxidase involved in innate immunity [50]. The *Il17c* and *Il22* genes
encode interleukins that work synergically to produce antibacterial peptides [51]. The
Ms4a17.a1 gene encodes a protein involved in adaptive immunity [52]. The *Muc2* gene
encodes a protein that protects the intestinal epithelium from pathogenic bacteria [53].
The *Irg1* gene encodes an enzyme that produces itaconic acid, which has antimicrobial
properties [54]. The *Usp18* gene encodes a protease that plays a role in adaptive
immunity [55]. The *Batf3* gene encodes a transcription factor that activates genes
involved in adaptive immunity [56]. The *Cyba* gene encodes an oxidase that is used to
kill microorganisms [57]. The *Tox2* gene encodes a transcription factor that regulates
Natural Killer (NK) cells of the innate immune system [58].

Fig 4. Upregulated immunity genes by postmortem time (h). A. zebrafish; B, mouse. Green, intermediate value; Red, maximum value. Some transcripts were represented by two different probes (e.g. *Il1b, Laao*).

Upregulation of immunity genes in the zebrafish occurred at different times with varying durations. While genes involved in adaptive immunity were upregulated at 0.1 to 0.3 h (*Egr*), 9 h (*Ms4a17.a1*) and 24 h (*Usp18, Batf3*) postmortem, genes involved in innate immunity were upregulated at 4 h (*Il1b*), 9 h (*Laao, Il17c*), 12 h (*Muc2, Irg1*) and 24 h (*Il22, Cyba, Tox2*) indicating a multi-pronged and progressive approach to deal with injury and the potential of microbial invasion.

In the mouse, upregulated antimicrobial genes included: Catalytic polypeptide-like 3G (*Apobec3g*), CRISPR-associated endonuclease (*Cas1*), Perforin-1 (*Prf1*), Immunoglobulin heavy variable 8-11 (*Ighv8-11*), C4b-binding protein (*C4b*), Complement component C7 (*C7*), T cell receptor alpha and delta chain (*Tcra/Tcrd*), High affinity immunoglobulin gamma Fc receptor I (*Fcgr1a*), Defensin (*Defb30*), Chemokine-4 (*Ccr4*), Interleukin-5 (*Il5*), NK cell receptor 2B4 (*Cd244*), Cluster of differentiation-22 (*Cd22*), Lymphocyte cytosolic protein 2 (*Lcp2*), Histocompatibility 2 O region beta locus (*H2ob*) and Interferon-induced transmembrane protein 1 (*Ifitm1*) (Fig 4). The *Apobec3g* gene encodes a protein that plays a role in innate anti-viral immunity [59]. The *Cas1* gene encodes a protein involved in regulating the activation of immune systems [60,61,62,63]. The *Prf1, C7,* and *Defb30* genes encode proteins that kill bacteria by forming pores in the plasma membrane of target cells [64,65,66]. The *Ighv8-11* gene encodes an immunoglobulin of uncertain function. The *C4b* gene encodes a protein involved in the complement system [67]. The *Tcra/Tcrd* genes encode proteins that play a role in the immune response [68]. The *Fcgr1a* gene encodes a protein involved in both innate and adaptive immune responses [69]. The *Ccr4* gene encodes a cytokine that attracts leukocytes to sites of infection [70]. The *Il5* gene encodes an interleukin involved in both innate and adaptive immunity [71,72]. The *Cd244* and *Cd22* genes encode proteins involved in innate immunity [73]. The *Lcp2* gene encodes a signal-transducing adaptor protein involved in T cell development and activation [74]. The

H2ob gene encodes a protein involved in adaptive immunity. The *Ifitm1* gene encodes a

protein that has antiviral properties [75].

Most immune response genes were upregulated within 1 h postmortem in the mouse

(*n*=14 out of 16 genes), indicating a more rapid response than that of the zebrafish.

Summary of immune response

The upregulated immune response genes in both organisms included innate and adaptive

immunity components. An interesting phenomenon observed in the mouse (but not

zebrafish) was that four genes (*C7, Tcra/Tcrd, Fcgr1a,* and *Defb30*) reached maximum

transcript abundance at two different postmortem times (i.e., 1 h and 12 h) while others

reached only one maximum. The variability in the gene transcript profiles suggests their

regulation is effected by feedback loops.

Inflammation response

We would anticipate the upregulation of inflammation genes in organismal death because

inflammation is an innate immunity response to injury. In the zebrafish, upregulated

inflammation genes included: *Egr1, Egr2, Il1b,* Tumor necrosis factor receptor

(*Tnfrsf19*), Heme oxygenase 1 (*Hmox1*), Tumor necrosis factor (*Tnf*), G-protein receptor

(*Gpr31*), Interleukin-8 (*Il8*), Tumor necrosis factor alpha (*Tnfa*), Nuclear factor (NF)

kappa B (*Nfkbiaa*), MAP kinase-interacting serine/threonine kinase 2b (*Mknk2b*), and

Corticotropin-releasing factor receptor 1 (*Crhr1*) (Fig 5). The *Egr1* and *Egr2* genes

encode transcription factors that are pro- and anti- inflammatory, respectively [47,48,76].

The *Il1b* gene encodes a pro-inflammatory cytokine that plays a key role in sterile

inflammation [77,78]. The *Tnfrsf19* gene encodes a receptor that has pro-inflammatory

functions [79]. The *Hmox1* gene encodes an enzyme that has anti-inflammatory

functions and is involved in Heme catabolism [80,81]. The *Tnf* and *Tnfa* genes encode

pro-inflammatory proteins. The *Gpr31* gene encodes a pro-inflammatory protein that

activates the NF-kB signaling pathway [82]. The *Il8* gene encodes a cytokine that has

pro-inflammatory properties [83]. The *Nfkbiaa* gene encodes a protein that integrates

multiple inflammatory signaling pathways including *Tnf* genes [84]. The *Mknk2b* gene

encodes a protein kinase that directs cellular responses and is pro-inflammatory [85].

The *Crhr1* gene modulates anti-inflammatory responses [86].

Fig 5. Upregulated inflammation genes by postmortem time (h). A. zebrafish; B, mouse. Inflammation, Pro-, + ; Anti, - . Green, intermediate value; Red, maximum value. The *Il1b* and *Mknk2b* genes were represented by two different probes.

The upregulation of pro-inflammatory *Egr1* gene at 0.1 h was followed by the upregulation of anti-inflammatory *Egr2* gene at 0.2 h, suggesting the transcription of one gene is affecting the regulation of another (Fig 5). Similarly, the upregulation of the pro-inflammatory *Il1b* gene at 4 h postmortem was followed by: upregulation of pro-inflammatory *Tnfrsf19, Tnf, Gpr31* and *Il8* genes and anti-inflammatory *Hmox1* gene at 9 h, the upregulation of pro-inflammatory *Tnfa, Nfkbiaa,* and *Mknk2b* genes at 12 h, and the upregulation of anti-inflammatory *Crhr1* gene at 24 h. Of note, while none of the pro-inflammatory genes were upregulated past 48 h, the anti-inflammatory *Crhr1* gene remained upregulated at 48 h. It should also be noted that the *Il1b, Il8,* and *Tnfa* genes have been reported to be upregulated in traumatic impact injuries in postmortem tissues from human brains [87].

In the mouse, upregulated inflammation genes included: mitogen-activated protein kinase (*Map3k2*), TNF receptors (*Tnfrsf9, Tnfrs14*), B-cell lymphoma 6 protein (*Bcl6*), C-C chemokine receptor type 4 (*Ccr4*), Prokineticin-2 (*Prok2*), and platelet-activating factor receptor (*Pafr*) (Fig 5). The *Map3k2* gene encodes a kinase that activates pro-inflammatory NF-kB genes [85]. The *Tnfrsf9* and *Tnfrs14* genes encode receptor proteins that have pro-inflammatory functions [79]. The *Bcl6* gene encodes a transcription factor that has anti-inflammatory functions [88]. The *Ccr4* gene encodes a cytokine receptor protein associated with inflammation [70]. The *Prok2 gene* encodes a cytokine-like molecule, while the *Pafr* gene encodes a lipid mediator; both have pro-inflammatory functions [89,90].

Most inflammation-associated genes were upregulated within 1 h postmortem and continued to be upregulated for 12 to 48 h. The anti-inflammatory *Bcl6* gene was upregulated at two different times: 0.5 to 6 h and at 24 h suggesting that it is presumably being regulated by a feedback loop. It should also be noted that pro-inflammatory *Map3k2* and *Tnfrs14* genes were not upregulated after 24 and 12 h, respectively, which also suggests regulation by a putative feedback loop from the *Bcl6* gene product.

Summary of inflammation response
In both organisms, some of the upregulated genes have pro-inflammatory functions while others have anti-inflammatory functions, presumably reflecting regulation by feedback loops. The putative feedback loops involve an initial inflammatory reaction followed by an anti-inflammatory reaction to repress it [91]. The variation in the upregulation of these inflammatory genes suggests an underlying regulatory network is involved in organismal death.

Apoptosis and related genes
Since apoptotic processes kill damaged cells for the benefit of the organism as a whole, we anticipated the upregulation of apoptosis genes in organismal death.

In the zebrafish, upregulated apoptosis genes included: Jun (*Jdp2, Jun*), Alkaline
ceramidase 3 (*Acer3*)*, Fos* (*Fosb, Fosab, Fosl1*), IAP-binding mitochondrial protein A
(*Diabloa*), Peroxiredoxin-2 (*Prdx2*), Potassium voltage-gated channel member 1
(*Kcnb1*), Caspase apoptosis-related cysteine peptidase 3b (*Casp3b*), DNA-damage-
inducible transcript 3 (*Ddit3*), BCL2 (B-cell lymphomas 2)-interacting killer (*Bik*), and
Ras association domain family 6 (*Rassf6*) (Fig 6). The *Jdp2* gene encodes a protein that
represses the activity of the transcription factor activator protein 1 (*AP-1*) [92]. The
Acer3 gene encodes an enzyme that maintains cell membrane integrity/function and
promotes apoptosis [93]. The *Fos* genes encode proteins that dimerize with *Jun* proteins
to form part of the *AP-1* that promotes apoptosis [94,95]. The *Diabloa* gene encodes a
protein that neutralizes inhibitors of apoptosis (IAP)-binding protein [95] and activates
caspases [96]. The *Prdx2* gene encodes antioxidant enzymes that control cytokine-
induced peroxide levels and inhibit apoptosis [97]. Although the *Kcnb1* gene encodes a
protein used to make ion channels, the accumulation of these proteins in the membrane
promotes apoptosis via cell signaling pathway [98]. The *Casp3b* encodes a protein that
plays a role in the execution phase of apoptosis [99]. The *Ddit3* gene encodes a
transcription factor that promotes apoptosis. The *Bik* gene encodes a protein that
promotes apoptosis [100]. The *Rassf6* gene encodes a protein that promotes apoptosis
[101].

Fig 6. Upregulated apoptosis genes by postmortem time (h). A, zebrafish; B, mouse.
Apopotosis, Pro, + ; Anti, - . Green, intermediate value; Red, maximum value. The
***Fosb* gene was represented by two different probes.**

In the zebrafish, both anti-apoptosis *Jdp2* and pro-apopotosis *Acer3* genes were
upregulated within 0.1 h postmortem (Fig 6). The upregulation of these genes was
followed by the upregulation of five pro-apoptosis genes and one anti-apoptosis gene
within 0.3 to 0.5 h. The transcriptional dynamics varied among the genes. Specifically,
(i) the *Fosb* transcript was at low abundance after 1 h, (ii) the *Diabloa* and *Fosab*
transcripts reached abundance maxima at 0.5 to 4 h and then were at low abundance after
9 h for the *Diabloa* and after 24 h for the *Fosab*, (iii) the *Jun* transcript reached two
maxima (one at 0.5 and another at 4 to 12 h) – and after 24 h was at low abundance, (iv)
the *Prdx2* transcript showed a continuous increase in abundance, reaching a maximum at
24 h and then was at low abundance. The remaining genes were pro-apoptosis and
upregulated after 1 to 24 h postmortem. The *Ddit3* and *Rassf6* genes were very different

from the other genes in that they were upregulated at one sampling time (12 h and 24 h,
respectively) and then the transcripts were at low abundance. Apparently the apoptosis
transcripts were at low abundance after 24 h in contrast to the transcripts in other
categories (e.g. stress and immunity genes were upregulated for 96 h postmortem).

In the mouse, upregulated apoptosis-associated genes included: BCL2-like protein 11
(*Bcl2L11*), Casein kinase IIa (*Csnk2a1*), Interleukin 15 receptor subunit a (*Il15ra*),
Myocyte enhancer factor 2 (Mef2a), F-box only protein 10 (*Fbxo10*), Sp110 nuclear body
protein (*Sp110*), TGFB-induced factor homeobox 1 (*Tgif1*), Intersectin 1 (*Itsm1*) gene,
the Ephrin type-B receptor 3 (*Ephb3*), and the p21 protein-activated kinase 4 (*Pak4*) (Fig
6). The *Bcl2L11* gene encodes a protein that promotes apoptosis [102]. The *Csnk2a1*
gene encodes an enzyme that phosphorylates substrates and promotes apoptosis [103].
The *Il15ra* gene encodes an anti-apoptotic protein [104]. The *Mef2a* gene encodes a
transcription factor that prevents apoptosis [105]. The *Fbxo10* gene encodes a protein
that promotes apoptosis [106]. The *Sp110* gene encodes a regulator protein that promotes
apoptosis [107]. The *Tgif1* gene encodes a transcription factor that blocks signals of the
transforming growth factor beta (*TGFβ*) pathway; and therefore, is pro-apoptosis [108].
The *Itsn1* gene encodes an adaptor protein that is anti-apoptosis [109]. The *Ephb3* gene
encodes a protein that binds ligands on adjacent cells for cell signaling and suppresses
apoptosis [104]. The *Pak4* gene encodes a protein that delays the onset of apoptosis
[110].

In the mouse, pro- and anti-apoptosis genes were upregulated within 0.5 h postmortem –
however, with exception to *Bcl2L11*, most reached transcript abundance maxima at 12 to
48 h postmortem (Fig 6). The *Bcl2L11* transcripts reached abundance maxima at 1 and 6
h postmortem.

Summary of apoptotic response
In both organisms, pro- and anti-apoptosis genes were upregulated in organismal death.
However, the timings of the upregulation, the transcript abundance maximum, and the
duration of the upregulation varied by organism. The results suggest the apoptotic genes
and their regulation are distinctly different in the zebrafish and the mouse, with the
mouse genes being upregulated up to 48 h postmortem, while zebrafish genes are
upregulated for 24 h. Nonetheless, the pro- and anti-apoptosis genes appear to be inter-
regulating each another.

Transport gene response

Transport processes maintain ion/solute/protein homeostasis and are involved in
influx/efflux of carbohydrates, proteins, signaling molecules, and nucleic acids across
membranes. We anticipated that transport genes should be upregulated in organismal
death in response to dysbiosis.

In the zebrafish, upregulated transport-associated genes included: Solute carrier family 26
Anion Exchanger Member 4 (*Slc26a4*), Potassium channel voltage gated subfamily H
(*Kcnh2*), Transmembrane Emp24 domain containing protein 10 (*Tmed10*), Leucine rich
repeat containing 59 (*Lrrc59*), the Nucleoprotein TPR (*Tpr*), Importin subunit beta-1
(*Kpnb1*), Transportin 1 (*Tnpo1*), Syntaxin 10 (*Stx10*) and Urea transporter 2 (*Slc14a2*)
(Fig 7). Of note, the four *Tmed10* transcripts shown in Fig 7, each represents a profile

targeted by an independent probe. The transcription profiles of this gene were identical,
indicating the high reproducibility of the Gene Meter approach. The *Slc26a4* gene
encodes prendrin that transports negatively-charged ions (i.e., Cl⁻, bicarbonate) across
cellular membranes [111]. The *Kcnh2* gene encodes a protein used to make potassium
channels and is involved in signaling [112]. The *Tmed10* gene encodes a membrane
protein involved in vesicular protein trafficking [113]. The *Lrrc59, Tpr, Tnpo1,* and
Kpnb1 genes encode proteins involved in trafficking across nuclear pores
[114,115,116,117]. The *Stx10* gene encodes a protein that facilitates vesicle fusion and
intracellular trafficking of proteins to other cellular components [118]. The *Slc14a2* gene
encodes a protein that transports urea out of the cell [119].

Fig 7. Upregulated transport genes by postmortem time (h) and stress category. A.
zebrafish; B, mouse. Green, intermediate value; Red, maximum value. The *Tmed10*
gene was represented by four different probes.

The *Slc26a4, Kcnh2, Lrrc59* and *Tpr* genes were initially upregulated within 0.3 h
postmortem and continue to be expressed for 12 to 24 h. The *Tnpo1* gene was
upregulated at two times: 4 h and 12 h suggesting putative regulation by a feedback loop.
The remaining genes were upregulated at 24 h. The upregulation of the *Slc14a2* gene
suggests a build up of urea in zebrafish cells at 24 to 96 h postmortem, which could be
due to the accumulation of urea under hypoxic conditions by the *Arg2* gene (see *Hsp*
stress response section).

In the mouse, the following transport-associated genes were upregulated: Calcium-bind
mitochondrial carrier protein (*Aralar2*), Sodium-coupled neutral amino acid transporter 4
(*Slc38a4*), SFT2 domain containing 1 (*Sft2d1*), Uap56-interacting factor (*Fyttd1*), Solute
carrier family 5 (sodium/glucose co-transporter) member 10 (*Slc5a10*), Mitochondrial
import receptor subunit (*Tom5*), 'Translocated promoter region' (*Tpr*), ATP-binding
cassette transporter 12 (*Abca12*), Multidrug resistant protein 5 (*Abc5*), LIM and SH3
domain-containing protein (*Lasp1*), Chromosome 16 open reading frame 62 (*C16orf62*),
Golgi transport 1 homolog A (*Golt1a*), ATP-binding cassette transporter transporter 17

(*Abca17*), Nucleotide exchange factor (*Sil1*), Translocase of inner mitochondrial
membrane 8A1 (*Timm8a1*), Early endosome antigen 1 (*Eea1*), and Potassium voltage-
gated channel subfamily V member2 (*Kcnv2*) (Fig 7). The *Aralar2* gene encodes a
protein that catalyzes calcium-dependent exchange of cytoplasmic glutamate with
mitochondrial aspartate across the mitochondrial membrane and may function in the urea
cycle [120]. The *Slc38a4* gene encodes a symport that mediates transport of neutral
amino acids and sodium ions [121]. The *Sft2d1* gene encodes a protein involved in
transporting vesicles from the endocytic compartment of the Golgi complex [122]. The
Fyttd1 gene is responsible for mRNA export from the nucleus to the cytoplasm [123].
The *Slc5a10* gene encodes a protein that catalyzes carbohydrate transport across cellular
membranes [124]. The *Tom5* gene encodes a protein that plays a role in importation to
proteins destined to mitochondrial sub-compartments [125]. The *Abca12*, *Abca17* and
Abc5 genes encode proteins that transport molecules across membranes [126,127,128].
The *Lasp1* gene encodes a protein that regulates ion-transport [129]. The *C16orf62* gene
encodes a protein involved in protein transport from the Golgi apparatus to the cytoplasm
[130]. The *Golt1a* gene encodes a vesicle transport protein [122]. The *Sil1* gene encodes
a protein involved in protein translocation into the endoplasmic reticulum [131]. The
Timm8a1 gene encodes a protein that assists importation of other proteins across inner
mitochondrial membranes [132]. The *Eea1* gene encodes a protein that acts as a
tethering molecule for vesicular transport from the plasma membrane to the early
endosomes [133]. The *Kcnv2* gene encodes a membrane protein involved in generating
action potentials [134].

Within 0.5 h postmortem, genes involved in: (i) ion and urea regulation (*Aralar*), (ii)
amino acid (*Slc38a4*), carbohydrate (*Slc5a10*), and protein (*Sft2d1, Tom5*) transport, (iii)
mRNA nuclear export (*Fyttd1, Tpr*), and (iv) molecular efflux (*Abca12, Abc5*) were
upregulated in the mouse. The transcription profiles of these genes varied in terms of
transcript abundance maxima and duration of the upregulation. While the transcripts of
Aralar, Sft2d1, Slc38a4, Fyttd1 and *Slc5a10* reached abundance maxima at 1 h, those of
Tom5, Tpr, Abca12, and *Abc5* reached maxima at 12 to 24 h postmortem. The duration
of the upregulation also varied for these genes since most were upregulated for 48 h
postmortem, while the *Sft2d1, Fyttd1* and *Slc5a10* were upregulated from 0.5 to 12 h or
more. The shorter duration of upregulation suggests prompt gene repression. The
transcript abundance of *Lasp1, C16orf62, Golt1a,* and *Abca17* increased at 1 h
postmortem and remained elevated for 48 h. The transcripts of *Sil1, Timm8a1, Eea1*
increased in abundance at 6 h, while those of *Kcnv2* increased at 24 h postmortem and
remained elevated for 48 h.

Summary of transport genes
The upregulation of transport genes suggests attempts by zebrafish and mice to
reestablish homeostasis. Although half of these genes were upregulated within 0.5 h
postmortem, many were upregulated at different times and for varying durations. While
most of the transport gene transcripts in the zebrafish were at low abundance after 24 h,
most transport gene transcripts in the mouse remained at high abundance at 24 to 48 h
postmortem.

Development genes

An unexpected finding in this study was the discovery of upregulated development genes in organismal death. Most development genes are involved at the early stage of development in the zebrafish and mouse; therefore, we did not anticipate their upregulation in organismal death.

In the zebrafish, upregulated development genes included: LIM domain containing protein 2 (*Limd2*), Disheveled-associated activator of morphogenesis 1 (*Daam1b*), Meltrin alpha (*Adam12*), Hatching enzyme 1a (*He1a*), Midnolin (*Midn*), Immediate early response 2 (*Ier2*), Claudin b (*Cldnb*), Regulator of G-protein signaling 4-like (*Rgs4*), Proline-rich transmembrane protein 4 (*Prrt4*), Inhibin (*Inhbaa*), Wnt inhibitory factor 1 precursor (*Wif1*), Opioid growth factor receptor (*Ogfr*), Strawberry notch homolog 2 (*Sbno2*), and Developing brain homeobox 2 (*Dbx2*) (Fig 8). The *Limd2* gene encodes a binding protein that plays a role in zebrafish embryogenesis [135]. The *Daam1b* gene regulates endocytosis during notochord development [136]. The *Adam12* gene encodes a metalloprotease-disintegrin involved in myogenesis [137]. The *He1a* gene encodes a protein involved in egg envelope digestion [138]. The *Midn* gene encodes a nucleolar protein expressed in the brain and is involved in the regulation of neurogenesis [139,140]. The *Ier2* gene encodes a protein involved in left-right asymmetry patterning in the zebrafish embryos [141]. The *Cldnb* gene encodes a tight junction protein in larval zebrafish [142]. The *Rgs4* gene encodes a protein involved in brain development [143]. The *Prrt4* gene encodes a protein that is predominantly expressed in the brain and spinal cord in embryonic and postnatal stages of development. The *Inhbaa* gene encodes a protein that plays a role in oocyte maturation [144]. The *Wif1* gene encodes a WNT inhibitory factor that controls embryonic development [145]. The *Ogfr* gene plays a role in embryonic development [146]. The *Sbno2* gene plays a role in zebrafish embryogenesis [147]. The *Dbx2* gene encodes a transcription factor that plays a role in spinal cord development [148].

Fig 8. Upregulated development genes by postmortem time (h) and stress category.
A. zebrafish; B, mouse. Green, intermediate value; Red, maximum value. The
Cldnb **gene was represented by two different probes.**

Although the abundances of *Limd2, Daam1, Adam12, and He1a* transcripts increased in
the zebrafish within 0.1 h postmortem, other gene transcripts in this category increased
from 0.3 to 24 h postmortem reaching abundance maxima at 24 h or more.

In the mouse, development genes were upregulated included: MDS1 and EVI1 complex
locus protein EVI1 (*Mecom*), MAM domain containing glycosylphosphatidylinositol
anchor 2 (*Mdga2*), FYVE, RhoGEF and PH domain containing 5 (*Fgd5*), RNA binding
motif protein 19 (*Rbm19*), Chicken ovalbumin upstream promoter (*Coup*), Single minded
homolog 2 (*Sim2*), Solute carrier family 38, member 4 (*Slc38a4*), B-cell lymphoma 6
protein (*Bcl6*), Sema domain transmembrane domain (TM) cytoplasmic domain
(semaphorin) 6D (*Sema6d*), RNA binding motif protein 45(*Rbm45*), Transcription factor
E2F4 (*E2f4*), Long chain fatty acid- CoA Ligase 4 (*Lacs4*), Kallikrein 1-related peptidase
b3 (*Klk1b3*), Sema domain, Immunoglobulin domain, transmembrane domain and short
cytoplasmic domain (*Sema4c*), TGFB-induced factor homeobox 1 (*Tgif1*), Interferon
regulatory factor 2-binding protein-like (*Irf2bpl*), Ephrin type-B receptor 3 (*Ephb3*),
Testis-specific Y-encoded-like protein 3 (*Tspyl3*), Protein ripply 3(*Ripply3*), Src kinase-
associated phosphoprotein 2 (Skap2), DNA polymerase zeta catalytic subunit (*Rev3l*),
MKL/Myocardin-Like 2 (*Mkl2*), and Protein phosphatase 2 regulatory subunit A
(*Ppp2r1a*) (Fig 8). The *Mecom* gene plays a role in embryogenesis and development
[149]. The *Mdga2* gene encodes immunoglobins involved in neural development [150].
The *Fgd5* gene is needed for embryonic development since it interacts with
hematopoietic stem cells [151]. The *Rbm19* gene is essential for preimplantation
development [152]. The *Coup* gene encodes a transcription factor that regulates

development of the eye [153] and other tissues [154]. The *Sim2* gene encodes a
transcription factor that regulates cell fate during midline development [155]. The
Slc38a4 gene encodes a regulator of protein synthesis during liver development and plays
a crucial role in fetal growth and development [156,157]. The *Bcl6* gene encodes a
transcription factor that controls neurogenesis [158]. The *Sema6d* gene encodes a protein
involved in retinal development [159]. The *Rbm45* gene encodes a protein that has
preferential binding to poly(C) RNA and is expressed during brain development [160].
The *E2f4* gene is involved in maturation of cells in tissues [161]. The *Lacs4* gene plays a
role in patterning in embryos [162]. The *Klk1b3* gene encodes a protein that plays a role
in developing embryos [163]. The *Sema4c* gene encodes a protein that has diverse
functions in neuronal development and heart morphogenesis [164, 165]. The *Tgif1* gene
encodes a transcription factor that plays a role in trophoblast differentiation [166]. The
Irf2bpl gene encodes a transcriptional regulator that plays a role in female
neuroendocrine reproduction [167]. The *Ephb3* gene encodes a kinase that plays a role in
neural development [168]. The *Tspyl3* gene plays a role in testis development [169].
The *Ripply3* gene encodes a transcription factor involved in development of the ectoderm
[170]. The *Skap2* gene encodes a protein involved in actin reorganization in lens
development [171]. The *Rev3l* gene encodes a polymerase that can replicate past certain
types of DNA lesions and is necessary for embryonic development [172]. The *Mkl2* gene
encodes a transcriptional co-activator is involved in the formation of muscular tissue
during embryonic development [173]. The *Ppp2r1a* gene plays a role in embryonic
epidermal development [174].

While the *Mecom, Mdga2, Fgd5, Rbm19, Coup, Sim2, Slc38a4, Bcl6, Sema6d, Rbm45,*
E2f4 and *Lacs4* genes were upregulated within 0.5 h postmortem, the other genes were
upregulated at 1 h to 48 h with most transcripts reaching abundance maxima at 12 h or
more.

Summary of development genes
In organismal death, there is progressive activation of developmental genes suggesting
that they are no longer silenced. A possible reason for this activation is that postmortem
physiological conditions resemble those during development.

Cancer genes
There are a number of databases devoted to cancer and cancer-related. Upon cross-
referencing the genes found in this study, we discovered a significant overlap. The genes
found in this search are presented below.

In the zebrafish, the following cancer genes were upregulated: *Jdp2,* xanthine
dehydrogenase (*Xdh*), *Egr1, Adam12*, myosin-IIIa (*Myo3a*), *Fosb, Jun,* Integrin alpha 6b
(*Itga6*), *Ier2, Tpr,* Dual specificity protein phosphatase 2 (*Dusp2*), Disintegrin and
metallopeptidase domain 28 (*Adam28*), *Tnpo1,* Ral guanine nucleotide dissociation
stimulator-like (*Rgl1*), Carcinoembryonic antigen-related cell adhesion molecule 5
(*Ceacam1*), *Fosl1, Il1b, Hif1a,* Serine/threonine-protein phosphatase 2A regulatory
(*Ppp2r5d*), DNA replication licensing factor (*Mcm5*), *Gadd45,* Myosin-9 (*Myh9*), *Casp3,*
Tnf, Il8, Cyclic AMP-dependent transcription factor (*Atf3*), small GTPase (*RhoA*),
Mknk2, Ephrin type-A receptor 7 precursor (*Epha7*), ETS-related transcription factor
(*Elf3*), *Nfkbia, Kpnb1, Wif1,* RAS guanyl-releasing protein 1 (*Rasgrp*), Ras association

domain-containing protein 6 (*Rassf6*), *Cyba,* DNA-damage-inducible transcript 3 (*Ddit3*),
Serine/threonine-protein kinase (*Sbk1*), and Tyrosine-protein kinase transmembrane
receptor (*Ror1*) (Fig 9).

Fig 9. Upregulated cancer genes by postmortem time (h). A, Zebrafish; B, Mouse.
Green, intermediate value; Red, maximum value. Bold gene name means it was
found in more than one cancer database. The *Rgl1* gene was represented by two
different probes.

In the mouse, the following cancer genes were upregulated: retinoblastoma-like protein 1
(*Rbl1*), Elongation factor RNA polymerase II (*Ell*), Bcl-2-like protein 11 (*Bcl2l11*), Sal-
like protein 1 (*Sall1*), *Map3k2, Bcl6, Tnfrsf9,* CK2 target protein 2 (*Csnk2a1*),
Transcription factor E2f4 (*E2f4*), Zinc finger DHHC-type containing 14 (*Zdhhc14*), *Tpr,*
RAS p21 protein activator 1 (*Rasa1*), *Gadd45,* prohibitin (*Phb2*), Serine/threonine-
protein phosphatase PP1-gamma catalytic (*Ppp1cc*), *Lasp1,* G protein-coupled receptor
kinase 4 (*Grk4*), LIM domain transcription factor (*Lmo4*), Protein phosphatase 1E
(*Ppm1e*), Protein sprouty homolog 1 (*Spry1*), Multiple PDZ domain protein (*Mpdz*),
Kisspeptin receptor (*Kiss1*), Receptor-type tyrosine-protein phosphatase delta precursor
(*Ptprd*), Small effector protein 2-like (*Cdc42*), AT-rich interactive domain-containing
protein 1A (*Arid1a*), Lymphocyte cytosolic protein 2 (*Lcp2*), DNA polymerase zeta
catalytic subunit (*Rev3l*), *Tnfrsf14,* Integrin beta-6 precursor (*Itgb6*), Triple functional
domain protein (*Trio*), ATPase class VI type 11C (*Atp11c*), and Serine/threonine-protein
phosphatase 2A regulatory (*Ppp2r1a*) (Fig 9).

Summary of cancer genes
These genes were classified as "cancer genes" in a Cancer Gene Database [7] (Fig 9).
The timing, duration and peak transcript abundances differed within and between
organisms. Note that some transcripts had two abundance maxima. In the zebrafish, this
phenomenon occurred for *Adam12, Jun, Tpr, Dusp2, Tnpo1,* and *Hif1a* and in the mouse,
Bcl6, Tnfrs9, Lasp1, Cdc42, and *Lcp2*, and is consistent with the notion that the genes are
being regulated through feedback loops.

Epigenetic regulatory genes
Epigenetic regulation of gene expression involves DNA methylation and histone
modifications of chromatin into active and silenced states [175]. These modifications
alter the condensation of the chromatin and affect the accessibility of the DNA to the
transcriptional machinery. Although epigenetic regulation plays important role in
development, modifications can arise stochastically with age or in response to
environmental stimuli [176]. Hence, we anticipated that epigenetic regulatory genes
should be involved in organismal death.

In the zebrafish, the following epigenetic genes were upregulated: Jun dimerization
protein 2 (*Jdp2*), Chromatin helicase protein 3 (*Chd3*), Glutamate-rich WD repeat-
containing protein 1 (*Grwd1*), Histone H1 (*Histh1l*), Histone cluster 1, H4-like
(*Hist1h46l3*) and Chromobox homolog 7a (*Cbx7a*) (Fig 10). The *Jdp2* gene is thought to
inhibit the acetylation of histones and repress expression of the c-Jun gene [177]. The
Chd3 gene encodes a component of a histone deacetylase complex that participates in the
remodeling of chromatin [178]. The *Grwd1* gene is thought to be a histone-binding
protein that regulates chromatin dynamics at the replication origin [179]. The *Histh1l*
gene encodes a histone protein that binds the nucleosome at the entry and exit sites of the
DNA and the *Hist1h46l3* gene encodes a histone protein that is part of the nucleosome
core [180]. The *Cbx7a* gene encodes an epigenetic regulator protein that binds non-
coding RNA and histones and represses gene expression of a tumor suppressor [181].

Fig 10. Upregulated epigenetic genes by postmortem time (h). A, Zebrafish; B, Mouse. Green, intermediate value; Red, maximum value. Bold gene name means it was found in more than one cancer database. The *Jdp2* gene was represented by two different probes.

Both *Jdp2* and *Chd3* genes were upregulated within 0.3 h postmortem, with their transcripts reaching abundance maxima at 0.5 h. Note that two different probes targeted the *Jdp2* transcript. The *Grwd1* gene was upregulated at 1 h and 24 h postmortem. The histone genes were upregulated at 4 h postmortem with their transcripts reaching abundance maxima at 24 h. The *Cbx7a* gene was upregulated at 12 h and its transcript reached an abundance maximum at 24 h. Transcripts of these genes were at low abundance after 24 h.

In the mouse, the following epigenetic genes were upregulated: Tubulin tyrosine ligase-like family member 10 (*Ttll10*), Histone cluster 1 H3f (*Hist1h3f*), Histone cluster 1 H4c (*Hist1h4c*), YEATS domain containing 2 (*Yeats2*), Histone acetyltransferase (*Kat7*), and Probable JmjC domain-containing histone demethylation protein 2C (*Jmjd1c*) (Fig 10). The *Ttll10* gene encodes a polyglycylase involved in modifying nucleosome assembly protein 1 that affects transcriptional activity, histone replacement, and chromatin remodeling [182]. The *Hist1h3f* and *Hist1h4c* genes encode histone proteins are the core of the nucleosomes [183]. The *Yeats2* gene encodes a protein that recognizes histone acetylations so that it can regulate gene expression in the chromatin [184]. The *Kat7* gene encodes an acetyltransferase that is a component of histone binding origin-of-replication complex, which acetylates chromatin and therefore regulates DNA replication and gene expression [185]. The *Jmjd1c* gene encodes an enzyme that specifically demethylates 'Lys-9' of histone H3 and is implicated in the reactivation of silenced genes [186].

The *Ttll10, Yeats2* and histone protein genes were upregulated 0.5 h postmortem but their transcripts reached abundance maxima at different times with the *Ttll10* transcript reaching a maximum at 1 to 6 h, the histone transcripts reaching maxima at 6 and 12 h postmortem, and the *Yeats2* transcript reaching maxima at 12 to 24 h postmortem (Fig 10). The *Kat7* and *Jmjd1c* genes were upregulated at 24 h and their transcripts reached abundance maxima at 48 h postmortem.

Summary of epigenetic regulatory genes
The upregulation of genes encoding histone proteins, histone-chromatin modifying proteins, and proteins involved in regulating DNA replication at the origin were common to the zebrafish and the mouse. These findings indicate that epigenetic regulatory genes

are modifying chromatin structure by regulating the accessibility of transcription factors
to the promoter or enhancer regions in organismal death.

Percentage of upregulated genes with postmortem time

The % of upregulated genes was defined as the number of upregulated genes at a specific
postmortem time over the total number of upregulated genes in a category. A
comparison of the % of upregulated genes by postmortem time of all upregulated genes
revealed similarities between the zebrafish and the mouse. Specifically, most genes were
upregulated between 0.5 to 24 h postmortem, and after 24 h, the upregulation of most
genes stopped (Fig 11, "All genes"). It should be noted that the same pattern was found
in stress, transport and development categories for both organisms. However, in the
zebrafish, the immunity, inflammation, apoptosis and cancer categories differed from the
mouse. Specifically, the genes in the immunity, inflammation, and cancer categories
were upregulated much later (1 to 4 h) in the zebrafish than the mouse, and the duration
of upregulation was much shorter. For example, while 90% of the genes in the immunity
and inflammation categories were upregulated in the mouse within 1 h postmortem, less
than 30% of the genes were upregulated in the zebrafish (Fig 11), indicating a slower
initial response. It should be noted that while the number of upregulated immunity genes
reaching transcript abundance maxima occurred at 24 h postmortem in both organisms,
the number of inflammation genes reaching transcript abundance maxima occurred at 1 to
4 h in the mouse, and 24 h in the zebrafish. The significance of these results is that the
inflammation response occurs rapidly and robustly in the mouse while in the zebrafish, it
takes longer to establish, which could be attributed to phylogenetic differences. There
were significant differences in the upregulation of apoptosis genes between the zebrafish
and the mouse. In the mouse, the number of upregulated apoptosis genes reached 100%
at 1 h postmortem and remained sustained for 48 h postmortem while the % of
upregulated genes in the zebrafish never reached 70% and gene upregulation was
abruptly stopped after 12 h.

Fig 11. % of upregulated genes by postmortem time and category. Number of total genes by organism and category are shown. "All genes" refers to the genes that significantly contributed to the ordination plots. Mouse is red; Zebrafish is black.

DISCUSSION

Active gene expression or residual transcription levels?

One could argue that our study identifies only residual transcription levels of pre-synthesized mRNA (rather than newly-synthesized mRNA) in dead tissues that happen to

be enriched with postmortem time. In other words, the observed upregulation of genes
may be viewed as an artifact merely reflecting the "enrichment of specific mRNA
transcripts" (e.g. stable mRNA) with time. The data, however, does not support this idea
because if it were true, one would expect stable transcripts to monotonically increase with
time, as they become more enriched (higher abundances) with postmortem time. The
data show that the transcripts of most upregulated genes did not display monotonic
behavior; rather, the transcripts reached abundance maximum (peak) or maxima (peaks)
at various postmortem times (Fig 2D and 2E). This finding should not be a surprise
because a statistical procedure was implemented to detect genes that were significantly
upregulated – which is essentially selecting gene transcriptional profiles that had peaks.
The statistics for the procedure was calibrated with more than a billion simulations. The
simulation process corrected for multiple comparisons. The residual transcription level
and enrichment idea is also not supported by transcriptional profiles displaying an up-,
down-, and up- regulation pattern, which putatively indicates feedback loops (Fig S3).
Similarly, simple differential decay rates of mRNAs would not display this pattern
because a transcript cannot be stable at one postmortem time, unstable at a subsequent
time, and then stable again. A mRNA transcript is either stable or it is not. In
conclusion, the gene profiles showing upregulation are the results of active transcription.
926
It should be noted that postmortem upregulation of genes has been previously reported in
cadavers. Using reverse transcription real-time quantitative PCR (RT-RTqPCR), a study
showed significant increases in expression of myosin light chain 3 (*Myl3*), matrix
metalloprotease 9 (*Mmp9*), and vascular endothelial growth factor A (*Vegfa*) genes in
body fluids after 12 h postmortem (187). Interestingly, postmortem upregulation of
myosin-related and matrix metalloprotease genes was also found in our study.
Specifically, the myosin-related genes included: Myosin-Ig (*Myo1g)* in the mouse*,* and
Myosin-IIIa (*Myo3a*) and Myosin-9 (*Myh9*) in the zebrafish. The matrix
metalloproteinase gene included the metalloproteinase-14 (*Mmp14b*) gene in the
zebrafish. The *Myo1g* gene encodes a protein regulating immune response (189), the
Myosin-IIIa (*Myo3a*) gene encodes an uncharacterized protein, the Myosin-9 (*Myh9*)
gene encodes a protein involved in embryonic development (190), and the *Mmp14b* gene
encodes an enzyme regulating cell migration during zebrafish gastrulation (188). The
Myo1g, Myh9 and *Mmp14b* transcripts increased right after death and reached abundance
maxima at 24 h postmortem, while the *Myo3a* transcript reached an abundance maximum
at 12 h postmortem. The significance of these results is two-fold: (i) two different
technologies (RT-RTqPCR and Gene Meters) have now demonstrated active postmortem
upregulation of genes and this expression has now been reported in three organisms
(human, zebrafish, and mouse), and (ii) there might be significant overlap in genes
upregulated in death as we have showed with myosin- and matrix metalloprotease genes,
which warrants further studies using other vertebrates. The purpose of such studies
would be to understand common mechanisms involved in the shutdown of highly ordered
biological systems.

Why study gene expression in death?
The primary motivation for the study was driven by curiosity in the processes of shutting
down a complex biological system – which has received little attention so far. While the

development of a complex biological system requires time and energy, its shutdown and
subsequent disassembly entails the dissipation of energy and unraveling of the complex
structures, which does not occur instantaneously and could provide insights into
interesting paths. Moreover, other fields of research have examined the shutdown of
complex systems (e.g., societies [191], government [192], electrical black outs [193]).
Yet, to our knowledge, no study has examined long-term postmortem gene expression of
vertebrates kept in their native conditions. The secondary motivation for the study was to
demonstrate the precision of Gene Meter technology for gene expression studies to
biologists who believe that high throughput DNA sequencing is the optimal approach.

Thermodynamic sinks

We initially thought that sudden death of a vertebrate would be analogous to a car driving
down a highway and running out of gas. For a short time, engine pistons will move up
and down and spark plugs will spark -- but eventually the car will grind to a halt and
"die". Yet, in our study we find hundreds of genes are upregulated many hours
postmortem, with some (e.g., *Kcnv2, Pafr, Degs2, Ogfod1, Ppp2rla, Ror1,* and *Iftm1*)
upregulated days after organismal death. This finding is surprising because in our car
analogy, one would not expect window wipers to suddenly turn on and the horn to honk
several days after running out of gas.

Since the postmortem upregulation of genes occurred in both the zebrafish and the mouse
in our study, it is reasonable to suggest that other multicellular eukaryotes will display a
similar phenomenon. What does this phenomenon mean in the context of organismal
life? We conjecture that the highly ordered structure of an organism – evolved and
refined through natural selection and self-organizing processes – undergoes a
thermodynamically driven process of spontaneous disintegration through complex
pathways, which apparently involve the upregulation of genes and feedback loops. While
evolution played a role in pre-patterning of these pathways, it does not play any role in its
disintegration fate. One could argue that some of these pathways have evolved to favor
healing or "resuscitation" after severe injury. For example, the upregulation of
inflammation response genes indicate that a signal of infection or injury is sensed by the
still alive cells after death of the body. Alternatively, the upregulation may be due to fast
decay of some repressors of genes or whole pathways (see below). Hence, it will be of
interest to study this in more detail, since this could, for example, provide insights into
how to better preserve organs retrieved for transplantation.

Chemical automator – on the way down to equilibrium

As one would expect, a living system is a collection of chemical reactions linked together
by the chemicals participating in them. Having these reactions to depend on one another
to a certain extent, we conjecture that the observed upregulation of genes is due to
thermodynamic and kinetic controls that are encountered during organismal death. For
example, epigenetic regulatory genes that were upregulated included histone modification
genes (e.g., *Histh1l*) and genes interacting with chromatin (e.g., *Grwd1, Chd3, Yeats,*
Jmjd1c) (Fig 10). It is possible that the activation of these genes was responsible for the
unraveling of the nucleosomes, which enabled transcription factors and RNA
polymerases to transcribe developmental genes that have been previously silenced since

embryogenesis. The energy barrier in this example is the tightly wrapped nucleosomes
that previously did not allow access to developmental genes. Other energy or entropy
barriers could be nucleopores that allow the exchange of mRNA and other molecules
between the mitochondria and the cytosol (e.g., *Tpr, Tnpo1, Lrrc59*), or the ion/solute
protein channels (e.g., *Aralar2, Slc38a4*) that control intracellular ions that regulate
apoptotic pathways [194,195].

The upregulation of genes indicates new molecules were synthesized. Hence, there was
sufficient energy and resources (e.g., RNA polymerase, dNTPs) in dead organisms to
maintain gene transcription to 96 h (e.g., *Zfand4, Tox2*, and *Slc14a2*) in the zebrafish and
to 48 h (e.g., *Deg2, Ogfod1,* and *Ifitm1*) in the mouse. Gene transcription was apparently
not prevented due to a lack of energy or resources. Several genes exhibited apparent
regulation by feedback loops in their transcriptional profiles (e.g., *Rbm45* and *Cdc42*
genes in the mouse (Fig 8 and 9, respectively; Fig S3)). Hence, an underlying regulatory
network appears to be still turning "on" and "off" genes in organismal death.

Interrupt the shutdown?

1010
A living biological system is a product of natural selection and self-organizing processes
[196]. Genes are transcribed and proteins translated in response to genetic and epigenetic
regulatory networks that sustain life. In organismal death, we assumed most of the
genetic and epigenetic regulatory networks operating in life would become disengaged
from the rest of the organism. However, we found that "dead" organisms turn genes on
and off in a non-random manner (Fig 2D and 2E). There is a range of times in which
genes are upregulated and transcript abundances are maximized. While most genes are
upregulated within 0.5 h postmortem (Fig 11), some are upregulated at 24 h and still
others at 48 h. A similar pattern occurs with peak transcript abundances and the timing
when upregulation is apparently stopped. These differences in timings and abundances
suggest some sort of global regulation network is still operating in both organisms. What
makes gene expression of life different from gene expression in death is that postmortem
upregulation of genes offers no obvious benefit to an organism. We argue that self-
organizing processes driven by thermodynamics are responsible for the postmortem
upregulation of genes. We emphasize that such postmortem conditions could allow
investigators to tease apart evolution from self-organizing processes that are typically
entangled in life.

Since our results show that the system has not reached equilibrium yet, it would be
interesting to address the following question: *what would happen if we arrested the*
process of dying by providing nutrients and oxygen to tissues? It might be possible for
cells to revert back to life or take some interesting path to differentiating into something
new or lose differentiation altogether, such as in cancer. We speculate that the recovering
cells will likely depend on the postmortem time – at least when such potentially
interesting effects might be seen.

Methodological validity

1035
The Gene Meter approach is pertinent to the quality of the microarray output obtained in
this study because conventional DNA microarrays yield noisy data [197,198]. The Gene
Meter approach determines the behavior of every microarray probe by calibration –

which is analogous to calibrating a pH meter with buffers. Without calibration, the
precision and accuracy of a meter is not known, nor can one know how well the
experimental data fits to the calibration (i.e., R^2). In the Gene Meter approach, the
response of a probe (i.e., its behavior in a dilution series) is fitted to either Freundlich or
Langmuir adsorption model, probe-specific parameters are calculated. The "noisy" or
"insensitive" probes are identified and removed from further analyses. Probes that
sufficiently fit the model are retained and later used to calculate the abundance of a
specific gene in a biological sample. The models take into consideration the non-linearity
of the microarray signal and the calibrated probes do not require normalization
procedures to compare biological samples. In contrast, conventional DNA microarray
approaches are biased because different normalizations can yield up to 20 to 30%
differences in the up- or down-regulation depending on the procedure selected [199-202].
We recognize that next-generation-sequencing (NGS) approaches could have been used
to monitor gene expression in this study. However, the same problems of normalization
and reproducibility (mentioned above) are pertinent to NGS technology [203]. Hence,
the Gene Meter approach is currently the most advantageous to study postmortem gene
expression in a high throughput manner.

Practical implications

The postmortem upregulation of genes in the mouse has relevance to transplantation
research. We observed clear qualitative and quantitative differences between two organs
(liver and brain) in the mouse in their degradation profiles (Fig 1). We also showed the
upregulation of immunity, inflammation and cancer genes within 1 h of death (Fig 11). It
would be interesting to explore if these differences are comparable to what occurs in
humans, and we wonder how much of the transplant success could be attributed to
differences in the synchronicity of postmortem expression profiles rather than
immunosuppression agents [204,205]. Our study provides an alternative perspective to
the fate of transplant recipients due to the upregulation of regulatory and response genes,
after the sample has been harvested from the donor.

Conclusion

This is the first study to demonstrate active, long-term expression of genes in organismal
death that raises interesting questions relative to transplantology, inflammation, cancer,
evolution, and molecular biology.

1071

1072 **Competing Interests**

1073 The authors declare that they have no competing interests.

1074

Authors' contributions

DT, RN, TDL and AEP designed the study. AEP, RN, TDL carried out the molecular genetic studies. BGL and AEP determined the statistically significant upregulated genes. SS and PAN annotated the genes. AEP and PAN conducted the statistical analyses and wrote the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We thank Till Sckerl for help with sacrificing the mice, Nicole Thomsen for help with tissue and RNA processing, and Elke Blohm-Sievers for microarray work. We thank Russell Bush for his helpful advice on thermodynamics and entropic barriers and M. Colby Hunter for proof-reading the manuscript.

Financial disclosure The work was supported by funds from the National Cancer Institute P20 Partnership grant (P20 CA192973 and P20 CA192976) and the Max-Planck-Society.

REFERENCES

1. Monod J. Genetic regulatory mechanisms in the synthesis of proteins. J Mol Biol. 1961;3: 318.

2. Zong WX, Thompson CB. Necrotic death as a cell fate. Genes Dev. 2006, 20: 1.

3. GalluzziL, Bravo-San Pedro JM, Kroemer G. Organelle-specific initiation of cell death. Nat Cell Biol. 2014;16: 728-736.

4. Ferriferri_2001 KF, Kroemer G. Organelle-specific initiation of cell death pathways. Nat Cell Biol. 2001;3: E255.

5. Syntichakisyntichaki_2002 P, Tavernarakis N. Death by necrosis. Uncontrollable catastrophe, or is there order behind the chaos? EMBO Rep. 2002;3: 604.

6. Pozhitkov AE, Noble PA, Bryk J, Tautz D. A revised design for microarray experiments to account for experimental noise and uncertainty of probe response. PLoS One. 2014;9: e91295.

7. Domazet-Loso T, Tautz D. Phylostratigraphic tracking of cancer genes suggests a link to the emergence of multicellularity in metazoa. BMC Biol. 2010;8: 66. doi: 10.1186/1741-7007-8-66.

8. Seear PJ, Sweeney GE. Stability of RNA isolated from post-mortem tissues of Atlantic salmon (Salmo salar L.). Fish Physiol Biochem. 2008;34: 19-24. doi: 10.1007/s10695-007-9141-x.

9. Bahar B, Monahan FJ, Moloney AP, Schmidt O, MacHugh DE, Sweeney T. Long-term stability of RNA in post-mortem bovine skeletal muscle, liver and subcutaneous adipose tissues. BMC Mol Biol. 2007;8: 108.

10. Inoue H, Kimura A, Tuji T. Degradation profile of mRNA in a dead rat body: basic semi-quantification study. Forensic Sci Int. 2002;130: 127-32.

11. Johnson SA, Morgan DG, Finch CE. Extensive postmortem stability of RNA from rat and human brain. J Neurosci Res. 1986;16: 267-80.

12. Catts VS, Catts SV, Fernandez HR, Taylor JM, Coulson EJ, Lutze-Mann LH. A microarray study of post-mortem mRNA degradation in mouse brain tissue. Brain Res Mol Brain Res. 2005;138: 164-77.

13. Heinrich M, Matt K, Lutz-Bonengel S, Schmidt U. Successful RNA extraction from various human postmortem tissues. Int J Legal Med. 2007;121: 136-42.

14. Neymotin, Athanasiadou R, Gresham D. Determination of in vivo RNA kinetics using RATE-seq. RNA. 2014;20: 1645.

15. Spilka R, Ernst C, Bergler H, Rainer J, Flechsig S, Vogetseder A, Lederer E, Benesch M, Brunner A, Geley S, Eger A, Bachmann F, Doppler W, Obrist P, Haybaeck J. eIF3a is over-expressed in urinary bladder cancer and influences its phenotype independent of translation initiation. Cell Oncol (Dordr). 2014;37: 253

16. Bortolotti GR, Tella JL, Baos R, Marchant TA. Stress response during development
predicts fitness in a wild, long lived vertebrate. Proc Natl Acad Sci USA. 2007;104:
8880-4.

17. Skaggs HS, Xing H, Wilkerson DC, Murphy LA, Hong Y, Mayhew CN, Sarge KD.
HSF1-TPR interaction facilitates export of stress-induced HSP70 mRNA. J Biol
Chem. 2007;282: 33902-7.

18. Snow CJ, Paschal BM. Roles of the nucleoporin Tpr in cancer and aging. Adv Exp
Med Biol. 2014;773: 309-22. doi: 10.1007/978-1-4899-8032-8_14.

19. David-Watine B. Silencing nuclear pore protein Tpr elicits a senescent-like
phenotype in cancer cells. PLoS One. 2011;6: e22423. doi:
10.1371/journal.pone.0022423.

20. Kadeba PI, Scammel JG, Cioffi DL. The chaperone heat shock protein 90 (Hsp90)
participates in the endothelial store operated calcium entry heterocomplex. FASEB J.
2013;27: 724.4

21. Mayer MP, Bukau B. Hsp70 chaperones: cellular functions and molecular
mechanism. Cell Mol Life Sci. 2005;62: 670-84.

22. Jakobsson ME, Davydova E, Małecki J, Moen A, Falnes, PØ. Saccharomyces
cerevisiae eukaryotic elongation factor 1A (eEF1A) is methylated at Lys-390 by a
METTL21-like methyltransferase. PLos One 2015;
doi:10.1371/journal.pone.0131426.t002.

23. Höhfeld J, Hartl FU. Requirement of the chaperonin cofactor Hsp10 for protein
sorting in yeast mitochondria. J Cell Biol/ 1994;126:305–315.

24. Wu BJ, Morimoto RI. Transcription of the human hsp70 gene is induced by serum
stimulation. Proc Natl Acad Sci USA. 1985;82: 6070-4.

25. Ngr di A, Domoki F, Dgi R, Borda S, Pkski M, Szab A, Bari F. Up-regulation of
cerebral carbonic anhydrase by anoxic stress in piglets. J Neurochem. 2003;85: 843-
50.

26. Lin KH, Kuo CH, Kuo WW, Ho TJ, Pai P, Chen WK, Pan LF, Wang CC, Padma VV,
Huang CY. NFIL3 suppresses hypoxia-induced apoptotic cell death by targeting the
insulin-like growth factor 2 receptor. J Cell Biochem. 2015;116: 1113-20. doi:
10.1002/jcb.25067

27. Zhang W, Zhang J, Kornuc M, Kwan K, Frank R, Nimer SD. Molecular cloning and
characterization of NF-IL3A, a transcriptional activator of the human interleukin-3
promoter. Mol Cell Biol. 1995;15: 6055-63.

28. Smith TG, Robbins PA, Ratcliffe PJ. The human side of hypoxia-inducible factor. Br
J Haematol. 2008, 141: 325-34. doi: 10.1111/j.1365-2141.2008.07029.x.

29. Ino Y, Yamazaki-Itoh R, Oguro S, Shimada K, Kosuge T, Zavada J, Kanai Y,
Hiraoka N. Arginase II expressed in cancer-associated fibroblasts indicates tissue
hypoxia and predicts poor outcome in patients with pancreatic cancer. PLoS One.
2013;8: e55146. doi: 10.1371/journal.pone.0055146.

30. Lafleur VN, Richard S, Richard DE. Transcriptional repression of hypoxia-inducible
factor-1 (HIF-1) by the protein arginine methyltransferase PRMT1. Mol Biol Cell.
2014;25: 925-35. doi: 10.1091/mbc.E13-07-0423.

31. Devlin CM, Lahm T, Hubbard WC, Van Demark M, Wang KC, Wu X, Bielawska A,
Obeid LM, Ivan M, Petrache I. Dihydroceramide-based response to hypoxia. J Biol
Chem. 2011;286: 38069-78. doi: 10.1074/jbc.M111.297994.

32. Hannun YA, Obeid LM. Principles of bioactive lipid signalling: lessons from
sphingolipids. Nat Rev Mol Cell Biol. 2008;9: 139-50. doi: 10.1038/nrm2329.

33. Mao C, Obeid LM. Ceramidases: regulators of cellular responses mediated by
ceramide, sphingosine, and sphingosine-1-phosphate. Biochim Biophys Acta.
2008;1781: 424-34. doi: 10.1016/j.bbalip.2008.06.002.

34. Rhee SG, Chae HZ, Kim K. Peroxiredoxins: a historical overview and speculative
preview of novel mechanisms and emerging concepts in cell signaling. Free Radic
Biol Med. 2005;38: 1543-52.

35. Salzano S, Checconi P, Hanschmann EM, Lillig CH, Bowler LD, Chan P, Vaudry D,
Mengozzi M, Coppo L, Sacre S, Atkuri KR, Sahaf B, Herzenberg LA, Herzenberg
LA, Mullen L, Ghezzi P. Linkage of inflammation and oxidative stress via release of
glutathionylated peroxiredoxin-2, which acts as a danger signal. Proc Natl Acad Sci
USA. 2014;111: 12157-62. doi: 10.1073/pnas.1401712111.

36. Landau G, Ran A, Bercovich Z, Feldmesser E, Horn-Saban S, Korkotian E, Jacob-
Hirsh J, Rechavi G, Ron D, Kahana C. Expression profiling and biochemical analysis
suggest stress response as a potential mechanism inhibiting proliferation of
polyamine-depleted cells. J Biol Chem. 2012; 287:35825-37. doi:
10.1074/jbc.M112.381335.

37. Fornace AJ Jr, Jackman J, Hollander MC, Hoffman-Liebermann B, Liebermann DA.
Genotoxic-stress-response genes and growth-arrest genes. gadd, MyD, and other
genes induced by treatments eliciting growth arrest. Ann N Y Acad Sci. 1992;663:
139-53.

38. Schmitz I. Gadd45 proteins in immunity. In: D.A. Liebermann and B. Hoffman (eds),
Gadd45 Stress Sensor Genes, Advances in Experimental Medicine and Biology 793,
Springer Science+Business Media New York 2013.

39. Lisowski P, Wieczorek M, Goscik J, Juszczak GR, Stankiewicz AM, Zwierzchowski
L, Swiergiel AH. Effects of chronic stress on prefrontal cortex transcriptome in mice
displaying different genetic backgrounds. J Mol Neurosci. 2013;50: 33-57. doi:
10.1007/s12031-012-9850-1.

40. Timpl P, Spanagel R, Sillaber I, Kresse A, Reul JM, Stalla GK, Blanquet V, Steckler
T, Holsboer F, Wurst W. Impaired stress response and reduced anxiety in mice
lacking a functional corticotropin-releasing hormone receptor 1. Nat Genet. 1998;19:
162-6.

41. Leikauf GD, Concel VJ, Bein K, Liu P, Berndt A, Martin TM, Ganguly K, Jang AS,
Brant KA, Dopico RA Jr, Upadhyay S, Cario C, Di YP, Vuga LJ, Kostem E, Eskin E,
You M, Kaminski N, Prows DR, Knoell DL, Fabisiak JP. Functional genomic

assessment of phosgene-induced acute lung injury in mice. Am J Respir Cell Mol Biol. 2013;49: 368-83. doi: 10.1165/rcmb.2012-0337OC.

42. Nice TJ, Deng W, Coscoy L, Raulet DH. Stress-regulated targeting of the NKG2D ligand Mult1 by a membrane-associated RING-CH family E3 ligase. J Immunol. 2010;185: 5369-76. doi: 10.4049/jimmunol.1000247.

43. Kokame K, Agarwala KL, Kato H, Miyata T. Herp, a new ubiquitin-like membrane protein induced by endoplasmic reticulum stress. J Biol Chem. 2000;275: 32846-53.

44. Coates PJ, Nenutil R, McGregor A, Picksley SM, Crouch DH, Hall PA, Wright EG. Mammalian prohibitin proteins respond to mitochondrial stress and decrease during cellular senescence. Exp Cell Res. 2001; 265:262-73.

45. Wehner KA, Schtz S, Sarnow P. OGFOD1, a novel modulator of eukaryotic translation initiation factor 2alpha phosphorylation and the cellular response to stress. Mol Cell Biol. 2010;30: 2006-16. doi: 10.1128/MCB.01350-09.

46. Rock KL, Lai JJ, Kono H. Innate and adaptive immune responses to cell death. Immunol Rev. 2011;243: 191-205. doi: 10.1111/j.1600-065X.2011.01040.x.

47. Crist SA, Elzey BD, Ahmann MT, Ratliff TL. Early growth response-1 (EGR-1) and nuclear factor of activated T cells (NFAT) cooperate to mediate CD40L expression in megakaryocytes and platelets. J Biol Chem. 2013;288: 33985-96. doi: 10.1074/jbc.M113.511881.

48. Li S, Miao T, Sebastian M, Bhullar P, Ghaffari E, Liu M, Symonds AL, Wang P. The transcription factors Egr2 and Egr3 are essential for the control of inflammation and antigen-induced proliferation of B and T cells. Immunity. 2012;37: 685-96. doi: 10.1016/j.immuni.2012.08.001.

49. Jayaraman P, Sada-Ovalle I, Nishimura T, Anderson AC, Kuchroo VK, Remold HG, Behar SM. IL-1β promotes antimicrobial immunity in macrophages by regulating TNFR signaling and caspase-3 activation. J Immunol. 2013;190: 4196-204. doi: 10.4049/jimmunol.1202688.

50. Hughes AL. Origin and diversification of the L-amino oxidase family in innate immune defenses of animals. Immunogenetics. 2010;62: 753-9. doi: 10.1007/s00251-010-0482-8.

51. Ramirez-Carrozzi V, Sambandam A, Luis E, Lin Z, Jeet S, Lesch J, Hackney J, Kim J, Zhou M, Lai J, Modrusan Z, Sai T, Lee W, Xu M, Caplazi P, Diehl L, de Voss J, Balazs M, Gonzalez L Jr, Singh H, Ouyang W, Pappu R. IL-17C regulates the innate immune function of epithelial cells in an autocrine manner. Nat Immunol. 2011;12: 1159-66 doi: 10.1038/ni.2156.

52. Zuccolo J, Bau J, Childs SJ, Goss GG, Sensen CW, Deans JP. Phylogenetic analysis of the MS4A and TMEM176 gene families. PLoS One. 2010, 5: e9369. doi: 10.1371/journal.pone.0009369.

53. Allen A, Hutton DA, Pearson JP. The MUC2 gene product: a human intestinal mucin. Int J Biochem Cell Biol. 1998;30: 797-801.

54. Michelucci A, Cordes T, Ghelfi J, Pailot A, Reiling N, Goldmann O, Binz T, Wegner
A, Tallam A, Rausell A, Buttini M, Linster CL, Medina E, Balling R, Hiller K.
Immune-responsive gene 1 protein links metabolism to immunity by catalyzing
itaconic acid production. Proc Natl Acad Sci USA. 2013;110: 7820-5. doi:
10.1073/pnas.1218599110.

55. Honke N, Shaabani N, Cadeddu G, Sorg UR, Zhang DE, Trilling M, Klingel K,
Sauter M, Kandolf R, Gailus N, van Rooijen N, Burkart C, Baldus SE, Grusdat M,
Lhning M, Hengel H, Pfeffer K, Tanaka M, Hussinger D, Recher M, Lang PA, Lang
KS. Enforced viral replication activates adaptive immunity and is essential for the
control of a cytopathic virus. Nat Immunol. 2011;13: 51-7. doi: 10.1038/ni.2169.

56. Martnez-Lpez M, Iborra S, Conde-Garrosa R, Sancho D. Batf3-dependent CD103+
dendritic cells are major producers of IL-12 that drive local Th1 immunity against
Leishmania major infection in mice. Eur J Immunol. 2015;45: 119-29. doi:
10.1002/eji.201444651.

57. Imai T, Baba M, Nishimura M, Kakizaki M, Takagi S, Yoshie O. The T cell-directed
CC chemokine TARC is a highly specific biological ligand for CC chemokine
receptor 4. J Biol Chem. 1997;272: 15036-42.

58. Vong QP, Leung WH1, Houston J1, Li Y1, Rooney B1, Holladay M1, Oostendorp
RA2, Leung W3. TOX2 regulates human natural killer cell development by
controlling T-BET expression. Blood. 2014;124: 3905-13. doi: 10.1182/blood-2014-
06-582965.

59. Takaori A. Antiviral defense by APOBEC3 family proteins. Uirusu. 2005;55: 267-72.

60. Arming S, Wipfler D, Mayr J, Merling A, Vilas U, Schauer R, Schwartz-Albiez R,
Vlasak R. The human Cas1 protein: a sialic acid-specific O-acetyltransferase?
Glycobiology. 2011;21: 553-64. doi: 10.1093/glycob/cwq153.

61. Crocker PR, Paulson JC, Varki A. Siglecs and their roles in the immune system. Nat
Rev Immunol. 2007;7: 255-66.

62. Crocker PR, Varki A. Siglecs in the immune system. Immunology. 2001;103: 137-45.

63. Shi WX, Chammas R, Varki NM, Powell L, Varki A. Sialic acid 9-O-acetylation on
murine erythroleukemia cells affects complement activation, binding to I-type lectins,
and tissue homing. J Biol Chem. 1996;271: 31526-32.

64. Praper T, Sonnen A, Viero G, Kladnik A, Froelich CJ, Anderluh G, Dalla Serra M,
Gilbert RJ. Human perforin employs different avenues to damage membranes. J Biol
Chem. 2011;286: 2946-55. doi: 10.1074/jbc.M110.169417.

65. Tschopp J, Masson D, Stanley KK. Structural/functional similarity between proteins
involved in complement- and cytotoxic T-lymphocyte-mediated cytolysis. Nature.
1986;322: 831-4.

66. White SH, Wimley WC, Selsted ME. Structure, function, and membrane integration
of defensins. Curr Opin Struct Biol. 1995;5: 521-7.

67. Blanchong CA, Chung EK, Rupert KL, Yang Y, Yang Z, Zhou B, Moulds JM, Yu
CY. Genetic, structural and functional diversities of human complement components

C4A and C4B and their mouse homologues, Slp and C4. Int Immunopharmacol.
2001;1: 365-92.

68. Chen L, Carico Z, Shih HY, Krangel MS. A discrete chromatin loop in the mouse
Tcra-Tcrd locus shapes the TCRδ and TCRα repertoires. Nat Immunol. 2015;16:
1085-93. doi: 10.1038/ni.3232.

69. Grage-Griebenow E, Flad HD, Ernst M, Bzowska M, Skrzeczyska J, Pryjma J.
Human MO subsets as defined by expression of CD64 and CD16 differ in phagocytic
activity and generation of oxygen intermediates. Immunobiology. 2000;202: 42-50.

70. Al-Banna NA, Vaci M, Slauenwhite D, Johnston B, Issekutz TB. CCR4 and CXCR3
play different roles in the migration of T cells to inflammation in skin, arthritic joints,
and lymph nodes. Eur J Immunol. 2014;44: 1633-43. doi: 10.1002/eji.201343995.
Epub 2014 Apr 23.

71. Ikutani M, Yanagibashi T, Ogasawara M, Tsuneyama K, Yamamoto S, Hattori Y,
Kouro T, Itakura A, Nagai Y, Takaki S, Takatsu K. Identification of innate IL-5-
producing cells and their role in lung eosinophil regulation and antitumor immunity. J
Immunol. 2012;188: 703-13. doi: 10.4049/jimmunol.1101270. Epub 2011 Dec 14.

72. Lillard JW Jr, Boyaka PN, Taub DD, McGhee JR. RANTES potentiates antigen-
specific mucosal immune responses. J Immunol. 2001;166: 162-9.

73. Kawasaki N, Rademacher C, Paulson JC. CD22 regulates adaptive and innate
immune responses of B cells. J Innate Immun. 2011;3: 411-9. doi:
10.1159/000322375.

74. Liu H, Thaker YR, Stagg L, Schneider H, Ladbury JE, Rudd CE. SLP-76 sterile α
motif (SAM) and individual H5 α helix mediate oligomer formation for microclusters
and T-cell activation. J Biol Chem. 2013;288: 29539-49. doi:
10.1074/jbc.M112.424846.

75. Feeley EM, Sims JS, John SP, Chin CR, Pertel T, Chen LM, Gaiha GD, Ryan BJ,
Donis RO, Elledge SJ, Brass AL. IFITM3 inhibits influenza A virus infection by
preventing cytosolic entry. PLoS Pathog. 2011;7: e1002337. doi:
10.1371/journal.ppat.1002337.

76. Saben J, Zhong Y, Gomez-Acevedo H, Thakali KM, Borengasser SJ, Andres A,
Shankar K. Early growth response protein-1 mediates lipotoxicity-associated
placental inflammation: role in maternal obesity. Am J Physiol Endocrinol Metab.
2013;305: e1-14. doi: 10.1152/ajpendo.00076.2013.

77. Ren K, Torres R. Role of interleukin-1beta during pain and inflammation. Brain Res
Rev. 2009;60: 57-64. doi: 10.1016/j.brainresrev.2008.12.020.

78. Chen GY, Nuñez G. Sterile inflammation: sensing and reacting to damage. Nat Rev
Immunol. 2010;10: 826-37. doi: 10.1038/nri2873.

79. Sedger LM, McDermott MF. TNF and TNF-receptors: From mediators of cell death
and inflammation to therapeutic giants - past, present and future. Cytokine Growth
Factor Rev. 2014;25: 453-72. doi: 10.1016/j.cytogfr.2014.07.016.

80. Lee TS, Chau LY. Heme oxygenase-1 mediates the anti-inflammatory effect of
interleukin-10 in mice. Nat Med. 2002;8: 240-6.

81. Piantadosi CA, Withers CM, Bartz RR, MacGarvey NC, Fu P, Sweeney TE, Welty-
Wolf KE, Suliman HB. Heme oxygenase-1 couples activation of mitochondrial
biogenesis to anti-inflammatory cytokine expression. J Biol Chem. 2011;286: 16374-
85. doi: 10.1074/jbc.M110.207738.

82. Guo Y, Zhang W, Giroux C, Cai Y, Ekambaram P, Dilly AK, Hsu A, Zhou S,
Maddipati KR, Liu J, Joshi S, Tucker SC, Lee MJ, Honn KV. Identification of the
orphan G protein-coupled receptor GPR31 as a receptor for 12-(S)-
hydroxyeicosatetraenoic acid. J Biol Chem. 2011;286: 33832-40. doi:
10.1074/jbc.M110.216564.

83. Bickel M. The role of interleukin-8 in inflammation and mechanisms of regulation. J
Periodontol. 1993;64: 456-60.

84. Alcock J, Franklin ML, Kuzawa CW. Nutrient signaling: evolutionary origins of the
immune-modulating effects of dietary fat. Q Rev Biol 2012;87: 187-223.

85. Pearson G, Robinson F, Beers Gibson T, Xu BE, Karandikar M, Berman K, Cobb
MH. Mitogen-activated protein (MAP) kinase pathways: regulation and physiological
functions. Endocr Rev. 2001;22: 153-83.

86. Tsatsanis C, Androulidaki A, Dermitzaki E, Gravanis A, Margioris AN. Corticotropin
releasing factor receptor 1 (CRF1) and CRF2 agonists exert an anti-inflammatory
effect during the early phase of inflammation suppressing LPS-induced TNF-alpha
release from macrophages via induction of COX-2 and PGE2. J Cell Physiol.
2007;210: 774-83.

87. Frugier T, Morganti-Kossmann MC, O'Reilly D, McLean CA. In situ detection of
inflammatory mediators in post mortem human brain tissue after traumatic injury. J
Neurotrauma. 2010;27: 497-507. doi: 10.1089/neu.2009.1120.

88. Sawant DV, Sehra S, Nguyen ET, Jadhav R, Englert K, Shinnakasu R, Hangoc G,
Broxmeyer HE, Nakayama T, Perumal NB, Kaplan MH, Dent AL. Bcl6 controls the
Th2 inflammatory activity of regulatory T cells by repressing Gata3 function. J
Immunol. 2012;189: 4759-69. doi: 10.4049/jimmunol.1201794.

89. Correa-Costa M, Andrade-Oliveira V, Braga TT, Castoldi A, Aguiar CF, Origassa
CS2, Rodas AC, Hiyane MI, Malheiros DM, Rios FJ, Jancar S, Cmara NO.
Activation of platelet-activating factor receptor exacerbates renal inflammation and
promotes fibrosis. Lab Invest. 2014;94: 455-66. doi: 10.1038/labinvest.2013.155.

90. Watson RP, Lilley E, Panesar M, Bhalay G, Langridge S, Tian SS, McClenaghan C,
Ropenga A, Zeng F, Nash MS. Increased prokineticin 2 expression in gut
inflammation: role in visceral pain and intestinal ion transport. Neurogastroenterol
Motil. 2012;24: 65-75 doi: 10.1111/j.1365-2982.2011.01804.x.

91. Pavlov VA, Wang H, Czura CJ, Friedman SG, Tracey KJ. The cholinergic anti-
inflammatory pathway: a missing link in neuroimmunomodulation. Mol Med. 2003;9:
125-34.

92. Lerdrup M, Holmberg C, Dietrich N, Shaulian E, Herdegen T, Jttel M, Kallunki T.
Depletion of the AP-1 repressor JDP2 induces cell death similar to apoptosis.
Biochim Biophys Acta. 2005;1745: 29-37.

93. Hu W, Xu R, Sun W, Szulc ZM, Bielawski J, Obeid LM, Mao C. Alkaline
ceramidase 3 (ACER3) hydrolyzes unsaturated long-chain ceramides, and its down-
regulation inhibits both cell proliferation and apoptosis. J Biol Chem. 2010;285:
7964-76. doi: 10.1074/jbc.M109.063586.

94. Preston GA, Lyon TT, Yin Y, Lang JE, Solomon G, Annab L, Srinivasan DG,
Alcorta DA, Barrett JC. Induction of apoptosis by c-Fos protein. Mol Cell Biol.
1996;16: 211-8.

95. Adrain C, Creagh EM, Martin SJ. Apoptosis-associated release of Smac/DIABLO
from mitochondria requires active caspases and is blocked by Bcl-2. EMBO J.
2001;20: 6627-36.

96. McNeish IA, Lopes R, Bell SJ, McKay TR, Fernandez M, Lockley M, Wheatley SP,
Lemoine NR. Survivin interacts with Smac/DIABLO in ovarian carcinoma cells but
is redundant in Smac-mediated apoptosis. Exp Cell Res. 2005;302: 69-82.

97. Zhao F, Wang Q. The protective effect of peroxiredoxin II on oxidative stress
induced apoptosis in pancreatic β-cells. Cell Biosci. 2012;2: 22. doi: 10.1186/2045-
3701-2-22.

98. Wu X, Hernandez-Enriquez B, Banas M, Xu R, Sesti F. Molecular mechanisms
underlying the apoptotic effect of KCNB1 K+ channel oxidation. J Biol Chem.
2013;288: 4128-34. doi: 10.1074/jbc.M112.440933.

99. Tang D, Lahti JM, Kidd VJ. Caspase-8 activation and bid cleavage contribute to
MCF7 cellular execution in a caspase-3-dependent manner during staurosporine-
mediated apoptosis. J Biol Chem. 2000;275: 9303-7.

100. Chinnadurai G, Vijayalingam S, Rashmi R. BIK, the founding member of the
BH3-only family proteins: mechanisms of cell death and role in cancer and
pathogenic processes. Oncogene. 2008;27: S20-9. doi: 10.1038/onc.2009.40.

101. Iwasa H, Kudo T, Maimaiti S, Ikeda M, Maruyama J, Nakagawa K, Hata Y. The
RASSF6 tumor suppressor protein regulates apoptosis and the cell cycle via MDM2
protein and p53 protein. J Biol Chem. 2013;288:30320-9. doi:
10.1074/jbc.M113.507384.

102. Liu X, Dong C, Jiang Z, Wu WK, Chan MT, Zhang J, Li H, Qin K, Sun X.
MicroRNA-10b downregulation mediates acute rejection of renal allografts by
derepressing BCL2L11. Exp Cell Res. 2015;333: 155-63. doi:
10.1016/j.yexcr.2015.01.018.

103. Sayed M, Pelech S, Wong C, Marotta A, Salh B. Protein kinase CK2 is involved
in G2 arrest and apoptosis following spindle damage in epithelial cells. Oncogene.
2001;20: 6994-7005.

104. Maddigan A, Truitt L, Arsenault R, Freywald T, Allonby O, Dean J, Narendran
A, Xiang J, Weng A, Napper S, Freywald A. EphB receptors trigger Akt activation

and suppress Fas receptor-induced apoptosis in malignant T lymphocytes. J Immunol.
2011;187: 5983-94. doi: 10.4049/jimmunol.1003482.

105. Okamoto S, Krainc D, Sherman K, Lipton SA. Antiapoptotic role of the p38
mitogen-activated protein kinase-myocyte enhancer factor 2 transcription factor
pathway during neuronal differentiation. Proc Natl Acad Sci USA. 2000;97: 7561-6.

106. Chiorazzi M, Rui L, Yang Y, Ceribelli M, Tishbi N, Maurer CW, Ranuncolo SM,
Zhao H, Xu W, Chan WC, Jaffe ES, Gascoyne RD, Campo E, Rosenwald A, Ott G,
Delabie J, Rimsza LM, Shaham S, Staudt LM. Related F-box proteins control cell
death in Caenorhabditis elegans and human lymphoma. Proc Natl Acad Sci USA.
2013;110: 3943-8. doi: 10.1073/pnas.1217271110.

107. Bloch DB, Nakajima A, Gulick T, Chiche JD, Orth D, de La Monte SM, Bloch
KD. Sp110 localizes to the PML-Sp100 nuclear body and may function as a nuclear
hormone receptor transcriptional coactivator. Mol Cell Biol. 2000;20: 6138-46.

108. Bengoechea-Alonso MT, Ericsson J. Tumor suppressor Fbxw7 regulates TGFβ
signaling by targeting TGIF1 for degradation. Oncogene. 2010;29: 5322-8. doi:
10.1038/onc.2010.278. Epub 2010 Jul 12.

109. Ma Y, Wang B, Li W, Ying G, Fu L, Niu R, Gu F. Reduction of intersectin1-s
induced apoptosis of human glioblastoma cells. Brain Res. 2010;1351: 222-8. doi:
10.1016/j.brainres.2010.05.028.

110. Gnesutta N, Qu J, Minden A. The serine/threonine kinase PAK4 prevents caspase
activation and protects cells from apoptosis. J Biol Chem. 2001;276: 14414-9.

111. Markovich D. Physiological roles and regulation of mammalian sulfate
transporters. Physiol Rev. 2001;81: 1499-533.

112. Guasti L, Crociani O, Redaelli E, Pillozzi S, Polvani S, Masselli M, Mello T,
Galli A, Amedei A, Wymore RS, Wanke E, Arcangeli A. Identification of a
posttranslational mechanism for the regulation of hERG1 K+ channel expression and
hERG1 current density in tumor cells. Mol Cell Biol. 2008;28: 5043-60. doi:
10.1128/MCB.00304-08.

113. Bremser M, Nickel W, Schweikert M, Ravazzola M, Amherdt M, Hughes CA,
Sllner TH, Rothman JE, Wieland FT. Coupling of coat assembly and vesicle budding
to packaging of putative cargo receptors. Cell. 1999;96: 495-506.

114. Zhen Y, Srensen V, Skjerpen CS, Haugsten EM, Jin Y, Wlchli S, Olsnes S,
Wiedlocha A. Nuclear import of exogenous FGF1 requires the ER-protein LRRC59
and the importins Kpnα1 and Kpnβ1. Traffic. 2012;13: 650-64. doi: 10.1111/j.1600-
0854.2012.01341.x.

115. Bangs P, Burke B, Powers C, Craig R, Purohit A, Doxsey S. Functional analysis
of Tpr: identification of nuclear pore complex association and nuclear localization
domains and a role in mRNA export. J Cell Biol. 1998;143: 1801-12.

116. Nakielny S, Siomi MC, Siomi H, Michael WM, Pollard V, Dreyfuss G.
Transportin: nuclear transport receptor of a novel nuclear protein import pathway.
Exp Cell Res. 1996;229: 261-6.

117.   Chi NC, Adam EJ, Adam SA. Sequence and characterization of cytoplasmic
nuclear protein import factor p97. J Cell Biol. 1995;130: 265-74.

118.   Tang BL, Low DY, Tan AE, Hong W. Syntaxin 10: a member of the syntaxin
family localized to the trans-Golgi network. Biochem Biophys Res Commun.
1998;242: 345-50.

119.   Chen G. Biochemical properties of urea transporters. Subcell Biochem. 2014;73:
109-26. doi: 10.1007/978-94-017-9343-8_7. PMID: 25298341

120.   Palmieri L, Pardo B, Lasorsa FM, del Arco A, Kobayashi K, Iijima M, Runswick
MJ, Walker JE, Saheki T, Satrstegui J, Palmieri F. Citrin and aralar1 are Ca(2+)-
stimulated aspartate/glutamate transporters in mitochondria. EMBO J. 2001;20: 5060-
9.

121.   Hatanaka T, Huang W, Ling R, Prasad PD, Sugawara M, Leibach FH, Ganapathy
V. Evidence for the transport of neutral as well as cationic amino acids by ATA3, a
novel and liver-specific subtype of amino acid transport system A. Biochim Biophys
Acta. 2001;1510: 10-7.

122.   Conchon S, Cao X, Barlowe C, Pelham HR. Got1p and Sft2p: membrane proteins
involved in traffic to the Golgi complex. EMBO J. 1999;18: 3934-46.

123.   Hautbergue GM, Hung ML, Walsh MJ, Snijders AP, Chang CT, Jones R, Ponting
CP, Dickman MJ, Wilson SA. UIF, a New mRNA export adaptor that works together
with REF/ALY, requires FACT for recruitment to mRNA. Curr Biol. 2009;19: 1918-
24. doi: 10.1016/j.cub.2009.09.041.

124.   Grempler R, Augustin R, Froehner S, Hildebrandt T, Simon E, Mark M,
Eickelmann P. Functional characterisation of human SGLT-5 as a novel kidney-
specific sodium-dependent sugar transporter. FEBS Lett. 2012;586: 248-53. doi:
10.1016/j.febslet.2011.12.027.

125.   Dietmeier K, Hönlinger A, Bömer U, Dekker PJ, Eckerskorn C, Lottspeich F,
Kübrich M, Pfanner N. Tom5 functionally links mitochondrial preprotein receptors to
the general import pore. Nature. 1997;388: 195-200.

126.   Annilo T, Shulenin S, Chen ZQ, Arnould I, Prades C, Lemoine C, Maintoux-
Larois C, Devaud C, Dean M, Denfle P, Rosier M. Identification and characterization
of a novel ABCA subfamily member, ABCA12, located in the lamellar ichthyosis
region on 2q34. Cytogenet Genome Res. 2002;98: 169-76.

127.   Moraes LA, Piqueras L, Bishop-Bailey D. Peroxisome proliferator-activated
receptors and inflammation. Pharmacol Ther. 2006;110: 371-85.

128.   Allikmets R, Gerrard B, Hutchinson A, Dean M. Characterization of the human
ABC superfamily: isolation and mapping of 21 new genes using the expressed
sequence tags database. Hum Mol Genet. 1996;5: 1649-55.

129.   Chew CS, Parente JA Jr, Chen X, Chaponnier C, Cameron RS. The LIM and SH3
domain-containing protein, lasp-1, may link the cAMP signaling pathway with
dynamic membrane restructuring activities in ion transporting epithelia. J Cell Sci.
2000;113: 2035-45.

130.   Phillips-Krawczak CA, Singla A, Starokadomskyy P, Deng Z, Osborne DG, Li H,
Dick CJ, Gomez TS, Koenecke M, Zhang JS, Dai H, Sifuentes-Dominguez L2, Geng
LN, Kaufmann SH, Hein MY, Wallis M, McGaughran J, Gecz J, Sluis Bv, Billadeau
DD, Burstein E. COMMD1 is linked to the WASH complex and regulates endosomal
trafficking of the copper transporter ATP7A. Mol Biol Cell. 2015;26: 91-103. doi:
10.1091/mbc.E14-06-1073.

131.   Tyson JR, Stirling CJ. LHS1 and SIL1 provide a lumenal function that is essential
for protein translocation into the endoplasmic reticulum. EMBO J. 2000;19: 6440-52.

132.   Sirrenberg C, Bauer MF, Guiard B, Neupert W, Brunner M. Import of carrier
proteins into the mitochondrial inner membrane mediated by Tim22. Nature.
1996;384: 582-5.

133.   Rubino M, Miaczynska M, Lipp R, Zerial M. Selective membrane recruitment of
EEA1 suggests a role in directional transport of clathrin-coated vesicles to early
endosomes. J Biol Chem. 2000;275: 3745-8.

134.   Zhu XR, Netzer R, Bhlke K, Liu Q, Pongs O. Structural and functional
characterization of Kv6.2 a new gamma-subunit of voltage-gated potassium channel.
Receptors Channels. 1999;6: 337-50.

135.   Toyama R, Kobayashi M, Tomita T, Dawid IB. Expression of LIM-domain
binding protein (ldb) genes during zebrafish embryogenesis. Mech Dev. 1998;71:
197-200.

136.   Kida YS, Sato T, Miyasaka KY, Suto A, Ogura T. Daam1 regulates the
endocytosis of EphB during the convergent extension of the zebrafish notochord.
Proc Natl Acad Sci USA. 2007;104: 6708-13.

137.   Gilpin BJ, Loechel F, Mattei MG, Engvall E, Albrechtsen R, Wewer UM. A
novel, secreted form of human ADAM 12 (meltrin alpha) provokes myogenesis in
vivo. J Biol Chem. 1998;273: 157-66.

138.   Sano K, Inohaya K, Kawaguchi M, Yoshizaki N, Iuchi I, Yasumasu S.
Purification and characterization of zebrafish hatching enzyme - an evolutionary
aspect of the mechanism of egg envelope digestion. FEBS J. 2008;275: 5934-46. doi:
10.1111/j.1742-4658.2008.06722.x.

139.   Hofmeister-Brix A, Kollmann K, Langer S, Schultz J, Lenzen S, Baltrusch S.
Identification of the ubiquitin-like domain of midnolin as a new glucokinase
interaction partner. J Biol Chem. 2013;288: 35824-39. doi:
10.1074/jbc.M113.526632.

140.   Tsukahara M, Suemori H, Noguchi S, Ji ZS, Tsunoo H. Novel nucleolar protein,
midnolin, is expressed in the mesencephalon during mouse development. Gene.
2000;254: 45-55.

141.   Hong SK, Dawid IB. FGF-dependent left-right asymmetry patterning in zebrafish
is mediated by Ier2 and Fibp1. Proc Natl Acad Sci USA. 2009;106: 2230-5. doi:
10.1073/pnas.0812880106.

142.   Kwong RW, Perry SF. The tight junction protein claudin-b regulates epithelial
1537   permeability and sodium handling in larval zebrafish, Danio rerio. Am J Physiol
1538   Regul Integr Comp Physiol 2013;304: R504-13. doi: 10.1152/ajpregu.00385.2012.

1539   143.   Ing T, Aoki Y. Expression of RGS2, RGS4 and RGS7 in the developing postnatal
1540   brain. Eur J Neurosci. 2002;15: 929-36.

1541   144.   Wu T, Patel H, Mukai S, Melino C, Garg R, Ni X, Chang J, Peng C. Activin,
1542   inhibin, and follistatin in zebrafish ovary: expression and role in oocyte maturation.
1543   Biol Reprod. 2000;62: 1585-92.

1544   145.   Malinauskas T, Aricescu AR, Lu W, Siebold C, Jones EY. Modular mechanism
1545   of Wnt signaling inhibition by Wnt inhibitory factor 1. Nat Struct Mol Biol. 2011;18:
1546   886-93. doi: 10.1038/nsmb.2081.

1547   146.   Zagon IS, Wu Y, McLaughlin PJ. Opioid growth factor and organ development in
1548   rat and human embryos. Brain Res. 1999;839: 313-22.

1549   147.   Takano A, Zochi R, Hibi M, Terashima T, Katsuyama Y. Expression of
1550   strawberry notch family genes during zebrafish embryogenesis. Dev Dyn. 2010;239:
1551   1789-96. doi: 10.1002/dvdy.22287.

1552   148.   Pierani A, Moran-Rivard L, Sunshine MJ, Littman DR, Goulding M, Jessell TM.
1553   Control of interneuron fate in the developing spinal cord by the progenitor
1554   homeodomain protein Dbx1. Neuron. 2001;29: 367-84.

1555   149.   Buonamici S, Chakraborty S, Senyuk V, Nucifora G. The role of EVI1 in normal
1556   and leukemic cells. Blood Cells Mol Dis. 2003;31: 206-12.

1557   150.   Litwack ED, Babey R, Buser R, Gesemann M, O'Leary DD. Identification and
1558   characterization of two novel brain-derived immunoglobulin superfamily members
1559   with a unique structural organization. Mol Cell Neurosci. 2004;25: 263-74.

1560   151.   Gazit R, Mandal PK, Ebina W, Ben-Zvi A, Nombela-Arrieta C, Silberstein LE,
1561   Rossi DJ. Fgd5 identifies hematopoietic stem cells in the murine bone marrow. J Exp
1562   Med. 2014; 211: 1315-31. doi: 10.1084/jem.20130428.

1563   152.   Zhang J, Tomasini AJ, Mayer AN. RBM19 is essential for preimplantation
1564   development in the mouse. BMC Dev Biol. 2008;8: 115. doi: 10.1186/1471-213X-8-
1565   115.

1566   153.   Tang K, Xie X, Park JI, Jamrich M, Tsai S, Tsai MJ. COUP-TFs regulate eye
1567   development by controlling factors essential for optic vesicle morphogenesis.
1568   Development. 2010;137: 725-34. doi: 10.1242/dev.040568.

1569   154.   Zhang P, Bennoun M, Gogard C, Bossard P, Leclerc I, Kahn A, Vasseur-Cognet
1570   M. Expression of COUP-TFII in metabolic tissues during development. Mech Dev.
1571   2002;119: 109-14.

1572   155.   Coumailleau P, Duprez D. Sim1 and Sim2 expression during chick and mouse
1573   limb development. Int J Dev Biol. 2009;53: 149-57. doi: 10.1387/ijdb.082659pc.

1574   156.   Kondou H, Kawai M, Tachikawa K, Kimoto A, Yamagata M, Koinuma T,
1575   Yamazaki M, Nakayama M, Mushiake S, Ozono K, Michigami T. Sodium-coupled

1576 neutral amino acid transporter 4 functions as a regulator of protein synthesis during
1577 liver development. Hepatol Res. 2013;43: 1211-23. doi: 10.1111/hepr.12069.

157. Li Z, Lai G, Deng L, Han Y, Zheng D, Song W. Association of SLC38A4 and
1578
1579 system A with abnormal fetal birth weight. Exp Ther Med. 2012;3: 309-313.

158. Tiberi L, van den Ameele J, Dimidschstein J, Piccirilli J, Gall D, Herpoel A,
1580
1581 Bilheu A, Bonnefont J, Iacovino M, Kyba M, Bouschet T, Vanderhaeghen P. BCL6
1582 controls neurogenesis through Sirt1-dependent epigenetic repression of selective
1583 Notch targets. Nat Neurosci. 2012;15: 1627-35. doi: 10.1038/nn.3264.

159. Matsuoka RL, Sun LO, Katayama K, Yoshida Y, Kolodkin AL. Sema6B,
1584
1585 Sema6C, and Sema6D expression and function during mammalian retinal
1586 development. PLoS One. 2013;8: e63207. doi: 10.1371/journal.pone.0063207.

160. Tamada H, Sakashita E, Shimazaki K, Ueno E, Hamamoto T, Kagawa Y, Endo H.
1587
1588 cDNA cloning and characterization of Drb1, a new member of RRM-type neural
1589 RNA-binding protein. Biochem Biophys Res Commun. 2002;297: 96-104.

161. Rempel RE, Saenz-Robles MT, Storms R, Morham S, Ishida S, Engel A, Jakoi L,
1590
1591 Melhem MF, Pipas JM, Smith C, Nevins JR. Loss of E2F4 activity leads to abnormal
1592 development of multiple cellular lineages. Mol Cell. 2000;6: 293-306.

162. Miyares RL, Stein C, Renisch B, Anderson JL, Hammerschmidt M, Farber SA.
1593
1594 Long-chain Acyl-CoA synthetase 4A regulates Smad activity and dorsoventral
1595 patterning in the zebrafish embryo. Dev Cell. 2013;27: 635-47. doi:
1596 10.1016/j.devcel.2013.11.011.

163. Louryan S, Biermans J, Flemal F. Nerve growth factor in the developing
1597
1598 craniofacial region of the mouse embryo. Eur J Morphol. 1995;33: 415-9.

164. Ko JA, Gondo T, Inagaki S, Inui M. Requirement of the transmembrane
1599
1600 semaphorin Sema4C for myogenic differentiation. FEBS Lett. 2005;579: 2236-42.

165. Wu H, Fan J, Zhu L, Liu S, Wu Y, Zhao T, Wu Y, Ding X, Fan W, Fan M.
1601
1602 Sema4C expression in neural stem/progenitor cells and in adult neurogenesis induced
1603 by cerebral ischemia. J Mol Neurosci. 2009;39: 27-39. doi: 10.1007/s12031-009-
1604 9177-8.

166. Pathirage NA, Cocquebert M, Sadovsky Y, Abumaree M, Manuelpillai U, Borg
1605
1606 A, Keogh RJ, Brennecke SP, Evain-Brion D, Fournier T, Kalionis B, Murthi P.
1607 Homeobox gene transforming growth factor β-induced factor-1 (TGIF-1) is a
1608 regulator of villous trophoblast differentiation and its expression is increased in
1609 human idiopathic fetal growth restriction. Mol Hum Reprod. 2013;19: 665-75. doi:
1610 10.1093/molehr/gat042.

167. Heger S, Mastronardi C, Dissen GA, Lomniczi A, Cabrera R, Roth CL, Jung H,
1611
1612 Galimi F, Sippell W, Ojeda SR. Enhanced at puberty 1 (EAP1) is a new
1613 transcriptional regulator of the female neuroendocrine reproductive axis. J Clin
1614 Invest. 2007;117: 2145-54.

168. Miao H, Strebhardt K, Pasquale EB, Shen TL, Guan JL, Wang B. Inhibition of
1615
1616 integrin-mediated cell adhesion but not directional cell migration requires catalytic

activity of EphB3 receptor tyrosine kinase. Role of Rho family small GTPases. J Biol Chem. 2005;280: 923-32.

169.   Vogel T, Boettger-Tong H, Nanda I, Dechend F, Agulnik AI, Bishop CE, Schmid M, Schmidtke J. A murine TSPY. Chromosome Res. 1998;6: 35-40.

170.   Janesick A, Shiotsugu J, Taketani M, Blumberg B. RIPPLY3 is a retinoic acid-inducible repressor required for setting the borders of the pre-placodal ectoderm. Development. 2012;139: 1213-24. doi: 10.1242/dev.071456.

171.   Zhou L, Zhang Z, Zheng Y, Zhu Y, Wei Z, Xu H, Tang Q, Kong X, Hu L. SKAP2, a novel target of HSF4b, associates with NCK2/F-actin at membrane ruffles and regulates actin reorganization in lens cell. J Cell Mol Med. 2011;15: 783-95. doi: 10.1111/j.1582-4934.2010.01048.x.

172.   Wittschieben J, Shivji MK, Lalani E, Jacobs MA, Marini F, Gearhart PJ, Rosewell I, Stamp G, Wood RD. Disruption of the developmentally regulated Rev3l gene causes embryonic lethality. Curr Biol. 2000;10: 1217-20.

173.   Selvaraj A, Prywes R. Megakaryoblastic leukemia-1/2, a transcriptional co-activator of serum response factor, is required for skeletal myogenic differentiation. J Biol Chem. 2003;278: 41977-87.

174.   O'Shaughnessy RF, Welti JC, Sully K, Byrne C. Akt-dependent Pp2a activity is required for epidermal barrier formation during late embryonic development. Development. 2009;136: 3423-31. doi: 10.1242/dev.037010.

175.   Jaenisch R, Bird A. Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. Nat Genet. 2003;33: 245-54.

176.   Gräff J, Kim D, Dobbin MM, Tsai LH. Epigenetic regulation of gene expression in physiological and pathological brain processes. Physiol Rev. 2011; 91: 603-49. doi: 10.1152/physrev.00012.2010.

177.   Pan J, Jin C, Murata T, Yokoyama KK. Histone modification activities of JDP2 associated with retinoic acid-induced differentiation of F9 cells. Nucleic Acids Symp Ser (Oxf). 2004;48: 189-90.

178.   Bowen NJ, Fujita N, Kajita M, Wade PA. Mi-2/NuRD: multiple complexes for many purposes. Biochim Biophys Acta. 2004;1677: 52-7.

179.   Sugimoto N, Maehara K, Yoshida K, Yasukouchi S, Osano S, Watanabe S, Aizawa M, Yugawa T, Kiyono T, Kurumizaka H, Ohkawa Y, Fujita M. Cdt1-binding protein GRWD1 is a novel histone-binding protein that facilitates MCM loading through its influence on chromatin architecture. Nucleic Acids Res. 2015;43: 5898-911. doi: 10.1093/nar/gkv509.

180.   Mariño-Ramírez L1, Kann MG, Shoemaker BA, Landsman D. Histone structure and nucleosome stability. Expert Rev Proteomics. 2005;2: 719-29.

181.   Yap KL, Li S, Muñoz-Cabello AM, Raguz S, Zeng L, Mujtaba S, Gil J, Walsh MJ, Zhou MM. Molecular interplay of the noncoding RNA ANRIL and methylated histone H3 lysine 27 by polycomb CBX7 in transcriptional silencing of INK4a. Mol Cell. 2010;38: 662-74. doi: 10.1016/j.molcel.2010.03.021.

182.   Ikegamii K, Horigome D, Mukai M, Livnat I, MacGregor GR, Setou M. TTLL10
is a protein polyglycylase that can modify nucleosome assembly protein 1. FEBS
Lett. 2008;582: 1129-34. doi: 10.1016/j.febslet.2008.02.079.

183.   Marzluff WF, Gongidi P, Woods KR, Jin J, Maltais LJ. The human and mouse
replication-dependent histone genes. Genomics. 2002;80: 487-98.

184.   Li Y, Wen H, Xi Y, Tanaka K, Wang H, Peng D, Ren Y, Jin Q, Dent SY, Li W,
Li H, Shi X. AF9 YEATS domain links histone acetylation to DOT1L-mediated
H3K79 methylation. Cell. 2014;159: 558-71. doi: 10.1016/j.cell.2014.09.049.

185.   Iizuka M, Stillman B. Histone acetyltransferase HBO1 interacts with the ORC1
subunit of the human initiator protein. J Biol Chem. 1999;274: 23027-34.

186.   Katoh M, Katoh M. Comparative integromics on JMJD1C gene encoding histone
demethylase: conserved POU5F1 binding site elucidating mechanism of JMJD1C
expression in undifferentiated ES cells and diffuse-type gastric cancer. Int J Oncol.
2007;31: 219-23.

187.   González-Herrera L, Valenzuela A, Marchal JA, Lorente JA, Villanueva E. Studies
on RNA integrity and gene expression in human myocardial tissue, pericardial fluid
and blood, and its postmortem stability. Forensic Sci Int. 2013; 232: 218-28.

188.   Coyle RC, Latimer A, Jessen JR. Membrane-type 1 matrix metalloproteinase
regulates cell migration during zebrafish gastrulation: evidence for an interaction
with non-canonical Wnt signaling. Exp Cell Res. 2008;314: 2150-62. doi:
10.1016/j.yexcr.2008.03.010.

189.   Gérard A, Patino-Lopez G, Beemiller P, Nambiar R, Ben-Aissa K, Liu Y, Totah FJ,
Tyska MJ, Shaw S, Krummel MF. Detection of rare antigen-presenting cells
through T cell-intrinsic meandering motility, mediated by *Myo1g*. Cell. 2014;15:
492-505. doi: 10.1016/j.cell.2014.05.044.

190.   Müller T, Rumpel E, Hradetzky S, Bollig F, Wegner H, Blumenthal A, Greinacher
A, Endlich K, Endlich N. Non-muscle myosin IIA is required for the development
of the zebrafish glomerulus. Kidney Int. 2011;80: 1055-63. doi:
10.1038/ki.2011.256.

191.   Ang Peng Hwa, Tekwani Shyam, Wang Guozhen. Shutting down the mobile phone
and the downfall of Nepalese society, economy and politics. Pacific Affairs. 2012,
85: 547-561.

192.   Sauer Hanno. Can't We All Disagree More Constructively? Moral Foundations,
Moral Reasoning, and Political Disagreement. Neuroethics. 2015, 8: 153-169.

193.   Pahwa S, Scoglio C, Scala A. Abruptness of cascade failures in power grids. Sci
Rep. 2014, 4: 3694. doi: 10.1038/srep03694.

194.   Mattson MP, Chan SL. Calcium orchestrates apoptosis. Nat Cell Biol. 2003, 5:
1041-3.

195.   Lang F, Föller M, Lang KS, Lang PA, Ritter M, Gulbins E, Vereninov A, Huber
SM. Ion channels in cell proliferation and apoptotic cell death. J Membr Biol. 2005,
205: 147-57.

196. Kauffman, Stuart A. The origins of order: self-organization and selection in
evolution. New York: Oxford University Press; 1993.

197. Tu Y,Stolovitzky G, Klein U.Quantitative noise analysis for gene expression
microarray experiments. Proc Natl Acad Sci USA. 2002, 99: 14031.

198. Jackson ES, Wayland MT, Fitzgerald W, Bahn S. A microarray data analysis
framework for postmortem tissues. Methods 2005, 37: 247-60.

199. Barash Y, Dehan E, Krupsky M, Franklin W, Geraci M, Friedman N, Kaminski N.
Comparative analysis of algorithms for signal quantitation from oligonucleotide
microarrays. Bioinform. 2004, 20: 839.

200. Seo J, Bakay M, Chen YW, Hilmer S, Shneiderman B, Hoffman EP. Interactively
optimizing signal-to-noise ratios in expression profiling: project-specific algorithm
selection and detection p-value weighting in Affymetrix microarrays. Bioinform.
2004, 20: 2534.

201. Harr B, Schlötterer C, Comparison of algorithms for the analysis of Affymetrix
microarray data as evaluated by co-expression of genes in known operons. Nucl
Acids Res. 2006, 34: e8.

202. Millenaar FF, Okyere J, May ST, van Zanten M, Voesenek LA, Peeters AJ. How to
decide? Different methods of calculating gene expression from short
oligonucleotide array data will give different results. BMC Bioinform. 2006, 7:
137.

203. Amend AS, Seifert KA, Bruns TD. Quantifying microbial communities with 454
pyrosequencing: does read abundance count? Mol Ecol. 2010, 19: 5555.

204. Aberg F, Pukkala E, Höckerstedt K, Sankila R, Isoniemi H. Risk of malignant
neoplasms after liver transplantation: a population-based study. Liver Transpl.
2008, 14: 1428-36.

205. Haagsma EB, Hagens VE, Schaapveld M, van den Berg AP, de Vries EG,
Klompmaker IJ, Slooff MJ, Jansen PL. Increased cancer risk after liver
transplantation: a population-based study. J Hepatol. 2001, 34: 84-91.

1727

Additional Files

Thanatotranscriptome: genes actively expressed after organismal death

Alex E. Pozhitkov, Rafik Neme, Tomislav Domazet-Lošo, Brian G. Leroux, Shivani Soni, Diethard Tautz and Peter A. Noble[*].

*Correspondence to: panoble@washington.edu

This PDF file includes:

Supplementary Text

Figs S1, S2 and S3

Table S1

Other Additional Files for this manuscript include the following:

Data files S1 to S8 as zipped archives:

File S1. MiceProbesParameters.txt

File S2. FishProbesParameters.txt

File S3. Mouse_liver_Log10_AllProfiles.txt

File S4. Mouse_brain_Log10_AllProfiles.txt

File S5. Fish_Log10_AllProfiles.txt

File S6. MiceProbesSeq.txt

File S7. FishProbesSeq.txt

File S8. Gene_annotation_lit_refs_v3.xls

SUPPLEMENTARY TEXT

Fig S1. Bioanalyzer results showing total mRNA from the zebrafish.

Fig S2. Bioanalyzer results showing total mRNA from the mouse.

Fig S3. Transcriptional profiles in the zebrafish (*Acer3* gene) and mouse (*Cdc42* and *Rbm45* genes) by postmortem time.

Table S1. Total mRNA extracted (ng/µl tissue extract) from zebrafish by time and replicate sample.

Table S2. Total mRNA extracted (ng/µl tissue extract) from mouse organ/tissue by time and replicate sample.

Table S3. % of global regulator genes and response genes.

Fig S1. Bioanalyzer results showing total mRNA from the zebrafish. Only one replicate per sampling time is shown. The dominant bands represent the 28S and 18S rRNAs.

Fig S2. Bioanalyzer results showing total mRNA from the mouse. All replicates per sampling time are shown. The dominant bands represent the 28S and 18S rRNAs.

Fig S3. Transcriptional profiles in the zebrafish (*Acer3* gene) and mouse (*Cdc42* and *Rbm45* genes) by postmortem time. Red arrows, up-regulation; blue arrows, down-regulation. One-way T-tests show significant differences between means. Results suggest that the differences in up- and down regulation by postmortem time are due to changes in regulation rather than changes in "residual transcription levels".

Table S1. Total mRNA extracted (ng/µl tissue extract) from zebrafish by time and replicate sample.

Time (h)	Repl#1	Repl#2
Live	1606	1648
0	1632	1481
0.25	1768	1573
0.5	1457	1481
1	1864	1514
4	1251	1651
8	1605	1499
12	1365	1422
24	1087	539
48	339	428
96	183	183

a, no replicate taken

Table S2. Total mRNA extracted (ng/µl tissue extract) from mouse organ/tissue by time and replicate sample.

Organ	Time (h)	Repl#1	Repl#2	Repl#3
Liver	0	432	387	443
Liver	0.5	533	651	569
Liver	1	421	426	601
Liver	6	541	624	528
Liver	12	839	450	845
Liver	24	1021	510	1066
Liver	48	1453	1197	-[a]
Brain	0	169	226	210
Brain	0.5	174	166	410
Brain	1	194	485	264
Brain	6	401	269	256
Brain	12	379	258	203
Brain	24	249	324	400
Brain	48	397	310	-

a, no replicate taken

Table S3. % of global regulator genes and response genes. Approx. 33% of all upregulated genes are involved in global regulation.

Global gene regulators and other response genes	Zebrafish	Mouse	Combined
Transcription factors and transcriptional regulatory genes	17	12	14
Cell signaling genes	24	16	19
Other response genes	59	73	67