Learning from critical care management of sheep receiving extra-corporeal membrane oxygenation

for smoke-induced acute lung injury as a tool for processing large clinical datasets

Chemonges Saul[1,2]

[1]Proteomics and Small Molecule Mass Spectrometry, Central Analytical Research Facility, Queensland

University of Technology, Gardens Point, Brisbane, QLD, Australia

[2]School of Veterinary Science, Building #8114, The University of Queensland, Gatton, QLD, Australia

Address correspondence to:

Chemonges Saul

Proteomics and Small Molecule Mass Spectrometry, Central Analytical Research Facility, Queensland

University of Technology, Level 6, P Block, Gardens Point Campus, 2 George Street, Brisbane QLD 4001,

Australia.

Telephone: +61 7 3138 9541; Fax: +61 7 3138 4438

E-mail: saul.chemonges@qut.edu.au

Abstract

Background: Many successful therapies developed for human medicine involve animal experimentation. With competition for public research funding and career advancement opportunities, animal studies focused on the translational potential may not sufficiently document unexpected outcomes. Such studies often have hastily developed methods with *ad hoc* modifications including the use of additional animals, leading to considerable amounts of idle, unprocessed data that could be used to advance veterinary science, or to refine the base animal model. Sheep, for example, are poorly understood models of intensive care and therefore, any experimental data arising from them should be interpreted with care. The hypothesis was that there is little information describing the development of methods of physiological data processing in multifaceted sheep models of intensive care and the author aimed to develop a suitable data processing method and to analyse the data, once processed.

Methods: Data from 19 adult mechanically ventilated ewes undergoing intensive care in a previous study evaluating a form of extracorporeal life support (treatment) for acute lung injury were used to develop a comprehensive method for processing manual and electronically gathered clinical observations. Eight sheep were injured by acute smoke inhalation prior to treatment (injured/treated), while another eight were not injured but treated (uninjured/treated). Two sheep were injured but not treated (injured/untreated), while one received room air instead of smoke as the injury, and was not treated (placebo/untreated). The data were then analysed for 11 physiological categories and compared between the two treated groups.

Results: The analysis revealed that compared with the baseline, treatment contributed to and exacerbated the deterioration of pulmonary pathology by reducing lung compliance and PaO_2/FiO_2 ratio. The oxygen extraction index changes mirrored those of the PaO_2/FiO_2 ratio. Decreasing coronary perfusion pressure predicted the severity of cardiopulmonary injury.

Conclusions: These novel observations could help in understanding similar pathology such as that which occurs in smoke inhalation animal victims of house and bush fires. A similar data processing method could be used when evaluating the effectiveness of other clinical interventions such as potentially reversible aspiration pneumonia secondary to tick paralysis in veterinary patients.

Keywords: Sheep, critical care, smoke-induced acute smoke inhalation injury, extra-corporeal life support, lung compliance, PaO_2/FiO_2 ratio

Background

During multifaceted experiments involving intensive care in large animal models in translational research,

information regarding animal monitoring is often collected with varying accuracy, scope, and end-user

applications. Data collection can be manual, electronic, or both [1-3]. Manually input data can include

subjectively scored end-points such as the plane of anaesthesia, and objective data such as heart rate or

breaths per minute. Depending on the goals of the study, some information may be used to validate or test

novel therapies, or to understand and refine existing treatments. In some cases, experimental information

may be gathered for scientific curiosity or for "classified" use, and outcomes may never be publicly available,

especially if the results are negative.

The source of data for this study was from a sheep model [2-4] being treated for smoke-induced

acute lung injury using veno-venous extracorporeal membrane oxygenation [2], a form of extracorporeal life

support (ECLS) developed to complement the treatment of acute lung injury in humans [5-7]. During this type

of ECLS, venous blood is carried from the patient to a gas exchange device where it becomes enriched with

oxygen, has carbon dioxide removed, and is returned to the patient's circulation in the right heart. This

method can be used for treatment, as respiratory support during lung transplantation, and in critically ill

patients with potentially reversible respiratory failure [7]. The multiple advanced cardiovascular [3],

respiratory, patient point-of-care procedures, and instrumentation associated with ECLS, even in animal

experimentation, is highly data- and equipment-intensive. This platform is useful for developing research and

methodological skills for *in vivo* animal instrumentation and the processing of large, real-time clinical data

sets from multifaceted animal studies that can be applied to similar intensive care scenarios. An opportunity

to develop these skills arose within the source study, which was an ongoing publicly funded animal

experimentation study (Queensland University of Technology Animal Ethics Approval No. 110000053). While

the objectives of the primary study had a separate focus, there were considerable amounts of redundant raw

data with potential use in veterinary science and other disciplines, once processed. The author hypothesised

that there was little information describing the development of methods of physiological data processing in

multifaceted large animal models of intensive care.

The overall goal was to provide useful information relevant to the sheep model, itself, and to those

interested in large animal experimentation and veterinary medicine, generally. The specific objectives were:

1) use the raw data from the sheep model study to create a data management system for tabulating large

data sets from human studies using animal models and, 2) analyse that data to provide biological information

that is not currently available for sheep receiving ECLS following smoke-induced acute lung injury.

Methods

The study was carried out at the purpose-built Medical Engineering Facility of Queensland University of

Technology (QUT-MERF) at the Prince Charles Hospital Campus of The University of Queensland [1]. In the

original study, sheep inhaled standardised cotton smoke generated by a device that combusts material in an

oxygen-deficient environment as previously described [8]. Briefly, 8 g of cotton towelling was combusted in a

chamber with transparent walls and 400 ml tidal volume. One tidal volume breath (approximately 10–12

ml/kg) of the smoke was delivered to the sheep via plastic tubing connected to a 1-m-long tracheostomy

tube. A fixed number (12) of breaths were given with each load of cotton over a period of approximately one

minute. Serial arterial blood gas samples were taken to assess the effect of smoke inhalation, starting at a

predetermined time point after the smoke breath cycles.

Physiological data management system

Raw data were obtained from a previous translational study of critical care monitoring of sheep undergoing

treatment for smoke-induced acute lung injury involving several separate projects. Data were collected prior

to 23 August 2013 and were obtained from two of the scientists who developed the base model [4] as part of

a Research Higher Degree project of the author [9, 10]. All data files were in Microsoft® Excel 97–2003

(Microsoft Corporation, Redmond, WA, USA) format and were grouped per sheep and date of the

experiment. Data consisted of separate files of real-time physiological data recorded on the hard drives of

the monitoring devices (electronically acquired data), and parameters manually recorded by those monitoring

the sheep under anaesthesia (manually acquired data), which included data from the electronic monitoring

equipment, as back-up if the electronic monitor malfunctioned.

The source study involved 64 sheep, comprising eight experimental groups of eight sheep based on

the study's multiple objectives, subsequent modifications, and later addition of experimental controls. The

experimental groups were classified based on: the duration of treatment (2 and 24 hours; E2H and E24H);

treatment after smoke inhalation (injury) for 2 and 24 hours (SE2H and SE24H); and treatment after smoke

inhalation and transfusion with fresh or stored (aged) blood (SEF24H or SEA24H), respectively. Two

additional groups included one group receiving smoke inhalation injury but no treatment (SC24H), and

another group that inhaled room air only as the injury (placebo) and no treatment (C24H). Data from sheep

involved in the treatment and transfusion studies were not included in the analysis in this study because

these data were beyond the scope of this study. Nineteen sheep were included in the present study; data

were analysed for 16 sheep with robust data (E24H and SE24H), and included, but not analysed, for three

from groups SC24H and C24H (early observational data). A systematic approach was developed for

processing the data.

Manually acquired physiological data workflow

A clone of the master manual data entry spreadsheet was created by removing the formatting and formulas.

Several members of the sheep ECMO research team inspected data repeatedly for errors to ensure that all

columns, rows, time points, and data points had been copied correctly, including number formats (Figure 1).

Redundant columns were removed and data were aligned to experimental time points (Figure 2). While

maintaining the same experimental time point header, data from the table in Figure 2 were split and grouped

into the following categories: ventilator settings, blood pressure and haemodynamics, fluids and urine output,

arterial blood gas values, activated clotting time, anaesthetics, anticoagulants, and ECLS circuit

observations.

Electronically acquired physiological data workflow

Raw electronically acquired physiologic monitoring data were inspected for completeness. The data

comprised 36 time points: ECLS pump time (min); time of day (h); electrocardiograph (heart rate); arterial

blood pressure (mean, systolic, diastolic, heart rate); central venous pressure (mean); pulmonary artery

pressure (mean, systolic, diastolic); oxygenator pressure (pre- and post-); capnography (end-tidal carbon

dioxide (etCO$_2$), respiratory rate); pulse oximetry (SpO$_2$, heart rate); ECLS pump (flow rate, speed); ventilator

(mode, frequency, oxygen, pressure control, inspiratory volume, expiratory volume, expiratory minute

volume, pressure maximum, mean pressure, positive end-expiratory pressure, plateau pressure, inspiratory

resistance, expiratory pressure, pulmonary compliance, inspiratory flow); mixed venous oxygen saturation

(SvO$_2$); and continuous cardiac output (CCO) (Figure 3). The yellow line in Figure 3 indicates the baseline

time point and the grey line represents the smoke inhalation time point. It is important to note that there may

or may not have been any data at any given point in time.

The electronically acquired physiological monitoring data were inspected for errors and cleaned to provide

data for downstream analysis (Figure 4).

Pre-data analysis checks

Data were then subjected to further integrity checks. An important step was to make a plot of data versus

time together with descriptive statistics for all data points in the grouped data. At the time of data processing,

the "Descriptive Statistics" tool of Microsoft® Excel 2010 (Microsoft Corporation) did not complete analysis

with missing values. Therefore, the data to be analysed were selected and then the "GoTo" tool (F5) was

used, and "Special", "Blanks", and thereafter, "OK" were selected to identify blanks. The blanks were deleted

by positioning the cursor in the blank cell and using the space bar to clear the cell (the "Delete" or

"Backspace" keys did not remove the blanks).

After artefact removal and integrity checks, data for individual sheep were placed into six categories:

activated clotting time; anaesthetics + inotropes and anticoagulants; arterial blood gas values; blood

pressure + ventilation and haemodynamic data; calculated respiratory + haemodynamic variables; and fluids

and urine production. Using specially written macros, data were extracted from each experiment and

grouped by parameter corresponding to experimental time points. All sheep treatment data were filed by

parameter.

Data from the 19 sheep from groups E24H, SE24H, C24H and SC24H (Figure 5) were processed

further. Data integrity checks were again performed and repeated by several sheep ECMO research team

members, and data were compiled as shown in Figure 6 (note the obliterated cells after removal of data

artefacts). The treatment timeline (a) comprised 22 time points for all experiments where sheep received

smoke inhalation acute lung injury (SE24H) (b). A trend plot (c) and descriptive statistics panel (d) were

useful in data quality control processes for suitability for downstream data analysis and end-user

applications.

Statistical methods

To meet the second objective, data from the groups, uninjured/treated and injured/treated groups were

analysed. The means, medians and standard deviations of the weights of the sheep, where applicable, were

tabulated. The physiological parameters of the groups were charted and compared against each other using

one-way analysis of variance (ANOVA), where appropriate. Parameters between groups were compared

using a paired two-tailed t-test. All p-values were two-sided and $p < 0.05$ was considered statistically

significant. All statistical calculations were performed using GraphPad PRISM 6 software (GraphPad

Software, La Jolla, CA, USA).

Results

The biodata of the sheep that were used in the current analysis are presented in Table 1. The weights of the

uninjured/treated sheep, unlike the injured/treated group, did not pass the D'Agostino–Pearson omnibus

normality test; however, there was no significant difference in the weights of the sheep between groups.

Ventilation

All animals were intubated and received mechanical ventilation as previously described [1]. Briefly, the initial

ventilator tidal volume was set to approximately 10 mL/kg with a respiratory rate of 15 breaths/min, positive

end expiratory pressure (PEEP) of 5 cm H_2O, and an initial F_iO_2 (fraction of inspired oxygen) of 1.0. These

settings were then titrated based on arterial blood gas results. A low tidal volume, high PEEP strategy was

used to minimise ventilator-induced lung injury.

Pulmonary compliance decreased in all of the sheep during the course of the experiments, with the

injured/treated (SE24H) animals having the most severe and drastic decrease followed by the

uninjured/treated (E24H), injured/untreated (SC24) and placebo/untreated (C24) sheep in that order (Figure

7). There was a significant difference ($p = 0.0013$) in pulmonary compliance between uninjured/treated and

injured/treated groups. The injured/treated sheep had consistently lower SpO_2 compared with the other

groups, but there was no significant difference in SpO_2 readings between the groups. There was an initial

increase in $etCO_2$ followed by a rapid decrease that lessened 15 minutes after the start of treatment. The

$etCO_2$ of the injured sheep continued to trend downward and plateaued in the uninjured groups. There was a

significant difference ($p = 0.0147$) in $etCO_2$ between the uninjured/treated and injured/treated groups.

Blood gases (arterial samples)

Blood pH varied between the groups. The placebo/untreated sheep had the highest pH while the

injured/treated group had the lowest. There was a significant difference in pH between the uninjured/treated

and injured/treated groups ($p = 0.0343$).

The pCO_2 in all but the uninjured/treated sheep increased initially before plummeting sharply,

forming a shallow trough corresponding to 1 hour after the start of treatment, followed by a slight increase

before stabilising in all sheep.

There was a gradual decrease in pO_2 in the treated groups of sheep from baseline before

decreasing dramatically at the start of treatment with the injured sheep having the most profound decrease

(Figure 8). However, there was no significant difference in pO_2 between the uninjured/treated and

injured/treated groups.

Haemoglobin [Hb] concentration decreased slightly from baseline before gradually increasing in the

injured sheep and remained relatively constant over time in the uninjured sheep. There was a significant

difference in [Hb] between the uninjured/treated and injured/treated ($p = 0.0131$) groups.

The fraction of oxyhaemoglobin (FO_2Hb) decreased sharply with the lowest reading at 5 minutes

post-injury before returning to near baseline levels within an hour of starting treatment. The injured/treated

sheep had a considerably deeper trough in FO_2Hb level and there was a significant difference ($p = 0.046$)

between troughs. There was no change in FO_2Hb for the uninjured sheep.

The fraction of carboxyhaemoglobin (FCOHb) increased sharply from baseline, peaking at

approximately 5 minutes post-injury and decreased sharply thereafter to the start of treatment before

gradually returning to near baseline levels at approximately 6 hours post-start of treatment in the injured

sheep. The injured/treated sheep had a higher peak FCOHb than the injured/untreated sheep, although the

difference was not significant. There was no change in FCOHb for the uninjured sheep.

The fraction of methaemoglobin (MetHb) increased gradually from baseline, peaking at

approximately 5 minutes post-injury and then gradually decreased to the start of treatment. This was

followed by a gradual return to near baseline levels at approximately 6 hours post-start of treatment in the

injured/treated sheep. There was no change in MetHb for the uninjured sheep. There was an initial subtle

decrease in calculated haematocrit (Hct) before a steady increase in the injured sheep and relatively flat

slopes for the uninjured sheep.

Electrolytes

The blood sodium concentration [Na^+] was relatively stable and there were no significant differences

between groups.

There was an initial decrease in the blood calcium [Ca^{2+}] level, with the lowest point at approximately

1 hour past the start of treatment before levelling out thereafter in all groups. There was a significant

difference in [Ca^{2+}] between the uninjured/treated and the injured/treated groups ($p = 0.0001$). The

placebo/untreated and injured/treated groups maintained the highest and lowest levels of [Ca^{2+}],

respectively, throughout the experiments.

Blood chloride [Cl^-] levels remained stable compared with baseline levels during the initial stages

and then increased gradually thereafter.

The blood potassium concentration [K^+] initially decreased compared with baseline levels, reaching a

minimum concentration 1 hour after starting treatment and then gradually increased with a peak at

approximately 12 hours post-treatment start in all experimental groups. Although the injured/untreated and

injured/treated sheep had higher [K^+] than the uninjured sheep, the differences were not significant.

Overall, the anion gap decreased gradually, reaching a relatively gentle slope at approximately 6

hours from the start of treatment and did not change significantly, thereafter. There was a gradual decrease

in anion gap from baseline during the course of the experiments and there was no significant difference in

anion gap between the uninjured/treated and injured/treated groups.

Metabolites

Although there was an increase in blood glucose level [Glu] for the injured/treated sheep after 6 hours of

treatment, the change was not significant.

There was an initial decrease in lactate levels [Lac] 6 hours after the start of treatment, followed by a

gradual increase for the injured sheep, especially for the injured/treated group. There was no significant

difference in [Lac] between the treated groups.

Acid-base balance

There was an increase in the blood base levels [Base (ecf)] that peaked 1 hour post-treatment, followed by a

gradual decrease in the untreated group. [Base (ecf)] in the treated groups remained at baseline levels to 1

hour post-start of treatment, before decreasing markedly in the injured/treated sheep. There was a significant

difference ($p = 0.0257$) in [Base (ecf)] between the uninjured/treated and injured/treated groups.

Blood bicarbonate concentrations [HCO_3^-] increased initially in the untreated groups before

decreasing gradually; however, levels remained higher compared with the treated sheep.

Haemodynamics

There was a gradual decrease in heart rate (HR) during the course of the experiments, with the

placebo/untreated groups maintaining a higher HR compared with the injured/untreated, injured/treated, and

uninjured/treated groups early in the experiments. There was no significant difference in HR between the

uninjured/treated and injured/treated groups.

The mean arterial blood pressure (MAP) decreased early in the experiments before subsequently

increasing gradually, peaking at approximately the treatment start time point, before gradually decreasing

again in all but the placebo/untreated sheep. The injured/treated groups had a consistently lower MAP

compared with the other groups and there was a significant difference in MAP ($p = 0.0058$) between the

uninjured/treated and injured/treated groups.

The mean pulmonary artery pressure (MPAP) increased gradually, with the injured/treated group

having a consistently higher MPAP. There was no significant difference in MPAP between the

uninjured/treated and injured/treated groups.

There was an initial, subtle increase in central venous pressure (CVP) that peaked at approximately

1 hour post-injury followed by a decrease that stabilised at approximately 1 hour post-start of treatment. CVP

levels in the injured/treated and placebo/untreated sheep were consistently higher and lower, respectively,

during the course of the experiments. Mixed venous oxygen saturation (SvO_2) had a lower baseline before

eventually rising to a relatively stable and higher level for the treated sheep, and a slightly lower level for the

untreated sheep. The injured/untreated sheep maintained a consistently lower SvO_2 compared with the other

groups.

Except for the placebo/untreated group, there was a decrease in continuous cardiac output (CCO)

from baseline to approximately 1 hour post-start of treatment. There was a significant difference ($p = 0.0009$)

in CCO between the uninjured/treated and injured/treated groups with CCO in the treated groups increasing

sharply before plateauing, especially in the uninjured/treated group. There was also a subsequent gradual

decrease in CCO in the injured/treated group.

Stroke volume (SV) began to increase 1 hour from the start of treatment for all groups, except for the

injured/untreated group where levels remained relatively constant. SV in the injured/treated group began to

decrease after 6 hours of treatment, while SV in the uninjured/treated and placebo/untreated sheep

increased steadily before decreasing or levelling out after 12 hours or more of treatment. There were no

significant differences in SV between.

Stroke volume index (SVI) began to increase 1 hour after of the start of treatment for all groups,

except for the injured/untreated group, for which SVI remained relatively constant. SVI in the injured/treated

group began to decrease after 6 hours of treatment while SVI in the uninjured/treated and placebo/untreated

groups increased before subsequently decreasing or levelling out after 12 hours or more of treatment. There

were no significant differences in SVI between groups.

While the cardiac index (CI) of the uninjured/treated and placebo/untreated groups remained

relatively close to baseline levels, CI the injured/treated and injured/untreated groups declined gradually over

the course of the experiments.

After an initial increase in systemic vascular resistance index (SVRI) to approximately 1 hour after

the start of treatment, SVRI began to decrease in all experimental groups before plateauing after 12 hours of

treatment followed by a gentle increasing trend until the end of the experiments. SVRI in the injured/treated

group was consistently below that of the other groups during treatment while that of the injured/untreated

group was correspondingly higher. There was no significant difference in SVRI between the groups.

Pulmonary vascular resistance index (PVRI) remained close to baseline levels for all of the groups

until 1 hour after the start of treatment when that of the injured groups progressively increased while that of

the uninjured groups remained lower with a subtle decrease to 6 hours post-treatment. PVRI in the

placebo/untreated sheep remained near baseline levels and the lowest throughout the course of the

experiment.

After a small peak corresponding to the start of treatment, right ventricular stroke work index

(RVSWI) in the uninjured sheep gradually increased while that of the injured sheep decreased. There was a

significant difference ($p = 0.0196$) in RVSWI gap between the uninjured/treated and injured/treated groups.

RVSWI in the placebo/untreated group remained high while that of the injured/treated group was consistently

the lowest.

Left ventricular stroke work index (LVSWI) gradually increased in the uninjured/treated and

placebo/untreated groups, before plateauing after 12 and 18 hours of treatment, respectively, while LVSWI in

the injured/untreated and injured/treated groups of sheep decreased before plateauing at 12 hours and

trending upward after 18 hours of treatment. LVSWI in the placebo/untreated group remained consistently

higher than in the other groups while that of the injured/treated group was consistently the lowest.

Following a decrease in the coronary perfusion pressure (CPP) from baseline in the smoke-injured

sheep, there was a subsequent increase in this parameter prior to a sustained decrease at 18 hours of

treatment, followed by another increase. There was a significant difference in CPP ($p = 0.0018$) between the

uninjured/treated and injured/treated groups and CCP in the placebo/untreated sheep remained relatively

stable after an initial, subtle increase.

There was an initial subtle decrease in arterial oxygen content (C_aO_2) from baseline in all groups

before a sustained increase in the injured/untreated group, a steady level in the placebo/untreated sheep,

and a sharp trough in the injured/treated and uninjured/treated groups. Following the trough, the C_aO_2 of the

injured/treated group gradually returned to baseline levels while that of the uninjured/treated group continued

on a downward trend. There was a significant difference ($p < 0.0085$) in C_aO_2 between the uninjured/treated

and injured/treated groups.

There was a slight decrease in the oxygen delivery index (DO_2I) in all groups to 1 hour of treatment

before a further marked decrease, except for the placebo/untreated sheep. There was a significant

difference ($p = 0.0013$) in DO_2I between the uninjured/treated and injured/treated groups. The injured/treated

group had the lowest DO_2I compared with the other groups while the placebo/untreated sheep maintained

the highest DO_2I profile.

The oxygen extraction index (O_2EI) decreased in all groups before plateauing at approximately 6

hours after the start of treatment. There was a significant difference ($p = 0.0247$) in O_2EI between the

injured/treated and uninjured/treated groups. O_2EI in the injured/treated and injured/untreated groups was

consistently lower and higher, respectively, compared with those of the other groups.

Fluid input and urine output

The volume of intravenous fluids administered to sheep in the different experimental groups varied. The

injured/treated sheep had the highest fluid requirements while the placebo/untreated sheep required the

least. There was a significant difference ($p < 0.0001$) in fluid requirements between uninjured/treated and

injured/treated sheep.

The injured/untreated and injured/treated groups produced the least and most urine on average,

respectively. There was no significant difference in urine output between the uninjured/treated and

injured/treated groups.

Anaesthetics

There was a significant difference ($p < 0.0001$) in the amount of alfaxalone required between the

uninjured/treated and injured/treated groups. The uninjured/treated group required more alfaxalone on

average and the injured/untreated group required the least amount on average. Ketamine requirements

differed between groups, with the injured/untreated group requiring the highest amount on average and the

injured/treated group requiring the least. There was no significant difference in the quantities of ketamine

required between the uninjured/treated and injured/treated groups but significant differences in midazolam

requirements occurred between the uninjured/treated and injured/treated groups ($p = 0.0067$).

Anticoagulation

There were no significant differences in heparin infusion doses between the uninjured/treated and

injured/treated sheep. Heparin requirements for the placebo/untreated group were the lowest. Activated

clotting time increased sharply from baseline during pre-treatment and peaked 1 hour after the start of

treatment before decreasing sharply and plateauing. There were no significant differences in activated

clotting time between groups.

ECLS circuit observations

There were significant differences in the ECLS pump speed, blood flow, and pressure differential between

the uninjured/treated and injured/treated groups. Pump speed, blood flow, and pressure differential were

significantly different (*p = 0.0022*), (*p = 0.0095*) and (*p = 0.0041*), respectively, between the two groups

receiving ECLS. These parameters in the uninjured/treated group were consistently higher than those of the

injured/treated group.

Body temperature

Body temperature in the untreated groups gradually increased from baseline levels and plateaued at

approximately 6 hours after starting treatment, and remained higher than for the treated groups. There was

no significant difference in body temperature between the treated groups.

Inflammatory cells and cytokines

Data were available in abstract form on inflammatory cell infiltration into the lung tissue with a trend toward

increased lung injury in the sheep that inhaled smoke, showing damage to the bronchiolar lining and

infiltration of inflammatory cells [11–13].

Discussion

The results of this study agree with and confirm earlier preliminary observations that ECLS causes a

decrease in pulmonary compliance over time [9]. It was expected that the injured sheep would have

relatively lower SpO_2 readings compared with the other groups because of episodes of hypotension with

hypoxemia, which can affect pulse oximeter function [14]. The relatively low $etCO_2$ in the injured sheep

suggested that the sheep may have hyperventilated, the causes of which were evaluated with respect to

reactive oxygen species or superoxide dismutase activity by a team from the source study [15, 16].

The relatively low blood pH in the injured/treated sheep suggested that the sheep tended to

metabolic acidosis as the same group of animals also had low $etCO_2$. This also means that there was no

respiratory component contributing to the observed acidosis. The low pCO_2 in the uninjured/treated sheep

could be a result of hyperventilation and the high pCO_2 in the injured/untreated sheep suggested that CO_2

clearance was curtailed by injury.

The treatment of the sheep contributed to lung injury by causing deterioration of pO_2. The low pO_2

translated to low partial arterial oxygen pressure/inspired oxygen fraction (PaO_2/FiO_2) ratio, which was much

worse in the injured sheep. This finding showed that ECLS contributed to the deterioration of the PaO_2/FiO_2

ratio in the injured/treated group of sheep, a novel finding that was also unexpected in the primary study.

The relatively higher levels of [Hb] in the injured sheep suggested that these animals could have

been dehydrated secondary to excessive fluid loss from inflammation and increased vascular permeability

[17] despite intravenous fluid replacement. However, blood total protein and albumin levels, better predictors

of dehydration in sheep [18], were not measured.

The inverse decrease in FO_2Hb relative to FCOHb following smoke injury was expected and agreed

with other studies [17, 19, 20]. It has recently been demonstrated that FCOHb is not correlated to the degree

of lung injury [17]. The gradual decrease in MetHb was probably caused by the enzymatic activity of

methaemoglobin reductase [21] and the higher Hct observed in the injured sheep could have been due to

dehydration because HCT was measured by an automated method.

The $[Ca^{2+}]$ was lower than the published normal level of 2.4 mmol/L [22]. Stress associated with

yarding of the sheep and phosphorus imbalance in feed are the most likely suggested causes of low $[Ca^{2+}]$

[23]. Fasting the sheep for 24 hours prior to the experimental procedures could also have contributed to the

relatively low $[Ca^{2+}]$.

The increase in [Cl⁻] beyond the normal range of 105 –110 mmol/L [22] during the experiments

suggests that the sheep may have developed respiratory alkalosis. Hyperventilation or metabolic acidosis

resulting from sustained salivary loss of sodium bicarbonate that was more severe in the injured/treated

group may have played a role hyperchloraemia, because Cl⁻ is known to replace HCO_3^- in the latter's loss

[24, 25]. Baseline $[K^+]$ in all of the sheep was below the published normal range of 4–5 mmol/L [22] and this

relative hypokalaemia may have been related to low $[K^+]$ in the diet [26-28]. The normal anion gap with

decreased HCO_3^- confirmed the presence of hyperchloraemic acidosis in all but the placebo/untreated

sheep. The cause of the hyperchloraemia was likely the prolonged administration of 0.9% NaCl.

Although normal [Glu] in ruminants is usually lower than for other species, its relative progressive

increase in the injured sheep may have been related to stress and severe pain associated with injury or the

development of enterotoxaemia [29, 30]. The relative increase in [Lac] beyond the reported normal range of

1–2 mmol/L in the injured sheep suggested dehydration, trauma, and sepsis [31]. Sepsis, in particular, is a

concern with sub-optimal rumen function leading to loss of its buffer effect and increasing numbers of

anaerobic bacteria with prolonged hypomotility, such as occurs during long-duration anaesthesia. Therefore,

the increases in both [Glu] and [Lac] are consistent with severe injury.

The elevated [Base (ecf)] above +2 mmol/L for most of the first 12 hours in the placebo/untreated

and injured/untreated sheep suggested that the sheep were metabolically alkalotic [29] before returning to

normal levels. The relatively low [Base (ecf)] (less than −2 mmol/L) was consistent with HCO_3^- loss and the

tendency to metabolic acidosis [29] in the injured/treated sheep. The marked decrease in [HCO_3^-] in the

injured sheep was consistent with metabolic acidosis and was more severe in the injured/treated group,

suggesting that ECLS was a contributing factor.

The resting HR of sheep is 50–80 beats/min [22]. In a study that instrumented conscious sheep, the

baseline heart rate was registered as 106 ± 9 beats/min [32]. In the present report, all of the sheep had a

relatively high HR, suggesting that stress and pain were contributing factors. The gradual decrease in HR

during the course of the experiments was consistent with the effects of anaesthesia [22].

In sheep, a mean arterial pressure below 60 mmHg indicates inadequate tissue perfusion [22].

Although the MAP values in the injured sheep were lower than for the uninjured, MAP values were still within

the published normal value of 70 mmHg [22]; the magnitude of injury was again a predictor of how low the

MAP was. Another predictor for the severity of the injury was the mean pulmonary artery pressure, which

was highest for the injured/treated sheep. The baseline values for MPAP were higher than the 17 ± 1 mmHg

reported in another study using sheep [32]. The baseline CVP in all of the sheep in the present report was >

10 mmHg, which was much higher than the 5.5 ± 1.2 mmHg reported elsewhere [32] in instrumented

conscious sheep and a novel finding in this study. The severity of injury and treatment contributed to the

CVP elevations in this study.

There was a benefit of ECLS treatment for SvO_2 as it remained high for both the injured/treated and

uninjured treated groups. The consistently low SvO_2 in the injured/untreated group was expected because of

the slightly reduced cardiac output in this group; however, this level of SvO_2 was still higher than that

reported in other studies [32]. Smoke injury was associated with a sustained decrease in cardiac output in all

of the sheep that were exposed to smoke. As in CCO changes, the SV, SVI and CI all had similar profiles for

the different groups, with the injured sheep having lower values. The decrease in SVRI in all of the sheep

later in the experiments suggested that there was systemic vasodilation. In contrast, the increase in PVRI in

the injured sheep suggested that vasoconstriction was caused by exposure to smoke injury. The exposure to

smoke injury worsened both RVSWI and LVSWI while there was an increase in both parameters in the

uninjured sheep. Reduced RVSWI is associated with poor functioning of the right ventricle [33, 34] and

LVSWI is a reliable parameter for left ventricular function [35].

The reduction in coronary perfusion pressure in the injured/treated, and to a certain extent the

uninjured/ treated sheep, suggested that ECLS contributed to the decrease in CPP, in addition to smoke

injury. CPP is an indicator of myocardial perfusion and has been proposed as a drug target during

resuscitation [36]. The observations in the present study support the suggestion that CPP could be used to

predict the severity of injury in sheep.

The apparent increase in CaO_2 in the injured sheep could have been due to the relative increase in

[Hb] secondary to dehydration. The low DO_2I in the injured/treated and uninjured/treated groups suggested

that ECLS had a contribution, in addition to smoke, based on the relatively higher DO_2I in the

injured/untreated sheep. Interestingly, the O_2EI had a comparable profile to that of the PaO_2/FiO_2 ratio, and

could also be used to predict the contribution of ECLS to smoke-related injury.

The smoke-injured sheep required considerable amounts of intravenous fluids to compensate for the

losses from pulmonary exudation and inflammation [17, 19]. The mean urine production in all groups was

marginally lower than the published normal of 1.2 mL/h [22] but still considered to be within the acceptable

range for this cohort of sheep. The amount of anaesthetic drugs used was considered adequate for the

experiments. Heparin infusion was indicated to prolong the activated clotting time to minimise the risk of

thrombosis during intravascular procedures [37].

The reduction in the ECLS pump speed, flow, and the pressure differential could have resulted from

systemic hypotension contributing to low amounts of blood to the pump. The ECLS was configured such that

the centrifugal pump pulled blood from the inferior vena cava and returned it into the right atrium; therefore, if

the circulating volume was low, the flow would decrease for a given pump speed and in this case, both rpm

and flow dropped. Centrifugal ECLS pumps are known to be preload dependent and afterload sensitive [38],

making rpm and flows directly proportional to each other. The reason for the systemic hypotension remains

undetermined. It is possible that an unknown pulmonary component or product produced in the smoke-

damaged lungs played a role. The body temperature of the sheep was generally within the physiological

range.

Certain observations about this study could affect the interpretation of red blood cell indices and their

derivatives. For instance, animals differ from humans in that estimated changes in plasma volume is

preferably determined by changes in packed cell volume (PCV) or haemoglobin concentration and total

plasma protein (TPP) [39–41]. Also, in animals, there is a wider range of normal PCV than TPP [42]. In

domestic animal critical care, the change in both PCV and TPP is most useful as a crude index of change in

plasma volume [43]. A centrifuge that spins minute amounts of blood for rapid, cost-effective determination

of PCV and TPP permits instant adjustments in the animal's fluid needs. However, measurements of PCV

and TPP were not made in the primary study. As with all data that are collected with different objectives, it

was considerably tedious to align certain time points with real-time observations made in the laboratory,

especially for the manually input data. There was also no information about pre-anaesthetic blood tests.

An additional limitation relates to the first objective of creating a data management system for tabulating

large data sets from human studies using animal models. Because the method has not been validated, it is

considered preliminary and further validation studies are required. Also, the numbers of sheep were low and

this was especially so in the injured/untreated and placebo/untreated groups, preventing comparisons

between the treated and untreated sheep. A further limitation is that cytokine levels, as predictors of lung

injury, were not quantified. Using ELISA assays to quantify cytokine levels proved difficult and the cost was

prohibitive in the present study. It is partly for this reason that pioneering studies for the development of

proteogenomic assays were proposed [44] as an alternative to ELISA to learn from circulating markers of

acute inflammation in injured sheep used as models of intensive care, to understand critical illness.

Conclusions

The results of this study demonstrated that this preliminary method of raw data processing was effective and

helped show that ECLS contributed to further worsening of pulmonary pathology by reducing lung

compliance and PaO_2/FiO_2 ratio. The O_2EI changes mirrored those of the PaO_2/FiO_2 ratio, and decreasing

CPP was a predictor of a greater magnitude of cardiopulmonary injury in sheep. These novel observations

further understanding of similar pathology in other patients; for instance, in the resuscitation of smoke-injured

animals in house or bush fires. A similar data processing approach could be used in evaluating the

effectiveness of a given experimental or clinical intervention to further the understanding of the clinical

condition being studied, and to aid in the formulation of treatments aimed at improving the survival of animal

patients. In veterinary medicine, albeit now a considerably expensive and remote option, ECLS knowledge

could complement the treatment of potentially reversible aspiration pneumonia, a secondary complication

associated with both *Ixodes holocyclus* toxicity and laryngeal paralysis, in valuable companion animals.

List of abbreviations

ANOVA: Analysis of variance

C24H: Control experiment for 24 hours

CaO2:   Arterial oxygen content

CCO: Continuous cardiac output

CI: Cardiac index

CO: Cardiac output

CPP: Coronary perfusion pressure

[Hb]:   Haemoglobin concentration

CVP: Central venous pressure

DO2I: Oxygen delivery index

E24H: Extracorporeal life support for 24 hours

E2H: Extracorporeal life support for 24 hours

ECLS: Extracorporeal life support

ECMO: Extracorporeal membrane oxygenation

etCO2:   End tidal carbon dioxide tension

FCOHb: Fraction of carboxyhaemoglobin

FiO_2:   Fraction of inspired oxygen

FO_2Hb: Fraction of oxyhaemoglobin

HR: Heart rate

LVSWI: Left ventricular stroke work index

MAP: Mean arterial pressure

MERF: Medical Engineering Facility

MetHb: Methaemoglobin

MPAP: Mean pulmonary artery pressure

O2EI: Oxygen extraction index

PaO2:   Arterial partial pressure of oxygen

PAP: Pulmonary artery pressure

pCO_2:   Partial pressure of carbon dioxide

PCV: Packed cell volume

PEEP: Positive end-expiratory pressure

pO_2:   Partial pressure of blood oxygen

PVRI: Pulmonary vascular resistance index

QUT: Queensland University of Technology

QUT-MERF: Queensland University of Technology Medical Engineering Research Facility

SC24:   Smoke control experiment for 24 hours

SD: Standard Deviation

SE24H: Smoke injury and extracorporeal life support for 24 hours

SEA24H: Smoke injury, stored blood transfusion and extracorporeal life support for 24 hours

SEF24H: Smoke injury, fresh blood transfusion and extracorporeal life support for 24 hours

SPO_2:   Blood oxygen saturation

SV: Stroke volume

SVI: Stroke volume index

SvO_2:   Mixed venous oxygen saturation

SVR: Systemic vascular resistance

SVRI: Systemic vascular resistance index

TPP: Total plasma protein

UQ: The University of Queensland

Competing interests

A previously undisclosed conflict of interest became apparent from a section of adjunct persons within the

research group when important early findings of this paper were first presented at an academic milestone

seminar at The University of Queensland in August 2013. Therefore, this report comprises work completed

during Research Higher Degree studies from September 2012 to 23 August 2013, only.

Author Contributions

The author (SC) was solely responsible for the study design, writing the manuscript, analysing and

interpreting the data, final approval of the manuscript, and is fully accountable for the work.

Acknowledgements

This study was funded by the Australian Government National Health and Medical Research Council

(NHMRC) and an Australian Postgraduate Award (APA) scholarship through the University of Queensland

for Research Higher Degree (RHD) studies. It was conducted at Queensland University of Technology

Medical Engineering Research Facility (QUT-MERF). Gratitude is extended to Dunster KR, Diab S and Platts

D for considerable assistance in the animal laboratory and sharing data, and also Passmore M, McDonald C,

Simonova G, Foley S, Hayes R, Shekar K, Fung Y, and Fraser JF for their various roles as members of the

sheep ECMO research team. Gratitude is also extended to Tung J-P, for the initial advice on data handling

and the staff at QUT-MERF for their invaluable assistance in the care of the sheep. The author thanks Dr.

Jane Charbonneau of Edanz Group, Ltd. for editorial support.

References

1. Chemonges S, Shekar K, Tung JP, Dunster KR, Diab S, Platts D, et al. Optimal Management of the Critically Ill: Anaesthesia, Monitoring, Data Capture, and Point-of-Care Technological Practices in Ovine Models of Critical Care. Biomed Res Int. 2014;2014:468309.

2. Platts DG, Fraser JF, Hilton A, Diab S, McDonald C, Tunbridge M, et al. A novel echocardiographic imaging technique, intracatheter echocardiography, to guide veno-venous extracorporeal membrane oxygenation cannulae placement in a validated ovine model. Intensive Care Med Exp. 2014;2:2.

3. Platts DG, Diab S, Dunster KR, Shekar K, Burstow DJ, Sim B, et al. Feasibility of Perflutren Microsphere Contrast Transthoracic Echocardiography in the Visualization of Ventricular Endocardium during Venovenous Extracorporeal Membrane Oxygenation in a Validated Ovine Model. Echocardiography. 2015;32:548-56.

4. Shekar K, Wallis SC, Smith MT, Roberts JA, Fraser JF, Fung YL, et al. Development of simulated and ovine models of extracorporeal life support to improve understanding of circuit-host interactions. Crit Care Resusc. 2012;14:105-11.

5. Rais-Bahrami K, Van Meurs KP. Venoarterial versus venovenous ECMO for neonatal respiratory failure. Semin Perinatol. 2014;38:71-7.

6. Brenner P, Zelger T, Kainzinger S, Sodian R, Überfuhr P, Mair H, et al. Prolonged extracorporeal membrane oxygenation (ECMO) in 231 patients after different indications with low cardiac output - a single-center experience. Thorac Cardiovasc Surg. 2012;60:V23.

7. Fraser JF, Platts DG, Mullany DV, Fung YL, Shekar K, Diab S, et al. ECMO – the clinician's view. ISBT Sci Ser. 2012;7:82-8.

8. Riedel T, Fraser JF, Dunster K, Fitzgibbon J, Schibler A. Effect of smoke inhalation on viscoelastic properties and ventilation distribution in sheep. J Appl Physiol. 2006;101:763-70.

9. Chemonges S. Critical care management of sheep receiving extra-corporeal membrane oxygenation due to smoke induced acute lung injury (ECMO S-ALI) and acute sepsis. In: Australian and New Zealand College of Veterinary Scientists: College Science Week. Surfers Paradise, Australia. ANZCVS; 2013.

10. Chemonges S. Contemporary data capture, anaesthesia monitoring and point-of-care technology in critical care research settings for animal models In: Australian and New Zealand College of Veterinary Scientists: College Science Week. Surfers Paradise, Australia. ANZCVS; 2013.

11. Passmore M, Dunster K, Chemonges S, Diab S, Minchinton R, Shekar K, et al. ECMO contributes to lung injury in an ovine model of veno-venous ECMO. In: Proceedings of the ANZICS/ACCCN Intensive Care Annual Scientific Meeting (ASM), Hotel Grand Chancellor Hobart; 2013.

12. Passmore M, Dunster K, Chemonges S, Diab S, Minchinton R, Shekar K, et al. ECMO contributes to lung injury in an ovine model of veno-venous ECMO. In: Proceedings of the 1st International Conference of the Asia-Pacific Chapter of the Extracorporeal Life Support Organization (APELSO) 11/2013. 2013; Beijing, China; 2013.

13. Passmore M, Fung YL, Dunster, K, Diab S, Shekar K, Fraser J. Lung Inflammatory Changes In An Ovine Model Of Venovenous Extracorporeal Membrane Oxygenation. In: American Thoracic Society International Conference Abstracts, B47 ACUTE RESPIRATORY DISTRESS SYNDROME. American Thoracic Society. 2014:A3064-A3064.

14. Quinn CT, Raisis AL, Musk GC. Evaluation of Masimo signal extraction technology pulse oximetry in anaesthetized pregnant sheep. Vet Anaesth Analg. 2013;40:149-56.

15. McDonald CI, Fung YL, Fraser JF. Antioxidant Trace Element Reduction in an In Vitro Cardiopulmonary Bypass Circuit. ASAIO J. 2012;58:217-22.

16. McDonald CI, Fung YL, Shekar K, Diab SD, Dunster KR, Passmore MR, et al. The impact of acute lung injury, ECMO and transfusion on oxidative stress and plasma selenium levels in an ovine model. J Trace Elem Med Biol. 2015;30:4-10.

17. Lange M, Cox RA, Traber DL, Hamahata A, Nakano Y, Traber LD, et al. No correlation between initial arterial carboxyhemoglobin level and degree of lung injury following ovine burn and smoke inhalation. Exp Lung Res 2014;40:99.

18. Combs MDA, Rendell D, Reed KFM, Mace WJ, Quinn JC. Evidence of dehydration and electrolyte disturbances in cases of perennial ryegrass toxicosis in Australian sheep. Aust Vet J. 2014;92:107-13.

19. Westphal M, Morita N, Enkhbaatar P, Murakami K, Traber DL, Traber L. Carboxyhemoglobin formation following smoke inhalation injury in sheep is interrelated with pulmonary shunt fraction. Biochem Biophys Res Commun. 2003;311:754-58.

20. Westphal M, Murakami K, Burke AS, Westphal-Varghese BB, Rudloff HE, Salsbury JR, et al. Combined burn and smoke inhalation injury impairs ovine hypoxic pulmonary vasoconstriction. Crit Care Med. 2006;34:1428-36.

21. Bradberry S. Methaemoglobinaemia. Medicine. 2007;35:552-3.

22. Adams D, McKinley M, Colditz I, Dart C. The Sheep. In: Australian and New Zealand Council for the Care of Animals in Research & Teaching (ANZCCART) Humane Science. ANZCCART Fact Sheet A9. The University of Adelaide: Australian and New Zealand Council for the Care of Animals in Research & Teaching; 2009.

23. McGrath SR, Lievaart JJ, Friend MA. Extent of utilisation of dual-purpose wheat for grazing by late-pregnant and lambing ewes and producer-reported incidence of health issues in southern New South Wales. Aust Vet J. 2013;91:432-6.

24. Handy JM, Soni N. Physiological effects of hyperchloraemia and acidosis. Br J Anaesth. 2008;101:141-50.

25. Peacock RE, Smart L. False hyperchloraemia in a dog secondary to ingestion of horse feed supplemented with potassium bromide. Aust Vet J. 2013;91:320-2.

26. Duhm J, Göbel BO. Na+-K+ transport and volume of rat erythrocytes under dietary K+ deficiency. Am J Physiol. 1984;246:C20-9.

27. Nguyen MT, Yang LE, Fletcher NK, Lee DH, Kocinsky H, Bachmann S, et al. Effects of K+-deficient diets with and without NaCl supplementation on Na+, K+, and H2O transporters' abundance along the nephron. Am J Physiol Renal Physiol. 2012;303:92-104.

28. Oh YT, Kim J, Youn JH. Role of pituitary in K+ homeostasis: impaired renal responses to altered K+ intake in hypophysectomized rats. Am J Physiol Regul Integr Comp Physiol. 2013;304:R1166-74.

29. Russell KE, Roussel AJ. Evaluation of the Ruminant Serum Chemistry Profile. Vet Clin North Am Food Anim Pract. 2007;23:403-26.

30. Sejrsen K, Hvelplund T, Nielsen MO. Ruminant physiology: digestion, metabolism and impact of nutrition on gene expression, immunology and stress. Wageningen: Wageningen Academic

Publishers; 2006.

31. Ewaschuk JB, Naylor JM, Zello GA. D-lactate in human and ruminant metabolism. J Nutr. 2005;135:1619-25.

32. Pohlmann JR, Akay B, Camboni D, Koch KL, Mervak BM, Cook KE. A low mortality model of chronic pulmonary hypertension in sheep. J Surg Res. 2012;175:44-8.

33. Frea S, Bovolo V, Bergerone S, D'Ascenzo F, Antolini M, Capriolo M, et al. Echocardiographic evaluation of right ventricular stroke work index in advanced heart failure: a new index? J Card Fail. 2012;18:886-93.

34. Cameli M, Bernazzali S, Lisi M, Tsioulpas C, Croccia MG, Lisi G, et al. Right ventricular longitudinal strain and right ventricular stroke work index in patients with severe heart failure: left ventricular assist device suitability for transplant candidates. Transplant Proc. 2012;44:2013.

35. Choi J-O, Park SW, Park PW, Lee S-C, Choi SH, Kim SM, et al. Noninvasive assessment of left ventricular stroke work index in patients with severe mitral regurgitation: correlation with invasive measurement and exercise capacity. Echocardiography. 2010;27:1161.

36. Raghunathan K, Barbeito A, MacLeod D. Goal-directed advanced cardiac life support: coronary perfusion pressure as a target during resuscitation. Crit Care Med. 2013;41:2817-18.

37. Oliver WC. Anticoagulation and coagulation management for ECMO. Semin Cardiothorac Vasc Anesth. 2009;13:154-75.

38. Jayanthkumar HS, Murugesan C, Rajkumar J, Harish BR, Muralidhar K. Our experience with implantation of VentrAssist left ventricular assist device. Indian J Anaesth. 2013;57:56-61.

39. Falzon LC, Menzies PI, Shakya KP, Jones-Bitton A, Vanleeuwen J, Avula J, et al. A longitudinal study on the effect of lambing season on the periparturient egg rise in Ontario sheep flocks. Prev Vet Med. 2013;110:467-80.

40. Hayes GM, Mathews K, Floras A, Dewey C. Refractometric total plasma protein measurement as a cage-side indicator of hypoalbuminemia and hypoproteinemia in hospitalized dogs. J Vet Emerg Crit Care. 2011;21:356-62.

41. White A, Reyes A, Godoy A, Martínez R. Effects of transport and racing on ionic changes in thoroughbred race horses. Comp Biochem Physiol A Physiol. 1991;99:343-6.

42. Tang Z, Lee JH, Louie RF, Kost GJ. Effects of different hematocrit levels on glucose measurements with handheld meters for point-of-care testing. Arch Pathol Lab Med. 2000;124:1135-40.

43. Stafford KJ, Mellor DJ, Todd SE, Gregory NG, Bruce RA, Ward RN. The physical state and plasma biochemical profile of young calves on arrival at a slaughter plant. N Z Vet J. 2001;49:142-9.

44. Chemonges S, Tung JP, Fraser JF. Proteogenomics of selective susceptibility to endotoxin using circulating acute phase biomarkers and bioassay development in sheep: a review. Proteome Sci. 2014;12:12.

Figure Legends

Figure 1. Uncleaned manually input extra-corporeal membrane oxygenation (ECMO) treatment monitoring

data in sheep

Figure 2. Cleaned and time-point annotated manually input extra-corporeal membrane oxygenation (ECMO)

treatment monitoring data in sheep

Figure 3. Uncleaned electronically acquired physiological monitoring and treatment data during extra-

corporeal membrane oxygenation (ECMO) in sheep

Figure 4. Cleaned electronically acquired physiological monitoring and treatment data during extra-corporeal

membrane oxygenation (ECMO) in sheep

Figure 5. Data integrity checks and artefact removal of physiological monitoring and treatment data during

extra-corporeal membrane oxygenation (ECMO) in sheep

Figure 6. Completed extra-corporeal membrane oxygenation (ECMO) treatment data sheet in sheep listed

by parameter for further data analysis

a: treatment timeline, b: all experiments where sheep received smoke inhalation acute lung injury

(SE24H), c: parameter trend plot, and d: descriptive statistics panel

Figure 7. Pulmonary compliance of smoke and non-smoke injured sheep receiving extracorporeal membrane

oxygenation (ECMO) support alongside untreated controls

Dotted lines represent error bar margins. Values are presented as mean ± standard deviation.

Figure 8. Arterial oxygen tension for injured and uninjured sheep receiving extracorporeal membrane

oxygenation (ECMO) support alongside untreated controls

Arterial oxygen tension (pO_2) values are presented as mean ± standard deviation with no error bars

shown.

Tables and captions

Table 1 Subject characteristics in treated and untreated groups (control experiments) of sheep

Experiment Group	Date	Sheep No.	Age (Y)	Weight (kg)	Length (m)	BSA
E24H	06/10/2011	E24H-01/390	2	50	110	1.29
	20/10/2011	E24H-02	2	47.6	110	1.25
	17/11/2011	E24H-03	2	51	110	1.31
	01/03/2012	E24H/4616	2	50	110	1.29
	29/03/2012	E24H-05/4627	2	47	110	1.24
	04/04/2012	E24H-06/4146	2	40	110	1.11
	12/04/2012	E24H-07/4032	2	52.5	110	1.34
	03/05/2012	E24H-08/4630	2	53	110	1.34
SE24H	02/02/2012	SE24H-01/4139	2	44	110	1.19
	09/02/2012	SE24H-02/4542	2	53	110	1.34
	16/02/2012	SE24H-03/4280	2	45.5	110	1.21
	23/02/2012	SE24H-04/4624	2	50	110	1.29
	17/05/2012	SE24H-05/4458	2	55	140	1.38
	24/05/2012	SE24H-06/8461	2	46	140	1.22
	24/01/2013	SE24H-07/09C8032	3	52	130	1.33
	21/02/2013	SE24H-09A0142	2	50	140	1.29
SC24H	18/06/2013	SC24H-01	2	51	140	1.31
	27/06/2013	SC24H-02	2	57	140	1.41
C24H	08/08/2013	C24H-01	2	53	140	1.34

719 **Key:** BSA=Body surface area; E24H=Uninjured sheep treated with extracorporeal life support (ECLS) for 24

720 hours (uninjured/treated); SE24H= Smoke-induced acute lung injured sheep treated with ECLS for 24 hours

721 (injured/treated); SC24H= Smoke-induced acute lung injured sheep monitored for 24 hours without ECLS

722 (injured/untreated); C24H=Sheep subjected to room air injury as a control for smoke and monitored for 24

723 hours without ECLS (placebo/untreated).

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
1	Sheep Number		390													
2	Date		10/06/2011													
3																
4					0:00	0:30	1:00	1:30	2:00	2:30	3:00	3:30	4:00	5:00	6:00	6:30
5																
6	Time Expected			Baseline	9:55	10:25	10:55	11:25	11:55	12:25	12:55	13:25	13:55	14:55	15:55	16:25
7	Time Actual			9.02	10:00	10:12	10:57	11:30	12:05				13:55	16:10		
8																
9	Temperature	38 - 39.5	°C	38.9	39			39	38.9		38.9		38.9	39	39	
10																
11	Ventilator Settings															
12	Mode			CMV	CMV			CMV	(S) CMV		(S) CMV		(S) CMV	(S) CMV	(S) CMV	
13	Ventilator Rate	/min		18	6			6	6		6		6	6	6	
14	Respiratory F 15 - 40	/min		0	0			6	6		6		6	6	6	
15	Ventilator Tidal Volume	/min		500	220			220	220		220		220	220	220	
16	Tidal Volume	mL		504	188			206	225		218		221	198	220	
17	Peak Inspirat <30	cmH2O		23	14			18	17		19		19	19	22	
18	Minute Volume	L/min		9.2	2.3			1.32	1.32		1.28		1.41	1.23	1.25	
19	FiO2 deep PaO2	%		100	40			40	40		40		40	21	21	
20	I:E Ratio			1:2	1:2			1:2					1:2	1:2	1:2	

Figure 1

	A	B	C	D	E	F	G	H	I	J	K	L	M	N	O	P
1	Sheep Number		390													
2	Date		10/06/2011													
3																
4	Blood Gas Values	Normal Values		Baseline	0 Hr of ECMO	0.25 Hr of ECMO	1 Hr of ECMO	1.5 Hrs of ECMO	2 Hrs of ECMO	4	6	6.5	7	8	10	12
5	pH	7.42 - 7.44		7.41	7.49	7.48	7.486	7.494	7.487	7.451	7.444	7.417	7.413	7.466	7.481	7.478
6	pCO2	35 - 44 (Ai mmHg		39.3	34.1	33.4	34.5	37.3	34.2	38.6	39.2	41.7	41.9	36.2	34.6	35.8
7	pO2	99-103 (>7 mmHg		553	576	277	193	142	230	205	110	106	110	94.5	100	114
8	Baro		mmHg	762	761	761	760	760	760	759	759	759	759	759	759	758
9																
10	Oximetry Values															
11	ctHb	9 - 15	g/dL	7.8	7.5	6.1	6.7	6.7	6.5	6.5	6.2	6	5.9	5.6	5.2	4.9
12	sO2		%													
13	F02Hb		%	99.1	99.1	98.3	98	97.7	97.9	98.4	96.8	96.8	96.8	96.3	96.9	97.3
14	FCO2Hb		%	4.6	4.8	4.8	4.8	4.9	5	4.8	5	4.8	4.9	5.1	5	5.1
15	FHHb		%													
16	FMetHb		%	1.7	1.7	2	2	1.8	2.1	1.7	1.9	1.8	1.8	1.8	1.7	1.6
17																
18	Electrolyte Values															
19	K	Pla 4 - 5	mmol/L	3.8	3.8	3	2.7	2.7	2.6	2.7	3.3	3.5	3.7	3.4	3.3	3.3
20	Na	141 - 149	mmol/L	143	141	141	140	141	141	140	139	140	140	140	141	141

Figure 2

Date	03-29-2012	ECG	Arterial				CVP	PA			Oxygenator		Capnograph		Oximeter		Pump		
Pump Time (mins)	Time of Day	Heart Rate	Mean	Systolic	Diastolic	Heart Rate	Mean	Mean	Systolic	Diastolic	Post	Pre	EtCO2	Resp Rate	SpO2	Heart Rate	Flow	Speed	Mode
-195	9:15:01				348	163	1	-5	7	-17			34	15					(S)CMV
-190	9:20:01	113			0		1	-6	-6	-6			35	15	95	111			(S)CMV
-185	9:25:01	118			0		1	-6	-5	-5			36	15	94	117			(S)CMV
-180	9:30:01	126			0		1	-6	-4	-7			36	15	95	126			(S)CMV
-175	9:35:01	127			0		1	-6	-6	-6			36	15	93	130			(S)CMV
-170	9:40:01	258			0		1	-6	-6	-6			36	15	96	131			(S)CMV
-165	9:45:01	259	126	143	115	129	1	-7	-7	-7			37	15	92	86			(S)CMV
-160	9:50:01	124	125	133	117	123	15	24	28	19			37	15					(S)CMV
-155	9:55:01	234	135	142	132	117	16	25	28	21			38	15					(S)CMV
-150	10:00:01	119	122	137	113	119	17	25	28	22			39	15	88	114			(S)CMV
-145	10:05:01	118	118	134	109	118	18	25	28	23			40	15	99	120			(S)CMV
-140	10:10:01	118	119	135	111	118	18	26	28	23				0	97	117			(S)CMV
-135	10:15:01	114	106	121	98	114	19	27	30	25	309	-16	43	15	99	113			(S)CMV
-130	10:20:01	112	116	120	114	0	18	27	29	24		-15	42	15	100	113			(S)CMV

Figure 18

Figure 4

	A	B	C	D	E	F	G
2	Date of start of Expt-->	2/02/2012	9/02/2012	16/02/2012	23/02/2012	17/05/2012	24/05/2012
3	Pump Time (mins)	SE24H-01	SE24H-02	SE24H-03	SE24H-04	SE24H-05	SE24H-06
4							
5	Baseline	120	106	121	126	135	110
6	-120	107	103	113	120	135	124
7	-115	106	103	114	120	134	124
8	-110	105	103	115	108	103	124
9	-105	102	102	115	121	110	119
10	-100	105	102	116	117	88	117
11	-95	110	104	119	122	86	119
12	-90	110	109	119	123	103	119
13	-85	109	109	118	123	115	169
14	-80	108	108	117	121	82	127
15	-75	110	109	121	122	91	125
16	-70	113	109	116	120	112	120
17	-65	111	107	111	121	109	115
18	-60	113	112	110	122	108	115
19	-55	113	111	112	122	85	103
20	-50	110	108	112	118	93	79
21	-45	110	108	114	147	115	126
22	-40	111	104	121	139	132	126
23	-35	111	104	112	120	122	129
24	-30	109	108	112	114	115	134
25	-25	119	105	114	183	107	131
26	-20	105	102	118	116	107	181
27	-15	100	98	111	156	106	123
28	-10	102	100	111	99	114	123

Statistics block (rows 25–28):

SE24H-01		SE24H-02		SE24H-03	
Mean	123.3986486	Mean	109.5828025	Mean	95.08598726
Standard Error	1.687118612	Standard Error	0.673284012	Standard Error	0.639207553
Median	116	Median	109	Median	95
Mode	110	Mode	100	Mode	95

Figure 5

Sheet tabs: ECG- Heart Rate - from KD | ECG- Heart Rate | MAP | SAP | DAP | HR (Arterial Line) | CVP (Mean) | PAP (Mean) | S-PAP | D-PAP | Pre-Oxygenator Pressure | Post-Oxygenator Pressure | EtCO2

Figure 6

ECMO Time (Hrs)	SE24H-01	SE24H-02	SE24H-03	SE24H-05	SE24H-06	SE24H-07	SE24H-08
Date of start of Expt-->	2/02/2012	9/02/2012	16/02/2012	23/02/2012	17/05/2012	24/05/2012	24/01/2013	21/02/2013
Baseline	461	521	514	493	558	554	566	503
Smoke Injury	513	449	413	459	507	550	467	585
5 Min Post Smoke	510	455	394	454	510	554	435	582
1 Hr Post Smoke	485	440	465	380	408	492	414	576
0 hr of ECMO	524	304	387	394	440	480	427	570
0.25 hr of ECMO	177	352	459	124	482		436	482
1 hr of ECMO	160	89.3	144	122	139	143	129	141
1.5 hrs of ECMO	134	80.6	122	81.1	151	155	140	157
2 hrs of ECMO	154	66.6	88.4		147	149	155	167
4	161	101		88	144	159	137	183
6	158	114	92	78	164	155	149	168
6.5	121	95.2	131	75.6	154	143	152	176
7		99.9	127	111	148	102	142	174
8	199	82.2	132		146	106	161	171
10	167	203	137	91.5	138	186	176	171
12	161	192	126	105	118	152	179	178
14	181	169	132	95.4	132	155	198	
16	140	170	127	109	80.8	163	172	210
18	163	180	134	89.2	90	159	131	197
20	150	147	137	80.9	120	151	185	199
22	153	158	145	203	140	156	176	188
24	110	164	166	203	75.1	155	185	222

Baseline and 24 Hours of ECMO in S-ALI Sheep

c

b

d

	SE24H-01		SE24H-02		SE24H-03
Mean	237.238095	Mean	210.581818	Mean
Standard Err	33.0276871	Standard Err	30.2840039	Standard Err
Median	161	Median	166.5	Median
Mode	161	Mode	#N/A	Mode

FIGURE 7. Pulmonary compliance (Mean ± SD) of smoke and non-smoke injured sheep receiving extracorporeal membrane oxygenation (ECMO) support alongside untreated controls. Dotted lines represent error bar margins

Figure 7

FIGURE 8. Arterial oxygen tension (pO$_2$) (Mean ± SD with no error bars shown) for smoke and non-smoke injured sheep receiving extracorporeal membrane oxygenation (ECMO) support alongside untreated controls

Figure 8