TITLE Disordered clusters of Bak dimers rupture mitochondria during apoptosis

AUTHORS/AFFILIATIONS

Rachel T. Uren[1,2], Martin O'Hely[1,2,3], Sweta Iyer[1,2], Ray Bartolo[1,4], Jason M. Brouwer[1,2], Amber E.
Alsop[1,2], Grant Dewson[1,2], Ruth M. Kluck*[1,2].

1. The Walter and Eliza Hall Institute of Medical Research, 1G Royal Parade, Parkville, Victoria,
3052, Australia

2. Department of Medical Biology, The University of Melbourne, 1G Royal Parade, Parkville,
Victoria, 3052, Australia

3. MOH Current address: Barwon Infant Study, Deakin University, PO Box 218, Geelong, Victoria,
3220, Australia

4. RB Current address: Murdoch Children's Research Institute, The Royal Children's Hospital,
Flemington Road, Parkville, Victoria, 3052, Australia

CONTACT

* Corresponding author Ruth M. Kluck, The Walter and Eliza Hall Institute of Medical Research,
1G Royal Parade, Parkville, Victoria, 3052, Australia, Tel: +61 3 9345 2487; Fax: +61 3 9347
0852; Email: kluck@wehi.edu.au

ABSTRACT

During apoptosis, Bak and Bax undergo major conformational change and form symmetric dimers

that coalesce to perforate the mitochondrial outer membrane via an unknown mechanism. We have

employed cysteine labelling and linkage analysis to the full length of Bak in mitochondria. This

comprehensive survey showed that in each Bak dimer the N-termini are fully solvent-exposed and

mobile, the core is highly structured, and the C-termini are flexible but restrained by their contact

with the membrane. Dimer-dimer interactions were more labile than the BH3:groove interaction

within dimers, suggesting there is no extensive protein interface between dimers. In addition,

linkage in the mobile Bak N-terminus (V61C) specifically quantified association between dimers,

allowing mathematical simulations of dimer arrangement. Together, our data show that Bak dimers

form disordered, compact clusters to generate lipidic pores. These findings provide a molecular

explanation for the observed structural heterogeneity of the apoptotic pore.

INTRODUCTION

The Bcl-2 family of proteins are the principal regulators of apoptotic cell death, with either Bak or Bax

required to permeabilise the mitochondrial outer membrane (Lindsten et al., 2000, Wei et al., 2001).

Bak and Bax are triggered to convert to their activated conformation by the binding of BH3-only

relatives such as Bid and Bim (Gavathiotis et al., 2010, Dai et al., 2011, Du et al., 2011, Czabotar et

al., 2013, Leshchiner et al., 2013, Brouwer et al., 2014), and once activated can be sequestered by

pro-survival relatives such as Mcl-1 and Bcl-x_L (Llambi et al., 2011).

The ability of Bak and Bax to change conformation and oligomerize in the mitochondrial outer

membrane appears crucial for their capacity to perforate this membrane (Westphal et al., 2014).

Both proteins comprise nine α-helices that adopt a globular fold in the non-activated state (Suzuki

et al., 2000, Moldoveanu et al., 2006). Non-activated Bak is anchored in the mitochondrial outer

membrane via the α9-helix forming a transmembrane domain (Iyer et al., 2015). In contrast, non-

activated Bax is largely cytosolic due to binding of α9 to its own hydrophobic groove (Wolter et al.,

1997, Gahl et al., 2014). Following binding of BH3-only proteins, both Bak and Bax undergo

similar conformation changes including α1 dissociation (Weber et al., 2013, Alsop et al., 2015) and

separation of the "latch" domain (α6-α8) (Czabotar et al., 2013, Brouwer et al., 2014) from the

"core" domain (α2-α5). The core and α6 then collapse onto the membrane surface and become

shallowly inserted into the membrane to lie in-plane (Aluvila, 2014, Bleicken et al., 2014, Westphal

et al., 2014). Symmetric homodimers then form when the exposed BH3 domain of each molecule is

re-buried in the hydrophobic groove of another activated molecule (Dewson et al., 2008, Dewson et

al., 2012, Czabotar et al., 2013, Brouwer et al., 2014). Bax oligomers may also form prior to

membrane insertion (Luo et al., 2014, Sung et al., 2015).

How symmetric homodimers of Bak or Bax then associate to porate the mitochondrial outer

membrane is still unknown. Although there is structural information for Bak and Bax homodimers,

no structures of higher order oligomers of Bak or Bax have been resolved. Bak and Bax have been

shown to coalesce into clusters at the mitochondria upon an apoptotic stimulus (Nechushtan et al.,

2001) and small clusters of Bax have been implicated in the rapid release of cytochrome *c* (Zhou

and Chang, 2008). The availability of full-length recombinant Bax makes biochemical assays more

amenable for Bax than for Bak. Thus, Bax complexes of various shapes and sizes have recently

been detected with fluorescence microscopy of recombinant Bax on artificial membranes (Subburaj

et al., 2015). Furthermore, super-resolution microscopy has shown Bax can be found in ring-like

structures, arcs and clusters at the mitochondrial outer membrane of apoptotic cells (Grosse et al.,

2016, Salvador-Gallego et al., 2016), although the orientation of Bax dimers in these structures

could not be visualised. Higher order oligomers of both Bak and Bax vary in size when assessed by

gel filtration, blue native PAGE or linkage (George et al., 2007, Dewson et al., 2012). Moreover,

linkage studies have identified several points at which dimers might associate, including

interactions at α-helices 1, 3, 5, 6, and 9 (Dewson et al., 2008, Dewson et al., 2009, Zhang et al.,

2010, Pang et al., 2012, Ma et al., 2013, Aluvila, 2014, Bleicken et al., 2014, Gahl et al., 2014, Iyer

et al., 2015, Zhang et al., 2016). Critically, it is not clear whether any of these interaction sites are

required for dimer-dimer association and assembly of the apoptotic pore. Here, we compare

linkages through the full length of Bak in cells and show that dimers associate in a disordered, lipid-

mediated fashion.

RESULTS

The Bak N-segment, α1 and α1- α2 loop are solvent-exposed in Bak oligomers

To resolve how Bak dimers coalesce to porate the mitochondrial outer membrane, we sought to

generate a more detailed biochemical map of the membrane topology of Bak dimers. This work

complemented our previous cysteine-accessibility analyses of Bak α5, α6 and α9 (Westphal et al.,

2014, Iyer et al., 2015), by analyzing the Bak N-terminus and additional residues in the α2-α5 core

and C-terminus. The two native cysteine residues of human Bak (C14 and C166) were first

substituted with serine to generate Bak Cys null (BakΔCys, i.e. C14S/C166S), and then a single

cysteine residue substituted at several positions throughout the molecule. Each Bak variant was

stably expressed in *Bak^{-/-}Bax^{-/-}* mouse embryonic fibroblasts (MEFs) and tested for function (Figure

S1). To convert Bak to the activated oligomeric form, membrane fractions enriched for

mitochondria were incubated with tBid, as previously (Dewson et al., 2008). To label solvent-

exposed cysteine residues, membrane fractions were treated with the thiol-specific labelling reagent

IASD (4-acetamido-4'-((iodoacetyl) amino)stilbene-2,2'-disulfonic acid). Two negative sulfonate

charges prevent IASD from accessing cysteine in hydrophobic environments (such as the

mitochondrial outer membrane or the hydrophobic protein core), and also allow isoelectric focusing

(IEF) to resolve IASD-labelled and IASD-unlabelled Bak (Tran et al., 2013, Westphal et al., 2014).

We assessed each cysteine-substituted Bak variant for labelling before, during or after incubation

with recombinant tBid or tBid^{Bax}, a tBid variant containing the Bax BH3 domain (Hockings et al.,

2015), that activates Bak analogous to tBid, as well as a control not exposed to IASD and another

fully exposed to labelling by denaturation and membrane solubilization (Figure 1A, Figure S2). The

approach thus monitors solvent-exposure of the residue as non-activated Bak converts to oligomeric

Bak, and may detect transient exposure during these conformational changes.

In the N-terminus, most tested residues were accessible to IASD even before tBid treatment (Figure

1B and S2), consistent with the crystal structure (2IMT) (Moldoveanu et al., 2006). Two residues in

α1 (R36C and Y41C) that were not fully accessible before tBid treatment became exposed

following incubation with tBid (Figures 1B and S2). Two additional α1 residues, Q44C and Q47C,

also showed a tendancy for increased labelling. These changes suggest that the α1-α2 loop separates

from α1, consistent with increased labelling of A54C in the α1-α2 loop that opposes α1. Thus,

increased labelling of cysteine residues placed throughout the N-segment, α1 and the α1-α2 loop,

together with exposure of several N-terminal antibody epitopes (Alsop et al., 2015), indicate that

the Bak N-terminus becomes completely solvent-exposed after activation (Figure 1D).

To complete our survey of the core and C-termini of Bak dimers (Westphal et al., 2014, Iyer et al.,

2015), we tested the cysteine accessibility of additional selected residues in the α2-α5 core and the

α7-α8 region (Figures 1B and S2). A summary of current and previous cysteine labelling data for

the full length of the Bak protein in Figure 1C provides the following picture of Bak dimer

topology. Certain cysteine residues placed in α2 became transiently exposed, consistent with

exposure of the BH3 domain and then its burial in the hydrophobic groove in the α2-α5 core dimer

(Figure 1C), as predicted by the published structure of the Bak BH3:groove homodimer (4U2V)

(Brouwer et al., 2014). Cysteine substitutions on the exposed surface of α3 were successfully

labelled with IASD before, during and after tBid treatment. In contrast, two buried residues in α4

were refractory to IASD labelling throughout. In the α4-α5 linker region, although I123C became

more exposed following activation, it was still incompletely labelled. This may reflect loss of

contact between I123C and two hydrophobic residues in the "latch" (I167 and W170), followed by

shallow membrane insertion of I123C as it resides on the hydrophobic face of the Bak dimer core.

Partial labelling of this residue is consistent with shallow membrane insertion as IASD can label

cysteine moieties up to 7.5 Å into the hydrocarbon core of a lipid bilayer (Grundling et al., 2000,

Westphal et al., 2014). Within the C-termini, the profile of IASD-labelling was consistent with the

amphipathic α6 helix lying in-plane on the membrane surface (Westphal et al., 2014) and the

hydrophobic α9 forming the sole transmembrane domain (Figure 1D).

In summary, Bak undergoes a series of conformational changes as it transitions from its non-

activated monomeric state, to an activated BH3-exposed monomeric intermediate, to the symmetric

dimer that is the building block of the Bak oligomer (Figure 1D). In the course of these structural

rearrangements, some regions of Bak display changes in solvent exposure (such as α1 which

becomes more exposed, or α2 which is transiently exposed) whilst others remain buried throughout

(such as the α9 transmembrane anchor).

The N- and C-termini of activated Bak are mobile

Linkage studies have provided significant insight into structural changes in Bak, as well as how the

activated proteins associate into high order oligomers (Dewson et al., 2008, Dewson et al., 2009,

Ma et al., 2013, Aluvila, 2014, Brouwer et al., 2014, Iyer et al., 2015). To directly compare reported

linkages, and to analyse the full length of Bak, we used our expanded library of single-cysteine Bak

variants. Each substituted cysteine was tested for the ability to disulphide bond to the same residue

in a neighbouring Bak molecule (or to a cysteine in a nearby protein) upon addition of the oxidant

copper phenanthroline (CuPhe). Linkage between Bak molecules was indicated by the presence of

2x complexes corresponding to twice the molecular weight of the Bak monomer on non-reducing

SDS-PAGE. It is important to note that these 2x complexes are the product of CuPhe-mediated

linkage between Bak molecules and do not necessarily represent the native Bak BH3:groove dimer

that will form even in the absence of linkage, as shown below on BNP. This system of single

cysteine substitutions offers an elegant screening approach for assessing the proximity of single

cysteine residues in neighbouring molecules in a point-to-point manner, but does not identify

interaction surfaces where a single cysteine substitution may not contact its counterpart (See

Discussion).

Even in the absence of tBid treatment, cysteine residues in flexible regions displayed some linkage

to neighbouring non-activated Bak molecules (Figure 2, upper panels). For example, some 2x

complexes were captured by cysteine substitutions in the flexible N-segment (G4C, C14, L19C,

S23C), and α1-α2 loop (G51C, D57C, P58C, Q66C, S68C, S69C), indicating some proximity of

Bak monomers in untreated mitochondria. Furthermore, disulphide linkage to proteins other than

Bak resulted in Bak complexes larger than the 2x species and was observed for cysteine

substitutions in the α2-α3 helices (D84C, R87C), α4-α5 loop (S121C, N124C), α8-α9 linker

(G186C, I188C) and C-terminal end of α9 (V205C). This is unsurprising as the MOM is a protein-

rich environment and there are documented examples of monomeric Bak associating with other

membrane proteins such as VDAC2 (Cheng et al., 2003, Lazarou et al., 2010, Ma et al., 2014).

After addition of tBid had activated Bak, each cysteine in the N-extremity (a region comprising the

N-terminus, α1 and α1-α2 loop) showed more efficient linkage to 2x complexes (Figure 2A, upper

panel). The freedom with which each cysteine in the N-extremity of Bak can link suggests this

region is highly mobile and does not engage in stable protein-protein or protein-lipid interactions

following activation. In contrast, cysteine residues positioned in the core (α2-α5) showed relatively

modest linkage to 2x complexes (Figure 2B, upper panel), consistent with lack of flexibility in the

α2-α5 core dimer (Brouwer et al., 2014) and its relative immobility due to association with the

membrane.

We next addressed linkage in the C-extremity encompassing the α6-α8 latch and α9 transmembrane

domain (Figure 2C, upper panel). After Bak activation, H164C in α6 showed increased linkage to

2x complexes, as reported previously (Dewson et al., 2009, Ma et al., 2013). Cysteine substitutions

in the α8-α9 loop and the beginning of the α9 transmembrane domain (up to residue N190C) also

showed linkage to 2x complexes (Figure 2C, upper panel). However, compared to the N-extremity,

linkage was less complete, possibly due to membrane association limiting mobility. Linkage of

membrane-buried α9 residues was even less efficient (Figure 2C). However, at a higher temperature

α9 residues show much stronger linkage (Figure S3) (Iyer et al., 2015), attributable to increased Bak

mobility or CuPhe penetration into the bilayer.

In summary, after Bak activation, the whole N-extremity is highly dynamic due to being fully

solvent-exposed; the core is constrained and membrane-associated; and the C-extremity is flexible,

but is anchored to the membrane.

Mobility of the N- and C-extremities permits linkage between dimers

We next used blue native PAGE (BNP) to consider two types of cysteine-mediated Bak linkage:

linkage *within* Bak dimers and linkage *between* Bak dimers. The Bak BH3:groove interaction

within dimers is an extensive protein-protein interface and is stable in 1% digitonin on BNP while

the interaction between dimers is not (Ma et al., 2013). Thus oligomerized Bak migrates only as

BH3:groove dimers on BNP. Addition of CuPhe to mitochondrial extracts prior to BNP can

stabilise high order complexes on BNP and does so by cysteine-mediated linkage *between* dimers

(Ma et al., 2013). Thus, if addition of CuPhe at the end of the mitochondrial incubation generated

4x and greater complexes on BNP, we conclude that the linkage was *between* dimers. If larger

complexes were absent and dimers predominated on BNP, any linkage was *within* the dimer. Thus,

parallel examination of CuPhe linked Bak species on BNP and non-reducing SDS PAGE permits

differentiation of three conditions: linkage *within* the dimer, linkage *between* dimers, or no cysteine

linkage (Figure 3A).

In the N-extremity (G4C to S69C), each introduced cysteine could link some dimers to larger

complexes on BNP (Figure 2A, lower panel). Upon tBid activation, SDS-PAGE showed an

increased proportion of 2x complexes, and higher order complexes (>10x) were observed on BNP

for all residues. Of note was a stretch of residues from E59 to T62 that, following tBid treatment,

exhibited very efficient linkage by SDS-PAGE (obvious 2x, negligible 1x). In this stretch, BNP

showed no dimer species and a concomitant shift to 4x species and higher order complexes,

indicating efficient linkage between dimers but no linkage within dimers. Residues closer to the α2-

α5 core (Q66C to S69C), showed progressively more 1x species (SDS-PAGE) and an increase in

dimers (BNP), indicative of becoming closer to the constrained core of the dimer. Together, these

linkage data further argue that dimers possess flexible N-extremities capable of efficient linkage

within and between dimers, with linkage becoming exclusively between dimers as residues

approach the constrained α2-α5 core (Figure 3B). Interestingly, because linkage of cysteine residues

in the E59-T62 region occurs *only* between dimers, and is very efficient, this linkage provides a

molecular sensor that can specifically quantitate dimer-dimer interactions, and is the first assay to

do so.

In the Bak C-terminus also, cysteines could link dimers to larger complexes on BNP, although a

striking laddering pattern was apparent (Figure 2C, lower panels). For example, the large (>10x)

complexes common at the highly mobile N-extremity were not apparent at the membrane-

associated C-extremity. Stronger ladders for certain residues may be explained by the membrane

associated α6-α8 allowing only certain residues to be presented laterally. Lastly, linkage in the C-

extremity may be only between dimers (rather than within dimers) as the proportion of 2x to 1x

species on SDS-PAGE is similar to the proportion of high order complexes to dimers on BNP.

Single-cysteine mutants in the rigid core capture some linkage, mostly at the lateral edges

Previous studies suggested that the structured α2-α5 core of the Bak (or Bax) dimer might form a

dominant, symmetric interface capable of driving dimers to form higher order complexes (Aluvila,

2014, Brouwer et al., 2014). For example, crystals of the Bak α2-α5 core dimer showed a side-by-

side association (4U2V)(Brouwer et al., 2014). However, this was attributed to crystal packing

effects, as chemical crosslinking between the α4-α5 loops was not detected (Brouwer et al., 2014).

Accordingly, in the present studies, disulphide bonding between the α4-α5 loops was not captured

with CuPhe (Figure 2B, lower). End-to-end associations were suggested by α3:α3' or α5:α5'

linkages (Aluvila, 2014, Brouwer et al., 2014), and disulphide bonding at the equivalent residues

was also evident in this study (H99C:H99C for α3:α3'; H145C:H145C' for α5:α5')(Figure 2B,

lower). Notably, this linkage was not as efficient as linkage observed for the N-extremity, and these

and other linkage-competent residues locate mostly to the lateral edges of the α2-α5 core dimer

when lying in-plane (Figure 3C). Thus, screening individual cysteine substitutions in the Bak dimer

core for linkage did not uncover any dominant sites of interaction between Bak dimers, but showed

widespread linkage of residues around the lateral edges of the dimer, consistent with the random

collision of Bak dimers.

These findings support a flexible extremity model for full-length Bak dimers in the MOM, outlined

in Figure 3D with side-on (left) and top-down views (right). At the centre is the α2-α5 core dimer

lying in-plane and partially submerged in the membrane surface. Extending from both sides of the

core dimer is a flexible latch (α6-α8) of ~50 residues that also lies in-plane and connects to the

transmembrane α9-helices. Extending from both ends of the dimer core is the solvent-exposed,

flexible and mobile N-extremity of ~70 residues. Each circle in Figure 3D illustrates the range of

possible positions for the flexible N- and C-extremities (blue and pink respectively). Overlap of the

largest blue circles (labelled G4C) indicates the potential for residues close to the N-terminus to link

within dimers. In contrast, the smaller blue circles (labelled V61C) do not overlap, indicating a lack

of linkage within the dimer at this residue. Notably, these linkage constraints are consistent with the

dimensions of the α2-α5 dimer core as determined from the recent crystal structure (4U2V) (Figure

3E).

These findings also reveal how dimers associate into high order oligomers. As linkage can be

induced between several regions of the Bak dimer, there is no dominant protein-protein interface

that mediates dimer assembly into high order oligomers. Rather, the linkage pattern is consistent

with transient collision of dimers in-plane on the membrane. Collisions of the α2-α5 core and α6-α8

latch are limited by their membrane-association, whereas the flexible, entirely solvent-exposed N-terminal region is free to link between dimers and does so very efficiently.

Inter-dimer interfaces are more labile to detergent than the intra-dimer (BH3:groove) interface

We next examined if the inter-dimer interactions identified above were stable in digitonin, as shown for the BH3:groove interface within dimers (Ma et al., 2013). Digitonin was added to the mitochondrial incubations after Bak had become oligomerized by incubation with tBid, but prior to disulphide bond formation induced by CuPhe (Figure 4A). As expected, digitonin did not prevent linkage *within* dimers as shown by 2x complexes of M71C/K113C on SDS-PAGE (Figure 4A, upper panel, lane 3). However, digitonin prevented all linkage *between* dimers on BNP of seven single-cysteine variants (Figure 4A, lower panel), indicating that inter-dimer interactions are mediated either by membrane or by weak protein-protein interactions.

These experiments also provided new insight into linkage within dimers. For example, two residues at the far N- and C-termini (L19C and the C-terminal extension GGSGGCK (Iyer et al., 2015)) could link within dimers to generate 2x complexes on SDS-PAGE after digitonin treatment (Figure 4A, upper panel). Thus, these residues were sufficiently distal from the core to reach their counterpart across the length of the structured α2-α5 core dimer and allow linkage within the dimer (Figure 3D, 3E). On BNP, dimers of these two residues (L19C, and to a lesser extent GGSGGCK) also migrated faster than dimers of other variants (Figure 4A, lower panel), suggesting that linkage within the dimer caused a more compact, faster migrating protein complex under these native conditions.

Bak dimers can become interspersed by Bax

To test if interactions between Bak dimers could be disrupted by other proteins, and thus test if Bak dimer aggregation was a dynamic, reversible process, we added recombinant proteins from the Bcl-2 protein family after Bak had formed dimers (Figure 4B). Dimer interactions were monitored by

linkage between the N-termini (V61C:V61C') or C-termini (H164C:H164C'), and compared to

BH3:groove linkage (M71C:K113C). To oligomerize Bak, mitochondria were first incubated with

tBid[M97A], a variant that binds poorly to prosurvival proteins such as Bcl-x$_L$ and Mcl-1 (Lee et al.,

2016). Mitochondria were then incubated with recombinant Bax, Bcl-x$_L$ or Mcl-1ΔNΔC. Finally,

CuPhe was applied and the extent of disulphide linkage assessed by non-reducing SDS-PAGE

(Figure 4B). Notably, Bax translocated to mitochondria and decreased linkage between Bak dimers

(e.g. V61C:V61C' and of H164C:H164C') but not within dimers (e.g. M71C:K113C) (Figure 4B).

In contrast, recombinant Bcl-x$_L$ and Mcl-1ΔNΔC translocated to the mitochondrial membrane but

did not interfere with linkage between Bak dimers (Figure 4B). Thus, Bax not only localised to the

same membrane microdomain as the pre-formed Bak oligomers, but became partially interspersed

with the dimers, indicating that dimer aggregation is dynamic.

V61C:V61C' linkage can monitor dimer-dimer interactions; stochastic simulations reveal a

disordered aggregate of Bak dimers

Our biochemical data support the random collision of Bak dimers at the mitochondrial outer

membrane during apoptosis. Indeed, several studies indicate that Bax and Bak can reside in a

variety of different structures at the mitochondrial outer membrane of apoptotic cells, including

clusters (Nechushtan et al., 2001, Zhou and Chang, 2008, Grosse et al., 2016, Nasu et al., 2016,

Salvador-Gallego et al., 2016). To test if random collisions between dimers could feasibly explain

the linkage patterns observed, simulations of the random contact between dimers were performed.

The efficient and exclusive linkage that occurs between dimers at residue V61C formed the basis of

our simulations (Figure 5).

To simulate the aggregation of Bak dimers on the mitochondrial outer membrane, this complex

biological system was simplified to limit the number of parameters required to mimic the

experimental outcome. We simulated the membrane as a flat surface (i.e. a 2D plane). Each subunit

representing a Bak dimer, was evenly distributed on a geometric grid imposed on this surface

(Figure 6A), and to avoid edge effects the grid was wrapped onto a torus. A hexagonal grid, rather

than square or triangular, was selected to best represent the close packing of objects on a plane with

the greatest degrees of freedom. The linkage potential of each subunit in the grid was based on the

known linkage constraints of the V61C residue; i.e. linkage occurred exclusively and efficiently

between dimers and each dimer could make at most 2 cysteine linkages, but possibly only 1 or 0

linkages if no other free cysteines were available. Furthermore, the simulation allowed linkages

only between immediately adjacent dimers, based on the short 4 Å limit of the CuPhe-mediated

disulphide linkage and the known dimensions of the Bak dimer. Thus, each hexagonal subunit had

the freedom to link with at most two neighbouring hexagons, in any direction, and this random

linkage between neighbouring dimers would result in a mixture of linkage states including double,

single or no linkages. With this hexagonal grid, four types of double linkage were possible;

opposite, obtuse, acute or reciprocal (Figure 6B). An equal probability for each of the double

linkages (opposite, obtuse, acute or reciprocal) was imposed, to reflect close but random packing of

dimers at the membrane. Furthermore, the simulation was allowed to proceed to complete linkage

of available cysteine residues, to reflect the near-complete linkage of V61C:V61C'. With these few

parameters, a complex biological system was elegantly reduced to a simplified simulation.

The simulation output was a frequency distribution of the number of Bak dimers present in each

linked complex (Figure S4). To approximate the BNP western blot densitometry, the counts were

then multiplied by the number of Bak dimer units (e.g. 1, 2, 3, 4 etc) in each of the linked Bak

complexes, and the predicted densitometry smoothed with a gaussian kernel to allow qualitative

comparison with the gaussian-like densitometry output from the BNP western blots.

Notably, the simulation produced outputs that closely matched the distinctive linkage pattern

observed for V61C, i.e. prominent 4x complexes and complexes greater than 10x the Bak molecular

weight (compare Figure 5A, 5B with Figure 6C, 6D). Thus, the simulations provided proof of

principle that the linkage pattern (V61C:V61C') observed in mitochondria can be explained by

random dimer arrangement. These logic arguments based on the linkage constraints of the V61C

residue have, for the first time, afforded single molecule resolution of Bak aggregation.

Simulations are robust to a reduction in linkage

To explore robustness of the simulations, we asked if the simulation could predict the linkage

pattern when linkage was limited due either to less V61C residues, or to cysteine residues at

positions other than V61C. In the first test, we performed mitochondrial experiments in which

~50% of the Bak molecules lacked cysteine residues. In those experiments, Bak V61C was co-

expressed with Bak Cys-null that was FLAG-tagged (Figure S5A, upper panel). As expected, only

half of the Bak molecules could link to 2x species after activation by tBid and induction of linkage

(Figure S5A, middle panel, lane 4). In addition, all molecules formed BH3:groove dimers after

incubation with tBid, as shown by BNP, but the capture of higher order oligomers mediated by

cysteine-linkage was greatly reduced (Figure S5A, lower panel, lane 4; quantified in Figure S5B).

In simulations of this system, we assumed equal expression of V61C and the Cys null variants, as

indicated by western blot analysis. We also assumed that this population of monomers forms dimers

in Mendelian proportions: 25% of dimers are V61C doublets and can link twice, 50% are

V61C:Cys null and can link once, and the remaining 25% are Cys null doublets and cannot link

(Figure S5C). The simulation output (Figure S5D) reflected that of the BNP densitometry (Figure

S5B) in terms of more dimers and fewer >10x complexes, indicating the simulation was

generalizable to a different distribution of linkage-competent Bak molecules.

We next examined if our simulations were also robust when approximating the linkage pattern of

residues other than V61C (Figure S6A). As noted above, because T62C, Q66C, S68C, and S69C

substitutions were closer than V61C to the constrained α2-α5 core, they did not link efficiently

(Figure 2A lower, quantified in Figure S6B). To simulate this steric hindrance caused by a shorter

distance from the constrained core, an *edge blocking* effect was introduced to the model; every edge

of every hexagon was prohibited from having a link form across it with this probability for the

duration of each simulation. As the edge blocking probability increased from 0 to 0.75, linkage

between dimers decreased to generate patterns comparable to BNP analysis of these core-proximal

residues (Figure S6C). Thus, modification of a single parameter could describe linkage of residues

closer than V61C to the Bak dimer core. The adaptability of this simple simulation further

illustrated the strength and feasibility of our model of random Bak dimer aggregation at the

mitochondrial outer membrane during apoptosis.

DISCUSSION

Here we investigated the topology of Bak dimers in the mitochondrial outer membrane, and how

dimers assemble into the high order oligomers thought necessary to form apoptotic pores in that

membrane. We found that Bak dimers are characterized by flexible N- and C-extremities flanking a

rigid α2-α5 core, and that these dimers aggregate in disordered, dynamic, compact clusters.

Critically, we found no evidence for a single, dominant protein-protein interface between dimers

and predict the lipid environment plays a crucial role in facilitating the aggregation of dimers and

subsequent membrane rupture.

We had previously proposed the in-plane model for Bak dimers (Westphal et al., 2014). Our current

data advances this model to show the N- and C-extremities are flexible, based on analysis of the

full-length of Bak when activated and dimerised at the mitochondrial outer membrane (Figure 3D).

The N-extremity (~70 residues C-terminal to α2) becomes fully solvent-exposed as shown by IASD

labelling, and fully mobile as indicated by linkage, consistent with exposure of N-terminal cleavage

sites and antibody epitopes in α1 and the α1-α2 loop (Griffiths et al., 1999, Weber et al., 2013,

Alsop et al., 2015). Failure of the N-extremity to re-engage in protein-protein or protein-lipid

interactions argues that it does not contribute to the assembly of dimers. The remainder of Bak is

membrane-associated, with the α2-α5 core and α6 (and possibly α7-α8) in-plane with the outer

mitochondrial membrane surface and partially embedded (Aluvila, 2014, Brouwer et al., 2014,

Westphal et al., 2014) while α9 forms a transmembrane domain (Iyer et al., 2015). The topology of

the α2-α5 region is consistent with the α2-α5 core dimer crystal structure (~35Å × 45Å)(Brouwer et

al., 2014), as V61C (~30Å from the core dimer) linked between dimers but not within dimers.

Flexibility of the C-terminal α6-α9 region, as indicated by multiple linkages between these regions

(Bleicken et al., 2014, Iyer et al., 2015, Zhang et al., 2016), implies that embedded dimers adopt a

range of conformations on the MOM surface (Figure 7A).

Linkage experiments have identified several possible interactions between dimers of Bak or Bax

(Dewson et al., 2008, Dewson et al., 2009, Zhang et al., 2010, Pang et al., 2012, Ma et al., 2013,

Aluvila, 2014, Bleicken et al., 2014, Gahl et al., 2014, Iyer et al., 2015, Zhang et al., 2016), with

many also shown in the present study. Notably, however, a dominant protein-protein interface

between dimers was not evident. Rather, linkage between the membrane-associated regions could

be detected at many positions, arguing for a disordered arrangement. As our single-cysteine

scanning approach can only examine the homotypic pairing of cysteine residues, it remains possible

that heterotypic pairing of cysteine substitutions may detect a more complex protein interface

between dimers, as seen for near complete linkage at the Bak BH3:groove interface (M71C:K113C,

Figure 4B) (Dewson et al., 2008). However, in support of the notion that there is no dominant

protein interface between dimers, the interaction between Bak dimers is lost when digitonin is

added, in contrast to the BH3:groove interface which persists in the presence of digitonin.

Moreover, lability of dimer-dimer association was highlighted by the capacity of Bax to intermingle

with pre-formed Bak dimers. Each of Bcl$_{xL}$, Mcl-1 and Bax localised to mitochondria with similar

efficiency, however a critical feature of Bax is that, like Bak, following activation by tBid it

collapses onto the membrane surface to form homodimers (Westphal et al., 2014). The large

membrane surface occupied by Bax dimers, together with disturbance of the outer leaflet, may

explain why Bax, but not the prosurvival proteins, could disrupt Bak dimer-dimer association in this

assay.

A variety of biophysical and imaging techniques have been deployed to better understand how Bak

(or Bax) dimers associate, yet it remains unclear. Well-characterised pore-forming proteins

typically reveal an extensive interface between subunits that facilitates pore formation (Song et al.,

1996, Mueller et al., 2009). Versions of this regular architecture have recently been considered for

Bak and Bax as a closed circuit of dimers lining a circular pore (Aluvila, 2014, Bleicken et al.,

2014, Brouwer et al., 2014). In contrast to these hypothetical linear dimer arrangements, our

biochemical data revealed that there is no single dominant interface between the α2-α5 core dimers

and support the random collision of Bak dimers in two dimensions at the mitochondrial outer

membrane during apoptosis. Here a novel approach of mathematical modelling was used to

simulate linkage between Bak dimers and showed the feasibility of random dimer association. For

example, as the V61C residue linked only between Bak dimers, and did so very efficiently, this

provided a powerful tool for testing different arrangements of Bak dimers on the membrane. A

minimalist set of assumptions were adopted for stochastic simulations of V61C:V61C' linkage

between Bak dimers: linkage is complete and irreversible but occurs only between dimers; only two

linkages are possible per dimer; and neighbouring dimers can only link if both have a free cysteine

available for disulphide bonding. Indeed, the simulation showed good concordance with different

sets of linkage data obtained from fully oligomerised Bak in mitochondria. From this we conclude

that Bak dimers are closely packed and aggregate with a random orientation. As pores no longer

form if Bak levels decrease by ~50% in these cells (Ferrer et al., 2012), our observations from this

whole population of Bak represent the species of Bak dimers that contribute to pore formation,

rather than a minor species not involved in pore formation.

A disordered cluster of dimers is reminiscent of the "carpet" model of pore formation by peptides

(reviewed by (Gilbert, 2016)). Amphipathic antimicrobial peptides such as melittin lie parallel to

the membrane plane, with accumulation causing strain in the outer layer of the membrane until at

higher concentrations the lamellar structure of the membrane is destabilised and non-lamellar

lipidic pores are formed (Lee et al., 2008, Lee et al., 2013). Like an amphipathic peptide, the Bak

and Bax α2-α5 core dimers and α6-helices penetrate the outer leaflet of the bilayer (Czabotar et al., 2013, Aluvila, 2014, Brouwer et al., 2014, Westphal et al., 2014). While very high concentrations of antimicrobial peptides can disintegrate membrane in a detergent-like manner, this does not appear to occur with near- physiological levels of Bak and Bax (Lee et al., 2008). Antimicrobial peptides also have membrane thinning attributes (Chen et al., 2003), as reported for Bax (Satsoura et al., 2012). Satsoura and colleagues discuss how membrane deformation by Bax may promote oligomerization mediated not by protein:protein interactions but by long-range changes in membrane tension (Satsoura et al., 2012). Thus, as illustrated in Figure 7B, we hypothesize that Bak dimers penetrate the outer leaflet to attract further dimers and in doing so increase the membrane disturbance. Enlarging clusters then destabilize the membrane sufficiently to form lipidic pores that release apoptogenic factors. The pores may then be stabilized by parts of the Bak and Bax dimers rearranging to line the pore and form proteolipidic pores (Terrones et al., 2004, Garcia-Saez, 2012, Westphal et al., 2014). Hence, our model for the disordered clustering of Bak dimers provides a holistic molecular explanation for the detection of Bak and Bax in a variety of formations including complete rings, arcs, lines and random clusters (Grosse et al., 2016, Nasu et al., 2016, Salvador-Gallego et al., 2016).

In conclusion, this study proposes a novel means of oligomerization by Bak and Bax in apoptotic cells: that dimers aggregate in dense clusters without a dominant interface and this dynamic association requires the involvement of the lipid membrane. We have described molecular tools that can precisely monitor the aggregation of Bak dimers in cells, and could also be used to examine how the pro-survival proteins inhibit Bak oligomerisation, which may impact on the development of cancer therapies that target these proteins. Critically, the combination of linkage data and mathematical simulations described here has offered, for the first time, insight into assembly of the apoptotic pore with single molecule resolution.

MATERIALS AND METHODS

Recombinant proteins

Recombinant proteins were prepared as previously described; caspase 8-cleaved human Bid (tBid) (as per (Kluck et al., 1999)), tBid^{Bax} and Bcl-x$_L$ full length (as per (Hockings et al., 2015)), Cys null Bax full length (as per (Czabotar et al., 2013)), Mcl-1ΔN151ΔC23 (as per (Chen et al., 2005). tBid^{M97A} was kindly provided by E. Lee (Lee et al., 2016).

Bak variant library cloning and expression

A library of human Bak mutants was generated by site-directed cysteine substitution (as per (Dewson et al., 2008). Human Bak with the endogenous cysteine residues mutated to serine (Bak Cys null C14S/C166S) was used as the template in overlap extension PCR to introduce single cysteine residues throughout the Bak sequence, and then cloned into the pMX-IRES-GFP retroviral vector (primer and vector sequences available on request). Mutants were introduced into SV40-immortalized *Bak$^{-/-}$Bax$^{-/-}$* mouse embryonic fibroblasts (MEFs) by retroviral infection (as per (Dewson et al., 2008).

Preparation of membrane fractions enriched for mitochondria

Bak$^{-/-}$Bax$^{-/-}$ MEFs expressing Bak cysteine mutants were permeabilised and membrane fractions containing the mitochondria were isolated as previously described (Dewson et al., 2008). Cells were washed in PBS and then re-suspended at a concentration of 1×10^7 cells/ml in wash buffer supplemented with digitonin (100 mM sucrose, 20 mM HEPES-NaOH pH 7.5, 100 mM KCl, 2.5 mM MgCl$_2$, 4 µg/ml Pepstatin A (Sigma-Aldrich, St. Louis, MO, USA), Complete protease inhibitors (Roche, Castle Hill, NSW, Australia) and 0.025% w/v digitonin (Calbiochem, Merck, Darmstadt, Germany). Cells were incubated for 10 min on ice, then membrane fractions were collected by centrifugation at 16,000 ×*g* for 5 min and the supernatant discarded. Pellets were re-

463   suspended in wash buffer (no digitonin), and permeabilisation of the cell membrane confirmed by

464   trypan blue uptake.

Activation of Bak by tBid or Etoposide

To assess the apoptotic function of each Bak variant in cells, $Bak^{-/-}Bax^{-/-}$ MEFs expressing Bak

cysteine mutants were treated with etoposide (10 μM) for 24 h, and cell death, as indicated by

propidium iodide (5 μg/ml) uptake, determined by flow cytometry (FACSCalibur, BD Biosciences,

San Jose, CA, USA). To activate Bak and induce mitochondrial outer membrane permeabilisation

in vitro, membrane fractions from $Bak^{-/-}Bax^{-/-}$ MEFs expressing Bak cysteine variants were treated

with 100 nM recombinant tBid for 30 min at 30°C (as described in (Dewson et al., 2008). Selected

samples were treated with the functionally equivalent protein tBid^{Bax} in which the BH3 domain of

Bid was substituted with the Bax BH3 domain (Hockings et al., 2015).

**Cysteine accessibility to IASD labelling, one-dimensional isoelectric focussing and IASD
quantification**

The cysteine residue of each Bak variant was assessed for solvent exposure by incubating

membrane fractions with the cysteine-labelling reagent IASD (Molecular Probes® Life

Technologies, Carlsbad, CA, USA) for 30 min at 30°C (as described in (Westphal et al., 2014)).

Membrane fractions were supplemented with 100 μM TCEP to prevent oxidization of cysteines that

would inhibit IASD labelling. Samples were unlabelled, or incubated with IASD before, during or

after tBid incubation or following denaturation with 1% w/v ASB-16 (Merck). IASD labelling was

quenched by the addition of 200 mM DTT and samples solubilised in 1% ASB-16 for 10 min at

22°C. Soluble supernatant fractions were isolated by centrifugation at 16,000 ×*g* for 5 min, and

added to an equal volume of IEF sample buffer (7 M urea, 2 M thiourea, 2% w/v CHAPS,

Complete protease inhibitors, 4 μg/ml pepstatin A, 1% w/v ASB-16 and 0.04% w/v bromophenol

blue). Samples were loaded onto 12-well Novex® pH 3-7 IEF gels (Life Technologies) and focused

with a Consort EV265 power supply with increasing voltage (100 V for 1 h, 200 V for 1 h and 500

V for 30 min). IEF gels were pre-soaked in SDS buffer (75 mM Tris/HCl, pH 6.8, 0.6% w/v SDS,

15% v/v glycerol), and then transferred to PVDF membrane for western blot analysis.

IASD labelling was quantified (ImageLab 4.1, Bio-RAD) from western blots by measuring signal

intensity for the unlabelled and labelled Bak band species in each lane to calculate the percentage of

the total signal from each lane attributed to the faster migrating IASD-labelled species. A global

background subtraction was applied to regions of interest from each western blot. Data were

presented as the mean ± SD (n≥3), or range (n=2) and the number of replicates indicated on the *x*-

axis. A two-tailed unpaired t-test was employed to determine significant differences ($p<0.05$) in the

percentage of IASD labelled Bak before versus after tBid treatment.

Oxidant-induced disulphide linkage (CuPhe)

The cysteine residue of each Bak variant was tested for disulphide linkage with proximal cysteine

residues on Bak (or other proteins) by incubation with the oxidant copper phenanthroline (CuPhe)

(as per (Dewson et al., 2008)). CuPhe was prepared as a stock solution of 30 mM $CuSO_4$ and 100

mM 1,10-phenanthroline in 4:1 water/ethanol. Following incubation of membrane fractions with

tBid at 30 ^{o}C, samples were pre-chilled on ice for 5 min, and then incubated with a 100-fold

dilution of the CuPhe solution on ice for 15 min. Note that (Iyer et al., 2015) performed CuPhe

linkage without the pre-chill incubation. Disulphide bond formation (linkage) was quenched by the

addition of 20 mM N-ethyl maleimide (NEM, to label any remaining free cysteines, 10 min on ice)

and 5 mM EDTA (to chelate copper, 5 min on ice) and samples analysed by non-reducing SDS

PAGE or blue native PAGE (BNP).

SDS PAGE and Blue Native PAGE

CuPhe-linked samples were analysed by SDS PAGE (12% TGX gels (Life Technologies)) in the

absence of reducing agents to preserve disulphide bonds. The efficiency of linkage between Bak

molecules was measured by a shift in Bak migration from 1x to 2x complexes. To discriminate

between linkages occurring within or between Bak dimers, CuPhe-linked samples were also

analysed in tandem by blue native PAGE (BNP) to preserve the native dimer interface (i.e. the

BH3:groove interface) in addition to any disulphide bonds between Bak molecules. Migration of

higher order linked complexes (i.e. greater than dimer) on BNP indicated the presence of disulphide

bonds between Bak dimers. Following quenching of CuPhe with NEM and EDTA, membrane

fractions were isolated by centrifugation at 16,000 $\times g$ for 5 min. Supernatants were discarded and

membrane pellets were solubilised in 20 mM Bis-Tris pH 7.4, 50 mM NaCl, 10% v/v glycerol, 1%

w/v digitonin, and incubated on ice for 1 h and insoluble material was removed by centrifugation at

16,000 $\times g$ for 5 min. The resulting supernatants were prepared for BNP by the addition of Native

Sample buffer (Life Technologies) and Coomassie Additive (Life Technologies), and then loaded

onto Novex® 4-16% Native PAGE 1.0 mm 10 well gels as per the manufacturer's instructions (Life

Technologies). Western blot transfer to PVDF was performed at 30 V for 150 min in transfer buffer

(25 mM Tris, 192 mM Glycine, 20% v/v Methanol) supplemented with 0.037 % w/v SDS. PVDF

membranes were de-stained with 10% v/v acetic acid 30% v/v ethanol, then further de-stained in

methanol and rinsed thoroughly with dH$_2$O before immunoblotting.

Immunoblotting

SDS PAGE and IEF membranes were immunoblotted for Bak using the rabbit polyclonal anti-Bak

aa23-38 (B5897, Sigma-Aldrich, Castle Hill, NSW, Australia). BNP membranes were

immunoblotted for Bak using an in-house anti-Bak monoclonal rat IgG (clone 7D10, WEHI,

(Dewson et al., 2009, Alsop et al., 2015)), except in the case of Bak mutants ranging from G51C to

P58C that were detected on BNP with the anti-Bak aa23-38. In-house monoclonal rat antibodies

were used to detect recombinant Bax (clone 49-F9, WEHI), Bcl-x$_L$ (clone 9C9, WEHI) and Mcl-1

(clone 19C 4-15, WEHI). Horseradish peroxidase conjugated IgG secondary antibodies were used;

anti-rabbit (4010-05, Southern Biotech) and anti-rat (3010-05, Southern Biotech). Immobilised

horseradish peroxidase was detected with Luminata Forte western HRP substrate (WBLUF0500,

Millipore, Billerica, MA, USA), images captured with the ChemiDoc XRS+ System (Bio-RAD,

Hercules, CA, USA) and signal intensity measured with Image Lab 4.1 software (Bio-RAD).

Blue Native PAGE densitometry

Images were transformed in Image Lab 4.1 (Bio-RAD) to correct any rotation of the image and

exported as .tif for densitometry in FIJI (ImageJ 1.47n) (Schindelin et al., 2012, Schneider et al.,

2012). The lookup table (LUT) was inverted and images rotated 90 degrees. Using the rectangular

selection tool, a box was drawn around the lane of interest to encompass the full range of Bak signal

(with fixed dimensions of 370 x 50 pixels). This box was then measured with the Dynamic ROI

profiler plugin. Plot values were exported for analysis in Prism (ver 6.0f, Graphpad Software Co.,

La Jolla, CA, USA), and normalised to a fraction of the total intensity for each lane of interest. A

clearer representation of multiple replicate plots of lane intensity was achieved by a linear rescaling

of each replicate's horizontal axis so that the maximum corresponding to Bak 4x species, and the

characteristic minimum (which is a highly reproducible artefact of our Bak BNP western blots) at

~720 kDa, coincided across replicates.

Simulation of linkage between dimers (V61C:V61C')

The formation of a disulphide linked oligomer is modelled by repeatedly (a) selecting an available

pair of cysteines on neighbouring dimers at random, (b) declaring them linked and (c) removing

those cysteines from the list of those subsequently available. We continue until no neighbouring

dimers have cysteines available for linkage, to reflect complete linkage of V61C:V61C'. We

consider a collection of Bak dimers on a two-dimensional surface (see Figure 6). Dimers are

arranged on a hexagonal grid, and linkage is permitted in all directions. Each dimer has two

cysteines available for disulphide linkage to a neighbouring dimer. We simulate such that linkage is

allowed in all directions with equal probability. Reduced linkage efficiency was modelled by

introducing an edge blocking effect. Every edge between neighbouring hexagonal cells was flagged

as eligible for a disulphide bond, independently at random with probability p constant across all

edges in a given simulation, for p=0, 0.25, 0.5 and 0.75. We modelled the experimental scenario

where Bak dimers were formed from a mixture of equal amounts of V61C and cys-null monomers

by assigning each dimer either zero, one or two available cysteines with probability 0.25, 0.5 and

0.25 respectively (see Figure S5C).

Predicted densitometry for each Bak dimer multiplicity are presented by taking the distribution of

oligomer sizes on completion of the simulation, multiplying by the number of Bak dimers in each

oligomer to obtain the theoretical density that would give rise to in the western blot, using an

exponential horizontal axis to mimic the nonlinear spacing of bands, and applying a gaussian kernel

estimator to approximate the smearing of bands and their running into one another at higher

molecular weights (see Figure S4).

AUTHOR CONTRIBUTIONS

Conceptualization, RTU, GD, RMK; Methodology, RTU, MOH, SI, GD, RMK; Software, MOH;

Investigation, RTU, SI, RB, JMB, AEA, GD; Formal analysis, RTU, MOH; Data curation, MOH;

Visualization, RTU, MOH, SI, RMK; Writing – Original Draft, RTU, RMK; Writing – Review &

Editing, RTU, MOH, SI, AEA, GD, RMK; Supervision, RTU, RMK

ACKNOWLEDGMENTS

The authors declare no conflict of interest. We thank Peter Colman, Peter Czabotar, John Markham,

Matthew Ritchie, Aline Oelgeklaus and Melissa Shi for critical comments on the manuscript. We

thank Colin Hockings for helpful discussions regarding simulations. We gratefully acknowledge the

IASD labelling assay development and analysis methods of Dana Westphal. Our work is supported

by NHMRC grants (637337 and 1016701), and the Victorian State Government Operational

Infrastructure Support and the Australian Government NHMRC IRIISS.

REFERENCES

ALSOP, A. E., FENNELL, S. C., BARTOLO, R. C., TAN, I. K., DEWSON, G. & KLUCK, R. M.
2015. Dissociation of Bak alpha1 helix from the core and latch domains is required for
apoptosis. *Nat Commun,* 6**,** 6841.
ALUVILA, S. M., T. HUSTEDT, E. FAJER, P. CHOE, JY. OH, KJ. 2014. Organization of the
mitochondrial apoptotic BAK pore: oligomerization of the Bak homodimers. *J Biol Chem,*
289**,** 2537-51.
BLEICKEN, S., JESCHKE, G., STEGMUELLER, C., SALVADOR-GALLEGO, R., GARCIA-
SAEZ, A. J. & BORDIGNON, E. 2014. Structural model of active bax at the membrane.
Mol Cell, 56**,** 496-505.
BROUWER, J. M., WESTPHAL, D., DEWSON, G., ROBIN, A. Y., UREN, R. T., BARTOLO, R.,
THOMPSON, G. V., COLMAN, P. M., KLUCK, R. M. & CZABOTAR, P. E. 2014. Bak
Core and Latch Domains Separate during Activation, and Freed Core Domains Form
Symmetric Homodimers. *Mol Cell,* 55**,** 938-46.
CHEN, F. Y., LEE, M. T. & HUANG, H. W. 2003. Evidence for membrane thinning effect as the
mechanism for peptide-induced pore formation. *Biophys J,* 84**,** 3751-8.
CHEN, L., WILLIS, S. N., WEI, A., SMITH, B. J., FLETCHER, J. I., HINDS, M. G., COLMAN,
P. M., DAY, C. L., ADAMS, J. M. & HUANG, D. C. 2005. Differential targeting of
prosurvival Bcl-2 proteins by their BH3-only ligands allows complementary apoptotic
function. *Molecular Cell,* 17**,** 393-403.
CHENG, E. H., SHEIKO, T. V., FISHER, J. K., CRAIGEN, W. J. & KORSMEYER, S. J. 2003.
VDAC2 inhibits BAK activation and mitochondrial apoptosis. *Science,* 301**,** 513-517.
CZABOTAR, P. E., WESTPHAL, D., DEWSON, G., MA, S., HOCKINGS, C., FAIRLIE, W. D.,
LEE, E. F., YAO, S., ROBIN, A. Y., SMITH, B. J., HUANG, D. C., KLUCK, R. M.,
ADAMS, J. M. & COLMAN, P. M. 2013. Bax Crystal Structures Reveal How BH3
Domains Activate Bax and Nucleate Its Oligomerization to Induce Apoptosis. *Cell,* 152**,**
519-31.
DAI, H., SMITH, A., MENG, X. W., SCHNEIDER, P. A., PANG, Y. P. & KAUFMANN, S. H.
2011. Transient binding of an activator BH3 domain to the Bak BH3-binding groove
initiates Bak oligomerization. *J Cell Biol,* 194**,** 39-48.
DEWSON, G., KRATINA, T., CZABOTAR, P., DAY, C. L., ADAMS, J. M. & KLUCK, R. M.
2009. Bak activation for apoptosis involves oligomerization of dimers via their alpha6
helices. *Mol Cell,* 36**,** 696-703.
DEWSON, G., KRATINA, T., SIM, H. W., PUTHALAKATH, H., ADAMS, J. M., COLMAN, P.
M. & KLUCK, R. M. 2008. To trigger apoptosis Bak exposes its BH3 domain and homo-
dimerizes via BH3:grooove interactions. *Mol Cell,* 30**,** 369-380.
DEWSON, G., MA, S., FREDERICK, P., HOCKINGS, C., TAN, I., KRATINA, T. & KLUCK, R.
M. 2012. Bax dimerizes via a symmetric BH3:groove interface during apoptosis. *Cell Death*
Differ, 19**,** 661-70.
DU, H., WOLF, J., SCHAFER, B., MOLDOVEANU, T., CHIPUK, J. E. & KUWANA, T. 2011.
BH3 domains other than Bim and Bid can directly activate Bax/Bak. *J Biol Chem,* 286**,** 491-
501.
FERRER, P. E., FREDERICK, P., GULBIS, J. M., DEWSON, G. & KLUCK, R. M. 2012.
Translocation of a Bak C-terminus mutant from cytosol to mitochondria to mediate
cytochrome C release: implications for Bak and Bax apoptotic function. *PLoS One,* 7**,**
e31510.
GAHL, R. F., HE, Y., YU, S. & TJANDRA, N. 2014. Conformational rearrangements in the pro-
apoptotic protein, Bax, as it inserts into mitochondria: a cellular death switch. *J Biol Chem,*
289**,** 32871-82.
GARCIA-SAEZ, A. J. 2012. The secrets of the Bcl-2 family. *Cell Death Differ,* 19**,** 1733-40.

GAVATHIOTIS, E., REYNA, D. E., DAVIS, M. L., BIRD, G. H. & WALENSKY, L. D. 2010.
BH3-Triggered Structural Reorganization Drives the Activation of Proapoptotic BAX.
Molecular Cell, 40**,** 481-92.
GEORGE, N. M., EVANS, J. J. & LUO, X. 2007. A three-helix homo-oligomerization domain
containing BH3 and BH1 is responsible for the apoptotic activity of Bax. *Genes Dev,* 21**,**
1937-1948.
GILBERT, R. J. 2016. Protein-lipid interactions and non-lamellar lipidic structures in membrane
pore formation and membrane fusion. *Biochim Biophys Acta,* 1858**,** 487-99.
GRIFFITHS, G. J., DUBREZ, L., MORGAN, C. P., JONES, N. A., WHITEHOUSE, J., CORFE,
B. M., DIVE, C. & HICKMAN, J. A. 1999. Cell damage-induced conformational changes
of the pro-apoptotic protein Bak in vivo precede the onset of apoptosis. *Journal of Cell*
Biology, 144**,** 903-914.
GROSSE, L., WURM, C. A., BRUSER, C., NEUMANN, D., JANS, D. C. & JAKOBS, S. 2016.
Bax assembles into large ring-like structures remodeling the mitochondrial outer membrane
in apoptosis. *EMBO J,* 35**,** 402-13.
GRUNDLING, A., BLASI, U. & YOUNG, R. 2000. Biochemical and genetic evidence for three
transmembrane domains in the class I holin, lambda S. *J Biol Chem,* 275**,** 769-76.
HOCKINGS, C., ANWARI, K., NINNIS, R. L., BROUWER, J., O'HELY, M., EVANGELISTA,
M., HINDS, M. G., CZABOTAR, P. E., LEE, E. F., FAIRLIE, W. D., DEWSON, G. &
KLUCK, R. M. 2015. Bid chimeras indicate that most BH3-only proteins can directly
activate Bak and Bax, and show no preference for Bak versus Bax. *Cell Death Dis,* 6**,**
e1735.
IYER, S., BELL, F., WESTPHAL, D., ANWARI, K., GULBIS, J., SMITH, B. J., DEWSON, G. &
KLUCK, R. M. 2015. Bak apoptotic pores involve a flexible C-terminal region and
juxtaposition of the C-terminal transmembrane domains. *Cell Death Differ*.
KLUCK, R. M., ESPOSTI, M. D., PERKINS, G., RENKEN, C., KUWANA, T., BOSSY-
WETZEL, E., GOLDBERG, M., ALLEN, T., BARBER, M. J., GREEN, D. R. &
NEWMEYER, D. D. 1999. The pro-apoptotic proteins, Bid and Bax, cause a limited
permeabilization of the mitochondrial outer membrane that is enhanced by cytosol. *J Cell*
Biol, 147**,** 809-22.
LAZAROU, M., STOJANOVSKI, D., FRAZIER, A. E., KOTEVSKI, A., DEWSON, G.,
CRAIGEN, W. J., KLUCK, R. M., VAUX, D. L. & RYAN, M. T. 2010. Inhibition of Bak
activation by VDAC2 is dependent on the Bak transmembrane anchor. *J Biol Chem,* 285**,**
36876-83.
LEE, E. F., GRABOW, S., CHAPPAZ, S., DEWSON, G., HOCKINGS, C., KLUCK, R. M.,
GRAY, D. H., WITKOWSKI, M. T., EVANGELISTA, M., PETTIKIRIARACHCHI, A.,
BOUILLET, P., CORY, S., VANDENBERG, C. J., LANE, R. M., CZABOTAR, P. E.,
COLMAN, P. M., SMITH, B. J., KILE, B. T. & FAIRLIE, W. D. 2016. Physiological
restraint of Bak by Bcl-xL is essential for cell survival. *Genes Dev,* In press.
LEE, M. T., HUNG, W. C., CHEN, F. Y. & HUANG, H. W. 2008. Mechanism and kinetics of pore
formation in membranes by water-soluble amphipathic peptides. *Proc Natl Acad Sci U S A,*
105**,** 5087-92.
LEE, M. T., SUN, T. L., HUNG, W. C. & HUANG, H. W. 2013. Process of inducing pores in
membranes by melittin. *Proc Natl Acad Sci U S A,* 110**,** 14243-8.
LESHCHINER, E. S., BRAUN, C. R., BIRD, G. H. & WALENSKY, L. D. 2013. Direct activation
of full-length proapoptotic BAK. *Proc Natl Acad Sci U S A,* 110**,** E986-95.
LINDSTEN, T., ROSS, A. J., KING, A., ZONG, W., RATHMELL, J. C., SHIELS, H. A.,
ULRICH, E., WAYMIRE, K. G., MAHAR, P., FRAUWIRTH, K., CHEN, Y., WEI, M.,
ENG, V. M., ADELMAN, D. M., SIMON, M. C., MA, A., GOLDEN, J. A., EVAN, G.,
KORSMEYER, S. J., MACGREGOR, G. R. & THOMPSON, C. B. 2000. The combined
functions of proapoptotic Bcl-2 family members Bak and Bax are essential for normal
development of multiple tissues. *Mol Cell,* 6**,** 1389-1399.

LLAMBI, F., MOLDOVEANU, T., TAIT, S. W., BOUCHIER-HAYES, L., TEMIROV, J., MCCORMICK, L. L., DILLON, C. P. & GREEN, D. R. 2011. A unified model of mammalian BCL-2 protein family interactions at the mitochondria. *Mol Cell,* 44**,** 517-31.

LUO, L., YANG, J. & LIU, D. 2014. Integration and oligomerization of Bax protein in lipid bilayers characterized by single molecule fluorescence study. *J Biol Chem,* 289**,** 31708-18.

MA, S., HOCKINGS, C., ANWARI, K., KRATINA, T., FENNELL, S., LAZAROU, M., RYAN, M. T., KLUCK, R. M. & DEWSON, G. 2013. Assembly of the Bak apoptotic pore: a critical role for the Bak protein alpha6 helix in the multimerization of homodimers during apoptosis. *J Biol Chem,* 288**,** 26027-38.

MA, S. B., NGUYEN, T. N., TAN, I., NINNIS, R., IYER, S., STROUD, D. A., MENARD, M., KLUCK, R. M., RYAN, M. T. & DEWSON, G. 2014. Bax targets mitochondria by distinct mechanisms before or during apoptotic cell death: a requirement for VDAC2 or Bak for efficient Bax apoptotic function. *Cell Death Differ*.

MOLDOVEANU, T., LIU, Q., TOCILJ, A., WATSON, M. H., SHORE, G. & GEHRING, K. 2006. The x-ray structure of a BAK homodimer reveals an inhibitory zinc binding site. *Molecular Cell,* 24**,** 677-688.

MUELLER, M., GRAUSCHOPF, U., MAIER, T., GLOCKSHUBER, R. & BAN, N. 2009. The structure of a cytolytic alpha-helical toxin pore reveals its assembly mechanism. *Nature,* 459**,** 726-30.

NASU, Y., BENKE, A., ARAKAWA, S., YOSHIDA, G. J., KAWAMURA, G., MANLEY, S., SHIMIZU, S. & OZAWA, T. 2016. In Situ Characterization of Bak Clusters Responsible for Cell Death Using Single Molecule Localization Microscopy. *Sci Rep,* 6**,** 27505.

NECHUSHTAN, A., SMITH, C. L., LAMENSDORF, I., YOON, S. H. & YOULE, R. J. 2001. Bax and Bak coalesce into novel mitochondria-associated clusters during apoptosis. *Journal of Cell Biology,* 153**,** 1265-1276.

PANG, Y. P., DAI, H., SMITH, A., MENG, X. W., SCHNEIDER, P. A. & KAUFMANN, S. H. 2012. Bak Conformational Changes Induced by Ligand Binding: Insight into BH3 Domain Binding and Bak Homo-Oligomerization. *Sci Rep,* 2**,** 257.

SALVADOR-GALLEGO, R., MUND, M., COSENTINO, K., SCHNEIDER, J., UNSAY, J., SCHRAERMEYER, U., ENGELHARDT, J., RIES, J. & GARCIA-SAEZ, A. J. 2016. Bax assembly into rings and arcs in apoptotic mitochondria is linked to membrane pores. *EMBO J,* 35**,** 389-401.

SATSOURA, D., KUCERKA, N., SHIVAKUMAR, S., PENCER, J., GRIFFITHS, C., LEBER, B., ANDREWS, D. W., KATSARAS, J. & FRADIN, C. 2012. Interaction of the full-length Bax protein with biomimetic mitochondrial liposomes: a small-angle neutron scattering and fluorescence study. *Biochim Biophys Acta,* 1818**,** 384-401.

SCHINDELIN, J., ARGANDA-CARRERAS, I., FRISE, E., KAYNIG, V., LONGAIR, M., PIETZSCH, T., PREIBISCH, S., RUEDEN, C., SAALFELD, S., SCHMID, B., TINEVEZ, J. Y., WHITE, D. J., HARTENSTEIN, V., ELICEIRI, K., TOMANCAK, P. & CARDONA, A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods,* 9**,** 676-82.

SCHNEIDER, C. A., RASBAND, W. S. & ELICEIRI, K. W. 2012. NIH Image to ImageJ: 25 years of image analysis. *Nat Methods,* 9**,** 671-5.

SONG, L., HOBAUGH, M. R., SHUSTAK, C., CHELEY, S., BAYLEY, H. & GOUAUX, J. E. 1996. Structure of staphylococcal alpha-hemolysin, a heptameric transmembrane pore. *Science,* 274**,** 1859-66.

SUBBURAJ, Y., COSENTINO, K., AXMANN, M., PEDRUEZA-VILLALMANZO, E., HERMANN, E., BLEICKEN, S., SPATZ, J. & GARCIA-SAEZ, A. J. 2015. Bax monomers form dimer units in the membrane that further self-assemble into multiple oligomeric species. *Nat Commun,* 6**,** 8042.

SUNG, T. C., LI, C. Y., LAI, Y. C., HUNG, C. L., SHIH, O., YEH, Y. Q., JENG, U. S. &
CHIANG, Y. W. 2015. Solution Structure of Apoptotic BAX Oligomer: Oligomerization
Likely Precedes Membrane Insertion. *Structure,* 23**,** 1878-88.
SUZUKI, M., YOULE, R. J. & TJANDRA, N. 2000. Structure of Bax: coregulation of dimer
formation and intracellular localization. *Cell,* 103**,** 645-654.
TERRONES, O., ANTONSSON, B., YAMAGUCHI, H., WANG, H. G., LIU, J., LEE, R. M.,
HERRMANN, A. & BASANEZ, G. 2004. Lipidic pore formation by the concerted action of
proapoptotic BAX and tBID. *J Biol Chem,* 279**,** 30081-30091.
TRAN, V. H., BARTOLO, R., WESTPHAL, D., ALSOP, A., DEWSON, G. & KLUCK, R. M.
2013. Bak apoptotic function is not directly regulated by phosphorylation. *Cell Death Dis,* 4**,**
e452.
WEBER, K., HARPER, N., SCHWABE, J. & COHEN, G. M. 2013. BIM-mediated membrane
insertion of the BAK pore domain is an essential requirement for apoptosis. *Cell Rep,* 5**,**
409-20.
WEI, M. C., ZONG, W. X., CHENG, E. H., LINDSTEN, T., PANOUTSAKOPOULOU, V.,
ROSS, A. J., ROTH, K. A., MACGREGOR, G. R., THOMPSON, C. B. & KORSMEYER,
S. J. 2001. Proapoptotic BAX and BAK: a requisite gateway to mitochondrial dysfunction
and death. *Science,* 292**,** 727-730.
WESTPHAL, D., DEWSON, G., MENARD, M., FREDERICK, P., IYER, S., BARTOLO, R.,
GIBSON, L., CZABOTAR, P. E., SMITH, B. J., ADAMS, J. M. & KLUCK, R. M. 2014.
Apoptotic pore formation is associated with in-plane insertion of Bak or Bax central helices
into the mitochondrial outer membrane. *Proc Natl Acad Sci U S A,* 111**,** E4076-85.
WOLTER, K. G., HSU, Y. T., SMITH, C. L., NECHUSHTAN, A., XI, X. G. & YOULE, R. J.
1997. Movement of Bax from the cytosol to mitochondria during apoptosis. *Journal of Cell*
Biology, 139**,** 1281-1292.
ZHANG, Z., SUBRAMANIAM, S., KALE, J., LIAO, C., HUANG, B., BRAHMBHATT, H.,
CONDON, S. G., LAPOLLA, S. M., HAYS, F. A., DING, J., HE, F., ZHANG, X. C., LI, J.,
SENES, A., ANDREWS, D. W. & LIN, J. 2016. BH3-in-groove dimerization initiates and
helix 9 dimerization expands Bax pore assembly in membranes. *EMBO J,* 35**,** 208-36.
ZHANG, Z., ZHU, W., LAPOLLA, S. M., MIAO, Y., SHAO, Y., FALCONE, M., BOREHAM,
D., MCFARLANE, N., DING, J., JOHNSON, A. E., ZHANG, X. C., ANDREWS, D. W. &
LIN, J. 2010. Bax forms an oligomer via separate, yet interdependent, surfaces. *J Biol*
Chem, 285**,** 17614-17627.
ZHOU, L. & CHANG, D. C. 2008. Dynamics and structure of the Bax-Bak complex responsible for
releasing mitochondrial proteins during apoptosis. *J Cell Sci,* 121**,** 2186-96.

Figure 1 Following oligomerization, the Bak N-segment, α1 and α1-α2 loop become fully solvent-
exposed in contrast to the partially exposed core (α2-α5) and latch (α6-α9).

(A) Solvent exposure of hBak cysteine mutants was assessed by IASD labelling before (lane 2),
during (lane 3) and after (lane 4) treatment with tBid. Controls of unlabelled (untreated, lane 1) and
fully labelled (denatured, lane 5) Bak were included for comparison. Example IEF western blots are
shown.

(B) Quantitation of IASD labelling before and after treatment with tBid for the panel of previously
untested Bak residues. Data are mean ± SD (n≥3), or range (n=2), with n for each residue labelled
on the x-axis. IASD labelling data for residue V194C were from (Westphal et al., 2014) (denoted
#). Residues for which there is a significant difference in IASD labelling before versus after tBid
are in bold and underlined ($p < 0.05$).

(C) Heat map overview of Bak IASD labelling with tBid treatment from (B) pooled with published
analyses from (Westphal et al., 2014)(denoted #) or (Iyer et al., 2015)(denoted ^) and from
treatment with the tBid[Bax] chimera (denoted *; see also Figure S2).

(D) Schematic of Bak structural rearrangement from its non-activated monomeric state to the
activated dimer. Helices are numbered for the non-activated Bak. Note the complete solvent-
exposure of α1 and the α1-α2 loop in the activated dimer.

Figure 2

A   N-extremity

B   α2-α5 core

C   C-extremity

Figure 2 The N- and C-extremities of oligomerized Bak are mobile relative to the α2-α5 core dimer. Mitochondrial fractions from cells expressing the indicated Bak cysteine mutants were incubated with tBid to oligomerize Bak, and oxidant (CuPhe) added to induce disulphide bonds. Aliquots were analyzed by non-reducing SDS PAGE (upper panels) and BNP (lower panels), and immunoblotted for Bak to detect linked species.

(A) Cysteine linkage occurs throughout the N-extremity.

(B) Cysteine linkage occurs, but is less complete, in the α2-α5 dimer core.

(C) Cysteine linkage occurs in the C-extremity. (See also α6:α6 linkage in (Dewson et al., 2008) and α9:α9 linkage in (Iyer et al., 2015)).

Figure 3 Linkage constraints for different regions of the active Bak dimer support a flexible
extremity model for full-length Bak dimers in the MOM.

(A) The oxidant CuPhe induces linkage *between* Bak monomers or dimers, or *within* Bak dimers.
Correlation of BNP with non-reducing SDS PAGE can differentiate linkage within and between
dimers.

(B) Summary of linkage outcomes within or between dimers for residues in the Bak dimer N-
extremity.

(C) Linkage at the lateral corners of the Bak α2-α5 core dimer is highlighted with residue labels and
sticks on the crystal structure (4U2V (Brouwer et al., 2014)).

(D) Schematic of the activated Bak dimer at the membrane in side and top views. The range of
movement of the N- and C-extremities is indicated with coloured ovals (blue and pink respectively).

(E) Our linkage data are consistent with the dimensions of the Bak core dimer. The dimensions of
the Bak α2-α5 core dimer 4U2V (Brouwer et al., 2014) are shown in side view and top view. In the
crystal structure, the distance from one edge of the core at S68 over the bended structure of the
symmetric core to the opposing S68' is ~75 Å. Thus, for residues in the mobile N-extremity to link
within the core dimer, the two homotypic cysteine residues must bridge a distance of ~75 Å. This is
not feasible for the residue V61C which is ~30 Å from the symmetric core.

A

B

Figure 4 Bak dimer-dimer interactions are disrupted by detergent or Bax.

(A) Digitonin prevents linkage between Bak dimers. Membrane fractions expressing the indicated Bak single-cysteine variants were first treated with tBid, and then supplemented, as indicated, with detergent (1% digitonin) prior to cysteine linkage. Two additional mutants were included to show linkage within dimers at the BH3:groove interface (M71C/K113C) and at extensions to the C-terminus (GGSGGCK).

(B) Bax can disrupt Bak dimer-dimer association. Membrane fractions first treated with tBid[M97A] were then incubated, as indicated, with Bax, Bcl-x$_L$, Mcl-1ΔN151ΔC23 or digitonin prior to cysteine linkage (upper panels). Supernatant and pellet fractions showed partial translocation of each recombinant protein to the membrane fraction (lower panels).

Figure 5 V61C:V61C' linkage is a marker of dimer-dimer interaction.

(A) BNP analysis of V61C:V61C' linkage revealed no dimers, and a high proportion of 4x and >10x species. In contrast, H164C:H164C' linkage revealed a ladder of linked species.

(B) Densitometry of V61C:V61C' BNP linkage in Bak oligomers shows negligible dimers but reproducible linked species at 4x, 6x, 8x, and 10x, and a large population of species >10x. Densitometry data (n=9) were normalised to the area under the curve after alignment as described in Experimental Procedures. To correct for small variations in electrophoretic migration, we employed a noise reduction algorithm in which the 4x peak (0) and a minima at ~720 kDa (1) were aligned between replicates.

Figure 6 Bak dimer arrangement examined by two-dimensional stochastic simulations: a random
arrangement is the best fit for the V61C:V61C' BNP linkage densitometry.

(A) A grid of 100x100 was designated for the 2D simulations. Each unit hexagon within that grid
represented a Bak dimer, with the capacity to link to 2 neighbouring hexagons. The direction of
linkage from each hexagon was randomised. An example visual output from a 100x100 hexagonal
array is shown: neighbouring subsets of linked Bak dimers within the grid are delineated by
different colours.

(B) Linkage possibilities between each dimer unit in a hexagonal 2D array.

(C,D) Overlay (C), and average (D) of 30 predicted densitometry plots.

Figure 7

A

B

     i   ii   iii   iv

Figure 7 Bak dimers adopt various conformations on the membrane surface and aggregate in compact, disordered clusters to disrupt the mitochondrial outer membrane.

(A) The top view of Bak dimers lying in-plane on the mitochondrial surface. The α2-α5 core is bounded by the large oval. Extending from the core are the membrane anchored C-terminal helices α6, α7 and α8. For simplicity, the α9 transmembrane domains that project into the membrane plane, and the flexible, solvent exposed N-termini of each dimer are not shown.

(B) Growing clusters of Bak dimers induce membrane tension to rupture mitochondria.

 (i) Upon activation, Bak dimers penetrate the outer leaflet of the membrane and accumulate in a compact, irregular cluster.

(ii) More dimers converge on the cluster, thus enlarging the patch of membrane disturbance.

(iii) Once the patch attains a critical area, a non-lamellar lipidic arrangement is generated, relieving membrane tension.

(iv) Lipids and Bak dimers rearrange to bury exposed hydrophobic surfaces, yielding a variety of proteolipid (toroidal) "pores". Our assays survey a mixture of Bak linkage products derived from stages (ii), (iii) and (iv).

Figure S1

Figure S1 Bak cysteine variants retain apoptotic function. *Bak$^{-/-}$Bax$^{-/-}$* MEF expressing Bak cysteine variants were treated with 10 μM etoposide for 24 h and the percentage cell death quantified by propidium iodide uptake. Data are for unpublished mutants only and are the mean ± SD. The number of replicates (n) for each residue is labelled on the x-axis.

Figure S2 Quantitation of Bak IASD labelling before, during and after Bak activation.

(A,B,C). Labelling before, during or after treatment with tBid, segregated by Bak N-extremity (A),
Core (B), and C-extremity (C). Previously untested Bak residues have been pooled with data from
previous studies (# denotes data from (Westphal et al., 2014), ^ denotes data from (Iyer et al.,
2015)). Data are mean ± SD, or range (n=2), with n for each residue labelled on the x-axis.
Residues for which there is a significant difference in IASD labelling before versus after tBid are in
bold and underlined ($p<0.05$).

(D) IASD labelling for selected Bak variants before, during and after treatment with tBid^{Bax}
chimera (Hockings et al., 2015). Data are mean ± SD (n ≥ 3), or range (n=2), with n for each
residue labelled on the x-axis. No significant differences (p value cutoff of 0.05) in IASD labelling
before versus after tBid^{Bax} were observed.

Figure S3

Figure S3 Higher temperature enhances disulphide bonding of cysteine residues in α6 (H164C) and the α9 transmembrane domain (L199C). Mitochondrial fractions from cells expressing the indicated Bak cysteine mutants were incubated with tBid to oligomerize Bak, and oxidant (CuPhe) added to induce disulphide bonds. Samples were incubated with CuPhe for 15 min either at 0°C or at 30°C. Aliquots were analyzed by non-reducing SDS PAGE (upper) and BNP (lower), and immunoblotted for Bak to detect linked species. Note that linkage outside the membrane (H164C:H164C') was efficient even at 0°C.

Figure S4 Adjustment to simulation output to allow qualitative comparison to western blot data. Mathematical simulations of Bak dimer linkage generated frequency distributions of the size of Bak complexes (multiples of Bak dimers). These frequency distributions were then transformed to a predicted densitometry by multiplying each count by the number of Bak dimer units present in each linked Bak complex. For instance, the count representing 2x complexes, is multiplied by 1, the count representing 4x complexes is multiplied by 2, the count representing 6x complexes is multiplied by 3, etc. This predicted densitometry approximates the total relative abundance of Bak molecules in each linked Bak complex, which was then transformed with an exponential horizontal axis to mimic the nonlinear spacing of bands, and a gaussian kernel estimator to approximate the smearing of bands (especially at higher molecular weights) to allow qualitative comparison to western blot densitometry outputs.

Figure S5 2D simulation with reduced efficiency linkage successfully models reduced linkage in
mitochondrial experiments.

(A) Co-expression of Bak V61C with Bak Cys null resulted in less high order linkage species on
BNP. Membrane fractions expressing Bak V61C, FLAG-Cys null, or both, were incubated with or
without tBid, and then subjected to CuPhe linkage and analysis by BNP, non-reducing SDS PAGE
or reducing SDS PAGE.

(B) Densitometry of BNP linkage data following tBid treatment for 3 replicates of the V61C:Flag-
Cys null co-expression. Data are normalised to the area under the curve.

(C) Diagram showing the four types of dimers that would form upon co-expression of V61C and
Cys null variants of Bak (i.e. a system in which 50 % of Bak molecules are unable to link). Note
that dimers will be capable of 0, 1 or 2 linkages.

(D) The 2D simulation was altered to incorporate 50% linkage-incompetent Bak molecules. This
simulation yielded a good concordance with the V61C:Flag Cys null densitometry.

Figure S6 2D simulation with edge blocking probability successfully models mitochondrial
experiments in which Bak linkage becomes constrained as cysteines are positioned closer to the
dimer core.

(A) Residues between V61C and the core of the Bak dimer have a progressively more limited range
of movement. On the left, blue circles are shown on a schematic of the Bak dimer to represent the
range of movement for G4C, V61C, T62C, Q66C, S68C and S69C. On the right, an illustration of
the reduced overlap of Bak dimer extremities (blue circles) is shown for residues with progressively
smaller ranges of movement (e.g. V61C>Q66C>S69C).

(B) Densitometry of BNP outputs from V61C, T62C, Q66C, S68C and S69C highlight the reduced
linkage efficiency (relative to V61C) as cysteine substitutions approach the constrained core of the
Bak dimer.

(C) The 2D simulation was altered to incorporate an "edge blocking probability" to mimic the
increasing linkage constraint experienced by cysteine substitutions as they approached the core. The
best approximation of each cysteine substitution mutant is indicated by arrows.