TITLE**:** Factors affecting genetic connectivity and diversity of the island night lizard.

AUTHORS: Stephen E. Rice[1,2,*] and Rulon W. Clark[1]

1. San Diego State University, Dept of Biology, 5500 Campanile Drive, San Diego,

California, 92182

2. University of California Riverside, Dept of Evolution, Ecology, and Organismal

Biology, Riverside, California, 92521

* CORRESPONDING AUTHOR:

E-MAIL ADDRESSES:

Stephen E. Rice: srice001@ucr.edu

Rulon W. Clark: rclark@mail.sdsu.edu

RUNNING TITLE: Landscape genetics of the San Clemente Island night lizard.

Abstract

Habitat loss and fragmentation is one of the most severe threats to global

biodiversity. Because human development often fragments natural areas into habitat

"islands", studies which characterize the genetic structure of species isolated on oceanic

islands may provide insight into the management of anthropogenic habitat islands. The

San Clemente Island night lizard, *Xantusia riversiana reticulata*, is endemic to two

California Channel Islands, each with a history of anthropogenic disturbance. We

genotyped 917 individuals from San Clemente Island and Santa Barbara Island at 23

microsatellite loci to quantify population structure and identify natural and anthropogenic

landscape features affecting intra-island connectivity. We found significant, but shallow,

population structure on each island with sites < 400 m apart identified as distinct

genepools. Landscape genetic analyses identified conductive habitat as California

boxthorn and prickly pear cactus on both islands. Landscape features which decreased

connectivity were unique to each island and included natural and human-mediated

features. These results can inform management plans on each island by identifying

habitat targets for mitigation and restoration efforts designed to improve connectivity.

Our results highlight the need for considering fine-scale features correlated with

contemporary and historical patterns of fragmentation, especially in small and isolated

habitats on the mainland that may be analogous to oceanic islands.

KEYWORDS: Landscape genetics, Channel Islands, Fragmentation, Conservation

Introduction

Habitat fragmentation is a leading cause of global biodiversity loss.
Fragmentation can decrease metapopulation connectivity, increase genetic divergence,
and increase extinction risk (e.g. Frankham 2005; Sumner 2005; Vandergast et al. 2016).
Fragmented and isolated habitat patches often form "habitat islands", patches of suitable
habitat embedded within an inhospitable matrix, such as a nature preserve surrounded by
human housing developments (e.g. Hokit et al. 1999; Driscoll 2004; Wang et al. 2009).
When this occurs, the ecological effects of fragmentation can be further exacerbated by
management practices, as isolated fragments may be managed independently by different
agencies, which may result in variable outcomes in terms of conservation goals, even
when different fragments share common species and habitats. Thus, one goal of modern
conservation is to identify generalizable approaches which can be readily implemented
across administrative boundaries to improve the management of fragmented and isolated
habitat patches. This approach requires examining genetic connectivity and diversity both
within and between fragments, in order to distinguish features shared in common among
fragments that can impede or enhance connectivity, as well as features that may be
specific to each fragment. Although these types of comparisons are often made in the
context of actual oceanic islands, they are less commonly applied to habitat islands. Here,
we conduct a management case study of a species distributed across two oceanic islands
managed by different agencies. We believe this and other case studies can serve as
valuable exemplars for conservation approaches that could also be applied to species
distributed across habitat islands.

The California Channel Islands are small oceanic islands off Southern

California's coast which share the Mediterranean climate of the mainland and have

experienced habitat degradation and fragmentation due to anthropogenic activities (e.g.

Fellers and Drost 1991; Fellers et al. 1998; Sturgeon 2000). Different islands are

managed by different government agencies, although they share several species of

conservation concern in common. Thus characterizing the contemporary genetic patterns

of Channel Island endemics will also provide critical data for monitoring and

management directed toward island endemics (Schwartz et al. 2007; reviewed in Stetz et

al. 2011) and serve as a case study for habitat islands.

Reptiles are an excellent group for studies of fragmentation due to their

intermediate dispersal distances, relatively high population densities, plasticity of

responses, thermal physiology, and potential vulnerability to habitat modifications (e.g.

Hokit et al. 1999; Jordan and Snell 2008; Wang et al. 2009; Tseng et al. 2015). One of the

few reptiles endemic to the Channel Islands is the island night lizard, *Xantusia riversiana*

(Cope 1883). This species was delisted from the Endangered Species Act in 2014,

contingent on continued monitoring (United States Fish and Wildlife Service (USFWS)

2014). Two subspecies are recognized (Smith 1946), *X. r. reticulata,* the San Clemente

Island night lizard, endemic to San Clemente Island and Santa Barbara Island (Figure 1)

and *X. r. riversiana* endemic to San Nicolas Island. These subspecies differ in both life

history traits and habitat use (Fellers and Drost 1991; Fellers et al. 1998). On San

Clemente Island lizards are found primarily in habitats dominated by California boxthorn

(*Lycium californicum*) and prickly pear cactus (*Opuntia littoralis*), and avoid deep

canyons, sand dunes, and canyon woodland (Mautz 1993). On Santa Barbara Island,

lizards are found primarily in habitats dominated by California boxthorn and prickly pear

cactus, and tend to avoid grasslands (Fellers and Drost 1991). In prime habitat *X. r.*

reticulata may reach densities in excess of 3,200 individuals/ha (Fellers and Drost 1991;

Mautz 1993).

Although the island night lizard can reach very high population densities in prime

habitat, this long-lived species may be vulnerable to habitat fragmentation and

disturbance due to a combination of life history traits and very low dispersal distances

(Fellers and Drost 1991; Mautz 1993). As part of an ongoing post-delisting monitoring

effort, we characterized the genetic structure and connectivity of *X. r. riversiana* across

its entire geographic range, sampling sites spread across both San Clemente and Santa

Barbara Islands. We also characterized habitat, anthropogenic structures, and other

geographic features that may serve as potential barriers to population connectivity. Based

on large population sizes and near-continuous distribution of viable habitat, we

hypothesized that population genetic structure would be minimal on each island but

influenced by roads and other anthropogenic barriers, as well as non-preferred habitat

types. We also hypothesized that detected patterns of genetic differentiation would be

primarily correlated with Euclidean distance, due to limited dispersal (Fellers and Drost

1991; Mautz 1993).

Materials and Methods

Sample Collection and Genotyping

We collected samples on San Clemente Island (Figure 2) from February through

November of 2013 and Santa Barbara Island (Figure 3) from May through September

2015. Island night lizards were captured by turning cover items and setting live traps

along well-worn trails. When multiple lizards were encountered, we made attempts to

capture and sample all lizards. Target sample size for each collection site was a minimum

of 30 individuals (Hale et al. 2012). Samples were also collected opportunistically from

encountered individuals. Captured lizards were processed following the United States

Geological Survey herpetofaunal monitoring protocols (Fisher et al. 2008) and GPS

coordinates were recorded with an estimated accuracy of 5 m. Tissue samples consisted

of toes that were clipped at the distal knuckle, which also served as a unique 4-digit

identification, and a 10 mm segment from the tail tip. Tissues were preserved in 95%

ethanol and stored at -20°C.

We extracted DNA with a standard salt digestion and genotyped individuals at 23

polymorphic microsatellite loci arranged into 8 reactions (Rice et al. 2016; Rice 2017

Appendix A, B). We scored alleles twice with GENEMAPPER for consistency; loci missing

data or with inconsistent allele calls were reamplified and rescored to generate a

consensus genotype. We assessed per-allele error rates (Pompanon et al. 2005) by

extracting and genotyping 35 individuals on San Clemente Island and 16 individuals on

Santa Barbara Island a second time.

Quality Control and Summary Statistics

We removed individuals from analyses if amplification was successful at fewer

than 16 loci. We removed first-order relatives (full siblings, parent-offspring) from data

sets for each island to prevent biases in detected population structure. We inferred

pairwise relationships for samples within each collection site when at least 2 of 3

methods were concordant. The methods we chose for inference were COLONY (Jones and

Wang 2010), CERVUS (Kalinowsi et al. 2007), and the DyadML estimator as calculated in

COANCESTRY (Wang 2011). Individuals were removed from each data set to retain the

largest sample size. Final data sets with relatives removed were used in all further

analyses.

We evaluated the quality of each locus for each island at the level of collection

sites following Selkoe and Toonen (2006). Loci were removed if amplification occurred

in fewer than 75% of individuals, when in linkage disequilibrium (LD) with another

locus, when null alleles occurred in a majority of sampled sites, or when the assumption

of neutrality was violated. We examined loci for LD and conformance to Hardy-

Weinberg and random mating proportions (HWE) in GENEPOP (Rousset 2008) with p-

values corrected for multiple comparisons through sequential Bonferroni testing (Holm

1979). We assessed the presence of null alleles with MICROCHECKER (Van Oosterhout et

al. 2004). We examined the putative neutrality of loci with LOSITAN (Beaumont and

Nichols 1996; Antao et al. 2008) under the infinite alleles model and neutral Fst

estimation. We used the package DIVERSITY (Keenan et al. 2013) in the R statistical

environment and GENEPOP to generate summary statistics for each collection site and

each island (Table 1).

Population Structure

We delineated genepools on each island following Waples and Gaggiotti (2006).

We conducted an exact test for genetic differentiation for all pairs of collection sites in

GENEPOP. We combined probabilities with Fisher's method (Fisher 1932) with 0.0001 as

the lower bound for individual test probabilities; combined probabilities were corrected

using sequential Bonferroni testing. Collection sites were grouped if no significant

difference was found and the procedure was repeated until all pairwise comparisons were

significant.

We estimated genetic differentiation between collection sites with ≥ 20

individuals with Weir and Cockerham's (1984) estimate of Fst with 95% bias corrected

confidence intervals generated by 1,000 bootstraps over individuals in DIVERSITY. We

defined differentiation as significant when the confidence interval did not overlap 0.

We used the program STRUCTURE (Pritchard et al. 2000; Falush et al. 2003) to

estimate the number of populations on each island (K) using data from all collection sites.

We used values of K from 1 to 18 for San Clemente Island and 1 to 10 for Santa Barbara

Island. Analyses for both island consisted of 20 runs with a 500,000 burn-in period and

1,000,000 MCMC iterations under the admixture model with uncorrelated and correlated

allele frequency models (Pritchard et al. 2010). We used the Evanno et al. (2005) method

as implemented in STRUCTURE HARVESTER (Earl and vonHoldt 2012) to estimate the

number of populations. We generated ancestry matrices with the greedy algorithm in the

program CLUMPP (Jakobsson and Rosenberg 2007) to produce barplots with the program

DISTRUCT (Rosenberg 2004) for the best supported value of K for each island.

Isolation by Distance

We examined patterns of isolation by distance (IBD, Wright 1943) at the level of

collection sites for each island. We used non-parametric rank-based multiple regression

on distance matrices (MRDM, Lichstein 2007) using the function *MRM* in the R package

ECODIST (Goslee and Urban 2007). As an extension of the Mantel test framework, results

of MRDM are comparable to Mantel results with a single predictor matrix. Analyses

were ran with 10,000 permutations, pairwise Fst as the response variable, and pairwise

Euclidean distance as the predictor.

Landscape Features Affecting Connectivity

We used the isolation by resistance framework (IBR, McRae 2006) as

implemented in the program CIRCUITSCAPE (McRae and Beier 2007) to model

conductance and resistance of landscape features on each island. Resistance and

conductance were modeled separately to clearly delineate landscape features with

positive and negative correlations with pairwise genetic distances. Raster cell values for

each feature ranged from 2 (low), 50 (moderate), and 100 (high). These values were

chosen to identify candidate landscape features and attribute a coarse value to each for

modelling purposes only and were not chosen to represent the true and unknown values

of landscape features. Collection sites were represented as focal regions composed of the

minimum-spanning convex hulls generated from capture coordinates.

We used non-parametric MRDM with 10,000 permutations to assess the

categorical features for each island. Models ranged from single features to full models of

all statistically supported features. Single features entered the model-building framework

if their inclusion was significant and the coefficient of determination (R^2) was greater

than the null model. The null model consisted of bounded Euclidean distance wherein all

terrestrial raster cells had a value of 1. In full models, raster cells had either the sum of

resistance values for all features within the cell or 1 when no focal feature was present.

195   We used a consensus approach for model selection which considered the ranked-

order of best models across MRDM, corrected Akaike information criterion (AICc), and

marginal R^2_β (Edwards et al. 2008; Van Strien et al. 2012). We used the *lmer* function in

the R package LME4 (Bates et al. 2015) to construct maximum-likelihood population

effects (MLPE) models (Clarke et al. 2002) fit with residual maximum likelihood

(REML) estimation with collection site as a random factor. We used the *KRmodcomp*

function in the R package PBKRTEST (Halekoh and Højsgaard 2014) to calculate the

marginal R^2_β from the MLPE models. We calculated AICc with the R package MuMIn

(Bartoń 2016) for MLPE models fit without REML estimation, as variables which

contributed the fixed effects differed between models (reviewed in Van Strien et al.

2012).

206   Analyses of San Clemente Island focused on three categorical GIS layers:

vegetation type (RECON, Inc 2012), roadways type (Naval Facilities and Engineering

Command South West (NAVFACSW), date unknown), and canyon length

(NAVFACSW, date unknown). Vegetation was parsed into 11 classes based on least-cost

transect analysis groupings. Roadways were divided into two classifications: paved roads,

which served as primary traffic conduits along the spine of the island and within more

populated areas, and secondary roads, which were generally unpaved access roads.

Canyons were divided into three length classes of small (0-499 ft), medium (500-999 ft),

and large (≥ 1,000 ft). We applied a 5 m buffer to roadways and canyons to eliminate

breaks in the linear features when rasterized and produced maps with a 100 m resolution

Analyses of Santa Barbara Island focused on three layers: vegetation type

(National Park Service (NPS), 2010), presence of western seagull (*Larus occidentalis*)

nesting colonies, and presence of a potential predator, barn owls (*Tyto alba)*. There are no

roads on Santa Barbara Island. Vegetation was classified into 14 groups based on

dominant vegetation alliances; due to observations of woolly seablite (*Suaeda taxifolia*)

dominated habitat transitioning seasonally into crystalline iceplant (*Mesembryanthemum*

crystallinum) dominated habitat (pers. comm., Rodriguez 2016) we grouped these

alliances into a single class. Western seagull nesting colony presence was included by

digitizing seabird nesting locations from 2015 (NPS 2015). Potential barn owl presence

was included by extracting and digitizing home ranges from Figure 1 of Thomsen et al.

(2014). Resolution on vegetation data was listed as 1 ft (NPS 2010); however raster maps

were generated with a cell size of 5 m for all layers due to the uncertainty in GPS

coordinates.

Results

Quality Control

We sampled 605 island night lizards from 18 collection sites on San Clemente

Island and 312 island night lizards from 7 collection sites and 3 smaller opportunistic

areas on Santa Barbara Island. We removed 18 individuals from the San Clemente Island

data set due to missing data; no individuals were removed from the Santa Barbara Island

data set for this reason. We identified 75 first-order relationships among 114 individuals

on San Clemente Island and 65 first-order relationships among 97 individuals on Santa

Barbara Island. We removed 53 individuals from the San Clemente Island data set and 41

individuals from the Santa Barbara Island data set due to these relationships. These

removals resulted in sample sizes of 534 individuals from San Clemente Island and 271

from Santa Barbara Island.

Final data sets consisted of 20 loci for San Clemente Island and 21 loci for Santa

Barbara Island. A single locus was removed due to null alleles in 12 of 18 collection sites

on San Clemente Island; no loci were removed from the Santa Barbara Island data set due

to null alleles. The program LOSITAN identified 2 loci as under selection for San

Clemente Island and 2 different loci as under selection for Santa Barbara Island. These

loci were removed from further analyses within each respective data set. The per-allele

error rates associated with genotyping were 0.062% for San Clemente Island and 0.272%

for Santa Barbara Island. Rates of non-robust amplification were 7.14% for San Clemente

Island and 3.80% for Santa Barbara Island.

Summary Statistics

Sample sizes for San Clemente Island collection sites ranged from 12-70

individuals. Rarefied allelic richness for collection sites with ≥ 20 samples ranged from

6.28 to 11.0. Observed heterozygosity ranged from 0.62 to 0.88 whereas expected

heterozygosity ranged from 0.69 to 0.87. Collection site 8, which is a coastal site

bordered by ocean and active sand dunes on all but one side was the only site which

deviated from HWE and also had the lowest allelic richness, observed heterozygosity,

and expected heterozygosity (Table 1).

Sample sizes for Santa Barbara Island collection sites ranged from 21-60

individuals and 8-12 individuals for three opportunistic collection areas. Rarified allelic

richness for the 7 main collection sites ranged from 7.17 to 8.65. Observed

heterozygosity ranged from 0.76 to 0.89. Expected heterozygosity ranged similarly from

0.78 to 0.83. No significant departures from HWE were detected (Table 1).

Population Structure

All collection sites on San Clemente Island were identified as distinct genepools

with combined p-values from functionally 0 to 1.02×10^{-5}. Pairwise Fst values between

collection sites revealed subtle but significant structure on San Clemente Island (Table

S1, supplemental) with values ranging from 0.0022 to 0.1278. The largest values for Fst

involved collection site 8. The best supported clustering solution found by STRUCTURE

for San Clemente Island was K=2 (Figure 2).

All collection sites on Santa Barbara Island, except two, were distinct genepools

with p-values ranging from functionally 0 to 1.71×10^{-7}. Summary statistics for the

grouped sites (LC and GC) are presented independently and together in Table 1. Since

LC was an opportunistic site with few samples, LC and GC were not aggregated for

further analyses. On Santa Barbara Island pairwise Fst also showed subtle but significant

structure (Table S2, supplemental) with values ranging from 0.0199 to 0.0590.

STRUCTURE analyses with correlated and uncorrelated allele frequencies converged on

solutions of K=4 and K=7. We chose the solution of K=4 (Figure 3) as the K=7 solution

displayed clusters with minimal contributions.

Factors Positively Affecting Connectivity

On San Clemente Island, conductance models (Table 2) that included California

boxthorn (boxthorn) and prickly pear cactus (prickly pear) had the strongest support

across methods. Four models were equivalent in significance (p=0.0001, R^2 range:

0.6433-0.6602). Ranking these models by AICc led to equivalence between two models

that differed only by the inclusion of grasslands as low conductance; the model with the

lowest AICc (-957.56) and highest R^2_β value (0.7257) contained both grasslands and

small canyons as low conductance (Figure 4).

On Santa Barbara Island, conductance models (Table 3) which included boxthorn

and prickly pear were also the best supported across methods. Eight models had AICc

values within 2 units of the best model (AICc range -151.976 to -153.318). The R^2 for

conductance models ranged from 0.8596 to 0.8877. The model with the greatest R^2 and

R^2_β (0.8718) had both boxthorn and prickly pear with high conductance and giant

coreopsis (*Leptosyne gigantean*) as low conductance (p=0.0008; Figure 5). Statistical

methods did not show congruence among model rankings after the best supported model.

Factors Negatively Affecting Connectivity

We detected significant patterns of IBD between collection sites with ≥ 20

samples on both San Clemente Island (p=0.001, R^2=0.2060; Figure S3, supplemental)

and Santa Barbara Island (p=0.0174, R^2=0.4258, Figure S4, supplemental). On San

Clemente Island, initial analyses resulted in no detection of IBD (p=0.145,

R^2=0.0249).We attributed this to the non-linear and non-monotonic relationship between

Fst and Euclidean distance resulting from the strong signal of genetic differentiation

between collection site 8 and all other sites. Removal of this site as an outlier resulted in a

significant finding of IBD among the remaining collection sites.

IBR analyses of San Clemente Island were also sensitive to the inclusion of

collection site 8. We evaluated 4 models to determine if the observed patterns could be

attributed to collection site 8 as an outlier or active sand dunes acting as a barrier. The

best supported model placed active sand dunes as a complete barrier (p=0.0005,

R^2=0.2602, Figure S3, supplemental), followed by a distance-only model with collection

site 8 removed (p=0.0020, R^2=0.1121), collection site 8 removed and active sand dunes

as a complete barrier (p=0.0096, R^2=0.0848), and a bounded Euclidean distance (IBR

null) model with all collection sites (p=0.0149, R^2=0.0681, Figure S3, supplemental).

On San Clemente Island, resistance models (Table 2) with cholla phase maritime

desert scrub (cholla), canyons > 500 ft, and secondary roads were best supported across

methods. The best models contained the same features, preventing model selection by

AICc (AICc: -972.94 to -974.47). In the top performing models cholla, medium canyons,

and secondary roadways were moderate resistance whereas large canyons were moderate

to high resistance. The model best supported by MRDM (p=0.0001, R^2=0.5125) and R^2_β

(0.8606) contained cholla, medium canyons, and secondary roads with moderate

resistance and large canyons with high resistance (Figure 4).

The IBR null model on Santa Barbara Island was significant (p=0.0214,

R^2=0.4804; Figure S4, supplemental). Resistance models which included woolly seablite

(seablite) and crystalline iceplant (iceplant) as moderately resistant were the best

supported models (Table 3). Seven models had equivalent R^2 (0.8582) and highly

significant p-values (p < 0.001). Four of these models had AICc values of -149.043 to -

150.066 and differed only in the resistance value assigned to barren ground and the

inclusion of needle goldfields (*Lasthenia gracilis*). The model with the greatest R^2_β and

within the AICc range contained seablite and iceplant as low resistance with barren

ground as moderate (R^2_β=0.8867); however, this model had a lower R^2 value (0.6919).

We chose the model in which combined seablite and iceplant with barren ground and

common fiddleneck had moderate resistance (Figure 5) as increased resistance and

additional habitat types failed to greatly improve the model (Table 3).

Discussion

We used a single marker set to conduct a detailed genetic assessment of *X.*

riversiana on San Clemente Island and Santa Barbara Island to inform conservation and

management approaches across these independently managed islands. We used island-

wide sampling with 23 microsatellite loci to estimate baseline parameters for each

collection site, quantify population structure, and identify landscape factors correlated

with genetic distance on each island. We found island night lizards were admixed on each

island with shallow, but significant, population structure. Each island also shared

common patterns of isolation by distance, with California boxthorn and prickly pear

cactus identified as the most conductive habitat. Active sand dunes were identified as a

barrier to gene flow on San Clemente Island but are absent on Santa Barbara Island.

Unlike conductive habitat, habitat resistance patterns were unique to each island and

included both natural and anthropogenic features.

Population Structure

STRUCTURE identified 2 populations on San Clemente Island associated with the

northern and southern ends and with strong admixture across the island (Figure 2).

Surprisingly, STRUCTURE identified 4 populations on Santa Barbara Island despite its

much smaller size. These populations were strongly associated with the western

collection site (WP), northern collection site (AP), mid-island collection sites, and the

south-eastern collection sites (Figure 3). The western and northern populations on Santa

Barbara Island displayed relatively little admixture whereas other populations were

admixed.

It is notable that the global Fst estimates for San Clemente Island and Santa

Barbara Island were remarkably similar (0.0341 and 0.0346 respectively) despite large

differences in island size and sampling scale. Our detection of population genetic

structure on each island could result from the clustered sampling of a ubiquitously

distributed organism characterized by limited dispersal (Schwartz and McKelvey 2009).

However, congruence between methods corroborates the finding that multiple genepools

exist across each island even at distances less than 400 m.

Factors Positively Affecting Connectivity

California boxthorn and prickly pear cactus support the greatest abundances of

island night lizards and are prime habitat types subject to continued monitoring (USFWS

2014). Our analyses identified prime habitat as highly conductive on both islands, which

is supported by ecological data (Fellers and Drost 1991; Mautz 1993). Surprisingly,

grasslands were identified as conductive on San Clemente Island even though previous

research indicated that grasslands are poor habitat (Fellers and Drost 1991). Grasslands

and small canyons on San Clemente Island may be conductive due to the presence of

foraging resources and cover items. Cover items are important landscape features for

thermoregulation (Regal 1968; Fellers and Drost 1991) and any habitat with suitable

cover and resources may be conductive. For example, island night lizards on San Nicolas

Island can be found in beach habitat with little vegetation, so long as cover rocks are

present (Fellers et al. 1998). The Santa Barbara Island conductance model also included

giant coreopsis as weakly conductive. The distribution of this habitat type on Santa

Barbara Island is generally contiguous with prime habitat and difficult to analyze

independently.

Factors Negatively Affecting Connectivity

No habitat features were identified as resistant on both San Clemente Island and

Santa Barbara Island. This finding probably reflects differences in topography, the extent

of certain vegetation types, and history of anthropogenic development. Each island is thus

characterized by unique factors which correlate with increased effective distances

between collection sites and may constrain dispersal and connectivity. Active sand dunes

on San Clemente Island were the only habitat we identified as an absolute barrier to

dispersal, which is supported by Mautz (1993). There are no sand dunes present on Santa

Barbara Island. It is likely that active sand dunes on San Nicolas Island act as barriers as

well, although this hypothesis has yet to be assessed.

On San Clemente Island, cholla cactus may act as a resistant surface as the spines

readily pierce flesh and could injure or debilitate fleeing or dispersing individuals

(personal obs.). The southern bombardment area of San Clemente Island is of particular

interest as increased disturbance may further propagate cholla increasing its density and

distribution. If further fragmentation of populations within the southern portion of San

Clemente Island becomes a management concern, we recommend controlling or

minimizing the propagation of cholla through habitat restoration or more targeted training

activities. Cholla habitat occurs in sparse and small patches on Santa Barbara Island

likely resulting in the lack of congruent detection across islands.

On Santa Barbara Island, multiple habitat types were identified as resistant:

common fiddleneck, needle goldfields, barren ground, and seablite grouped with iceplant.

The identification of common fiddleneck and needle goldfields within similarly

supported resistance models are not consistent with the ecological study of Fellers and

Drost (1991) and could represent biases in the statistical optimization process. Notably,

grassland habitats were absent from the final resistance model despite Fellers and Drost

(1991) finding that grasslands required equivalent capture efforts to seablite and iceplant

habitats. It is likely that when cover is dense enough to cause refugial fissures in the soil

or supplemented by cover items, resistance may be minimized. Iceplant may render areas

unsuitable for other vegetation (Vivrette and Muller 1977) and may, as with barren

ground, limit the cover necessary for foraging, thermoregulation, and dispersal (Fellers

and Drost 1991). However, we could not determine the independent contributions of

seablite and iceplant habitats due to the spatial and temporal transitions between them

(pers. comm., Rodriguez 2016). Thus, on Santa Barbara Island, we recommend habitat

remediation efforts which prioritize iceplant removal and revegetation with prime habitat.

410   Canyons >500 ft in length and secondary roadways were also detected as resistant

411 on San Clemente Island. The association of canyons with increased genetic

412 differentiation between collection sites is supported by ecological research, and may be

413 due to both difficulty in crossing canyons and a lack of resources within them (Mautz

414 1993). Secondary roadways form extensive networks which fragment prime habitat of

415 boxthorn and prickly pear with barren ground, particularly along the western escarpment

416 of the island. Notably, the primary roadway was not identified as a resistant surface, even

417 though it is paved and has more traffic. However, this primary road is a relatively short

418 road running down the middle of the island and is unlikely to significantly influence

419 connectivity patterns. Thus, any efforts to mitigate fragmentation should be applied to

420 secondary roadways.

421 *Management Implications*

422   Management of species that persist on oceanic islands needs to examine the

423 conductance and resistance of habitat within islands and compare between them to derive

424 generalizable actions. As many management actions are spatially explicit, it is important

425 to characterize genetic patterns using spatially explicit methods (Shaffer et al. 2015). The

426 workflow used in our study can be generalized to examine connectivity patterns within

427 mainland ecosystems that are fragmented into habitat islands to inform management

428 actions within and between independently managed habitat fragments.

429   An exemplar region which could benefit from this approach is the peninsular

430 region of Southern California, which is separated into several administrative units tasked

431 with management and monitoring activities. This region may experience increased

extinction risks of plant and animal species (e.g. Lawson 2011) and genetic studies of

terrestrial vertebrates have revealed patterns of significant genetic differentiation and

decreased genetic diversity (Luckau 2015; Lion 2016). The modeling frameworks we

present can be applied to support management efforts of peninsular species through the

identification of fine-scale resistant features, for targeted mitigation efforts within each

management area, while also identifying key landscape features within and between

management areas which support connectivity. This combined approach may better

identify management strategies aimed at sustaining local populations, such as increasing

the area of prime habitat and improving the quality of the intervening matrix (e.g. Fahrig

2001).

The contemporary genetic patterns we identify for *X. riversiana* can inform

management and monitoring practices on each independently managed island. Common

management strategies across islands include restoring invasive vegetation to prime

habitat to support intra-island connectivity and increase population sizes. Improving

intra-island connectivity and population size may also increase species resilience to

potential threats of climate change or other catastrophic events by increasing genetic

diversity and decreasing the effects of demographic stochasticity. We encourage the

comparison of fine-scale patterns within and between regional management units to

identify common strategies and unique factors to inform mitigation efforts.

Funding

This work was supported by United States Department of Defense (W9126G-12-2-0060);

and the Southern California Research and Learning Center (S18309).

Acknowledgements

All capture data, individual genetic profiles, and protocols used to generate

microsatellite data may be found in SR's dissertation (Rice 2017). We would like to

thank the National Park Service for access to Santa Barbara Island and providing

landscape-level GIS layers, San Clemente Naval Base for access to San Clemente Island

and providing landscape-level GIS layers, and our field assistants. In addition we would

like to thank Drs. A. Bohonak, H. Regan, K. Anderson, J. Gatesy, and R. Rose for

comments received during manuscript preparation.

Literature Cited

Antao T, Lopes A, Lopes RJ, Beja-Pereira A, Luikart G (2008) Lositan: A workbench to detect molecular adaptation based on a Fst-outlier method. BMC Bioinformatics, 9, 323.

Bartoń K (2016) MuMIn: multi-model inference. R package, version 1.15.6. Available online: http://r-forge.r-project.org/projects/mumin/

Bates D, Maechler M, Bolker B, Walker S (2015) Fitting Linear Mixed-Effects Models Using lme4. Journal of Statistical Software, 67:1-48.

Beaumont MA, Nichols RA (1996) Evaluating loci for use in the genetic analysis of population structure. Proceedings of the Royal Society of London. Series B: Biological Sciences, 263, 1619-1626.

Clarke RT, Rothery P, Raybould AF (2002) Confidence limits for regression relationships between distance matrices: estimating gene flow with distance. Journal of Agricultural Biological and Environmental Statistics, 7, 361–372.

Cope ED (1883) Note on a species of Xantusia. In: Proceedings of the Academy of Natural Sciences of Philadelphia 1884 (ed Nolan EJ), pp. 29-32. Academy of Natural Sciences of Philadelphia, PA.

Driscoll D (2004) Extinction and outbreaks accompany fragmentation of a reptile community. Ecological Applications, 14, 220-240.

Earl DA, vonHoldt BM (2012) Structure Harvester: a website and program for visualizing structure output and implementing the Evanno method. Conservation Genetics Resources, 4, 359-361.

Edwards LJ, Muller KE, Wolfinger RD, Qaqish BF, Schabenberger O (2008) An R^2 statistic for fixed effects in the linear mixed model. Statistics in Medicine, 27, 6137-6157.

Evanno G, Regnaut S, Goudet J (2005) Detecting the number of clusters of individuals using the software structure: a simulation study. Molecular Ecology, 14, 2611-2620.

Fahrig L (2001) How much habitat is enough? Biological Conservation, 100, 65-74.

Falush D, Stephens M, Pritchard JK (2003) Inference of population structure using multilocus genotype data: linked loci and correlated allele frequencies. Genetics, 164, 1567-1587.

Fellers GM, Drost CA (1991) Ecology of the island night lizard, *Xantusia riversiana*, on Santa Barbara Island, California. Herpetological Monographs, 5, 28-78.

Fellers GM, Drost CA, Mautz WJ, Murphey T (1998) Ecology of the island night lizard, *Xantusia riversiana*, on San Nicolas Island, California. Prepared for the US Navy by the USGS, Western Ecological Research Center, Pt. Reyes Field Station, Pt. Reyes National Seashore, CA.

Fisher RA (1932) *Statistical Methods for Research Workers*. Oliver and Boyd, Edinburgh.

Fisher R, Stokes D, Rochester C, Brehme C, Hathaway S, Case T (2008) *Herpetological monitoring using a pitfall trapping design in southern California*. U. S. Geological Survey Techniques and Methods 2-A5.

Frankham R (2005) Genetics and extinction. Biological Conservation 126:131-140.

Goslee SC, Urban DL (2007) The ecodist package for dissimilarity-based analysis of ecological data. Journal Statistical Software, 22, 1-19.

Hale ML, Burg TM, Steeves TE (2012) Sampling for microsatellite-based population genetic studies: 25 to 30 individuals per population is enough to accurately estimate allele frequencies. PLoS One, 7, e45170.

Halekoh U, Højsgaard S (2014) A Kenward-Roger Approximation and Parametric Bootstrap Methods for Tests in Linear Mixed Models – The R Package pbkrtest. Journal Statistical Software, 59, 1-30.

Hokit GD, Stith BM, Branch LC (1999) Effects of landscape structure in Florida scrub: a population perspective. Ecological Applications, 9, 124-134.

Holm S (1979) A simple sequentially rejective multiple test procedure. Scandinavian Journal of Statistics, 6, 65-70.

Jakobsson M, Rosenberg NA (2007) Clumpp: a cluster matching and permutation program for dealing with label switching and multimodality in analysis of population structure. Bioinformatics, 23, 1801-1806.

Jones OR, Wang J (2010) Colony: a program for parentage and sibship inference from multilocus genotype data. Molecular Ecology Resources, 10, 551-555.

Jordan MA, Snell HL (2008) Historical fragmentation of islands and genetic drift in populations of Galapagos lava lizards (*Microlophus albemarlensis* complex). Molecular Ecology, 17, 1224-1237.

Kalinowski ST, Taper ML, Marshall TC (2007) Revising how the computer program CERVUS accommodates genotyping error increases success in paternity assignment. Molecular Ecology, 16, 1099-1106.

Keenan K, McGinnity P, Cross TF, Crozier WW, Prodohl PA (2013) Diversity: an R package for the estimation and exploration of population genetics parameters and their associated errors. Methods in Ecology and Evolution, 4, 782-788.

Lawson DM (2011) Multi-species conservation in the context of global change. Ph.D. Dissertation, University of California, Davis.

Lichstein JW (2007) Multiple regression on distance matrices: a multivariate spatial analysis tool. Plant Ecology, 188, 117-131.

Lion K (2016) A comparative study of genetic patterns in two closely related and sympatric Peromyscus species. MSc Thesis, San Diego State University.

Luckau TK (2015) Comparative conservation genetics of two sympatric lizard species across multiple landscapes in San Diego Country. MSc Thesis, San Diego State University.

Mautz WJ (1993) Ecology and energetics of the island night lizard, *Xantusia riversiana*, on San Clemente Island, California. In: Third California Islands symposium:

Recent advances in research on the California Islands (ed Hochberg FG), pp. 417-428. Santa Barbara Museum of Natural History, Santa Barbara, CA.

McRae, BH (2006) Isolation by resistance. Evolution, 60, 1551-1561.

McRae BH, Beier P (2007) Circuit theory predicts gene flow in plant and animal populations. Proceedings of the National Academy of Sciences USA, 104, 19885–19890.

Naval Facilities and Engineering Command South West (NAVFACSW)(Date Unknown) Spatial data developed by NAVFACSW.

National Park Service (NPS) (2010) Channel Islands National Park, Santa Barbara Island vegetation map final draft.

National Park Service (NPS)(2015) Channel Islands National Park, Santa Barbara Island seabird nesting locations.

Pompanon F, Bonin A, Bellemain E, Taberlet P (2005) Genotyping errors: causes, consequences and solutions. Nature Reviews Genetics, 6, 847-859.

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using multilocus genotype data. Genetics, 155, 945-959.

Pritchard JK, Wen X, Falush D (2010) Documentation for structure software: version 2.3.

RECON Environmental, Inc (2012) San Clemente Island vegetation mapping: How GIS helped two botanists map 36,000 acres in five field days. Poster accessible online: http://www.recon-us.com/services/biological-res/san_clemente_island/san_clemente_poster.pdf.

Regal PJ (1968) An analysis of heat seeking in a lizard. Ph.D. Dissertation, University of California, Los Angeles CA.

Rice SE (2017) Conservation Genetic Assessment of the Island Night Lizard, *Xantusia riversiana*, Under Contemporary and Future Environmental Conditions. Ph.D. Dissertation, San Diego State University, San Diego and University of California Riverside, Riverside CA.

Rice SE, Beasley RR, Lance SL, Jones KL, Clark RW (2016) Development of 24 polymorphic microsatellite markers for the Island Night Lizard (*Xantusia riversiana*) in Microsatellite records for volume 8, issue 2. Conservation Genetics Resources, 8, 169-196 doi: 10.1007/s12686-016-0549-4

Rosenberg NA (2004) Distruct: a program for the graphical display of population structure. Molecular Ecology Notes, 4, 137-138.

Rousset F (2008) Genepop'007: a complete re-implementation of the genepop software for Windows and Linux. Molecular Ecology Resources, 8, 103-106.

Shaffer HB, Gidiş M, McCartney-Melstad E, Neal KM, Oyamaguchi HM, Tellez M, Toffelmier EM (2015) Conservation genetics and genomics of amphibians and reptiles. Annual Review of Animal Biosciences, 3, 113-138.

Schwartz MK, McKelvey KS (2009) Why sampling scheme matters: the effect of sampling scheme on landscape genetic results. Conservation Genetics, 10, 441-452.

Schwartz MK, Luikart G, Waples R (2007) Genetic monitoring as a promising tool for conservation and management. Trends in Ecology and Evolution, 22, 25-33.

Selkoe KA, Toonen RJ (2006) Microsatellites for ecologists: a practical guide to using and evaluating microsatellite markers. Ecology Letters, 9, 615-629.

Smith HM (1946) A subspecies of the lizard Xantusia riversiana. Journal of the Washington Academy of Sciences, 36, 392–393.

Stetz, JB, Kendall KC, Vojta CD, GeM Working Group (2011) Genetic monitoring for managers: a new online resource. Journal of Fish and Wildlife Management, 2, 216-219.

Sturgeon WJ (2000) San Clemente Island: A chronological military history (1934-2000). Online: http://www.scisland.org/aboutsci/history/SCI-History.pdf

Sumner J (2005) Decreased relatedness between male prickly forest skinks (*Gnypetoscincus queenslandiae*) in habitat fragments. Conservation Genetics, 6, 333-340.

Thomsen, SK, Kroeger CE, Bloom PH, Harvey AL (2014) Space use and home-range size of barn owls on Santa Barbara Island. Monographs of the Western North American Naturalist, 7, 339-347.

Tseng S-P, Wang C-J, Li S-H, L S-M (2015) Within-island speciation with an exceptional case of distinct separation between two sibling lizard species divided by a narrow stream. Molecular Phylogenetics and Evolution, 90, 164-175.

United States Fish and Wildlife Service (USFWS) (2014) Island night lizard (*Xantusia riversiana*) final Post-Delistment Monitoring Plan. U.S. Fish and Wildlife Service, Carlsbad Fish and Wildlife Office, Carlsbad, California.

Van Oosterhout C, Hutchinson WF, Wills DP, Shipley P (2004) Micro-checker: software for identifying and correcting genotyping errors in microsatellite data. Molecular Ecology Notes, 4, 535-538.

Van Strien MJ, Keller D, Holderegger R (2012) A new analytical approach to landscape genetic modelling: least-cost transect analysis and linear mixed models. Molecular Ecology, 21,4010-4023.

Vandergast AG, Wood DA, Thompson AR, Fisher M, Barrows CW, Grant TJ (2016) Drifting to oblivion? Rapid genetic differentiation in an endangered lizard following habitat fragmentation and drought. Diversity and Distributions, 22, 344-357.

Vivrette NJ, Muller CH (1977) Mechanism of invasion and dominance of coastal grassland by *Mesembryanthemum crystallinum*. Ecological Monographs, 47, 301-318.

Wang J (2011) Coancestry: a program for simulating, estimating and analyzing relatedness and inbreeding coefficients. Molecular Ecology Resources, 11, 141-145.

Wang Y, Zhang J, Feeley KJ, Jiang P, Ding P (2009) Life-history traits associated with fragmentation vulnerability of lizards in the Thousand Island Lake, China. Animal Conservation, 12, 329-337.

Waples RS, Gaggiotti O (2006) What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. Molecular Ecology, 15, 1419-1439.

Weir BS, Cockerham CC (1984) Estimating F-statistics for the analysis of population structure. Evolution, 38, 1358-1370.

Wright S (1943) Isolation by distance. Genetics, 28, 114-138.

Figure 1. Location of Channel Islands inhabited by the island night lizard. The island night lizard is endemic to 3 Channel Islands: San Clemente Island, Santa Barbara Island, and San Nicolas Island.

Figure 2. San Clemente Island collection sites with coancestry proportions. All collection sites on San Clemente Island are shown. Pie charts represent the site-level coancestry proportions for the STRUCTURE solution K=2 (barplot).

Figure 3. Santa Barbara Island collection sites with coancestry proportions. All collection sites on Santa Barbara Island are shown. Pie charts represent the site-level coancestry proportions for the STRUCTURE solution K=4 (barplot).

Figure 4. San Clemente Island landscape effects. The landscape-level features correlated with genetic distance incorporated as a composite image. Active sand dunes were identified as a barrier (black). Landscape features in gray were neutral with respect to connectivity. Conductant features (California boxthorn, prickly pear cactus, grasslands, and canyons < 500ft in length) are shown in white. Resistant features (cholla phase maritime desert scrub secondary roadways, canyons > 500ft in length) are shown in dark gray.

Figure 5. Santa Barbara Island landscape effects. The landscape-level features correlated with genetic distance incorporated as a composite image. Landscape features in gray were neutral with respect to connectivity. Conductant features (California boxthorn, prickly pear cactus, and giant coreopsis) are shown in white. Resistant habitat (woolly seablite with crystalline iceplant, barren ground, and common fiddleneck) are shown in dark gray.

Table 1. Summary statistics for each island and collection site. Collection sites are listed under their respective islands with the total samples for that location (N) and those in the reduced data set (Nrr). Summary statistics were: rarified allelic richness for all collection sites (Ar), rarified allelic richness for only those sites with ≥ 20 individuals in the reduced data set (Ar (≥20)), expected heterozygosity (He), observed heterozygosity (Ho), HWE probabilities (HWE) with probabilities <0.0001 set to 0.0001 at the locus level. Island-level comparisons were generated from the reduced data sets, analyzed at the 18 shared loci. The inbreeding coefficient (Fis) was calculated by allele identity within GENEPOP overall and for each site. NA denotes analysis not applicable.

Site	N	Nrr	Ar	Ar (≥20)	He	Ho	HWE	Fis
San Clemente Island	609	534	21.47	NA	0.89	0.84	<0.0001	0.0459
1	33	31	8.98	10.35	0.86	0.87	0.8981	0.0137
2	39	33	8.78	10.08	0.86	0.86	0.2835	0.0148
3	86	71	9.31	11.0	0.86	0.87	0.1168	0.0027
4	30	25	8.53	9.77	0.86	0.87	0.6173	0.0019
5	28	24	8.59	9.92	0.86	0.88	0.3311	0.0018
6	31	29	8.24	9.50	0.84	0.86	0.4165	0.0044
HR	36	28	8.07	9.25	0.84	0.86	0.3063	-0.0008
7	32	30	8.21	9.49	0.84	0.83	0.2200	0.0317
8	27	20	5.46	6.28	0.69	0.62	0.0005	0.1157
10	30	30	9.20	10.80	0.86	0.87	0.1464	-0.0022
11	31	28	8.55	9.94	0.85	0.82	0.0089	0.0508
13	29	28	9.07	10.69	0.87	0.88	0.1188	0.0089
C	13	12	7.34	NA	0.82	0.86	0.4647	-0.0056
20	36	34	8.27	9.57	0.84	0.85	0.0838	0.0023
29	29	27	8.17	9.28	0.85	0.86	0.0488	0.0062
37	34	27	8.26	9.63	0.84	0.83	0.3470	0.0319
38	30	28	9.21	10.83	0.87	0.86	0.9637	0.0275
39	31	29	8.31	9.65	0.84	0.85	0.2572	0.0097
Santa Barbara Island	312	271	13.82	NA	0.86	0.82	<0.0001	0.0473
AP	28	21	5.85	7.20	0.80	0.80	0.2612	0.0292
CB	8	8	5.62	NA	0.79	0.83	0.8888	-0.0051
CC	31	24	6.44	8.18	0.83	0.83	0.3197	0.0086
GC	39	38	6.65	8.65	0.83	0.84	0.6265	0.0134
LC	9	8	5.97	NA	0.81	0.89	0.9830	-0.0208
NP	69	60	6.49	8.42	0.82	0.82	0.0227	0.0134
PP	37	33	6.28	7.87	0.82	0.81	0.1433	0.0251
S	13	12	5.92	NA	0.79	0.77	0.1239	0.0581
SE	35	31	6.56	8.36	0.83	0.85	0.6892	0.0026
WP	43	36	5.67	7.17	0.78	0.76	0.1836	0.0453
Aggregate LC+GC	48	46	6.77	8.88	0.84	0.84	0.3103	0.0089

Table 2. San Clemente Island competing conductance and resistance models. The best-supported conductance and resistance models are shown with the associated p-value (p) and coefficient of determination from non-parametric rank-based regressions (R^2) from the resulting MRDM analysis. The corrected AIC value (AICc) and R_β^2 value are also presented and were used in model selection. Model components are listed with the value assigned to each factor in parentheses.

	P	R^2	AICc	R_β^2
Null	0.0005	0.2602	-912.35	0.2825
Conductance Models				
Null + California boxthorn (100) + Prickly pear cactus (100)	0.0001	0.6554	-951.04	0.5630
Null + California boxthorn (100) + Prickly pear cactus (100) + Grassland (2)	0.0001	0.6433	-952.10	0.6769
Null + California boxthorn (100) + Prickly pear cactus (100) + Canyons < 500 ft (2)	0.0001	0.6602	-955.75	0.6116
Null + California boxthorn (100) + Prickly pear cactus (100) + Grassland (2) + Canyons < 500 ft (2)	0.0001	0.6461	-957.56	0.7257
Resistance Models				
Null + Cholla (50) + Secondary Roads (50) + Canyons 500-1000 ft (50) + Canyons > 1000 ft (50)	0.0002	0.5039	-974.47	0.8607
Null + Cholla (50) + Secondary Roads (50) + Canyons 500-1000 ft (50) + Canyons > 1000 ft (100)	0.0001	0.5125	-974.39	0.8606
Null + Cholla (50) + Secondary Roads (50) + Canyons 500-1000 ft (50) + Canyons > 1000 ft (50) + Canyons < 500 ft (2)	0.0002	0.5137	-973.19	0.8445
Null + Cholla (50) + Secondary Roads (50) + Canyons 500-1000 ft (50) + Canyons > 1000 ft (100) + Canyons < 500 ft (2)	0.0004	0.5175	-972.94	0.8427

Table 3. Santa Barbara Island competing conductance and resistance models. The best-supported conductance and resistance models are shown with the associated p-value (p) and coefficient of determination from non-parametric rank-based regressions (R^2) from the resulting MRDM analysis. The corrected AIC value (AICc) and R_β^2 value are also presented and were used in model selection. Model components are listed with the value assigned to each factor in parentheses.

	P	R^2	AICc	R_β^2
Null	0.0212	0.4804	-140.12	0.6911
Conductance Models				
Null + California Boxthorn(100)+ Prickly pear cactus (50) + Common fiddleneck (2)	0.0013	0.8611	-151.98	0.8614
Null + California Boxthorn(100)+ Prickly pear cactus (50) + Needle goldfields (2)	0.0003	0.8801	-151.98	0.8624
Null + California Boxthorn(100)+ Prickly pear cactus (2)	0.0002	0.8653	-152.12	0.8667
Null + California Boxthorn(50)+ Prickly pear cactus (2)	0.0012	0.8617	-152.36	0.8683
Null + California Boxthorn(100)+ Prickly pear cactus (50) + Giant Coreopsis (2)	0.0005	0.8677	-153.13	0.8716
Null + California Boxthorn(100)+ Prickly pear cactus (100)+ Giant Coreopsis (2)	0.0008	0.8877	-153.19	0.8718
Null + California Boxthorn(100)+ Prickly pear cactus (50) + Oatgrass (2)	0.0006	0.8693	-153.23	0.8709
Null + California Boxthorn(100)+ Prickly pear cactus (100) + Oatgrass (2)	0.0010	0.8693	-153.32	0.8713
Resistance Models				
Null + Seablite with iceplant (50) + Barren ground (100) + Common fiddleneck (50)	0.0003	0.8582	-149.69	0.8076
Null + Seablite with iceplant (50) + Barren ground (50) + Common fiddleneck (50)	0.0005	0.8582	-149.39	0.7490
Null + Seablite with iceplant (50) + Barren ground (100) + Common fiddleneck (50) + Needle goldfields (100)	0.0004	0.8582	-149.04	0.8371
Null + Seablite with iceplant (50) + Barren ground (50) + Common fiddleneck (50) + Needle goldfields (100)	0.0005	0.8582	-150.07	0.8704
Null + Seablite with iceplant (2) + Barren ground (50)	0.0027	0.6919	-150.22	0.8867