Genomic insights into temperature-dependent transcriptional responses of *Kosmotoga olearia*, a deep-biosphere bacterium that can grow from 20ºC to 79ºC

Stephen M. J. Pollo[1], Abigail A. Adebusuyi[1#], Timothy J. Straub[2], Julia M. Foght[1], Olga Zhaxybayeva[2,3] and Camilla L. Nesbø[1,4*]

[1]Department of Biological Sciences, University of Alberta, Edmonton, Alberta, Canada

[2]Department of Biological Sciences, Dartmouth College, Hanover, NH, USA

[3]Department of Computer Science, Dartmouth College, Hanover, NH, USA

[4]Centre for Ecological and Evolutionary Synthesis, Department of Biology, University of Oslo, Oslo, Norway

* Correspondence:

Corresponding Author

nesbo@ualberta.ca

[#]**Current address:** Vancouver Island University, 900 Fifth Street, Nanaimo, BC. Canada V9R 5S5

Keywords: Thermotogae, RNA-Seq, lateral gene transfer, temperature response, cold response, thermal adaptation

Abstract

Temperature is one of the defining parameters of an ecological niche. Most organisms thrive within a

temperature range that rarely exceeds ~ 30°C, but the deep subsurface bacterium *Kosmotoga olearia*

can grow over a temperature range of 59°C (20°C - 79°C). To identify genes correlated with this

flexible phenotype, we compared transcriptomes of *K. olearia* cultures grown at its optimal 65°C to

those at 30°C, 40°C, and 77°C. The temperature treatments affected expression of 573 of 2,224 *K.*

olearia genes. Notably, this transcriptional response elicits re-modeling of the cellular membrane and

changes in metabolism, with increased expression of genes involved in energy and carbohydrate

metabolism at high temperatures and up-regulation of amino acid metabolism at lower temperatures.

At sub-optimal temperatures, many transcriptional changes were similar to those observed in

mesophilic bacteria at physiologically low temperatures, including up-regulation of typical cold

stress genes and ribosomal proteins. Comparative genomic analysis of additional Thermotogae

genomes, indicate that one of *K. olearia*'s strategies for low temperature growth is increased copy

number of some typical cold response genes through duplication and/or lateral acquisition. At 77°C

one third of the up-regulated genes are of hypothetical function, indicating that many features of high

temperature growth are unknown.

Introduction

Microorganisms are capable of growing over an impressive temperature range, at least from -15°C to 122°C (Takai et al. 2008;Mykytczuk et al. 2013), and temperature is one of the most important physical factors determining their distribution, diversity, and abundance (Schumann 2009). However, individual microbial species grow only within a much narrower temperature interval. For example, *Escherichia coli* O157:H7 thrives in the laboratory between 19°C and 41°C (Raghubeer and Matches 1990), while *Geobacillus thermoleovorans* has a growth range of 37°C to 70°C (Dinsdale et al. 2011). Microorganisms with temperature ranges >50°C are rare and, to date, research into the few that have ranges >40°C has focused on psychrophiles (e.g. Mykytczuk et al. 2013). *Kosmotoga olearia* TBF 19.5.1 (hereafter referred to as *K. olearia*) is an anaerobic thermophile from the bacterial phylum Thermotogae with a growth range that spans almost 60°C (DiPippo et al. 2009). How does this organism achieve such physiological flexibility, and what are the evolutionary advantages and implications of this capability?

Fluctuations in temperature induce broad physiological changes in cells, including growth rate, alterations to cell wall and membrane composition, translation, and energy metabolism (Barria et al. 2013;Pollo et al. 2015;Schumann 2009). These physiological changes can be classified into two broad types of cellular response. Cold or heat *shock* designates the changes observed *immediately* after the shift of a culture to a lower or higher temperature, while *prolonged growth* at a specific lower or higher temperature elicits an *acclimated* low- or high-temperature response (Barria et al. 2013;Schumann 2009). Most studies of prokaryotes have focused on temperature shock responses rather than acclimated growth. Among the Thermotogae, responses to both heat shock and prolonged growth at high temperatures have been studied in the hyperthermophile *Thermotoga maritima*, which can grow between 55°C and 90°C (Pysz et al. 2004;Wang et al. 2012). During prolonged high-temperature growth *T. maritima* strongly up-regulates central carbohydrate metabolism genes and

expresses a few typical heat shock protein genes (Wang et al. 2012). Little is known about how *T.*

maritima responds to sub-optimal temperatures, although it encodes some genes implicated in cold

shock response. For example, its family of cold shock proteins (Csp), which are nucleic acid

chaperones known to be induced during cold shock and cold acclimation in mesophilic bacteria

(Barria et al. 2013;Phadtare 2004), exhibits nucleic acid melting activity at physiologically low

temperatures (Phadtare et al. 2003). Similarly, responses to cold shock in a few other thermophiles

involve many of the genes implicated in mesophilic cold shock response (e.g. Boonyaratanakornkit et

al. 2005;Mega et al. 2010). In this study, we systematically assessed bacterial physiological changes

associated with response to prolonged growth at both high and low temperature using *K. olearia* as a

model system. Such changes can reflect not only response to temperature itself, but also growth rate

effects and general responses to stress. By conflating these factors, our study examined overall

changes in *K. olearia*'s gene expression in the environments defined by a specific temperature.

K. olearia was isolated from a deep subsurface oil reservoir with *in situ* temperature of 68°C

(DiPippo et al. 2009). Cultivation experiments, 16S rRNA surveys, and metagenomic analyses have

confirmed that the principal natural habitat of *K. olearia* is deep subsurface sediments and that these

bacteria are particularly abundant in high temperature petroleum reservoirs (e.g. Vigneron et al.

2017;Duncan et al. 2009;Gittel et al. 2009;Pollo et al. 2016;Nesbø et al. 2010;Kotlar et al. 2011).

However, *K. olearia* has also been either isolated or detected via 16S rRNA sequencing in oil fields

(Nesbø et al. 2010), in an underground salt cavern plant used to store oily sand (Bordenave et al.

2013), and in engineered systems (Duncan et al. 2014;Oko et al. 2017) with *in situ* temperatures of

20°C–50°C, indicating that this bacterium occupies an ecological niche defined by a broad

temperature range.

The *K. olearia* genome (NC_012785) has 2,302,126 bp and is predicted to encode 2,224

genes (Swithers et al. 2011). Within the Thermotogae, genome size, intergenic region size, and

number of predicted coding regions correlate with the optimal growth temperature of an isolate

(Zhaxybayeva et al. 2012), with hyperthermophilic Thermotogae genomes being the most compact.

Phylogenetically, the Thermotogae order Kosmotogales comprises the genera *Kosmotoga* and

Mesotoga spp., the latter being the only described mesophilic Thermotogae lineage (Pollo et al.

2015). Assuming a hyperthermophilic last common ancestor of the Thermotogae (Zhaxybayeva et al.

2009), the Kosmotogales can be hypothesized to have acquired wide growth temperature tolerance

secondarily by expanding its gene repertoire. Moreover, it is likely that the ability of the

Kosmotogales common ancestor to grow at low temperatures enabled the evolution of mesophily in

Mesotoga (Pollo et al. 2015).

Such adaptations to novel environments can be greatly facilitated by lateral gene transfer

(LGT), since genes already "adapted" to the new conditions are readily available in the microbial

communities of the new environment (Boucher et al. 2003). For instance, LGT has been implicated

in adaptation to high temperature growth in hyperthermophilic bacteria, including *Thermotoga* spp.,

and to low temperature growth in archaea (López-García et al. 2015;Pollo et al. 2015;Boucher et al.

2003). Genome analysis of the mesophilic *Mesotoga prima* revealed that it laterally acquired 32% of

its genes after it diverged from other Thermotogae (Zhaxybayeva et al. 2012). Many of the predicted

gene donors are mesophiles, supporting the importance of lateral acquisition of genes already adapted

to mesophilic conditions in the evolution of *Mesotoga*.

To further gain insights into mechanisms of bacterial temperature response we sequenced 13

transcriptomes from isothermal cultures of *K. olearia* and examined transcriptional differences at

temperatures spanning its wide growth range (30°C, 40°C, 65°C, and 77°C). We also investigated the

importance of gene family expansion for adaptation of *K. olearia* to growth over a wide temperature

range via comparative genomic and phylogenetic analyses of identified temperature responsive genes

and their homologs in two newly sequenced *Kosmotoga* isolates, as well as in genomes of other

thermophilic and mesophilic Thermotogae.

Results

Temperature shifts and isothermic conditions elicit different growth patterns in *K. olearia*.

Under laboratory conditions in liquid anaerobic medium we observed growth of *K. olearia* at

temperatures as low as 25°C and as high as 79°C, with optimal growth at 65°C, defined as the

temperature affording the fastest growth rate (Fig. 1 and Fig. S1). Using a non-linear regression

model (Ratkowsky et al. 1983) we estimate a growth-permissive temperature range of 20.2 – 79.3°C,

consistent with the previously reported wide growth range of this isolate (DiPippo et al. 2009).

Interestingly, we were not able to cultivate *K. olearia* at temperatures near its range boundaries (30°C

and 77°C) by direct transfer from 65°C cultures. Instead, the growth temperature had to be changed

sequentially in ≤10°C increments. Particularly at the extremes, even small temperature shifts caused

both a longer lag phase and a slower growth rate compared to isothermal cultures (Fig. 1 and Fig.

S1). This phenomenon has been previously noted for mesophilic bacteria, especially for transitions

from high to low temperature (Swinnen et al. 2004). Our observations suggest that cells shifted to a

new temperature need to undergo large physiological changes that require time (i.e. an 'acclimation'

period (Barria et al. 2013)), and that these physiological challenges are too great to overcome when

temperature changes are large.

Architecture of the *K. olearia* transcriptome

Analysis of transcription start and stop sites predicted a minimum of 916 transcriptional units (TU) in

K. olearia (Supplemental material and Table S2), 52% of which consist of a single gene. This

fraction of single-gene TUs lies between the 65% reported for *E. coli* (Cho et al. 2009) and the 43%

recorded for *T. maritima*, which has also been shown to have a streamlined genome and a low-

complexity transcriptome (i.e. few sub-operonic transcripts and few genes with multiple start sites)

(Latif et al. 2013). The average TU length of ~2.39 genes in *K. olearia* is less than the 3.3 genes per

transcript of *T. maritima* (Latif et al. 2013) but closer to 2.2 genes per transcript in the mesophilic

firmicute *Geobacter sulfurreducens* (Qiu et al. 2010) and 1-2 genes per transcript in bacteria in

general (e.g. Cho et al. 2009)). Given that the *K. olearia* genome has more intergenic DNA than *T.*

maritima's genome (the ratio of the nucleotides located in non-coding vs. coding regions is 0.13 in *K.*

olearia and 0.06 in *T. maritima*), the shorter TU lengths in *K. olearia* may point to more flexible

transcriptional regulation.

Consistent energy generation across different temperature conditions

K. olearia produces ATP from pyruvate using a biochemically well-understood fermentation

pathway that generates hydrogen, carbon dioxide and acetate ((DiPippo et al. 2009); Fig. 2 and data

not shown). Since pyruvate was the carbon and energy source provided in all experiments, we

surveyed 51 genes predicted to be involved in core energy metabolism during growth on pyruvate.

Based on expression levels, we constructed a model that accounts for all major end products during

growth at 65°C (Fig. 2) and contains 15 genes with consistently high expression across all

temperature treatments (Fig. S2a and Table S3). In addition to indirectly validating the previously

known functional annotations of these genes, we furthermore propose that the most highly expressed

ABC-transporter gene cluster, Kole_1509 – 1513, encodes a pyruvate transporter (Fig. 2). Its current

annotation as a peptide ABC transporter may be erroneous, since most of the peptide ABC

transporters predicted in *T. maritima* using bioinformatics have been shown instead to bind and

transport sugars (Nanavati et al. 2006). Further functional studies of the transporter (e.g. binding

assays, expression with different substrates) are needed to confirm this hypothesis.

Identification of temperature-related transcriptional responses in *K. olearia*

Hierarchical clustering and Principal Component Analysis (PCA) separated the transcriptomes

according to temperature treatment (Fig. 3, Fig. S3 and Supplemental material), suggesting that the

observed changes in transcription are due to the culture growth temperature. Several genes with a

high correlation between their expression level and a specific growth temperature (Table S4 and

vectors in Fig. 3) are known to be involved in temperature response (Pollo et al. 2015). For example,

expression of the heat shock serine protease Kole_1599 positively correlated with the 77°C

transcriptomes (Fig. 3), where high expression of this protease was expected based on its

involvement in heat shock response in *T. maritima* (Pysz et al. 2004). Similarly, expression of the

cold shock protein genes Kole_0109 and Kole_2064 positively correlated with low temperature

growth (Fig. 3). Lastly, some observed changes presumably were due to the expected decreased

metabolic activity of the culture at sub- and supra-optimal temperatures. This can be exemplified by

the high expression and strong correlation of the alcohol dehydrogenase Kole_0742 (see

Supplemental material for further discussion of the expression pattern of this gene) and the central

carbon metabolism gene glyceraldehyde-3-phosphate dehydrogenase (Kole_2020) with the 65°C

transcriptomes.

Putative temperature-responsive genes were identified by pairwise comparisons of Illumina

transcriptomes from each temperature treatment to the optimal condition of 65°C (i.e. 30°C vs 65°C,

40°C vs 65°C, and 77°C vs 65°C). Across all comparisons 573 genes fulfilled our criteria for

temperature responsiveness ($\geq$ 2-fold difference in expression, > 20 reads per transcript, False

Discovery Rate < 0.05) with 430, 115, and 169 genes detected in the 30°C vs 65°C, 40°C vs 65°C,

and 77°C vs 65°C comparisons, respectively (Table S5). Expression of 306 of the 573 putative

temperature-responsive genes correlated with growth rate (r > |0.7|; Table S5). We used ANCOVA to

untangle the effect of growth rate from temperature, which revealed that growth rate was a significant

factor for only 60 of the 306 genes (p < 0.05; Table S5). Moreover, for 26 of these 60 genes

temperature was also a significant factor, and the simpler growth-rate-only model was rejected for all

but 13 of the 60 genes (Likelihood Ratio Test, FDR-corrected p-value < 0.05; Table S5). In an

alternative comparison based on relative effect size of temperature vs. growth rate, growth rate had

larger influence on expression of 51 of the 306 genes (Table S5). Among these genes were the

glyceraldehyde-3-phosphate dehydrogenase (Kole_2020) and the iron-containing alcohol

dehydrogenase (Kole_0742) mentioned above. Therefore, although we retained the designation of

"putatively temperature-responsive" for all 573 genes, below we focus on genes not primarily

affected by growth rate.

Distribution of temperature responsive genes across functional categories suggests a regulated

response to temperature

Most of the temperature-responsive genes were up-regulated when compared to the expression at

65°C (155 genes are down-regulated and 559 up-regulated; Table S5). One notable exception to this

overall trend were genes involved in carbohydrate and energy metabolism (Clusters of Orthologous

Groups [COG] categories C and G) where 32 genes were down-regulated at 30°C compared to 15

genes up-regulated. In transcriptomes at all non-optimal temperatures the list of putative temperature-

responsive genes was depleted in genes involved in translation (COG category J) and nucleotide

metabolism (COG category F) (Fig. 4) and conversely enriched in genes involved in signal

transduction (COG category T), and replication, recombination and repair (COG category L,

particularly at 30°C). Differential expression of the signal transduction genes (COG category T)

suggests the importance of these systems for regulating cellular responses at all tested temperatures.

Most of the identified COG category L genes are either mobile elements or CRISPR-associated

proteins, hinting at an increased activity of mobile genetic elements – a known common feature of

stress responses (Foster 2007). Additionally, at both 30°C and 77°C many genes encoding

transcription regulators (COG category K) are up-regulated, implying that prolonged growth at sub-

and supra-optimal temperatures results in detectable changes in transcriptional gene regulation in *K.*

olearia.

At 40°C there are pronounced differences in membrane fatty acid composition but no signs of

cold stress

Although the growth rate of *K. olearia* at 40°C is only one-third of that at the optimum 65°C (Fig. 1

and Fig. S1), clustering analysis suggested that the 40°C transcriptome was most similar to that at

65°C (Fig. 3 and Fig. S3). However, at 40°C even the four most highly expressed temperature-

responsive genes, including the growth-rate dependent alcohol dehydrogenase (Kole_0742), had

significantly lower expression levels (Table S5), reflecting the slower metabolic rate at lower

temperature growth. Yet, 94 of 115 putative temperature responsive genes were up-regulated (Table

S5, Fig. S2b), suggesting that slower metabolism is not the only explanation for the observed

transcriptional response to growth at 40°C.

Lipid metabolism (COG category I) appears to be particularly important at 40°C. For

instance, all the predicted fatty acid synthesis genes showed the highest expression at 40°C (Fig. S4),

with significantly higher expression of two genes involved in synthesis of unsaturated fatty acids

(Kole_0968) and initiation of fatty acid synthesis (Kole_0969). Biochemical analyses of total fatty

acids at 40°C and 65°C showed a much greater diversity of fatty acids at 40°C (Table S6), which

may explain the higher demand for the products of these genes at lower temperatures. Interestingly,

there was increased expression of a phosphate ABC transporter (Kole_0707 – Kole_0711, Table S5,

Fig. S2b), which may be linked to increased production of polar membrane lipids at moderately low

temperatures.

227   An enrichment of differentially expressed genes in "post-translational modification, protein

turnover, and chaperone function" (COG category O; the category that harbors genes related to

cellular stress) was due to both up- and down-regulation of the involved genes (Fig. 4). For instance,

K. olearia has three temperature-responsive peptidylprolyl isomerase (PPIase) genes: two PpiC-type

genes (Kole_1682 and Kole_0383) that are both highly expressed at 40°C, and one FKBP-type gene

(Kole_1745) that shows high expression at all temperatures except 77°C (Table S5). At lower

temperatures (e.g. 37°C), these enzymes catalyze proline isomerization, which happens

spontaneously at higher temperatures (Godin-Roulling et al. 2015). However, the enzymes known to

assist protein folding under cellular stress, chaperones (GroEL [Kole_1627] and Hsp70 [Kole_0886])

and protease Do (Kole_1599), were significantly down-regulated at 40°C (Table S5 and Fig. S2b).

Among other typical cold stress-related proteins, only one of *K. olearia*'s three cold shock proteins

(Kole_0109) showed significantly higher expression at 40°C, but its up-regulation was merely

moderate when compared to its expression levels at 30°C (Table S5 and Fig. S2b). Taken together,

the observed expression patterns of known cold stress-related genes are consistent with the cells

being in a non-stressed state at 40°C.

K. olearia is in cold stress at 30°C

Among the three most highly expressed up-regulated genes at 30°C are two Csp-encoding genes

(Kole_0109 and Kole_2064, Table S5, Fig. S2c), suggesting that the cells were in a cold-stressed

state during growth at 30°C. In support of this there was also significant up-regulation of other genes

previously linked to bacterial cold response (e.g. Alreshidi et al. 2015;Barria et al. 2013), including a

DEAD/DEAH-box RNA helicase (Kole_0922), *rbfA* (Kole_2103), *nusA* (Kole_1529) and ribosomal

proteins L10 (Kole_1840) and L7/L12 (Kole_1839). Genes encoding several additional ribosomal

proteins and ribosomal RNA (rRNA) methyltransferases, *rmlH* (Kole_1718) and *rmlL* (Kole_0897),

were also up-regulated (Fig. 3 and Table S5).

To detect a decrease in environmental temperature and elicit an appropriate regulatory

response, some bacteria have evolved two-component cold sensors (de Mendoza 2014). These signal

transduction systems consist of a sensor, a membrane-integrated protein with a kinase domain that

detects changes in the fluidity of the cell membrane, and the cytoplasmic response regulator, a

protein that induces expression of cold-responsive genes. In *K. olearia*, a histidine kinase with two

predicted transmembrane domains (Kole_1017) and two response regulators (Kole_1015 and

Kole_1016) showed a steady increase in expression as temperatures decreased from 65°C, but no

significant change in expression at 77°C (Table S5), leading us to hypothesize that these genes

encode a cold-sensing two-component system.

At 30°C, and to a lesser extent at 40°C, we also observed an over-representation of highly

expressed genes involved in amino acid metabolism (COG category E). Specifically, several genes in

the arginine (Kole_0092 – Kole_0097) and lysine (Kole_0104 – Kole_0107, 30°C only) biosynthesis

pathways were up-regulated, suggesting the potential for accumulation of peptides and amino acids

(or their intermediates) at lower temperatures. Interestingly, while the cells may accumulate peptides

at 30°C, at 40°C there was increased expression of an oligo-peptidase (Kole_1190) and genes

involved in lysine degradation (Kole_0958, Kole_0963 – Kole_0966). Such distinguishably different

metabolic responses to moderately low (40°C) and low (30°C) temperatures suggest a fine-tuned

temperature-dependent peptide turnover.

***K. olearia* is in heat stress at 77°C**

Since 77°C is near the upper limit for *K. olearia* growth under our laboratory conditions, we

hypothesize that the observed differences in expression profiles at 65 and 77°C would reflect a cell-

wide heat stress response. Of the 169 significantly differentially expressed genes, 119 showed

increased expression at 77°C (Table S5 and Fig. S2d). Among the most highly expressed genes were

those encoding the structural RNAs *ffs* (Kole_R0010), *ssrA* (Kole_R0006), and *rnpB* (Kole_R0049)

(Fig. 3), suggesting an increased rate of RNA turnover at supra-optimal temperature. Moreover,

genes involved in carbohydrate and energy metabolism (COG C and G categories) were over-

represented and up-regulated at 77°C (Fig. 4, Table S5). However, only two of the known heat *shock*

response genes (Pysz et al. 2004), the extreme heat stress sigma factor-24 (*rpoE*, Kole_2150) and the

heat shock protease (Kole_1599), were up-regulated, and 41 of 119 genes (34%) are annotated as

"hypothetical proteins", indicating that adaptation to growth at sustained high temperature remains

largely uncharacterized. Putative functions could be inferred for some of the encoded proteins of

these 41 genes of unknown function. Seventeen of the 41 proteins have predicted signal peptides,

suggesting that they may be secreted, and four have two or more predicted transmembrane regions,

suggesting they are membrane-associated. Kole_0652 carries a PrcB-domain, which interacts with

and stabilizes PrtP protease (Godovikova et al. 2010). Kole_1314 (and its paralog Kole_1297)

contains an AbiEii-toxin domain, and may be part of a toxin-antitoxin system. Furthermore, the 41

genes could be grouped into 34 transcriptional units, each containing at most three of these genes

(Table S2) and scattered across the *K. olearia* genome. Two of the genes, Kole_1430 and Kole_1431,

are co-transcribed with genes from a two-component system (Kole_1428 and Kole_1429), suggesting

they may be involved in environmental sensing or signaling. Three other co-transcribed genes

(Kole_1266, Kole_1267, Kole_1270) are found in a cluster containing CRISPR-genes, and two of

them (Kole_1266 and Kole_1270) contain RAMP-domains, suggesting CRISPR-related function

(Makarova et al. 2011). The majority of the 41 genes of unknown function have homologs in

genomes of other *Kosmotoga* spp. (N=38), *Mesotoga* spp. (N=23), and other Thermotogae (N=26).

Global regulators of temperature response

The transcriptional changes seen at the sub- and supra-optimal temperatures are likely to be

controlled by one or a few global regulators (Balleza et al. 2009), such as some temperature sensors

(de Mendoza 2014) and sigma factors needed for transcription initiation (Buck et al. 2000). The two-

component cold sensor up-regulated at low temperatures (Kole_1015 - Kole_1017) may represent

one such global regulator. Two different sigma factors (Kole_2150 and (Kole_1408) were

significantly up-regulated at 77°C and at 30°C and 40°C, respectively (Table S5), hinting at existence

of temperature-specific sigma factors. Kole_2150 belongs to the sigma-24 ECF subfamily, which is

activated in response to environmental stress (Balleza et al. 2009). Kole_1408 belongs to the sigma-

54 family, which is involved in enhancer-dependent transcription (Buck et al. 2000), introducing the

possibility that this sigma factor may be a target of the two-component cold sensor. In general, at

both 30°C and 77°C the differentially expressed genes were enriched in genes involved in

transcriptional regulation (COG category K) (Fig. 4), leaving a possibility of additional global

regulators.

Conservation of *K. olearia*'s temperature-responsive genes within the *Kosmotoga* genus

All genes that are required for adaptation and response of *K. olearia* strain TBF 19.5.1 to a wide

range of growth temperatures are expected to be present in other *K. olearia* isolates, whereas some

may be absent from *Kosmotoga* species that have a narrower spectrum of growth temperature.

Therefore, we contrasted the *K. olearia* genome to the genomes of *Kosmotoga* sp. DU53 and

Kosmotoga arenicorallina (Pollo et al. 2016). When compared to *K. olearia*, *Kosmotoga* sp. DU53

has a similar growth temperature range (25°C - 79°C, Table S7) and >99% average nucleotide

identity (ANI), while *K. arenicorallina* exhibits a narrower growth temperature range (35°C - 70°C,

Table S7) and has only 84% ANI.

Indeed, the *Kosmotoga* sp. DU53 genome lacks only 10 of the 573 *K. olearia* putative

temperature-responsive genes (BLASTP and TBLASTN searches, E-value $<10^{-3}$, Table S5). All 10

genes were expressed in *K. olearia* at relatively low levels, suggesting that they are unlikely to be

essential for high or low temperature growth. On the other hand, the *K. arenicorallina* genome does

not have detectable homologs of 103 of the 573 *K. olearia*'s putative temperature-responsive genes

(BLASTP and TBLASTN searches, E-value $<10^{-3}$; Table S5). The list of absent genes includes

several of the arginine and lysine biosynthesis genes that are up-regulated in *K. olearia* during

growth at 30°C, and seven of the genes of unknown function up-regulated at 77°C. Therefore, we

hypothesize that a subset of these 103 genes may play a role in extending the growth range of *K.*

olearia to ≤35°C and ≥70°C.

Many *key* temperature-responsive genes are laterally acquired

Obtaining "pre-adapted" genes from other genomes is one way prokaryotes adjust to new

environmental conditions (Boucher et al. 2003). Using HGTector (Zhu et al. 2014) we predicted that

354 of *K. olearia*'s 2,118 protein coding genes have been acquired laterally by *K. olearia* or the

Kosmotogales (i.e. *Kosmotoga* and *Mesotoga* genera), presumably representing LGT events

occurring after the divergence of Kosmotogales from other Thermotogae (Table S8). Eighty-eight of

the 354 genes were temperature responsive (Table S5, Fig. S5A and S5B), including several above-

discussed highly expressed genes (Table 1), and 37 of the 88 genes are shared with *Mesotoga* (Table

S5). Notably, 76% of these 37 genes are upregulated at 30°C (Fig. S5C), suggesting that their

acquisition may have been important in adaptation to low temperature growth. Among these are the

earlier mentioned rRNA methyltransferase genes (Kole_1718 and Kole_0897). The fatty acid

synthesis genes (Kole_0969 - Kole_0973) that are up-regulated at 40°C, as well as their

Kosmotogales homologs, form a distantly related sister clade to other Thermotogae lineages (Fig.

S6A), implying that these genes may have been acquired from an un-sampled lineage. Similarly, the

Csp-encoding gene highly expressed at 30°C (Kole_0109) is placed outside of the Thermotogae

clade (Fig. S6B).

Notably, some putative lateral gene acquisitions by *K. olearia* do not have homologs in

Mesotoga. These include genes encoding the predicted cold temperature sensor (Kole_1015 –

Kole_1017), one of the PPIase genes (Kole_1745), as well as the canonical cold response enzyme

DEAD/DEAH box RNA helicase (Kole_0922). Absence of these genes in *Mesotoga* suggests their

potential importance for *K. olearia*'s ability to grow over a wide temperature range.

Discussion

How can *K. olearia* grow at such a wide range of temperature?

Examination of *K. olearia*'s transcriptional response to sustained exposure to a non-optimal

temperature revealed both high expression of core metabolic genes and differential expression of

hundreds of other genes, with selected features highlighted in Fig. 5. At each tested temperature, core

metabolism genes for pyruvate utilization show high relative expression, which strongly suggests that

K. olearia uses the same proteins for its core energy metabolism and that these proteins can function

across its wide growth temperature range. In contrast, genes involved in regulatory functions showed

significant changes in expression at all temperatures, particularly at 30°C and 77°C (COG category K

and T in Fig. 4), suggesting that regulating gene expression is important in response to a temperature

shift. Among these genes we identified putative global temperature regulators: the two-component

cold sensor and temperature-specific sigma factors.

Close to *K. olearia*'s growth temperature maximum of 77°C, carbohydrate and energy

metabolism genes (COG categories C and G) were up-regulated (Fig. 4). It is unclear, however, if the

underlying cause is the increased turnover of enzymes at elevated temperatures, a demand for more

enzymes due to increased carbohydrate catabolism, or a combination of these factors. Increased

carbohydrate metabolism in response to prolonged growth at supra-optimal temperature has been

observed previously in *T. maritima* (Wang et al. 2012), and therefore may be a common adaptation to

high temperature growth in the Thermotogae. As observed for *K. olearia,* the prolonged supra-

optimal temperature growth of *T. maritima* also did not involve up-regulation of typical heat-shock

response proteins (Wang et al. 2012). This highlights the difference between cellular response to an

immediate heat-shock and to prolonged growth at supra-optimal temperature (Balleza et al. 2009),

and in general justifies classifying the cellular response to temperature into these two distinct

categories.

At the moderately sub-optimal growth temperature of 40°C, *K. olearia* cells face

physiological challenges of proper protein folding (Godin-Roulling et al. 2015), and of maintenance

of a functional cell membrane (de Mendoza 2014). Our observation that at 40°C, despite the lower

growth rate, lipid metabolism genes were among the most highly expressed genes suggests that

changes to the cell membrane composition are one of the most important adaptations for survival of

K. olearia at lower temperatures. Proper protein folding may require enzymatic assistance (Godin-

Roulling et al. 2015), which may be achieved via *K. olearia*'s three PPIases. The significant up-

regulation of the PPIase genes at 40°C suggest that they are particularly important at moderately low

temperatures where the cells are still relatively active. However, the overall lack of induction of

typical stress-related genes at 40°C, especially when compared to 30°C and 77°C, suggests that 40°C

is within the "Goldilocks" temperature range for *K. olearia.*

At 30°C *K. olearia* is clearly under cold stress (Fig. 5), as evidenced by expression of genes

known to be implicated in cold response. One of the strategies for maintenance of proper cellular

function at non-optimal temperatures (Pollo et al. 2015) is accumulation of compatible solutes.

Specifically, re-modelling of amino acid metabolism and possible accumulation of amino acids as

compatible solutes at low temperatures has been observed in bacteria (e.g. (Dahlsten et al.

2014;Ghobakhlou et al. 2015)). The up-regulation of many genes involved in amino acid metabolism

suggests that *K. olearia* may also accumulate amino acids or their intermediates for this purpose,

especially at 30°C. At 30°C there was also significant up-regulation of a citrate synthase gene

(Kole_1230), suggesting that *K. olearia* cells may accumulate citrate, as was observed in

Staphylococcus aureus during prolonged cold stress (Alreshidi et al. 2015). Alternatively, citrate

synthase, together with isocitrate dehydrogenase (Kole_1227), may be involved in converting

pyruvate or acetyl-CoA to 2-oxoglutarate, a precursor for several amino acids including arginine,

which has been suggested to accumulate in for instance *Clostridium* (Dahlsten et al. 2014).

Interestingly, the genome of strictly mesophilic *M. prima* encodes more genes involved in amino acid

metabolism than the genomes of *K. olearia* and other thermophilic Thermotogae (Zhaxybayeva et al.

2012). This supports the notion that the shift towards more amino acid metabolism in *K. olearia* may

be an adaptation to low temperature growth and suggests that the mesophilic *Mesotoga* have taken

this a step further by also *acquiring* more amino acid metabolism genes. Moreover, amino acid

metabolism genes are among the most numerous bacterial genes laterally acquired by mesophilic

archaea, which is hypothesized to reflect archaeal adaptation to low temperature growth (López-

García et al. 2015).

Ribosomes also need to be fine-tuned (i.e. to adjust the stoichiometric proportions of

accessory proteins) to function properly at low temperature (Barria et al. 2013). Consistently, we

observed a change in expression of several genes encoding ribosomal proteins. The most dramatic

differential expression, however, was observed for a ribosomal protein gene not yet connected to cold

response (L34; Kole_0258, Fig. 3, Table S5). L34, a bacteria-specific ribosomal protein hypothesized

to be a relatively recent addition to the evolving ribosome (Fox 2010), is required for proper

ribosome formation (Akanuma et al. 2014). A *Bacillus subtilis* mutant lacking the L34 gene showed

particularly slow growth at low temperature (Akanuma et al. 2012), suggesting a role for L34 in this

condition. We also observed significant increased expression of rRNA methyltransferases, and

methylation of rRNAs has been associated with responses to environmental stress, including

temperature (Baldridge and Contreras 2014). Therefore, we hypothesize that *K. olearia* modifies its

ribosome by changing stoichiometry of the ribosome components and by methylating rRNA. Time

required for such ribosomal adjustments could also explain the longer lag phase following

temperature shifts (Fig. S1).

Expansion of cold-responsive gene families may represent a common strategy for low

temperature adaptation, as has been noted in many bacteria, especially in psychrophiles (e.g. (Piette

et al. 2010). *K. olearia* exhibits the same trend. For instance, when compared to other Thermotogae,

all three analyzed *Kosmotoga* genomes harboured more copies of Csp-encoding genes (Table S9).

The observed gene family expansions, however, might be important not necessarily for low

temperature growth alone, but instead for growth over a wide temperature interval. *Mesotoga*

functions with only a single *csp* gene, demonstrating that having more copies of this gene is not

required for low temperature growth. Having several versions of a gene could make differential

regulation under different growth temperatures easier. For example, extra homologs (Kole_0111 and

Kole_0110) of the earlier-discussed putative cold sensor system do not show coordinated temperature

responses: Kole_0110 is up-regulated at 40°C, while Kole_0111 is up-regulated at 77°C (Table 1).

Therefore, these additional homologs may encode sensors tuned to different temperatures.

Evolutionary mechanisms that drive adaptive changes in Kosmotogales genomes

Gene family expansions can be achieved via within-lineage gene duplication or through LGT, and a

combination of these mechanisms appears to be at work in *K. olearia*. For example, even though

several Thermotogae genomes contain as many copies of PPIase genes as do *Kosmotoga* genomes

(Table S9), phylogenetic analysis suggests that in the Kosmotogales this gene family has only

recently been expanded by both LGT (the FKBP-type, Table 1) and duplication (the PpiC-type, Fig.

S6C). Similar conclusions can be made from the phylogenetic analyses of *csp* genes (Fig. S6B).

LGT appears to be a significant factor in evolution of low temperature growth in

Kosmotogales. In *K. olearia* although the proportion of transferred genes among the temperature

responsive genes is similar to what is observed in the genome (Fig. S5), many of the genes identified

here as key for low temperature growth have been acquired by LGT. For instance, the fatty acid

synthesis genes with high expression at 40°C were acquired by LGT into the Kosmotogales and are

likely important for low temperature growth in both *Mesotoga* and *Kosmotoga*. Other key LGT genes

including the DEAD/DEAH box RNA helicase, the temperature sensor, and the low temperature

induced *csp* genes, are only found in *Kosmotoga* spp. and may therefore be more important for its

wide temperature growth (Table 1). The predicted acquisition of fatty acid synthesis and *csp* genes by

(now mesophilic) archaea (López-García et al. 2015) additionally argues for the importance of these

genes in adaptation to low temperature growth.

Despite the importance of expanded gene families in low temperature adaptation, the role of

mutation and consequent natural selection on specific genes in response to changing environmental

conditions should not be neglected. For example, typical cold response genes *rbfA* (Kole_2103) and

nusA (Kole_1529) were not laterally acquired, but nevertheless show high expression only at 30°C.

Deciphering adaptive changes that occurred in such genes compared to thermophilic homologs may

elucidate molecular mechanisms of low temperature adaptation.

Why maintain the capacity for growth over such a wide temperature range?

Most bacteria are under selection to eradicate extraneous DNA (and genes) from their genomes

(Graur 2016), and among free-living bacteria Thermotogae in general have very compact genomes.

Kosmotogales, however, have notably larger genomes than other thermophilic Thermotogae (Pollo et

al. 2015;Zhaxybayeva et al. 2012), raising the possibility that expanded genomes are advantageous in

K. olearia's habitat. As discussed above, many of the genes in *K. olearia*, such as the cold-sensor

system, were expressed only at specific sub- or supra-optimal temperatures, and do not seem to be

important for growth at other temperatures (Table 1 and Table S5). The regulated response to low

temperatures and the preservation of the laterally acquired genes specifically expressed at 40°C and

30°C suggest that *K. olearia* encounters environments with very different temperatures frequently

enough to maintain these genes in its genome. Such environments may include oil reservoirs located

at different depths, as well as marine sediments influenced by the mixing of cold deep-sea water and

hydrothermal fluids (Sievert and Vetriani 2012). Perhaps, *K. olearia* migrates between such locations

via subsurface fluids and, as a result, may have been selected to become a temperature generalist.

Indeed, the environmental conditions of the subsurface environments and marine hydrothermal vents

from which *Kosmotoga* spp. have been detected and/or isolated vary substantially (DiPippo et al.

2009;Nunoura et al. 2010;L'Haridon et al. 2014;Vigneron et al. 2017;Duncan et al. 2009;Duncan et

al. 2014;Gittel et al. 2009;Kotlar et al. 2011;Pollo et al. 2016;Oko et al. 2017;Bordenave et al.

2013;Nesbø et al. 2010). Moreover, *Kosmotoga* sp. DU53, which is most similar to *K. olearia*, was

isolated from an oil reservoir with an *in situ* temperature of ~50°C, while *K. arenicorallina* was

obtained from hydrothermal sediments with a temperature of ~40°C (Nunoura et al. 2010). Notably,

K. olearia was also identified as a major constituent of a metagenome from a deep subsurface oil

reservoir with *in situ* temperature of 85°C and pressure of 25MPa (Kotlar et al. 2011). While the

reservoir temperature is higher than the maximum *K. olearia* growth temperature reported here,

elevated pressure could extend *K. olearia*'s temperature maximum, as has been demonstrated for

some archaea (e.g. Takai et al. 2008). Therefore, *K. olearia*'s growth temperature range under natural

conditions may be even broader than 20-79°C.

Conclusions

The present study demonstrates that even bacteria with relatively small genomes can use

transcriptional changes to respond effectively to large changes in temperature (Fig. 5). We showed

that *K. olearia*'s response to sustained exposure to a non-optimal temperature includes up-regulation

of hundreds of genes. Several key genes with known temperature-related functions apparently have

been acquired laterally, suggesting that LGT is an evolutionarily successful strategy for expansion of

temperature tolerance. However, gene duplication and subsequent neo-functionalization of the

paralogs likely also play an important adaptive role.

The ability of *K. olearia* to inhabit both high and low temperature environments suggests that

members of this lineage encounter environments with large temperature fluctuations and/or migrate

across ecological niches within the deep biosphere (e.g. between deep and shallow subsurface oil

reservoirs). Therefore, the subsurface environments, as well as their microbial populations, might be

viewed as a connected archipelago instead of isolated islands. As a corollary, we speculate that *K.*

olearia-like ecological generalists could also facilitate LGT among seemingly isolated deep

biosphere microbial communities adapted to a narrower ecological niche. For example, we have

previously demonstrated high levels of gene flow among hyperthermophilic *Thermotoga* populations

in subsurface oil reservoirs and marine hydrothermal vents (Nesbø et al. 2015), environments that are

separated by non-thermophilic surroundings but are hydrologically linked. The mechanism of such

gene flow is not yet known, but *K. olearia*-like Thermotogae capable of growing both in subsurface

oil reservoirs and adjacent marine sediments could serve as mediators of gene exchange.

Although some of the identified 573 temperature-responsive genes are already known to be

expressed in bacteria and archaea grown at high or low temperatures, most of the up-regulated genes

have not previously been implicated in temperature response and are in need of better functional and

biochemical characterization. Moreover, the majority of the *K. olearia* genes responsive to elevated

temperature encode proteins of unknown functions. Versatile proteins that work across a broad range

of temperatures also warrant further biochemical and evolutionary analyses, as understanding of their

enzymatic flexibility can aid the design of commercially important thermostable proteins.

Material and Methods

Bacterial culturing

K. olearia TBF 19.5.1 (DSM 21960(T), ATCC BAA-1733(T), Genbank accession number

NC_012785) was grown at different temperatures (30°C, 40°C, 65°C, and 77°C), but otherwise

optimal conditions, in *Kosmotoga* medium (KTM) using pyruvate as growth substrate as described in

(DiPippo et al. 2009). Cultures used as inocula were stored at 4°C, except those used for experiments

at ≥77°C. Actively growing cultures at temperatures ≥77°C had to be used directly as inoculum

because the cultures would not grow from inocula stored at either 4°C or room temperature (~22°C).

Replicate cultures received the same volume of inoculum; however, variable inoculum volume was

used at different temperatures (Table S1), as larger inoculum volumes were required to achieve

growth at the non-optimal temperature treatments.

Measurement of *K. olearia* growth at different temperatures

Growth curves were constructed from optical density measurements at 600 nm (OD_{600}) using an

Ultrospec 3100 pro. For cultures grown at 40°C, 65°C, 77°C, and 78°C two sets of triplicate bottles,

inoculated from the same inoculum 12 h apart, were monitored for a 12 h period per day to generate

the growth curves. The cultures for isothermic growth at 40°C, 65°C, 77°C, and 78°C were

monitored hourly, while the cultures for shifted growth at these temperatures were monitored every

1-5 hours. At 30°C one set of triplicate bottles was monitored once daily. Isothermic growth curves

were calculated from six replicates, except for the curves for 30°C and 77°C that had three and 12

replicates, respectively. All shifted growth curves consisted of six replicate cultures, except for 40°C

and 30°C which had four and three replicates, respectively. To determine growth rates (Fig. 1), for

each culture the $lnOD_{600}$ was plotted against growth time and the curve was fitted with a linear trend

line. The growth rate was defined as the slope at the log phase. To determine the time span of each

growth phase (Fig. S1), full composite growth curves were constructed using pooled replicate data.

For each curve, OD_{600} for all replicates was plotted against growth time and a polynomial regression

trend line was fitted.

Cultivation of *Kosmotoga* sp. DU53 and *K. arenicorallina*, and confirmation of their growth

temperature ranges

Kosmotoga sp. DU53 was grown in KTM as described above for *K. olearia*. *K. arenicorallina* was

also grown in KTM; however, maltose was used as substrate (2.5 mL and 0.5 mL 10% maltose was

added to serum bottles and Hungate tubes, respectively). One mL of culture was used as inoculum for

all cultures (bottles and tubes). The temperature range of each strain was assessed by examination of

culture turbidity as a proxy for growth. Starting from cultures grown at optimal growth temperature

(~ 65°C for *Kosmotoga* sp. DU53 and 60°C for *K. arenicorallina*), new cultures were shifted in

≤10°C increments. If growth was observed after a shift, then that culture was used to initiate a new

culture. The shifting procedure was terminated when growth was no longer evident at a given

temperature.

RNA isolation and processing

Cultures used for RNA extraction were inoculated from cultures that had been grown under the same

temperature conditions for at least three transfers. The time at which a culture was expected to be in a

desired growth phase was determined from the composite growth curves (Fig. S1), and was used as a

cell harvesting time (listed in Table S1). This procedure avoided exposure of the cultures to the lower

ambient temperatures in the laboratory during subsampling. For the 30°C cultures, OD_{600} was

additionally measured 24 h before harvesting to ensure the culture was in mid log phase. In order to

stabilize the transcripts and to avoid any transcriptional response to the change in temperature, an

equal volume of "stop solution" (10% phenol in ethanol) was added to the sealed cultures via syringe

immediately upon removal from the incubator. For each temperature treatment, RNA was extracted

in mid-log to late-log phase, using the Zymo Research Fungal/Bacterial RNA MiniPrep Kit

(Cedarlane Laboratories, Ltd.; Burlington, Ontario) and following the manufacturer's protocols

(Table S1).

Following recommendations in (Haas et al. 2012), we aimed to sequence ~3 million non-

ribosomal-RNA reads per sample. Ribosomal RNA (rRNA) depletion was performed on all samples

using the Ribo-Zero rRNA Removal Kit (Gram-Positive Bacteria) Magnetic Kit (MRZGP126,

Epicentre). On average, two aliquots containing 5ug total RNA (the maximum amount allowed by

the Ribo-Zero kit) needed to be rRNA-depleted and then pooled to generate sufficient input RNA

(200-500 ng for Ion Torrent PGM and 10-400 ng for Illumina MiSeq), although some samples

required as many as five rRNA depletions. Quality and quantity of the total RNA, as well as

efficiency of rRNA depletion, were assessed on an Agilent 2100 Bioanalyzer RNA Nano chip or

RNA Pico chip following the manufacturer's instructions for "Prokaryote Total RNA". RNA

successfully depleted of rRNA was used to construct RNA-Seq libraries following the

manufacturer's instructions, and sequenced on either an Ion Torrent PGM (RNA-Seq kit V2;

transcriptomes are labeled with an "IT" suffix) or an Illumina MiSeq (TruSeq RNASeq v2 2x100 bp).

The platform and RNA extraction technique used for each transcriptome are summarized in Table S1.

The transcriptomes are available in the Sequence Read Archive (http://www.ncbi.nlm.nih.gov/sra)

under the accession number SRP075860.

RNA-Seq analysis

For each transcriptome, sequenced reads were analyzed using the "RNA-Seq" function in CLC

Genomics Workbench version 7.0.4 (https://www.qiagenbioinformatics.com/). Briefly, reads were

first trimmed allowing no more than 2 ambiguous nucleotides and a maximum base-calling error

probability of 0.05. To remove sequences that matched remaining rRNA transcripts, the trimmed

reads were subjected to a relaxed mapping protocol: only rRNA genes were used as a "reference

genome", and reads were mapped only to the reference sequences if at least half of the alignment

(length fraction = 0.5) had at least 50% identity to the reference (similarity fraction = 0.5); all other

parameters were set to default (maximum number of hits for a read = 10, map to both strands,

mismatch cost = 2, insertion cost = 3, deletion cost = 3, auto-detect paired distances. The mapped

reads were designated as rRNA and were removed from further analysis.

The remaining reads were subjected to an RNA-Seq protocol with strict mapping parameters:

at least 95% of the alignment had to have at least 95% identity (i.e. similarity fraction = 0.95; length

fraction = 0.95) to the reference (the *K. olearia* TBF 19.5.1 annotated genome), mapping to

intergenic regions was allowed, and all other default settings as described above. Unmapped reads

were discarded. Expression levels for every gene were estimated using "Reads Per Kilobase of

transcript per Million mapped reads" (RPKM) values.

RPKM values for all genes are listed in Table S4. Differentially expressed genes were

identified by doing pairwise comparisons of Illumina transcriptomes of the isothermically grown

cultures at 30°C, 40°C, and 77°C to the cultures grown at the optimal temperature of 65°C. The

analyses used the "Empirical Analysis of DGE" function in CLC Genomics Workbench, which

employs the "Exact Test" for two-group comparisons (Robinson and Smyth 2008). A gene was

considered differentially expressed in a pairwise comparison if it had (i) > 20 reads in at least one of

the two transcriptomes, (ii) a statistically significant difference in the RPKM values (corrected for

multiple testing using False Discovery Rate [FDR] < 0.05), and (iii) a difference in RPKM values at

least two-fold in magnitude. Principal Component Analysis (PCA) and biplot visualization were

performed using R packages *ade4* and *bpca* respectively (Dray et al. 2007;Faria et al. 2013). Each

gene was assigned to a Clusters of Orthologous Groups (COG; (Galperin et al. 2015)) functional

category using the Integrated Microbial Genomes (IMG) portal (Markowitz et al. 2014). Genes

assigned to more than one COG category were counted in all assigned categories. Signal peptides and

membrane domains were identified in Geneious v.9.

Batch culture cannot selectively discern gene expression that is exclusively influenced by

temperature from expression that is solely growth rate-dependent. In theory, continuous culture

conducted at a single growth rate could afford such discrimination. However, given the extremely

slow growth of *K. olearia* near its temperature maxima and minima (Fig. 1), it was not feasible to use

anaerobic bioreactors to cultivate cells at this marginal growth rate across the temperature range.

Hence, in order to assess how differences in growth rates influence the expression patterns, we

examined correlations of the expression pattern of the putative temperature responsive genes with

growth rates calculated at each temperature. Pearson correlation of expression values and growth

rates were calculated in Microsoft Excel. ANCOVA, linear regression, and likelihood ratio tests were

carried out in R. The "growth rate only" model was the linear regression (expression = growth rate).

In the ANCOVA model growth rates was set as the quantitative variable and temperature as

qualitative variable (expression = growth rate + temperature). The growth rates used were 0.006 for

30°C, 0.087 for 40°C, 0.274 for 65°C and 0.107 for 77°C (see Fig. 1). When comparing the most

significant temperature coefficient to the growth rate coefficient, the latter was scaled by the average

growth rate. The growth rate effect was defined as being greater than the temperature rate effect if

|temperature/(growth rate * 0.115)| < 1.

Identification of genes involved in growth on pyruvate

K. olearia genes predicted to be involved in pathways for pyruvate conversion to acetate, CO_2, H_2

and ATP were retrieved from the KEGG (Kanehisa et al. 2014) and BioCyc (Caspi et al. 2014)

databases. Genes encoding hydrogenases were taken from (Schut et al. 2012), and genes encoding the

F-type ATPase subunits were identified using IMG (Markowitz et al. 2014).

Fatty acids analysis

Total lipids were extracted from *K. olearia* grown at 40°C to early stationary phase and 65°C to mid-

log, early stationary, late stationary and death phase by using methanol-chloroform (1:1 v/v). Fatty

acid methyl esters (FAME) were prepared from total lipids extracts using mild alkaline methanolysis

(Guckert et al. 1985). Dried FAME samples were re-dissolved in 300 µl chloroform (HPLC grade,

Fisher Scientific) and analyzed by gas chromatography with mass spectrometry (GC-MS) on an

Agilent 6890N gas chromatograph with a model 5973 inert mass selective detector (Agilent) fitted

with an Agilent HP-5MS capillary column (30 m × 0.25 mm ID, 0.25 µm film thickness; J + W

Scientific). Helium was used as the carrier gas with a temperature program of 130°C increasing at

3°C min^{-1} to 290°C and held for 2 min. Sample peaks were identified by comparison to Bacterial

Acid Methyl Ester Mix standards (Supelco, Sigma Aldrich) or on the basis of mass spectra and

expressed as a percentage of the total FAME quantified in each sample.

Comparative analyses of three *Kosmotoga* spp. genomes

The genome of *K. olearia* was compared to genomes of *Kosmotoga* sp. DU53 (accession number

JFHK00000000) and *K. arenicorallina* (accession number JGCK00000000) (Pollo et al. 2016) using

the IMG portal (Markowitz et al. 2014) and Geneious v.9. Specifically, genes were declared

homologous if they were significantly similar in BLASTP and TBLASTN (Altschul et al. 1997)

searches (E-value < 10^{-3}). For phylogenetic analyses, additional homologs of *K. olearia* genes were

retrieved from the NCBI non-redundant (*nr*) protein database and the IMG databases via BLASTP

searches, retaining the 100 top-scoring matches with E-value < 10^{-3}. Sequences were aligned using

MAFFT (Katoh et al. 2002), and phylogenetic trees were reconstructed using RAxML (Stamatakis

2006), as implemented in Geneious v. 9.1.3. Homologs from the recently released genome of

Kosmotoga pacifica (NZ_CP011232) (L'Haridon et al. 2014) were included in gene-specific

phylogenetic analyses. Pairwise Average Nucleotide Identity (ANI) (Goris et al. 2007) was

calculated using the Enveomics Toolbox (Rodriguez-R and Konstantinidis 2016).

Detection of laterally transferred genes

A customized version of HGTector (Zhu et al. 2014) (available through https://github.com/ecg-

lab/hgtector) was used to identify putatively transferred protein-coding genes in the *K. olearia*

genome. Homologs of each annotated protein-coding open reading frame (ORF) in the NC_012785

GenBank record were retrieved from a local copy of the NCBI *nr* database (downloaded November

21, 2014) using the BLASTP program from BLAST 2.2.28+ (Altschul et al. 1997). Sequences were

first filtered for low complexity regions using the *seg* algorithm. Then, only matches with E-value

$<10^{-5}$ and sequence coverage $\geq 70\%$ were retained. Database matches were expanded according to

the *MultispeciesAutonomousProtein2taxname* file from RefSeq release 68. This was necessary as

some genes across various taxonomic ranks were combined into a single entry in RefSeq, which

artificially decreased the representation of these genes in Close and Distal groups (see below), and

would confound downstream analysis. Taxonomic affiliation of each match was assigned using the

NCBI Taxonomy database (downloaded on November 21, 2014). Only 500 top-scoring matches

(after filtering for sequence coverage) were used as input for HGTector. The "Self" group was

defined as TaxID 651456 (genus *Kosmotoga*), and the "Close" group was defined as either TaxID

1184396 (genus *Mesotoga*, a sister group) or TaxID 2419 (order Thermotogales, comprising

Thermotoga, *Mesotoga*, and *Kosmotoga*). In either case, the "Distal" group comprised the remaining

taxonomic groups. The conservative cutoff (the median between the zero peak and the first local

minimum) was used for both the "Close" and "Distal" groups. A gene was designated as putatively

transferred if its "Close" score was below the cutoff and its "Distal" score was above the cutoff.

Putatively transferred genes with no top-scoring match in Thermotogae were designated as recent

transfer events into *K. olearia* (labelled "K" in Table S8). Putatively transferred genes for which the

difference between Close(Thermotoga) and Close(Mesotoga) scores was <1 were designated as gene

transfer events into Kosmotogales (i.e. *Kosmotoga* and *Mesotoga*; labelled "K+M" in Table S8).

Acknowledgments

This work was supported by a Research council of Norway award (project no. 180444/V40) to

C.L.N.; a Simons Investigator award from the Simons Foundation, a Dartmouth College Walter and

Constance Burke Research Initiation Award, and Dartmouth College start-up funds to O.Z.; and an

NSERC Alexander Graham Bell Canada Graduate Scholarship CGS-M to S.M.J.P. We thank

Rhianna Charchuk for technical assistance, Dr. Amanda J. Lohan for advice and discussion of RNA-

Seq methodology, Dr. Karen Budwill for head-space gas analyses, and Jennifer Franks for initial

discussions of lateral gene transfer detection methods.

References

Akanuma G, Nanamiya H, Natori Y, Yano K, Suzuki S, Omata S, Ishizuka M, Sekine Y, Kawamura

F (2012) Inactivation of ribosomal protein genes in *Bacillus subtilis* reveals importance of each

ribosomal protein for cell proliferation and cell differentiation. J Bacteriol 194: 6282-6291

Akanuma G, Kobayashi A, Suzuki S, Kawamura F, Shiwa Y, Watanabe S, Yoshikawa H, Hanai R,

Ishizuka M (2014) Defect in the formation of 70S ribosomes caused by lack of ribosomal protein L34

can be suppressed by magnesium. J Bacteriol 196: 3820-3830

Alreshidi MM, Dunstan RH, Macdonald MM, Smith ND, Gottfries J, Roberts TK (2015)

Metabolomic and proteomic responses of *Staphylococcus aureus* to prolonged cold stress. Journal of

proteomics 121: 44-55

Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997) Gapped

BLAST and PSI-BLAST: a new generation of protein database search programs. Nucleic Acids Res

25: 3389-3402

Baldridge KC, Contreras LM (2014) Functional implications of ribosomal RNA methylation in

response to environmental stress. Crit Rev Biochem Mol Biol 49: 69-89

Balleza E, López-Bojorquez LN, Martínez-Antonio A, Resendis-Antonio O, Lozada-Chávez I,

Balderas-Martínez YI, Encarnación S, Collado-Vides J (2009) Regulation by transcription factors in

bacteria: beyond description. FEMS Microbiol Rev 33: 133-151

Barria C, Malecki M, Arraiano CM (2013) Bacterial adaptation to cold. Microbiology 159: 2437-

2443

Boonyaratanakornkit BB, Simpson AJ, Whitehead TA, Fraser CM, El-Sayed NMA, Clark DS (2005)

Transcriptional profiling of the hyperthermophilic methanarchaeon *Methanococcus jannaschii* in

response to lethal heat and non-lethal cold shock. Environ Microbiol 7: 789-797

Bordenave S, Chatterjee I, Voordouw G (2013) Microbial community structure and microbial

activities related to CO_2 storage capacities of a salt cavern. International Biodeterioration and

Biodegradation 81: 82-87

Boucher Y, Douady CJ, Papke RT, Walsh DA, Boudreau MER, Nesbø CL, Case RJ, Doolittle WF

(2003) Lateral gene transfer and the origins of prokaryotic groups. Annu Rev Genet 37: 283-328

Buck M, Gallegos M, Studholme DJ, Guo Y, Gralla JD (2000) The bacterial enhancer-dependent

σ^{54} (σ^N) transcription factor

. J Bacteriol 182: 4129-4136

Caspi R, Altman T, Billington R, Dreher K, Foerster H, Fulcher CA, Holland TA, Keseler IM,

Kothari A, Kubo A, Krummenacker M, Latendresse M, Mueller LA, Ong Q, Paley S, Subhraveti P,

Weaver DS, Weerasinghe D, Zhang P, Karp PD (2014) The MetaCyc database of metabolic

pathways and enzymes and the BioCyc collection of pathway/genome databases. Nucleic Acids Res

42: D459-D471

Cho B, Zengler K, Qiu Y, Park YS, Knight EM, Barrett CL, Gao Y, Palsson BØ (2009) The

transcription unit architecture of the *Escherichia coli* genome. Nat Biotechnol 27: 1043-1049

Dahlsten E, Isokallio M, Somervuo P, Lindström M, Korkeala H (2014) Transcriptomic analysis of

(group I) *Clostridium botulinum* ATCC 3502 cold shock response. PloS one 9: e89958

de Mendoza D (2014) Temperature sensing by membranes. Annu Rev Microbiol 68: 101-116

Dinsdale AE, Halket G, Coorevits A, van Landschoot A, Busse HJ, de Vos P, Logan NA (2011)

Emended descriptions of *Geobacillus thermoleovorans* and *Geobacillus thermocatenulatus*. Int J

Syst Evol Microbiol 61: 1802-1810

DiPippo JL, Nesbø CL, Dahle H, Doolittle WF, Birkland N, Noll KM (2009) *Kosmotoga olearia*

gen. nov., sp. nov., a thermophilic, anaerobic heterotroph isolated from an oil production fluid. Int J

Syst Evol Microbiol 59: 2991-3000

Dray S, Dufour AB, Chessel D (2007) The ade4 Package - II : Two-table and K-table methods. R

News 7: 47-52

Duncan KE, Gieg LM, Parisi VA, Tanner RS, Tringe SG, Bristow J, Suflita JM (2009) Biocorrosive

thermophilic microbial communities in Alaskan North Slope oil facilities. Environmental Science

and Technology 43: 7977-7984

Duncan KE, Perez-Ibarra BM, Jenneman G, Harris JB, Webb R, Sublette K (2014) The effect of

corrosion inhibitors on microbial communities associated with corrosion in a model flow cell system.

Appl Microbiol Biotechnol 98: 907-918

Faria JC, Demétrio CGB, Allaman IB (2013) bpca: Biplot of multivariate data based on principal

components analysis. UESC and ESALQ, Ilheus, Bahia, Brasil and Piracicaba, Sao Paulo, Brasil

Foster PL (2007) Stress-induced mutagenesis in bacteria. Crit Rev Biochem Mol Biol 42: 373-397

Fox GE (2010) Origin and evolution of the ribosome. In: Deamer D, Szostak JW (eds) The Origins

of Life. Cold Spring Harb Perspect Biol, pp 2:a003483

Galperin MY, Makarova KS, Wolf YI, Koonin EV (2015) Expanded microbial genome coverage and

improved protein family annotation in the COG database. Nucleic Acids Res 43: D261-D269

Ghobakhlou A, Johnston A, Harris L, Antoun H, Laberge S (2015) Microarray transcriptional

profiling of Arctic *Mesorhizobium* strain N33 at low temperature provides insights into cold adaption

strategies. BMC Genomics 16: 383-396

Gittel A, Sørensen KB, Skovhus TL, Ingvorsen K, Schramm A (2009) Prokaryotic community

structure and sulfate reducer activity in water from high-temperature oil reservoirs with and without

nitrate treatment. Appl Environ Microbiol 75: 7086-7096

Godin-Roulling A, Schmidpeter PAM, Schmid FX, Feller G (2015) Functional adaptations of the

bacterial chaperone trigger factor to extreme environmental temperatures. Environ Microbiol 17:

2407-2420

Godovikova V, Wang HT, Goetting-Minesky MP, Ning Y, Capone RF, Slater CK, Fenno JC (2010)

Treponema denticola PrcB is required for expression and activity of the PrcA-PrtP (dentilisin)

complex. J Bacteriol 192: 3337-3344

Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM (2007) DNA-DNA

hybridization values and their relationship to whole-genome sequence similarities. Int J Syst Evol

Microbiol 57: 81-91

Graur D (2016) Prokaryotic Genome Evolution. In: Anonymous Molecular and Genome Evolution.

Sinauer Associates, Sunderland, MA, pp 451

Guckert JB, Antworth CP, Nichols PD, White DC (1985) Phospholipid, ester-linked fatty acid

profiles as reproducible assays for changes in prokaryotic community structure of estuarine

sediments. FEMS Microbiol Lett 31: 147-158

Haas BJ, Chin M, Nusbaum C, Birren BW, Livny J (2012) How deep is deep enough for RNA-Seq

profiling of bacterial transcriptomes? BMC Genomics 13: 734-745

Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M (2014) Data, information,

knowledge and principle: back to metabolism in KEGG. Nucleic Acids Res 42: D199-D205

Katoh K, Misawa K, Kuma K, Miyata T (2002) MAFFT: a novel method for rapid multiple sequence

alignment based on fast Fourier transform. Nucleic Acids Res 30: 3059-3066

Kotlar HK, Lewin A, Johansen J, Throne-Holst M, Haverkamp T, Markussen S, Winnberg A,

Ringrose P, Aakvik T, Ryeng E, Jakobsen K, Drabløs F, Valla S (2011) High coverage sequencing of

DNA from microorganisms living in an oil reservoir 2.5 kilometres subsurface. Environmental

Microbiology Reports 3: 674-681

Latif H, Lerman JA, Portnoy VA, Tarasova Y, Nagarajan H, Schrimpe-Rutledge AC, Smith RD,

Adkins JN, Lee D, Qiu Y, Zengler K (2013) The genome organization of *Thermotoga*

maritima reflects its lifestyle. PLoS genetics 9: e1003485

L'Haridon S, Jiang L, Alain K, Chalopin M, Rouxel O, Beauverger M, Xu H, Shao Z, Jebbar M

(2014) *Kosmotoga pacifica* sp. nov., a thermophilic chemoorganoheterotrophic bacterium isolated

from an East Pacific hydrothermal sediment. Extremophiles 18: 81-88

López-García P, Zivanovic Y, Deschamps P, Moreira D (2015) Bacterial gene import and mesophilic

adaptation in archaea. Nature reviews.Microbiology 13: 447-456

Makarova KS, Aravind L, Wolf YI, Koonin EV (2011) Unification of Cas protein families and a

simple scenario for the origin and evolution of CRISPR-Cas systems. Biology direct 6: 38-65

Markowitz VM, Chen IA, Palaniappan K, Chu K, Szeto E, Pillay M, Ratner A, Huang J, Woyke T,

Huntemann M, Anderson I, Billis K, Varghese N, Mavromatis K, Pati A, Ivanova NN, Kyrpides NC

(2014) IMG 4 version of the integrated microbial genomes comparative analysis system. Nucleic

Acids Res 42: D560-7

Mega R, Manzoku M, Shinkai A, Nakagawa N, Kuramitsu S, Masui R (2010) Very rapid induction

of a cold shock protein by temperature downshift in *Thermus thermophilus*. Biochem Biophys Res

Commun 399: 336-340

Mykytczuk NCS, Foote SJ, Omelon CR, Southam G, Greer CW, Whyte LG (2013) Bacterial growth

at -15°C; molecular insights from the permafrost bacterium *Planococcus halocryophilus* Or1. The

ISME journal 7: 1211-1226

Nanavati DM, Thirangoon K, Noll KM (2006) Several archaeal homologs of putative oligopeptide-

binding proteins encoded by *Thermotoga maritima* bind sugars. Appl Environ Microbiol 72: 1336-

1345

Nesbø CL, Kumaraswamy R, Dlutek M, Doolittle WF, Foght J (2010) Searching for mesophilic

Thermotogales bacteria: "Mesotogas" in the wild. Appl Environ Microbiol 76: 4896-4900

Nesbø CL, Swithers KS, Dahle H, Haverkamp TH, Birkeland N, Sokolova T, Kublanov I,

Zhaxybayeva O (2015) Evidence for extensive gene flow and *Thermotoga* subpopulations in

subsurface and marine environments. The ISME Journal 9: 1532-1542

Nunoura T, Hirai M, Imachi H, Miyazaki M, Makita H, Hirayama H, Furushima Y, Yamamoto H,

Takai K (2010) *Kosmotoga arenicorallina* sp. nov. a thermophilic and obligately anaerobic

heterotroph isolated from a shallow hydrothermal system occurring within a coral reef, southern part

of the Yaeyama Archipelago, Japan, reclassification of *Thermococcoides shengliensis* as *Kosmotoga*

shengliensis comb. nov., and emended description of the genus *Kosmotoga*. Arch Microbiol 192:

811-819

Oko BJ, Tao Y, Stuckey DC (2017) Dynamics of two methanogenic microbiomes incubated in

polycyclic aromatic hydrocarbons, naphthenic acids, and oil field produced water. Biotechnology for

Biofuels 10: 123

Phadtare S (2004) Recent developments in bacterial cold-shock response. Curr Issues Mol Biol 6:

125-136

Phadtare S, Hwang J, Severinov K, Inouye M (2003) CspB and CspL, thermostable cold-shock

proteins from *Thermotoga maritima*. Genes to Cells 8: 801-810

Piette F, D'Amico S, Struvay C, Mazzucchelli G, Renaut J, Tutino ML, Danchin A, Leprince P,

Feller G (2010) Proteomics of life at low temperatures: trigger factor is the primary chaperone in the

Antarctic bacterium *Pseudoalteromonas haloplanktis* TAC125. Mol Microbiol 76: 120-132

Pollo SMJ, Charchuk R, Nesbø CL (2016) Draft genome sequences of *Kosmotoga* sp. strain DU53

and *Kosmotoga arenicorallina* S304. GenomeA 4: e00570-16

Pollo SMJ, Zhaxybayeva O, Nesbø CL (2015) Insights into thermoadaptation and the evolution of

mesophily from the bacterial phylum *Thermotogae*. Can J Microbiol 61: 655-670

Pysz MA, Ward DE, Shockley KR, Montero CI, Conners SB, Johnson MR, Kelly RM (2004)

Transcriptional analysis of dynamic heat-shock response by the hyperthermophilic bacterium

Thermotoga maritima. Extremophiles 8: 209-217

Qiu Y, Cho BK, Park YS, Lovley D, Palsson B, Zengler K (2010) Structural and operational

complexity of the *Geobacter sulfurreducens* genome. Genome Res 20: 1304-1311

Raghubeer EV, Matches JR (1990) Temperature range for growth of *Escherichia coli* serotype

0157:H7 and selected coliforms in *E. coli* medium. J Clin Microbiol 28: 803-805

Ratkowsky DA, Lowry RK, McMeekin TA, Stokes AN, Chandler RE (1983) Model for bacterial

culture growth rate throughout the entire biokinetic temperature range. J Bacteriol 154: 1222-1226

Robinson MD, Smyth GK (2008) Small-sample estimation of negative binomial dispersion, with

applications to SAGE data. Biostatistics 9: 321-332

Rodriguez-R LM, Konstantinidis KT (2016) The enveomics collection: a toolbox for specialized

analyses of microbial genomes and metagenomes. Peer J Preprints 4: e19001

Schumann W (2009) Temperature sensors of eubacteria. Advances in applied microbiology 67: 213-

256

Schut GJ, Boyd ES, Peters JW, Adams MWW (2012) The modular respiratory complexes involved

in hydrogen and sulfur metabolism by heterotrophic hyperthermophilic archaea and their

evolutionary implications. FEMS Microbiol Rev 37: 182-203

Sievert S, Vetriani C (2012) Chemoautotrophy at Deep-Sea Vents: Past, Present, and Future.

Oceanography 25: 218-233

Stamatakis A (2006) RAxML-VI-HPC: Maximum likelihood-based phylogenetic analyses with

thousands of taxa and mixed models. Bioinformatics 22: 2688-2690

Swinnen IAM, Bernaerts K, Dens EJJ, Geeraerd AH, Van Impe J. F. (2004) Predictive modelling of

the microbial lag phase: A review. Int J Food Microbiol 94: 137-159

Swithers KS, DiPippo JL, Bruce DC, Detter C, Tapia R, Han S, Goodwin LA, Han J, Woyke T,

Pitluck S, Pennacchio L, Nolan M, Mikhailova N, Land ML, Nesbø CL, Gogarten JP, Noll KM

(2011) Genome sequence of *Kosmotoga olearia* strain TBF 19.5.1, a thermophilic bacterium with a

wide growth temperature range, isolated from the Troll B oil platform in the North Sea. J Bacteriol

193: 5566-5567

Takai K, Nakamura K, Toki T, Tsunogai U, Miyazaki M, Miyazaki J, Hirayama H, Nakagawa S,

Nunoura T, Horikoshi K (2008) Cell proliferation at 122°C and isotopically heavy CH_4 production

by a hyperthermophilic methanogen under high-pressure cultivation. Proc Natl Acad Sci U S A 105:

10949-10954

Vigneron A, Alsop EB, Lomans BP, Kyrpides NC, Head IM, Tsesmetzis N (2017) Succession in the

petroleum reservoir microbiome through an oil field production lifecycle. The ISME Journal : 1-14

Wang Z, Tong W, Wang Q, Bai X, Chen Z, Zhao J, Xu N, Liu S (2012) The temperature dependent

proteomic analysis of *Thermotoga maritima*. PloS one 7: e46463

Zhaxybayeva O, Swithers KS, Foght J, Green AG, Bruce D, Detter C, Han S, Teshima H, Han J,

Woyke T, Pitluck S, Nolan M, Ivanova N, Pati A, Land ML, Dlutek M, Doolittle WF, Noll KM,

Nesbø CL (2012) Genome sequence of the mesophilic Thermotogales bacterium *Mesotoga prima*

MesG1.Ag.4.2 reveals the largest Thermotogales genome to date. Genome Biol Evol 4: 700-708

Zhaxybayeva O, Swithers KS, Lapierre P, Fournier GP, Bickhart DM, DeBoy RT, Nelson KE, Nesbø

CL, Doolittle WF, Gogarten JP, Noll KM (2009) On the chimeric nature, thermophilic origin, and

phylogenetic placement of the *Thermotogales*. Proc Natl Acad Sci U S A 106: 5865-5870

Zhu Q, Kosoy M, Dittmar K (2014) HGTector: An automated method facilitating genome-wide

discovery of putative horizontal gene transfers. BMC Genomics 15: 717

Table 1. Gene expression in selected laterally acquired temperature-responsive genes. At each

temperature, the listed RPKM values represent the average expression levels across replicates.

Values that are significantly different from 65°C are shown in bold font. None of the temperature

responsive expression patterns show significant effect of growth rate and, where tested (i.e. |R| > 0.7),

the growth-rate-only model was rejected.

Locus Tag	Functional annotation	RPKM values				Identified by
		30°C[a]	40°C	65°C	77°C	
Kole_0109	Cold shock protein	**5602**	**892**	222	119	Phylogenetic analysis[a]
Kole_0110	Histidine kinase	175	**333**	144	312	Phylogenetic analysis
Kole_0111	Response regulator	166	204	173	**446**	HGTector
Kole_0505	Glycerol dehydrogenase	721	**2668**	752	1242	HGTector
Kole_0506	Hypothetical protein	559	**2037**	461	783	Phylogenetic analysis
Kole_0507	Hypothetical protein	555	**2193**	521	809	HGTector
Kole_0508	Poly (3-hydroxybutyrate) depolymerase-like protein	212	423	200	314	HGTector
Kole_0897	Ribosomal RNA methyltransferase, rmlL	**503**	498	228	232	HGTector
Kole_0922	DEAD/DEAH box helicase	**755**	**288**	89	102	HGTector
Kole_0969	3-oxoacyl-ACP synthase	**2386**	**3063**	939	1424	HGTector

III , FabH

Kole_0970	enoyl-ACP reductase II , fabK	2226	3243	1486	1641	HGTector[a]
Kole_0971	malonyl CoA-acyl carrier protein transacylase, fabD	2304	4211	2303	2647	HGTector
Kole_0972	acyl carrier protein	6531	12601	4850	4241	HGTector
Kole_0973	3-oxoacyl-ACP synthase II, fabF	4815	9257	4753	4498	HGTector
Kole_1015	Response regulator	**1289**	**515**	95	130	HGTector
Kole_1016	Response regulator	**783**	**280**	54	72	HGTector
Kole_1017	Histidine kinase	**697**	**275**	59	90	Phylogenetic analysis
Kole_1718	Ribosomal RNA methyltransferase, rmlH	**531**	332	211	203	HGTector
Kole_1745	PPIase FKBP-type	2783	2382	1541	**430**	HGTector

893   a) Phylogenetic tree is shown in Fig. S6.

    Fig. Legends:

Fig. 1. Growth rate of *K. olearia* as a function of temperature. Isothermic growth curves were

generated at each temperature from an inoculum grown at that temperature for at least three transfers

(except for 25°C and 80°C, for which an inoculum from the same temperature could not be

generated; see materials and methods). Up-shifted and down-shifted growth curves were generated

from an inoculum that was grown at lower and higher temperatures, respectively. Red squares,

growth temperature up-shifted from 65°C to 77°C or from 40°C to 65°C; Blue circles, growth

temperature down-shifted from 77°C to 65°C, 65°C to 40°C, or 40°C to 30°C. Data points represent

the mean of replicate cultures (see materials and methods); error bars represent standard error.

Fig. 2. Model of energy generation pathway in *K. olearia* during growth on pyruvate. The model

includes genes likely involved in conversion of pyruvate to acetate, CO_2, H_2, and ATP. The genes

were selected from the list of genes highly expressed across all temperature conditions (Fig. S2a,

Table S3). Acetate transport is not shown. The dashed box indicates hydrogenase activity. The two

highly expressed hydrogenases are shown, but their potential interactions with each other or with the

membrane are not known. Increased expression of citrate synthase at low temperature, which could

redirect acetyl-CoA away from acetate production, is shown in grey. The assignment of the ABC

transporter as pyruvate transporter (Kole_1509-1513) is based on its high expression level, but

experiments are needed to confirm its involvement in pyruvate transport. The model also explains the

observed lower ratio of carbon dioxide to hydrogen produced by growth on maltose vs. pyruvate (not

shown), because during growth on maltose reduced electron carriers would be generated from the

conversions of maltose to pyruvate.

Fig. 3. Biplot of the principal component analysis of 8 transcriptomes. The plot is based on

expression values from all genes. Each transcriptome is denoted by a point and genes are represented

by vectors. The 20 longest (i.e. most highly correlated) gene vectors are shown. Coordinates and

vector length for all genes can be found in Table S4. It should be noted that the *ffs* (Kole_R0010)

transcript is only 115 nt, and may not have been fully represented in every transcriptome due to our

isolation protocol which selects against small RNA (<200 nucleotides). Also, the high expression of

the alcohol dehydrogenase (Kole_0742) is probably due to the addition of stop solution before RNA

isolation (see Supplemental material).

Fig. 4. Difference between observed and expected number of temperature responsive genes

across functional categories. Functional categories were assigned using the Clusters of Orthologous

Groups (COG) database as implemented in IMG (Markowitz et al. 2014) and are denoted by one-

letter abbreviations along the X-axis. For each temperature treatment (30°C, 40°C and 77°C) only the

temperature-responsive fraction of the *K. olearia* genome was considered. If the temperature-

responsive genes were randomly distributed across functional categories we would expect the same

fraction of temperature-responsive genes in each COG category. The difference (in percent) between

the observed and expected number of temperature responsive genes is plotted on the Y-axis with

positive and negative values referring to over- and under-representation of the temperature-

responsive genes, respectively. COG category abbreviations: J: Translation, ribosomal structure and

biogenesis, K: Transcription, L: Replication, recombination and repair, D: Cell cycle control, cell

division, chromosome partitioning, V: Defense mechanisms, T: Signal transduction mechanisms, M:

Cell wall/membrane/envelope biogenesis, U: Intracellular trafficking, secretion, and vesicular

transport, O: Post-translational modification, protein turnover, and chaperones, C: Energy production

and conversion, G: Carbohydrate transport and metabolism, E: Amino acid transport and metabolism,

F: Nucleotide transport and metabolism, H: Coenzyme transport and metabolism, I: Lipid transport

and metabolism, P: Inorganic ion transport and metabolism, Q: Secondary metabolites biosynthesis,

transport, and catabolism, R: General function prediction only, S: Function unknown, NC: <u>N</u>ot in

<u>C</u>OG database.

Fig. 5. Schematic of *K. olearia*'s major temperature-induced transcriptional responses. Major

responses outlined in the text occurring in the three states observed are shown. The number of

chaperone, protease, and cold-shock symbols reflects their relative expression at each temperature.

Chaperones include groEL (Kole_1627), groES (Kole_1626), and dnaK (Kole_0886). Proteases refer

to protease Do (Kole_1599) and protease La (Kole_0536). Cold-shock genes consist of Kole_2064

(dark green squares) and Kole_0109 (green trapezoids). The different coloured ribosomes represent

changes in ribosomal protein composition at sub-optimal temperatures (see text for discussion). The

putative two-component regulatory system (Kole_1015 – 1017) that had a dramatic increase in

expression under cold stress is shown. Prominent changes in functional categories (COGs) for each

temperature condition are also shown. Arrows indicate relative gene expression when compared to

growth at optimal temperature. Core functions occurring at all temperatures are shown in the white

box. Basic membrane lipoprotein and the major toga anchor protein refer to Kole_1554 and

Kole_1500, respectively. See Fig. 2 for genes involved in pyruvate metabolism.

30°C ■   40°C ●   65°C ▲   77°C ◆

axis 2, 24% of variance

Kole_R0006 ssrA RNA
Kole_R0010 ffs RNA
Kole_R0049 rnpB RNA
Kole_1599 protease Do
Kole_1431 hypothetical protein
Kole_1511 extracellular solute-binding protein

axis 1, 66% of variance

Kole_0972 acyl carrier
Kole_0109 Csp
Kole_2064 Csp
Kole_1840 rpl L10
Kole_1907 30S ribosomal protein S12
Kole_0258 rpl L34

Kole_0742 alcohol dehydrogenase
Kole_1501 OmpB toga protein
Kole_1627 chaperonin groEL
Kole_1904 EF-Tu
Kole_2020 GAPDH
Kole_1554 basic membrane lipoprotein
Kole_1626 groES
Kole_0175 extracellular solute-binding protein family 1

Optimal

30°C
Cold Stress

40°C
Intermediate

65°C

77°C
Heat Stress

Ribosome

Ribosome

Ribosome

Ribosome

Chaperones

Chaperones

Chaperones

Chaperones

Proteases

Proteases

Proteases

Proteases

Cold-shock Proteins

Fewer Cold-shock Proteins

Fewer Cold-shock Proteins

Induction of Stress Sigma Factor

σ^E

Two-Component System and other Signal Transduction Genes

Helicase

RNA Turnover

Hypothetical Proteins

CRISPRs and Transposases

Acetyl-CoA

Citrate   Acetate

Fatty Acid Synthesis

Energy Production and Conversion

Trafficking and Secretion

Carbohydrate Metabolism and Transport

Amino Acid Metabolism

Amino Acid Metabolism

Core Functions (At all temperatures):

Pyruvate

↓

Acetate

PPIases

PPIases

Basic membrane lipoprotein

Toga Proteins