# Tradict enables accurate prediction of eukaryotic transcriptional states from 100 marker genes

Surojit Biswas[1,*], Konstantin Kerner[2], Paulo José Pereira Lima Teixeira[3,4], Jeffery L. Dangl[3-7,†], Vladimir Jojic[8,†], Philip A. Wigge[9,†,*]

[1] Department of Biomedical Informatics, Harvard Medical School, Boston, MA 02115, USA
[2] Botanical Institute, Biocenter, University of Cologne, D-50674 Cologne, Germany.
[3] Howard Hughes Medical Institute and [4] Department of Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA.
[5] Carolina Center for Genome Science, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
[6] Department of Microbiology and Immunology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
[7] Curriculum in Genetics and Molecular Biology, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
[8] Department of Computer Science, University of North Carolina at Chapel Hill, Chapel Hill, NC 27599, USA
[9] Sainsbury Laboratory, University of Cambridge, Cambridge CB2 1LR, UK

[†] Contributed equally
* Correspondence: surojitbiswas@g.harvard.edu, Philip.Wigge@slcu.cam.ac.uk

## Abstract

Transcript levels are a critical determinant of the proteome and hence cellular function. Because the transcriptome is an outcome of the interactions between genes and their products, it may be accurately represented by a subset of transcript abundances. We developed a method, Tradict (transcriptome predict), capable of learning and using the expression measurements of a small subset of 100 marker genes to predict transcriptome-wide gene abundances and the expression of a comprehensive, but interpretable list of transcriptional programs that represent the major biological processes and pathways of the cell. By analyzing over 23,000 publicly available RNA-Seq datasets, we show that Tradict is robust to noise and accurate. Coupled with targeted RNA sequencing, Tradict may therefore enable simultaneous transcriptome-wide screening and mechanistic investigation at large scales.

## Introduction

As the critical determinant of the proteome and therefore cellular status, the transcriptome represents a key node of regulation for all life[1]. Transcriptional control is managed by a finely tuned network of transcription factors that integrate environmental and developmental cues in order to actuate the appropriate responses in gene expression[2–4]. Importantly, the transcriptomic state space is constrained. Pareto efficiency constraints suggest that no gene expression profile or phenotype can be optimal for all tasks, and consequently, that some expression profiles or phenotypes must come at the expense of others[5,6]. Furthermore, across all major studied kingdoms of life, cellular networks demonstrate remarkably conserved scale-free properties that are topologically characterized by a small minority of highly connected regulatory nodes that link the remaining majority of sparsely connected nodes to the network[7–9]. These theories suggest that the effective dimension of the transcriptome should be far less than the total number of genes it contains. If true to a large enough extent, it may be possible to faithfully compress and prospectively summarize entire transcriptomes by measuring only a small, carefully chosen subset of it.

**Figure 1. The primary drivers of variation in our training transcriptome collection are developmental stage and tissue**. a) *A. thaliana*, b) *M. musculus*. Also shown are plots of PC3 vs. PC1 to provide another perspective.

Indeed, previous studies have exploited this reduced dimensionality to perform gene expression imputation for microarray data for missing or corrupted values[10–12]. Others have extended these intuitions to predict expression from probe sets containing a few hundred genes[13,14]. However, prediction accuracies have been modest and usually limited to 4,000 target probes/genes. Recently, several studies examined the transcriptomic information recoverable by shallow sequencing especially as it applies to single-cell experiments[15–18]. Jaitin *et al.* (2014) and Pollen *et al.* (2014) demonstrated that only tens of thousands of reads are required per cell to learn and classify cell types *ab initio*[16,18]. Heimberg *et al.* (2016) extended these findings and demonstrated that the major principal components of a typically sequenced mouse bulk or single-cell expression dataset may be estimated with little error at even 1% of the depth[15]. Though these approaches advance the notion of strategic transcriptome undersampling, they only recover broad transcriptional states and are restricted to measuring only the most abundant genes. During sample preparation -- arguably the most expensive cost of a multiplexed sequencing experiment -- shallow sequencing-based approaches still utilize protocols meant for sampling the entire transcriptome and therefore consume more resources than necessary. Furthermore, given that the expression of even the most abundant genes is highly skewed, sequencing effort is wastefully distributed compared to an approach that chooses which genes to measure more wisely. Finally, it is still not clear from sample sizes and biological contexts previously studied whether the low dimensionality of the transcriptome may be leveraged unconditionally (or nearly so) across organism and application.

In this work, we introduce Tradict (<u>tra</u>nscriptome pre<u>dict</u>), a robust-to-noise and probabilistically sound algorithm, for inferring gene abundances transcriptome-wide, and predicting the expression of a transcriptomically comprehensive, but interpretable list of *transcriptional programs* that represent the major biological processes and pathways of the cell. Tradict makes its predictions using only the expression measurement of a single, context-independent, machine-learned subset of 100 marker genes. Importantly, Tradict's predictions are formulated as posterior distributions over unmeasured genes and programs, and therefore simultaneously provide point and credible interval estimates over predicted expression. Using a representative sampling of over 23,000 publicly available, transcriptome-wide RNA-Seq datasets for *Arabidopsis thaliana* and *Mus musculus*, we show Tradict prospectively models program expression with striking accuracy. Our work demonstrates the development and large-scale application of a probabilistically reasonable multivariate count/non-negative data model, and highlights the power of directly modeling the expression of a comprehensive list of transcriptional programs in a supervised manner. Consequently, we believe that Tradict, coupled with targeted RNA sequencing[19–24], can rapidly illuminate biological mechanism and improve the time and cost of performing large forward genetic, breeding, or chemogenomic screens.

**Figure 2. Tradict's algorithmic workflow.** a) During training, the transcriptome is first quantitatively summarized in terms of a collection of a few hundred, biologically comprehensive transcriptional programs. These are then decomposed into a subset of marker genes using an adaptation of the Simultaneous Orthogonal Matching Pursuit algorithm. A Multivariate Normal Continuous-Poisson hierarchical model is used as a predictive model to capture covariance relationships between markers, transcriptional programs, and all genes. b) During prediction, Tradict predicts the expression of transcriptional programs and all genes in the transcriptome using the expression measurements of the marker genes. c) The Multivariate Normal Continuous-Poisson hierarchy enables Tradict to efficiently model statistical coupling between the non-negative expression measurements typical of sequencing experiments. This is done by assuming that associated with each observed, noisy TPM measurement, there is an unmeasured (denoised), latent abundance the logarithm of which comes from a Multivariate Normal distribution over all genes and transcriptional programs.

86 **Results**
87
**Assembly of a comprehensive training collection of transcriptomes**
We downloaded all available Illumina sequenced publicly deposited RNA-Seq samples
(transcriptomes) on NCBI's Sequence Read Archive (SRA). Among samples with at least 4 million
reads, we successfully downloaded and quantified the raw sequence data of 3,621 and 27,450
transcriptomes for *A. thaliana* and *M. musculus,* respectively. After stringent quality filtering, we
retained 2,597 (71.7%) and 20,847 (76.0%) transcriptomes comprising 225 and 732 unique SRA
submissions for *A. thaliana* and *M. musculus*, respectively. An SRA 'submission' consists of multiple,
experimentally linked samples submitted concurrently by an individual or lab. We defined 21,277 (*A.*
*thaliana*) and 21,176 (*M. musculus)* measurable genes with reproducibly detectable expression given
our tolerated minimum sequencing depth and mapping rates (see Supplemental Information "Materials
and Methods" for further information regarding data acquisition, transcript quantification, quality filtering,
and expression filtering). We hereafter refer to the collection of quality and expression filtered
transcriptomes as our *training transcriptome collection*.
In order to assess the quality and comprehensiveness of our training collection, we performed a
deep characterization of the expression spaced spanned by these transcriptomes. We found that the
transcriptome of both organisms was highly compressible and that the primary drivers of variation were

tissue and developmental stage (Figure 1a-b, Figure S1), with many biologically realistic trends within
each cluster (Supplemental Analysis 1). We additionally examined the distribution of submissions
across the expression space, compared inter-submission variability within and between tissues,
inspected expression correlations among genes with well-established regulatory relationships, and
assessed the evolution of the expression space across time. These investigations revealed our training
collection is of high and reproducible technical quality, reflective of known biology, and stable
(Supplemental Analysis 1, Figures S2-S4). Given additionally the diversity of tissues, genetic
perturbations, and environmental stimuli represented in the SRA, these results, taken together, suggest
that our training collection is an accurate and representative sampling of the transcriptomic state space
that is of experimental interest for both organisms.
114
115     **Tradict - algorithm overview**
116             Given a training transcriptome collection, Tradict encodes the transcriptome into a single subset
of globally representative *marker* genes and learns their predictive relationship to the expression of a
comprehensive collection of transcriptional programs (e.g. pathways, biological processes) and to the
rest of the genes in the transcriptome. Tradict's key innovation lies in using a Multivariate Normal
Continuous-Poisson (MVN-CP) hierarchical model to model marker latent abundances -- rather than
their measured, noisy abundances -- jointly with the expression of transcriptional programs and the
abundances of the remaining non-marker genes in the transcriptome. In so doing, Tradict is able to 1)
efficiently capture covariance structure within the non-negative, right-skewed output typical of
sequencing experiments, and 2) perform robust inference of gene set and non-marker expression even
in the presence of significant noise.
126             Figure 2 illustrates Tradict's general workflow. Estimates of expression are noisy, especially for
low to moderately expressed genes. Given samples are often explored unevenly and that the *a priori*
abundance of each gene differs, the level of noise in a gene's measured expression for a given sample
varies, but it can be estimated. Therefore, during training, Tradict first learns the log-latent, denoised
abundances for each gene in every sample in the training collection using the lag transformation[25]. It
then collapses this latent transcriptome into a collection of predefined, comprehensive collection of
*transcriptional programs* that represent the major processes and pathways of the cell related growth,
development, and response to the environment (Supplemental Tables 3-4). In this work, we focus on
creating a Gene Ontology derived panel of transcriptional programs, in which the first principal
component of all genes contained within an appropriately sized and representative GO term is used to
define an accordingly named transcriptional program[26,27]. The expression values of these programs are
then encoded using an adapted version of the Simultaneous Orthogonal Matching Pursuit algorithm
into a small subset of marker genes selected from the transcriptome[28,29]. Tradict finally stores the mean
and covariance relationships between the log-latent expression of the selected markers, the
transcriptional programs, and the log-latent expression of the remaining non-marker genes at the
Multivariate Normal layer of the MVN-CP hierarchical model (Figure 2a).
142             Prospectively, only the expression of these marker genes needs to be measured and the
expression of genes and/or transcriptional programs can be inferred as needed. During prediction,
Tradict uses the observed marker measurements as well as their log-latent mean and covariance
learned during training, to estimate -- via MCMC sampling -- the posterior distribution over the log-latent
abundances of the markers[30]. Though a simply a consequence of proper inference of our model, this
denoising step adds considerable robustness to Tradict's predictions. From this estimate, Tradict uses
covariance relationships learned during training to estimate the conditional posterior distributions over
the remaining non-marker genes and transcriptional programs (Figure 2b).  From these distributions,
the user can derive point estimates (e.g. posterior mean or mode), as well as measures of confidence
(e.g. credible intervals). A complete description of the entire algorithm can be found in the "Tradict
algorithm" section of the Materials and Methods in the Supplemental Information.
153

**Figure 3. Tradict prospectively predicts gene and transcriptional program expression with superior and robust accuracy.** Tradict's prospective prediction accuracy during 20-fold cross validation of the entire training collection for both organisms. a) Heatmaps illustrating test-set reconstruction performance of all transcriptional programs for *A. thaliana*. Shown is the reconstruction performance for all samples in our transcriptome collection when they were in the test-set. b) Density plots of predicted vs. actual test-set expression for all genes (left) and transcriptional programs (right) for *A. thaliana*, after controlling for inter-sumbission biological signal. The intra-submission expression of each gene and transcriptional program was z-score transformed to make their expression comparable. c & d) Same as a & b, but for *M. musculus*. e) Comparison of Tradict's performance vs. several baselines: SR (structured regression), LWA (locally weighted averaging), and Tradict Shallow-Seq. f) A posterior predictive check illustrating the concordance between Tradict's posterior predictive distribution and the distribution of test-set expression values for genes (left) and transcriptional programs (right). Plotted is the percent of test-set observations contained within a credible interval vs. the size of the credible interval. A unique credible interval is derived for each gene/program. The "x=y" line is illustrated as a dotted black line. Shaded error bands depict the sampling distribution of this analysis across test-sets from a 20-fold cross validation on the *A. thaliana* dataset.

## Tradict prospectively predicts gene and transcriptional program expression with superior and robust accuracy

To understand Tradict's prospective predictive performance, we performed 20-fold cross validation on the training transcriptome collections for both *A. thaliana* and *M. musculus* and evaluated Pearson correlation coefficients (PCC) between predicted and actual expression for each fold when the remaining 95% of folds were used for training. To make this experiment as reflective of reality as possible, folds were divided by submission so that samples from the same set of experiments would not

appear both in training and test sets. Because submissions to the SRA span a broad array of biological
contexts, the total biological signal contained in any given test set likely exceeds that of what would be
expected for typical application, which in turn would lead to overly optimistic estimates of prediction
accuracy. We therefore evaluated *intra-submission accuracy*, in which PCC calculations are performed
on 'submission-adjusted' expression values. To do this, for each gene and program, each test-set
submission's mean expression was subtracted from the expression values for all samples associated
with that submission. In effect, this regresses out between-submission effects, and allows us to assess
Tradict's predictive performance one experiment at a time, as one would do in practice.
Figures 3a and 3c illustrate that the reconstruction performance for transcriptional programs in
both organisms is strikingly accurate across all collected submissions. Quantitatively speaking, the
average intra-submission PCCs for transcriptional programs are 0.94 and 0.93 for *A. thaliana* and *M.*
*musculus*, respectively. This is despite lower prediction performance on gene expression (Figures 3b
and 3d). Intuitively, this is because transcriptional programs are measured as linear combinations of the
log-latent TPMs of the genes that comprise them, effectively smoothing over the orthogonal noise
present in each gene's expression prediction.
We also found Tradict's performance to be superior to several baselines. These include two
successful approaches developed in Donner *et al.* (2012) for microarray[14], and a version of Tradict that
uses the 100 most abundant genes as its selected markers (Figure 3e, Figure S5, Supplemental
Analysis 2). The former baselines rely on structured regression (SR) and locally weighted averaging
(LWA), linear/parametric and non-parametric methods, respectively. The latter baseline examines the
utility of simple shallow sequencing by using the most abundant genes as markers for Tradict (see
Supplemental Analysis 2 for a more detailed description). Figure 3e illustrates test-set intra-submission
performance of each method as a function of the number of markers entered into the model. LWA
demonstrates the quickest performance gain, but then saturates after 10 markers. This is likely
because a non-linear kernel based approach makes the most efficient use of a few markers, but is
adversely impacted by the curse of dimensionality as more markers are added. The parametric
methods (Tradict, SR) navigate this dimensionality increase more efficiently and ultimately realize
better performance for a still reasonable number of markers. Tradict outperforms SR and Tradict
Shallow-Seq, ultimately obtaining a PCC between predicted and actual expression of 0.71 for genes
and 0.96 for transcriptional programs. This suggests Tradict's probabilistic framework is more
reasonable than SR's and that Tradict's marker selection is more optimal than picking the most
abundant genes. We additionally found Tradict's predictions were robust to noise in the form of low
sequencing depth and/or corrupt marker measurements (Figure S5 and S10, Supplemental Analysis 2
and 5), which we attribute to its probabilistic framework, in which training and prediction are performed
in the space of denoised latent abundances.
To further assess the validity of Tradict's modeling assumptions, we examined how Tradict's
posterior predictive distribution matched the distribution of test-set gene and program expression
values. Specifically, we performed a posterior predictive check in which we asked what percent of test-
set gene or program expression values fall within an X% credible interval, where a unique interval is
defined for each gene/program[30]. If Tradict's posterior predictive distribution is reasonable then X% of
the true expression values should fall within this interval for any X. Figure 3f illustrates the results of this
analysis as performed on disjoint test-sets from a 20-fold cross validation on the *A. thaliana* dataset. On
average, the X% credible interval captures X% of test-set observations for any choice of X. The
posterior predictive distribution for transcriptional programs may be slightly too wide at moderate
interval sizes (30-70%), which would make Tradict more conservative (higher type II error rate) than it
should be. However, in practice it is accurate (p-value = 0.24, *t*-test) for larger, more standard interval
sizes (e.g. 95%). We conclude that Tradict's probabilistic modeling assumptions capture unseen data
well.
We next characterized Tradict's limitations through error, power, program annotation robustness
analyses, and a timing and memory analysis (Supplemental Analysis 3-4). These analyses revealed
that training-set expression variance and mean abundance correlated positively with both program and

gene expression prediction performance (Supplemental Analysis 3.I, Supplemental Tables 3-4).
Combined with program size as another predictor, these variables could account for most of the error
(60% of total variance) in program expression prediction. A power analysis revealed that for both *A.*
*thaliana* and *M. musculus*, 1,000 samples -- comprising approximately 100 submissions -- is sufficient
for optimum performance (Supplemental Analysis 3.II). An examination of how gene-to-program mis-
annotation rates influence predictive performance revealed that program expression prediction
perfromance was robust up to a 20% mis-annotation rate and that gene expression prediction
performance was completely robust to any level of mis-annotation. The latter result is a consequence of
a statistical decoupling between gene and program expression prediction (Supplemental Analysis 3.III).
Finally, Tradict's training time and peak memory requirements scaled linearly with training set size, and
increased 0.26 seconds and 1.1 Mb per sample. Prediction time was limited by MCMC sampling from
the conditional posterior distributions of gene and program expression, and required 3.1 seconds per
sample on average (Supplemental Analysis 4). Taken together, we conclude that the causes of
Tradict's errors are well understood and intuitive, and that Tradict's sample requirements are
reasonable, especially for major model organisms (Supplemental Table 5, Supplemental Analysis 3.II).
Furthermore, Tradict is robust to noisily defined transcriptional programs, and its computational
requirements scale well to large datasets.
231
**Case studies reveal the power of predicting and studying transcriptional program expression**
To demonstrate how Tradict may be applied in practice, we focused on two case studies related
to innate immune signaling -- one performed using bulk *A. thaliana* seedlings (detailed below), and the
other using primary immune *M. musculus* cell lines (detailed in Supplemental Analysis 5; Figure S10).
We trained Tradict on our full collection of training transcriptomes for each organism to produce two
organism-specific Tradict models. Each was based on the selection of 100 markers learned from the
full training transcriptome collection (Supplemental Tables 7-8) that we assert are globally
representative, and context-independent. The case study samples do not, of course, appear in the
collection of training transcriptomes.
241
**Tradict accurately predicts temporal expression patterns for a diverse panel of *A. thaliana***
**immune signaling mutants under different hormone perturbations -** The hormones salicylic acid
(SA) and jasmonic acid (JA) play a major, predominantly antagonistic regulatory role in the activation of
plant defense responses to pathogens. Yang *et al.* (2016) investigated the effect of a transgenically
expressed bacterial effector, HopBB1, on immune signaling in *A. thaliana*[31]. In their study, they
performed a time course experiment, treating plants with MeJA (a JA response inducer), BTH (an SA
mimic and SA response inducer), or mock buffer and monitored the transcriptome of bulk seedlings at 0
hr, 1 hr, 5 hr, and 8 hr post treatment. These experiments included several immune signaling mutants
with differing degrees of response efficiency to MeJA and BTH treatment. Among other findings, they
conclude that HopBB1 enhances the JA response, thereby repressing the SA response and facilitating
biotrophic pathogen infection.
We asked to what extent strategic undersampling of the transcriptome and application of Tradict
could quantitatively recapitulate the findings of Yang *et al.* (2016). Given Tradict's near perfect accuracy
on predicting the expression of transcriptional programs, we took a top down, but hypothesis driven
approach to our analysis which first examined the expression of all transcriptional programs. Figure 4a
illustrates the actual and predicted expression of all transcriptional programs in *A. thaliana* as a function
of time and treatment. Here, Tradict reconstructs the expression of all transcriptional programs with an
average PCC of 0.91.

**Figure 4. Tradict accurately predicts transcriptional responses across time in response to hormone perturbation in an *A. thaliana* innate immune signaling dataset.** After being trained on the full *A. thaliana* training transcriptome collection, the selected set of 100 globally representative and context-independent markers were used to predict the expression of transcriptional programs and all genes for the transcriptomes presented in Yang *et al.* (2016). a) Actual vs. predicted heatmaps for the expression of all 150 transcriptional programs in *A. thaliana* across genotype, time, and hormone treatment. b) Predicted vs. actual expression of i) the JA response transcriptional program, and ii) the genes involved in the JA response program. c i-ii) Same as b, but for the SA response transcriptional program. d) Hypothesis free, differential transcriptional program expression analysis as performed on the actual expression of transcriptional programs vs those predicted by Tradict. Blue circles represent the actual and orange represent the predicted. All heatmaps are clustered in the same order across time, treatment, genotype, and between predicted and actual.

Recall that the genes participating in each of our transcriptional programs are pre-defined, in
this work, by a carefully chosen, interpretable, but maximally representative set of GO biological
processes. Therefore, given the goals of this study, we next examined the expression of the "response
to jasmonic acid" and "response to salicylic acid" transcriptional programs. Figure 4b shows the
expression behavior for the "response to jasmonic acid" transcriptional program across all the
genotypes and time points upon MeJA treatment. More specifically, part (i) shows that the predicted
expression and actual expression are qualitatively and quantitatively in agreement, both in magnitude
and rank across the different genotypes. For example, as expected, *coi1-16*, which cannot sense JA,
does not respond to the MeJA stimulus, while wildtype Col-0 does. However, even more subtle
expression dynamics are captured by Tradict's predictions. For example, *eds16-1* and *npr1-1* -- slightly

and strongly impaired SA responders, respectively -- are slightly and strongly hyper-responsive to
MeJA, respectively -- just as expected from the JA-SA antagonism. Furthermore, as demonstrated in
Yang *et al.* (2016), the *35S::HopBB1* transgenic line exhibits a prolonged and sustained JA response
for both the actual and predicted expression for this transcriptional program. Part (ii) of Figure 4b
illustrates the expression of all the MeJA responsive genes in this transcriptional program. Again
Tradict's predictions are in good agreement with actuality, achieving a PCC of 0.72, and it's visually
clear that the expression magnitude of these genes positively correlates with the registered expression
magnitude of the "response to jasmonic acid" transcriptional program. Figure 4c parts (i) and (ii) are
presented in the same light as Figure 4b, but are instead illustrated for the SA response transcriptional
program and constituent genes under BTH treatment. Again predictions match actuality, and the
observed trends make biological sense[32].
In order to illustrate Tradict's use in hypothesis-free investigation, we performed a differential
transcriptional program expression analysis for transcriptional programs affected by MeJA or BTH
treatment (Figure 4d, see Methods). Differentially expressed transcriptional programs based on
Tradict's predictions versus actual measurements were highly concordant and biologically reasonable.
Transcriptional programs differentially expressed with respect to MeJA treatment included "response to
bacterium," "defense response to fungus", "response to wounding," and "response to jasmonic acid" as
expected.  Transcriptional programs differentially expressed with respect to BTH treatment included
various abiotic stress responses, "defense response to fungus", "response to jasmonic acid" (via
antagonism), and "response to salicylic acid," again, as expected.
290
**Discussion**
Tradict is an accurate, robust-to-noise method for predicting the expression of a comprehensive,
but interpretable list of transcriptional programs that represent the major biological processes and
pathways of the cell. Given the comprehensiveness, stability, and exponentially growing size of the
training datasets we have assembled from publicly available sources, and as evidenced by our cross
validation experiments, which were performed over thousands of samples deposited over the span of
six years, the 100 markers Tradict learns are likely to be predictive independent of most contexts and
applications. As illustrated through our case studies, examining the expression of these predicted
transcriptional programs makes intuitive sense and provides a neat summary of underlying gene
expression patterns.
Tradict additionally provides expression predictions for all genes in the transcriptome. Though
its current gene expression prediction accuracy is less than ideal for more sensitive applications,
Tradict's performance is superior to previous efforts and is improving logarithmically in the number of
samples. We attribute Tradict's performance gains over previous methods first to improved
measurement technology. Previous methods were developed for microarray, a substantially more noisy
technology than RNA-Sequencing[10–14]. Consequently, training efficiency was lower as well as
measurement accuracy of true expression, thus leading to modest prediction accuracy. By contrast,
Tradict is meant to interface with sequencing based readouts of gene expression, a data type that is
popular and proliferating exponentially as the time and price of sequencing continues to fall. Second,
we believe Tradict's probabilistic framework goes a step beyond previous efforts by modeling marker-
gene and marker-program relationships not at the level of measured abundances, which are noisy, but
at the level of latent abundances. Working in this denoised space naturally improves accuracy and
affords robustness.
Taken together, we believe that Tradict coupled with target RNA sequencing can enable
transcriptome-wide screening cheaply and at scale. Well-established commercial[19,20] and non-
commercial[21,33] methods exist for targeted RNA sequencing in a multiplexed manner, and they are able
to measure the expression of 10's-100's of genes with accuracy, making their use immediately
compatible with Tradict. One method in particular, RASL-Seq[22–24], does so cheaply with high precision
and multiplexibility by directly probing total RNA and making efficient use of dual-indexing. We estimate
that Tradict coupled with a time and resource efficient targeted RNA-sequencing protocol such as

RASL-Seq could bring the cost of obtaining actionable transcriptome-wide information simultaneously
for thousands to tens of thousands of samples to close to $1 per sample.
This scale could greatly benefit high-throughput breeding and screening applications. Forward
genetic screens in most eukyarotic organisms require assaying $10^3$-$10^4$ mutants. Small molecule, or
more generally chemogenomic, drug screens often require screening thousands of molecules against
multiple cell lines in multiple conditions. Agricultural screens -- whether for breeding or field
phenotyping -- also require measuring thousands of individuals. Though in these cases a screen is
made cheap and scalable by monitoring an easily selectable phenotype, new phenotyping architectures
must be developed and optimized for each new screen (e.g. reporter lines, imaging hardware/software).
Given the ubiquity of RNA, a transcriptome-wide screening approach would not suffer from such a
drawback. Furthermore, and more importantly, though quickly interpretable, the phenotype being
screened for is usually a uni-dimensional datum that offers little immediate insight into mechanism. In
contrast, using Tradict to help perform transcriptome-wide screening could couple the process of
hypothesis generation and mechanistic investigation. Here, we argue that the scalable monitoring of the
expression of a comprehensive list of just a few hundred transcriptional programs affords an attractive
balance of nuance and interpretability. Consequently, this efficient investigation, largely facilitated by
Tradict, could greatly accelerate the pace of genetic dissection, breeding, and drug discovery.
338

**Acknowledgements**
We thank Brian Cleary, Aviv Regev, Amaro Taylor-Weiner, Craig Bohrson, and Derek Lundberg for
valuable discussions in preparing this manuscript. SB was supported by a Churchill Scholarship from
the Winston Churchill Foundation of the United States and by a training grant from the NHGRI/NIH (T32
HG002295). PJPLT was supported by a fellowship from the Pew Latin American Fellows Program in
the Biomedical Sciences. This work was additionally funded, in part, by a fellowship to PW from the
Gatsby Foundation (GAT2373/GLB) and by grants to JLD from the National Institutes of Health (1RO1
GM107444), the Gordon and Betty Moore Foundation (GBMF3030), and the HHMI. JLD is an
Investigator of the Howard Hughes Medical Institute.
348

**Code availability**
A MATLAB implementation of Tradict is available at https://github.com/surgebiswas/tradict. All code to
perform data downloads, analysis, and generate figures are available at
https://github.com/surgebiswas/transcriptome_compression. Note that in order to make Tradict open-
source and reach a wider audience we are developing a user-friendly, unit-tested Python version of
Tradict that will be made available before publication.

**References**
1. Jacob, F. & Monod, J. Genetic regulatory mechanisms in the synthesis of proteins. *J. Mol. Biol.* **3,** 318–356 (1961).
2. Kaufmann, K., Pajoro, A. & Angenent, G. C. Regulation of transcription in plants: mechanisms controlling developmental switches. *Nat. Rev. Genet.* **11,** 830–842 (2010).
3. Mitchell, P. J. & Tjian, R. Transcriptional Regulation in Mammalian Cells by DNA Binding Proteins. *Science (80-. ).* **245,** 371–378 (1989).
4. Segal, E. *et al.* Module networks: identifying regulatory modules and their condition-specific regulators from gene expression data. *Nat. Genet.* **34,** (2003).
5. Hart, Y. *et al.* Inferring biological tasks using Pareto analysis of high-dimensional data. *Nat. Methods* **12,** (2015).
6. Shoval, O. *et al.* Evolutionary Trade-Offs, Pareto Optimality, and the Geometry of Phenotype Space. *Science (80-. ).* **336,** 1157–1160 (2012).
7. Jeong, H., Tombor, B., Albert, R., Oltvai, Z. N. & Barabási, a L. The large-scale organization of metabolic networks. *Nature* **407,** 651–654 (2000).

8.  Albert, R., Lee, J. H. & Barabási, A.-L. Error and attack tolerance of complex networks. *Nature* **406,** 378–382 (2000).
9.  Barabási, A.-L. & Albert, R. Emergence of Scaling in Random Networks. *Science (80-. ).* **286,** 509–513 (1999).
10. Troyanskaya, O. *et al.* Missing value estimation methods for DNA microarrays. **17,** 520–525 (2001).
11. Liew, A. W., Law, N. & Yan, H. Missing value imputation for gene expression data : computational techniques to recover missing data from available information. **12,** 498–513 (2010).
12. Celton, M., Malpertuy, A., Lelandais, G. & Brevern, A. G. De. Comparative analysis of missing value imputation methods to improve clustering and interpretation of microarray experiments. (2010).
13. Ling, M. H. T. & Poh, C. L. A predictor for predicting Escherichia coli transcriptome and the effects of gene perturbations. *BMC Bioinformatics* **15,** 140 (2014).
14. Donner, Y., Feng, T., Benoist, C. & Koller, D. Imputing gene expression from selectively reduced probe sets. *Nat. Methods* **9,** (2012).
15. Heimberg, G., Bhatnagar, R., El-samad, H. & Thomson, M. Low Dimensionality in Gene Expression Data Enables the Accurate Extraction of Transcriptional Programs from Shallow Sequencing. *Cell Syst.* **2,** 239–250 (2016).
16. Pollen, A. A. *et al.* Low-coverage single-cell mRNA sequencing reveals cellular heterogeneity and activated signaling pathways in developing cerebral cortex. *Nat. Biotechnol.* **32,** (2014).
17. Kliebenstein, D. J. Exploring the shallow end; estimating information content in transcriptomics studies. *Front. Plant Sci.* **3,** 1–10 (2012).
18. Jaitin, D. A. *et al.* Massively Parallel Single-Cell RNA-Seq for Marker-Free Decomposition of Tissues into Cell Types. *Science (80-. ).* **343,** 776–779 (2014).
19. ThermoFisher Scientific. Targeted RNA Sequencing by Ion Torrent Next-Generation Sequencing. at <http://www.thermofisher.com/us/en/home/life-science/sequencing/rna-sequencing/targeted-rna-sequencing-ion-torrent-next-generation-sequencing.html>
20. Illumina. TruSeq Targeted RNA Expression Kits. at <http://www.illumina.com/products/truseq-targeted-rna-expression-kits.html>
21. Mercer, T. R. *et al.* Targeted sequencing for gene discovery and quantification using RNA CaptureSeq. *Nat. Protoc.* **9,** 989–1009 (2014).
22. Li, H., Qiu, J. & Fu, X. RASL-seq for Massively Parallel and Quantitative Analysis of Gene Expression. *Curr. Protoc. Mol. Biol.* 1–9 (2012). doi:10.1002/0471142727.mb0413s98
23. Larman, H. B. *et al.* Sensitive, multiplex and direct quantification of RNA sequences using a modified RASL assay. *Nucleic Acids Res.* 1–12 (2014). doi:10.1093/nar/gku636
24. Scott, E. R. *et al.* RASLseqTools: open-source methods for designing and analyzing RNA-mediated oligonucleotide Annealing, Selection, and, Ligation sequencing (RASL-seq) experiments. *bioRxiv* (2016).
25. Biswas, S. The latent logarithm. *arXiv* 1–11 (2016).
26. Ma, S. & Kosorok, M. R. Identification of differential gene pathways with principal component analysis. *Bioinformatics* **25,** 882–889 (2009).
27. Fan, J. *et al.* Characterizing transcriptional heterogeneity through pathway and gene set overdispersion analysis. *Nat. Methods* **13,** 241–244 (2016).
28. Tropp, J. a & Gilbert, A. C. Signal Recovery From Random Measurements Via Orthogonal Matching Pursuit. *IEEE Trans. Inf. Theory* **53,** 4655–4666 (2007).
29. Tropp, J. a., Gilbert, A. C. & Strauss, M. J. Algorithms for simultaneous sparse approximation. Part I: Greedy pursuit. *Signal Processing* **86,** 572–588 (2006).
30. Gelman, A. *et al. Bayesian Data Analysis.* (Chapman & Hall, 2013).
31. Yang, L. *et al.* The Pseudomonas syringae type III effector HopBB1 fine tunes pathogen virulence by gluing together host transcriptional regulators for degradation. *Submitted* (2016).

32. Jones, J. D. G. & Dangl, J. L. The plant immune system. *Nature* **444,** 323–9 (2006).
33. Fu, G. K. *et al.* Molecular indexing enables quantitative targeted RNA sequencing and reveals poor efficiencies in standard library preparations. **111,** 1891–1896 (2014).