1

Cell fusion proteins can rejuvenate adult dendritic trees

Veronika Kravtsov[1,#], Meital Oren-Suissa[1,2,#], and Benjamin Podbilewicz[1†]

[1]Department of Biology, Technion- Israel Institute of Technology, Haifa 32000, Israel

[#]These authors contributed equally to this work

[†]Correspondence to: podbilew@technion.ac.il

Additional footnotes:

[2]Present address: Columbia University Medical Center, Department of Biochemistry &

Molecular Biophysics, New York, 10032, USA

Abstract

The convoluted architecture of dendritic arbors poses a challenge to understanding age-

dependent alterations and regeneration following injury. Here, we show that induction of

cellular fusogens can remodel and facilitate regeneration of dendrites in polymodal PVD

neurons of aging *Caenorhabditis elegans*. Using whole-animal live imaging, we find that the

PVD dendritic trees, composed of repetitive "menorah" units, show age-dependent

hyperbranching, disorganization, and loss of self-avoidance. These processes, while

independent of canonical lifespan-regulating pathways, can be partially rescued by ectopic

expression of the fusogen EFF-1. Furthermore, the decreased capacity of old animals to

repair laser-induced severed dendrites via auto-fusion can be restored by reducing DAF-2

(Insulin/IGF-1Receptor) function or by ectopic expression of the EFF-1 paralog AFF-1. Our

findings demonstrate that fusogens are sufficient to maintain the dendritic arbor structure and

increase its regeneration potential in aging animals. These antiaging strategies can be

potentially applied to other organisms to protect them from neurodegeneration.

Introduction

Aging is the primary risk for neuronal diseases and general cognitive decline in humans [1],

yet, our understanding of the process of neuronal aging at the molecular and cell biological

levels is still limited. In particular, very few studies have investigated the fate of complex

dendritic arbors during aging, and their regenerative capacity following injury. *C. elegans* is

a powerful system to study the genetics of neuronal aging and regeneration [2-4]. Signaling via

the DAF-2 Insulin/IGF-1 receptor is the most prominent and conserved pathway that controls

aging and longevity of *C. elegans*, flies, and mammals [2]. Normally, when the DAF-2/IGF-1

receptor is activated it induces a conserved PI3K/AKT kinase cascade, which in turn inhibits

the DAF-16/FOXO transcription factor from entering into the nucleus. Reduction of *daf-2*

function doubles life span, and long-lived worms are considered to stay healthy for longer [2,5].

In these animals, DAF-16/FOXO affects transcription of different genes that encode heat-

shock proteins, antimicrobial proteins, antioxidants, and other molecules, which leads

ultimately to extended lifespan [6,7]. Recently, aging-associated axonal morphological

alterations and decline in regenerative capacity were described in *C. elegans*. *daf-2*

mutations delayed these age-related morphological changes and improved regeneration of

aged severed axons in a *daf-16*-dependent manner [8-11]. Moreover, in the case of GABA

motor neurons, as adult animals age there is a reduction in axon growth, retraction and

regrowth in response to injury. Surprisingly, the decline in regeneration is controlled by *daf-*

16 in a cell-autonomously fashion and independently of lifespan [11].

Different invertebrates, including nematodes and crustaceans, use membrane fusion as

an alternative mechanism for repair of injured axons [12-15]. Cell fusion events were also

observed in the brains of mammals both spontaneously and as a result of injury such as stroke

[16-18], but the role of these events has remained unclear. In *C. elegans*, axonal regeneration via

auto-fusion is mediated by the fusogen EFF-1 [12,13,15]. EFF-1 is the first *bona fide* eukaryotic

developmental cell-cell fusion protein. It is expressed in different cell types including

neurons, and mediates fusion between cells by a homotypic mechanism [19-21].

We have been studying the role of fusion proteins in the PVD neuron, a polymodal

nociceptor that senses harsh touch to the body, cold temperature, and proprioception [22-25].

The PVD neuron exhibits an elaborate and invariant network of dendrites, which is composed

of a repetitive unit that is termed "menorah" **(Figure 1A)** [23]. Menorahs arise and are

maintained through several intrinsic and extrinsic genetic pathways [23-33].

The PVD develops in a stereotypic fashion from the L2 larval stage to the young adult

[23]. The fusion protein EFF-1 mediates dendrite retraction and auto-fusion of excess branches

by a novel cell-autonomous pruning mechanism [23]. Thus, this fusogen maintains the

menorah structure by trimming excess branching during normal developmental arborization

[23]. AFF-1, a paralog of EFF-1, is a second *C. elegans* fusogen displaying a more restricted

tissue distribution pattern [34]; AFF-1 was not found to be involved in PVD remodeling during

development [35]. AFF-1 and EFF-1 fusion proteins also auto-fuse epithelial and myoepithelial

cells to form tubes and reshape glial cells [36-38]. Moreover, it has recently been demonstrated

that in vertebrates auto-fusion takes place in the development of the vascular endothelium,

where it leads to pruning of excess blood vessels [39] in a process that remarkably resembles

EFF-1-mediated PVD pruning [23]. It is unknown whether PVD structure and function is

affected by aging and by the activity of fusion proteins.

Here, we use the PVD polymodal neuron as a model to study how aging affects the

morphology, function, and regeneration of dendrites following injury. Our study

demonstrates the age-related progressive morphological alterations of intricate dendritic

arbors in a living organism. We found that the fusogen EFF-1, when expressed in the PVD

neuron, simplifies and therefore rejuvenates aged dendritic trees. In contrast, insulin/IGF-1

receptor mutations (*daf-2*) fail to inhibit the progressive aging of dendrites and do not prevent

the decline in response to harsh touch during aging. We also discovered that PVD aging is

characterized by a decline in regenerative potential of dendrites following experimental laser

dendrotomy. Furthermore, the regeneration of transected dendritic trees can be restored in old

animals by DAF-2 insulin/IGF-1 receptor mutations, and can be differentially reestablished

by ectopic AFF-1 fusogen expression. Thus, fusogens ectopically expressed in the PVD and

mutations in DAF-2/IGF-1R, differentially rejuvenate some aspects of dendritic architecture

and regeneration potential in aging *C. elegans*.

Results

Progressive dendritic remodeling during aging is independent from the Insulin/IGF-1 pathway

To determine how aging affects the complex arborized dendritic structure, we analyzed the *C. elegans* PVD dendritic branching patterns from the fourth larval stage (L4) to ten-day-old adults. We found that during aging PVD's dendrites and menorah structure undergo disorganization and hyperbranching, which is particularly evident in regions closer to the cell body **(Figure 1A and 1C)**. Remarkably, we found that these age-dependent morphological changes of the PVD dendritic pattern were not affected in long-lived animals carrying a mutation in *daf-2* **(Figure 1A-1H, and Figure 1-figure supplement 1)**. Adjacent menorahs normally avoid each other [40], and we found that as the animal ages menorahs lose their self-avoidance properties **(Figure 1C, 1D, and 1I)**. Consistently, this age-dependent dendritic pattern did not improve in *daf-2* mutant animals; young *daf-2* animals at the L4 stage exhibited significantly more *daf-16*-dependent self-avoidance deficiencies in comparison to wild-type **(Figure 2)**. Thus, taken together our results reveal that aging causes a *daf-2*-independent increase in disorganized branching and loss of menorah self-avoidance.

To further understand the nature of the age-dependent dendritic tree remodeling, we followed individual animals over time and analyzed time-lapse movies. We found that 5-day adults still exhibited some plasticity of the dendritic tree, with dynamic growth and retraction events **(Figure 3A** and **Movie S1)**; however, both growth and retraction were approximately two times slower **(Figure 3B)** compared to younger L4 animals. To shed more light on the link between structure and function of dendrites during aging, we performed a functionality test. PVD responds to harsh stimulations, and we used a classic harsh touch assay in *mec-4* mutant animals to specifically test PVD activity without the background light touch response

mediated by the six light-touch mechanosensory neurons [23,41]. First, we found that although

mec-4 animals have shorter life spans [10], their dendrites looked similar to wild-type between

the ages L4-5d **(Figure 4)**, which further demonstrates that the morphological alterations we

see are lifespan-independent. Second, we found that the PVD functionality decreased with

aging, with 5d adults presenting a reduced harsh touch response in comparison to 1d adults,

in both *mec-4* as well as *mec-4;daf-2* double mutants **(Figure 4F)**. The specific components

in the PVD circuit affected in 5d adults have not been identified and it is possible that

multiple components of the sensorimotor circuit contribute to the age-dependent decline in

PVD activities. Thus, our results reveal that the morphological and behavioral hallmarks of

aging in PVD dendritic arbors are independent from the canonical IGF-1 pathway that affects

lifespan.

Age-dependent dendrites remodeling can be modulated by EFF-1

The fusogen EFF-1 is essential to fuse some injured axons in *C. elegans* [12,15], and is involved

in PVD's dendrite pruning in a cell-autonomous and dosage-dependent manner during

development and in young adults [23]. When EFF-1 is overexpressed in the PVD, a strong

gradient of arborization is seen in L4s and young adults, with almost complete lack of

branches in areas that are distal from the cell body. However, in areas around the cell body

the PVD menorahs appear similar to those in the wild-type [23] **(Figure 5A** and **5B)**. Since

EFF-1 retracts and simplifies dendritic arbors in young animals [23] and is expressed in the

PVD throughout adulthood **(Figure 5-figure supplement 1)**, we hypothesized that EFF-1

overexpression will be able to retract dendrites in aged adults. Indeed we found that EFF-1

overexpression in the PVDs simplified the hyperbranching around the cell body at 9-10 days

adults **(Figure 5C-5E)**; in particular, the quaternary branch order was decreased **(Figure 5F)**.

A trend of reduction in fifth and higher order branches also appeared in aged animals

overexpressing EFF-1 **(Figure 5G)**. Thus, overexpression of the fusogen EFF-1 in the PVD

neuron is sufficient to simplify aged menorahs.

Age-dependent decline in dendrite regeneration is dependent on Insulin/IGF-1

Mammalian axons regenerate better in younger individuals than in adults [42]. Similarly, axonal

regenerative ability declines drastically as nematodes advance through development and age

[11,43]. Our knowledge is still poor on the regenerative capacity of the dendrite, and how aging

affects this process of neuronal repair. To study dendritic regeneration we severed the

primary dendrites of PVD neuron in aging adults. Typically, the PVDs show robust

regeneration at L4 larval stage, consisting in dendrite sprouting from the proximal fragment

still attached to the cell body and reconnection via fusion with the separated distal dendrite

fragment **(Figure 6A, Movie S2)**. To directly measure reconnection by auto-fusion, we used

the photoconvertible reporter Kaede **(Figure 6-figure supplement 1 and Movie S3)**. If

fusion fails to occur, the detached distal part eventually degenerates. We found that 2-3 day-

old wild-type adults respond more slowly to laser dendrotomy in comparison to L4 and

young adults (~70% of the young animals presented regeneration, whereas at the age of 2-3d

neither regeneration nor degeneration occurred within 3-6 hours **(Figure 6E and Movie S4)**.

At the age of 5 days, the ability to regenerate by dendrite auto-fusion was almost completely

lost **(Figure 6C** and **6F)**. Remarkably, long-lived *daf-2* mutants showed similar regeneration

to wild-type at L4 stage **(Figure 6B** and **6F)**, whereas at older age (5d) *daf-2* mutants had a

much higher regenerative ability than wild-types (70% successful regeneration in *daf-2*

versus 12.5% in wild-type) **(Figure 6D** and **6F)**. Similarly to axonal response to injury [11],

we found that DAF-2 inhibits regeneration of aged dendrites through inhibition of DAF-16,

as *daf-16* mutants and *daf-2;daf-16* double mutants showed regenerative decline during aging

similar to wild-type **(Figure 6F)**. In conclusion, our results reveal that dendrite regeneration

following transection declines with aging, a phenotype that is dependent on DAF-2/IGF-1R

and its target DAF-16/FOXO.

AFF-1 mediates and restores dendrite regeneration in aged animals

A second fusogen in *C. elegans* is AFF-1 (Anchor cell Fusion Failure 1), a transmembrane

protein related to EFF-1 that executes several fusion events during development [34]. *aff-1*

mutants have no evident morphological phenotypes in the PVD neuron and expression has

not been detected in the PVD (**Figure 5-figure supplement 1**); moreover, when AFF-1 is

ectopically expressed in the PVD in an *eff-1* mutant background it does not retract excess

branching, demonstrating that it is unable to rescue the pruning defects of *eff-1* **(Figure 7-**

figure supplement 1). However, AFF-1 is required for dendrite fusion in response to injury

in L4s and young adults [35]. Adults carrying loss-of-function mutations in *aff-1* have severe

egg laying defects, shorter life spans and excretory system defects that prevented us from

studying regeneration in aging *aff-1* mutant animals.

We next tested whether AFF-1 can restore the regenerative ability of dendrites in

older animals by overexpressing AFF-1 specifically in the PVD (PVDp::AFF-1; **Figure 7-**

figure supplement 2). We found that menorahs of young animals expressing PVDp::AFF-1

appeared morphologically wild-type and responded similarly to dendrotomy **(Figure 7A,**

Movie S5). However, when 5-day old PVDp::AFF-1 animals were dendrotomized, the

percentage of regenerating worms by dendrite fusion was significantly higher compared to

wild-type animals (60-80% and 13%, respectively; **Figure 7B and 7C, Movie S6)**.

Thus, AFF-1-specific overexpression in the PVD, as well as systemic *daf-2* reduction

of function, enables dendrite regeneration in older animals **(Figures 6F and 7C)**.

Differential rejuvenation of fusion potential of transected dendrites by *daf-2(-)* and AFF-1(+)

We wished to further compare the regenerative modalities of AFF-1 overexpression and *daf-2* mutation following dendritic laser-induced severing. Dendritic fusion following injury can occur in three possible ways: 1) tertiary branch fusions that bypass the injury site ("menorah-menorah" fusion); 2) fusion of the proximal primary (1ry) branch to the detached distal 1ry; 3) both menorah-menorah fusion and 1ry-1ry fusion **(Figure 8A)**. We found that in wild-type and PVDp::AFF-1 animals at the L4 stage, the most prevalent mechanism of repair is menorah-menorah fusion, whereas in *daf-2* mutants menorah-menorah together with 1ry-1ry fusion (outcome 3) increased compared to wild-type **(Figure 8B-8D)**. In wild-type, we found degeneration in 50% (2-4d) and 80% (5-6d) of the dendrotomized animals **(Figure 8E and 8H)**. Adult *daf-2* mutant animals (5-6d) presented a response to injury that resembled that observed in L4 wild-type animals, with 70% of the animals showing regeneration, mainly via menorah-menorah fusions **(Figure 8F and 8I)**. In contrast, in PVDp::AFF-1 animals the response to injury followed a different mechanism, with more regeneration via enhanced 1ry-1ry fusion **(Figure 8G and 8J)**. However, the AFF-1 effect was unrelated to longevity as PVDp::AFF-1 animals had normal lifespans **(Figure 8-figure supplement 1)**. In summary, rejuvenation of the dendritic arbors uncovered here involves three distinct activities: i) EFF-1 pruning of old hyperbranched dendrites, ii) negative regulation of dendrite regeneration by DAF-2/ IGF-1R, and iii) AFF-1-induced auto-fusion of old transected primary dendrites **(Figure 9)**.

Discussion

Age-related deterioration of dendritic trees is insulin/IGF1-independent

Precise dendritic arborization is critical for proper functioning of neuronal networks, while defective dendritic development and maintenance lead to neuropathologies [44,45]. There is evidence from mammalian CNS showing increase in number and length of terminal dendritic arbors during aging, and reduced arborization in senile dementia [46]. However, it remains unclear whether there is a direct link between altered dendritic structure and reduced function in mammals. Axons of aging neurons also show altered morphologies in diverse species. In *C. elegans* these axonal alterations are not affected by organismal longevity [11,42,43,47]. Here we also found that dendritic alterations during aging are not affected by chronological age; however, unlike the PVD dendrites, in axons the *daf-2* mutation was found to delay the morphological alterations of aged animals [8-10]. Nevertheless, the *daf-2* effect in age-related axonal branching was reported to be uncoupled of its role in extending the lifespan of the worms [8,9]. It thus appears that DAF-2 is specifically involved in cell-autonomous pathways that maintain axonal, but not dendritic morphology. While progressive age-related hyperbranching of the PVD arbors was independent of insulin/IGF-1 pathway, overexpression of EFF-1 in the PVD was sufficient to partially rescue this phenotype in old animals (**Figure 5**). EFF-1 appears to act cell-autonomously to simplify the dendritic trees via its pruning activity mediated by branch retraction [23] (**Figures 5 and 9**).

AFF-1 restores regeneration of old broken dendrites via plasma membrane fusion

We further found that dendrite regeneration following PVD dendrotomy decreases with age. This decline was "rescued" in *daf-2* mutants (**Figure 6**). Axonal regeneration is known to decline with age in different organisms [11,42,43,47-49]. In GABA neurons of *C. elegans* the decline in axonal regeneration is also delayed in *daf-2* mutants [11]. However, these neurons do

not engage in self-fusion during regeneration, whereas in PVD dendrites the main outcome of

regeneration is auto-fusion. Recently, an important role of auto-fusion during axonal

regeneration was demonstrated in the PLM sensory neurons of *C. elegans*, a process

mediated by the fusogen EFF-1 [12,15]. In the PVD EFF-1 is not necessary for dendrite

reconnection following injury [35]. In the PVD dendrites, this role is fulfilled by the EFF-1

paralog AFF-1. Significantly, when AFF-1 is ectopically expressed, this fusogen plays a role

in maintenance of regenerative potential during aging (**Figures 7 and 9**). These findings

highlight fusion mechanisms as potential target for pharmacological intervention in

neuropathologies that result from both injury and aging.

Acknowledgments

We thank D. Cassel, M. Hilliard, and A. Sapir for critically reading this manuscript. M.

Heiman and C. Yip for providing CHB392, C. Smith and D. Miller for NC1841, and T.

Gattegno for BP709. We also want to thank R. Kishony and all members of the Podbilewicz

laboratory for discussion. This work was supported by Israel Science Foundation 443/12 and

ERC ELEGANSFUSION 268843 grants to BP. Some strains were provided by the CGC,

which is funded by NIH Office of Research Infrastructure Programs (P40 OD010440).

Author contributions: V.K., M.O.S. and B.P. designed the experiments. V.K. performed

most experiments. M.O.S. developed the system to study neuronal degeneration and

regeneration following dendrotomy using the PVD neurons. M.O.S. generated the

PVDp::AFF-1 and PVDp::EFF-1 transgenics and M.O.S. and V.K. tested the effects of EFF-

1 and AFF-1 over-expression on regenerating animals. B.P. supervised this work. V.K. and

B.P. wrote the paper with input from M.O.S.

The authors declare no conflict of interest.

This article contains supporting information including six figures and six movies

Materials and Methods

Nematode strains

Animals were maintained at 20°C, unless otherwise stated, according to standard protocols [50].
All experiments were performed on hermaphrodites. N2 served as wild-type, and the
following mutants and transgenic strains were used:

NC1841 *[wdIs52(F49H12.4::GFP); rwIs1(pmec-7::RFP)]* [24], outcrossed 4X, CHB392
[hmnEx133(ser-2prom3::kaede)], kindly provided by Candice Yip and Max Heiman [51],
BP709 *[hmnIs133 (ser-2prom3::kaede)]*, kindly provided by Tamar Gattegno, MF190
[hmIs4(des-2::gfp, pRF4)] [23]. CB1338 *[mec-3(e1318)IV]*, *Caenorhabditis* Genetics Center
(CGC), CB1611 *[mec-4(e1611)X]*, (CGC), BP176 *[hyEx23(des-2::eff-1, des-2::gfp, pRF4)]*
[23], BP177 *[hyEx392(des-2::eff-1, des-2::gfp, pRF4)]*, BP488 *[eff-1(hy21)II; hmIs4;*
hyEx39(des-2p::AFF-1,myo-2::GFP,KS)], BP906 *[daf-2(e1370)III; wdIs52; rwIs1]*, BP911
[hyEx39; hmnEx133], BP915 *[hyEx39; hmnIs133]*, BP551 *[hyEx391(des-2p::AFF-1, myo-*
2::GFP, KS); hmnEx133], BP919 *[daf-2(e1370)III; daf-16(mu86)I; hmnIs133]*, BP923
[daf-16(mu86)I; hmnIs133], BP924 *[daf-2(e1370)III; hmnIs133]*, BP925 *[mec-4(e1611)X;*
hmnIs133], BP926 *[daf-2(e1370)III;mec-4(e1611)X; hmnIs133]*, BP1056 *hyEx68[AFF-*
1::TY1::EGFP::3xFLAG, pRF4, KS]; dzIs53[F49H12.4p::mCherry], AFF-
1::TY1::EGFP::3xFLAG was kindly provided by transgenome project [52], BP500 *zzIs22*
[pJdC41(eff-1p::EFF-1::GFP), pRF4]; dzIs53 (pF49H12.4::mCherry) zzIs22 was kindly
provided by William Mohler [53].

Molecular Biology

The *ser-2prom3::Kaede* (pCY2 plasmid) is a kind gift from Candice Yip and Max Heiman [51].

It was constructed by digesting *ser-2prom3* with SbfI and AscI and Kaede with AscI and

NotI and then ligated together.

The *des-2p::AFF-1* construct (pME4 plasmid) was constructed by inserting an AFF-1

genomic fragment, digested from *hsp16.2::AFF-1* [34] with NheI and KpnI and cloned into

pME2 [23] cut with the same enzymes. *des-2p::AFF-1* was injected at a concentration of 0.1ng/ml

into *eff-1(hy21)* animals and three independent lines were obtained. For dendrotomy experiments,

two lines (*hyEx39* and *hyEx391*) were crossed into a wild-type background expressing the PVD

marker *ser-2p3::Kaede.*

Live imaging of worms

Time-lapse imaging and short imaging (one time point) of worms by Nomarski optics and

fluorescence microscopy was performed using a Nikon eclipse Ti inverted microscope with

Yokogawa CSU-X1 spinning disk or using the Zeiss Laser Scanning Microscope (LSM) 510

META. Animals were anesthetized using 0.1% tricaine and 0.01% tetramisole in M9

solution for 20-30 minutes, and then they were transferred to a 3% agar slide with an eyelash

attached to a toothpick. For short time imaging worms were often mounted on 3% agar slides

containing 5–10 mM NaN_3 instead. Image acquisition was done using Andor iQ or

Metamorph software, when using the spinning disk confocal (SDC), and Zen software when

using the LSM 510 meta microscope. Z-stacks were taken with PlanApochromat 60x oil

NA=1.4 objective using the SDC or 63x NA=1.4 objective using the LSM. Excitation of

GFP and green Kaede was done with 488 nm wavelength laser and 525 filter (6-15%, 50 ms

exposure time), RFP and red Kaede was excited with 561 nm wavelength laser and 607 filter

(15-20%, 50-100 ms exposure time). When using the sCMOS (Andor) camera z-stacks were

taken with ~0.23 μm z-step. With iXon EMCCD camera (Andor) z-stacks were taken with

~0.5 μm z- step, gain was ~100. When the LSM 510 meta was used, z-step was ~0.8 μm.

Multidimensional data was reconstructed as maximum intensity projections using the FIJI

software (NIH Image). Images were prepared using Imaris, FIJI and Adobe Photoshop CS5.

Final figures were constructed using Adobe Illustrator CS5.1.

Laser dendrotomies

Micropoint, pulsed nitrogen laser, or the tunable Chameleon Ultra Ti-Sapphire laser system

(two-photon), were used to sever the primary dendrite of the PVD. The Micropoint system

was used on the Nikon eclipse Ti inverted microscope with Yokogawa CSU-X1 spinning

disk. In order to transect neurons, we used 405 beamsplitter and 365nm dye cell and either

IQ or Metamorph software. We used highest levels with the attenuator plate (all the way in),

while the software controlled attenuator was adjusted between 80-90% when using IQ, and

30-50% when using Metamorph. Roughly 15 pulses at 10 Hz were administered for each cut

in IQ and Metamorph. For all worms the primary dendrite was injured anterior to cell body.

Animals were imaged and a z-stack was collected immediately after cut to confirm that the

injury was successful.

The two-photon system was used on the Zeiss LSM 510 META microscope. The

laser produces 200fs short pulses at 113MHz repetition rate and energy of 5nJ. In order to

cut neurons we used 820 nm wavelength and 20-30% laser power using the Zen software.

Worms were imaged immediately after cut to confirm that the injury was successful, and z-

stacks were taken using the spinning disk confocal system or the LSM 510 META as

described for live imaging of worms. After surgery, animals were recovered on NGM agar

plates seeded with OP50 in a drop of M9 and imaged again later, or time-lapse movies were

immediately acquired.

Regeneration was defined as continuation of the fluorescent signal between the distal and

proximal ends or using Kaede photoconversion (**Figure 6-figure supplement 1**; see below).

Significant differences between ages and genotypes were determined using *Fisher's exact*

test.

Kaede photoconversion

In order to verify that dendrites fuse as response to injury we used the photoconvertible

protein Kaede driven by a PVD specific promoter *ser-2prom3*. Irreversible photoconversion

of the green Kaede to red Kaede was achieved using Mosaic system on the Nikon eclipse Ti

inverted microscope with Yokogawa CSU-X1 spinning disk, at 405 nm, 20-50 ms exposure

time with 10-20 repeats across the region of interest, which was always the cell body, using

either IQ or Metamorph software.

Morphological quantitation of the PVD

Branch count and disorganization of menorahs was counted in the 100 μm around cell body

(unless otherwise stated), as previously described [23]. Lack of self-avoidance between

adjacent menorahs was determined in the same region as previously described [40]. Z-stack

maximum intensity projections of time-lapse movies were analyzed manually, using FIJI

software. Factorial analysis of variance (ANOVA) and *t-tests* were performed to compare

between genotypes and ages.

Harsh touch assay

Harsh touch assay was performed as previously described [23,41] and the experimenter was

blind to the genotype. The experiments were done in light touch mutant background (*mec-*

4(e1611)) [54]. *mec-3(e1338)* worms were used as negative control [41]. Significant differences

between the ages were determined by the $\chi 2$ test.

Life-span assay

Lifespan assays were carried out at 20°C as previously described [55]. A population of worms

was synchronized using hypochlorite and NaOH solution and ~100 animals at the L4

"Christmas tree" stage were placed on NGM plates containing 49.5 μM 5-Fluoro-2′-

deoxyuridine (FUDR, Sigma, F0503) seeded with OP50 *E. coli*, at a density of 20 worms per

6 cm plate. Adult day 1 was designated as the day after the L4 larval stage that served as

time 0. Animals were scored as dead or alive every 2-3 days until all animals died. An

animal was scored dead when it did not move or react at all to prodding with a platinum wire.

Animals that crawled off of the plate, were trapped in fungal infection, "exploded" (i.e.,

showed extruded tissue from the vulva), or showed internal hatching ("bagging"), were

excluded. If plates were contaminated, animals were transferred to fresh plates, using

platinum wire. Statistical significance was determined using the Log-Rank (Mantel-Cox)

test.

RNA Isolation, Reverse Transcription, and Quantitative Real-Time PCR.

Total RNA from Wild-type worms and des-2p::AFF-1 transgenic worms was isolated using

the RNeasy Micro kit following the manufacturer's protocol (Qiagen). RNA was used for

cDNA synthesis using the qScript cDNA Synthesis Kit (Quanta BioSciences) with random

primers. Real-time PCR was performed using AFF-1 and Fast SYBR® Green Master Mix

(Applied Biosystems). The PCR amplification conditions were as follows: 40 cycles of 95 °C

for 20 s and 60 °C for 20 s. Each PCR was run in triplicate. Data analysis and quantification

were performed using StepOne software V2.2 supplied by Applied Biosystems. To account

for the variability in the initial concentration of the RNA, results were normalized to the

ACT-1 housekeeping gene.

Statistical analysis

Statistical analysis was performed using the GraphPad online tool:

http://www.graphpad.com/quickcalcs/ for *t-tests, Fisher's exact test*s and χ^2 tests. Analysis of

variance (ANOVA) that was performed when more than two groups were compared and Log-

Rank survival test, using SPSS. *Freeman-Halton* extension of the *Fisher's exact test* (for 2x4

contingency table) was performed using the Vassarstats.net online tool, to compare between

the distributions of response to injury between different genotypes and ages.

References

1 Yankner, B. A., Lu, T. & Loerch, P. in *Annual Review of Pathology-Mechanisms of Disease* Vol. 3
Annual Review of Pathology-Mechanisms of Disease 41-66 (Annual Reviews, 2008).
2 Kenyon, C. J. The genetics of ageing. *Nature* **464**, 504-512, doi:10.1038/nature08980 (2010).
3 Hammarlund, M. & Jin, Y. Axon regeneration in C. elegans. *Current Opinion in Neurobiology* **27**,
199-207, doi:10.1016/j.conb.2014.04.001 (2014).
4 Ghosh-Roy, A. & Chisholm, A. D. Caenorhabditis elegans: A new model organism for studies of axon
regeneration. *Developmental Dynamics* **239**, 1460-1464, doi:10.1002/dvdy.22253 (2010).
5 Kenyon, C., Chang, J., Gensch, E., Rudner, A. & Tabtiang, R. A C-elegans mutant the lives twice as
long as wild- type *Nature* **366**, 461-464, doi:10.1038/366461a0 (1993).
6 Lin, K., Hsin, H., Libina, N. & Kenyon, C. Regulation of the Caenorhabditis elegans longevity protein
DAF-16 by insulin/IGF-1 and germline signaling. *Nature Genetics* **28**, 139-145, doi:10.1038/88850
(2001).
7 Murphy, C. T., McCarroll, S. A., Bargmann, C. I., Fraser, A., Kamath, R. S., Ahringer, J., Li, H. &
Kenyon, C. Genes that act downstream of DAF-16 to influence the lifespan of Caenorhabditis elegans.
Nature **424**, 277-284, doi:10.1038/nature01789 (2003).
8 Tank, E. M. H., Rodgers, K. E. & Kenyon, C. Spontaneous age-related neurite branching in
Caenorhabditis elegans. *J. Neurosci.* **31**, 9279-9288, doi:10.1523/jneurosci.6606-10.2011 (2011).
9 Toth, M. L., Melentijevic, I., Shah, L., Bhatia, A., Lu, K., Talwar, A., Naji, H., Ibanez-Ventoso, C.,
Ghose, P., Jevince, A., Xue, J., Herndon, L. A., Bhanot, G., Rongo, C., Hall, D. H. & Driscoll, M.
Neurite sprouting and synapse deterioration in the aging Caenorhabditis elegans nervous system. *J.*
Neurosci. **32**, 8778-8790, doi:10.1523/jneurosci.1494-11.2012 (2012).
10 Pan, C.-L., Peng, C.-Y., Chen, C.-H. & McIntire, S. Genetic analysis of age-dependent defects of the
Caenorhabditis elegans touch receptor neurons. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 9274-9279,
doi:10.1073/pnas.1011711108 (2011).
11 Byrne, A. B., Walradt, T., Gardner, K. E., Hubbert, A., Reinke, V. & Hammarlund, M. Insulin/IGF1
signaling inhibits age-dependent axon regeneration. *Neuron* **81**, 561-573,
doi:10.1016/j.neuron.2013.11.019 (2014).
12 Neumann, B., Coakley, S., Giordano-Santini, R., Linton, C., Lee, E. S., Nakagawa, A., Xue, D. &
Hilliard, M. A. EFF-1-mediated regenerative axonal fusion requires components of the apoptotic
pathway. *Nature* **517**, 219-U259, doi:10.1038/nature14102 (2015).

13   Neumann, B., Nguyen, K. C. Q., Hall, D. H., Ben-Yakar, A. & Hilliard, M. A. Axonal Regeneration
Proceeds Through Specific Axonal Fusion in Transected C. elegans Neurons. *Developmental*
Dynamics **240**, 1365-1372, doi:10.1002/dvdy.22606 (2011).
14   Hoy, R. R., Bittner, G. D. & Kennedy, D. Regeneration in Crustacean motoneurons - evidence for
axonal fusion *Science* **156**, 251-&, doi:10.1126/science.156.3772.251 (1967).
15   Ghosh-Roy, A., Wu, Z. L., Goncharov, A., Jin, Y. S. & Chisholm, A. D. Calcium and cyclic AMP
promote axonal regeneration in Caenorhabditis elegans and require DLK-1 kinase. *J. Neurosci.* **30**,
3175-3183, doi:10.1523/jneurosci.5464-09.2010 (2010).
16   Paltsyn, A., Komissarova, S., Dubrovin, I. & Kubatiev, A. Increased cell fusion in cerebral cortex may
contribute to poststroke regeneration. *Stroke research and treatment* **2013**, 869327-869327,
doi:10.1155/2013/869327 (2013).
17   Alvarez-Dolado, M., Pardal, R., Garcia-Verdugo, J. M., Fike, J. R., Lee, H. O., Pfeffer, K., Lois, C.,
Morrison, S. J. & Alvarez-Buylla, A. Fusion of bone-marrow-derived cells with Purkinje neurons,
cardiomyocytes and hepatocytes. *Nature* **425**, 968-973,
doi:http://www.nature.com/nature/journal/v425/n6961/suppinfo/nature02069_S1.html (2003).
18   Johansson, C. B., Youssef, S., Koleckar, K., Holbrook, C., Doyonnas, R., Corbel, S. Y., Steinman, L.,
Rossi, F. M. V. & Blau, H. M. Extensive fusion of haematopoietic cells with Purkinje neurons in
response to chronic inflammation. *Nature Cell Biology* **10**, 575-583, doi:10.1038/ncb1720 (2008).
19   Mohler, W. A., Shemer, G., del Campo, J. J., Valansi, C., Opoku-Serebuoh, E., Scranton, V., Assaf, N.,
White, J. G. & Podbilewicz, B. The type I membrane protein EFF-1 is essential for developmental cell
fusion. *Developmental Cell* **2**, 355-362, doi:10.1016/s1534-5807(02)00129-6 (2002).
20   Podbilewicz, B., Leikina, E., Sapir, A., Valansi, C., Suissa, M., Shemer, G. & Chernomordik, L. V.
The C-elegans developmental fusogen EFF-1 mediates homotypic fusion in heterologous cells and in
vivo. *Developmental Cell* **11**, 471-481, doi:10.1016/j.devcel.2006.09.004 (2006).
21   Shemer, G., Suissa, M., Kolotuev, I., Nguyen, K. C. Q., Hall, D. H. & Podbilewicz, B. EFF-1 Is
Sufficient to Initiate and Execute Tissue-Specific Cell Fusion in C. elegans. *Current Biology* **14**, 1587-
1591, doi:10.1016/j.cub.2004.07.059 (2004).
22   Tsalik, E. L., Niacaris, T., Wenick, A. S., Pau, K., Avery, L. & Hobert, O. LIM homeobox gene-
dependent expression of biogenic amine receptors in restricted regions of the C-elegans nervous
system. *Developmental Biology* **263**, 81-102, doi:10.1016/s0012-1606(03)00447-0 (2003).
23   Oren-Suissa, M., Hall, D. H., Treinin, M., Shemer, G. & Podbilewicz, B. The fusogen EFF-1 controls
sculpting of mechanosensory dendrites. *Science* **328**, 1285-1288, doi:10.1126/science.1189095 (2010).
24   Smith, C. J., Watson, J. D., Spencer, W. C., O'Brien, T., Cha, B., Albeg, A., Treinin, M. & Miller, D.
M., III. Time-lapse imaging and cell-specific expression profiling reveal dynamic branching and
molecular determinants of a multi-dendritic nociceptor in C. elegans. *Dev. Biol.* **345**, 18-33,
doi:10.1016/j.ydbio.2010.05.502 (2010).
25   Chatzigeorgiou, M., Yoo, S., Watson, J. D., Lee, W. H., Spencer, W. C., Kindt, K. S., Hwang, S. W.,
Miller, D. M., Treinin, M., Driscoll, M. & Schafer, W. R. Specific roles for DEG/ENaC and TRP
channels in touch and thermosensation in C. elegans nociceptors. *Nat. Neurosci.* **13**, 861-U106,
doi:10.1038/nn.2581 (2010).
26   Salzberg, Y., Diaz-Balzac, C. A., Ramirez-Suarez, N. J., Attreed, M., Tecle, E., Desbois, M.,
Kaprielian, Z. & Buelow, H. E. Skin-derived cues control arborization of sensory dendrites in
Caenorhabditis elegans. *Cell* **155**, 308-320, doi:10.1016/j.cell.2013.08.058 (2013).
27   Dong, X., Liu, O. W., Howell, A. S. & Shen, K. An extracellular adhesion molecule complex patterns
dendritic branching and morphogenesis. *Cell* **155**, 296-307, doi:10.1016/j.cell.2013.08.059 (2013).
28   Dong, X., Shen, K. & Bülow, H. E. Intrinsic and Extrinsic Mechanisms of Dendritic Morphogenesis.
Annual Review of Physiology **77**, null, doi:doi:10.1146/annurev-physiol-021014-071746 (2015).
29   Liu, O. W. & Shen, K. The transmembrane LRR protein DMA-1 promotes dendrite branching and
growth in C. elegans. *Nat. Neurosci.* **15**, 57-U74, doi:10.1038/nn.2978 (2012).
30   Maniar, T. A., Kaplan, M., Wang, G. J., Shen, K., Wei, L., Shaw, J. E., Koushika, S. P. & Bargmann,
C. I. UNC-33 (CRMP) and ankyrin organize microtubules and localize kinesin to polarize axon-
dendrite sorting. *Nat. Neurosci.* **15**, 48-U66, doi:10.1038/nn.2970 (2012).
31   Cohen, E., Chatzigeorgiou, M., Husson, S. J., Steuer-Costa, W., Gottschalk, A., Schafer, W. R. &
Treinin, M. Caenorhabditis elegans nicotinic acetylcholine receptors are required for nociception.
Molecular and Cellular Neuroscience **59**, 85-96, doi:10.1016/j.mcn.2014.02.001 (2014).
32   Husson, S. J., Costa, W. S., Wabnig, S., Stirman, J. N., Watson, J. D., Spencer, W. C., Akerboom, J.,
Looger, L. L., Treinin, M., Miller, D. M., III, Lu, H. & Gottschalk, A. Optogenetic Analysis of a
Nociceptor Neuron and Network Reveals Ion Channels Acting Downstream of Primary Sensors.
Current Biology **22**, 743-752, doi:10.1016/j.cub.2012.02.066 (2012).

33   Wei, X., Howell, A. S., Dong, X., Taylor, C. A., Cooper, R. C., Zhang, J., Zou, W., Sherwood, D. R. &
Shen, K. The unfolded protein response is required for dendrite morphogenesis. *eLife* **4**, e06963,
doi:10.7554/eLife.06963 (2015).
34   Sapir, A., Choi, J., Leikina, E., Avinoam, O., Valansi, C., Chernomordik, L. V., Newman, A. P. &
Podbilewicz, B. AFF-1, a FOS-1-regulated fusogen, mediates fusion of the anchor cell in C-elegans.
Dev. Cell **12**, 683-698, doi:10.1016/j.devcel.2007.03.003 (2007).
35   Oren-Suissa, M., Gattegno, T., Kravtsov, V. & Podbilewicz, B. Extrinsic repair of injured dendrites as
a paradigm for regeneration by fusion. *Submitted to eLife* (2016).
36   Procko, C., Lu, Y. & Shaham, S. Glia delimit shape changes of sensory neuron receptive endings in C.
elegans. *Development* **138**, 1371-1381, doi:10.1242/dev.058305 (2011).
37   Stone, C. E., Hall, D. H. & Sundaram, M. V. Lipocalin signaling controls unicellular tube development
in the Caenorhabditis elegans excretory system. *Developmental Biology* **329**, 201-211,
doi:10.1016/j.ydbio.2009.02.030 (2009).
38   Rasmussen, J. P., English, K., Tenlen, J. R. & Priess, J. R. Notch signaling and morphogenesis of
single-cell tubes in the C-elegans digestive tract. *Developmental Cell* **14**, 559-569,
doi:10.1016/j.devcel.2008.01.019 (2008).
39   Lenard, A., Daetwyler, S., Betz, C., Ellertsdottir, E., Belting, H.-G., Huisken, J. & Affolter, M.
Endothelial cell self-fusion during vascular pruning. *PLoS biology* **13**, e1002126-e1002126,
doi:10.1371/journal.pbio.1002126 (2015).
40   Smith, C. J., Watson, J. D., VanHovene, M. K., Colon-Ramos, D. A. & Miller, D. M., III. Netrin
(UNC-6) mediates dendritic self-avoidance. *Nat. Neurosci.* **15**, 731-737, doi:10.1038/nn.3065 (2012).
41   Way, J. C. & Chalfie, M. The mec-3 gene of Caenorhabditis-elegans requires its own product for
maintained expression and is expressed in 3 neuronal cell-types *Genes Dev.* **3**, 1823-1833,
doi:10.1101/gad.3.12a.1823 (1989).
42   Verdu, E., Ceballos, D., Vilches, J. J. & Navarro, X. Influence of aging on peripheral nerve function
and regeneration. *J. Periph. Nerv. Syst.* **5**, 191-208, doi:10.1046/j.1529-8027.2000.00026.x (2000).
43   Zou, Y., Chiu, H., Zinovyeva, A., Ambros, V., Chuang, C.-F. & Chang, C. Developmental decline in
neuronal regeneration by the progressive change of two intrinsic timers. *Science* **340**, 372-376,
doi:10.1126/science.1231321 (2013).
44   Jan, Y. N. & Jan, L. Y. Branching out: mechanisms of dendritic arborization. *Nat. Rev. Neurosci.* **11**,
316-328, doi:10.1038/nrn2836 (2010).
45   Koleske, A. J. Molecular mechanisms of dendrite stability. *Nat. Rev. Neurosci.* **14**, 536-550,
doi:10.1038/nrn3486 (2013).
46   Buell, S. J. & Coleman, P. D. Dendritic growth in the aged human-brain and failure of growth in senile
dementia *Science* **206**, 854-856, doi:10.1126/science.493989 (1979).
47   Hammarlund, M., Nix, P., Hauth, L., Jorgensen, E. M. & Bastiani, M. Axon regeneration requires a
conserved MAP kinase pathway. *Science* **323**, 802-806, doi:10.1126/science.1165527 (2009).
48   Gabel, C. V., Antonie, F., Chuang, C.-F., Samuel, A. D. T. & Chang, C. Distinct cellular and molecular
mechanisms mediate initial axon development and adult-stage axon regeneration in C. elegans.
Development **135**, 1129-1136, doi:10.1242/dev.013995 (2008).
49   Pestronk, A., Drachman, D. B. & Griffin, J. W. Effects of aging on nerve sprouting and regeneration
Exp. Neurol. **70**, 65-82, doi:10.1016/0014-4886(80)90006-0 (1980).
50   Brenner, S. Genetics of Caenorhabditis-elegans *Genetics* **77**, 71-94 (1974).
51   Yip, Z. C. & Heiman, M. G. Duplication of a Single Neuron in C. elegans Reveals a Pathway for
Dendrite Tiling by Mutual Repulsion. *Cell reports*, doi:10.1016/j.celrep.2016.05.003 (2016).
52   Sarov, M., Schneider, S., Pozniakovski, A., Roguev, A., Ernst, S., Zhang, Y., Hyman, A. A. & Stewart,
A. F. A recombineering pipeline for functional genomics applied to Caenorhabditis elegans. *Nat*
Methods **3**, 839-844, doi:10.1038/nmeth933 (2006).
53   del Campo, J. J., Opoku-Serebuoh, E., Isaacson, A. B., Scranton, V. L., Tucker, M., Han, M. &
Mohler, W. A. Fusogenic activity of EFF-1 is regulated via dynamic localization in fusing somatic
cells of C. elegans. *Curr Biol* **15**, 413-423 (2005).
54   Chalfie, M. & Sulston, J. Developmental genetics of the mechanosensory neurons of Caenorhabditis-
elegans *Dev. Biol.* **82**, 358-370, doi:10.1016/0012-1606(81)90459-0 (1981).
55   Apfeld, J. & Kenyon, C. Cell nonautonomy of C-elegans daf-2 function in the regulation of diapause
and life span. *Cell* **95**, 199-210, doi:10.1016/s0092-8674(00)81751-1 (1998).
540

541 **Figures**

543
544
Figure 1.
PVD's dendritic alterations during aging

(A-D) PVD in wild-type (wt; **A** and **C**) and *daf-2* mutants (**B** and **D**) at L4 stage and 9 days
of adulthood (9d). Upper panels, magnified boxed areas with "menorah" colored according to
the scheme **(A).** c, cell body. Scale bars, 50 μm and 20 μm in magnified images. Anterior is
left and ventral is down in all the figures. Filled arrows, self-avoidance defects. Empty
arrows, disorganized menorahs. **(E-I)** Quantitation of phenotypes shown in images.
Percentages are for 100 μm of length around cell body. Error bars, ± s.e.m. *p* values from *t*
tests: **p<0.01, **p<0.001*. Number of animals analyzed: *daf-2* n ≥ 8 and wild-types n ≥ 5.

Figure 2.

DAF-2-DAF-16 control menorah self-avoidance at L4 stage

(A) Image showing four wild-type menorahs labeled with GFP, two of them do not overlap
(filled arrow) and two show defective self-avoidance (empty arrow).

(B) Percentage of defects in self-avoidance in 100 μm of length around PVD cell body at L4
stage in different genotypes, n ≥ 7. Defective self-avoidance increased in *daf-2* mutants in a
daf-16-dependent manner. Error bars are ± s.e.m. *p* value from *t* test: * *p<0.05*.

Figure 3.

Adults show a reduction in dendritic plasticity

(A) Time-lapse confocal projections of 5 days of adulthood (5d) wild-type animal. Boxed
areas are enlarged in the lower panel and reveal dynamic growth (arrows) and retraction
(arrow heads) of branches. Scale bars, 10 μm.

(B) Growth and retraction rates in μm/minute of branches at L4 stage and 5d as measured
from time-lapse movies. Number of branches analyzed ≥ 18 from 4 L4 animals and 2 5d
animals. Error bars are ± s.e.m. *p* values from *t* tests: * *p<0.01, ** p<0.001*. See also **Movie**
S1.

577
578

Figure 4.

Decline in response to harsh touch during aging in both wild-type and *daf-2*

(A-D) PVD's cell-body region of wt and light-touch insensitive *mec-4* mutants at the L4
stage and at 5 days of adulthood (5d). Scale bars are 20 μm.

(E) Total number of branches counted in 100 μm of length around cell-body in wt and *mec-4*
mutants at L4 and 5d (no significant differences between wt and *mec-4)*. Error bars are ±
s.e.m.

(F) Percentage of animals responding to harsh touch by escaping away from the stimulus. At
the age of 5d, the response declined in *mec-4* mutants and in *daf-2;mec-4* double mutants. *p*
values from χ2 test as compared between 1d to other ages (3d or 5d) for each genotype: * *p*
<0.05. Number of animals: *mec-3* ≥ 50; *mec-4* ≥ 70; *daf-2;mec-4* ≥ 40; wt ≥ 26.

Figure 5.

EFF-1 overexpression in the PVD simplifies "aged" menorahs

(A-D) Inverted fluorescence images of PVD neurons.

(**A** and **C)** Represent wild-type neurons from L4 and 9 days of adulthood (9d).

(**B** and **D)** Represent EFF-1 overexpression under PVD specific promoter (PVDp) at L4 and
9d. In each panel one menorah (boxed) is enlarged and colored (see **Figure 1A**). Scale bars,
20 μm and 10 μm in the enlarged images.

(E-G) Graphs showing number of branches in 100 μm of length around cell body. Error bars,
± s.e.m. *p* values from *t* tests: * *p<0.05,* ** *p<0.001,* *** *p<0.0001* Number of animals: n ≥
4. PVDp::EFF-1 line 1 and 2 correspond to worms carrying the extrachromosomal arrays
hyEx392 and *hyEx23,* respectively.

Figure 6.

daf-2(-) restores regenerative ability of aged animals via *daf-16*

(A-D) PVD neurons immediately after cut and 24 h later in wild-type and *daf-2* mutants at L4
and 5 days of adulthood (5d), as indicated. Schematic illustrations below negative images.
Red lightings, injury sites. Blue arrows, fusion sites (successful regeneration). Scale bars, 20
μm.

(E) Response to injury of wild-types at L4-young adult (YA) and 2-3d within a short time (3-
6 h) after injury. Number of animals: n=7 (L4 –YA) and n=4 (2-3d).

(F) Percentage of successfully regenerating animals within 24-48 h post injury. *p* values from
Fisher's exact tests: * *p<0.05, ** p<0.01, *** p<0.001*. Number of animals: n ≥ 8. See also
Figure 6-figure supplement 1 and **Movies S2-S4**.

Figure 7.

AFF-1 enhances regeneration by dendrite fusion in 5d-old animals

(A-B) Inverted fluorescence images of PVD neurons immediately after cut and 24 hours (h)
later in animals overexpressing AFF-1 in the PVD (PVDp::AFF-1), at L4 and 5 days of
adulthood (5d), as indicated on each image (schematic drawings below each image). Red
lightning marks the injury site and blue arrows point at fusion sites (menorah-menorah
fusion)**.** Scale bars, 20 μm.

(C) Percentage of successfully regenerating animals within 24-48h after injury. *p* value from
Fisher's exact test: * *p<0.05; ** p<0.001*. Number of animals: n ≥ 10. See also **Figure 7-**
figure supplement 1 and **Movies S5-S6**. PVDp::AFF-1 line 1 and 2 correspond to worms
carrying the extrachromosomal arrays *hyEx390* and *hyEx391,* respectively.

Figure 8.

daf-2(-) and AFF-1(+) rejuvenate fusion potential of broken dendrites

(A) Cartoon representing four possible outcomes of dendrotomy. Dendrite auto-fusion indicates successful regeneration (outcomes 1-3), if fusion does not occur degeneration takes place (outcome 4).

(B-J) Percentages of wild-type worms (wt), *daf-2* mutants and AFF-1 overexpressing animals (PVDp::AFF-1) at different ages, divided into the four different types of response to injury as described in (A). Genotype and age are listed above each plot. *p* values from *Fisher's exact tests*: * *p<0.05*, ** *p<0.01*, *** *p<0.001*. Additional significant differences were found between: (C and F), (C and I) and between (B and J) (*p<0.05*). See also **Figure 8-figure supplement 1.**

Figure 9.

Model of morphological and regenerative antiaging activities on dendritic arbors

Morphological aging of PVD menorahs is a progressive and dynamic process that results in
loss of self-avoidance, disorganization and hyperbranching. When the fusogen EFF-1 is
ectopically expressed in the PVD neuron it has an antiaging activity that involves pruning of
hyperbranched dendritic trees.

Aged menorahs lose their *regenerative potential* because following laser-induced
dendrotomy old neurons usually fail to auto-fuse broken dendrites and undergo degeneration.
DAF-2/ IGF-1R negatively regulates the regeneration process. When the fusogen AFF-1 is
ectopically expressed in the PVD neuron it has an antiaging activity that promotes auto-
fusion of old transected primary dendrites.

661
662 **Supplementary Figures**
663

Figure 1-figure supplement 1.

Anterior-posterior branching gradient sharpens as animals age

(A-B) Average number of branches per 100 μm of body length (A4-P2; A, anterior; P,
posterior; CB, cell body; d, days of adulthood) in wild-type (wt) and *daf-2* mutants. Gradient
sharpening is delayed in *daf-2*. We performed three way analysis of variance (ANOVA) to
compare between wt (**A**) and *daf-2* (**B**). The contribution of body position and age to
branching were significant (p=0.0001 and p=1.4X10^{-6}, respectively). Age*genotype, body
position*genotype, body position*age and genotype alone were significant (p=0.007,
p=0.001, p=0.043 and p=0.023, respectively). Number of animals: wt n ≥ 2; *daf-2* n ≥ 3.

(C-H) Each bar is a 100 μm long PVD region. Bars and schematic menorahs on the bottom:
2ry, magenta; 3ry, red; 4ry, green; 5ry, orange; 6ry, yellow; 7ry, brown.

678
679
Figure 5-figure supplement 1.

AFF-1 and EFF-1 expression pattern in L4 and aging worms

(A) A 30kb fosmid-based reporter for AFF-1 shows expression in the epithelial seam-cells at
the L4 stage (marked with arrowheads), and in 5-day adults. AFF-1 is not expressed at the
PVD in any stage. Multiple transgenic lines were analyzed and a similar expression pattern
was observed in all.

(B) EFF-1 expression at the L4 stage and 5-days adults was analyzed using an EFF-1::GFP
translational chimera: 7.5 kb *eff-1* promoter driving the full-length *eff-1* genomic coding
sequence was fused to GFP [53]. EFF-1 is expressed in the seam-cells (arrowheads) and the
PVD cell body (arrows and inset) in vesicles. Expression persists after 5 days.

690
691
Figure 6-figure supplement 1.

Fusion is a crucial step in regeneration of injured dendrites

(A) Scheme describing the fusion assay using a photoconvertible fluorescent marker Kaede.
Primary dendrite is injured using a laser, then the animal is recovered and imaged again after
~24 hours in green and red channels. Green Kaede is photoconverted using a U.V. laser
focused to the cell body. After ~1 hour the spread of red Kaede is observed again.

(B) Upper panel: confocal reconstructions of wild-type L4 animal immediately after
dendrotomy, green fluorescence 24 hours after injury and before photoconversion, red
fluorescence before photoconversion and red fluorescence an hour after photoconversion of
the cell-body. In the lower panel are illustrations. Red kaede passed into the distal area,
meaning that the broken dendrite fused to the proximal part.

(C) A negative control showing L4 wild-type animal in which the primary branch did not
regenerate within 24 hours. Proximal menorahs are fused (arrows), but it did not bridge the
gap between the distal and proximal stumps (arrowheads). Indeed red kaede did not spread
into the detached distal stump. The order of images is the same as described in **(B)**.
Lightings point at injury sites, arrows point at fusion sites. Scale bars, 20 μm.

eff-1(hy21); PVDp::AFF-1

708
709
Figure 7-figure supplement 1.

AFF-1 does not retract PVD's dendrites. *eff-1(hy21)* animals expressing AFF-1 in the
PVD (PVDp::AFF-1) show excess branching, similarly to *eff-1* mutants

Thus the fusogen AFF-1 cannot rescue *eff-1* loss-of-function phenotype when expressed in
the PVD. Arrow points to hypodermal cells expressing GFP after ectopic PVD neurite-
epidermal fusion. Scale bar, 10 μm.
716

717
718
719 **Figure 7-figure supplement 2.**

720 Real-time RT-PCR analysis of AFF-1 mRNA expression in wild-type and transgenic worms
721 over expressing AFF-1 in the PVD. Bar graph representing the fold changes of mRNA levels
722 quantified by normalization to the *act-1* gene as an internal control. *p* value from *t* test
723 ****p<0.0001.*

Figure 8-figure supplement 1.

AFF-1 overexpressing animals exhibit wild-type life-span

Life-span curve of wild-type (blue) and AFF-1 overexpression in the PVD (PVDp::AFF-1, red) . Median life-span is 14 days of adulthood for wild-types and 13 for PVDp::AFF-1 containing animals.

Legends for movies

Movie S1. Dendritic plasticity of aged wild-type

Inverted fluorescence confocal maximum intensity projections time lapse movie of wild-type

PVD marked with GFP at 5 days of adulthood. Some branches show dynamic growth and

retraction (arrow). The counter in this and other movies are in Hours: minutes.

Movie S2. Young wild-types regenerate rapidly after dendrotomy

Inverted fluorescence confocal maximum intensity projections time lapse movie of two L4

wild-type animals. PVD marked with Kaede. Both animals regenerated within 2 hours after

laser induced dendrotomy via menorah-menorah fusion (blue arrows). Red lightnings mark

the sites of injury.

Movie S3. Kaede photoconversion confirms auto-fusion as part of regeneration

Confocal time lapse movie of wild-type L4 worm expressing the photoconvertible protein

Kaede in the PVD. The 1ry branch of this worm was dendrotomized 24 hours prior to the

movie, then green kaede was photoconverted in the cell body, using U.V. laser (marked with

purple asterisk). In the left panel the red channel is seen, with red for the photoconverted

kaede protein traveling from the proximal part to the fused distal part (blue arrows). In the

right panel the green channel is shown. Injury site is indicated by yellow lightning. The

photoconverted red-Kaede was transferred to the distal part (top of the image) by menorah-

menorah fusion and there was no 1ry-1ry fusion.

Movie S4. Adult wild-type responds slowly to injury

Inverted fluorescence confocal maximum intensity projections time lapse movie of wild-type animal at the age of 2 days of adulthood. PVD is marked by Kaede. There is neither regeneration nor degeneration within the first 3 hours after cut. Red lightning points at injury site.

Movie S5. PVDp::AFF-1 L4 regenerates similarly to wild-type

Inverted fluorescence confocal maximum intensity projections time lapse movie of PVDp::AFF-1 (AFF-1 overexpression in the PVD) L4 nematode. The worm was injured and imaged immediately after. In this movie, it rapidly regenerates (within 3 hours from cut) via menorah-menorah fusion, as marked by blue arrows and by novel outgrowth that fuses to the distal primary and by that bridges the gap (yellow arrow). Red lightning points to injury site. Red arrowhead marks a menorah that degenerated during the movie.

Movie S6. PVDp::AFF-1 5d old animal reconnects after cut

Confocal time lapse movie of 5 days old PVDp::AFF-1 transgenic animal, beginning 5 hours post injury. Growth is seen near site of dendrotomy. The primary branch grows toward the distal primary and reconnects with it 10 hours after injury (blue arrows). After this time point the worm was recovered and imaged 15 hours later (25 hours after cut), where regeneration via novel outgrowth from 1ry to distal fragment of 1ry branch fusion and menorah-menorah fusion is indicated by blue arrows. Red lightning points at injury site.