Title: Detecting the impact of temperature on transmission of Zika, dengue and

chikungunya using mechanistic models

Short title: Temperature predicts Zika, dengue, and chikungunya transmission

Authors: Erin A. Mordecai[a*], Jeremy M. Cohen[b], Michelle V. Evans[c], Prithvi Gudapati[a], Leah

R. Johnson[b], Catherine A. Lippi[c], Kerri Miazgowicz[d], Courtney C. Murdock[d,e], Jason R. Rohr[b],

Sadie J. Ryan[c,f,g,h], Van Savage[i,j], Marta S. Shocket[a,k], Anna Stewart Ibarra[l], Matthew B.

Thomas[m], Daniel P. Weikel[n]

Affiliations:

[a]Biology Department, Stanford University, 371 Serra Mall, Stanford, CA 94305

[b]Department of Integrative Biology, University of South Florida, 4202 East Fowler Ave,

SCA110 Tampa, FL 33620

[c]Department of Geography, University of Florida, PO Box 117315, Turlington Hall, Gainesville,

FL 32611

[d]Center for Tropical and Emerging Global Disease, Department of Infectious Diseases,

University of Georgia College of Veterinary Medicine, 501 D.W. Brooks Drive, Athens, GA

30602

[e]Odum School of Ecology, University of Georgia, 140 E. Green St., Athens, GA

30602[f]Emerging Pathogens Institute, University of Florida, P.O. Box 100009, 2055 Mowry Road

Gainesville, FL 32610

[g]Center for Global Health and Translational Science, Department of Microbiology and

Immunology, Weiskotten Hall, SUNY Upstate Medical University, Syracuse, NY 13210

[h]School of Life Sciences, College of Agriculture, Engineering, and Science, University of

KwaZulu Natal, Private Bag X01, Scottsville, 3209, KwaZulu Natal, South Africa

[i]Department of Ecology and Evolutionary Biology, University of California Los Angeles and

Department of Biomathematics, University of California Los Angeles, Los Angeles, CA 90095

[j]Santa Fe Institute, 1399 Hyde Park Rd, Santa Fe, NM 87501

[k]Department of Biology, Indiana University, 1001 E. 3rd St., Jordan Hall 142, Bloomington, IN

47405

[l]Center for Global Health and Translational Sciences, SUNY Upstate Medical University,

Syracuse, NY13210

[m]Department of Entomology and Center for Infectious Disease Dynamics, Penn State University,

112 Merkle Lab, University Park, PA 16802

[n]Department of Biostatistics, University of Michigan, 1415 Washington Heights, Ann Arbor, MI

48109

[*]Correspondence to: Erin Mordecai. Biology Department, Stanford University, 371 Serra Mall,

Stanford, CA 94305. (650) 497-7447. emordeca@stanford.edu

Keywords: Zika, dengue, chikungunya, temperature, vector transmission, *Aedes aegypti*, *Aedes*

albopictus

Abstract

Recent epidemics of Zika, dengue, and chikungunya have heightened the need to understand the

seasonal and geographic range of transmission by *Aedes aegypti* and *Ae. albopictus* mosquitoes.

We use mechanistic transmission models to derive predictions for how the probability and

magnitude of transmission for Zika, chikungunya, and dengue change with mean temperature,

and we show that these predictions are well matched by human case data. Across all three

viruses, models and human case data both show that transmission occurs between 18-34°C with

maximal transmission occurring in a range from 26-29°C. Controlling for population size and

two socioeconomic factors, temperature-dependent transmission based on our mechanistic model

is an important predictor of human transmission occurrence and incidence. Risk maps indicate

that tropical and subtropical regions are suitable for extended seasonal or year-round

transmission, but transmission in temperate areas is limited to at most three months per year even

if vectors are present. Such brief transmission windows limit the likelihood of major epidemics

following disease introduction in temperate zones.

Author Summary (150-200 words)

Understanding the drivers of recent Zika, dengue, and chikungunya epidemics is a major public

health priority. Temperature may play an important role because it affects mosquito

transmission, affecting mosquito development, survival, reproduction, and biting rates as well as

the rate at which they acquire and transmit viruses. Here, we measure the impact of temperature

on transmission by two of the most common mosquito vector species for these viruses, *Aedes*

aegypti and *Ae. albopictus*. We integrate data from several laboratory experiments into a

mathematical model of temperature-dependent transmission, and find that transmission peaks at

26-29°C and can occur between 18-34°C. Statistically comparing model predictions with recent

observed human cases of dengue, chikungunya, and Zika across the Americas suggests an

important role for temperature, and supports model predictions. Using the model, we predict that

most of the tropics and subtropics are suitable for transmission in many or all months of the year,

but that temperate areas like most of the United States are only suitable for transmission for a

few months during the summer (even if the mosquito vector is present).

Main Text

Epidemics of dengue, chikungunya, and Zika are sweeping through the Americas, and are part of a global public health crisis that places an estimated 3.9 billion people in 120 countries at risk [1]. Dengue virus (DENV) distribution and intensity in the Americas has increased over the last three decades, infecting an estimated 390 million people (96 million clinical) per year [2]. Chikungunya virus (CHIKV) emerged in the Americas in 2013, causing 1.8 million suspected cases from 44 countries and territories (www.paho.org). In the last two years, Zika virus (ZIKV) has spread throughout the Americas, causing 714,636 suspected and confirmed cases, with many more unreported (http://ais.paho.org/phip/viz/ed_zika_cases.asp, as of January 5, 2017). The growing burden of these diseases (including links between Zika infection and both microcephaly and Guillain-Barré syndrome [3]) and potential for spread into new areas creates an urgent need for predictive models that can inform risk assessment and guide interventions such as mosquito control, community outreach, and education.

Predicting transmission of DENV, CHIKV, and ZIKV requires understanding the ecology of the vector species. For these viruses the main vector is *Aedes aegypti*, a mosquito that prefers and is closely affiliated with humans, while *Ae. albopictus*, a peri-urban mosquito, is an important secondary vector [4,5]. We expect one of the main drivers of the vector ecology to be the climate, particularly temperature. For that reason, mathematical and geostatistical models that incorporate climate information have been valuable for predicting and responding to *Aedes* spp. spread and DENV, CHIKV, and ZIKV outbreaks [5–10].

The effects of temperature in ectotherms are largely predictable from fundamental metabolic and ecological processes. Survival, feeding, development, and reproductive rates predictably respond to temperature across a variety of ectotherms, including mosquitoes [11,12].

Because these traits help to determine transmission rates, the effects of temperature on

transmission should also be broadly predictable from mechanistic models that incorporate

temperature-dependent traits. Here, we introduce a model based on this framework that

overcomes several major gaps that currently limit our understanding of climate suitability for

transmission. Specifically, we develop models of temperature-dependent transmission for *Ae.*

aegypti and *Ae. albopictus* that are (a) mechanistic, facilitating extrapolation beyond the current

disease distribution, (b) parameterized with biologically accurate unimodal thermal responses for

all mosquito and virus traits that drive transmission, and (c) validated against human dengue,

chikungunya, and Zika case data across the Americas.

We synthesize available data to characterize the temperature-dependent traits of the

mosquitoes and viruses that determine transmission intensity. With these thermal responses, we

develop mechanistic temperature-dependent virus transmission models for *Ae. aegypti* and *Ae.*

albopictus. We then ask whether the predicted effect of temperature on transmission is consistent

with patterns of actual human cases over space and time. To do this, we validate the models with

DENV, CHIKV, and ZIKV human incidence data at the country scale from the Americas from

2014-2016. To isolate temperature dependence, we also statistically controlled for population

size and two socioeconomic factors that may influence transmission. If temperature

fundamentally limits transmission potential, transmission should only occur at actual

environmental temperatures that are predicted to be suitable, and conversely, areas with low

predicted suitability should have low or zero transmission (i.e., false negative rates should be

low). By contrast, low transmission may occur even when temperature suitability is high because

other factors like vector control can limit transmission (i.e., the false positive rate should be

higher than the false negative rate). Finally, if the simple mechanistic model accurately predicts

climate suitability for transmission, then we can use it to map climate-based transmission risk of

DENV, CHIKV, ZIKV, and other emerging pathogens transmitted by *Ae. aegypti* and *Ae.*

albopictus seasonally and geographically.

Results

Temperature-dependent transmission

Data gathered from the literature [9,13–15,15–21,21–30] revealed that all mosquito traits

relevant to transmission—biting rate, egg-to-adult survival and development rate, adult lifespan,

and fecundity—respond strongly to temperature and peak between 23°C and 34°C for the two

mosquito species (*Ae. aegypti* in Fig. 1 and *Ae. albopictus* in Fig. S1). DENV extrinsic

incubation and vector competence peak at 35°C [31–37] and 31-32°C [31,32,34,38],

respectively, in both mosquitoes—temperatures at which mosquito survival is low, limiting

transmission potential (Figs. 1, S1). Appropriate thermal response data were not available for

CHIKV and ZIKV extrinsic incubation and vector competence.

Fig. 1. Thermal responses of *Ae. aegypti* and DENV traits that drive transmission (data sources

listed in Table S2). Informative priors based on data from additional *Aedes* spp. and flavivirus

studies helped to constrain uncertainty in the model fits (see Materials and Methods; Table S3).

Points and error bars indicate the data means and standard errors (for display only; models were

fit from the raw data). Black solid lines are the mean model fits; red dashed lines are the 95%

credible intervals. Thermal responses for *Ae. albopictus* are shown in Fig. S1.

We estimated the posterior distribution of $R_0(T)$ and used it to calculate key temperature

values that indicate suitability for transmission: the mean and 95% credible intervals (95% CI)

on the critical thermal minimum, maximum, and optimum temperature for transmission by the

two mosquito species. At constant temperature, *Ae. aegypti* transmission peaked at 29.1°C (95%

CI: 28.4 – 29.8°C), and declined to zero below 17.8°C (95% CI: 14.6 – 21.2°C) and above

34.6°C (95% CI: 34.1 – 35.6°C) (Fig. 2). *Ae. albopictus* transmission peaked at 26.4°C (95% CI:

25.2 – 27.4°C) and declined to zero below 16.2°C (95% CI: 13.2 – 19.9°C) and above 31.6°C

(95% CI: 29.4 – 33.7°C) (Fig. 2). Overall, the thermal response curve for *Ae. albopictus* is

shifted towards lower temperatures than *Ae. aegypti,* so *Ae. albopictus* transmission is better

suited to colder environments. For a more realistic scenario in which daily temperature ranged

over 8°C, the transmission peak, minimum, and maximum were slightly lower for both *Ae.*

aegypti (28.5°C, 13.5°C, 34.2°C, respectively) and *Ae. albopictus* (26.1°C, 11.9°C, and 28.3°C,

respectively). The lower thermal maximum under fluctuating temperatures occurs because we

incorporated empirically supported irreversible lethal effects of temperatures that exceed thermal

maxima for survival (see Materials and Methods).

Fig. 2. Relative R_0 across constant temperatures (°C; top) for *Ae. albopictus* (light blue) and *Ae.*

aegypti (dark blue), and histograms of the posterior distributions of the critical thermal minimum

(bottom left), temperature at peak transmission (bottom middle), and critical thermal maximum

(bottom right; all in °C). Solid lines: mean posterior estimates; dashed lines: 95% credible

intervals. R_0 curves normalized to a 0-1 scale for ease of comparison and visualization.

The posterior distribution of $R_0(T)$ allows us to evaluate uncertainty in key temperature

values that define the transmission range, including critical thermal minimum, maximum, and

optimum. Uncertainty was higher for the critical thermal minimum for transmission than for the

maximum or optimum, and the two mosquito species overlapped most for this outcome (Fig. 2,

bottom panels). This occurred because several trait thermal responses increase gradually from

low to mid temperatures but decline more steeply at high temperatures (Fig. 1), so uncertainty is

greatest at low temperatures. *Ae. aegypti* has a substantially higher optimum and maximum

temperature than *Ae. albopictus* (Fig. 2) due to its greater rates of adult survival at high

temperatures (see Supplementary Materials for sensitivity analyses).

Model validation

We used generalized linear models (GLM) to ask whether the predicted relationship

between temperature and transmission, $R_0(T)$, was consistent with observed human cases of

DENV, CHIKV, and ZIKV. Specifically, we assessed whether $R_0(T)$ was an important predictor

of the probability of autochthonous transmission occurring and of the incidence given that

transmission occurred. We also controlled for human population size, virus species, and two

socioeconomic factors. (Note that we focused on testing the $R_0(T)$ model, rather than on

constructing the best possible statistical model of human case data.) To do this, we used the

version of the *Ae. aegypti* $R_0(T)$ model that includes 8°C daily temperature range, along with

country-scale weekly case reports of DENV, CHIKV, and ZIKV in the Americas and the

Caribbean between 2014-2016. We first addressed the fact that countries with larger populations

have greater opportunities for (large) epidemics by creating two predictors that incorporate

scaled $R_0(T)$ and population size. In the models of the probability of autochthonous transmission

occurring we used the product of the posterior probability that $R_0(T) > 0$ (which we notate as

GR_0) and the log of population size (p) to give $log(p)*GR_0$. In the models of incidence given that

transmission does occur we used the log of the product of the posterior mean of $R_0(T)$ and

population size, $log(p*R_0(T))$. To control for several socioeconomic factors that might obscure

the impact of temperature, we also included log of gross domestic product (GDP) and log percent

of GDP in tourism (using logs to improve normality). These are potential indicators of

investment in and/or success of vector control and infrastructure improvements that prevent

transmission. By comparing models that included the $R_0(T)$ metric alone, socioeconomic factors

alone, or both, we tested whether $R_0(T)$ was an important predictor of observed transmission

occurrence and incidence (see Table S4). Note that $R_0(T)$ is out of sample because it is derived

and calculated strictly from laboratory data on mosquitoes, and we perform a validation analyses

for $R_0(T)$ using independent case incidence reports. For this validation step we assessed model

adequacy for the transmission data in two ways. First we used the full dataset for case incidence

reports to select the best model (Table S4) and determine whether or not our predicted value of

relative $R_0(T)$ based on laboratory data was included in the model ("within sample" analysis).

Second we used a bootstrapping approach where models were fit on subsets of the case incidence

data that were randomly sampled and then predictive accuracy of the competing models (Table

S4) was assessed on left-out data ("out of sample" analysis).

For the probability of autochthonous transmission occurring, the model that included both

the $R_0(T)$ predictor and socioeconomic predictors had overwhelming support based on Bayesian

Information Criterion (BIC; model PA5 relative probability = 1, Table S4). Based on deviance

explained, the models that included $R_0(T)$, with or without the socioeconomic predictors out-

performed the model that did not include $R_0(T)$ (Table S4; Figs. 3A, S2). In analyses of out-of-

sample accuracy, models that included the $R_0(T)$ metric (with or without the socioeconomic

factors) were surprisingly accurate. They predicted the probability of transmission with 86-91%

out-of-sample accuracy for DENV (Table S4). For CHIKV and ZIKV, models that included the

$R_0(T)$ metric or population alone had 66-69% out-of-sample accuracy (Table S4). There were no

significant differences in out-of-sample accuracy between the top four models but for both

DENV and CHIKV/ZIKV the best model was significantly better than the worst model (see

Supplementary Code for full results). The lower out-of-sample accuracy for CHIKV and ZIKV

likely reflects the much lower frequency of positive values and the lower total sample size of this

dataset. All results were similar for a set of models that separated GR_0 from population size, so

for simplicity we show the model predictors that combines GR_0 and population size here (see

Table S4 and Supplementary Code for results of other models). Further, from a biological

perspective, the combined model better describes what we know about disease systems: if either

the probability of $R_0(T)$ being greater than zero is small or population size is very small,

transmission is unlikely to occur. Together, these analyses suggest that $R_0(T)$ is an important

predictor of transmission occurrence, but that CHIKV and ZIKV need further data to better

explain the probability of transmission occurrence (Figs. 3A, S2).

Fig. 3. *Ae. aegypti $R_0(T)$* and population size predict the probability and magnitude of

transmission of DENV, CHIKV, and ZIKV across the Americas. A, *log(p)*GR_0* (the posterior

probability that $R_0(T) > 0$ times the log of population size) versus the probability of local

transmission in the data. B, *log(p*R_0(T))* (log of $R_0(T)$ times the population size) versus the log

of incidence, given that it exceeds the threshold for local transmission. Tick-marks and points:

human transmission occurrence and incidence data, respectively, by country-week in the

Americas and Caribbean. Lines and shaded areas: mean and 95% CI from GLM fits for DENV

(blue) and CHIKV and ZIKV (red). For simplicity, we show the models that only include the

covariates $log(p)*GR_0$ or $log(p*R_0(T))$, respectively, and do not include the socioeconomic

covariates (models PA6 and IM4 in Table S4). For each case report data point, $log(p)*GR_0$ and

$log(p*R_0(T))$ were calculated at the mean temperature 10 weeks prior to the reporting week [39].

$R_0(T)$ was also an important predictor of incidence, given that autochthonous

transmission did occur. Within-sample, incidence was best predicted by the model that included

both $R_0(T)$ and the socioeconomic predictors (model IM5 in Table S4) based on BIC (relative

probability = 1). The models that included $R_0(T)$ out-performed those that did not based on

deviance explained (Table S4). In out-of-sample validation, the models that included $R_0(T)$

explained the magnitude of incidence based on mean absolute percentage error (85-86%

accuracy versus 83% accuracy for models that did not include $R_0(T)$; Table S4), but this

difference was not statistically significant. For illustration, we show the simpler model that only

contains the $R_0(T)$ predictor in the main text (Fig. 3B; model IM1 in Table S4). Notably, the

models that contained $R_0(T)$ predicted incidence well for all three viruses, despite the lower

incidence of CHIKV and ZIKV.

Although predicted $R_0(T)$ correlated with the observed occurrence and magnitude of

human incidence for all three viruses, these observed incidence metrics were higher for DENV

than for CHIKV and ZIKV. While the reason for this difference is unclear, the most likely

explanation is that DENV is much more established in the region, so it is more likely to be

detected, diagnosed, and reported. Because ZIKV and CHIKV are newly emerging, they may not

have fully saturated the region at this early stage.

The ability of the model to explain the probability and magnitude of transmission is

notable given the coarse scale of the human incidence versus mean temperature data (i.e.,

country-scale means), the lack of CHIKV- and ZIKV-specific trait thermal response data to

inform the model, the nonlinear relationship between transmission and incidence, and all the

well-documented factors other than temperature that influence transmission. Together, these

analyses show simple mechanistic models parameterized with laboratory data on mosquitoes and

dengue virus are consistent with observed temperature suitability for transmission. Moreover, the

similar responses of human incidence of ZIKV, CHIKV, and DENV to temperature suggest that

the thermal ecology of their shared mosquito vectors is a key determinant of outbreak location,

timing, and intensity.

Mapping climate suitability for transmission

The validated model can be used to predict where transmission is not excluded (posterior

probability that $R_0(T) > 0$, a conservative estimate of transmission risk). Considering the number

of months per year at which mean temperatures do not prevent transmission, large areas of

tropical and subtropical regions, including Puerto Rico and parts of Florida and Texas, are

currently suitable year-round or seasonally (Fig. 4). These regions are fundamentally at risk for

DENV, CHIKV, ZIKV, and other *Aedes* arbovirus transmission during a substantial part of the

year (Fig. 4). Indeed, DENV, CHIKV, and/or ZIKV local transmission has occurred in Texas,

Florida, Hawaii, and Puerto Rico (www.cdc.gov). On the other hand, many temperate regions

experience temperatures suitable for transmission three months or less per year (Fig. 4), and the

virus incubation periods in humans and mosquitoes restrict the transmission window even

further. Temperature thus limits the potential for the viruses to generate extensive epidemics in

temperate areas even where the vectors are present. Moreover, many temperate regions with

seasonally suitable temperatures currently lack *Ae. aegypti* and *Ae. albopictus* mosquitoes,

making vector transmission impossible (Fig. 4, black line). The posterior distribution of $R_0(T)$

also allows us to map months of risk with different degrees of uncertainty (e.g., 97.5%, 50%, and

2.5% posterior probability that that $R_0 > 0$), ranging from the most to least conservative (Fig.

S4).

Fig. 4. Map of predicted temperature suitability for virus transmission by *Ae. albopictus* and *Ae.*

aegypti. Color indicates the consecutive months in which temperature is permissive for

transmission (predicted $R_0 > 0$) for *Aedes* spp. transmission based on the minimum likely range

(> 97.5% posterior probability that $R_0 > 0$). Black lines indicate the CDC estimated range for the

two *Aedes* spp. in the United States. Model suitability predictions combine temperature mean

and 8°C daily variation and are informed by laboratory data (Figs. 1, S1) and validated against

field data (Fig. 3).

Discussion

Temperature is an important driver of—and limitation on—vector transmission, so

accurately describing the temperature range and optimum for transmission of DENV, CHIKV,

and ZIKV is critical for predicting their geographic and seasonal patterns of spread [12,40]. We

directly estimated the temperature – transmission relationship using mechanistic transmission

models for each mosquito species (Fig. 2). These models are built using empirical estimates of

the (unimodal) effects of temperature on mosquito and pathogen traits that drive transmission,

including survival, development, reproduction, and biting rates (Figs. 1, S1). Because these trait

thermal responses are unimodal across the majority of ectotherm taxa and traits, and because the

traits combine nonlinearly to drive transmission, the emergent relationship between temperature

and transmission is difficult to infer directly from field data or from individual trait responses.

Here, we present a model of temperature-dependent DENV, CHIKV, and ZIKV transmission

that advances on previous models because it is mechanistic, fitted from experimental trait data,

and validated against independent human case data at a broad geographic scale (Fig. 3).

Mechanistic understanding is valuable for extrapolating beyond the current spatial and

temporal range of transmission (Fig. 4), as compared to environmental niche models, for

example [5,41,42]. Of the six previous mechanistic temperature-dependent models of DENV,

CHIKV, or ZIKV transmission by *Ae. aegypti* and *Ae. albopictus* that we were able to reproduce,

three had similar thermal optima [7,43,44] while the other three had dramatically higher optima

(3-6°C) [9,45] (Fig. S5). Two models predicted much greater suitability for transmission at low

temperatures [45], four predicted greater suitability at high temperatures [7,9,45], and two were

very similar to ours [43,44] (Fig. S5). Only one of these previous models was (like ours)

statistically validated against independent data not used to estimate model parameters, and its

predictions were very similar to those of our model [43]. Other mechanistic and environmental

niche models could not be directly compared with ours [5,10,40–42], either because fully

reproducible equations, parameters, and/or code were not provided or because their predicted

marginal effects of temperature were not displayed. Visually, our maps are similar to maps based

on a previous model of *Ae. aegypti* and *Ae. albopictus* persistence suitability indices [40]. Recent

environmental niche models of Zika distribution have shown similar but more constrained

predicted distributions of environmental suitability, in part because these models include not just

temperature suitability but also further environmental, socioeconomic, and demographic

constraints [5,41,42,46].

Even though the thermal response data are imperfect—for example, CHIKV and ZIKV

thermal response data are missing—and the human case data are reported at a coarse spatial

scale, the validation analyses suggest that $R_0(T)$ is an important predictor of both the probability

of transmission occurring and the magnitude of incidence for DENV, CHIKV, and ZIKV. This

has several key implications. First, temperature-dependent transmission is pervasive enough to

be detected at a coarse spatial scale. Second, dynamics of the mosquito predict transmission for a

suite of *Ae. aegypti*-transmitted viruses, without additional virus-specific information. Third,

climate and socio-economic factors combine to shape variation in incidence across countries.

Finally, these simple predictors explain a substantial proportion of the variance in both the

probability and intensity of transmission.

Predicting arbovirus transmission at a higher spatial resolution and precision will require

more detailed information on factors like the exposure and susceptibility of human populations,

environmental variation (e.g., oviposition habitat availability, seasonal and daily temperature

variation), and socioeconomic factors. However, as a first step our mechanistic model provides

valuable insight because it makes broad predictions about suitable environmental conditions for

transmission, it is mechanistic and grounded in experimental trait data, it is validated against

independent human case data, and its predictions are applicable across three different viruses.

Using these thermal response models as a scaffold, additional drivers could be incorporated to

obtain more precise and specific predictions about transmission dynamics, which could in turn be

used for public health and vector control applications. For this purpose, all code and data used in

the models are available as Supplementary Files.

The socio-ecological conditions that enabled CHIKV, ZIKV, and DENV to become the

three most important emerging vector-borne diseases in the Americas make the emergence of

additional *Aedes*-transmitted viruses likely (potentially including Mayaro, Rift Valley fever,

yellow fever, Uganda S, or Ross River viruses). Efforts to extrapolate and to map temperature

suitability (Fig. 4) will be critical for improving management of both ongoing and future

emerging epidemics. Mechanistic models like the one presented here are useful for extrapolating

the potential geographic range of transmission beyond the current envelope of environmental

conditions in which transmission occurs (e.g., under climate change and for newly invading

pathogens). Accurately estimating temperature-driven transmission risk in both highly suitable

and marginal regions is critical for predicting and responding to future outbreaks of these and

other *Aedes*-transmitted viruses.

Materials and Methods

<u>Temperature-sensitive R_0 models</u>

We constructed temperature-dependent models of transmission using a previously

developed R_0 framework. We modeled transmission rate as the basic reproduction rate, R_0—the

number of secondary infections that would originate from a single infected individual introduced

to a fully susceptible population. In previous work on malaria, we adapted a commonly used

expression for R_0 for vector transmission to include the temperature-sensitive traits that drive

mosquito population density :

$$R_0(T) = \left(\frac{a(T)^2\, b(T)\, c(T)\, e^{-\mu(T)/PDR(T)}\, EFD(T)\, p_{EA}(T)\, MDR(T)}{N\, r\, \mu(T)^3} \right)^{1/2} \tag{1}$$

Here, (*T*) indicates that the trait is a function of temperature, *T*; *a* is the per-mosquito biting rate,

b is the proportion of infectious bites that infect susceptible humans, *c* is the proportion of bites

on infected humans that infect previously uninfected mosquitoes (i.e., $b*c$ = vector competence),

μ is the adult mosquito mortality rate (lifespan, $lf = 1/\mu$), PDR is the parasite development rate

(i.e., the inverse of the extrinsic incubation period, the time required between a mosquito biting

an infected host and becoming infectious), EFD is the number of eggs produced per female

mosquito per day, p_{EA} is the mosquito egg-to-adult survival probability, MDR is the mosquito

immature development rate (i.e., the inverse of the egg-to-adult development time), N is the

density of humans, and r is the human recovery rate. For each temperature-sensitive trait in each

mosquito species, we fit either symmetric (Quadratic, $-c(T - T_0)(T - T_m)$) or asymmetric (Brière,

$cT(T - T_0)(T_m - T)^{1/2}$) unimodal thermal response models to the available empirical data [47]. In

both functions, T_0 and T_m are respectively the minimum and maximum temperature for

transmission, and c is a positive rate constant.

We consider a normalized version of the R_0 equation such that it is rescaled to range from

zero to one with the value of one occurring at the unimodal peak. Although absolute values of R_0

that are used to determine when transmission is stable depend on additional factors not captured

in our model, the minimum and maximum temperatures for which $R_0 > 0$ map exactly onto our

normalized equations, allowing us to accurately calculate whether or not transmission should be

possible at all. Empirical estimates of absolute values of R_0 are difficult to obtain in any case, but

it is much easier to determine whether transmission is occurring and for how long. While

different model formulations for predicting R_0 versus temperature can produce results with

different magnitudes and potentially different overall shapes [48], the temperatures for which R_0

is above or below zero (or one) are mostly model independent. For instance, two competing

models differ only by whether or not the formula in equation (1) is squared, but the square of a

number (e.g., an absolute R_0 value) greater than one is always greater than one, and the square of

a number less than one is always less than one. Therefore, the threshold temperatures at which

absolute $R_0 > 0$ or absolute $R_0 > 1$ will be exactly the same for either choice of formula (Fig. S6).

Similarly, because different expressions for R_0, including the square of equation (1), map

monotonically onto our function, they will produce identical estimates for the temperatures at

which transmission declines to zero and peaks (Fig. S6). Consequently, our use of relative R_0

adequately describes the nonlinear relationship between mosquito and virus traits and

transmission.

We fit the trait thermal responses in equation (1) based on an exhaustive search of

published laboratory studies that fulfilled the criterion of measuring a trait at three or more

constant temperatures, ideally capturing both the rise and the fall of each unimodal curve (Tables

S1-S2). Constant-temperature laboratory conditions are required to isolate the direct effect of

temperature from confounding factors in the field and to provide a baseline for estimating the

effects of temperature variation through rate summation [49]. We attempted to obtain raw data

from each study, but if they were not available we collected data by hand from tables or digitized

data from figures using WebPlotDigitizer [50]. We obtained raw data from Delatte [19] and Alto

[21] for the *Ae. albopictus* egg-to-adult survival probability (*pEA*), mosquito development rate

(MDR), gonotrophic cycle duration (GCD, which we assumed was equal to the inverse of the

biting rate) and total fecundity (TFD) (Table S2). Data did not meet the inclusion criterion for

CHIKV or ZIKV vector competence (*b, c*) or extrinsic incubation period (*EIP*) in either *Ae.*

albopictus or *Ae. aegypti*. Instead, we used DENV EIP and vector competence data, combined

with sensitivity analyses.

Following Johnson *et al.* [51], we fit a thermal response for each trait using Bayesian

models. We first fit Bayesian models for each trait thermal response using uninformative priors

($T_0 \sim$ Uniform (0, 24), $T_m \sim$ Uniform (25, 45), $c \sim$ Gamma (1, 10) for Brière and $c \sim$ Gamma (1,

1) for Quadratic fits) chosen to restrict each parameter to its biologically realistic range (i.e., $T_0 <$

T_m and we assumed that temperatures below 0°C and above 45°C were lethal). Any negative

values for all thermal response functions were truncated at zero, and thermal responses for

probabilities (p_{EA}, b, and c) were also truncated at one. We modeled the observed data as arising

from a normal distribution with the mean predicted by the thermal response function calculated

at the observed temperature, and the precision τ, ($\tau = 1/\sigma$), distributed as $\tau \sim$ Gamma (0.0001,

00001). We fit the models using Markov Chain Monte Carlo (MCMC) sampling in JAGS, using

the R [52] package *rjags* [53]. For each thermal response, we ran five MCMC chains with a

5000-step burn-in and saved the subsequent 5000 steps. We thinned the posterior samples by

saving every fifth sample and used the samples to calculate R_0 from 15-40°C, producing a

posterior distribution of R_0 versus temperature. We summarized the relationship between

temperature and each trait or overall R_0 by calculating the mean and 95% highest posterior

density interval (HPD interval; a type of credible interval that includes the smallest continuous

range containing 95% of the probability, as implemented in the *coda* package [54]) for each

curve across temperatures.

428   We fit a second set of models for each mosquito species that used informative priors to

reduce uncertainty in R_0 versus temperature and in the trait thermal responses. In these models,

we used Gamma-distributed priors for each parameter T_0, T_m, c, and τ fit from an additional

'prior' dataset of *Aedes* spp. trait data that did not meet the inclusion criteria for the primary

dataset (Table S3). We found that these initial informative priors could have an overly strong

influence on the posteriors, in some cases drawing the posterior distributions well away from the

primary dataset, which was better controlled and met the inclusion criteria. We accounted for our

lower confidence in this data set by increasing the variance in the informative priors, by

multiplying all hyperparameters (i.e., the parameters of the Gamma distributions of priors for T_0,

T_m, and c) by a constant k to produce a distribution with the same mean but $1/k$ times larger

variance. We chose the value of k based on our relative confidence in the prior versus main data.

Thus we chose $k = 0.5$ for $b, c,$ and PDR and $k = 0.01$ for lf. This is the main model presented in

the text (Fig. 2). It is comparable to some but not all previous mechanistic models for *Ae. aegypti*

and *Ae. albopictus* transmission (Fig. S5). Results of our main model, fit with informative priors,

did not vary substantially from the model fit with uninformative priors (Figs. S7-S8).

<u>Incorporating daily temperature variation in transmission models</u>

Because organisms do not typically experience constant temperature environments in

nature, we incorporated the effects of temperature variation on transmission by calculating a

daily average R_0 assuming a daily temperature range of 8°C, across a range of mean

temperatures. This range is consistent with daily temperature variation in tropical and subtropical

environments but lower than in most temperate environments. At each mean temperature, we

used a Parton-Logan model to generate hourly temperatures and calculate each temperature-

sensitive trait on an hourly basis [55]. We assumed an irreversible high-temperature threshold

above which mosquitoes die and transmission is impossible [56,57]. We set this threshold based

on hourly temperatures exceeding the critical thermal maximum (T_m in Tables S1-S2) for egg-to-

adult survival or adult longevity by any amount for five hours or by 3°C for one hour. We

averaged each trait over 24 hours to obtain a daily average trait value, which we used to calculate

relative R_0 across a range of mean temperatures. We used this model in the validation against

human cases (Fig. 3) and the risk map (Fig. 4).

<u>Model validation with DENV, CHIKV, and ZIKV incidence data</u>

To validate the model, we used data on human cases of DENV, CHIKV, and ZIKV at the

country scale and mean temperature during the transmission window. Using statistical models

(as described below), we estimated the effects of predicted $R_0(T)$ on the probability of local

transmission and the magnitude of incidence, controlling for population size and several

socioeconomic factors. We downloaded and manually entered Pan American Health

Organization (PAHO) weekly case reports for DENV and CHIKV for all countries in the

Americas (North, Central, and South America and the Caribbean Islands) from week 1 of 2014

to week 8 of 2015 for CHIKV and from week 52 of 2013 to week 47 of 2015 for DENV

(www.paho.org). ZIKV weekly case reports for reporting districts (e.g., provinces) within

Colombia, Mexico, El Salvador, and the US Virgin Islands were available from the CDC

Epidemic Prediction Initiative (https://github.com/cdcepi/) from November 28, 2015 to April 2,

2016. We aggregated the ZIKV data into country-level weekly case reports to match the spatial

resolution of the DENV, CHIKV, and covariate data.

Temperature data collection

We matched the DENV, CHIKV, and ZIKV incidence data with temperature using daily

temperature data from METAR stations in each country, averaged at the country level by

epidemic week. A previous study found a six-week lagged relationship between temperature and

oviposition for *Aedes aegypti* in Ecuador [39]. Assuming that the subsequent transmission,

disease development, medical care-seeking, and case reporting in humans takes an additional

four weeks, we assumed *a priori* a ten-week lag between temperature and incidence (i.e., mean

temperature for the week that is ten weeks prior to each case report). METAR stations are

internationally standardized weather reporting stations that report hourly temperature and

precipitation measures. Outlier weather stations were excluded if they reported a daily maximum

temperature below 5°C or a daily minimum temperature above 40°C during the study period,

extremes that would certainly eliminate the potential for transmission in a local area. Because

case data are reported at the country level, we needed a collection of weather stations in each

country that accurately represent weather conditions in the areas where transmission occurs,

excluding extreme areas where transmission is unlikely. For the study period of October 1, 2013

through April 30, 2016, we downloaded daily temperature data for each station from Weather

Underground using the *weatherData* package in R [58]. We removed all data from Chile because

it spans so much latitude and the terrain is so diverse that its country-level mean is unlikely to be

very representative of the temperature where an outbreak occurred.

Socioeconomic covariate data

We accessed available data on projected 2016 gross domestic product (GDP) for

countries of interest via the International Monetary Fund's World Economic Outlook Database

(http://www.imf.org/external/ns/cs.aspx?id=28). The direct and total contributions of tourism to

GDP in 2016 were compiled from World Travel and Tourism Council economic impact reports

(http://www.wttc.org/research/economic-research/economic-impact-analysis/country-

reports/#undefined). We retrieved population size data for 2013-2015 from the United Nations

Population Division (https://esa.un.org/unpd/wpp/Download/Standard/Population/) and averaged

them across the three years for each country. Throughout the analyses below, unless otherwise

specified, we used the natural log of the population size and of GDP as our predictors. We have

two reasons for this choice. The first is that, intuitively, the relative order of magnitude of the

population/GDP is more important in determining observed outbreak sizes or probabilities than

their absolute sizes. Second, population sizes and GDPs across countries tend to exhibit clumped

patterns with a few outliers that are much larger than the others. From a statistical perspective,

using the un-transformed populations (or GDPs) results in those few large/rich countries having

very high leverage in the analysis, and thus potentially skewing the results. Taking a log of the

population better balances these predictors and is the standard accepted approach when using

these kinds of predictors in regression models.

<u>Validation analyses with human incidence versus temperature datasets</u>

To validate the $R_0(T)$ model while controlling for population and socio-economic factors,

we used generalized linear regression on the weekly case count data. Importantly, we focused on

testing whether the case counts were consistent with the transmission – temperature relationship

predicted from our model, rather than on maximizing the variation explained in the statistical

model. We are more specifically interested in understanding autochthonous transmission (i.e.,

locally acquired, not just imported cases). We set country-level thresholds for the number of

cases defining autochthonous transmission for our three diseases separately, based on current

transmission understanding: seven cases of CHIKV, 70 cases of DENV, and three cases of

ZIKV. We derived these thresholds in the following way. First, we looked for data on outbreaks

of travel related cases in countries that are not expected to experience any local transmission. For

instance, in 2014 Canada experienced 320 confirmed, travel-related cases of chikungunya

(http://www.phac-aspc.gc.ca/publicat/ccdr-rmtc/15vol41/dr-rm41-01/rapid-eng.php), equivalent

to an average of more than six cases per week. Thus, to be conservative in our estimates, we set

the threshold of transmission as seven cases/week for CHIKV. The reported weekly cases of

DENV transmission in our study sample are considerably higher than for CHIKV (mean DENV

incidence was nearly 100 times higher mean CHIKV incidence). We chose a moderately high

threshold of 70 cases in a week (i.e., 10 times higher than the CHIKV threshold based on

Canadian cases) to reflect higher overall incidence and increased potential for travel related

cases. We examined the sensitivity of the results to choice of threshold by varying it from 25 to

100, and we found qualitatively similar results for all thresholds that we tested. As ZIKV is not

as well established as either CHIKV or DENV at this time, smaller numbers of cases may

indicate autochthonous transmission. Consequently, we chose a threshold of three cases for

ZIKV (approximately half the CHIKV threshold). Further, the results were fairly sensitive to the

ZIKV threshold as many locations have small numbers of cases. Since higher thresholds exclude

a very large proportion of available case data making analysis impossible, we used the slightly

less conservative threshold of three cases for autochthonous transmission of ZIKV. The resulting

data consisted of zeros for no transmission and positive case counts when transmission is

presumed to be occurring. To model these data, we used a hurdle model that first uses logistic

regression on the presence/absence of local transmission data to understand the factors correlated

with local transmission occurring or not (PA analysis). Then we modeled the log of incidence

(number of new cases per reporting week) for positive values with a gamma generalized linear

models (GLM; i.e., incidence analysis).

We were interested in understanding whether $R_0(T)$ was an important predictor of human

transmission occurrence and incidence, after controlling for potentially confounding factors like

population size and socioeconomic conditions. To do this, we fit a series of models with different

subsets of predictors that included $R_0(T)$, the socioeconomic variables with population, or both

(see Table S4 for full models). To control for human population size, we created new metrics

based on $R_0(T)$ and population size to use for validation against the PAHO incidence data. We

define GR_0, which is the posterior probability that $R_0(T) > 0$. We use $log(p)*GR_0$, where p is the

population size, as the relevant R_0-based predictor for the PA analysis. For the incidence

analysis, we instead use $log(p*R_0(T))$ as the predictor. In all cases log refers to the natural

logarithm. For simplicity, we refer to these as the $R_0(T)$ metrics hereafter and in the Results.

In both the PA and incidence analyses, we first used the full data sets to examine which

of the candidate models best described the data. Randomized quantile residuals indicated that the

logistic and gamma GLM models were performing adequately. We compared the approximate

model probabilities, calculated from the BIC scores, as well as the proportion of deviance

explained (D^2) from each model. Next we examined the performance of the models in predicting

out of sample, for both PA and incidence analyses. To do this we created 1000 random

partitions, where 90% of the data were used to train the model and 10% were used for testing. In

the PA analyses we classified each partition based on presence/absence, with separate

classification thresholds for DENV versus CHIKV/ZIKV as these grouping had much different

probabilities of occurrence. We assessed the performance of the model for the PA analysis based

on the mean misclassification rate. In the incidence analyses we assessed the model performance

based on the predictive mean absolute percentage error (MAPE). Since differences in prediction

success between the models in both the PA and incidence analyses were not statistically

significant, we present the simpler models that only include the $R_0(T)$ metrics in the main text

(Fig. 3) and the models that additionally include socioeconomic covariates in the Supplementary

Information (Figs. S2-S3). We plotted the model predictions as a function of the $R_0(T)$ metrics

together with the observed data for the PA and incidence analyses using the R package *visreg*

[59].

The residuals of the incidence model exhibit "inverse trumpeting," in which residual

variation is larger at low than high predicted incidence (Fig. S9). This occurs in part because we

forced the model to go through the origin, i.e., no transmission when $R_0(T)$ or the population size

is equal to zero. However, the data did sometimes show transmission where we did not expect it,

potentially because of imported cases, errors in reporting, or small pockets of transmission

suitability in countries or times that are otherwise unsuitable on average. More local-scale case

reporting that separates autochthonous from travel-associated cases would be needed to tease

apart the source of this error.

Mapping temperature suitability for transmission

Using our validated model, we were interested in where the temperature was suitable for

Ae. aegypti and/or *Ae. albopictus* transmission for some or all of the year to predict the potential

geographic range of outbreaks in the Americas. We visualized the minimum, median, and

maximum extent of transmission based on probability of occurrence thresholds from the R_0

models for both mosquitoes. We calculated the number of consecutive months in which the

posterior probability of $R_0 > 0$ exceeds a threshold of 0.025, 0.5, or 0.975 for both mosquito

species, representing the maximum, median, and minimum likely ranges, respectively. The

minimum range is shown in Fig. 4 and all three ranges are overlaid in Fig. S4. This analysis

indicates the predicted seasonality of temperature suitability for transmission geographically, but

does not indicate its magnitude. To generate the maps, we cropped monthly mean temperature

rasters from 1950-2000 for all twelve months (Worldclim; www.worldclim.org/) to the Americas

(*R*, *raster* package, *crop* function) and assigned cells values of one or zero depending on whether

the probability that $R_0 > 0$ exceeded the threshold at the temperatures in those cells. We then

synthesized the monthly grids into a single raster that reflected the maximum number of

consecutive months where cell values equaled one. The resulting rasters were plotted in ArcGIS

10.3, overlaying the three cutoffs (Fig. S4). We repeated this process for both mosquito species.

Acknowledgments

Barry Alto, Krijn Paaijmans, Francis Ezeakacha, and Helene Delatte kindly provided raw data

used in the analyses. We gratefully acknowledge the Centers for Disease Control and Prevention

Epidemic Predictions Initiative (CDC EPI) for collating and sharing the Zika incidence data on

GitHub (https://zenodo.org/record/48946#.Vz-EM2bb8ys).

Supporting Information Legends

S1 Appendix. Supplementary Results, References, Figures S1-S15, and Tables S1-S3.

S2 Appendix. Table S4. Model validation results.

S3 Appendix. Online code, data, and analyses. Available as a ZIP file on Figshare:

https://figshare.com/s/b79bc7537201e7b5603f, DOI:

https://dx.doi.org/10.6084/m9.figshare.4563928.

References:

1. Brady OJ, Gething PW, Bhatt S, Messina JP, Brownstein JS, Hoen AG, et al. Refining the global spatial limits of dengue virus transmission by evidence-based consensus. PLOS Negl Trop Dis. 2012;6: e1760. doi:10.1371/journal.pntd.0001760

2. Bhatt S, Gething PW, Brady OJ, Messina JP, Farlow AW, Moyes CL, et al. The global distribution and burden of dengue. Nature. 2013;496: 504–507. doi:10.1038/nature12060

3. Rasmussen SA, Jamieson DJ, Honein MA, Petersen LR. Zika virus and birth defects — reviewing the evidence for causality. N Engl J Med. 2016;374: 1981–1987. doi:10.1056/NEJMsr1604338

4. Scott TW, Takken W. Feeding strategies of anthropophilic mosquitoes result in increased risk of pathogen transmission. Trends Parasitol. 2012;28: 114–121. doi:10.1016/j.pt.2012.01.001

5. Messina JP, Kraemer MU, Brady OJ, Pigott DM, Shearer FM, Weiss DJ, et al. Mapping global environmental suitability for Zika virus. eLife. 2016;5: e15272. doi:10.7554/eLife.15272

6. Magori K, Legros M, Puente ME, Focks DA, Scott TW, Lloyd AL, et al. Skeeter Buster: A stochastic, spatially explicit modeling tool for studying Aedes aegypti population replacement and population suppression strategies. PLOS Negl Trop Dis. 2009;3: e508. doi:10.1371/journal.pntd.0000508

7. Johansson MA, Powers AM, Pesik N, Cohen NJ, Staples JE. Nowcasting the spread of chikungunya virus in the Americas. PLoS ONE. 2014;9: e104915. doi:10.1371/journal.pone.0104915

8. Perkins TA, Metcalf CJE, Grenfell BT, Tatem AJ. Estimating drivers of autochthonous transmission of chikungunya virus in its invasion of the Americas. PLoS Curr. 2015;7. doi:10.1371/currents.outbreaks.a4c7b6ac10e0420b1788c9767946d1fc

9. Morin CW, Monaghan AJ, Hayden MH, Barrera R, Ernst K. Meteorologically driven simulations of dengue epidemics in San Juan, PR. PLoS Negl Trop Dis. 2015;9: e0004002. doi:10.1371/journal.pntd.0004002

10. Zhang Q, Sun K, Chinazzi M, Pastore-Piontti A, Dean NE, Rojas DP, et al. Projected spread of Zika virus in the Americas. bioRxiv. 2016; 066456. doi:10.1101/066456

11. Dell AI, Pawar S, Savage VM. Systematic variation in the temperature dependence of physiological and ecological traits. Proc Natl Acad Sci. 2011;108: 10591–10596.

12. Mordecai EA, Paaijmans KP, Johnson LR, Balzer CH, Ben-Horin T, de Moor E, et al. Optimal temperature for malaria transmission is dramatically lower than previously predicted. Ecol Lett. 2013;16: 22–30.

13. Focks DA, Haile DG, Daniels E, Mount GA. Dynamic life table model for Aedes aegypti (Diptera: Culicidae): analysis of the literature and model development. J Med Entomol. 1993;30: 1003–1017.

14. Yang HM, Macoris MLG, Galvani KC, Andrighetti MTM, Wanderley DMV. Assessing the effects of temperature on the population of Aedes aegypti, the vector of dengue. Epidemiol Infect. 2009;137: 1188–1202. doi:10.1017/S0950268809002040

15. Rueda L, Patel K, Axtell R, Stinner R. Temperature-dependent development and survival rates of Culex quinquefasciatus and Aedes aegypti (Diptera: Culicidae). J Med Entomol. 1990;27: 892–898.

16. Tun-Lin W, Burkot TR, Kay BH. Effects of temperature and larval diet on development rates and survival of the dengue vector Aedes aegypti in north Queensland, Australia. Med Vet Entomol. 2000;14: 31–37. doi:10.1046/j.1365-2915.2000.00207.x

649 17. Kamimura K, Matsuse IT, Takahashi H, Komukai J, Fukuda T, Suzuki K, et al. Effect of temperature on the
650    development of Aedes aegypti and Aedes albopictus. Med Entomol Zool. 2002;53: 53–58.

651 18. Eisen L, Monaghan AJ, Lozano-Fuentes S, Steinhoff DF, Hayden MH, Bieringer PE. The impact of
652    temperature on the bionomics of Aedes (Stegomyia) aegypti, with special reference to the cool geographic
653    range margins. J Med Entomol. 2014;51: 496–516. doi:10.1603/ME13214

654 19. Delatte H, Gimonneau G, Triboire A, Fontenille D. Influence of temperature on immature development,
655    survival, longevity, fecundity, and gonotrophic cycles of Aedes albopictus, vector of chikungunya and dengue
656    in the Indian Ocean. J Med Entomol. 2009;46: 33–41. doi:10.1603/033.046.0105

657 20. Muturi EJ, Lampman R, Costanzo K, Alto BW. Effect of temperature and insecticide stress on life-history traits
658    of Culex restuans and Aedes albopictus (Diptera: Culicidae). J Med Entomol. 2011;48: 243–250.

659 21. Alto BW, Juliano SA. Temperature effects on the dynamics of Aedes albopictus (Diptera: Culicidae)
660    populations in the laboratory. J Med Entomol. 2001;38: 548–556.

661 22. Westbrook CJ, Reiskind MH, Pesko KN, Greene KE, Lounibos LP. Larval environmental temperature and the
662    susceptibility of Aedes albopictus Skuse (Diptera: Culicidae) to chikungunya virus. Vector-Borne Zoonotic
663    Dis. 2010;10: 241–247. doi:10.1089/vbz.2009.0035

664 23. Briegel H, Timmermann SE. Aedes albopictus (Diptera: Culicidae): Physiological aspects of development and
665    reproduction. J Med Entomol. 2001;38: 566–571. doi:10.1603/0022-2585-38.4.566

666 24. Calado DC, Navarro-Silva MA. Influência da temperatura sobre a longevidade, fecundidade e atividade
667    hematofágica de Aedes (Stegomyia) albopictus Skuse, 1894 (Diptera, Culicidae) sob condições de laboratório.
668    Rev Bras Entomol. 2002;46: 93–98. doi:10.1590/S0085-56262002000100011

669 25. Beserra EB, Fernandes CRM, Silva SA de O, Silva LA da, Santos JW dos. Efeitos da temperatura no ciclo de
670    vida, exigências térmicas e estimativas do número de gerações anuais de Aedes aegypti (Diptera, Culicidae).
671    Iheringia Sér Zool. 2009; Available: http://agris.fao.org/agris-search/search.do?recordID=XS2010500501

672 26. Westbrook CJ. Larval ecology and adult vector competence of invasive mosquitoes Aedes albopictus and
673    Aedes aegypti for Chikungunya virus [Internet]. University of Florida. 2010. Available:
674    http://etd.fcla.edu/UF/UFE0041830/westbrook_c.pdf

675 27. Couret J, Dotson E, Benedict MQ. Temperature, Larval Diet, and Density Effects on Development Rate and
676    Survival of Aedes aegypti (Diptera: Culicidae). PLoS ONE. 2014;9: e87468.
677    doi:10.1371/journal.pone.0087468

678 28. Ezeakacha N. Environmental impacts and carry-over effects in complex life cycles: the role of different life
679    history stages. Dissertation. 2015; Available: http://aquila.usm.edu/dissertations/190

680 29. Teng H-J, Apperson CS. Development and Survival of Immature Aedes albopictus and Aedes triseriatus
681    (Diptera: Culicidae) in the Laboratory: Effects of Density, Food, and Competition on Response to
682    Temperature. J Med Entomol. 2000;37: 40–52. doi:10.1603/0022-2585-37.1.40

683 30. Wiwatanaratanabutr S, Kittayapong P. Effects of temephos and temperature on Wolbachia load and life history
684    traits of Aedes albopictus. Med Vet Entomol. 2006;20: 300–307. doi:10.1111/j.1365-2915.2006.00640.x

685 31. Xiao F-Z, Zhang Y, Deng Y-Q, He S, Xie H-G, Zhou X-N, et al. The effect of temperature on the extrinsic
686    incubation period and infection rate of dengue virus serotype 2 infection in Aedes albopictus. Arch Virol.
687    2014;159: 3053–3057. doi:10.1007/s00705-014-2051-1

32. Watts DM, Burke DS, Harrison BA, Whitmire RE, Nisalak A. Effect of temperature on the vector efficiency of
Aedes aegypti for dengue 2 virus. Am J Trop Med Hyg. 1987;36: 143–152.

33. McLean DM, Clarke AM, Coleman JC, Montalbetti CA, Skidmore AG, Walters TE, et al. Vector capability of
Aedes aegypti mosquitoes for California encephalitis and dengue viruses at various temperatures. Can J
Microbiol. 1974;20: 255–262. doi:10.1139/m74-040

34. Carrington LB, Armijos MV, Lambrechts L, Scott TW. Fluctuations at a low mean temperature accelerate
dengue virus transmission by Aedes aegypti. PLoS Negl Trop Dis. 2013;7: e2190.
doi:10.1371/journal.pntd.0002190

35. Davis NC. The effect of various temperatures in modifying the extrinsic incubation period of the yellow fever
virus in Aedes aegypti. Am J Epidemiol. 1932;16: 163–176.

36. McLean DM, Miller MA, Grass PN. Dengue virus transmission by mosquitoes incubated at low temperatures.
Mosq News. 1975; Available: http://agris.fao.org/agris-search/search.do?recordID=US19760088008

37. Focks DA, Daniels E, Haile DG, Keesling JE. A simulation model of the epidemiology of urban dengue fever:
literature analysis, model development, preliminary validation, and samples of simulation results. Am J Trop
Med Hyg. 1995;53: 489–506.

38. Alto BW, Bettinardi D. Temperature and dengue virus infection in mosquitoes: independent effects on the
immature and adult stages. Am J Trop Med Hyg. 2013;88: 497–505. doi:10.4269/ajtmh.12-0421

39. Stewart Ibarra AM, Ryan SJ, Beltrán E, Mejía R, Silva M, Muñoz Á. Dengue vector dynamics (Aedes aegypti)
influenced by climate and social factors in Ecuador: implications for targeted control. PLoS ONE. 2013;8:
e78263. doi:10.1371/journal.pone.0078263

40. Brady OJ, Golding N, Pigott DM, Kraemer MUG, Messina JP, Reiner Jr RC, et al. Global temperature
constraints on Aedes aegypti and Ae. albopictus persistence and competence for dengue virus transmission.
Parasit Vectors. 2014;7: 338. doi:10.1186/1756-3305-7-338

41. Carlson CJ, Dougherty ER, Getz W. An Ecological Assessment of the Pandemic Threat of Zika Virus. PLoS
Negl Trop Dis. 2016;10: e0004968. doi:10.1371/journal.pntd.0004968

42. Samy AM, Thomas SM, Wahed AAE, Cohoon KP, Peterson AT, Samy AM, et al. Mapping the global
geographic potential of Zika virus spread. Mem Inst Oswaldo Cruz. 2016;111: 559–560. doi:10.1590/0074-
02760160149

43. Wesolowski A, Qureshi T, Boni MF, Sundsøy PR, Johansson MA, Rasheed SB, et al. Impact of human
mobility on the emergence of dengue epidemics in Pakistan. Proc Natl Acad Sci. 2015; 201504964.
doi:10.1073/pnas.1504964112

44. Liu-Helmersson J, Stenlund H, Wilder-Smith A, Rocklöv J. Vectorial Capacity of Aedes aegypti: Effects of
Temperature and Implications for Global Dengue Epidemic Potential. PLoS ONE. 2014;9: e89783.
doi:10.1371/journal.pone.0089783

45. Caminade C, Turner J, Metelmann S, Hesson JC, Blagrove MSC, Solomon T, et al. Global risk model for
vector-borne transmission of Zika virus reveals the role of El Niño 2015. Proc Natl Acad Sci. 2017;114: 119–
124. doi:10.1073/pnas.1614303114

46. Bogoch II, Brady OJ, Kraemer MUG, German M, Creatore MI, Kulkarni MA, et al. Anticipating the
international spread of Zika virus from Brazil. The Lancet. 2016;387: 335–336. doi:10.1016/S0140-
6736(16)00080-5

47. Briere J-F, Pracros P, Le Roux A-Y, Pierre J-S. A novel rate model of temperature-dependent development for
arthropods. Environ Entomol. 1999;28: 22–29.

48. Li J, Blakeley D, Smith? RJ. The Failure of R0. Comput Math Methods Med. 2011;2011: e527610.
doi:10.1155/2011/527610

49. Lambrechts L, Paaijmans KP, Fansiri T, Carrington LB, Kramer LD, Thomas MB, et al. Impact of daily
temperature fluctuations on dengue virus transmission by Aedes aegypti. Proc Natl Acad Sci. 2011;108:
7460–7465. doi:10.1073/pnas.1101377108

50. Rohatgi A. WebPlotDigitizer [Internet]. 2015. Available: http://arohatgi.info/WebPlotDigitizer

51. Johnson LR, Ben-Horin T, Lafferty KD, McNally A, Mordecai E, Paaijmans KP, et al. Understanding
uncertainty in temperature effects on vector-borne disease: a Bayesian approach. Ecology. 2015;96: 203–213.
doi:10.1890/13-1964.1

52. R Development Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna,
Austria: R Foundation for Statistical Computing; 2014. Available: http://www.R-project.org

53. Plummer M. rjags: Bayesian Graphical Models using MCMC [Internet]. 2016. Available: http://CRAN.R-
project.org/package=rjags

54. Plummer M, Best N, Cowles K, Vines K. CODA: Convergence Diagnosis and Output Analysis for MCMC.
2006.

55. Parton WJ, Logan JA. A model for diurnal variation in soil and air temperature. Agric Meteorol. 1981;23: 205–
216. doi:10.1016/0002-1571(81)90105-9

56. Paaijmans KP, Heinig RL, Seliga RA, Blanford JI, Blanford S, Murdock CC, et al. Temperature variation
makes ectotherms more sensitive to climate change. Glob Change Biol. 2013;19: 2373–2380.
doi:10.1111/gcb.12240

57. Vasseur DA, DeLong JP, Gilbert B, Greig HS, Harley CDG, McCann KS, et al. Increased temperature variation
poses a greater risk to species than climate warming. Proc R Soc Lond B Biol Sci. 2014;281: 20132612.
doi:10.1098/rspb.2013.2612

58. Narasimhan R. weatherData: Get Weather Data from the Web [Internet]. 2014. Available: https://cran.r-
project.org/web/packages/weatherData/index.html

59. Breheny P, Burchett W. visreg: Visualization of Regression Models [Internet]. 2016. Available: https://cran.r-
project.org/web/packages/visreg/index.html

Figure 1

Figure 2

765   **Figure 3**

Figure 4

A. *Aedes albopictus*   ## B. *Aedes aegypti*