Title: Temperature determines Zika, dengue and chikungunya transmission potential in

the Americas

Authors: Erin A. Mordecai[a*], Jeremy M. Cohen[b], Michelle V. Evans[c], Prithvi Gudapati[a], Leah

R. Johnson[b], Kerri Miazgowicz[d], Courtney C. Murdock[c,d], Jason R. Rohr[b], Sadie J. Ryan[e,f,g,h],

Van Savage[i,j], Marta Shocket[k], Anna Stewart Ibarra[l], Matthew B. Thomas[m], Daniel P. Weikel[n]

Affiliations:

[a]Biology Department, Stanford University, 371 Serra Mall, Stanford, CA 94305

[b]Department of Integrative Biology, University of South Florida, 4202 East Fowler Ave,

SCA110 Tampa, FL 33620

[c]Odum School of Ecology, University of Georgia, 140 E. Green St., Athens, GA 30602

[d]Center for Tropical and Emerging Global Disease, Department of Infectious Diseases,

University of Georgia College of Veterinary Medicine, 501 D.W. Brooks Drive, Athens, GA

30602

[e]Department of Geography, University of Florida, PO Box 117315, Turlington Hall, Gainesville,

FL 32611

[f]Emerging Pathogens Institute, University of Florida, P.O. Box 100009, 2055 Mowry Road

Gainesville, FL 32610

[g]Center for Global Health and Translational Science, Department of Microbiology and

Immunology, Weiskotten Hall, SUNY Upstate Medical University, Syracuse, NY 13210

[h]School of Life Sciences, College of Agriculture, Engineering, and Science, University of

KwaZulu Natal, Private Bag X01, Scottsville, 3209, KwaZulu Natal, South Africa

[i]Department of Ecology and Evolutionary Biology, University of California Los Angeles and

Department of Biomathematics, University of California Los Angeles, Los Angeles, CA 90095

[j]Santa Fe Institute, 1399 Hyde Park Rd, Santa Fe, NM 87501

[k]Department of Biology, Indiana University, 1001 E. 3rd St., Jordan Hall 142, Bloomington, IN

47405

[l]Center for Global Health and Translational Sciences, SUNY Upstate Medical University,

Syracuse, NY13210

[m]Department of Entomology and Center for Infectious Disease Dynamics, Penn State University,

112 Merkle Lab, University Park, PA 16802

[n]Department of Biostatistics, University of Michigan, 1415 Washington Heights, Ann Arbor, MI

48109

[*]Correspondence to: Erin Mordecai. Biology Department, Stanford University, 371 Serra Mall,

Stanford, CA 94305. (650) 497-7447. emordeca@stanford.edu

Classification: BIOLOGICAL SCIENCES: Ecology

Keywords: Zika, dengue, chikungunya, temperature, vector transmission, *Aedes aegypti*, *Aedes*

albopictus

Abstract

Recent epidemics of Zika, dengue, and chikungunya have heightened the need to understand virus transmission ecology for *Aedes aegypti* and *Ae. albopictus* mosquitoes. An estimated 3.9 billion people in 120 countries are at risk for these diseases. Temperature defines the fundamental potential for vector-borne disease transmission, yet the potential for transmission in sub-tropical and temperate regions remains uncertain. Using mechanistic transmission models fit to mosquito and virus physiology data and validated with human case data, we show that mean temperature accurately bounds transmission risk for Zika, chikungunya, and dengue in the Americas. Transmission occurs between 18-34°C and peaks at 29°C for *Ae. aegypti* (between 11-28°C with a peak at 26°C for *Ae. albopictus*). As predicted, high relative incidence of Zika, dengue, and chikungunya in humans occurs between 23-32°C, peaks at 27-29°C, and is very low outside the predicted range. As a proxy for infrastructure and vector control effort, economic reliance on tourism explains some departures from areas otherwise suitable for high rates of transmission. Nonetheless, the temperature-based models alone provide fundamental eco-physiological measures of transmission potential. Tropical and subtropical regions are suitable for extended seasonal or year-round transmission by *Ae. aegypti* and *Ae. albopictus*. By contrast, potential transmission in temperate areas is constrained to at most three months per year even if vectors are present (which is currently not the case for large parts of the US). Such brief transmission windows limit the likelihood and potential extent of epidemics following disease introduction in temperate zones.

Significance statement

Viruses transmitted by *Aedes aegypti* and *Ae. albopictus* mosquitoes, including Zika, dengue, and chikungunya, present one of the most rapidly growing infectious disease threats, putting an

estimated 3.9 billion people in 120 countries at risk. Understanding the relationship between

transmission and climate, particularly temperature, is critical for predicting and responding to

potential spread into sub-tropical and temperate areas. Using models informed by laboratory

experiments and tested against actual human infection data, we show that transmission potential

of these three viruses is substantial between 23-32°C and peaks at 27-29°C. This implies that

while year-round transmission is likely in tropical and sub-tropical areas, temperate areas are at

risk for at most seasonal transmission, given that the necessary mosquito species are present.

Introduction

Epidemics of dengue, chikungunya, and Zika are sweeping through the Americas, part of a public health crisis that places an estimated 3.9 billion people in 120 countries at risk globally (1). Dengue virus (DENV) distribution and intensity in the Americas has increased over the last three decades, infecting an estimated 390 million people (96 million clinical) per year (2). Chikungunya virus (CHIKV) emerged in the Americas in 2013, causing 1.8 million suspected cases from 44 countries and territories (www.paho.org). In the last year, Zika virus (ZIKV) has spread throughout the Americas, causing nearly 500,000 suspected and confirmed cases, with many more unreported (http://ais.paho.org/phip/viz/ed_zika_cases.asp, as of July 7, 2016). The growing burden of these diseases (including links between Zika infection and microcephaly and Guillain-Barré syndrome (3)) and potential for spread into new areas creates an urgent need for predictive models that can inform risk assessment and guide interventions such as mosquito control, community outreach, and education.

Aedes aegypti, a mosquito with a close affiliation with and preference for humans, is the main vector of DENV, CHIKV, and ZIKV, while *Ae. albopictus*, a peri-urban mosquito, is an important secondary vector (4, 5). Predicting transmission of these viruses requires understanding the ecology of the vector species. Mathematical and geostatistical models that incorporate climate information have been immensely valuable for predicting and responding to *Aedes* spp. spread and DENV, CHIKV, and ZIKV outbreaks (5–9). Yet we currently lack a validated mechanistic model of the relationship between key climatic factors and transmission for *Ae. aegypti, Ae. albopictus*, and the arboviruses they transmit. We expect the role of climate, particularly temperature, to be predictable from fundamental ecological responses of mosquitoes and viruses to environmental conditions. Survival, feeding, development, and reproductive rates

predictably respond to temperature across a variety of ectotherms, including mosquitoes (10, 11).

Because these traits determine transmission rates, the effects of temperature on transmission

should also be broadly predictable from mechanistic models that incorporate temperature-

dependent traits.

Here, we synthesize available data to characterize the temperature-dependent traits of the

mosquitoes and viruses that determine transmission intensity. With these thermal responses, we

develop mechanistic temperature-dependent virus transmission models for *Ae. aegypti* and *Ae.*

albopictus, and validate them with DENV, CHIKV, and ZIKV human incidence data from the

Americas from 2014-2016. Because the relationship between transmission and incidence is

nonlinear, we better approximated transmission by dividing incidence by the cumulative number

of cases. We confined our analyses to the increasing portion of each epidemic, before susceptible

individuals become limiting and when transmission is most dependent on environmental

conditions. In validating the temperature-dependent mechanistic models, we examine the rates of

false positive and false negative predictions. If temperature fundamentally limits transmission

potential, high transmission should only occur under temperatures predicted to be highly

suitable, and areas with low predicted suitability should have low transmission (i.e., false

negative rates should be low). By contrast, low transmission may occur even when temperature

suitability is high because other factors like vector control can limit transmission (i.e., the false

positive rate should be higher than the false negative rate).

Results

Data gathered from the literature (9, 12–23) revealed that all mosquito traits relevant to

transmission peak between 23°C and 34°C for the two mosquito species (Figs. S1-S2). Humidity

linearly increases survival at all mosquito life stages, but does not interact with temperature (Fig.

S3). DENV extrinsic incubation and vector competence peak at 35°C (24–27) and 31-32°C (24,

25, 27, 28), respectively, in both mosquitoes—temperatures at which mosquito survival is low,

limiting transmission potential (Figs. S1-S2). The parameterized models predict optimal

transmission at 29°C by *Ae. aegypti* and 26°C by *Ae. albopictus* (Fig. 1). Transmission potential

is zero below 14-18°C or above 34-35°C for *Ae. aegypti*, and below 11-16°C or above 28-32°C

for *Ae. albopictus* (ranges depend on the amount of daily temperature variation; Figs. 1, S4-S5).

Weekly relative incidence of DENV, CHIKV, and ZIKV across countries in the

Americas and the Caribbean is consistent with mechanistic model predictions (Fig. 1). The

highest observed incidence occurs at 27°C for CHIKV and DENV and at 29°C for ZIKV, and the

majority of transmission occurs between 23-32°C (i.e., false negative rate was low; Fig. 1). At a

local scale, DENV relative incidence reported across 14 years in Iquitos, Peru and 24 years in

San Juan, Puerto Rico also matches the predicted thermal optimum (Fig. S6). The accuracy of

the model in bounding transmission potential is notable given the coarse scale of the human

incidence versus mean temperature data (i.e., country-scale means for the DENV and CHIKV),

the lack of CHIKV- and ZIKV-specific trait thermal response data to inform the model, the

nonlinear relationship between transmission and incidence, and all the well-documented factors

other than temperature that influence transmission. Together, these analyses show that

temperature suitability for transmission is predictable from simple mechanistic models

parameterized with laboratory data on the two mosquito species and dengue virus. Moreover, the

similar responses of human incidence of ZIKV, CHIKV, and DENV to temperature suggest that

the thermal ecology of their shared mosquito vectors is a key determinant of outbreak location,

timing, and size.

High transmission only occurred where predicted by the model (Fig. 1) but as expected,

some sites with high potential for transmission had low actual transmission (i.e., false positives

far exceeded false negatives). Socio-ecological factors such as vector control effort might

explain these false positives. For example, many countries in Latin America and the Caribbean

economically depend on tourism, and are likely to invest heavily in preventing outbreaks. We

therefore expect these countries to have lower transmission than predicted by the model. Indeed,

while the temperature-dependent model predicted maximum incidence well for data from

countries with low to medium dependence on tourism, countries with the heaviest reliance on

tourism had no correlation between expected and observed transmission (10^{th}, 50^{th}, and 90^{th}

percentile, respectively, in proportion of GDP that comes from tourism; Fig. 2). In sum, mean

temperature alone predicts where transmission and large epidemics can and cannot occur, while

factors correlated with economic reliance on tourism explain lower than expected transmission in

some sites with highly suitable temperatures.

The validated model predicts where transmission is not excluded (a conservative estimate

of transmission risk). Considering the number of months per year at which mean temperatures do

not prevent transmission (i.e., $R_0 > 0$; since our model is relativized, we cannot estimate where

absolute $R_0 > 1$), large areas of tropical and subtropical regions, including Puerto Rico and parts

of Florida and Texas, are currently suitable year-round or seasonally (Fig. 3). These regions are

fundamentally at risk for DENV, CHIKV, ZIKV, and other *Aedes* arbovirus transmission during

a substantial part of the year (Fig. 3). On the other hand, many temperate regions experience

temperatures suitable for transmission 3 months or less per year (Fig. 3), and the virus incubation

periods in humans and mosquitoes restrict the transmission window even further. Temperature

thus limits the potential for the viruses to generate extensive epidemics in temperate areas even

where the vectors are present. Moreover, many temperate regions with seasonally suitable

temperatures currently lack *Ae. aegypti* and *Ae. albopictus* mosquitoes, making vector

transmission impossible (Fig. S7).

Discussion

Predicting the geographic and seasonal potential for transmission of Zika is particularly

important in light of the upcoming Olympic games in Brazil in August 2016. Over the last 10

years, the mean temperature in Rio de Janeiro in August has been 22.2°C, which is at the low

end of the predicted suitable range for transmission (Fig. 1, lines). Observed transmission rates

of ZIKV, DENV, and CHIKV have been low at this temperature (Fig. 1, points). The

temperature suitability for transmission during the Olympics is relatively low compared with the

Brazilian summer (e.g., during Carnival in February, when the 10-year mean temperature is

27.6°C, nearly optimal for transmission). Moreover, transmission risk during the 2016 Olympics

also depends on mosquito abundance and biting rate, control efforts, infrastructure, and the

number of susceptible visitors. Together, low temperature suitability, vector control efforts, and

infrastructure improvements may limit ZIKV transmission risk during the Rio Olympics. The

possibility of new epidemics arising from cases exported from the 2016 Olympics is an

additional risk. Regions with longer transmission seasons (e.g., >3 months) could have suitable

temperatures beyond the August 21, 2016 closing ceremony (e.g., the Southeastern US), but

much of the temperate northern hemisphere is expected to have at most a short transmission

window following the Olympics (Fig. S7).

The socio-ecological conditions that enabled CHIKV, ZIKV, and DENV to become the

three most important emerging vector-borne diseases in the Americas make further *Aedes*-

transmitted virus emergence likely (e.g., Rift Valley fever virus, yellow fever virus, Uganda S

virus, Ross River Virus). Efforts to extrapolate and to map temperature suitability (Fig. 3) will be

critical for improving management of both ongoing and future emerging epidemics. Whereas

statistical models are well suited for accurate interpolation from existing distribution ranges (5),

mechanistic models are useful for extrapolating beyond current distributions. These mechanistic

models are especially important for effects of temperature because thermal responses are well

known to be nonlinear (Figs. S1-S2) and because thermal responses of multiple traits must be

integrated to accurately predict temperature suitability for transmission and thus to extrapolate

beyond the current distribution of the viruses. Strikingly, we show that our model based just on

temperature is able to bound human incidence for DENV, CHIKV, and ZIKV (Fig. 1). Socio-

ecological factors such as vector control effort and infrastructure (for which the proportion of

GDP in tourism might be a proxy) explain additional variation in transmission in suitable areas

(Fig. 2). Accurately estimating temperature-driven transmission risk in both highly suitable and

marginal regions is critical for predicting and responding to future outbreaks of these and other

Aedes-transmitted viruses.

Materials and Methods

<u>Temperature-sensitive R_0 models</u>

We constructed temperature-dependent models of transmission using a previously

developed R_0 framework. We modeled transmission rate as the basic reproduction rate, R_0—the

number of secondary infections that would originate from a single infected individual introduced

to a fully susceptible population. In previous work on malaria, we adapted a commonly used

expression for R_0 for vector transmission to include the temperature-sensitive traits that drive

mosquito population density (11):

$$207 \quad R_0(T) = \left(\frac{a(T)^2\, b(T)\, c(T)\, e^{-\mu(T)/PDR(T)}\, EFD(T)\, p_{EA}(T)\, MDR(T)}{N\, r\, \mu(T)^3} \right)^{1/2} \quad\quad (1)$$

Here, (*T*) indicates that the trait is a function of temperature, *T*; *a* is the per-mosquito biting rate,

b is the proportion of infectious bites that infect susceptible humans, *c* is the proportion of bites

on infected humans that infect previously uninfected mosquitoes (i.e., *b*c* = vector competence),

μ is the adult mosquito mortality rate (lifespan, *lf* = *1/μ*), *PDR* is the parasite development rate

(i.e., the inverse of the extrinsic incubation period, the time required between a mosquito biting

an infected host and becoming infectious), *EFD* is the number of eggs produced per female

mosquito per day, p_{EA} is the mosquito egg-to-adult survival probability, *MDR* is the mosquito

immature development rate (i.e., the inverse of the egg-to-adult development time), *N* is the

density of humans, and *r* is the human recovery rate. For each temperature-sensitive trait in each

mosquito species, we fit either symmetric (Quadratic, $c(T - T_0)(T - T_m)$) or asymmetric (Brière,

$cT(T - T_0)(T_m - T)^{1/2}$) unimodal thermal response models to the available empirical data (29). In

both functions, T_0 and T_m are respectively the minimum and maximum temperature for

transmission, and *c* is a rate constant. We normalized the R_0 equation to take values between 0-1

because absolute values of R_0 depend on additional factors not captured in our model. Therefore,

$R_0 > 0$ is an absolute threshold for whether or not transmission is possible, but the model does

not predict when transmission is stable (i.e., absolute $R_0 > 1$).

We fit the trait thermal responses in equation (1) based on an exhaustive search of

published laboratory studies that fulfilled the criterion of measuring a trait at three or more

constant temperatures, ideally capturing both the rise and the fall of each unimodal curve (Tables

S1-S2). Constant-temperature laboratory conditions are required to isolate the direct effect of

temperature from confounding factors in the field and to provide a baseline for estimating the

effects of temperature variation through rate summation (30). We attempted to obtain raw data

from each study, but if they were not available we collected data by hand from tables or digitized

data from figures using WebPlotDigitizer (31). We obtained raw data from Delatte (18) and Alto

(20) for the *Ae. albopictus* egg-to-adult survival probability (*pEA*), mosquito development rate

(MDR), gonotrophic cycle duration (GCD) and total fecundity (TFD) (Table S2). Data did not

meet the inclusion criterion for CHIKV or ZIKV extrinsic incubation period (EIP) in either *Ae.*

albopictus or *Ae. aegypti*. Instead, we used DENV EIP data, combined with sensitivity analyses.

Following Johnson *et al.* (32), we fit a thermal response for each trait using Bayesian

models. We first fit Bayesian models for each trait thermal response using uninformative priors

($T_0 \sim$ Uniform (0, 24), $T_m \sim$ Uniform (25, 45), $c \sim$ Gamma (1, 10) for Brière and $c \sim$ Gamma (1,

1) for Quadratic fits) chosen to restrict each parameter to its biologically realistic range (i.e., $T_0 <$

T_m and we assumed that temperatures below 0°C and above 45°C were lethal). Any negative

values for all thermal response functions were truncated at zero, and thermal responses for

probabilities (p_{EA}, *b,* and *c*) were also truncated at one. We modeled the observed data as arising

from a normal distribution with the mean predicted by the thermal response function calculated

at the observed temperature, and the precision τ, ($\tau = 1/\sigma$), distributed as $\tau \sim$ Gamma (0.0001,

00001). We fit the models using Markov Chain Monte Carlo (MCMC) sampling in JAGS, using

the R (33) package *rjags* (34). For each thermal response, we ran five MCMC chains with a

5000-step burn-in and saved the subsequent 5000 steps. We thinned the posterior samples by

saving every fifth sample and used the samples to calculate R_0 from 15-35°C, producing a

posterior distribution of R_0 versus temperature. We summarized the relationship between

temperature and each trait or overall R_0 by calculating the mean and 95% highest posterior

density interval (HPD interval; a type of credible interval that includes the smallest continuous

range containing 95% of the probability, as implemented in the *coda* package (35)) for each

curve across temperatures.

We fit a second set of models for each mosquito species that used informative priors, for

two purposes: 1) to reduce uncertainty in R_0 versus temperature and in the trait thermal

responses, and 2) to ensure that our results were not overly dependent on the particular set of

data available. In these models, we used Gamma-distributed priors for each parameter T_0, T_m, c,

and τ fit from an additional 'prior' dataset of *Aedes* spp. trait data that did not meet the inclusion

criteria for the primary dataset (Table S3). We found that these initial informative priors could

have an overly strong influence on the posteriors, in some cases drawing the posterior

distributions well away from the primary dataset, which was better controlled and met the

inclusion criteria. We accounted for our lower confidence in this data set by increasing the

variance in the informative priors, by multiplying all hyperparameters (i.e., the parameters of the

Gamma distributions of priors for T_0, T_m, and c) by a constant k to produce a distribution with the

same mean but 1/k times larger variance. We chose the value of k based on our relative

confidence in the prior versus main data. Thus we chose k = 0.5 for *b, c,* and *PDR* and k = 0.01

for *lf*. This is the main model presented in the text. It is comparable to some but not all previous

mechanistic models for *Ae. aegypti* and *Ae. albopictus* transmission (Fig. S8). Results of our

main model, fit with informative priors, did not vary substantially from the model fit with

uninformative priors (Figs. S9-S10).

<u>Effects of humidity on dengue R_0</u>

Like temperature, humidity is expected to affect vector transmission via its effects on

mosquito populations. Specifically, we expected humidity to affect egg-to-adult survival and

adult survival. Using the methods described above, we extracted laboratory data on these traits at

constant humidity in the laboratory. We searched for *Aedes* spp. survival probability data across

temperature and humidity. We obtained data on *Aedes aegypti* egg hatching probability (36) and

Anopheles gambiae lifespan (37), which we used in absence of *Aedes* spp. because they have a

similar (tropical, anthrophilic) life history. We used linear regression to estimate the relationship

between these traits and relative humidity (%), plugged them into the $R_0(T)$ equation (assuming

the probability of egg hatching and egg-to-adult survival have the same relationship with

humidity), and plotted R_0 versus temperature across a range of relative humidity values (Fig. S3).

For the dataset that included variation in temperature and humidity independently, there was no

interaction between the unimodal temperature response and the linear humidity response (i.e.,

humidity did not affect the relationship between lifespan and temperature) (37). The resulting

relationship between R_0 and relative humidity was exponential.

Incorporating daily temperature variation in transmission models

Because organisms do not typically experience constant temperature environments in the

field, we incorporated the effects of temperature variation on transmission by calculating a daily

average R_0 assuming a daily temperature range of 8°C, across a range of mean temperatures.

This range is consistent with daily temperature variation in tropical and subtropical environments

but lower than in most temperate environments. At each mean temperature, we used a Parton-

Logan model to generate hourly temperatures and calculate each temperature-sensitive trait on an

hourly basis (38). We assumed an irreversible high-temperature threshold above which

mosquitoes die and transmission is impossible (39, 40). We set this threshold based on hourly

temperatures exceeding the critical thermal maximum (T_m in Tables S1-S2) for egg-to-adult

survival or adult longevity by any amount for five hours or by 3°C for one hour. We averaged

each trait over 24 hours to obtain a daily average trait value, which we used to calculate relative

R_0 across a range of mean temperatures. We used this model to predict human relative incidence

(Fig. 1).

DENV, CHIKV, and ZIKV incidence data

To validate our model, we plotted the rate of new human cases of DENV, CHIKV, and

ZIKV against mean temperature during the transmission window. We downloaded and manually

entered Pan American Health Organization (PAHO) weekly case reports for DENV and CHIKV

for all countries in the Americas (North, Central, and South America and the Caribbean Islands)

from week 1 of 2014 to week 8 of 2015 for CHIKV and from week 52 of 2013 to week 47 of

2015 for DENV (www.paho.org). ZIKV weekly case reports for reporting districts (e.g.,

provinces) within Colombia, Mexico, El Salvador, and the US Virgin Islands were available

from the CDC Epidemic Prediction Initiative (https://github.com/cdcepi/) from November 28,

2015 to March 1, 2016. Additional serotype-specific DENV case data (matched with temperature

and other covariates) were available from the Dengue Forecasting Challenge

(www.dengueforecasting.noaa.gov/) for Iquitos, Peru from 2000-2013 and for San Juan, Puerto

Rico from 1990-2013. We manually selected separate epidemics of different DENV serotypes by

examining the weekly incidence data plotted over time.

Incidence of human cases has a nonlinear relationship with transmission potential (R_0 or

the number of infectious mosquitoes per person). During the rising phase of an epidemic, new

cases arise more quickly as the epidemic gets larger, while during the declining phase, the

exhaustion of the susceptible population leads to declines in transmission (41). We controlled for

this nonlinearity in two ways. First, we calculated relative incidence as the number of new cases

divided by the total number of cases to date. Second, we removed the declining portion of each

epidemic from analyses. We did this by discarding all data following the first week that relative

incidence was less than the average over the preceding ten weeks.

Temperature data collection

We matched the PAHO DENV and CHIKV incidence data with temperature using daily

temperature data from METAR stations in each country, averaged at the country level by

epidemic week. We assumed a two-week lag between temperature and incidence (i.e., mean

temperature for the week that is two weeks prior to each case report). METAR stations are

internationally standardized weather reporting stations that report hourly temperature and

precipitation measures. Outlier weather stations were excluded if they reported a daily maximum

temperature below 5°C or a daily minimum temperature below 40°C during the study period,

extremes that would certainly eliminate the potential for transmission in a local area. Because

case data are reported at the country level, we needed a collection of weather stations in each

country that accurately represent weather conditions in the areas where transmission occurs,

excluding extreme areas where transmission is unlikely. For the study period of October 1, 2013

through January 10, 2016, we downloaded daily temperature data for each station from Weather

Underground using the *weatherData* package in R (42). This totaled 523 stations for 49

countries. We removed all data from Chile because it spans so much latitude and the terrain is so

diverse that its country-level mean is unlikely to be very representative of the temperature where

an outbreak occurred. For ZIKV, we manually selected weather stations in or near each reporting

area (department within country). El Salvador and the US Virgin Islands each had only two

weather stations, which we used for all reporting locations within those countries.

Validation analyses with human incidence versus temperature datasets

To validate our model predictions for transmission, we plotted each time point for

relative incidence against mean temperature from two weeks earlier for DENV, CHIKV, and

ZIKV. Within a set of reports with similar temperatures, we expected the sites with the highest

relative incidence to represent those where transmission is least limited by vector control and

other socio-ecological factors. We selected these maxima within a set of temperature bins for

each incidence dataset (PAHO DENV, CHIKV, and ZIKV). To determine the size of each bin,

we took the subset of each dataset with non-zero relative incidence and created bins of equal size

for a total of 20 bins. These maxima are highlighted with filled circles in Fig. 1.

We were then interested whether relative incidence maxima would be highest at the most

suitable temperatures predicted by the model (~26-29°C), and lower at temperatures predicted to

be less suitable. To examine this relationship, we plotted predicted transmission rate against

observed relative incidence for the maximum points only, using the best-fit prediction from

either the *Ae. aegypti* or the *Ae. albopictus* R_0 model because the mosquito vector responsible for

transmission often is not known (5).

Even for the maximum relative incidence at each temperature, we expect some variation

to be explained by socio-ecological conditions beyond temperature. In particular, countries that

rely heavily on tourism should be highly motivated to prevent epidemics, and thus the proportion

of GDP in tourism might be a good proxy for vector control effort and infrastructure that

prevents human-mosquito contact (air conditioning and window screens). Based on this

motivation, we conducted a linear mixed-effects model with observed maximum incidence data

(from the 20 bins for each virus) as the Gaussian response variable, country as a random effect,

and virus species, predicted R_0 from the mechanistic model, log of percent of GDP in tourism,

and a predicted R_0-by-GDP interaction as predictors. We plotted the expected versus observed

transmission relationship based on whether the outbreak origin was in the 10[th], 50[th], or 90[th]

percentile of countries in percent of GDP in tourism, using the R package *visreg* (43). Each point

in this partial residual plot (Fig. 2) appears only once in the panel to which it is closest. However,

the least squares estimates from the mixed model are not equivalent to drawing a line through the

partial residuals because they draw on the pooled data that are shown in separate panels (43).

Mapping temperature suitability for transmission

Using our validated model, we were interested in where the temperature was suitable for

Ae. aegypti and/or *Ae. albopictus* transmission for some or all of the year to predict the potential

geographic range of outbreaks in the Americas. We visualized the minimum, median, and

maximum extent of transmission based on probability of occurrence thresholds from the R_0

models for both mosquitoes. We calculated the number of consecutive months in which the

posterior probability of $R_0 > 0$ exceeds a threshold of 0.025, 0.5, or 0.975 for both mosquito

species, representing the maximum, median, and minimum likely ranges, respectively. This

analysis indicates the predicted seasonality of temperature suitability for transmission

geographically, but does not indicate its magnitude. To generate the maps, we cropped monthly

mean temperature rasters for all twelve months (Worldclim; www.worldclim.org/) to the

Americas (*R*, *raster* package, *crop* function) and assigned cells values of 1 or 0 depending on

whether the probability that $R_0 > 0$ exceeded the threshold at the temperatures in those cells

(Table S3). We then synthesized the monthly grids into a single raster that reflected the

maximum number of consecutive months where cell values equaled 1. For plotting, rasters were

overlaid on a global country polygon map (*rworldmap* package (44)). We repeated this process

for each combination of mosquito species and posterior probability thresholds.

Acknowledgments

EAM, MBT, VS, SJR, LRJ, ASI, JRR, MS, JC, and DPW were supported by the National Science Foundation (DEB-1518681). J.R.R. was supported by the NSF (EF-1241889), National Institutes of Health (R01GM109499 and R01TW010286-01), US Department of Agriculture (2009-35102-0543) and US Environmental Protection Agency (CAREER 83518801). EAM and CCM were supported by the NSF (DEB-1640780). Barry Alto, Krijn Paaijmans, Francis Ezeakacha, and Helene Delatte kindly provided raw data used in the analyses. We gratefully acknowledge the Centers for Disease Control and Prevention Epidemic Predictions Initiative (CDC EPI) for collating and sharing the Zika incidence data on GitHub (https://zenodo.org/record/48946#.Vz-EM2bb8ys). The authors declare no competing interests.

References:

1. Brady OJ, et al. (2012) Refining the global spatial limits of dengue virus transmission by evidence-based consensus. *PLOS Negl Trop Dis* 6(8):e1760.

2. Bhatt S, et al. (2013) The global distribution and burden of dengue. *Nature* 496(7446):504–507.

3. Rasmussen SA, Jamieson DJ, Honein MA, Petersen LR (2016) Zika virus and birth defects — reviewing the evidence for causality. *N Engl J Med* 374:1981–1987.

4. Scott TW, Takken W (2012) Feeding strategies of anthropophilic mosquitoes result in increased risk of pathogen transmission. *Trends Parasitol* 28(3):114–121.

5. Messina JP, et al. (2016) Mapping global environmental suitability for Zika virus. *eLife* 5:e15272.

6. Magori K, et al. (2009) Skeeter Buster: A stochastic, spatially explicit modeling tool for studying *Aedes aegypti* population replacement and population suppression strategies. *PLOS Negl Trop Dis* 3(9):e508.

7. Johansson MA, Powers AM, Pesik N, Cohen NJ, Staples JE (2014) Nowcasting the spread of chikungunya virus in the Americas. *PLoS ONE* 9(8):e104915.

8. Perkins TA, Metcalf CJE, Grenfell BT, Tatem AJ (2015) Estimating drivers of autochthonous transmission of chikungunya virus in its invasion of the Americas. *PLoS Curr* 7. doi:10.1371/currents.outbreaks.a4c7b6ac10e0420b1788c9767946d1fc.

9.   Morin CW, Monaghan AJ, Hayden MH, Barrera R, Ernst K (2015) Meteorologically driven

simulations of dengue epidemics in San Juan, PR. *PLoS Negl Trop Dis* 9(8):e0004002.

10. Dell AI, Pawar S, Savage VM (2011) Systematic variation in the temperature dependence

of physiological and ecological traits. *Proc Natl Acad Sci* 108(26):10591–10596.

11. Mordecai EA, et al. (2013) Optimal temperature for malaria transmission is dramatically

lower than previously predicted. *Ecol Lett* 16:22–30.

12. Focks DA, Haile DG, Daniels E, Mount GA (1993) Dynamic life table model for *Aedes*

aegypti (Diptera: Culicidae): analysis of the literature and model development. *J Med*

Entomol 30(6):1003–1017.

13. Yang HM, Macoris MLG, Galvani KC, Andrighetti MTM, Wanderley DMV (2009)

Assessing the effects of temperature on the population of *Aedes aegypti*, the vector of

dengue. *Epidemiol Infect* 137(8):1188–1202.

14. Rueda L, Patel K, Axtell R, Stinner R (1990) Temperature-dependent development and

survival rates of *Culex quinquefasciatus* and *Aedes aegypti* (Diptera: Culicidae). *J Med*

Entomol 27(5):892–898.

15. Tun-Lin W, Burkot TR, Kay BH (2000) Effects of temperature and larval diet on

development rates and survival of the dengue vector *Aedes aegypti* in north Queensland,

Australia. *Med Vet Entomol* 14(1):31–37.

16. Kamimura K, et al. (2002) Effect of temperature on the development of *Aedes aegypti* and

Aedes albopictus. *Med Entomol Zool* 53(1):53–58.

17.   Eisen L, et al. (2014) The impact of temperature on the bionomics of *Aedes (Stegomyia)*

aegypti, with special reference to the cool geographic range margins. *J Med Entomol*

51(3):496–516.

18.   Delatte H, Gimonneau G, Triboire A, Fontenille D (2009) Influence of temperature on

immature development, survival, longevity, fecundity, and gonotrophic cycles of *Aedes*

albopictus, vector of chikungunya and dengue in the Indian Ocean. *J Med Entomol*

46(1):33–41.

19.   Muturi EJ, Lampman R, Costanzo K, Alto BW (2011) Effect of temperature and insecticide

stress on life-history traits of *Culex restuans* and *Aedes albopictus* (Diptera: Culicidae). *J*

Med Entomol 48(2):243–250.

20.   Alto BW, Juliano SA (2001) Temperature effects on the dynamics of *Aedes albopictus*

(Diptera: Culicidae) populations in the laboratory. *J Med Entomol* 38(4):548–556.

21.   Westbrook CJ, Reiskind MH, Pesko KN, Greene KE, Lounibos LP (2010) Larval

environmental temperature and the susceptibility of *Aedes albopictus* Skuse (Diptera:

Culicidae) to chikungunya virus. *Vector-Borne Zoonotic Dis* 10(3):241–247.

22.   Briegel H, Timmermann SE (2001) *Aedes albopictus* (Diptera: Culicidae): Physiological

aspects of development and reproduction. *J Med Entomol* 38(4):566–571.

23.   Calado DC, Navarro-Silva MA (2002) Influência da temperatura sobre a longevidade,

fecundidade e atividade hematofágica de *Aedes (Stegomyia) albopictus* Skuse, 1894

(Diptera, Culicidae) sob condições de laboratório. *Rev Bras Entomol* 46(1):93–98.

24. Xiao F-Z, et al. (2014) The effect of temperature on the extrinsic incubation period and

infection rate of dengue virus serotype 2 infection in *Aedes albopictus*. *Arch Virol*

159(11):3053–3057.

25. Watts DM, Burke DS, Harrison BA, Whitmire RE, Nisalak A (1987) Effect of temperature

on the vector efficiency of *Aedes aegypti* for dengue 2 virus. *Am J Trop Med Hyg*

36(1):143–152.

26. McLean DM, et al. (1974) Vector capability of *Aedes aegypti* mosquitoes for California

encephalitis and dengue viruses at various temperatures. *Can J Microbiol* 20(2):255–262.

27. Carrington LB, Armijos MV, Lambrechts L, Scott TW (2013) Fluctuations at a low mean

temperature accelerate dengue virus transmission by *Aedes aegypti*. *PLoS Negl Trop Dis*

7(4):e2190.

28. Alto BW, Bettinardi D (2013) Temperature and dengue virus infection in mosquitoes:

independent effects on the immature and adult stages. *Am J Trop Med Hyg* 88(3):497–505.

29. Briere J-F, Pracros P, Le Roux A-Y, Pierre J-S (1999) A novel rate model of temperature-

dependent development for arthropods. *Environ Entomol* 28(1):22–29.

30. Lambrechts L, et al. (2011) Impact of daily temperature fluctuations on dengue virus

transmission by Aedes aegypti. *Proc Natl Acad Sci* 108(18):7460–7465.

31. Rohatgi A (2015) *WebPlotDigitizer* Available at: http://arohatgi.info/WebPlotDigitizer.

32. Johnson LR, et al. (2015) Understanding uncertainty in temperature effects on vector-borne

disease: a Bayesian approach. *Ecology* 96(1):203–213.

33.   R Development Core Team (2014) *R: A Language and Environment for Statistical*

Computing (R Foundation for Statistical Computing, Vienna, Austria) Available at:

http://www.R-project.org.

34.   Plummer M (2016) *rjags: Bayesian Graphical Models using MCMC* Available at:

http://CRAN.R-project.org/package=rjags.

35.   Plummer M, Best N, Cowles K, Vines K (2006) *CODA: Convergence Diagnosis and*

Output Analysis for MCMC.

36.   Dickerson CZ (2007) The effects of temperature and humidity on the eggs of *Aedes aegypti*

(L.) and *Aedes albopictus* (skuse) in Texas. Doctoral (Texas A & M University, College

Station, Texas).

37.   Bayoh MN (2001) Studies on the development and survival of *Anopheles gambiae* sensu

stricto at various temperatures and relative humidities. Doctoral (Durham University).

Available at: http://etheses.dur.ac.uk/4952/ [Accessed October 15, 2014].

38.   Parton WJ, Logan JA (1981) A model for diurnal variation in soil and air temperature.

Agric Meteorol 23:205–216.

39.   Paaijmans KP, et al. (2013) Temperature variation makes ectotherms more sensitive to

climate change. *Glob Change Biol* 19(8):2373–2380.

40.   Vasseur DA, et al. (2014) Increased temperature variation poses a greater risk to species

than climate warming. *Proc R Soc Lond B Biol Sci* 281(1779):20132612.

41.   Keeling MJ, Rohani P (2008) *Modeling Infectious Diseases in Humans and Animals*

501   (Princeton University Press).

502   42.   Narasimhan R (2014) *weatherData: Get Weather Data from the Web* Available at:

503   https://cran.r-project.org/web/packages/weatherData/index.html [Accessed May 25, 2016].

504   43.   Breheny P, Burchett W (2016) *visreg: Visualization of Regression Models* Available at:

505   https://cran.r-project.org/web/packages/visreg/index.html [Accessed June 8, 2016].

506   44.   South A (2016) *rworldmap: Mapping Global Data* Available at: https://cran.r-

507   project.org/web/packages/rworldmap/index.html [Accessed June 10, 2016].

508   45.   Wesolowski A, et al. (2015) Impact of human mobility on the emergence of dengue

509   epidemics in Pakistan. *Proc Natl Acad Sci*:201504964.

510   46.   Liu-Helmersson J, Stenlund H, Wilder-Smith A, Rocklöv J (2014) Vectorial capacity of

511   *Aedes aegypti*: Effects of temperature and implications for global dengue epidemic

512   potential. *PLoS ONE* 9(3):e89783.

513   47.   Focks DA, Haile DG, Daniels E, Mount GA (1993) Dynamic life table model for *Aedes

514   aegypti* (Diptera: Culicidae): analysis of the literature and model development. *J Med

515   Entomol* 30(6):1003–1017.

516   48.   Ezeakacha N (2015) Environmental impacts and carry-over effects in complex life cycles:

517   the role of different life history stages. *Dissertation*. Available at:

518   http://aquila.usm.edu/dissertations/190.

519   49.   Beserra EB, Fernandes CRM, Silva SA de O, Silva LA da, Santos JW dos (2009) Efeitos da

520   temperatura no ciclo de vida, exigências térmicas e estimativas do número de gerações

521   anuais de *Aedes aegypti* (Diptera, Culicidae). *Iheringia Sér Zool*. Available at:

522   http://agris.fao.org/agris-search/search.do?recordID=XS2010500501 [Accessed September

523   10, 2015].

524   50.   Joshi DS (1996) Effect of fluctuating and constant temperatures on development, adult

525   longevity and fecundity in the mosquito *Aedes krombeini*. *J Therm Biol* 21(3):151–154.

526   51.   Westbrook CJ (2010) Larval ecology and adult vector competence of invasive mosquitoes

527   *Aedes albopictus* and *Aedes aegypti* for Chikungunya virus. Dissertation (University of

528   Florida). Available at: http://etd.fcla.edu/UF/UFE0041830/westbrook_c.pdf [Accessed

529   October 23, 2013].

530   52.   Wiwatanaratanabutr S, Kittayapong P (2006) Effects of temephos and temperature on

531   Wolbachia load and life history traits of *Aedes albopictus*. *Med Vet Entomol* 20(3):300–

532   307.

533   53.   Teng H-J, Apperson CS (2000) Development and Survival of Immature *Aedes albopictus*

534   and *Aedes triseriatus* (Diptera: Culicidae) in the Laboratory: Effects of Density, Food, and

535   Competition on Response to Temperature. *J Med Entomol* 37(1):40–52.

536   54.   Tun-Lin W, Burkot T, Kay B (2000) Effects of temperature and larval diet on development

537   rates and survival of the dengue vector *Aedes aegypti* in north Queensland, Australia. *Med

538   Vet Entomol* 14(1):31–37.

55. Jalil M (1972) Effect of temperature on larval growth of *Aedes triseriatus*. *J Econ Entomol*

65(2):625–626.

56. Calado DC, Silva MAN da (2002) Avaliaçäo da influência da temperatura sobre o

desenvolvimento de *Aedes albopictus*. *Rev Saúde Pública* 36(2):173–179.

57. Couret J, Dotson E, Benedict MQ (2014) Temperature, larval diet, and density effects on

development rate and survival of *Aedes aegypti* (Diptera: Culicidae). *PLoS ONE*

9(2):e87468.

58. Davis NC (1932) The effect of various temperatures in modifying the extrinsic incubation

period of the yellow fever virus in *Aedes aegypti*. *Am J Epidemiol* 16(1):163–176.

59. Focks DA, Daniels E, Haile DG, Keesling JE (1995) A simulation model of the

epidemiology of urban dengue fever: literature analysis, model development, preliminary

validation, and samples of simulation results. *Am J Trop Med Hyg* 53(5):489–506.

60. McLean DM, Miller MA, Grass PN (1975) Dengue virus transmission by mosquitoes

incubated at low temperatures. *Mosq News*. Available at: http://agris.fao.org/agris-

search/search.do?recordID=US19760088008 [Accessed August 26, 2015].

61. Reisen WK, Fang Y, Martinez VM (2006) Effects of temperature on the transmission of

West Nile virus by *Culex tarsalis* (Diptera: Culicidae). *J Med Entomol* 43(2):309–317.

Figure Captions

Fig. 1. Relative R_0 (assuming an 8°C daily temperature range) versus mean temperature (°C) for

Ae. albopictus (light blue line) and *Ae. aegypti* (dark blue line). For validation, weekly relative

incidence of DENV (dark blue points, N = 1,195), CHIKV (gray points, N = 1,642), and ZIKV

(red points, N = 10,741) are plotted against the mean temperature two weeks prior to each

reporting date. Filled circles represent the maximum relative incidence within each temperature

bin for each virus (20 bins per virus). Both R_0 curves and the relative incidence of the three

viruses are each normalized to a 0-1 scale for ease of comparison and visualization.

Fig. 2. A partial residual plot showing the relationship between the predicted transmission rate

(relative R_0) from the mechanistic model and the observed maximum relative incidence (rate of

new cases divided by cumulative cases; filled circles in Fig. 1, N = 59) for dengue, chikungunya,

and Zika as a function of the proportion of a country's gross domestic product (GDP) that comes

from tourism. The percent of GDP from tourism is used as a proxy for vector control effort and

infrastructure that prevents mosquito contacts (air conditioning and window screens). Lines and

95% confidence intervals are based on the entire dataset, even though the data are plotted in

three discrete percentile groups (see Materials and Methods and Supplemental Results for

details). Predicted R_0 from the mechanistic model ($X^2 = 10.26$, $df = 1$, $p = 0.001$) and its

interaction with the log of percent of GDP in tourism ($X^2 = 4.14$, $df = 1$, $p = 0.042$) significantly

predicted the observed maximum relative incidence.

Fig. 3. Map of predicted temperature suitability for virus transmission by *Ae. albopictus* and *Ae.*

aegypti. Color indicates the consecutive months in which temperature is permissive for

transmission (predicted $R_0 > 0$) for *Aedes* spp. transmission. Red, minimum likely range (>

97.5% probability that $R_0 > 0$), purple, median likely range (> 50% probability that $R_0 > 0$), teal,

maximum likely range (> 2.5% probability that $R_0 > 0$). Model suitability predictions combine

temperature mean and 8°C daily variation and are informed by laboratory data (Figs. S1-S2) and

validated against field data (Figs. 1-2).

Figure 1

Figure 2

Figure 3