Title

A hidden Markov model approach for simultaneously estimating local ancestry and admixture time using next generation sequence data in samples of arbitrary ploidy

Short Title

Estimating local ancestry and admixture time in samples of arbitrary ploidy

Authors

Russell Corbett-Detig[1,2,*] and Rasmus Nielsen[2,3]

[1]Department of Biomolecular Engineering, UC Santa Cruz, Santa Cruz, CA
[2]Department of Integrative Biology, UC Berkeley, Berkeley, CA
[3]The Natural History Museum of Denmark, University of Copenhagen, Denmark.
*Correspondence to: russcd@gmail.com

Keywords

Local Ancestry Inference, Admixture, Hidden Markov Model, *Drosophila melanogaster*, Pool-seq

Abstract

Admixture—the mixing of genomes from divergent populations—is increasingly

appreciated as a central process in evolution. To characterize and quantify patterns of

admixture across the genome, a number of methods have been developed for local ancestry

inference. However, existing approaches have a number of shortcomings. First, all local

ancestry inference methods require some prior assumption about the expected ancestry

tract lengths. Second, existing methods generally require genotypes, which is not feasible to

obtain for many next-generation sequencing projects. Third, many methods assume

samples are diploid, however a wide variety of sequencing applications will fail to meet this

assumption. To address these issues, we introduce a novel hidden Markov model for

estimating local ancestry that models the read pileup data, rather than genotypes, is

generalized to arbitrary ploidy, and can estimate the time since admixture during local

ancestry inference. We demonstrate that our method can simultaneously estimate the time

since admixture and local ancestry with good accuracy, and that it performs well on

samples of high ploidy—*i.e.* 100 or more chromosomes. As this method is very general, we

expect it will be useful for local ancestry inference in a wider variety of populations than

what previously has been possible. We then applied our method to pooled sequencing data

derived from populations of *Drosophila melanogaster* on an ancestry cline on the east coast

of North America. We find that regions of local recombination rates are negatively

correlated with the proportion of African ancestry, suggesting that selection against foreign

ancestry is the least efficient in low recombination regions. Finally we show that clinal

outlier loci are enriched for genes associated with gene regulatory functions, consistent

with a role of regulatory evolution in ecological adaptation of admixed *D. melanogaster*

populations. Our results illustrate the potential of local ancestry inference for elucidating

fundamental evolutionary processes.

Author Summary

When divergent populations hybridize, their offspring obtain portions of their genomes

from each parent population. Although the average ancestry proportion in each descendant

is equal to the proportion of ancestors from each of the ancestral populations, the

contribution of each ancestry type is variable across the genome. Estimating local ancestry

within admixed individuals is a fundamental goal for evolutionary genetics, and here we

develop a method for doing this that circumvents many of the problems associated with

existing methods. Briefly, our method can use short read data, rather than genotypes and

can be applied to samples with any number of chromosomes. Furthermore, our method

simultaneously estimates local ancestry and the number of generations since admixture—

the time that the two ancestral populations first encountered each other. Finally, in

applying our method to data from an admixture zone between ancestral populations of

Drosophila melanogaster, we find many lines of evidence consistent with natural selection

operating to against the introduction of foreign ancestry into populations of one

predominant ancestry type. Because of the generality of this method, we expect that it will

be useful for a wide variety of existing and ongoing research projects.

Introduction

Characterizing the biological consequences of admixture—the mixing of genomes from divergent ancestral populations—is a fundamental and important challenge in evolutionary genetics. Admixture has been reported in a variety of natural populations of animals [1,2], plants [3-5] and humans [6,7], and theoretical and empirical evidence suggests that admixture may affect a diverse suite of evolutionary processes. Individuals' ancestry can affect disease susceptibility in admixed populations, and inferring and correcting for sample population ancestries is a common practice in human genome wide association studies [8-10]. More generally, admixture has the potential to influence patterns of genetic variation within populations [11,12], to introduce novel adaptive [13,14] and deleterious variants [7,15,16], as well as to disrupt epistatic gene networks [17,18]. Therefore, developing a comprehensive understanding of the extent of admixture in natural populations and resulting mosaic genome structures is essential to furthering our understanding of a variety of evolutionary processes.

Estimating genome-wide ancestry proportions has become a common practice in population genetic inference. For example, the program STRUCTURE [19], originally released in 2000, uses a Bayesian framework to model the ancestry proportions of individuals derived from any number of source populations based on genotype data at a set of unlinked genetic markers. More recently, this model for ancestry proportion estimation has been extended to cases where individual genotypes are not known, but can be studied probabilistically using low-coverage sequencing short read sequencing data [20], which is an important step towards accommodating modern sequencing practices. Additionally,

Bergland *et. al.* [21] developed a method for estimating ancestry proportions in pooled

population samples of relatively high ploidy (*i.e.* 40-250 distinct chromosomes) from short

read sequencing data. In general, it is straightforward to estimate genome-wide ancestry

proportions using a number of sequencing strategies and applications.

It is substantially more challenging to accurately estimate local ancestry (LA) at markers

distributed along the genome of a sample. Nonetheless, analyses of LA have the potential to

yield more nuanced insights into our understanding of the evolutionary processes affecting

ancestry proportions across the genome. One of the first LA inference (LAI) methods was

an extension of the STRUCTURE [19] framework that modeled the correlation in ancestry

among markers due to linkage. Because the ancestry at each locus is not observed, Falush

et al. [22] suggested that a hidden Markov model (HMM) is a straightforward means of

inferring the ancestry states at each site in the genome (which are unobserved) based on

observed genotype data distributed along a chromosome. Most subsequent LAI methods

have also used an HMM framework, and the majority are geared towards estimating LA in

admixed human populations (*e.g.* [23,24]). Consequently, most existing LAI methods are

limited to diploid genomes with high quality genotype calls. Furthermore, many methods

require phased reference panels [24,25], and require the user to provide an estimate of, or

make implicit assumptions about, the number of generations since the initial admixture

event [2,23-25]. This is straightforward with human population genomic samples, where

abundant high quality genotyped samples are available and for which well-documented

demographic histories are sometimes known. However for most other species,

demographic histories are less well characterized, and assumptions about admixture times

may bias the result of LAI methods.

A number of approaches exist to estimate the time since admixture based on well

characterized ancestry tract length distributions [26-29] but in general, these parameters

are unknown prior to LAI. Conversely, another class of methods can be used to estimate the

time of admixture based on the decay of linkage disequilibrium without performing LAI

[30-32]; however as with LAI procedure above, these approaches are also limited to diploid

genotype data. We may therefore expect to improve LAI by simultaneously estimating LA

and demographic parameters (*e.g.* admixture time). Furthermore, in the majority of

sequencing applications, relatively low individual sequencing coverage is often used to

probabilistically estimate individual and population allele frequencies (*e.g.* [33]) but these

data are often not sufficient to determine high confidence genotypes that are required for

existing LAI applications. Hence, there is a clear need for a general LAI method that can

accommodate genotype uncertainty and requires less advanced knowledge of admixed

populations' demographic histories.

Here, we introduce a framework for simultaneously estimating LA using short read pileup

data and the time of admixture within a population. Briefly, as with many previously

proposed LAI methods, we model ancestry across the genome of a sample as a HMM. We

estimate LA by explicitly modeling read counts as a function of sample allele frequencies

within an admixed population. Our method is generalized to accommodate arbitrary

sample ploidies, and is therefore applicable to haploid (or inbred), diploid, tetraploid, as

well as pooled sequencing applications. We show that this approach accurately infers the

time since admixture when data are simulated under the assumed model. Furthermore, our

method yields accurate LA estimates for simulated datasets, including samples of high

sample ploidy and including evolutionary scenarios that violate the assumptions of the

neutral demographic model. In comparisons between ours and an existing LAI method,

WINPOP [34], we find that our approach offers a significant improvement and is accurate

over longer time scales. Furthermore, we demonstrate, using a published dataset, that even

state-of-the-art LAI methods can be significantly impacted by assumptions about the time

since admixture, and that our method provides a solution to this problem.

Finally, we apply this method to a *Drosophila melanogaster* ancestry cline on the east coast

of North America. This species originated in sub-Saharan Africa, and approximately

10,000—15,000 years ago a subpopulation expanded out of the ancestral range. During

this expansion, the derived subpopulation experienced a population bottleneck that

resulted in decreased nucleotide polymorphism, extended linkage disequilibrium within

the derived population and substantial genetic differentiation between ancestral and

derived populations [2,35-39]. Hereafter, the ancestral population will be referred to as

"African" and the derived population as "Cosmopolitan". Following this bottleneck,

descendant populations of African and Cosmopolitan *D. melanogaster* have admixed in

numerous geographic regions [2,11,21]. Of particular relevance to this work, North

America was colonized recently by a population descendent from African individuals from

the South, and by a population descendent from cosmopolitan *D. melanogaster* in the North

[11,21,38]. Where these populations encountered each other in eastern North America,

they form an ancestry cline where southern populations have a greater contribution of

African ancestry than northern populations [21].

Previous work on these ancestry clines has shown that ancestry proportions vary across

populations with increasing proportions of cosmopolitan alleles in more temperate

localities. Evidence suggests spatially varying selection affects the distribution of genetic

variants [40-45]. Furthermore, strong epistatic reproductive isolation barriers partially

isolate individuals from northern and southern populations along this ancestry cline

[46,47]. This may be generally consistent with recent observations of ancestry-associated

epistatic fitness interactions within a *D. melanogaster* population in North Carolina [17],

and with the observation of widespread fitness epistasis between populations of this

species more generally [48]. There is therefore good reason to believe that natural

selection has acted to shape LA clines that are tightly linked to selected mutations in these

D. melanogaster populations.

Here, we show that African ancestry in North American *D. melanogaster* populations is

negatively correlated with recombination rates, consistent with more efficient selection

against foreign ancestry in high recombination rate regions of the genome. We also find

that the X chromosome displays a higher rate of LA outlier loci, potentially consistent with

a greater role of the X chromosome in clinal adaptation. Clinal loci are disproportionately

likely to be associated with high level gene regulatory protein complexes, and may play

important roles in ecological divergence between African and Cosmopolitan *D.*

melanogaster populations. Furthermore, we identify numerous loci with decreased African

ancestry across all populations, which suggests that these alleles that are disfavored on

predominantly cosmopolitan genetic backgrounds. This subset of loci is enriched for genes

related to oogenesis, potentially consistent with epistatic interactions that affect female

reproductive success in these populations.

Results and Discussion

The Model

Although admixed populations often are diploid, we derived a general model of ploidy in

which the individual has n gene copies at each locus, i.e. for diploid species $n = 2$. In

practice, sequences are often obtained from fully or partially inbred individuals (e.g.

[39,49]), which represent only a single uniquely derived chromosome. It is also common to

pool individuals prior to sequencing for allele frequency estimation, so called pool-seq (e.g.

[21,40,42,50-53]). If the pooling fractions are exactly equal, such a sample of b diploid

individuals can be treated as a sample from a single individual with ploidy $n = 2b$. Although

that requirement is restrictive, pool-seq has been experimentally validated as a method for

accurate allele frequency estimation—*i.e.* alleles are approximately binomially sampled

from the sample allele frequencies [54]. We therefore aimed to derive a model that can

accommodate arbitrary sample ploidies. In the model, we assumed that the focal

population was founded following a single discrete admixture event between two ancestral

subpopulations, labeled 0 and 1, with admixture proportions 1-m and m, respectively, at a

time t generations in the past. We modeled emission probabilities such that the method

can work directly on read pileup data, rather than high quality known genotypes. Briefly,

in our model, we specify an HMM $\{H_v\}$ with state space $S = \{0,1,...,n\}$, where $H_v = i$, $i \in S$,

indicates that in the vth position i chromosomes are from population 0 and $n - i$

chromosomes are from population 1. In other words, this HMM enables one to estimate

what ancestry frequencies are present at a given site along a chromosome within a sample.

Importantly, we designed this method to simultaneously estimate the time of admixture,

which is related to the correlation between ancestry informative markers along a

chromosome. See Methods for a complete description of the HMM including the emissions

and transition probability calculations. The source code and manual are available at

https://github.com/russcd/Ancestry_HMM. For this model, it is assumed that the number

of chromosomes present in a sample, n, is known and that the global ancestry proportion,

m, is known. As there are many methods for accurately estimating m in a wide variety of

contexts implemented in standard population genetic analysis pipelines [19,20], we believe

this assumption is not too restrictive.

Admixture Simulation Framework

In order to test our method with data of known provenance, we also developed an

approach for simulating chromosomes sampled from admixed populations. Briefly, we first

simulated genetic diversity consistent with ancestral populations using a coalescent

simulation method [55]. We then generated ancestry tracts consistent with admixture

models developed to test our inference method using the forward-time admixture

simulation program, SELAM [56]. We retained a portion of each coalescent-generated

population to serve as a reference panel for allele frequency and LD estimation. We then

took the remaining chromosomes and placed them on the appropriate ancestry tracts in

admixed chromosomes. Finally, we generated read counts for these chromosomes, or pools

of chromosomes for samples with ploidy greater than one, via binomial sampling from the

genotype frequencies of the sample. Implicitly, this procedure assumes that the allele

frequencies in the reference panel and the admixed individuals whose ancestry is from a

given reference panels are equivalent. For large, well-mixed populations such as those of *D.*

melanogaster, this is likely to be a reasonable assumption. Nonetheless, below we assess

the impact of differences in the ancestral allele frequencies for plausible demographic

models in this species.

Dependence on Ancestral Linkage Disequilibrium

Within an admixed population, there are two sources of LD. LD that is induced due to the

correlation of alleles from the same ancestry type (*i.e.* admixture LD), and LD that is

present within each of the ancestral populations (ancestral LD). Admixture LD, is the signal

of LA that we seek to detect using the HMM. The second type, ancestral LD, limits the

independence of the ancestral information captured by each marker, and is expected to

confound HMM-based analyses, particularly as we aimed to estimate the time since

admixture within this framework. We therefore sought to quantify the effect of ancestral

LD by discarding one of each pair of sites in LD within either ancestral population. We

found that ancestral LD tends to increase admixture time estimates obtained using our

method, and we decreased the cutoff of the LD parameter, $|r|$, by 0.1 until the time

estimates obtained for single chromosomes were unbiased with respect to the true time

since admixture. We found that $|r| \leq 0.4$ fit this criterion, although for relatively ancient

admixture events with highly skewed ancestry proportions—*i.e.* $m < 0.1$ or $m > 0.9$—some

residual bias was apparent in the estimates of admixture time (Figure 1). This reflects the

fact that the SMC' ancestry tract distribution performs poorly with highly skewed ancestry

proportions and especially for long times since admixture [57].

Figure 1 also reveals a striking difference between otherwise equivalently skewed

admixture proportions. For example when m = 0.1, there was a much larger effect of

ancestral LD than when m = 0.9. This is due to differences in the variability of LD within the

ancestral populations. That is, due to the strong population bottleneck, cosmopolitan *D.*

melanogaster populations have substantially more LD and fewer polymorphic sites than

African *D. melanogaster* populations. Because the time estimation procedure appears to be

sensitive to the amount of ancestral LD present in the data, simulations of the type we

described here may be necessary to determine what $|r|$ cutoffs are required to produce

unbiased time estimates given the ancestral LD of the populations in a given analysis using

this method.

Accuracy and Applications to Diploid and Pooled Samples

We next sought to quantify the accuracy of our approach across varying sample ploidies

and times since admixture (Figure 2). Especially for moderate and short admixture times

(*i.e.* 0—500 generations), our method performed well for all ploidies considered and we

were able to accurately recover the correct admixture time with relatively little bias.

However, as true admixture time increases, the time estimates for pooled samples become

significantly less reliable and show a clear negative bias. Nonetheless, across the range of

times presented in Figure 2, samples of ploidy one and two showed little bias, and we

therefore believe our method will produce sufficiently accurate admixture time estimates

for a wide variety of applications.

All measures of accuracy decrease with increasing time since admixture (Figure 2).

However, even for relatively long times since admixture—2000 generations—and for large

sample ploidies, the mean posterior error remained relatively low for all ancestry

proportions and for long times since admixture. This indicates that this approach may be

sufficiently accurate for a wide variety of applications, sequencing depths, and sample

ploidies. Nonetheless, the proportion of sites within the 95% credible interval decreased

with larger pool sizes and it is clear that for larger pools the posterior credible interval

tends to be too narrow. Therefore, correcting for this bias may be necessary for

applications that are sensitive to the accuracy of the credible interval.

An important consideration is that estimates of t will be reliable only if the local

recombination rates are known with reasonably high accuracy [58]. In many species, an

accurate broad-scale map is available. However, fine-scale variation in recombination rates

has only been documented for a few model species. Therefore, for relatively short to

moderate times since admixture, error in the genetic map is expected to have a limited

impact on date estimates. However, for longer times since admixture, this factor has the

potential to bias estimates of t [58], particularly in species with large variance in local

recombination rates (*e.g.* due to hotspots). Since *D. melanogaster* has one of the best

recombination maps currently available in any species [59] and because we do not aim to

estimate time in our applications, we do not believe this will heavily impact the analyses

we present below. However, for most applications, it will be necessary to consider the

impact of error in the assumed genetic map to accurately interpret estimates of t obtained

using this method. We emphasize that this challenge is not unique to this application, but

will impact virtually all ancestry estimation methods that rely on a genetic map for

estimating the time since admixture.

Non-Independence Among Ancestry Tracts

As described above, estimates of the time of admixture demonstrate an apparent bias in

pools of higher ploidy (Figure 2). Specifically, time tends to be slightly overestimated for

relatively short admixture times and underestimated at relatively long admixture times.

This is particularly apparent at highly skewed ancestry proportions. Given that this bias is

primarily evident in pools of 10 to 20 individuals, we hypothesized that it might be due to

the non-independence of ancestry tracts among chromosomes, which should tend to

disproportionately affect samples of higher ploidy because all ancestry breakpoints are

assumed to be independent in our model. To test this, we simulated genotype data from

independent and identically distributed exponential tract lengths as is assumed by our

model. When we ran our HMM on this dataset, we found that no bias is evident for

simulations of up to 2000 generations (Supplemental Figure S1), indicating that the

primary cause of this bias was violations in the real data of the independence of ancestry

tracts that we assumed when computing the transition probabilities. However, it should be

possible to quantify and correct for this bias in applications of this method that aim to

estimate the time since admixture.

Robustness to Unknown Population Size

The transition probabilities of this HMM depend on knowledge of the population size. In

practice, this parameter is unlikely to be known with certainty. Hence, to assess the impact

of misspecification of the population size, we performed simulations using a range of

population sizes that span three orders of magnitude (N=100, 1000, 10000, and 100000).

All analyses presented here were conducted by applying our HMM to haploid and diploid

samples, but qualitatively similar results hold for samples of larger ploidy. We then

analyzed these data assuming the default population size, 10000, is correct. For relatively

short times since admixture, there was not a clear bias for any of the true population sizes

considered. However, at longer true admixture times, estimated admixture times for both

N=100 and N=1000 asymptote at a number of generations near to the population sizes.

This result reflects the fact that smaller populations will tend to coalesce at a portion of the

loci in the genome relatively quickly, and ancestry tracts cannot become smaller following

coalescence. Nonetheless, the accuracy of LAI remained high even when time estimates

were unreliable (Supplemental Figure S2) for the tested marker densities and patterns of

LD. Furthermore, in some cases it should be straightforward to determine if a population

has coalesced to either ancestry state at a large portion of the loci in the genome,

potentially obviating this issue.

A more subtle departure from the expectation was evident for population sizes that are

larger than we assumed in analyzing these data (Supplemental Figure S2). This likely

reflects the fact that the probability of back coalescence to the previous marginal genealogy

to the left after a recombination event is inversely related to the population size. Hence, the

rate of transition between ancestry types is actually slightly higher in larger populations

where back coalescence is less likely than we assumed during the LAI procedure. This

produced a slight upward bias in the estimates of admixture time when the population was

assumed to be smaller than it is in reality. However, this bias appears to be relatively

minor, and we expect that time estimates obtained using this method will be useful so long

as population sizes can be approximated to within an order of magnitude. Of course, this

bias is not unique to our application, and it will affect methods that aim to estimate

admixture time after LAI as well. That is, estimating the correct effective population size is

an inherent problem for all admixture demographic inference methods.

Application to Ancient Admixture

Although it is clear that accurately estimating relatively ancient admixture times is

challenging in higher ploidy samples, we sought to determine the limits of our approach for

LAI and time estimation for longer admixture times for haploid sequence data. Because of

rapid coalescence in smaller samples (see above), we performed admixture simulations

with a diploid effective population size of 100,000. It is clear that there is a limit to the

inferences that can be made directly using our method. Like the higher ploidy samples,

time estimates for haploid samples departed from expectations shortly after 2,000

generations since admixture (Supplemental Figure S3). Nonetheless, the magnitude of this

bias is slight, and it is likely that it could be corrected for when applying this method even

for very ancient admixture events. For all admixture times considered, LAI remained

acceptably accurate despite the slight bias in time estimates (Supplemental Figure S3).

Reference Panel Size

One question is what effect varying the reference panel sizes will have on LAI inference using this method. We therefore compared results from reference panels of size 10 chromosomes with those from panels of size 100 chromosomes (Supplemental Figure S4). As with results obtained for reference panels of size 50, panels of size 100 were sufficient to accurately estimate admixture time and LA over many generations since admixture. Whereas, when panel sizes were just 10 chromosomes, time estimates were clearly biased and the result was variable across ancestry proportions (Supplemental Figure S4). However, since there was a strong correlation between true and estimated admixture times even with relatively small panel sizes, it may therefore be possible to infer the correct time by quantifying this bias through simulation and correcting for it. Furthermore, although LAI is clearly less reliable with smaller panels, these results are not altogether discouraging and this approach, in conjunction with modest reference panels may still be effective for some applications.

Allele Frequency Differences Between Ancestral and Admixed Populations

Ultimately, there are three reasons why allele frequencies in the reference panels and in the admixed population panel would be expected to differ beyond that expected from binomial samples with the same mean. First, some amount of genetic drift may have occurred in the ancestral population and in the admixed population in the time since the admixed population was founded. Second, in some cases, it is infeasible to sample the ancestral population of an admixed group, and a genetically divergent population must suffice as the reference panel if this method is to be used. Third, divergent selection may

quickly modify allele frequencies between admixed and ancestral populations. Hence,

genetic divergence between reference and admixed populations may be an important

challenge for this method.

To address this, we simulated the second scenario, where increasingly divergent

populations are used as the reference panels to study admixed populations. In order to

make this relevant to the application to *D. melanogaster* populations, below, we selected

times for divergence that might be consistent with differences across continental

populations in Sub-Saharan Africa and in Cosmopolitan populations. Although time

estimates obtained using this approach are weakly positively biased with increasing

divergence between the ancestral population and reference panels, the accuracy of this LAI

method is largely unaffected (Supplemental Figure S5). Hence, for biological scenarios

potentially consistent with those of *D. melanogaster* ancestral populations, we do not

expect this challenge to strongly bias our method. Nonetheless, in applications to other

populations, with potentially differently structured ancestral populations, it would be

necessary to examine the effects of this bias in detail.

High Sample Ploidy

In a wide variety of pool-seq applications, samples are pooled in larger groups than we

have considered above (*e.g.* [40,50,52]). We are therefore interested in determining how

our method will perform on pools of 100 individuals. Towards this, we performed

simulations as before, but we designed our parameters to resemble those of the pooled

sequencing data that we analyze in the application of this method below. Specifically, we

simulated data with a mean sequencing depth of 25, a time since admixture of 1500

generations, and an ancestry proportion of 0.8. Consistent with results for ploidy 20, we

found that time tends to be dramatically underestimated (*i.e.* the mean estimate of

admixture time was 680 generations). However, when we provided the time since

admixture, our method produced reasonably accurate LAIs for these samples. Although the

posterior credible interval was again too narrow, the mean posterior error was just 0.053

when expressed as an ancestry frequency, indicating that this approach can produce LA

estimates that are close to their true values for existing sequencing datasets (*e.g.* Figure 3).

However, the HMM's run time increases dramatically for higher ploidy samples and higher

sequencing depths, a factor that may affect the utility of this program for some analyses.

Nonetheless, for more than 36,000 markers, a sample ploidy of 100 and a mean sequencing

depth of 25, the average runtime was approximately 42 hours. In contrast, for the same set

of parameters, but where individuals are sequenced and analyzed as diploids, the mean

runtime was just 8 minutes (See Supplemental Table S1 for a comparison of run times

across many parameter sets).

Robustness to Deviations From the Neutral Demographic Model

An important concern is that many biologically plausible admixture models would violate

the assumptions of this inference method. In particular, continuous migration and selection

acting on alleles from one parental population are two potential causes of deviation from

the expected model in the true data. To assess the extent of this potential bias, we

performed additional simulations. First, we considered continuous migration at a constant

rate that began *t* generations prior to sampling. In simulations with continuous migration,

additional non-recombinant migrants enter the population each generation. Relative to a

single pulse admixture model, this indicates that the ancestry tract lengths will tend to be

longer than those under a single pulse admixture model in which all individuals entered at

time t. Indeed, we found that admixture times tended to be underestimated with models of

continuous migration. However, the accuracy of LAI remained high across all situations

considered here (Table 1), indicating that the LAI aspect of this approach may be robust to

alternative demographic models.

Table 1. Parameter estimation and LAI when admixture occurs at a constant rate, rather

than in a single pulse.

Admixture Time	Migration Rate	Sample Ploidy	Estimated Time	Proportion in 95% CI	Mean 95% CI Width	Mean Posterior Error	Proportion MLE Correct
100	0.0005	1	53	1.000	0.002	0.001	1.000
		2	49	1.000	0.006	0.001	0.998
		10	129	0.963	0.305	0.017	0.839
		20	98	0.545	0.328	0.033	0.353
	0.001	1	55	1.000	0.004	0.001	0.999
		2	53	1.000	0.013	0.002	0.997
		10	156	0.951	0.558	0.028	0.727
		20	90	0.551	0.719	0.043	0.179
	0.002	1	54	1.000	0.006	0.002	0.999
		2	52	0.999	0.019	0.003	0.996
		10	123	0.949	0.758	0.035	0.671
		20	74	0.679	1.115	0.045	0.176
	0.004	1	43	1.000	0.008	0.002	0.998
		2	54	0.999	0.035	0.005	0.993
		10	91	0.955	1.085	0.044	0.605
		20	75	0.860	1.788	0.045	0.248
500	0.0005	1	254	0.999	0.033	0.010	0.993
		2	250	0.997	0.121	0.018	0.974
		10	331	0.956	1.395	0.027	0.557
		20	333	0.882	2.321	0.051	0.261
	0.001	1	266	0.999	0.049	0.014	0.990
		2	268	0.996	0.198	0.027	0.962

	10	325	0.967	1.887	0.063	0.521
	20	366	0.926	3.049	0.055	0.294
0.002	1	294	0.999	0.055	0.016	0.989
	2	297	0.996	0.238	0.032	0.956
	10	352	0.977	2.076	0.064	0.542
	20	370	0.951	3.238	0.054	0.336
0.004	1	346	0.999	0.038	0.010	0.993
	2	350	0.997	0.164	0.021	0.973
	10	403	0.989	1.634	0.045	0.692
	20	462	0.979	2.773	0.041	0.473

In the second set of simulations, we considered additive selection on alleles that are

perfectly correlated with local ancestry in a given region (*i.e.* selected sites with

frequencies 0 in population 0 and frequency 1 in population 1), and experience relatively

strong selection (selective coefficients were between 0.005 and 0.05). We placed selected

sites at 2, 5, 10 and 20 loci distributed randomly across the simulated chromosome, where

admixture occurred through a single pulse. Ancestry tracts tend to be longer immediately

surrounding selected sites, and we therefore expected admixture time to be

underestimated when selection is widespread. When the number of selected loci was small,

time estimates were nearly unbiased (Table 2), suggesting that our approach can yield

reliable admixture time estimates despite the presence of a small number of selected loci

(*i.e.* 2 selected loci on a chromosome arm). However, with more widespread selection on

alleles associated with local ancestry, time estimates showed a downward bias that

increased with increasing numbers of selected loci. This is likely because selected loci will

tend to be associated with longer ancestry tracts due to hitchhiking. However, the accuracy

of the LAI remains high for all selection scenarios that we considered here, further

indicating that our method can robustly delineate LA, even when the data violate

assumptions of the inference method (Table 1,2).

Table 2. Parameter estimation and LAI when a subset of loci experience natural selection

in the admixed population.

Admixture Time	Migration Rate	Sample Ploidy	Estimated Time	Proportion in 95% CI	Mean 95% CI Width	Mean Posterior Error	Proportion MLE Correct
100	0.0005	1	53	1.000	0.002	0.001	1.000
		2	49	1.000	0.006	0.001	0.998
		10	129	0.963	0.305	0.017	0.839
		20	98	0.545	0.328	0.033	0.353
	0.001	1	55	1.000	0.004	0.001	0.999
		2	53	1.000	0.013	0.002	0.997
		10	156	0.951	0.558	0.028	0.727
		20	90	0.551	0.719	0.043	0.179
	0.002	1	54	1.000	0.006	0.002	0.999
		2	52	0.999	0.019	0.003	0.996
		10	123	0.949	0.758	0.035	0.671
		20	74	0.679	1.115	0.045	0.176
	0.004	1	43	1.000	0.008	0.002	0.998
		2	54	0.999	0.035	0.005	0.993
		10	91	0.955	1.085	0.044	0.605
		20	75	0.860	1.788	0.045	0.248
500	0.0005	1	254	0.999	0.033	0.010	0.993
		2	250	0.997	0.121	0.018	0.974
		10	331	0.956	1.395	0.027	0.557
		20	333	0.882	2.321	0.051	0.261
	0.001	1	266	0.999	0.049	0.014	0.990
		2	268	0.996	0.198	0.027	0.962
		10	325	0.967	1.887	0.063	0.521
		20	366	0.926	3.049	0.055	0.294
	0.002	1	294	0.999	0.055	0.016	0.989
		2	297	0.996	0.238	0.032	0.956
		10	352	0.977	2.076	0.064	0.542
		20	370	0.951	3.238	0.054	0.336
	0.004	1	346	0.999	0.038	0.010	0.993
		2	350	0.997	0.164	0.021	0.973
		10	403	0.989	1.634	0.045	0.692
		20	462	0.979	2.773	0.041	0.473

Comparison to WinPop

We next compared the results of our method to those of WinPop [34]. Because WinPop accepts only diploid genotypes, we provided this program diploid genotype data. However, for these comparisons, we still ran our method on simulated read pileups with the mean depth equal to 2. WinPop was originally designed for local ancestry inference in very recently admixed populations. As expected, WinPop performed acceptably for very short admixture times, but rapidly decreased in performance with increasing time (Supplemental Figure S6). However, by default, WinPop removes sites in strong LD within the admixed samples, which includes ancestral LD, but also admixture LD—the exact signal LAI methods use to identify ancestry tracts.

We therefore reran WinPop, but instead of pruning LD within the admixed population, we removed sites in strong LD within the ancestral populations as described above in our method. With this modification, WinPop performs nearly as well as our method, but remains slightly less accurate especially at longer admixture times (Supplemental Figure S6). This difference presumably reflects the windowed-based approach of WinPop. At longer times since admixture a given genomic window may overlap a breakpoint between ancestry tracts. Although the performance is nearly comparable with this modification, we emphasize that our method enables users to estimate the time since admixture, where this must be supplied for WinPop, and allows for LAI on read pileups, therefore incorporating genotype uncertainty into the LAI procedure. Indeed our method is more accurate at longer timescales even when supplied with considerably lower quality read data. However WinPop supports LAI with multiple ancestral populations, which our method currently

does not (but see Conclusions). Furthermore many LAI algorithms utilize haplotype

information, which may be particularly valuable in populations where LD extends across

large distances as in *e.g.* human populations.

Assessing Applications to Human Populations

Given the strong interest in studying admixture and local ancestry in human populations

(*e.g.* [22-25]), it is useful to ask if our method can be applied to data consistent with

admixed populations of humans. Towards that goal, we simulated data similar to what

would be observed in admixture between modern European and African lineages and

applied our HMM to estimate admixture times and LA. We found that our method can

accurately estimate admixture times for relatively short times since admixture, however,

substantially more stringent LD pruning in the reference panels is necessary to produce

unbiased estimates (Figure 4). This may be expected given that linkage disequilibrium

extends across longer distances in human populations than it does in *D. melanogaster*. In

other words, the scales of ancestral LD and admixture LD become similar rapidly in

admixed human populations. Furthermore, this approach yields accurate time estimates

for shorter times since admixture than with genetic data consistent with *D. melanogaster*

populations. For a relatively short time since admixture, around 100 generations, it is

possible to obtain accurate and approximately unbiased estimates of the admixture time

over a wide range of ancestry proportions, indicating that this method may be applicable to

recently admixed human populations as well (Figure 4). Nonetheless, this result

underscores the need to examine biases associated with LD pruning in this approach prior

to application to a given dataset.

Bias in LAI due to Uncertainty in Time of Admixture

To demonstrate that assumptions about the number of generations since admixture have

the potential to bias LAI, we analyzed a SNP-array dataset from Greenlandic Inuits [60,61].

The authors had previously noted a significant impact of t on the LAI results produced

using RFMix [24], which we were able to reproduce here for chromosome 10

(Supplemental Figure S7). Indeed, even for comparisons between $t = 5$ and $t = 20$, both of

which may be biologically plausible for these populations, the mean difference in posterior

probabilities between samples estimated using RFMix was 0.0903 (Supplemental Figure

S7). However, when we applied our method to these data, a clear optimum from t was

obtained at approximately 6-7 generations prior to the present, which is close to the

plausible times of admixture for these populations (Supplemental Figure S7). This

comparison therefore demonstrates that even relatively minor changes in assumptions of t

have the potential to strongly impact LAI results, and underscores the importance of

simultaneously performing LAI while estimating t.

However, these results also indicate that our method may not be robust in situations where

the background LD is high and ancestry informative markers are neither common nor

distributed evenly across the genome. When we compared the results of our method at $t =$

5 and at $t = 20$, we also obtained differences in the mean posterior among individuals as

with RFMix. However, one notable difference is that the mean posterior difference using

RFMix has a particularly high variance and therefore higher mean error (Supplemental

Figure S7), but actually a lower median difference than we found using our method. There

are likely two causes for differences observed in the mean ancestry posterior among

individuals. First, the datasets considered were generated with a metabochip SNP-chip

[62], which contains a highly non-uniform distribution of markers across the genome.

Second, the ancestral LD in the Inuit population is extensive [61], and we could only retain

a relatively small proportion of the markers after LD pruning in the reference panels. These

results therefore also underscore the challenges of LAI when the signal to noise ratio is low

as may be the case in some human populations, for which LD is extensive, and for some

sequencing strategies.

Bias due to Incorrect Estimates of t and m

Although in general it is straightforward to estimate m from genome-wide data, in some

cases this parameter may be misestimated prior to LAI. We therefore sought to quantify

this potential effect by performing LAI after supplying incorrect values of m. In general, we

found that values close to the true range, *i.e.* within 0.05 of the true m, tend to yield

reasonably accurate time estimates. However, increasingly incorrect values produce

sharply downwardly biased time estimates and this effect is especially pronounced for

highly skewed true m (Supplemental Figure S8). As could be expected given the robustness

of LAI to many perturbations (above), when the incorrect t is supplied to the program, the

LA results remain reasonable. However it is worth noting that the penalty appears to be

greatest when t is too small rather than too large (Supplemental Figure S9).

Estimating Confidence Intervals for t

Although this is not a primary focus for this work, for some users it may be of interest to

construct confidence intervals for estimates of t. We recommend the block bootstrap as the

preferred method for estimating confidence interval for t, and we have written a script that

will produce these (available on the github page for this project:

https://github.com/russcd/Ancestry_HMM). Simulations confirm that this can produce

confidence intervals overlapping the true t (Supplemental Figure S10), but bias in t

estimates for higher ploidy samples may still be apparent in some cases.

Patterns of LA on Inversion Bearing Chromosomes in *D. melanogaster*

Given their effects suppressing recombination in large genomic regions, chromosomal

inversions may be expected to strongly affect LAI [2,63]. Although we attempted to limit

the impact of chromosomal inversions by eliminating known polymorphic arrangements

from the reference panels (see methods), many known inversions are present within the

pool-seq samples we aimed to analyze [64]. We therefore focused on known inverted

haplotypes within the DGPR samples [63,65-67], which are comprised of inbred

individuals, and therefore phase is known across the entire chromosome.

In comparing LA estimates between inverted and standard arrangements, it is clear that

chromosomal inversions can substantially affect LA across the genomes (Figure 5). In

general, the chromosomal inversions considered in this work originated in African

populations of *D. melanogaster* [63], and consistent with this observation, most inversion

bearing chromosomes showed evidence for elevated African ancestry. This was

particularly evident in the regions surrounding breakpoints, where recombination with

standard arrangement chromosomes is most strongly suppressed. Importantly, this pattern

continued outside of inversion breakpoints as well, consistent with numerous observations

that recombination is repressed in heterokaryotypes in regions well outside of the

inversion breakpoints in *Drosophila* (*e.g.* [2,63,68]). In(3R)Mo is an exception to this

general pattern of elevated African ancestry within inverted arrangements (Figure 5). This

inversion originated within a cosmopolitan population [63], and has only rarely been

observed within sub-Saharan Africa [69,70]. Consistent with these observations, In(3R)Mo

displayed lower overall African ancestry than chromosome arm 3R than standard

arrangement chromosomes.

Although chromosomal inversions may affect patterns of LA in the genome on this ancestry

cline, we believed including chromosomal inversions in the pool-seq datasets would not

heavily bias our analysis of LA clines. Inversions tend to be low frequency in most

populations studied [64], and because they affect LA in broad swaths of the genome—

sometimes entire chromosome arms—including inversions is unlikely to affect LA cline

outlier identification which appears to affect much finer scale LA (below). Furthermore,

inversion breakpoint regions were not enriched for LA cline outliers in our analysis

(Supplemental Table S2), suggesting that inversions have a limited impact on overall

patterns of local ancestry on this cline. Nonetheless, the LAI complications associated with

chromosomal inversions should be considered when testing selective hypotheses for

chromosomal inversions as genetic differentiation may be related to LA, rather than

arrangement-specific selection in admixed populations such as those found in North

America.

Application to *D. melanogaster* Ancestry Clines

Finally, we applied our method to ancestry clines between cosmopolitan and African

ancestry *D. melanogaster*. Genomic variation across two ancestry clines have been studied

previously [21,38,40,52]. In particular, the cline on the east coast of North America has

been sampled densely by sequencing large pools of individuals to estimate allele

frequencies, and previous work has shown that the overall proportion of African ancestry

is strongly correlated with latitude [21]. Consistent with this observation, we found a

significant negative correlation for all chromosome arms between proportion of average

African ancestry and latitude (rho = -0.891, -0.561, -0.912, -0.913, and -0.755, for 2L, 2R,

3L, 3R, and X respectively).

Although global ancestry proportions have previously been investigated in populations on

this ancestry cline [21,38], these analyses neglected the potentially much richer

information in patterns of LA across the genome. We therefore applied our method to these

samples. Because of the relatively recent dual colonization history of these populations and

subsequent mixing of genomes, a genome-wide ancestry cline is expected [21]. However,

loci that depart significantly in clinality from the genome-wide background levels may

indicate that natural selection is operating on a site linked to that locus.

LA is Correlated with Recombination Rate

Previously Pool (2015) found that regions of low recombination are disproportionately

enriched for African ancestry in the Raleigh, NC population [17]. Here, we find a similar

pattern and we further find that is replicated across all populations that were assayed on

this ancestry cline. Specifically, in all populations studied the proportion of African ancestry

is significantly negatively correlated with local recombination rates (Figure 6). Ultimately,

this correlation may have two causes. First, if selection is more efficient at purging African

alleles in high recombination regions, these loci will tend to be removed preferentially in

those genomic regions. An alternative explanation is that introgressing African alleles that

are favored by selection would tend to bring larger linkage blocks along with them in the

predominantly low recombination regions. Regardless of the specific source of natural

selection, a neutral admixture model would not predict this robust correlation between LA

and recombination rates within all populations, indicating that natural selection has played

an important role in shaping LA on this ancestry cline.

Robustness of LAI to Genomic Heterogeneity

Previous studies have found that heterogeneity in the genome with respect to ancestry

informative markers may impact the accuracy of LAI [71]. To assess this possibility, we

computed the mean difference between posterior mean estimates for the two samples from

Florida and between the two samples from Maine. Importantly, because these pooled

samples were created using different isofemale lines [40], this is a conservative test of our

method since there will be true biological differences as well as stochastic sequencing

differences between replicates from each population. We found no correlation between the

mean difference of the posterior means and local recombination rates (P = 0.2353 and P =

0.7529, Spearman's rank correlation for Florida and Maine respectively), indicating that

the correlation observed between local recombination rates and LA is unlikely to be an

artifact of differential accuracy of LAI in different genomic regions. However, it should be

acknowledged that in some genomic regions it maybe challenging to unambiguously infer

LA [17,71].

Outlier LA Clines

Selection within admixed populations may take several distinct forms. On the one hand,

loci that are favorable in the admixed population—either because they are favored on an

admixed genetic background, enhance reproductive success in an admixed population, or

are favorable in the local environment—will tend to achieve higher frequencies, and we

would expect these sites to have a more positive correlation with latitude than the genome-

wide average. Conversely, loci that are disfavored within the admixed population may be

expected to skew towards a more negative correlation with latitude.

Although it is not possible to distinguish between these hypotheses directly, a majority of

evidence suggests that selection has primarily acted to remove African ancestry from the

largely Cosmopolitan genetic backgrounds found in the Northern portion of this ancestry

cline. First, abundant evidence suggests pre-mating isolation barriers between some

African and cosmopolitan populations [72-74]. Second, there is strong post-mating

isolation between populations on the ends of this cline [46,47]. Third, we report here a

strong negative correlation between LA frequency and local recombination rates (above).

Finally, circumstantially, the local environment on the east coast of North America is

perhaps most similar to the environment of Cosmopolitan compared to African ancestral

populations, which further suggests that Cosmopolitan alleles are likely favored through

locally adaptive mechanisms. For these reasons, we therefore examined loci that are

outliers for a negative partial correlation with latitude, as this is the expected pattern for

African alleles that are disfavored in more temperate populations. In other words, the

outlier regions show a significantly stronger negative correlation between local African

ancestry and latitude than the chromosome arm does on average.

There is an ongoing debate about the relative merits of an outlier approach versus more

sophisticated models for detecting and quantifying selection in genome-wide scans. We

believe that the difficulties of accurately estimating demographic parameters for this

ancestry cline make the outlier approach most feasible for our purposes. Using our outlier

approach, we identified 80 loci that showed the expected negative partial correlation with

latitude (Figure 7). Although the specific statistical threshold that we employed is

admittedly arbitrary, given the strength of evidence indicating widespread selection on

local ancestry in this species (above), we expected that the tail of the LA cline distribution

would be enriched for the genetic targets of selection.

Differences Among Chromosome Arms

Due to the differences in inheritance, evolutionary theory predicts that selection will

operate differently on the X chromosome relative to autosomal loci. Of specific relevance to

this work, the large-X effect [75,76] is the observation that loci on the X chromosome

contribute to reproductive isolation at a disproportionately high rate. Additionally, and

potentially the cause of the large-X effect, due to the hemizygosity of X-linked loci, the X

chromosome is expected to play a larger role in adaptive evolution, the so-called faster-X

effect [77]. There is therefore reason to believe that the X chromosome will play a

significant role in genetically isolating Cosmopolitan and African *D. melanogaster*.

Consistent with a larger role for the sex chromosomes in generating reproductive isolation

or selective differentiation between *D. melanogaster* ancestral populations, we found that

that the X chromosome has a lower mean African ancestry proportion than the autosomes

in all populations. Furthermore, the X displays a stronger correlation between local

recombination rates and the frequency of African ancestry than the autosomes in all 14

populations samples, potentially indicating that selection has had a disproportionately

strong effect shaping patterns of local ancestry on this chromosome than on the autosomes.

In addition, the X has a significantly higher rate of outlier LA clinal loci than the autosomes

(23 LA outliers on the X, 57 on the Autosomes, $p = 0.0341$, one-tailed exact Poisson test).

Although consistent with evolutionary theory, differences between autosomal arms and the

X chromosome may also be explained in part by differences in effective recombination

rates on the X chromosome than the autosomes, differences in power to identify LA clines

associated with chromosome arm specific patterns, or by the disproportionately larger

number of chromosomal inversions on the autosomes than on the X chromosome in these

populations [64,69]. Distinguishing between this hypothesis and confounding factors will

be central to determining whether key results from speciation research are replicated in

much more recently diverged populations.

Biological Properties of Outlier LA Clinal Loci

We next applied gene ontology analysis to the set of outlier genes to identify common

biological attributes that may suggest more specific organismal phenotypes underlying LA

clinal outliers. In total, we identified seven GO terms that remained significant after

applying a 5% FDR correction (Table S3). These GO terms reflect the presence of two

primary clusters of genes. The first, which corresponds broadly to histone acetylation, may

be related to chromatin remodeling and therefore is expected to effect gene expression

levels across a large number of loci. Previous work focused on this ancestry cline has

identified chromatin remodeling genes as a potentially important component locally

adaptive variation on this ancestry cline [78]. This may indicate that this previous efforts to

identify spatially varying selection in these populations may have been detecting selection

on local ancestry components associated with ecological adaptation in ancestral

populations. The second GO cluster, eukaryotic translation initiation factor 2 complex, also

appears to implicate a central role of clinal LA outliers on the regulation of gene expression.

One plausible explanation of these observations is that gene expression, particularly high

level regulation of gene expression, may be especially likely to contribute to epistatic

interactions as these proteins will inherently interact with a diverse set of loci throughout

the genome. Given that two distinct gene clusters related to gene expression are identified

by this analysis, gene expression would appear to be a plausible candidate phenotype to

investigate in future work on ecological divergence and isolating factors in admixed *D.*

melanogaster populations. Testing this prediction empirically through expression profiling

may therefore offer fruitful grounds of understanding the earliest stages of reproductive

isolation.

Regions of Decreased African Ancestry

Another subset of loci that we may wish to identify using these data are those that

contribute to reproductive isolation between African and Cosmopolitan *D. melanogaster*

populations and would therefore be removed by selection from most populations on this

ancestry cline. Although it is possible that Cosmopolitan alleles would be disfavored in an

admixed background as well, because these populations are predominantly Cosmopolitan,

we expect that the majority of selection on negatively epistatically interacting loci would

remove African alleles from populations. To identify these loci, we first computed the mean

African ancestry across all populations, and we then identified the subset of loci that were

in the lowest 5% tail. From those loci, we selected the loci minima from adjacent genomic

windows (see Methods, Figure 8), and we obtained a total of 84 local ancestry outliers.

As with the clinal outlier analysis above, to identify commonalities in the types of loci

identified by this analysis, we performed GO analysis on the set of loci that are outliers for

the mean proportion of African ancestry. After a 5% FDR correction, there are again several

gene clusters that are significantly enriched in this set of outlier loci (Supplemental Table

S4). Of particular interest is the GO term oogenesis, which may indicate that female

reproduction is affected during admixture between cosmopolitan and African populations

of *D. melanogaster*. This finding is particularly interesting in light of the fact that female

fertility is strongly affected when autosomal chromosomes from one end of this ancestry

cline are made homozygous on a genetic background carrying the X chromosome from the

other end of this ancestry cline [47]. Hence, the effects of combining divergence ancestry

types on female fertility, and specifically the genetic basis of oogenesis, may be an

appealing phenotype to characterize in detail in attempting to clarify the genetic effects

that isolate African and Cosmopolitan *D. melanogaster* populations.

Candidate Behavioral Reproductive Isolation Genes

Given the abundance of evidence supporting a role for pre-mating isolation barriers

between African and Cosmopolitan flies [72-74], we are interested in highlighting genes

potentially related to behavioral isolation between ancestral populations of *D.*

melanogaster. Consistent with this observation, one of the strongest LA cline outliers, *egh*,

has been conclusively linked to strong effects on male courtship behavior using a variety of

genetic techniques [79]. Additionally, gene knockouts of CG43759, another LA cline outlier

locus, have strong effects on inter-male aggressive behavior [80], and may also contribute

to behavioral differences between admixed individuals. These loci are therefore appealing

candidate genes for functional follow-up analyses, and illustrate the power of this LAI

approach for identifying candidate genes that are potentially associated with well

characterized phenotypic differences between ancestral populations.

Little Evidence for Seasonal LA Outliers

The Pennsylvania population included in this study has been sampled extensively,

including several paired fall and spring samples across three consecutive years. Previously,

Bergland et al. [40] identified numerous SNPs that showed recurrent and rapid seasonal

frequency changes in the Pennsylvania populations included in this study. They concluded

that these sites are experiencing recurrent selection associated with recurrent

environmental seasonal changes. To determine if LA across the *D. melanogaster* genome

might also experience selection associated with seasonal frequency shifts, we searched for

loci that showed a strong recurrent seasonal shift in LA. However, we identified fewer

significantly seasonal sites than we would expect to by chance (the proportion of

significant site at the alpha = 0.05 level of significance is 0.041). Furthermore, after

applying a false discovery rate correction [81], there are no sites that are significantly

seasonal at the q = 0.1 level. Collectively, these results indicate that LA within the

Pennsylvania populations of *D. melanogaster* remains remarkably stable during seasonal

environmental cycles.

Although this observation may, to a first approximation, appear to be at odds with the

results reported in Bergland et al. [40], we believe that it is consistent with the model

proposed in that work. Specifically, the authors suggested that long term balancing

selection may maintain these seasonally favorable polymorphisms in diverse *D.*

melanogaster populations and even in the ancestors of *D. melanogaster* and *D. simulants*

[40]. We therefore may expect that these polymorphisms will be maintained at similar

frequencies in African and Cosmopolitan populations. Hence, although the seasonal SNPs

change rapidly in frequency between spring and fall [40], the LA at these sites can remain

stable during seasonal fluctuations.

Conclusion

A growing number of next-generation sequencing projects produce low coverage data that

cannot be used to unambiguously assign individual genotypes, but which can be analyzed

probabilistically to account for uncertainty in individual genotypes [82-84]. However, most

existing LAI methods require genotype data derived from diploid individuals. Hence, there

is an apparent disconnect between existing LAI approaches and the majority of ongoing

sequencing efforts. In this work, we developed the first framework for applying LAI to

pileup read data, rather than genotypes, and we have generalized this model to arbitrary

sample ploidies. This method therefore has immediate applications to a wide variety of

existing and ongoing sequencing projects, and we expect that this approach and extensions

thereof will be valuable to a number of researchers. Although evaluating this application is

beyond the scope of this work, one particularly enticing potential use of this method is LAI

in ancient DNA samples for which sequencing depths often preclude accurate genotype

calling. Importantly, it would be straightforward to model site-specific errors in this

framework, which could be particularly important for ancient DNA applications [6].

For many applications, a parameter of central biological interest is the time since

admixture began (t). A wide variety of approaches have been developed that aim to

estimate t and related parameters in admixed populations [26,28-31,85,86]. Many of these

methods are based on an inferred distribution of tract lengths, however, inference of the

ancestry tract length distribution is associated with uncertainty that is typically not

incorporated in currently available methods for estimating t. Furthermore, incorrect

assumptions regarding t have the potential to introduce biases during LAI. Hence, it is

preferable to estimate demographic parameters such as the admixture time during the LAI

procedure. Nonetheless, as noted above, although LAI using our method is robust to many

deviations from the assumed model, admixture time estimates are sensitive to a variety of

potential confounding factors and examining the resulting ancestry tract distributions after

LAI may be necessary to confirm that the assumed demographic model provides a

reasonable fit to the data.

To our knowledge, this is the first method that attempts to simultaneously link LAI and

population genetic parameter estimation directly, and we can envision many extensions of

this approach that could expand the utility of this method to a broad variety of applications.

For example, it is straightforward to accommodate additional reference populations (*e.g.*

by assuming multinomial rather than binomial read sampling). Alternatively, any

demographic or selective model that can be approximated as a Markov process could be

incorporated—in particular, it is feasible to accommodate two-pulse admixture models and

possibly models including ancestry tracts that are linked to positively selected sites. Such

methods can be used to construct likelihood ratio tests of evolutionary models and for

providing improved parameter estimates.

Methods

Constructing Emissions Probabilities

We model the ancestry using an HMM $\{H_v\}$ with state space $S = \{0,1,...,n\}$, where $H_v = i$,

$i \in S$, indicates that in the vth position i chromosomes are from population 0 and $n - i$

chromosomes are from population 1. In the following, to simplify the notation and without

loss of generality, we will omit the indicator for the position in the genome as the structure

of the model is the same for all positions of equivalent ploidy. We assume each variant site

is biallelic, with two alleles A and a, and the availability of reference panels from source

populations 0 and 1 with total allelic counts C_{0a}, C_{1a}, C_{0A}, and C_{1A}, where the two subscripts

refer to population identity and allele, respectively. Also, $C_0 = C_{0A} + C_{0a}$ and $C_1 = C_{1A} + C_{1a}$.

Finally, we also assume we observe a pileup of r reads from the focal population, with r_A

and r_a reads for alleles A and a respectively ($r = r_A + r_a$). The emission probability of state

$i \in S$ of the process is then defined as $e_i = \Pr(r_A, C_{0A}, C_{1A} \mid r, C_0, C_1, H = i, \varepsilon)$, where

ε is an error rate. This probability can be calculated by summing over all possible

genotypes in the admixed sample and over all possible population identities of the reads, as

explained in the following section.

The probability of obtaining r_0 (=$r - r_1$) reads, in the admixed population, from

chromosomes of ancestry 0, given r and the hidden state $H = i$, and assuming no mapping or

sequencing biases, is binomial,

$$(1) \quad r_0 \mid H = i, n, r \sim \mathrm{Bin}(r, i/n)$$

These probabilities are pre-computed in our implementation for all possible values of $i \in S$

and $r_0, 0 \le r_0 \le r$. Similarly, for the reference populations, for $j = 0, 1$,

$$(2) \quad C_{jA} \mid C_j, f_j \sim \mathrm{Bin}(C_j, f_j)$$

where f_j is the allele frequency of allele A in population j. The analogous allelic counts in the

admixed population, denoted $C_{M0a}, C_{M1a}, C_{M0A}$, and C_{M1A}, are unobserved (only reads are

observed for the admixed population), but are also conditionally binomially distributed,

i.e.:

$$(3) \quad C_{M0A} \mid H = i, f_0 \sim \mathrm{Bin}(i, f_0) \text{ and } C_{M1A} \mid H = i, n, f_1 \sim \mathrm{Bin}(n-i, f_1)$$

Finally, in the absence of errors, and assuming no sequencing or mapping biases, the

conditional probability of obtaining r_{0A} reads of allele A in the admixed population is

$$(4) \quad r_{0A} \mid H = i, r_0, C_{M0A} \sim \mathrm{Bin}(r_0, C_{M0A} / i)$$

It should be noted that because we are explicitly modeling the process of sampling alleles

from the population (Equation 3) and the process of sampling reads conditional on the

sample allele frequencies (Equation 4), that this approach corrects for the increased

variance associated with two rounds of binomial sampling in poolseq applications that has

been reported previously (*e.g.*, in [52]).

This probability can be expanded to include errors, in particular assuming a constant and

symmetric error rate ☐ between major and minor allele, and assuming all reads with

nucleotides that are not defined as major or minor are discarded, we have

$$(5) \quad \square_{0\square} \mid \square = \square, \square_\square, \square_{\square 0\square}, \square \sim \mathrm{Bin}(\square_0, (1-\square)\square_{\square 0\square}/\square + \square(1-\square_{\square 0\square}/\square)).$$

Using these expressions, and integrating over allele frequencies in the source populations,

we have

$$(6)\ \Pr\left(\square_{0\square},\square_{0\square,}\middle|\square_0,\square_0,\square,\square=\square,\varepsilon\right)=$$

$$\int_0^1 \sum_{\square=0}^{\square}\Pr(\square_{0\square}|\square=\square,\square_0,\square_{\square0\square}=\square,\square)\Pr(\square_{\square0\square}=\square|\square=\square,\square_0)\square(\square_0)\square\square_0 \quad =$$

$$(7)\ \frac{\square_0!\,\square!}{(\square_0-\square_{0\square})!\,\square_{0\square}!\,(\square_0+\square+1)!}\sum_{\square=0}^{\square}\Pr(\square_{0\square}|\square=\square,\square_0,\square_{\square0\square}=\square,\square)\frac{(\square_0-\square_{0\square}+\square-\square)!\,(\square_{0\square}+\square)!}{(\square-\square)!\,\square!}$$

907 assuming a uniform [0, 1] distribution for f_0. A similar expression is obtained for

908 $\Pr\left(\square_{1\square},\square_{1\square,}\middle|\square_1,\square_1,\square,\square=\square,\varepsilon\right)$, assuming $f_1 \sim U[0,1]$, and these expressions combine

909 multiplicatively to give

$$(8)\ \Pr\left(\square_{\square},\square_{1\square,},\square_{0\square,}\middle|\square_0,\square_0,\square_1,\square_1,\square,\square=\square,\varepsilon\right)=$$

$$\sum_{\square_{0\square}=\max\{0,\square_{\square}-\square_1\}}^{\min\{\square_0,\square_{\square}\}}\Pr\left(\square_{0\square},\square_{0\square,}\middle|\square_0,\square_0,\square,\square=\square,\varepsilon\right)\Pr\left(\square_{1\square}=\right.$$

$$\square_{\square}-\square_{0\square},\square_{1\square,}\middle|\square_1,\square_1,\square,\square=\square,\varepsilon),$$

915 and the emission probabilities become

917 (9) $\Pr(r_A, C_{0A}, C_{1A} \mid r, C_0, C_1, H = i, \varepsilon) =$

$$\sum_{\square_0=0}^{\square}\Pr(\square_0|\square=\square,\square,\square)\,\Pr\left(\square_{\square},\square_{1\square,},\square_{0\square,}\middle|\square_0,\square_0,\square_1=\square-\square_0,\square_1,\square,\square=\square,\varepsilon\right)$$

919 Alternatively, if the sample genotypes are known with high confidence, i.e. $C_{MA} = C_{M0A} + C_{M1A}$

920 is observed, the emission probabilities are the defined as

$$(10)$$

923
$$\Pr\left(C_{MA}, C_{0A}, C_{1A} \mid C_0, C_1, n, H = i\right) =$$

$$\binom{C_0}{C_{0A}}\binom{C_1}{C_{1A}} \sum_{k=\max\{C_{MA}-i,0\}}^{\min\{n-i,C_{MA}\}} \int_0^1 \binom{n-i}{k}(f_0)^{C_{0A}+k}(1-f_0)^{C_0+n-i-C_{0A}-k} df_0 \int_0^1 \binom{i}{C_{MA}-k}(f_1)^{C_{MA}-k+C_{1A}}(1-f_1)^{C_1+i-C_{1A}-C_{MA}+k} df_1$$

$$= \sum_{k=\max\{C_{MA}-i,0\}}^{\min\{n-i,C_{MA}\}} \frac{C_0! C_1! i!(n-i)!(C_{MA}+C_{1A}-k)!(C_{0A}+k)!(C_1-C_{MA}-C_{1A}+i+k)!(C_0-C_{0A}-i-k+n)!}{(C_0-C_{0A})! C_{0A}!(C_1-C_{1A})! C_{1A}!(C_{MA}-k)! k!(k+i-C_{MA})!(n-k-i)!(n-i+C_0+1)!(i+C_1+1)!}$$

These emissions probabilities are sometimes substantially faster to compute than those for

short read pileups, especially when sequencing depths are high. However, the genotypes must

be estimated with high accuracy for this approach to be valid. For applications with low read

coverage, or with ploidy >2 for which many standard genotype callers are not applicable, it is

usually preferable to use the pileup-based approach described above.

Constructing Transition Probabilities

We assume an admixed population, of constant size, with N diploid individuals, in which a

proportion m of the individuals in the population where replaced with migrants t

generations before the time of sampling. Given these assumptions, and an SMC' model of

the ancestral recombination graph [87], the rate of transition from ancestry 0 to 1, along

the length of a single chromosome, is

938
$$\lambda_0 = 2Nm\left(1 - e^{\frac{-t}{2N}}\right)$$

$$(11)$$

940   per Morgan [57]. Similarly, the rate of transition from ancestry 1 to 0 on a single

941   chromosome is

943
$$\lambda_1 = 2N(1-m)\left(1 - e^{\frac{-t}{2N}}\right)$$
(12)

per Morgan. Importantly, because these expressions are based on a coalescence model,

they account for the possibility that a recombination event occurs between two tracts of

the same ancestry type and the probability that the novel marginal genealogy will back-

coalesce with the previous genealogy [57]. Both events are expected to decrease the

number of ancestry switches along a chromosome and ignoring their contribution will

cause overestimation of the rate of change between ancestry types between adjacent

markers.

The transition rates are in units per Morgan, but can be converted to rates per bp, by

multiplying with the recombination rate in Morgans/bp, r_{bp} within a segment. The

transition probabilities of the HMM for a single chromosome, $\mathbf{P}(l) = \{P_{ij}(l)\}, i, j \in S$, between

two markers with a distance l between each other, is then approximately

958
$$\mathbf{P}(l) = \begin{bmatrix} 1 - \lambda_0 r_{bp} & \lambda_0 r_{bp} \\ \lambda_1 r_{bp} & 1 - \lambda_1 r_{bp} \end{bmatrix}^l$$
(13)

960 using discrete distances, or

963 $$\mathbf{P}(l) = \begin{bmatrix} \dfrac{\lambda_1}{\lambda_0 + \lambda_1} + \dfrac{\lambda_0}{\lambda_0 + \lambda_1} e^{-r_{bp}l(\lambda_0 + \lambda_1)} & \dfrac{\lambda_0}{\lambda_0 + \lambda_1} - \dfrac{\lambda_0}{\lambda_0 + \lambda_1} e^{-r_{bp}l(\lambda_0 + \lambda_1)} \\ \dfrac{\lambda_0}{\lambda_0 + \lambda_1} + \dfrac{\lambda_1}{\lambda_0 + \lambda_1} e^{-r_{bp}l(\lambda_0 + \lambda_1)} & \dfrac{\lambda_1}{\lambda_0 + \lambda_1} - \dfrac{\lambda_1}{\lambda_0 + \lambda_1} e^{-r_{bp}l(\lambda_0 + \lambda_1)} \end{bmatrix}$$

(14)

using continuous distances along the chromosome. Here, we use the continuous

representation for calculations. We emphasize that the assumption of a Markovian process

is known to be incorrect [57], in fact admixture tracts tend to be more spatially correlated

than predicted by a Markov model, and the degree and structure of the correlation depends

on the demographic model [57]. Deviations from a Markovian process may cause biases in

the estimation of parameters such as t.

The Markov process defined above is applicable to a single chromosome. We now want to

approximate a similar process for a sample of n chromosomes from a single sequencing

pool. The true process is quite complicated, and we choose for simplicity to approximate

the process for n chromosomes sampled from one population, as the union of n

independent chromosomal processes. We will later quantify biases arising due to this

independence assumption using simulations. Under the independence assumption, the

transition probability from i to j is simply the probability of l transitions from state 1 to

state 0 in the marginal processes and $j - i + l$ transitions from state 0 to state 1, summed

over all admissible values of l, i.e.,

$$\Pr\left(H_{v+k} = j \mid H_v = i\right) = \sum_{l=\max\{0,i-j\}}^{\min\{n-j,i\}} \binom{n-i}{j-i+l} \left(P_{01}(k)\right)^{j-i+l} \left(1 - P_{01}(k)\right)^{n-j+i-l} \binom{i}{l} \left(P_{10}(k)\right)^{l} \left(1 - P_{10}(k)\right)^{i-l}$$
   (15)

Although this procedure can be computationally expensive when there are many markers,

read depths are high, and especially when n is large, in our implementation, we reduce the

compute time by pre-calculating and storing all binomial coefficients.

Estimating Time Since Admixture

A parameter of central biological interest, that is often unknown in practice, is the time

since the initial admixture event (t). We therefore use the HMM representation to provide

maximum likelihood estimates of t using the forward algorithm to calculate the likelihood

function. As this is a single parameter optimization problem for a likelihood function with a

single mode, optimization can be performed using a simple golden section search [88].

Default settings for this optimization in our software, including the search range maxima

defaults, t_{max} and $t_{min,}$ are documented in the C++ HMM source code provided at

https://github.com/russcd/Ancestry_HMM.

Posterior Decoding

After either estimating or providing a fixed value of the time since admixture to the HMM,

we obtained the posterior distribution for all variable sites considered in our analysis using

the forward-backward algorithm, and we report the full posterior distribution for each

marker along the chromosome.

1003

Simulating Ancestral Polymorphism

To validate our HMM, we generated sequence data for each of two ancestral populations

using the coalescent simulator MACS [55]. We sought to generate data that could be

consistent with that observed in Cosmopolitan and African populations of *D. melanogaster*,

which has been studied previously in a wide variety of contexts [2,11,35-37]. We used the

command line "macs 400 10000000 -i 1 -h 1000 -t 0.0376 -r 0.171 -c 5 86.5 -I 2 200 200 0 -

en 0 2 0.183 -en 0.0037281 2 0.000377 -en 0.00381 2 1 -ej 0.00382 2 1 -eN 0.0145 0.2" to

generate genotype data. This will produce 200 samples of ancestry 0 and 200 samples of

ancestry 1 on a 10mb chromosome—*i.e.* this should resemble genotype data for about half

of an autosomal chromosome arm in *D. melanogaster*. Unless otherwise stated below, we

then sampled the first 50 chromosomes from each ancestral population as the ancestral

population reference panel, whose genotypes are assumed to be known with low error

rates. The sample size was chosen because it is close to the size of the reference panel that

we obtained in our application of this approach to *D. melanogaster* (below).

1018

To evaluate the performance of our method on data consistent with human populations, we

simulated data that could be consistent with that observed for modern European and

African human populations. Specifically, we simulated the model of [89] using the

command line "macs 200 1e8 -I 3 100 100 0 -n 1 1.682020 -n 2 3.736830 -n 3 7.292050 -eg

0 2 116.010723 -eg 1e-12 3 160.246047 -ma x 0.881098 0.561966 0.881098 x 2.797460

0.561966 2.797460 x -ej 0.028985 3 2 -en 0.028986 2 0.287184 -ema 0.028987 3 x

7.293140 x 7.293140 x x x x x -ej 0.197963 2 1 -en 0.303501 1 1 -t 0.00069372 -r

0.00069372". Admixture between ancestral populations was then simulated as described

below.

1028

Simulating Admixed Populations

Although it is commonly assumed that admixture tract lengths can be modeled as

independent and identically distributed exponential random variables (e.g. [26,29] and in

this work, above), this assumption is known to be incorrect as ancestry tracts are neither

exponentially distributed, independent across individuals, nor identically distributed along

chromosomes [57]. We therefore aim to determine what bias violations of this assumption

will have on inferences obtained from this model. Towards this, we used SELAM [56] to

simulate admixed populations under the biological model described above. Because this

program simulated admixture in forward time, it generates the full pedigree-based

ancestral recombination graph, and is therefore a conservative test of our approach

relative to the coalescent which is known to produce incorrect ancestry tract distributions

for short times [57]. Briefly, we initialized each admixed population simulation with a

proportion, m, of ancestry from ancestral population 1, and a proportion 1-m ancestry from

ancestral population 0. Unless otherwise stated, all simulations were conducted with

neutral admixture and a hermaphroditic diploid population of size 10,000.

1044

1045   We then assigned the additional, non-reference chromosomes from the coalescent

1046   simulations, to each ancestry tract produced in SELAM simulations according to their local

1047 ancestry along the chromosome. In this way, each chromosome is a mosaic of the two

1048 ancestral subpopulations. See, *e.g.* [2], for a related approach for simulating genotype data

1049 of admixed chromosomes.

1050

Pruning Ancestral Linkage Disequilibrium

Correlations induced by LD between markers within ancestral populations violates a

central assumption of the Markov model framework. Although it may be feasible to

explicitly model linkage within ancestral populations (*e.g.*, [24,25]), when ancestral

populations have relatively little LD, such as those of *D. melanogaster*, another effective

approach is to discard sites that are in strong LD in the ancestral populations. Hence, to

avoid this potential confounding aspect of the data, we first computed LD between all pairs

of markers within each reference panel that are within 0.01 centimorgans of one another.

We then discarded one of each pair of sites where $|r|$ in either reference panel exceeded a

particular threshold, and we decreased this threshold until we obtained an approximately

unbiased estimate of the time since admixture estimates of the HMM. This approach differs

from a previous method, WinPop [34], where LD is pruned from within admixed samples

(see also below).

1064

Simulating Sequence Data

We first identified all sites where the allele frequencies of the ancestral populations differ

by at least 20% within the reference panels. We excluded weakly differentiated sites to

decrease runtime and because these markers carry relatively little information about the

LA at a given site. Then, to generate data similar to what would be produced using Illumina

sequencing platforms, we simulated allele counts for each sample, by first drawing the

depth at a given site from a Poisson distribution. In most cases and unless otherwise stated,

the mean of this distribution is set to be equal to the sample ploidy. We did this to ensure

equivalent sequencing depth per chromosome regardless of pooling strategy, and because

this depth is sufficiently low that high quality genotypes cannot be determined. We then

generated set of simulated aligned bases via binomial sampling from the sample allele

frequency and included a uniform error rate of 1% for both alleles at each site.

1077

Unless otherwise stated, we simulated a total of 40 admixed chromosomes. The HMM can

perform LAI on more than one sample at a time, and we therefore included all samples

when running it. Hence, we used 40 haploid, 20 diploid, 4 pools of 10 chromosomes, and 2

pools of 20 chromosomes for most comparisons of accuracy reported below, unless

otherwise stated. It is worth noting that it is possible to jointly analyze distinct samples

from the same subpopulation that have been sequenced at different ploidies.

1084

Simulating Divergent Ancestral Populations

To investigate the effects of allele frequency differences between reference populations and

admixed populations, we performed coalescent simulations using the software MACS [55],

using the command line "./macs 500 10000000 -i 1 -h 1000 -t 0.0376 -r 0.171 -c 5 86.5 -I 8

100 100 50 50 50 50 50 50 0 -en 0 2 0.183 -en 0.0037281 2 0.000377 -en 0.00381 2 1 -ej

0.00382 2 1 -eN 0.0145 0.2 -ej 0.0005 3 2 -ej 0.000500001 4 1 -ej 0.001 5 2 -ej

0.001000001 6 1 -ej 0.002 7 2 -ej 0.002000001 8 1". This might be expected to produce

populations that are differentiated similarly to how populations of *D. melanogaster* would

be across European populations or between populations in Central Africa. We then

substituted the increasingly divergent populations for the reference panel. All allele

frequency differences and LD pruning were performed as described above on each of the

substitute reference panels.

1097

Accuracy Statistics

To evaluate the performance of the HMM, we computed four statistics. First, we compute

the proportion of sites where the true state is within the 95% posterior credible interval,

where ideally, this proportion would be equal to or greater than 0.95. As this HMM has

discrete states, there are many ways the 95% credible interval could be defined. In light of

the fact that the credible interval tends to be narrow (Results), we defined the interval to

include all states that are overlapped, by any amount, in the 95% confidence interval of the

posterior distribution. Second, we compute the mean posterior error, the average distance

between the posterior distribution of hidden states and the true state

1107

1108
$$E = \frac{\sum_{v=0}^{S}\sum_{i=0}^{n} p(H_v = i \mid \mathbf{r})|i - I_v|}{Sn}$$

1109

1110 Here S is the total number of sites, I_v is the true state at site v, and $\mathbf{r}$ is all the combined read

1111 data. Third, we also report the proportion of sites where the maximum likelihood estimate

1112 of the hidden state is equal to the true ancestry state. Finally, as an indicator of the

1113 specificity of our approach, we also report the average width of the 95% credible interval.

1114

Deviations from the Assumed Neutral Demographic Model

A potential issue with this framework is that the assumptions underlying the transition matrixes and related time of admixture estimation procedure is likely to be violated in a number of biologically relevant circumstances. We therefore simulated populations wherein individuals of ancestral population 1 began entering a population entirely composed of individuals from ancestral population 0, at a time t generations before the present, at a constant rate that is sustained across all subsequent generations until the time of sampling. That is, additional unadmixed individuals of ancestry 1 migrate each generation from t until the present.

Natural selection acting on admixed genetic regions has been inferred in a wide variety of systems (*e.g.* [5,7,13,17,18]), and is expected to have pronounced effects on the distribution of LA among individuals within admixed populations. Here again, this aspect of biologically realistic populations will tend to violate central underlying assumptions of the model assumed in this work. Towards this, we simulated admixed populations with a single pulse of admixture t generations prior to the time of sampling. We then incorporated selection at 2,5,10, and 20 loci at locations uniformly distributed along the length of the chromosome arm. All selected loci were assumed to be fixed within each ancestral population. Selection was additive and selective coefficients were assigned based on a uniform [0.005, 0.05] distribution to either ancestry 0 or 1 alleles with equal probability. As above, these simulations were conducted using SELAM [56].

1137 For both selected and continuous migration simulations, we then performed the genotype

1138 and read data simulation procedure, and reran our HMM as described above. We

1139 performed 10 simulations for each treatment.

1140

Comparisons to WinPop

We next sought to compare our method to a commonly used local ancestry inference

method, WinPop [34]. Towards this, we again simulated data using MACS and SELAM as

described above. For these comparisons, the initial ancestry contribution was 0.5 and the

number of generations since admixture varied between 5 and 1000. For comparison, we

supplied WinPop and our program the correct time since admixture and ancestry

proportions, as these are required parameters for WinPop. We also supplied the program

with genotypes rather than read counts, another requirement of WinPop, whereas we

supplied our HMM with read data simulated as described above. We then ran WinPop

under default parameters, and we also reran WinPop using LD pruning within the

reference panels, as we do in our method, instead of the default LD pruning implemented in

WinPop.

1153

Analysis of Inuit Genotype Data

To demonstrate that LAI methods can be biased by the arbitrary selection of the time since

admixture, we analyzed a dataset of SNP-array genotype data from Greenlandic Inuits.

These data are described in detail elsewhere [60,61]. This population has received some

admixture from a European source population, and the authors had previously used RFMix

[24] to perform LAI, and found some sensitivity to the assumed time since admixture (J.

Crawford *pers. Comm.*). We analyzed data from chromosome 10 using RFMix v1.5.4 [24] as

described in Moltke *et al.* [61] assuming admixture occurred either 5 or 20 generations ago.

We subsequently analyzed chromosome 10 using our HMM including the genotype-

analysis emissions probabilities and assuming a genotype error rate of 0.2%. For our

analysis we identified the LD cutoff that is appropriate for these data as described above.

1165

Generating *D. melanogaster* Reference Populations

To generate reference panels, we used a subset of the high quality *D. melenaogaster*

assemblies that have been described previously in Pool *et al.* (2012) and Lack *et al.* (2015).

As in the local ancestry analysis of Pool (2015), we used the French population. For our

African reference panel, we selected a subset of the Eastern and Western African

populations (CO, RG, RC, NG, UG, GA, GU) and grouped them into a single population for the

purposes of our analysis. We elected to combine populations so that we would have a

larger reference panel of African populations for this analysis, this solution may be justified

because these *D. melanogaster* populations are only weakly genetically differentiated

[2,21,90], particularly after common inversion-bearing chromosomes are removed from

analyses. Specific individuals were selected for inclusion in the African reference panel if

previous work found they have relatively little cosmopolitan ancestry (*i.e.,* below 0.2

genome-wide in [2]).

1179

Because of their powerful effects on recombination, chromosomal inversions are known to

have substantial impacts on the distribution of genetic variants on chromosomes

containing chromosomal inversions in *D. melanogaster* [2,63]. For this reason, we removed

all common inversion-bearing chromosome arms from the reference populations [91].

Nonetheless, it is clear that chromosomal inversions are present in the pool-seq samples

[64]. Although the inversions certainly violate key assumptions of our model—particularly

the transmission probabilities—given that our approach is robust to a many perturbations,

we expect the LA within inverted haplotypes can be estimated with reasonable confidence,

and the overall LAI procedure will still perform adequately with low frequencies of

chromosomal-inversion bearing chromosomes present within these samples.

1190

Although these reference populations are believed to have relatively little admixture, some

admixture is likely to remain within these samples [2]. To mitigate this potential issue, we

first applied our HMM to each reference population using the genotype-based emissions

probabilities (above). Calculated across all individuals, we found that our maximum

likelihood ancestry estimates were identical with those of Pool *et al.* (2012) at 96.2% of

markers considered in our analysis. The differences between the results of these methods

may reflect differences in the methodology of LAI or differences in the reference panels.

Nonetheless, the broad concordance suggests the two methods are yielding similar overall

results. We masked all sites where the posterior probability of non-native ancestry was

greater than 0.5 within each reference individual's genome. These masked sequences were

then used as the reference panel for the analyses of pool-seq data below.

1202

Ancestry Cline Sequence Data Analysis

We acquired pooled sequencing data from six populations from the east coast of the United

States. The generation of these samples, sequencing data, and accession numbers are

described in detail in [21,40]. Briefly, the samples are comprised of individuals drawn from

natural populations and sequenced in relatively large pools of 66-232 chromosomes. We

aligned all data using BWA v0.7.9a-r786 [92] using the 'MEM' function and the default

program parameters. For all alignments, we used version 5 of the *D. melanogaster*

reference genome [93] in order to make our analysis and coordinates compatible with the

Drosophila genome nexus [91]. We then realigned all reads using the indelrealigner tool

within the GATK package [84], and we extracted the sequence pileup using samtools

mpileup v1.1 [94] using the program's default parameters.

1214

We extracted sites at ancestry informative positions within the reference panels, where we

required that the reference panel have a minimum of 50% of individuals with a high quality

genotype call in both Cosmopolitan and African reference populations. As above, ancestry

informative sites were defined as those with a minimum of 20% difference in allele

frequencies between the reference panels used, and we retained only ancestry informative

sites for our analyses. We then produced global ancestry estimates for each chromosome

arm separately for each sample using the method of Bergland *et al.* (2016). We ran our

HMM for each chromosome arm and each population, and we provided the program this

estimate of the ancestry proportion and the time since admixture, 1593 generations [17].

We elected to provide the time since admixture because we have found that this parameter

is difficult to estimate in relatively large pools (see Results). However, the program can

accurately estimate LA in high ploidy samples even when the time since admixture cannot

be estimated correctly (see Results).

1228

Correlation with Local Recombination Rates

To assess the correlation between local recombination rates and LA in the genome, we

computed Spearman's rank sum correlation between the proportions African ancestry and

the local recombination rates in windows of 100 ancestry informative markers. As above,

we used the recombination rate estimates of [59]. We estimated confidence intervals using

1000 block-bootstrap samples using window sizes of 100 SNPs.

Robustness of LAI to Genomic Heterogeneity

To determine if there are systematic biases in LAI across the genome, we computed the

mean difference in genomic windows between LA estimates for two samples form Maine

and between two samples from Florida. We assessed evidence for systematic biases

through the correlation between local recombination rates and differences in local ancestry

inference using Spearman's rank sum correlation.

Identifying LA Cline Outliers

To detect loci that show evidence for steeper ancestry clines than the genomic average, we

first computed the Spearman's rank correlation between mean ancestry proportions and

latitude for each chromosome arm separately. Then, for each site for which we obtained a

posterior ancestry distribution for all samples, we computed the partial Spearman's rank

correlation between the posterior ancestry mean and latitude while correcting for the

correlation between latitude and the overall ancestry proportion. We then computed the

probability of obtaining the observed partial correlation in R, which implements the

approach of [95], and we retained those sites where the probability of the partial

correlation between local ancestry and latitude was less than 0.005 as significant in our

analysis. Although this cutoff is arbitrary, given the strong evidence for local adaptation

and reproductive isolation in these populations [46,47,96], the tail of the LA cline

distribution will likely be enriched for sites experiencing selection on this ancestry

gradient. Due to linkage, adjacent sites show strong autocorrelation. We therefore selected

the local optima for a given clinally significant LA segment (*i.e.* a tract where all positions

are significantly correlated with latitude at our threshold) and retained these for analyses

of outlier loci. Finally, to further reduce the effect of autocorrelation, we retained only

those local optima for which no other optimum had a stronger correlation with latitude

within 100,000bp on either side on the site.

1262

1263 **Identifying Low African Ancestry Outlier Loci**

1264 To identify loci with a disproportionately low proportion of African ancestry across this

1265 ancestry cline, we computed the mean African ancestry across all populations. We then

1266 selected those sites in the lowest 5% tail on each chromosome arm and selected only the

1267 local minima within 100kb windows on either side of a selected locus.

1268

Gene Ontology Analyses

We performed Gene-ontology (GO) analyses on outlier SNPs using Gowinda [97], where the

background set of SNPs was all positions at which we obtained a posterior distribution in

all samples (i.e. the set on which we obtained estimates of the posterior probability of

African ancestry). We ran the program using default parameters, except that we included

all genes within 10000bp of a focal SNP, and we performed 1e6 total GO simulations.

1275

1276 **Seasonality of LA in the Pennsylvania Populations**

1277 To identify recurrent seasonal changes in the local ancestry, we followed an approach

1278 similar to [40]. Specifically, we fit a generalized linear model of the form

1279

1280 $$Mean\ Posterior\ Ancestry \sim Season + \varepsilon$$

1281

1282 We then recorded the estimated effect size, and probability of the observed correlation for

1283 each site in the genome at which we obtained a posterior ancestry distribution in all

1284 samples considered. To correct for multiple testing, we applied a false discovery rate

1285 correction [81] to the resulting p-value distribution.

1286

1287 **Acknowledgements**

1288 We thank three anonymous reviewers, Tyler Linderoth, Shelbi Russell and members of the

1289 Nielsen and Slatkin labs for helpful comments. We also thank Jacob Crawford for providing

1290 Inuit polymorphism data and for providing the observation that LAI inference depended

1291 heavily on *t.*

1292

References

1. Kronforst MR, Young LG, Blume LM, Gilbert LE. Multilocus analyses of admixture and
introgression among hybridizing *Heliconius* butterflies. Evolution. 2006;60: 1254–16.
doi:10.1554/06-005.1

2. Pool JE, Corbett-Detig RB, Sugino RP, Stevens KA, Cardeno CM, Crepeau MW, et al.
Population genomics of sub-Saharan Drosophila melanogaster: African diversity and
non-African admixture. PLoS Genet. 2012;8: e1003080–24.
doi:10.1371/journal.pgen.1003080

3.   Hufford MB, Lubinksy P, Pyhäjärvi T, Devengenzo MT, Ellstrand NC, Ross-Ibarra J.
The genomic signature of crop-wild introgression in maize. PLoS Genet. 2013;9:
e1003477–13. doi:10.1371/journal.pgen.1003477

4.   Brandvain Y, Kenney AM, Flagel L, Coop G, Sweigart AL. Speciation and introgression
between Mimulus nasutus and Mimulus guttatus. PLoS Genet. 2014;10: e1004410–
15. doi:10.1371/journal.pgen.1004410

5.   Rieseberg LH, Raymond O, Rosenthal DM, Lai Z, Livingstone K, Nakazato T, et al.
Major ecological transitions in wild sunflowers facilitated by hybridization. Science.
2003;301: 1211–1216. doi:10.1126/science.1086949

6.   Green RE, Krause J, Briggs AW, Maricic T, Stenzel U, Kircher M, et al. A draft sequence
of the neandertal genome. Science. 2010;328: 710–722.
doi:10.1126/science.1188021

7.   Sankararaman S, Mallick S, Dannemann M, Prüfer K, Kelso J, Pääbo S, et al. The
genomic landscape of Neanderthal ancestry in present-day humans. Nature.
2014;507: 354–357. doi:10.1038/nature12961

8.   Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal
components analysis corrects for stratification in genome-wide association studies.
Nat Genet. 2006;38: 904–909. doi:10.1038/ng1847

9.   Lohmueller KE, Pearce CL, Pike M, Lander ES, Hirschhorn JN. Meta-analysis of genetic
association studies supports a contribution of common variants to susceptibility to
common disease. Nat Genet. 2003;33: 177–182. doi:10.1038/ng1071

10.   Freedman ML, Reich D, Penney KL, McDonald GJ, Mignault AA, Patterson N, et al.
Assessing the impact of population stratification on genetic association studies. Nat
Genet. 2004;36: 388–393. doi:10.1038/ng1333

11.   Caracristi G, Schlotterer C. Genetic differentiation between American and European
Drosophila melanogaster populations could be attributed to admixture of African
alleles. Mol Biol Evol. 2003;20: 792–799. doi:10.1093/molbev/msg091

12.   Kolbe JJ, Glor RE, Schettino LR, Lara AC, Larson A. Genetic variation increases during
biological invasion by a Cuban lizard. Nature. 2004;431: 177–181.

13.   Consortium THG, Consortium G. Butterfly genome reveals promiscuous exchange of
mimicry adaptations among species. Nature. 2012;487: 94–98.
doi:10.1038/nature11041

14.   Racimo F, Sankararaman S, Nielsen R, Huerta-Sanchez E. Evidence for archaic
adaptive introgression in humans. Nat Genet. 2015;16: 359–371.
doi:10.1038/nrg3936

15.   Juric I, Aeschbacher S, Coop G. The strength of selection against Neanderthal
1337   introgression. bioRxiv. 2015 Oct pp. 1–24. doi:10.1101/030148

1338   16.   Harris K, Nielsen R. The genetic cost of neanderthal introgression. Genetics.
1339   2016;203: 881–891. doi:10.1534/genetics.116.186890

1340   17.   Pool JE. The mosaic ancestry of the Drosophila Genetic Reference Panel and the D.
1341   melanogaster reference genome reveals a network of epistatic fitness interactions.
1342   Mol Biol Evol. 2015;: msv194–16. doi:10.1093/molbev/msv194

1343   18.   Schumer M, Cui R, Powell DL, Dresner R, Rosenthal GG, Andolfatto P. High-resolution
1344   mapping reveals hundreds of genetic incompatibilities in hybridizing fish species.
1345   eLife. 2014;3: 610–21. doi:10.7554/eLife.02535

1346   19.   Pritchard JK, Stephens M, Donnelly P. Inference of population structure using
1347   multilocus genotype data. Genetics. 2000;155: 945–959.

1348   20.   Skotte L, Korneliussen TS, Albrechtsen A. Estimating individual admixture
1349   proportions from next generation sequencing data. Genetics. 2013;195: 693–702.
1350   doi:10.1534/genetics.113.154138/-/DC1

1351   21.   Bergland AO, Tobler R, González J, Schmidt P, Petrov D. Secondary contact and local
1352   adaptation contribute to genome-wide patterns of clinal variation in Drosophila
1353   melanogaster. Mol Ecol. 2016;25: 1157–1174. doi:10.1111/mec.13455

1354   22.   Falush D, Stephens M, Pritchard JK. Inference of population structure using
1355   multilocus genotype data: linked loci and correlated allele frequencies. Genetics.
1356   2003;164: 1567–1587.

1357   23.   Sankararaman S, Sridhar S, Kimmel G, Halperin E. Estimating local ancestry in
1358   admixed populations. Am J Hum Genet. 2008;82: 290–303.
1359   doi:10.1016/j.ajhg.2007.09.022

1360   24.   Maples BK, Gravel S, Kenny EE, Bustamante CD. RFMix: a discriminative modeling
1361   approach for rapid and robust local-ancestry inference. Am J Hum Genet. 2013;93:
1362   278–288. doi:10.1016/j.ajhg.2013.06.020

1363   25.   Baran Y, Pasaniuc B, Sankararaman S, Torgerson DG, Gignoux C, Eng C, et al. Fast and
1364   accurate inference of local ancestry in Latino populations. Bioinformatics. 2012;28:
1365   1359–1367. doi:10.1093/bioinformatics/bts144

1366   26.   Pool JE, Nielsen R. Inference of historical changes in migration rate from the lengths
1367   of migrant tracts. Genetics. 2009;181: 711–719. doi:10.1534/genetics.108.098095

1368   27.   Koopman W, Li Y, Coart E. Linked vs. unlinked markers: multilocus microsatellite
1369   haplotype-sharing as a tool to estimate gene flow and introgression. Mol Ecol.
1370   2007;16: 243–256. doi:10.1111/j.1365-294X.2006.03137.x

28. Patterson N, Hattangadi N, Lane B. Methods for high-density admixture mapping of
disease genes. Am J Hum Genet. 2004;74: 979–1000.

29. Gravel S. Population Genetics Models of Local Ancestry. Genetics. 2012;191: 607–
619. doi:10.1534/genetics.112.139808

30. Moorjani P, Patterson N, Hirschhorn JN, Keinan A, Hao L, Atzmon G, et al. The history
of African gene flow into southern Europeans, Levantines, and Jews. PLoS Genet.
2011;7: e1001373–13. doi:10.1371/journal.pgen.1001373

31. Loh PR, Lipson M, Patterson N, Moorjani P. Inferring admixture histories of human
populations using linkage disequilibrium. Genetics. 2013;193: 1233–1254.
doi:10.1534/genetics.112.147330/-/DC1

32. Hellenthal G, Busby GBJ, Band G, Wilson JF, Capelli C, Falush D, et al. A Genetic Atlas
of Human Admixture History. Science. 2014;343: 747–751.
doi:10.1126/science.1243518

33. Consortium 1GP, BGI-Shenzhen, European Bioinformatics Institute, Illumina,
Hospital BAW, College B, et al. An integrated map of genetic variation from 1,092
human genomes. Nature. 2012. doi:10.1038/nature11632

34. Pasaniuc B, Sankararaman S, Kimmel G, Halperin E. Inference of locus-specific
ancestry in closely related populations. Bioinformatics. 2009;25: i213–i221.
doi:10.1093/bioinformatics/btp197

35. Haddrill PR, Thornton KR, Charlesworth B, Andolfatto P. Multilocus patterns of
nucleotide variability and the demographic and selection history of Drosophila
melanogaster populations. Genome Res. 2005;15: 790–799. doi:10.1101/gr.3541005

36. Thornton K, Andolfatto P. Approximate Bayesian inference reveals evidence for a
recent, severe bottleneck in a Netherlands population of Drosophila melanogaster.
Genetics. 2006;172: 1607–1619. doi:10.1534/genetics.105.048223

37. Li H, Stephan W. Inferring the demographic history and rate of adaptive substitution
in Drosophila. PLoS Genet. 2006;2: e166. doi:10.1371/journal.pgen

38. Duchen P, Živković D, Hutter S, Stephan W. Demographic inference reveals African
and European admixture in the North American Drosophila melanogaster
population. Genetics. 2013;193: 291–301. doi:10.1534/genetics.112.145912/-/DC1

39. Langley CH, Stevens K, Cardeno C, Lee YCG, Schrider DR, Pool JE, et al. Genomic
variation in natural populations of Drosophila melanogaster. Genetics. 2012;192:
533–598. doi:10.1534/genetics.112.142018

40. Bergland AO, Behrman EL, O'Brien KR, Schmidt PS, Petrov DA. Genomic evidence of
rapid and stable adaptive oscillations over seasonal time scales in Drosophila. PLoS

Genet. 2014;10: e1004775–19. doi:10.1371/journal.pgen.1004775

41.   Schrider DR, Hahn MW, Begun DJ. Parallel evolution of copy-number variation across
continents in Drosophila melanogaster. Mol Biol Evol. 2016;33: 1308–1316.
doi:10.1093/molbev/msw014

42.   Reinhardt JA, Kolaczkowski B, Jones CD, Begun DJ, Kern AD. Parallel geographic
variation in Drosophila melanogaster. Genetics. 2014;197: 361–373.
doi:10.1534/genetics.114.161463/-/DC1

43.   Fabian DK, Kapun M, Nolte V, Kofler R, Schmidt PS, Schlotterer C, et al. Genome-wide
patterns of latitudinal differentiation among populations of Drosophila
melanogasterfrom North America. Mol Ecol. 2012;21: 4748–4769.
doi:10.1111/j.1365-294X.2012.05731.x

44.   Oakeshott JG, Gibson JB, Anderson PR, Knibb WR. Alcohol dehydrogenase and
glycerol-3-phosphate dehydrogenase clines in Drosophila melanogaster on different
continents. Evolution. 1982;36: 86–96.

45.   Berry A, Kreitman M. Molecular analysis of an allozyme cline: alcohol dehydrogenase
in Drosophila melanogaster on the east coast of North America. Genetics. 1993;134:
869–893.

46.   Yukilevich R, True JR. African morphology, behavior, and phermones underlie
incipient sexual isolation between US and Caribbean Drosophila melanogaster.
Evolution. 2008;62: 2807–2828. doi:10.1111/j.1558-5646.2008.00488.x

47.   Lachance J, True JR. X-autosome incompatibilities in Drosophila melanogaster: test of
Haldane's rue and geographic patterns within species. Evolution. 2010;64: 3035–
3046. doi:10.1111/j.1558-5646.2010.01028.x

48.   Corbett-Detig RB, Zhou J, Clark AG, Hartl DL, Ayroles JF. Genetic incompatibilities are
widespread within species. Nature. 2013;504: 135–137. doi:10.1038/nature12678

49.   Langley CH, Crepeau M, Cardeno C, Corbett-Detig R, Stevens K. Circumventing
heterozygosity: sequencing the amplified genome of a single haploid Drosophila
melanogaster embryo. Genetics. 2011;188: 239–246.
doi:10.1534/genetics.111.127530

50.   Kofler R, Betancourt AJ, Schlotterer C. Sequencing of Pooled DNA Samples (Pool-Seq)
Uncovers Complex Dynamics of Transposable Element Insertions in Drosophila
melanogaster. PLoS Genet. 2012;8: e1002487–16.
doi:10.1371/journal.pgen.1002487

51.   terWengel PO, Kapun M, Nolte V, Kofler R, Flatt T, Schlotterer C. Adaptation of
Drosophila to a novel laboratory environment reveals temporally heterogeneous
trajectories of selected alleles. Mol Ecol. 2012;21: 4931–4941. doi:10.1111/j.1365-

294X.2012.05673.x

52.   Kolaczkowski B, Kern AD, Holloway AK, Begun DJ. Genomic differentiation between
temperate and tropical Australian populations of Drosophila melanogaster. Genetics.
2011;187: 245–260. doi:10.1534/genetics.110.123059

53.   Kapun M, Schalkwyk H, McAllister B, Flatt T, Schlotterer C. Inference of chromosomal
inversion dynamics from Pool-Seq data in natural and laboratory populations of
Drosophila melanogaster. Mol Ecol. 2014;23: 1813–1827. doi:10.1111/mec.12594

54.   Zhu Y, Bergland AO, González J, Petrov DA. Empirical validation of pooled whole
genome population re-sequencing in Drosophila melanogaster. PLoS ONE. 2012;7:
e41901. doi:10.1371/journal.pone.0041901

55.   Chen GK, Marjoram P, Wall JD. Fast and flexible simulation of DNA sequence data.
Genome Res. 2008;19: 136–142. doi:10.1101/gr.083634.108

56.   Corbett-Detig R, Jones M. SELAM: simulation of epistasis and local adaptation during
admixture with mate choice. Bioinformatics. 2016;btw365.

57.   Liang M, Nielsen R. The lengths of admixture tracts. Genetics. 2014;197: 953–967.
doi:10.1534/genetics.114.162362

58.   Sankararaman S, Patterson N, Li H, Pääbo S, Reich D. The Date of Interbreeding
between Neandertals and Modern Humans. PLoS Genet. 2012;8: e1002947–9.
doi:10.1371/journal.pgen.1002947

59.   Comeron JM, Ratnappan R, Bailin S. The many landscapes of recombination in
Drosophila melanogaster. PLoS Genet. 2012;8: e1002905.
doi:10.1371/journal.pgen.1002905

60.   Moltke I, Grarup N, Jørgensen ME, Bjerregaard P, Treebak JT, Fumagalli M, et al. A
common Greenlandic TBC1D4 variant confers muscle insulin resistance and type 2
diabetes. Nature. 2014;512: 190–193. doi:10.1038/nature13425

61.   Moltke I, Fumagalli M, Korneliussen TS. Uncovering the genetic history of the
present-day Greenlandic population. Am J Hum Genet. 2015;96: 54–69.
doi:10.1016/j.ajhg.2014.11.012

62.   Voight BF, Kang HM, Ding J, Palmer CD, Sidore C, Chines PS, et al. The metabochip, a
custom genotyping array for genetic studies of metabolic, cardiovascular, and
anthropometric traits. PLoS Genet. 2012;8: e1002793–12.
doi:10.1371/journal.pgen.1002793

63.   Corbett-Detig RB, Hartl DL. Population genomics of inversion polymorphisms in
Drosophila melanogaster. 2012;8: e1003056–15. doi:10.1371/journal.pgen.1003056

64.　Kapun M, Fabian DK, Goudet J, Flatt T. Genomic evidence for adaptive inversion
clines in Drosophila melanogaster. Mol Biol Evol. 2016;33: 1317–1336.
doi:10.1093/molbev/msw016

65.　Mackay TFC, Richards S, Stone EA, Barbadilla A, Ayroles JF, Zhu D, et al. The
Drosophila melanogaster Genetic Reference Panel. Nature. 2012;482: 173–178.
doi:10.1038/nature10811

66.　Huang W, Massouras A, Inoue Y, Peiffer J, Ramia M, Tarone AM, et al. Natural
variation in genome architecture among 205 Drosophila melanogaster Genetic
Reference Panel lines. Genome Res. 2014;24: 1193–1208.
doi:10.1101/gr.171546.113

67.　Corbett-Detig RB, Cardeno C, Langley CH. Sequence-based detection and breakpoint
assembly of polymorphic inversions. Genetics. 2012;192: 131–137.
doi:10.1534/genetics.112.141622

68.　Kulathinal RJ, Stevison LS, Noor MAF. The genomics of speciation in Drosophila:
diversity, divergence, and introgression estimated using low-coverage genome
sequencing. PLoS Genet. 2009;5: e1000550–7. doi:10.1371/journal.pgen.1000550

69.　Krimbas CB, Powell JR. Drosophila Inversion Polymorphism. CRC Press; 1992.

70.　Aulard S, David JR, Lemenieur F. Chromosomal inversion polymorphism in
Afrotropical populations of *Drosophila melanogaster*. Genet Res. 2002;79: 49–63.
doi:10.1017/S0016672301005407

71.　Pasaniuc B, Sankararaman S, Torgerson DG, Gignoux C, Zaitlen N, Eng C, et al.
Analysis of Latino populations from GALA and MEC studies reveals genomic loci with
biased local ancestry estimation. Bioinformatics. 2013;29: 1407–1415.

72.　Ting CT, Takahashi A, Wu CI. Incipient speciation by sexual isolation in Drosophila:
concurrent evolution at multiple loci. PNAS. 2001;98: 6709–6713.

73.　Hollocher H, Ting CT, Wu ML, Wu CI. Incipient speciation by sexual isolation in
Drosophila melanogaster: extensive genetic divergence without reinforcement.
Genetics. 1997;147: 1191–1201.

74.　Hollocher H, Ting CT, Pollack F, Wu CI. Incipient speciation by sexual isolation in
Drosophila melanogaster: variation in mating preference and correlation between
sexes. Evolution. 1997;51: 1175–1181.

75.　Coyne JA. Genetics and speciation. Nature. 1992;335: 511–515.

76.　Coyne JA, Orr HA. Patterns of speciation in Drosophila. Evolution. 1989;43: 362–381.

77.　Charlesworth B, Coyne JA, Barton NH. The relative rates of evolution of sex

chromosomes and autosomes. American Naturalist. 1987;130: 113–146.

78.   Levine MT, Begun DJ. Evidence of Spatially Varying Selection Acting on Four
1512   Chromatin-Remodeling Loci in Drosophila melanogaster. Genetics. 2008;179: 475–
1513   485. doi:10.1534/genetics.107.085423

1514   79.   Ellis LL, Carney GE. Socially-Responsive Gene Expression in Male Drosophila
1515   melanogaster Is Influenced by the Sex of the Interacting Partner. Genetics. 2011;187:
1516   157–169. doi:10.1534/genetics.110.122754

1517   80.   Edwards AC, Zwarts L, Yamamoto A, Callaerts P, Mackay TF. Mutations in many genes
1518   affect aggressive behavior in Drosophila melanogaster. BMC Biol. 2009;7: 29–13.
1519   doi:10.1186/1741-7007-7-29

1520   81.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and
1521   powerful approach to multiple testing. Journal of the Royal Statistical Society.
1522   1995;57: 289–300.

1523   82.   Nielsen R, Korneliussen T, Albrechtsen A, Li Y, Wang J. SNP calling, genotype calling,
1524   and sample allele frequency estimation from new-generation sequencing data. PLoS
1525   ONE. 2012;7: e37558–11. doi:10.1371/journal.pone.0037558

1526   83.   Fumagalli M, Vieira FG, Linderoth T, Nielsen R. ngsTools: methods for population
1527   genetics analyses from next-generation sequencing data. Bioinformatics. 2014;30:
1528   1486–1487.

1529   84.   DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework
1530   for variation discovery and genotyping using next-generation DNA sequencing data.
1531   Nat Genet. 2011;43: 491–498. doi:10.1038/ng.806

1532   85.   Baird SJE, Barton NH, Etheridge AM. The distribution of surviving blocks of an
1533   ancestral genome. Theoretical Population Biology. 2003;64: 451–471.
1534   doi:10.1016/S0040-5809(03)00098-4

1535   86.   Henn BM, Botigué LR, Gravel S, Wang W, Brisbin A, Byrnes JK, et al. Genomic ancestry
1536   of north Africans supports back-to-Africa migrations. PLoS Genet. 2012;8:
1537   e1002397–11. doi:10.1371/journal.pgen.1002397

1538   87.   Marjoram P, Wall JD. Fast "coalecsent" simulation. BMC Genet. 2006;7: 16–9.
1539   doi:10.1186/1471-2156-7-16

1540   88.   Kiefer J. Sequential minimax search for a maximum. Proceedings of the American
1541   Mathematical Society. 1953;4: 502–506.

1542   89.   Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD. Inferring the joint
1543   demographic history of multiple populations from multidimensional SNP frequency
1544   data. PLoS Genet. 2009;5: e1000695–11. doi:10.1371/journal.pgen.1000695

90. Pool JE, Aquadro CF. History and Structure of Sub-Saharan Populations of Drosophila
melanogaster. Genetics. 2006;174: 915–929. doi:10.1534/genetics.106.058693

91. Lack JB, Cardeno CM, Crepeau MW, Taylor W, Corbett-Detig RB, Stevens KA, et al. The
Drosophila Genome Nexus: a population genomic resource of 623 Drosophila
melanogaster genomes, including 197 from a single ancestral range population.
Genetics. 2015;199: 1229–1241. doi:10.1534/genetics.115.174664/-/DC1

92. Li H. Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM. arXiv preprint arXiv:13033997. 2013.

93. Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, Amanatides PG. The genome
sequence of Drosophila melanogaster. Science. 2000;287: 2185–2195.

94. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–2079.
doi:10.1093/bioinformatics/btp352

95. Best DJ, Roberts DE. Algorithm AS 89: the upper tail probabilities of Spearman's rho.
Journal of the Royal Statistical Society Series C. 1975;24: 377–379.

96. Kao JY, Lymer S, Hwang SH, Sung A, Nuzhdin SV. Postmating reproductive barriers
contribute to the incipient sexual isolation of the United States and Caribbean
Drosophila melanogaster. Ecol Evol. 2015;5: 3171–3182. doi:10.1002/ece3.1596

97. Kofler R, Schlötterer C. Gowinda: unbiased analysis of gene set enrichment for
genome-wide association studies. Bioinformatics. 2012;28: 2084–2085.
doi:10.1093/bioinformatics/bts315

1566

Figure Legends

Figure 1. The effect of increasing stringency with ancestral LD pruning. From left to right,

ancestry proportions are 0.1, 0.25, 0.5, 0.75 and 0.9. |r| cutoffs are: none (red), 1.0 (orange),

0.9 (yellow), 0.8 (green), 0.7 (dark blue), 0.6 (cyan), 0.5 (indigo), and 0.4 (violet). The solid

line indicates the expectation for unbiased time estimation. All read data were simulated

with ploidy = 1. True admixture time was drawn from a uniform (0, 2000) distribution.

1573

Figure 2. Time estimates and accuracy statistics for samples of varying ploidies. From left

to right, ancestry proportions are 0.1, 0.25, 0.5, 0.75 and 0.9. Each sample ploidy is

represented by one point color with ploidy one (black), two (red), ten (blue) and twenty

(green). From top to bottom, each row is the estimated time in generations, the proportion

of sites where the true state is within the 95% credible interval, the width of the 95%

credible interval, the mean posterior error, and the proportion of sites where the maximum

likelihood estimate is equal to the true state.

1581

1582 **Figure 3.** Accuracy of the HMM for samples of high ploidy. The 95% credible interval

1583 (shaded blue region), and the posterior mean (red) contrasted with the true ancestry

1584 frequencies (black). Simulated data were generated with an admixture time of 1500

1585 generations, an ancestry proportion of 0.2, a sample ploidy of 100, and a mean sequencing

1586 depth of 25.

1587

1588 **Figure 4.** Admixture time estimates for simulated data consistent with variation present in

1589 modern European and African populations. From left to right, m = 0.1, m = 0.25, m = 0.5, m

1590 = 0.75, m = 0.9. The top row is completely phased chromosomes and the bottom row is

1591 unphased diploid data.

1592

1593 **Figure 5.** Local ancestry of inversion bearing chromosomes (red) compared with those of

1594 standard arrangement chromosomes (black) for the same chromosome arm. Positions of

1595 inversion breakpoints, as reported in [63,67] are shown as vertical dashed lines.

1596

Figure 6. The relationship between the proportion of African ancestry proportion and local

recombination rates in 100 ancestry informative SNP windows within the Raleigh, NC

population (left). The correlation between the proportion of African ancestry proportion

and local recombination rates in 100 ancestry informative SNP windows in all populations

assayed (right). Lines indicate the 95% confidence interval obtained via block bootstrap

replicates (see Methods).

1603

1604   **Figure 7.** The partial correlation between LA and latitude with correction for chromosome-

1605   wide ancestry proportions. Sites for which the probability of the observed clinal

1606   relationship was less than 0.005 were retained as significant (red). Inversion breakpoints

1607   for inversions that are at polymorphic frequencies on this ancestry cline are shown as

1608   dotted blue lines.

1609

1610   **Figure 8.** The mean African ancestry proportion across all populations on the ancestry

1611   cline for chromosome arms 2L, 2R, 3L, 3R, and X (top to bottom). Local minima outlier loci

1612   are shown in red (see Methods).

1613

1614   **Supplemental Figure S1.** Comparison between LAI using the full ancestral recombination

1615   graph via forward-time simulations (red) with those from independent and identically

1616   distributed draws from the SMC' distribution (black). Simulations were conducted using an

1617   ancestry proportion of 0.25 and population size of 10,000 hermaphroditic individuals.

1618

Supplemental Figure S2. Effects of unknown admixed population sizes on LAI. All LAI was

conducted assuming the true population size was 10,000. Simulated population sizes were

100 (black), 1,000 (red), 10,000 (blue) and 100,000 (green). Ploidy 1 on the right, ploidy 2

on the left. From top to bottom, rows are the estimated time of admixture, the proportion of

sites where the true state is within the 95% credible interval, the width of the 95% credible

interval, the mean posterior error, and the proportion of times that the maximum

likelihood estimate is equal to the true state. For all simulations, the ancestry proportion

was equal to 0.5.

1627

Supplemental Figure S3. LAI accuracy when admixture times are increasingly ancient.

Here, ancestry proportions are 0.5 (black), 0.25 (blue), 0.1 (violet), 0.75 (orange) and 0.9

(red). From top to bottom, statistics plotted are estimated time, the proportion of sites

where the true ancestry frequency is within the 95% credible interval, the mean 95%

credible interval width, mean posterior error, and the proportion of times that the

maximum likelihood estimate is correct.

1634

Supplemental Figure S4. The effects of reference panel size on LAI and time estimation

using the HMM. Here, we compare reference panels of size 100 (blue) with reference

panels of size 10 (black). From left to right, ancestry proportions are 0.1, 0.25, 0.5, 0.75 and

0.9. From top to bottom the plotted statistics are estimated time, proportion in the 95%

credible interval, the average width of the 95% credible interval, the mean posterior error,

and the proportion of sites where the maximum likelihood ancestry estimate is correct.

1641

Supplemental Figure S5. Accuracy of time estimation and LAI when reference populations

are increasingly divergent from the source of the admixture pulses. In columns are

divergence times between ancestral populations (in units of 4Ne) of 0, 0.0005, 0.001, 0.002.

From top to bottom the plotted statistics are estimated time, proportion in the 95%

credible interval, the average width of the 95% credible interval, the mean posterior error,

and the proportion of sites where the maximum likelihood ancestry estimate is correct.

1648

Supplemental Figure S6. Comparison of the proportion of sites where the maximum

likelihood ancestry estimate of local ancestry is correct between WinPop and our method.

WinPop was run with default parameters (black), and with LD pruned in the ancestral

populations, but not in the admixed population (red). Our method was run with default

parameters (blue), but with the time since admixture and correct ancestry proportion

supplied to our program as these parameters are required by WinPop.

1655

Supplemental Figure S7. Bias in LAI due to uncertainty in t. The posterior probability

estimated using RFMix of European ancestry at a given site in the genome assuming t = 5

(black) and assuming t = 20 (red) for a sample representative of the average difference (top

left) and a more extreme example (top right). The distribution of differences in mean Inuit

ancestry for all samples (bottom left) using RFMix. The log likelihood of each time since

admixture as computed using our method (bottom right), which shows a clear optimum at

6-7 generations since admixture. All analyses were restricted to SNPs on chromosome 10.

1663

Supplemental Figure S8. Bias in LAI and time estimation due to incorrect estimation of m.

On the left, true m is 0.1and on the right true m is 0.5. Supplied m varies across 0.05 to 0.95.

From top to bottom, the plotted statistics are estimated t, proportion in the 95% confidence

interval, mean 95% confidence interval width, mean posterior error and the proportion of

sites where the maximum likelihood estimate is correct. All plots include ploidy one

(back), ploidy two (red), ploidy ten (blue), and ploidy twenty (green).

1670

Supplemental Figure S9. Bias in LAI and time estimation due to incorrect assumptions of

t. On the left, true t is 100 and on the right true t is 1000. Supplied t varies across 100 to

2000 generations. From top to bottom, the plotted statistics are estimated t, proportion in

the 95% confidence interval, mean 95% confidence interval width, mean posterior error

and the proportion of sites where the maximum likelihood estimate is correct. All plots

include ploidy one (back), ploidy two (red), ploidy ten (blue), and ploidy twenty (green).

1677

1678 **Supplemental Figure S10.** Estimates of t obtained from block bootstrap replicates for

1679 populations that have admixed for 1000 (top), and 2000 (bottom) generations. From left to

1680 right, sample ploidies are 1, 2, 10, and 20. For both simulations, $m = 0.5$.

1681

1682 **Supplemental Table S1.** Comparison of run times for various demographic models and

1683 sample ploidies using this method.

1684

1685 **Supplemental Table S2.** LA clinality in the genomic intervals immediately surrounding

1686 breakpoints of known polymorphic inversions.

1687

1688   **Supplemental Table S3.** Results of GO analysis of 80 identified LA clinal outlier loci.

1689

1690   **Supplemental Table S4.** Results of GO analysis of 84 identified low African ancestry

1691   outlier loci.