1

Impact of Sample Type and DNA Isolation Procedure on Genomic

Inference of Microbiome Composition

Berith E. Knudsen[a], Lasse Bergmark[a*], Patrick Munk[a], Oksana Lukjancenko[a], Anders

Priemé[2], Frank M. Aarestrup[a], Sünje J. Pamp[a#]

[a]National Food Institute, Technical University of Denmark, Kgs. Lyngby, Denmark

[b]Department of Biology, University of Copenhagen, Copenhagen, Denmark

B.E.K and L.B. contributed equally to this work.

[*]Present address: Novo Nordisk, Bagsværdvej, 2880 Bagsværd, Denmark

[#]Address correspondence to: Sünje J. Pamp, sjpa@food.dtu.dk

22 Running title: DNA Isolation Methodology for Microbiome Genomics

23 Keywords: microbial ecology, microbiome, DNA isolation, 16S rDNA gene sequencing,

24 metagenomics

ABSTRACT

Explorations of complex microbiomes using genomics greatly enhance our understanding about their diversity, biogeography, and function. The isolation of DNA from microbiome specimens is a key prerequisite for such examinations, but challenges remain in obtaining sufficient DNA quantities required for certain sequencing approaches, achieving accurate genomic inference of microbiome composition, and facilitating comparability of findings across specimen types and sequencing projects. These aspects are particularly relevant for the genomics-based global surveillance of infectious agents and antimicrobial resistance from different reservoirs. Here, we compare a total of eight DNA extraction procedures for three specimen types (human feces, pig feces, hospital sewage), assess DNA extraction using spike-in controls and different types of beads for bead-beating facilitating cell lysis. We evaluate DNA concentration, purity, and stability, and microbial community composition using 16S rRNA gene sequencing and for selected samples using shotgun metagenomic sequencing. Our results suggest that inferred community composition was dependent on inherent specimen properties as well as DNA extraction method. We further show that bead-beating or enzymatic treatment can increase the extraction of DNA from Gram-positive bacteria. Final DNA quantities could be increased by isolating DNA from a larger volume of cell lysate compared to standard protocols. Based on this insight, we have designed an improved DNA isolation procedure optimized for microbiome genomics that can be used for the three examined specimen types and potentially also for other biological specimens.

IMPORTANCE

Sequencing-based analyses of microbiomes may lead to a breakthrough in our understanding of the microbial world associate with humans, animals, and the environment. Such insight could further the development of innovative ecosystem management approaches for the protection of our natural resources, and the design of more effective and sustainable solutions to prevent and control infectious diseases. Genome sequence information is an organism- (pathogen-) independent language that can be used across sectors, space, and time. Harmonized standards, protocols, and workflows for sample processing and analysis can facilitate the generation of such actionable information. In this study, we assessed several procedures for the isolation of DNA for next-generation sequencing. Our study highlights several important aspects to consider in the design and conduction of sequence-based analysis of microbiomes. We provide a standard operating procedure for the isolation of DNA from a range of biological specimens particularly relevant in clinical diagnostics and epidemiology.

INTRODUCTION

Microbial communities fulfill central roles in biological systems, such as in human, animal, and environmental ecosystems. Genomics-based interrogations of these communities can provide unprecedented insight into their composition and function, and reveal general principles and rules about their ecology and evolution (1-4). Genomics-based microbiome analyses can also have important practical implications, such as for the diagnosis and management of infectious diseases. Together with relevant metadata and appropriate bioinformatics and statistical approaches, genomic sequencing data could enable the global surveillance of emerging and re-emerging infectious diseases, and teach us about the reservoirs and transmission pathways of pathogens (5-7). Ultimately, genomics-based information about infectious disease epidemiology may help us to predict, prevent, and control infectious diseases faster, more precisely, and more sustainably.

In order to facilitate large-scale microbiome analyses, harmonized standards for sample handling and data analysis need to be ensured. To be able to establish pathogen reservoirs and transmission pathways, specimens from different sources, such as from humans, animals, and the environment, will need to be examined. For genomics analysis, the DNA needs to be isolated from the specimens for DNA sequencing. However, DNA isolation methods are often only evaluated and established in the context of specimens from an individual source (e.g. human fecal specimens), and seldom across a variety of specimen types (8-12), which is addressed in the present study. Current sequencing technologies, such as Illumina MiSeq and Hiseq, PacBio, IonTorrent, and Nanopore systems, still require large initial DNA template quantities,

particularly from the perspective of PCR-free metagenomics-based analysis. In contrast,

16S rRNA gene profiling can reveal a bacterial and archaeal composition for samples

with low initial DNA template quantities. In metagenomics, low quantities of input DNA

can result in low sequencing data output, and impact the inferred microbial community

composition (13). Hence, modified DNA isolation protocols for increasing DNA quantities

obtained from different types of specimens are desirable.

Here, we examine three specimen types (human feces, animal feces, and sewage), a

total of eight commercially available DNA isolation kits, and a number of protocol

modifications in regard to output DNA (quantity, purity, stability) and microbiome

composition (16S rRNA gene profiling, metagenomics). Our results suggest that both,

the specimen itself as well as the DNA isolation procedure, can affect DNA quantity and

quality, and inferred microbiome composition. Based on the insight gained, we

developed an improved laboratory protocol that can be used for DNA isolations from a

variety of biological specimens.

RESULTS

DNA concentration, purity, and stability depend on the type of specimen and DNA isolation method. We extracted DNA from human feces, pig feces, and hospital sewage, using seven commonly used DNA isolation kits and determined DNA concentration, purity, and stability of the isolated DNA (Fig. 1A and Table 1). The DNA concentrations varied greatly (Fig. 1B; see also Table S1A in the supplemental material). For human feces, the highest DNA concentrations were obtained using the EasyDNA, MagNAPure, and QIAStool procedure, for pig feces using the EasyDNA, QIAStool, and QIAStool+BB procedures, and for sewage using the MagNAPure and EasyDNA procedure, while for three methods the DNA concentration from sewage was below the detection limit. On average across the three types of specimen, the highest DNA concentrations were obtained using EasyDNA (44.96 ng/µl +/- 20.99 SEM) and QIAStool 27.88 ng/µl +/- 2.55 SEM), and the lowest using the PowerSoil.HMP (1.55 ng/µl +/-0.31 SEM) and InnuPREP (7.77 ng/µl +/- 5.54 SEM) methods.

With regard to DNA purity, the best results for human and pig feces were obtained using the EasyDNA, QIAStool, and QIAStool+BB procedure (see Table S1A in the supplemental material). The DNA was generally stable for at least 7 days when stored at room temperature (22°C) with some exceptions (see Table S1A in the supplemental material). A decrease in DNA concentration over time was observed for example for the human feces when extracted with EasyDNA (57% decease in DNA concentration) or MagNAPure (21% decrease in DNA concentration), suggesting the presence of DNases in these extracts. In some cases, an increase in DNA concentration over time was observed, such as for the pig feces when extracted with EasyDNA (32% increase in

DNA concentration). An increase in DNA concentration over time at room temperature

was previously shown to be related to the hyperchromicity of DNA, and dependent on

the DNA concentration and ionic strength of the solution (14).

Microbial richness and diversity are influenced by DNA isolation procedure. For

the human fecal specimen, the highest bacterial Operational Taxonomic Unit (OTU)

richness and diversity were detected using the QIAStool+BB and FastDNA methods,

followed by InnuPURE and PowerSoil as assessed by 16S rRNA gene profiling (Fig. 1C;

see also Table S1B in the supplemental material). In comparison, the determined

richness and diversity for the EasyDNA method was low, and the relative abundance of

Ruminococcaceae and Bifidobacteriaceae dominated the composition compared to the

extracts from the other methods (Fig. 1C; see also Fig. S1A in the supplemental

material). Thirty-nine samples (human feces, pig feces, and sewage) with high DNA

concentration were selected and examined using metagenomic sequencing. In this

assessment, the species richness and diversity for human feces was highest for the

EasyDNA procedure, and a high relative abundance of Ruminococcaceae and

Bifidobacteriaceae was apparent in this analysis as well (see Fig. S1A in the

supplemental material).

For the pig fecal specimen, the highest bacterial richness and diversity were detected

using the PowerSoil.HMP and MagNAPure methods, followed by QIAStool+BB (Fig. 1C;

see also Table S1B in the supplemental material). Similarly, richness and diversity were

highest using the MagNAPure and EasyDNA methods when assessed using

metagenomics. Based on 16S rRNA gene profiling, the richness and diversity for the

FastDNA method were lower compared to all other methods, and the relative

abundance of Clostridiaceae and Turicibacteraceae was higher and the abundance of

Prevotellaceae and Ruminococcaceae lower using this method, compared to the other

methods (Fig. 1C; see also Fig. S1A in the supplemental material).

For the sewage specimen, the highest bacterial richness and diversity was detected

using the InnuPURE method, followed by PowerSoil.HMP and QIAStool+BB, and similar

levels were achieved using the other methods (Fig. 1C; see also Table S1B in the

supplemental material). The relative abundance of Clostridiaceae was highest in the

samples extracted using EasyDNA, and the abundance of Enterobacteriales highest in

the samples extracted using PowerSoil.HMP.

Overall, the relative abundance of predicted Gram-positive bacteria was highest in the

human and sewage specimens when extracted with the EasyDNA method, and highest

in the pig specimen when extracted using the FastDNA method (see Fig. S2 in the

supplemental material). The abundance of predicted Gram-positive bacteria was lowest

using MagNAPure and QIAStool, the two methods that did neither include a bead-

beating step nor specific enzymatic cell-wall digestion.

Microbial community composition depends on the choice of DNA isolation

procedure. The microbial communities from the three types of specimen clustered

separately according to specimen type when examined in PCoA Bray-Curtis ordination,

and not according to DNA isolation procedure (see Fig. S3 in the supplemental

material), indicating that the largest differences between these samples are driven by

the inherent microbiota composition. Bray-Curtis dissimilarity distance analysis carried

out separately for each of the three specimens revealed that the samples largely

clustered according to DNA isolation procedure (Fig. 2A-C). For the human fecal

specimen, the bacterial community composition derived from the EasyDNA isolation

differed from the communities obtained using all other methods (Fig. 2A), which is in

agreement with the observations on microbial richness (above). The Bray-Curtis

distances between the samples from InnuPURE, MagNAPure, FastDNA,

PowerSoil.HMP, QIAStool, and QIAStool+BB DNA isolations were on average 0.337 +/-

0.012 SEM, whereas the distances between these and the ones derived from the

EasyDNA procedure were on average 0.825 +/- 0.014 SEM.

For the pig fecal specimen, the bacterial communities derived from the FastDNA

isolation differed from all other communities (Fig. 2B). The average Bray-Curtis distance

between the samples originating from all but the FastDNA procedure was on average

0.473 +/- 0.008 SEM, whereas the distance between these and the ones derived from

the FastDNA procedure was on average 0.877 +/- 0.007 SEM.

For the hospital sewage specimen, the bacterial communities originating from the

EasyDNA method differed from all others (average Bray-Curtis distance 0.600 +/- 0.006

SEM) (Fig. 2C), similar to the human fecal matrix (Fig. 2A). In addition, the communities

originating from the QIAStool DNA isolation differed from all others (average Bray-Curtis

distance 0.514 +/- 0.009 SEM), whereas the average Bray-Curtis distance between all

but the QIAStool and EasyDNA samples was 0.460 +/-0.11 SEM on average.

Distinct taxa account for the differences observed between DNA isolation

methods. To quantify the effect of DNA isolation method on microbial community

composition we tested for differential abundance of taxa between the communities

derived from the different DNA isolation methods using DESeq2 analyses. In pairwise

comparisons, significant differences between the DNA isolation methods were observed

(Fig. 2D-F; see also Table S2 in the supplemental material).

The most abundant family on average in the human fecal specimen was Prevotellaceae

(Bacteroidetes), and its abundance was significantly lower in the samples extracted with

EasyDNA as compared to all other methods (e.g. 18.3-fold lower in EasyDNA vs.

QIAStool, adjusted p-value 1.91^{-6}) (Fig. 2D; see also Table S2 in the supplemental

material). Similarly, the abundance of Bacteroidaceae (Bacteroidetes),

Porphyromonadaceae (Bacteroidetes), Alcaligenaceae (β-Proteobacteria), and

Pasteurellaceae (γ-Proteobacteria) was lower in the samples from the EasyDNA

isolation compared to the other methods. In contrast, the abundance of

Bifidobacteriaceae (Actinobacteria) was higher in the samples originating from the

EasyDNA procedure compared to all other methods (e.g. 770-fold higher in EasyDNA

vs. QIAStool, adjusted p-value 7.49^{-57}). The abundance of Verrucomicrobiaceae

(Verrucomicrobia) was significantly lower in the samples from the QIAStool+BB and

PowerSoil.HMP DNA isolations (e.g. 4.15-fold lower in QIAStool+BB vs. QIAStool,

adjusted p-value 0.001).

The most abundant family on average in the pig fecal specimen was Prevotellaceae

(Bacteroidetes), and its abundance differed significantly between the DNA isolation

procedures (e.g. 2.3-fold lower in EasyDNA vs. PowerSoil.HMP, adjusted p-value 1.28^{-5})

(Fig. 2E; see also Table S2 in the supplemental material). The abundance of

Clostridiaceae (Clostridia), the on average fourth most abundant family in the pig feces,

was significantly higher in the samples extracted by the FastDNA method (e.g. 166-fold

higher in FastDNA vs. EasyDNA, adjusted p-value 7.35^{-110}).

Moraxellaceae (γ-Proteobacteria) was the most abundant family on average in the

hospital sewage, and its abundance was significantly higher in the samples from the

EasyDNA isolation compared to other DNA isolation methods (e.g. 2.6-fold higher in

EasyDNA vs. PowerSoil.HMP, adjusted p-value 3.82^{-5}) (Fig. 2F; see also Table S2 in

the supplemental material). Ruminococcaceae (Clostridia), the on average third most

abundant family in sewage, were also significantly more abundant in the samples from

the EasyDNA isolation compared to other DNA isolation procedures (e.g. 7.3-fold higher

in EasyDNA vs. FastDNA, adjusted p-value 4.28^{-17}).

DNA isolation procedure affects the abundance of taxa differently across

specimens. Given that differential taxa abundances were observed for the different

DNA isolation procedures for the three specimen types, we investigated whether the

abundance differed in the same way between DNA isolation procedures across

specimens. For example, we were asking: If taxon A is observed at a higher abundance

upon DNA isolation with method X compared to method Y in specimen type 1, is this

taxon also observed at a higher abundance upon DNA isolation with method X

compared to method Y in specimen type 2? We examined taxa that were detected in all

three specimen types, and selected representative families from different phyla (Fig. 3).

Similar patterns of differential abundance were observed for certain taxa across

specimen types, with exceptions, including two families from the Bacteroidetes phylum.

The abundance of Prevotellaceae and Bacteroidaceae was significantly lower when

human fecal specimen were extracted with EasyDNA compared to other methods. In

contrast, these two families were observed at a significantly higher abundance when

sewage was extracted with EasyDNA compared to other methods (Fig. 3).

Likewise, Ruminococcaceae of the phylum Clostridia were observed at a significantly

higher abundance in human fecal and hospital sewage samples but not in pig fecal

samples when extracted with the EasyDNA method compared to other methods. The

same pattern was however not observed for all families of the phylum Clostridia.

Clostridiaceae abundance appeared higher in human and pig feces when extracted with

FastDNA compared to other methods, and Clostridiaceae abundance appeared higher

in sewage when extracted using the EasyDNA method compared to other methods (Fig.

3).

Thus, we found significant differences in the abundance of certain families according to

specimen type, which sometimes depend on the DNA isolation procedure. Some of the

differential abundance patterns were similar across the three types of specimens, while

others differed.

Detection of spiked bacteria is dependent on DNA isolation procedure and

specimen type. In order to quantify DNA isolation efficiency, we spiked the three

specimen with known numbers of two bacterial representatives, namely *Salmonella*

enterica serotype Typhimurium DT104 (Gram-negative) and *Staphylococcus aureus*

ST398 (Gram-positive) in a CFU ratio of 1.02. Both, *S. enterica* and *S. aureus* were

present in negligible numbers in the three specimens before spiking. DNA was isolated

from these samples using the seven different DNA isolation methods, and the

abundance of the two strains determined using 16S rRNA gene profiling, and for some

samples also using metagenomics. Based on 16S rRNA gene profiling, the spiked

organisms accounted for an average abundance of 1.0% (+/-0.29 SEM)

Enterobacteriaceae, and 0.29% (+/-0.11 SEM) Staphylococcaceae across the three

types of specimen.

Using QIAStool, a DNA isolation method that does not involve a bead-beating step, the

abundance of Enterobacteriaceae was higher in the spiked human fecal specimen than

expected, with an Enterobacteriaceae/Staphylococcaceae (E/S) ratio of 13.9 (Fig. 4A).

This ratio was lower in the spiked human fecal specimen using InnuPURE, FastDNA,

PowerSoil.HMP, and QIAStool+BB, which are all methods that involve a bead-beating

step (E/S ratio range 0.3-2.3). The EasyDNA method involves an additional enzymatic

lysis step, and using this method the determined E/S ratio was 3.7. Using the

MagNAPure method no or lower read numbers assigned to Staphylococcaceae were

detected in the spiked samples compared to not spiked samples in the human fecal

specimen, and hence the ratio resulted in negative values (Fig. 4A). A similar result was

obtained when the samples were examined using metagenomics (see Fig. S4 in the

supplemental material).

Overall, most DNA isolation methods exhibited a similar tendency across the three types

of specimen. For example, for all three specimen types, the E/S ratio was higher using

the QIAStool method, compared to the other methods (except MagNAPure for sewage).

However, when the strain mix, composed of *S. enterica* and *S. aureus* only, was

extracted using the seven DNA isolation procedures, their determined E/S ratio was in

almost all cases similar to the expected ratio of 1.02, including the QIAStool method.

This suggests that quantification of distinct organisms from complex specimens is more

challenging when the organisms are present at lower abundance levels. Inherent

specimen properties may influence the DNA isolation efficiency leading to a biased

pattern of microbial community composition.

Protocol modifications for increasing DNA concentration. One goal in genomics is

to obtain a predicted pattern of microbial community composition that closely resembles

the actual composition of microorganisms in a particular environment. Another challenge

is to obtain sufficient DNA for metagenome sequencing. To address this aspect, we

examined the effect of modifications to standard protocols on output DNA concentration

(modifications are described in detail in the Supplemental Materials & Methods section).

We chose the QIAStool method as a starting point, as we obtained DNA extracts using

this method that were of high purity and stability (see Table S1A in the supplemental

material). Another concern is processing time and costs for DNA isolation procedures,

particularly for large-scale microbiome projects. The protocol of the QIAamp Fast DNA

Stool Mini kit (QIAFast), a kit that became available at the time the present study was

carried out, suggested reduced processing time compared to the QIAStool method.

When we compared the QIAStool and QIAFast methods using metagenomic

sequencing, we obtained a similar richness, diversity, and microbial community

composition with these two methods (see Fig. S5 in the supplemental material).

Furthermore, given that our previous results suggested that including a bead-beading

step might result in a predicted community composition that was more similar to the

community of known composition than without this step (Fig. 4), we included a bead-

beating step and examined the effect of beads of differing types and cost (Table 1). We

obtained a higher DNA concentration using pig feces and the QIAStool kit, when bead

beating was applied and the double amount of volume after cell lysis was transferred

(Fig. 5A). Similarly, for the QIAFast method, we obtained an on average 2.6-fold higher

DNA concentration by including a bead beading step and transferring the double amount

of volume after cell lysis, compared to DNA isolations without these modifications (Fig.

5A). Both, DNA purity and stability were in the expected range (see Table S3 in the

supplemental material). Even though the DNA concentration was higher with these

protocol modifications, the richness, diversity and community composition did not

significantly differ when assessed by 16S rRNA gene profiling (Fig. 5A).

A particular DNA isolation method did not however lead to the highest DNA

concentrations for each of the three types of specimen. Whereas the highest DNA

concentration for sewage was achieved using the QIAFast+BB.GBT+2Trans method

(27.30 ng/ul +/- 4.5 SEM), the highest DNA concentration for human feces was obtained

using the QIAStool+BB.LMA method (22.50 ng/ul +/- 4.77 SEM) (Fig. 5B). For pig feces,

the highest DNA concentrations were obtained using the QIAStool+BB.LMA (15.43 ng/ul

+/-3.48 SEM) and QIAStool (14.57 ng/ul +/-3.62 SEM) methods. On average across the

three types of specimen, the highest DNA concentrations were obtained using the

QIAFast+BB.GBT+2Trans (17.66 ng/ul +/- 4.82 SEM) and QIAStool+BB.LMA (17.46

ng/ul +/- 2.54 SEM) methods.

DISCUSSION

Genomics-based investigations of complex microbiomes greatly enhance our

understanding about microbial community composition and function relevant to human,

animal, and plant health, infectious diseases, environmental pollution, agriculture, and

food safety. One current ambitious goal is to establish a global surveillance system for

infectious agents and antimicrobial resistance based on next-generation DNA

sequencing approaches (15). Given that infectious agents occupy various ecological

habitats, DNA needs to be extracted from various types of specimen using standardized

approaches in a time- and cost-efficient manner. It is advantageous, if a range of

different specimens can be processed using the same standard operating procedure. In

light of these considerations, we compared a total of eight DNA isolation procedures,

and based on the findings developed an improved protocol using the QIAamp® Fast

DNA Stool Mini kit.

Overall, the amounts of DNA obtained from each DNA isolation method differed greatly,

and there was no significant correlation between increasing DNA amount and increase

in community diversity or richness. The taxonomic microbiome composition appeared to

be dependent on both, the specimen and DNA isolation method. For example, the

EasyDNA procedure preferentially extracted DNA from Gram-positive bacteria from the

human feces and hospital sewage, while the FastDNA procedure preferentially extracted

DNA from Gram-positive bacteria from pig feces. Methods that did not include a bead-

beating or enzymatic treatment step generally extracted less DNA from Gram-positive

bacteria.

When using a particular procedure we found some similar abundance patterns of

specific bacterial families between the three specimen types. However, we also

observed several differences. Hence, one cannot conclude that the DNA from a

particular bacterial family will be extracted preferentially using one specific DNA isolation

method across different types of specimens. This could be due to different inherent

cellular properties of the taxa belonging to a specific family, affecting mechanical and

enzymatic cell lysis. Moreover, the chemical and physical composition of the specimen

could influence DNA isolation and downstream procedures. For example, it is well

known that certain compounds, such as humic acid, polysaccharides, and bilirubin can

affect PCR (16). Furthermore, fecal sample consistency, reflecting differences in water

content and activity, can impact on microbial community composition (17).

Our observations from 16S rRNA gene profiling and metagenomics generally agreed,

but the taxonomic patterns also exhibited some differences. One reason could be the

known primer biases towards certain taxa in 16S rRNA gene based analysis (18). An

additional reason could be differences in the composition of the reference databases

used for the two sequence-based strategies. While 16S rRNA gene databases are

composed of 16S rRNA gene sequences from a large diversity of taxa, the

metagenomic sequence databases are based on whole and draft genome sequences

from fewer and less diverse taxa. Both strategies complement each other, and efforts

are ongoing in developing harmonized analytical workflows for sequence-based

microbial community analysis.

Based on the insight gained in this study, we have developed an improved DNA

isolation method based on the QIAamp® Fast DNA Stool Mini kit. This procedure

includes a bead beading step to obtain DNA from both, Gram-positive and Gram-

negative taxa, and a step in which the double amount of cell lysate is transferred to the

column to increase the DNA quantity. A standard operating procedure for this DNA

isolation method is available from https://dx.doi.org/10.6084/m9.figshare.3475406.v1,

and which can be used for different specimen types relevant to projects like EFFORT-

against-AMR, COMPARE-Europe, the International Microbiome Initiative, and

International Human Microbiome Standards.

In summary, our findings provide new insight into the effect of different specimen types

and DNA isolation methods on DNA quantities and genomic-based inference of

microbiome composition. We offer an optimized strategy for the DNA isolation for

different sample types providing a representative insight into community composition,

and which can be conducted in a time- and cost-efficient manner.

MATERIAL AND METHODS

Specimen Collection and Handling

Human fecal specimens were collected from a healthy individual. Pig fecal specimens

were collected from animals at a conventional pig production farm in Denmark.

Untreated sewage was collected from the sewage inlet of the Herlev hospital waste

water treatment plant, Denmark. For details regarding sample handling and processing,

see Supplemental Materials and Methods.

Spiking with strain mix

Subsequent to specimen collection, about half of the aliquots from the human, pig, and

sewage were spiked with a representative of Gram-positive and Gram-negative bacteria,

namely *Staphylococcus aureus* ST398 (strain S0385) and *Salmonella enterica* serotype

Typhimurium DT104. For details regarding the preparation of the strain mix, see

Supplemental Materials and Methods.

DNA isolation

In a first step, seven DNA isolation procedures were examined, namely: InnuPure® C16,

Analytic Jena AG (InnuPURE); MagNA Pure LC DNA isolation Kit III, Roche

(MagNAPure); Easy-DNA™ gDNA Purification Kit, Invitrogen (EasyDNA); MP

FastDNA™ Spin Kit, MP Biomedicals (FastDNA); PowerSoil® DNA Isolation kit, MoBio

(PowerSoil.HMP); QIAamp® DNA Stool Mini Kit, Qiagen (QIAStool); QIAamp® DNA

Stool Mini Kit +Bead Beating, Qiagen (QIAStool+BB) (see Table 1, and details below).

In a second step, a variety of modifications to two Qiagen kits were examined, namely

the QIAamp® DNA Stool Mini Kit (QIAStool), and QIAamp® Fast DNA Stool Mini Kit

(QIAFast). The standard operating procedure for an improved DNA isolation method (i.e.

QIAamp Fast DNA Stool Modified, corresponding to QIAFast+BB.GBT+2Trans

described here) can be found at https://dx.doi.org/10.6084/m9.figshare.3475406.v1. For

details regarding the individual DNA isolation procedures, see Supplemental Materials

and Methods.

DNA quantitation and quality assessment

Subsequent to DNA isolation, the DNA was portioned into 10-µl aliquots to prevent

repeated freeze-thawing cycles, and stored at -20°C. DNA concentrations were

measured using Qubit® dsDNA BR Assay Kit on a Qubit® 2.0 Fluorometer (Invitrogen,

Carlsbad, CA). As DNA extracts can contain contaminants, such as proteins and other

organic molecules that can affect downstream procedures such as DNA amplifications in

PCR, we determined the DNA purity by measuring the ratios of absorbance at 260/280

and 260/230, respectively, using a NanoDrop 1000 Spectrophotometer (Thermo

Scientific, Pittsburgh, USA). DNA extracts with a 260/280 ratio between ~1.7 to ~ 2.0,

and 260/230 ration between ~2.0 to ~2.2 are regarded as "pure". The stability of the

DNA in the extracts was determined by measuring the DNA concentration after 2 and 7

days incubation at 22°C. A decrease in DNA concentration over time can indicate the

presence of DNases in the extract.

16S rRNA gene profiling

16S rRNA gene amplicon libraries were generated using a two-step protocol similar as

described in Part # 15044223 Rev. B by Illumina. In a first PCR, the V4 region of the

16S rRNA genes were amplified using the universal primers (515f 5'-

TGCCAGCAGCCGCGGTAATAC (19) and 806r 5'-GGACTACNNGGGTATCTAAT (20).

The samples were pooled in equal concentrations, and concentrated using 'DNA clean

and concentrator-5 kit' (Zymo Research, Orange, CA). Paired-end 2 × 250 bp

sequencing of barcoded amplicons was performed on a MiSeq machine running v2

chemistry (Illumina Inc., San Diego, CA, USA). The sequences were processed using

the UPARSE pipeline (21) and a OTU x sample contingency table was created. Using

QIIME1.8.0 (22), taxonomy was assigned with uclust using assign_taxonomy.py based

on the Greengenes 13.8 reference database. Ecological diversity estimates and

microbial community comparisons were performed using the relevant scripts provided by

QIIME, phyloseq, and R (22-24). For details regarding the 16S rRNA gene-based

microbial community analysis, see Supplemental Materials and Methods.

Metagenomics

A subset of the DNA extracts was subjected to metagenomic sequencing. The samples

were prepared and sequenced following the Nextera XT DNA Library Preparation Guide

for the MiSeq system Part # 15031942 Rev. D, using paired-end v2 2×250bp

sequencing. The taxonomic microbiome compositions were determined through the use

of the MGmapper pipeline (25). The MGmapper package is available for download at

www.cbs.dtu.dk/public/MGmapper/. For details regarding the metagenomics-based

microbial community analysis, see Supplemental Materials and Methods.

Differential abundance analysis

In order to test for the differential abundance of taxa that may drive the differences

observed between the communities derived from the different DNA isolation procedures,

we performed DESeq2 analyses. The read count tables from the 16S rRNA gene

profiling and metagenomics sequence analysis, respectively, were aggregated to the

family level in R (v. 3.2.3, 64bit) (24) We performed an analysis that allows for varied

sequencing depth, similar as suggested previously (26), and carried out two-sided Wald

tests as implemented in the DESeq2 (v. 1.10.1) package (27). The size factors were

determined by DESeq2 from the read count tables. For details regarding the differential

abundance analysis, see Supplemental Materials and Methods.

Quantification of strain mix

The samples that were spiked with the strain mix composed of *S. enterica* Typhimurium

DT104 and *S. aureus* ST398 were extracted, sequenced, and analyzed together with

the non-spiked samples. For each type of specimen and isolation method, the

abundance of Enterobacteriaceae and Staphylococcaceae for 16S rRNA gene profiling

and metagenomics, respectively, were determined. The ratios between

Enterobacteriaceae and Staphylococcaceae was determined for each sample matrix

and isolation method, and compared to the *S. enterica* Typhimurium DT104 / *S. aureus*

ST398 ratio of CFU that were added to the original samples. For details regarding the

quantification of the strain mix, see Supplemental Materials and Methods.

Ethics

The collection of human and pig fecal specimens as well as sewage was non-invasive,

and were performed in accordance with the Declaration of Helsinki, and complied with

Danish and European directives (86/609/EEC). The collection of specimens was

conducted in accordance with the act on research ethics of heath research projects as

administrated and confirmed by the National Committee on Health Research Ethics of

Denmark (Region Hovedstaden), Journal nr. H-14013582.

Nucleotide sequences

The 16S rRNA gene sequences are available from the European Nucleotide Archive

(ENA) at the European Bioinformatics Institute (EBI) under accession number

PRJEB12431, and the metagenomic sequences from ENA at EBI under accession

number PRJEB14814.

ACKNOWLEDGEMENTS

We thank Karin Vestberg (University of Copenhagen), Christina A. Svendsen (Technical

University of Denmark) and Jacob D. Jensen (Technical University of Denmark) for

technical assistance related to DNA sequencing. Marie S. Jensen (Technical University

of Denmark) is acknowledged for the collection of pig fecal samples.

FUNDING INFORMATION

This work was supported by the European Unions's Seventh Framework Programme,

FP7 (613754), the Framework Programme for Research and Innovation, Horizon2020

(643476), and The Villum Foundation (VWR023052). Sünje J. Pamp was supported by a

grant from Carlsbergfondet (2013_01_0377). The funders had no role in study design,

data collection and interpretation, or the decision to submit the work for publication.

AUTHOR CONTRIBUTIONS

B.E.K., L.B., F.M.A., and S.J.P. designed the research; B.E.K., L.B., O.L., and P.M.

performed the research; B.E.K., L.B., O.L., P.M., A.P., F.M.A., and S.J.P. contributed

510   analytic tools; B.E.K., L.B., O.L., P.M., and S.J.P. analyzed the data; B.E.K., L.B., and

511   S.J.P. wrote the manuscript; and O.L., P.M., and A.P. edited the manuscript.

REFERENCES

1. **Ding T**, **Schloss PD**. 2015. Dynamics and associations of microbial community types across the human body. Nature **509**:357–360.

2. **Schloissnig S**, **Arumugam M**, **Sunagawa S**, **Mitreva M**, **Tap J**, **Zhu A**, **Waller A**, **Mende DR**, **Kultima JR**, **Martin J**, **Kota K**, **Sunyaev SR**, **Weinstock GM**, **Bork P**. 2013. Genomic variation landscape of the human gut microbiome. Nature **493**:45–50.

3. **Guidi L**, **Chaffron S**, **Bittner L**, **Eveillard D**, **Larhlimi A**, **Roux S**, **Darzi Y**, **Audic S**, **Berline L**, **Brum JR**, **Coelho LP**, **Espinoza JCI**, **Malviya S**, **Sunagawa S**, **Dimier C**, **Kandels-Lewis S**, **Picheral M**, **Poulain J**, **Searson S**, **Stemmann L**, **Not F**, **Hingamp P**, **Speich S**, **Follows M**, **Karp-Boss L**, **Boss E**, **Ogata H**, **Pesant S**, **Weissenbach J**, **Wincker P**, **Acinas SG**, **Bork P**, **de Vargas C**, **Iudicone D**, **Sullivan MB**, **Raes J**, **Karsenti E**, **Bowler C**, **Gorsky G**. 2016. Plankton networks driving carbon export in the oligotrophic ocean. Nature **532**:465–470.

4. **Costello EK**, **Stagaman K**, **Dethlefsen L**, **Bohannan BJM**, **Relman DA**. 2012. The application of ecological theory toward an understanding of the human microbiome. Science **336**:1255–1262.

5. **Relman DA**. 2014. Actionable Sequence Data on Infectious Diseases in the Clinical Workplace. Clinical Chemistry **61**:38–40.

6. **Petersen TN**, **Rasmussen S**, **Hasman H**, **Carøe C**, **Bælum J**, **Schultz AC**, **Bergmark L**, **Svendsen CA**, **Lund O**, **Sicheritz-Ponten T**, **Aarestrup FM**. 2015. Meta-genomic analysis of toilet waste from long distance flights; a step towards global surveillance of infectious diseases and antimicrobial resistance. Sci Rep 1–9.

7. **Lipkin WI**. 2013. The changing face of pathogen discovery and surveillance. Nature Reviews Microbiology **11**:133–141.

8.   **Wesolowska-Andersen A**, **Bahl MI**, **Carvalho V**, **Kristiansen K**, **n TS-P**, **Gupta R**,

Licht TR. 2014. Choice of bacterial DNA extraction method fromfecal material influences

community structure asevaluated by metagenomic analysis. Microbiome **2**:1–11.

9.   **Mackenzie BW**, **Taylor MW**, **Waite DW**. 2015. Evaluating variation in human gut

microbiota profiles dueto DNA extraction method and inter-subject differences 1–11.

10.   **Henderson G**, **Cox F**, **Kittelmann S**, **Miri VH**, **Zethof M**, **Noel SJ**, **Waghorn GC**,

Janssen PH. 2013. Effect of DNA Extraction Methods and Sampling Techniques on the

Apparent Structure of Cow and Sheep Rumen Microbial Communities. PLoS ONE

8:e74787.

11.   **Yuan S**, **Cohen DB**, **Ravel J**, **Abdo Z**, **Forney LJ**. 2012. Evaluation of Methods for the

Extraction and Purification of DNA from the Human Microbiome. PLoS ONE **7**:e33865.

12.   **Albertsen M**, **Karst SM**, **Ziegler AS**, **Kirkegaard RH**, **Nielsen PH**. 2015. Back to Basics

– The Influence of DNA Extraction and Primer Choice on Phylogenetic Analysis of

Activated Sludge Communities. PLoS ONE **10**:e0132783.

13.   **Bowers RM**, **Clum A**, **Tice H**, **Lim J**, **Singh K**, **Ciobanu D**, **Ngan CY**, **Cheng J-F**,

Tringe SG, **Woyke T**. 2015. Impact of library preparation protocols and template quantity

on the metagenomic reconstruction of a mock microbial community. BMC Genomics

16:1–12.

14.   **Gupta Das NN**, **Basu S**, **Bagchi B**. 1967. Conformational Changes in Denaturated DNA,

pp. 663–688. *In* Ramachandran, GN (ed.), Conformation of Biopolymers. Academic

Press, London and New York.

15.   **Aarestrup FM**, **Koopmans MG**. 2016. Sharing Data for Global Infectious Disease

Surveillance and Outbreak Detection. Trends in Microbiology **24**:241–245.

16. **Radstrom P**, **Knutsson R**, **Wolffs P**, **Lovenklev M**, **Lofstrom C**. 2004. Pre-PCR

processing: strategies to generate PCR-compatible samples. Mol Biotechnol **26**:133–146.

17. **Vandeputte D**, **Falony G**, **Vieira-Silva S**, **Tito RY**, **Joossens M**, **Raes J**. 2016. Stool

consistency is strongly associated with gut microbiota richness and composition,

enterotypes and bacterial growth rates. Gut **65**:57–62.

18. **Klindworth A**, **Pruesse E**, **Schweer T**, **Peplies J**, **Quast C**, **Horn M**, **Glockner FO**.

2012. Evaluation of general 16S ribosomal RNA gene PCR primers for classical and next-

generation sequencing-based diversity studies. Nucleic Acids Research **41**:e1–e1.

19. **Yu Y**, **Lee C**, **Kim J**, **Hwang S**. 2005. Group-specific primer and probe sets to detect

methanogenic communities using quantitative real-time polymerase chain reaction.

Biotechnol Bioeng **89**:670–679.

20. **Sundberg C**, **Al-Soud WA**, **Larsson M**, **Alm E**, **Yekta SS**, **Svensson BH**, **Sørensen SJ**,

Karlsson A. 2013. 454 pyrosequencing analyses of bacterial and archaeal richness in 21

full-scale biogas digesters. FEMS Microbiology Ecology **85**:612–626.

21. **Edgar RC**. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon

reads. Nature Methods **10**:996–998.

22. **Caporaso JG**, **Kuczynski J**, **Stombaugh J**, **Bittinger K**, **Bushman FD**, **Costello EK**,

Fierer N, **Peña AG**, **Goodrich JK**, **Gordon JI**, **Huttley GA**, **Kelley ST**, **Knights D**,

Koenig JE, **Ley RE**, **Lozupone CA**, **McDonald D**, **Muegge BD**, **Pirrung M**, **Reeder J**,

Sevinsky JR, **Turnbaugh PJ**, **Walters WA**, **Widmann J**, **Yatsunenko T**, **Zaneveld J**,

Knight R. 2010. QIIME allows analysis of high-throughput community sequencing data.

Nature Methods **7**:335–336.

23. **McMurdie PJ**, **Holmes S**. 2013. phyloseq: An R Package for Reproducible Interactive

Analysis and Graphics of Microbiome Census Data. PLoS ONE **8**:e61217.

24.   **R Development Core Team**. 2014. R: A Language and Environment for Statistical

Computing. R Foundation for Statistical Computing, Vienna, Austria.

25.   **Petersen TN**, **Bælum J**, **Lukjancenko O**, **Geertz-Hansen HM**, **Thomsen MCF**,

Sperotto MM, **Lund O**, **Aarestrup FM**, **Sicheritz-Ponten T**. 2015. MGmapper: an

automated pipeline for mapping and stratification of metagenomics sequence data.

Submitted.

26.   **McMurdie PJ**, **Holmes S**. 2014. Waste Not, Want Not: Why Rarefying Microbiome Data

Is Inadmissible. PLOS Computational Biology **10**:e1003531.

27.   **Love MI**, **Huber W**, **Anders S**. 2014. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biology **15**:31.

FIGURES

FIG 1. Comparison of DNA extraction methods. (A) Experimental design. Human feces, pig feces, and hospital sewage were extracted using seven different DNA extraction methods (see also Table 1): InnuPure® C16, MagNA Pure LC DNA isolation Kit III, Easy-DNA™ gDNA Purification Kit, MP FastDNA™ Spin Kit, PowerSoil® DNA

Isolation kit, QIAamp® DNA Stool Mini Kit, QIAamp® DNA Stool Mini Kit + Bead Beating (For details see Materials and Methods). DNA concentration, purity, and stability were examined, and microbial community composition determined using 16S rRNA gene profiling and metagenomics (selected samples). (B) DNA from each method was dissolved in 100 ul solution and DNA concentrations were determined using Qubit® dsDNA BR Assay Kit measurements. Values represent averages from duplicate or triplicate DNA extractions (See also Supplemental Table S1A). (C) Ecological richness (Chao 1) and diversity (Shannon index) were determined based on contingency tables from 16S rRNA gene profiling and metagenomic sequencing data at OTU and species levels, respectively (See also Supplemental Table S1B).

FIG 2. Microbial community dissimilarity. The dissimilarity between the microbiotas from the human, pig, and sewage samples based on DNA extraction methods was

examined using Principal Coordinates Analysis of Bray-Curtis distances (A-C) and differential abundance analysis using DESeq2 (D-F) from 16S rRNA amplicon data. (A-F) For the PCoA Bray-Curtis ordination analysis only samples with 800 or more reads were included. (D-F) For the differential abundance analysis pairwise testing by DNA extraction method was performed, and bacterial families were considered significantly differentially abundant if their adjusted P-value was <0.1 (see also Table S2 in the supplemental material). Examples for differentially abundant families are shown that are among the top10 most abundant taxa found in the sample, respectively. For each family, the total number of DNA isolation procedures, that exhibit significantly different abundance values compared to a particular DNA isolation procedure, are indicated above the plot, respectively.

FIG 3. Differential abundance of bacterial families. Pairwise testing by DNA extraction method was performed using DESeq2, and the log2-fold difference displayed (column vs. rows) for selected families present in all sample matrices if their adjusted P-value was <0.1 (see also Table S2 in the supplemental material). The rank abundance position for each family per sample matrix type is noted according to their regularized log abundance. The baseMean (bM) indicates the mean of negative-binominal-based normalized read counts.

FIG 4. Detection of spiked bacteria. The human fecal (A), pig fecal (B), and hospital sewage (C) samples were spiked with a strain mix composed of *Salmonella enterica* serotype Typhimurium DT104 and *Staphylococcus aureus* ST398 in a CFU ratio of 1.02. The three sample matrices, as well as aliquots of the strain mix (D) were extracted using seven different DNA extraction methods. The two strains were detected by 16S rRNA gene profiling, and their ratios determined. For details, see Materials and Methods. An asterisk in (D) indicates that the values for the particular DNA extraction of the strain mix are based on single measurements. All other values are based on averages from duplicate or triplicate DNA extractions. The dashed line indicates the ratio of the strain mix based on CFU determinations.

FIG 5. Effect of protocol modifications. A) Pig feces was extracted using standard as well as modified protocols based on the QIAamp® DNA Stool Mini and QIAamp® Fast DNA Stool Mini kits. The modifications included bead beading, pre-treatment of the sample, and transfer of the double amount of volume after cell lysis. In the bead-beating step, different bead types were examined (For details, see Materials and Methods, and Table 1). The alpha diversity (Chao 1 and Shannon index) was determined at OTU-level, and the microbial community composition examined at family-level based on 16S rRNA gene profiling. B) Selected standard and modified DNA extraction protocols were employed to extract DNA from human feces, pig feces, and sewage and their DNA concentration was displayed in a star plot. The values indicate the averages from duplicate extractions.

TABLE 1. Overview of DNA extraction procedures

Extraction Method	Sample amount (g)	Cell lysis method	Bead type	DNA separation	Cost pr. extraction (€)[a]	Processing time for 20 samples (h)
Step 1: Seven commonly used DNA extraction kits						
InnuPure® C16 (Analytic Jena AG) [A]	0.1	Chemical, Mechanical, Heat	Ceramic	Magnetic beads	7.3	4
MagNA Pure LC DNA isolation Kit III (Roche) [A]	0.25	Chemical, Heat	-	Magnetic beads	2.6[b]	2.5
Easy-DNA™ gDNA Purification Kit (Invitrogen)	0.25	Chemical, Enzymatic	None	Phenol:Chloroform, Precipitation	4.5	8.8
MP FastDNA™ Spin Kit (MP Biomedicals)	0.5	Chemical, Mechanical	Ceramic & Garnet	Silica membrane-based columns	14.1[c]	5
PowerSoil® DNA Isolation kit (MoBio)	0.25	Chemical, Mechanical, Heat	Garnet	Silica membrane-based columns	5.3	5.5
QIAamp® DNA Stool Mini Kit (Qiagen)	0.2	Chemical, Heat	-	Silica membrane-based columns	5.3	4
QIAamp® DNA Stool Mini Kit (Qiagen) +BB (Lysing Matrix A, MP Biomedicals)	0.2	Chemical, Mechanical, Heat	Ceramic & Garnet	Silica membrane-based columns	12.7	4
Step 2: New and modified DNA extraction procedures						
QIAamp® DNA Stool Mini Kit (Qiagen) +BB (Garnet Bead Tubes, MoBio)	0.2	Chemical, Mechanical, Heat	Garnet	Silica membrane-based columns	8.5	3
QIAamp Fast DNA Stool Mini	0.2	Chemical, Mechanical, Heat	-	Silica membrane-based columns	6.2	2.6
QIAamp Fast DNA Stool Mini +BB (Lysing Matrix A, MP Biomedicals)	0.2	Chemical, Mechanical, Heat	Ceramic & Garnet	Silica membrane-based columns	13.6	3
QIAamp Fast DNA Stool Mini +BB	0.2	Chemical, Mechanical,	Glass	Silica membrane-based columns	10	3

(Pathogen Lysis Tubes S, Qiagen)		Heat				
QIAamp Fast DNA Stool Mini +BB (Pathogen Lysis Tubes L, Qiagen)	0.2	Chemical, Mechanical, Heat	Glass	Silica membrane-based columns	10	3
QIAamp Fast DNA Stool Mini +BB (Garnet Bead Tubes, MoBio)	0.2	Chemical, Mechanical, Heat	Garnet	Silica membrane-based columns	8.5	3
QIAamp Fast DNA Stool Mini +BB (Bead Beating Tubes, A&A Biotechnology)	0.2	Chemical, Mechanical, Heat	Zirconia / Silica	Silica membrane-based columns	8.2	3

[A] Automated procedure

BB Bead beating

[a]Calculations do not include costs for additional laboratory supply, such as pipette tips and reaction tubes.

[b]Excluding costs for special pipette tips and plastic cartridges required for the robot.

[c]Based on price in the USA, excluding general sales tax that is being added in other countries.

SUPPLEMENTAL MATERIAL

I) Supplemental Materials and Methods

Specimen Collection and Handling
Human fecal specimens were collected from a healthy individual at three time points over a single day. The specimens were kept at 4°C, and transported to the laboratory within 24 hours. Upon arrival, the three samples were pooled and homogenized. For this study, fecal specimens from an infant were chosen, as infant fecal samples often contain a high proportion of Actinobacteria (e.g. Bifidobacteria), from which genomic DNA can be difficult to isolate. Pig fecal specimens were collected from animals at a conventional pig production farm in Denmark. Samples from individual animals were obtained directly after defecation, stored in a cooling box, and transported to the laboratory within four hours. Upon arrival, three random samples were pooled and homogenized. Untreated sewage was collected from the sewage inlet of the Herlev hospital waste water treatment plant, Denmark. Specimens were stored in a cooling box and transported to the laboratory within two hours. Upon arrival 24 x 40 ml sewage samples were sedimented for 10 minutes at 8000xg in an Eppendorf 5810R centrifuge. The sewage pellets were pooled and homogenized. For all three types of specimen (human feces, pig feces, sewage), the homogenized samples were separated into 0.5 g aliquots, respectively. A subset of aliquots for each specimen type was spiked with two bacterial strains (see details below). The individual sample aliquots with and without strain mix were stored at -80°C until further processing.

Spiking with strain mix
Subsequent to specimen collection, about half of the aliquots from the human feces, pig feces, and sewage were spiked with a representative of Gram-positive and Gram-negative bacteria, namely *Staphylococcus aureus* ST398 (strain S0385) and *Salmonella enterica* serotype Typhimurium DT104. The strains were cultivated in Luria-Bertani (LB) broth at 37°C. Cells were harvested when the culture reached late exponential growth phase at OD_{600} ~0.9. The strain mix was prepared by mixing equal volumes of the bacterial cultures. To determine the number of cells of *S. aureus* ST398 and *Salmonella* Typhimurium DT104 in the two cultures, dilutions of these were plated on LB agar, the plates incubated overnight at 37°C, and colony forming units (CFU) determined the following day. The strain mix was added at about 5% of the volume of the aliquot, and the added cell numbers of *S. aureus* and *S. enterica* Thyphimurium were calculated based on the CFU determinations.

DNA isolation
In a first step, seven DNA isolation procedures were examined, namely: InnuPure® C16 (Analytic Jena AG), MagNA Pure LC DNA isolation Kit III (Roche), Easy-DNA™ gDNA Purification Kit (Invitrogen), MP FastDNA™ Spin Kit (MP Biomedicals), PowerSoil® DNA Isolation kit (MoBio), QIAamp® DNA Stool

Mini Kit (Qiagen), QIAamp® DNA Stool Mini Kit (Qiagen) +Bead Beating (see Table 1, and details below). These methods were selected because they are widely used and represent a variety of isolation procedures involving manual or automated DNA isolation, DNA separation using filter-columns or magnetic beads, chemical or mechanical lysis, and phenol/chloroform-based or non-chloroform based isolations. Bead-beating steps were performed in a Qiagen TissueLyser II if not stated otherwise, and centrifugation steps were carried out in an Ole Dich 157.MP Microcentrifuge (Denmark). DNA isolation was performed on duplicate or triplicate aliquots, dependent on specimen availability. One to two isolation controls were included at each round of isolation.

InnuPure® C16, Analytic Jena AG (InnuPURE)
Automatic isolation with the InnuPURE–C16 robot using the InnuPURE Stool DNA Kit–IP–C16 according to the manufacture's instructions. Prior to the automatic isolation, a lysis step was performed according to the protocol for lysis of bacterial DNA from stool samples using a SpeedMILL PLUS provided by the manufacturer. The cell disruption process was carried out two times for 30 sec at 50 Hz ($50\ s^{-1}$). The DNA was eluted in 100 ul of buffer supplied with the kit.

MagNA Pure LC DNA isolation Kit III, Roche (MagNAPure)
Automatic isolation with the MagNA Pure LC instrument using the DNA Isolation Kit III (Bacteria, Fungi) according to the manufacture's instructions. The pre-isolation step for stool samples described in the protocol was performed before transferring the samples to the MagNA Pure LC. The protocol states a starting amount of a peanut-size sample, and in order to ensure consistency across isolations a starting amount of 0.25 g was chosen. The DNA was eluted in 100 ul of buffer supplied with the kit.

Easy-DNA™ gDNA Purification Kit, Invitrogen (EasyDNA)
The DNA isolation was performed according to the manufacturer's instructions with minor modifications. Pretreatment of the samples were performed following the protocol for small amounts of cells, tissues, or plant leaves. Initially, 0.25 g sample aliquots were dissolved in 1.5 ml 0.9% NaCl, respectively. The samples were centrifuged at 600xg for 3 minutes. The supernatant was transferred to new tubes and centrifuged at 8000xg for 10 minutes. After centrifugation, the supernatant was discarded and the pellet was resuspended in 200 µl PBS. 30 µl lysozyme (10 mg/ml) and 15 µl lysostaphin (10 mg/ml) were added and the samples were incubated at 37°C for 20 minutes shaking at 550 ppm, before adding 30 µl 10% SDS. The final pretreatment step included the addition of 15 µl proteinase K (20 mg/ml) and incubation at 37°C for 20 minutes. The final step in the isolation protocol was prolonged to an incubation for 1.5 (instead of 0.5) hours at 37°C. The DNA was eluted in 100 ul of buffer supplied with the kit.

MP FastDNA™ Spin Kit, MP Biomedicals (FastDNA)
The DNA isolation was performed according to the manufacturer's instructions with minor modifications. A centrifugation step at 3000xg for 2 minutes was

included to ensure proper settling of the silica matrix. The protocol suggested eluting the DNA in 50-100 µl DNase/Pyrogen-Free water, and here the DNA was eluted in 100 µl.

PowerSoil® DNA Isolation kit, MoBio Laboratories Inc. (PowerSoil.HMP)
The DNA isolation was performed according to the protocol employed in the Human Microbiome Project (HMP Protocol # 07-001 version 12), with a minor modification to the initial protocol step. The HMP protocol states to resuspend 2 ml fecal sample in 5 ml MoBio lysis buffer. Here, we resuspended 0.5 g sample in 1.25 ml MoBio lysis buffer (i.e. same ratio). Subsequently, the samples were centrifuged according to the HMP protocol and 1 ml of supernatant transferred to a garnet bead tube containing 0.75 ml MoBio buffer. The samples were heated at 65°C for 10 minutes followed by an additional heating step at 95°C for 10 minutes. The samples were processed further according to the HMP protocol including the modification at step 12, where the centrifugation step was prolonged to 2 minutes. The DNA was eluted in 100 ul of buffer supplied with the kit.

QIAamp® DNA Stool Mini Kit, Qiagen (QIAStool)
The DNA isolation was performed according to the manufacturer's protocol for isolation of DNA from stool for pathogen detection with minor modifications. In the lysis step, the samples were first heated at 70°C for 5 minutes and subsequently at 95°C for 5 minutes. The DNA was eluted in 100 µl elution buffer.

QIAamp® DNA Stool Mini Kit, Qiagen +Bead Beating (QIAStool+BB)
The DNA isolation was performed according to the manufacturer's protocol for isolation of DNA from stool for pathogen detection with minor modifications that included a bead-beating step. Sample aliquots of 0.2 g were mixed with 1.4 ml ASL buffer, respectively, and added to Lysing matrix A bead beating tubes (MP Biomedicals) and were briefly homogenized. The samples were treated in a Qiagen TissueLyser II at 30 f/s (Hz) three times for 30 seconds, with placement of the samples on ice in between bead beating steps. Subsequently, the samples were heated at 95°C for 15 minutes. The remaining steps were carried out according to the manufacturer's recommendations, and the DNA eluted in 100 ul elution buffer.

In a second step, a variety of modifications to two Qiagen kits were examined, namely the QIAamp® DNA Stool Mini Kit (QIAStool), and QIAamp® Fast DNA Stool Mini Kit (QIAFast). The latter was released to the market during the course of this study. The main difference between the QIAStool and the QIAFast kits relies in the way inhibitor compounds are being removed. In the QIAStool kit, InhibitEX tablets are being dissolved in the samples that are adsorbing the inhibitors and together are removed via centrifugation. The QIAFast kit contains an InhibitEX buffer to remove inhibitor compounds, and no tablets are required.

QIAamp® DNA Stool Mini Kit, Qiagen and Modifications

Five different protocols based on the QIAamp$^®$ DNA Stool Mini Kit were examined. i) QIAStool: see above, ii) QIAStool+BB.LMA: QIAStool+BB procedure using Lysing matrix A tubes (see above), iii) QIAStool+BB.LMA+2Trans: QIAStool+BB procedure using Lysing matrix A tubes (see above) with modifications. To reduce the loss of sample, the double amount of supernatant was transferred to proteinase K (i.e. 400 µl instead of 200 µl). The volumes of Proteinase K, buffer AL and ethanol were doubled, respectively. Due to the increased volume, the passing of the sample through the spin columns is performed in two centrifugation steps. The DNA was washed twice before elution in 100 µl elution buffer. iv) QIAStool+PreT+BB.LMA: QIAStool+BB procedure using Lysing matrix A tubes (see above) with modifications. An increased starting sample amount was used and pre-treated. 0.5g of sample was mixed with 1.5 ml 0.9% NaCl solution. After homogenization by vortexing, the samples were centrifuged at 600 x g for 3 minutes to settle large particles. The supernatant was centrifuged at 8000 x g for 10 minutes to pellet microbial cells. The pellet was resuspended in 200 µl PBS and transferred to Lysing Matrix A bead beating tubes. v) QIAStool+PreT+BB.LMA+2Trans: QIAStool+BB procedure using Lysing matrix A tubes (see above) with modifications described in iii) and iv).

QIAamp$^®$ Fast DNA Stool Mini Kit, Qiagen and Modifications
Six different protocols (i–vi) based on the QIAamp$^®$ Fast DNA Stool Mini Kit were examined using five different bead types (ii–vi). i) QIAFast: The DNA isolation was performed according to the manufacturer's protocol for isolation of DNA from stool for pathogen detection with minor modifications. In the lysis-step, the samples were first heated at 70°C for 5 minutes and subsequently at 90°C for 5 minutes. The DNA was eluted in 100 µl elution buffer for 2 minutes. ii) QIAFast+BB.LMA+2Trans: The DNA isolation was performed according to the manufacturer's protocol for isolation of DNA from stool for pathogen detection with minor modifications that included a bead-beating step. Sample aliquots of 0.2 g were mixed with 1 ml InhibitEX buffer, respectively, and added to Lysing matrix A bead beating tubes (MP Biomedicals) and are briefly homogenized. The samples are treated in a Qiagen TissueLyser II at 30 f/s (Hz) three times for 30 seconds, with placement of the samples on ice in between bead beating steps. Subsequently, the samples are heated at 95°C for 7 minutes. Similar to the modifications described above, following the bead-beating and heating steps, the double amount of supernatant was transferred to proteinase K (i.e. 400 µl instead of 200 µl). The volumes of proteinase K, Buffer AL and ethanol were also doubled. The passing of the sample through the filter columns were subsequently carried out in two centrifugation steps rather than one, to accommodate the increased sample volume. The remaining steps were carried out according to the manufacturer's recommendations, and the DNA eluted in 100 ul elution buffer. A laboratory protocol for this procedure can be found at https://dx.doi.org/10.6084/m9.figshare.3475406.v1. iii) QIAFast+BB.LTS+2Trans: Same procedure as described in ii) with Pathogen Lysis Tubes S (Qiagen). iv) QIAFast+BB.LTL+2Trans: Same procedure as described in ii) with Pathogen Lysis Tubes L (Qiagen). v) QIAFast+BB.GBT+2Trans: Same procedure as

described in ii) with Garnet Bead Tubes (MoBio). vi) QIAFast+BB.AAB+2Trans: Same procedure as described in ii) with A&A Bead Tubes (A&A Biotechnology, Gdynia, Poland).

Together, the evaluation and improvements of DNA isolation methods were carried out in a step-wise approach. In the first step, seven DNA extraction kits were evaluated using human feces, pig feces, and hospital sewage (Figures 1–4, Supplemental Figures S1-S4, and Supplemental Tables S1+S2). The standard and modified procedures based on the QIAStool and QIAFast methods were tested using a second set of pig fecal samples (Figure 5A, and Supplemental Figure S5, and Supplemental Table S3). Upon evaluation of the different DNA isolation methods, promising procedures were selected and examined using a new set of human feces, pig feces, and hospital sewage (Figure 5B).

DNA quantitation and quality assessment
Subsequent to DNA isolation, the DNA was portioned into 10 µl aliquots to prevent repeated freeze-thawing cycles, and stored at -20°C. DNA concentrations were measured using Qubit® dsDNA BR Assay Kit on a Qubit® 2.0 Fluorometer (Invitrogen, Carlsbad, CA). As DNA extracts can contain contaminants, such as proteins or other organic molecules that can affect downstream procedures such as DNA amplifications in PCR, we determined the DNA purity by measuring the ratios of absorbance at 260/280 and 260/230, respectively, using a NanoDrop 1000 Spectrophotometer (Thermo Scientific, Pittsburgh, USA). DNA extracts with a 260/280 ratio between ~1.7 to ~ 2.0, and 260/230 ration between ~2.0 to ~2.2 are regarded as "pure". The stability of the DNA in the extracts was determined by measuring the DNA concentration after 2 and 7 days incubation at 22°C. A decrease in DNA concentration over time can indicate the presence of DNases in the extract.

16S rRNA gene profiling
16S rRNA amplicon libraries were generated using a two-step protocol similar as described in Part # 15044223 Rev. B by Illumina (http://www.illumina.com/con tent/dam/illumina-support/documents/documentation/chemistry_documenta tion/16s/16s-metagenomic-library-prep-guide-15044223-b.pdf). In a first PCR, the V4 region of the 16S rRNA genes were amplified using the universal primers (515f 5'-TGCCAGCAGCCGCGGTAATAC (1) and 806r 5'-GGACTACNNGGGTATCTAAT (2). Each 20-µl PCR reaction contained 2 µl of 10 x AccuPrime PCR Buffer II (15mM MgCl2, Invitrogen), 1 µl (10 µM) of the primers, 0.12 µl AccuPrime Taq DNA polymerase (2 units/µl, Invitrogen), 1 µl template DNA and 14.88 µl ddH2O. PCR conditions: denaturation at 94°C for 2 min; 30 cycles at 94°C for 20 s, 56°C for 20 s, 68°C for 30 s; followed by 68°C for 5 min, and 3 min at 70 °C. Subsequently, the PCR products were placed on ice to prevent hybridization between PCR products and nonspecific amplicons. Samples were quantified using Quant-iT™ PicoGreen® dsDNA Assay Kit (Invitrogen, Carlsbad, CA) on a Lightcycler 96 (Roche, Mannheim, Germany) and adjusted to equal concentrations. In the second PCR the same conditions were

used as in the first round, except the PCR was reduced to 15 cycles and the primers had a unique adaptor/linker/index sequence per sample (3). The PCR products were purified using Agencourt AmPure XP beads (Beckman Coulter Inc, A63881), and concentrations were measured using Quant-iT™ PicoGreen® dsDNA Assay Kit on a Lightcycler 96. The samples were pooled in equal concentrations, and concentrated using 'DNA clean and concentrator-5 kit' (Zymo Research, Orange, CA). Paired-end 2 × 250 bp sequencing of barcoded amplicons was performed on a MiSeq machine running v2 chemistry (Illumina Inc., San Diego, CA, USA) at University of Copenhagen, Section of Microbiology.

The primer sequences were trimmed, quality filtering performed, and paired sequences assembled using the UPARSE pipeline (4). Low quality reads were removed with a maximum expected error threshold of 0.5 (maxee). Sequences were barcoded and pooled before dereplication and removal of duplicates (-minseqlength 64). Prior to clustering of the OTUs the dereplicated reads were sorted according to abundance, and singeltons where removed. Chimera filtering was performed using UCHIME (5) with rdp_gold.fa as reference database. The reads were mapped back to OTUs, including singletons, at a 97% identity level and an OTU-table was generated using uc2otutab.py. Using QIIME1.8.0 (6), taxonomy was assigned with uclust using assign_taxonomy.py based on the Greengenes 13.8 reference database. The average number of reads per sample was 192965, and the read length was between 186-251 bp. The average number of reads in the isolation controls was 34063 and the majority of these reads were affiliated with the two strains used for spiking (Enterobacteriaceae and Staphylococcaceae), as well as dominant taxa that were present in the complex samples, such as Ruminococcaceae, Prevotellaceae, and Bacteroidales. Ecological diversity estimates and microbial community comparisons were performed using the relevant scripts provided by QIIME, phyloseq, and R (6-8). For the estimation of bacterial diversity and richness (Fig. 1C, and Fig. 5A), and principal coordinate analysis (Fig. 2A-C, and Fig. S3) the samples were rarefied to 800 reads per sample. The abundance of Gram-positive and Gram-negative bacteria was predicted at order levels based on information from the literature. For some bacteria (mainly Firmicutes), the Gram status could not be assigned at this level, and for those the family level was used instead.

Metagenomics
A subset of thirty-nine DNA extracts was subjected to metagenomic sequencing. The samples were prepared and sequenced following the Nextera XT DNA Library Preparation Guide for the MiSeq system, Part # 15031942 Rev. D (http://support.illumina.com/content/dam/illumina-support/documents/documentation/chemistry_documentation/samplepreps_nextera/nextera-xt/nextera-xt-library-prep-guide-15031942-01.pdf), using paired-end v2 2×250bp sequencing. The taxonomic microbiome compositions were determined through the use of the MGmapper pipeline (9). The MGmapper package is available for download at www.cbs.dtu.dk/public/MGmapper/. The analysis consisted of three main steps: i) Pre-processing and quality trimming of raw reads, ii) Mapping of reads to

reference sequence databases, and iii) Analysis of read count data. In the first step, cutadapt (10) was employed for adapter sequence removal, trimming of low-quality bases from the ends of the reads (-q 30), and removal of reads that were shorter than 30 bp. In a second step, the remaining paired-end reads were mapped in chain-mode to four databases: 1. complete bacterial genomes, 2. draft bacterial genomes, 3. MetaHit Assembly (http://www.sanger.ac.uk/resources/downloads/bacteria/metahit/, July 2014), and 4. Human Microbiome assemby (http://www.hmpdacc.org/resources/data_browser.php, July 2014) using the BWA-MEM algorithm (http://bio-bwa.sourceforge.net). For the analysis in the present study only the reads mapping to the two primary bacterial databases (complete and draft bacterial genomes) were considered. These two databases were composed of 2685 complete and 22224 draft bacterial and archaeal genomes obtained from Genbank on July 2014 and December 2014, respectively. The order by which the databases are specified in chain-mode is important, as reads that exhibit a significant hit to the previous reference database are removed before mapping to the next database. Samtools (10) was used to remove singletons and all reads that did not map as pairs. An alignment of a read pair with a region in a genome was considered a hit only if the sum of the alignment scores (SAS) was higher than any SAS values from other hits in the database. In the third step, the alignments were filtered based on the Fraction of Matches+Mismatches (FMM) threshold, i.e. the fraction of a read that should align. Here, the default FMM threshold of 80% was used. From 96 155 142 raw read pairs, 7 567 574 read pairs mapped genomes in the two reference databases. The final read count table was composed of 9436 bacterial and archaeal reference strains with an average of 69952 mapped read pairs per sample. For each sample, the read counts were normalized according to the genome length of the respective genomes in the database, and for sequencing depth using total sum scaling.

Differential abundance analysis
In order to test for the differential abundance of taxa that may drive the differences observed between the communities derived from the different DNA isolation procedures, we performed DESeq2 analyses. The (unnormalized) read count tables from the 16S rRNA gene profiling and metagenomics sequence analysis, respectively, were aggregated to the family level in R (v. 3.2.3, 64bit) (8) We performed an analysis that allows for varied sequencing depth, similar as suggested previously (12), and carried out two-sided Wald tests as implemented in the DESeq2 package (v. 1.10.1) (13). The size factors were determined by DESeq2 from the read count tables. An example for such an analysis is available from https://figshare.com/s/99fc2ac5e1ed54fa641c.
When testing the effect of added strain mix, we included the samples to which the strain mix was added as well as the corresponding samples to which no strain mix was added and accounted for DNA isolation method and sample matrix type. When testing the effects of the DNA isolation method, we analyzed the data from the three types of fecal specimen separately and extracted results from all two-wise comparisons of DNA isolation methods. For each DESeq2 test,

p-values were adjusted for the false-discovery rate (FDR) using the Benjamini-Hochberg procedure (14). As recommended by DESeq2, comparisons with an FDR below 0.1 were considered significant. For the visualization of the data, the read count data were variance-stabilized using the DESeq2 regularized log (rlog) transformation. This transformation also accounts for sequencing depth differences, allowing inter-sample comparisons of taxa.

Quantification of strain mix
The samples that were spiked with the strain mix composed of *S. enterica* Typhimurium DT104 and *S. aureus* ST398 were extracted, sequenced, and analyzed together with the non-spiked samples. For each type of specimen and isolation method, the relative abundance of Enterobacteriaceae and Staphylococcaceae for 16S rRNA gene profiling and metagenomics, respectively, were determined. Our differential abundance analysis using DESeq2 confirmed, that these two strains were present in significantly higher abundance in the spiked samples than in the not spiked samples for 16S rRNA gene profiling: Enterobacteriaceae adjusted P-value 3.08^{-30} and Staphylococcaceae adjusted P-value 2.13^{-10}; and for metagenomics: Enterobacteriaceae adjusted P-value 1.74^{-77} and Staphylococcaceae adjusted P-value 1.07^{-4}. The average relative abundance values from the samples without added strain mix were subtracted from the corresponding samples to which the strain mix was added. Subsequently, the 16S rRNA gene copy numbers of the two added strains were taken into account with 5 for *S. aureus* and 7 for *S. enterica* (for 16S rRNA gene profiling). The ratios between Enterobacteriaceae and Staphylococcaceae were determined for each sample matrix and isolation method, and compared to the *S. enterica* Typhimurium DT104 / *S. aureus* ST398 ratio of CFU that were added to the original samples.

II) Supplemental Figures

FIG S1. Microbial community composition. The Top 10 most abundant families for the human fecal (A), pig fecal (B), and hospital sewage (C) samples based on (i) 16S rRNA gene profiling, (ii) metagenomics analysis that include normalization based on reference genome size, and (iii) metagenomics analysis without normalization according

to genome size. For details regarding sequence data analysis and normalization see Materials and Methods.

FIG S2. Microbial community composition based on predicted Gram-staining. Gram-positive and Gram-negative affiliations were assigned at the order-level based on information found in the literature. For some taxa the Gram-staining status was unknown.

FIG S3. Microbial community dissimilarity. The dissimilarity between the microbiotas from the human, pig, and sewage samples was examined using Principal Coordinates Analysis of Bray-Curtis distances based on the 16S rRNA gene count data. For the PCoA Bray-Curtis ordination analysis only samples with a minimum of 800 reads were included.

FIG S4. Detection of spiked bacteria using metagenomics. The human fecal (A), and pig fecal (B) samples were spiked with a strain mix composed of *Salmonella enterica* serotype Typhimurium DT104 and *Staphylococcus aureus* ST398 in a CFU ratio of 1.02. These two sample matrices, as well as aliquots of the strain mix (C) were extracted using three different DNA extraction methods. The two strains were detected by metagenomics analysis, and their ratios determined. For details, see Materials and Methods. An asterisk indicates that the values for the particular DNA extraction of the strain mix (D) are based on single measurements. All other values are based on averages from duplicate or triplicate measurements. The dashed line indicates the ratio of the strain mix based on CFU determinations.

FIG 5. Comparison between QIAStool and QIAFast DNA extraction methods by metagenomics. Pig feces was extracted using the QIAamp® DNA Stool Mini and QIAamp® Fast DNA Stool Mini kits, and analyzed using metagenomics. The alpha diversity (Chao 1 and Shannon index) was determined at species-level. The microbial community composition was examined at genus-level and the relative abundance of the Top 10 most abundant taxa are shown here.

Supplemental References

1. **Yu Y**, **Lee C**, **Kim J**, **Hwang S**. 2005. Group-specific primer and probe sets to detect methanogenic communities using quantitative real-time polymerase chain reaction. Biotechnol Bioeng **89**:670–679.

2. **Sundberg C**, **Al-Soud WA**, **Larsson M**, **Alm E**, **Yekta SS**, **Svensson BH**, **Sørensen SJ**, **Karlsson A**. 2013. 454 pyrosequencing analyses of bacterial and archaeal richness in 21 full-scale biogas digesters. FEMS Microbiology Ecology **85**:612–626.

3. **Riber L**, **Poulsen PHB**, **Al-Soud WA**, **Skov Hansen LB**, **Bergmark L**, **Brejnrod A**, **Norman A**, **Hansen LH**, **Magid J**, **Sørensen SJ**. 2014. Exploring the immediate and long-term impact on bacterial communities in soil amended with animal and urban organic waste fertilizers using pyrosequencing and screening for horizontal transfer of antibiotic resistance. FEMS Microbiology Ecology **90**:206–224.

4. **Edgar RC**. 2013. UPARSE: highly accurate OTU sequences from microbial amplicon reads. Nature Methods **10**:996–998.

5. **Edgar RC**, **Haas BJ**, **Clemente JC**, **Quince C**, **Knight R**. 2011. UCHIME improves sensitivity and speed of chimera detection. Bioinformatics **27**:2194–2200.

6. **Caporaso JG**, **Kuczynski J**, **Stombaugh J**, **Bittinger K**, **Bushman FD**, **Costello EK**, **Fierer N**, **Peña AG**, **Goodrich JK**, **Gordon JI**, **Huttley GA**, **Kelley ST**, **Knights D**, **Koenig JE**, **Ley RE**, **Lozupone CA**, **McDonald D**, **Muegge BD**, **Pirrung M**, **Reeder J**, **Sevinsky JR**, **Turnbaugh PJ**, **Walters WA**, **Widmann J**, **Yatsunenko T**, **Zaneveld J**, **Knight R**. 2010. QIIME allows analysis of high-throughput community sequencing data. Nature Methods **7**:335–336.

7. **McMurdie PJ**, **Holmes S**. 2013. phyloseq: An R Package for Reproducible Interactive Analysis and Graphics of Microbiome Census Data. PLoS ONE **8**:e61217.

8. **Team RDC**. 2014. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.

9. **Petersen TN**, **Bælum J**, **Lukjancenko O**, **Geertz-Hansen HM**, **Thomsen MCF**, **Sperotto MM**, **Lund O**, **Aarestrup FM**, **Sicheritz-Ponten T**. 2015. MGmapper: an automated pipeline for mapping and stratification of metagenomics sequence data. Submitted.

10. **Martin M**. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. EMBnetjournal **17**:10–12.

11. **Li H**, **Handsaker B**, **Wysoker A**, **Fennell T**, **Ruan J**, **Homer N**, **Marth G**, **Abecasis G**, **Durbin R**, **1000 Genome Project Data Processing Subgroup**. 2009. The Sequence Alignment/Map format and SAMtools. Bioinformatics

25:2078–2079.

12. **McMurdie PJ**, **Holmes S**. 2014. Waste Not, Want Not: Why Rarefying Microbiome Data Is Inadmissible. PLOS Computational Biology **10**:e1003531.

13. **Love MI**, **Huber W**, **Anders S**. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology **15**:31.

14. **Benjamini Y**, **Hochberg Y**. 1995. Controlling the False Discovery Rate: a Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B (Methodological) **57**:289–300.

Table S1A. DNA concentration, purity, and stability.

		HUMAN										PIG										SEWAGE									
		Nucleic acid quantitation						DNA stability at 22 degree C****				Nucleic acid quantitation						DNA stability at 22 degree C****				Nucleic acid quantitation						DNA stability at 22 degree C****			
		DNA concentration		DNA Purity				DNA conc. after 2 days		DNA conc. after 7 days		DNA concentration		DNA Purity				DNA conc. after 2 days		DNA conc. after 7 days		DNA concentration		DNA Purity				DNA conc. after 2 days		DNA conc. after 7 days	
	DNA extraction method*	DNA conc. [ng/ul]	SE	DNA A260/280**	SE	DNA A260/230***	SE	DNA conc. [ng/ul]	SE	DNA conc. [ng/u	SE	DNA conc. [ng/ul]	SE	DNA A260/280**	SE	DNA A260/230***	SE	DNA conc. [ng/ul]	SE	DNA conc. [ng/u	SE	DNA conc. [ng/ul]	SE	DNA A260/280**	SE	DNA A260/230***	SE	DNA conc. [ng/ul]	SE	DNA conc. [ng/u	SE
A	InnuPURE Stool DNA [A]	2.23	0.73	2.15	0.23	1.19	0.54	ND	ND	ND	ND	13.3	0.7	1.99	0.03	1.81	0.01	ND	ND	ND	ND	<0.01	-	<0	3.22	1.28	0.36	ND	ND	ND	ND
E	MagNA Pure LC DNA isolation Kit [A]	42.45	9.45	1.52	0.01	0.75	0.02	33.75	3.15	33.35	0.65	4.66	4.65	1.62	0.14	0.83	0.09	9.88	0.22	9.69	0.28	13.75	1.55	1.76	0.16	1.12	0.11	13.35	0.75	14.15	0.55
G	Easy-DNA™ gDNA Purification Kit	78.27	4.48	1.97	0.01	2.02	0.02	61.93	15.3	33.47	6.3	50.43	8.29	1.77	0.02	1.55	0.06	68.7	5.07	74.57	4.14	6.19	1.32	1.77	0.03	0.53	0.07	4.85	0.92	4.5	0.71
C	MP FastDNA™ Spin Kit	11.7	0.4	2	0.03	0.36	0.05	ND	ND	ND	ND	14	0.4	1.92	0.03	0.98	0.09	ND	ND	ND	ND	1.5	0.07	2.4	0.27	0.02	0	ND	ND	ND	ND
B	PowerSoil® DNA Isolation kit	1.85	0.1	1.46	0.02	1.46	0.02	ND	ND	ND	ND	1.24	0.24	1.52	0.1	0.41	0.005	ND	ND	ND	ND	<0.01	-	1.96	0.34	0.53	0.06	ND	ND	ND	ND
F	QIAamp® DNA Stool Mini Kit	30.43	1.22	2.08	0.02	1.73	0.07	28.33	0.24	28.07	0.55	25.33	0.64	1.98	0.02	1.38	0.07	23.6	0.72	23.5	0.78	<0.01	-	2.25	0.07	0.26	0.02	<0.01	-	<0.01	-
D	QIAamp® DNA Stool Mini Kit +BB	13.35	1.25	1.98	0.12	0.75	0.12	ND	ND	ND	ND	26.75	13.03	2.06	0.08	1.03	0.18	ND	ND	ND	ND	1.2	0.04	2.48	0.24	0.17	0.02	ND	ND	ND	ND

*Duplicate sample aliquods were extracted using procedure A, B, C, D, E. Triplicate sample aliquods were extracted using procedure F, G.

** Expected ~1.7 to ~ 2.0

*** Expected ~2.0 to ~2.2

****Expected >95% after 2-7 days incubation at room temperature

ND=Not determined

Table S1B. Microbiome richness and diversity.

	DNA extraction procedure*	HUMAN								PIG								SEWAGE							
		16S rRNA gene profiling				Metagenomics				16S rRNA gene profiling				Metagenomics				16S rRNA gene profiling				Metagenomics			
		Richness		Diversity		Richness		Diversity		Richness		Alpha Diversity		Richness		Alpha Diversity		Richness		Alpha Diversity		Richness		Alpha Diversity	
		Chao1	SE	Shannon index	SE	Chao1	SE	Shannon index	SE	Chao1	SE	Shannon index	SE	Chao1	SE	Shannon index	SE	Chao1	SE	Shannon index	SE	Chao1	SE	Shannon index	SE
A	InnuPURE Stool DNA [A]	151.25	0.58	4.56	0.07	ND	-	ND	-	320.78	9.46	5.82	0.08	ND	-	ND	-	577.67	-	7.09	-	ND	-	ND	-
E	MagNA Pure LC DNA isolation Kit [A]	134.59	9.21	3.62	0.36	250.34	12.44	2.26	0.14	387.82	22.04	6.52	0.02	616.45	45.79	4.84	0.11	455.71	7.69	6.78	0.09	ND	-	ND	-
G	Easy-DNA™ gDNA Purification Kit	90.52	14.66	3.88	0.18	309.43	9.75	3.90	0.02	302.63	20.15	5.88	0.14	593.61	12.94	4.52	0.11	501.08	18.01	6.83	0.11	ND	-	ND	-
C	MP FastDNA™ Spin Kit	161.82	5.33	5.02	0.14	ND	-	ND	-	109.9	7.52	3.37	0.06	ND	-	ND	-	499.51	7.48	6.96	0.02	ND	-	ND	-
B	PowerSoil® DNA Isolation kit	160.9	-	4.38	-	ND	-	ND	-	431.11	17.86	6.77	0.16	ND	-	ND	-	505.42	47.02	6.71	0.12	ND	-	ND	-
F	QIAamp® DNA Stool Mini Kit	117.04	2.34	3.2	0.02	243.49	8.37	1.83	0.02	355.07	1.14	6.33	0.02	508.55	4.61	3.48	0.11	483.96	7.48	6.7	0.01	ND	-	ND	-
D	QIAamp® DNA Stool Mini Kit +BB	169.69	8.62	4.59	0.01	ND	-	ND	-	373.97	-	6.36	-	ND	-	ND	-	503.87	18.27	6.97	0.18	ND	-	ND	-

*Duplicate sample aliquods were extracted using procedure A, B, C, D, E. Triplicate sample aliquods were extracted using procedure F, G.

ND=Not determined

Table S2A. Difference abundance of families - Human fecal microbial community

baseMean	log2FoldChange	lfcSE	stat	pvalue	padj	family	compare
8087.80815	9.58965882	0.59563182	16.0999775	2.55E-58	7.49E-57	Bifidobacteriaceae	EasyDNA vs QIAStool
261.894528	9.6387835	0.59914643	16.0875255	3.12E-58	7.49E-57	[Mogibacteriaceae]	EasyDNA vs QIAStool
261.894528	6.9943178	0.50812435	13.7649727	4.14E-43	2.19E-41	[Mogibacteriaceae]	EasyDNA vs InnuPURE
261.894528	5.98959231	0.47453074	12.6221376	1.59E-36	9.88E-35	[Mogibacteriaceae]	EasyDNA vs FastDNA
261.894528	5.68437867	0.45525408	12.4861675	8.88E-36	4.26E-34	[Mogibacteriaceae]	EasyDNA vs QIAStool+BB
8087.80815	8.06288475	0.6580903	12.251943	1.64E-34	4.35E-33	Bifidobacteriaceae	EasyDNA vs InnuPURE
8087.80815	8.10323384	0.66343913	12.2139824	2.62E-34	1.52E-32	Bifidobacteriaceae	EasyDNA vs MagNAPure
261.894528	7.46735031	0.63949976	11.6768617	1.67E-31	4.85E-30	[Mogibacteriaceae]	EasyDNA vs MagNAPure
195.592384	7.11547535	0.63330982	11.2353782	2.73E-29	4.37E-28	Coriobacteriaceae	EasyDNA vs QIAStool
8087.80815	8.22252999	0.81577634	10.0793925	6.81E-24	3.13E-22	Bifidobacteriaceae	EasyDNA vs PowerSoil.HMP
118.367167	4.14939619	0.43086714	9.6303379	5.95E-22	7.14E-21	Streptococcaceae	EasyDNA vs QIAStool
261.894528	6.50010878	0.67196014	9.67335475	3.91E-22	9.00E-21	[Mogibacteriaceae]	EasyDNA vs PowerSoil.HMP
195.592384	6.52674567	0.70556721	9.25035285	2.24E-20	3.95E-19	Coriobacteriaceae	EasyDNA vs InnuPURE
8087.80815	6.05120873	0.6598584	9.17046551	4.71E-20	1.70E-18	Bifidobacteriaceae	FastDNA vs QIAStool
750.272969	-5.5100567	0.62548793	-8.8092135	1.26E-18	1.67E-17	Porphyromonadaceae	EasyDNA vs InnuPURE
750.272969	-4.8876587	0.57173354	-8.5488402	1.24E-17	1.19E-16	Porphyromonadaceae	EasyDNA vs QIAStool
750.272969	-5.3905037	0.62605815	-8.6102285	7.29E-18	1.41E-16	Porphyromonadaceae	EasyDNA vs MagNAPure
195.592384	6.37027321	0.76210202	8.35881943	6.33E-17	9.19E-16	Coriobacteriaceae	EasyDNA vs MagNAPure
195.592384	5.59941114	0.66873123	8.37318631	5.61E-17	1.35E-15	Coriobacteriaceae	EasyDNA vs QIAStool+BB
14192.9873	4.1205321	0.50198036	8.20855245	2.24E-16	1.79E-15	Ruminococcaceae	EasyDNA vs QIAStool
3174.43554	-3.2374316	0.4150328	-7.8004235	6.17E-15	9.84E-14	Veillonellaceae	FastDNA vs QIAStool
195.592384	5.37934136	0.69281817	7.76443455	8.20E-15	9.84E-14	Coriobacteriaceae	FastDNA vs QIAStool
8087.80815	5.04874196	0.65458206	7.71292439	1.23E-14	1.97E-13	Bifidobacteriaceae	EasyDNA vs QIAStool+BB
195.592384	7.26194948	0.9934254	7.31000989	2.67E-13	4.10E-12	Coriobacteriaceae	EasyDNA vs PowerSoil.HMP
188.732855	-5.3331196	0.7367495	-7.2387149	4.53E-13	5.25E-12	[Barnesiellaceae]	EasyDNA vs MagNAPure
3052.94009	-4.8906164	0.69105732	-7.0770053	1.47E-12	1.56E-11	Bacteroidaceae	EasyDNA vs InnuPURE
3052.94009	-4.7809223	0.6911788	-6.9170558	4.61E-12	4.46E-11	Bacteroidaceae	EasyDNA vs MagNAPure
680.114734	-5.3983716	0.79094213	-6.8252422	8.78E-12	7.75E-11	Pasteurellaceae	EasyDNA vs InnuPURE
3052.94009	-4.2181346	0.62453115	-6.754082	1.44E-11	9.86E-11	Bacteroidaceae	EasyDNA vs QIAStool
750.272969	-5.1128	0.75653447	-6.7581851	1.40E-11	1.61E-10	Porphyromonadaceae	EasyDNA vs PowerSoil.HMP
3174.43554	-3.1274017	0.45366454	-6.8936438	5.44E-12	2.23E-10	Veillonellaceae	FastDNA vs MagNAPure
118.367167	3.1468372	0.46933577	6.70487405	2.02E-11	2.42E-10	Streptococcaceae	EasyDNA vs QIAStool+BB
188.732855	-4.4807982	0.6822668	-6.5675161	5.12E-11	3.07E-10	[Barnesiellaceae]	EasyDNA vs QIAStool
50.0848862	-6.0497647	0.91762758	-6.5928322	4.32E-11	3.58E-10	[Odoribacteraceae]	EasyDNA vs MagNAPure
8087.80815	-4.5409169	0.65998484	-6.8803351	5.97E-12	5.73E-10	Bifidobacteriaceae	QIAStool vs QIAStool+BB
12.0964552	6.19822102	0.96500816	6.42297263	1.34E-10	7.13E-10	Actinomycetaceae	EasyDNA vs QIAStool
118.367167	3.04916807	0.47421688	6.429902	1.28E-10	9.67E-10	Streptococcaceae	EasyDNA vs InnuPURE
750.272969	-4.0416602	0.62575935	-6.4588091	1.06E-10	1.01E-09	Porphyromonadaceae	EasyDNA vs QIAStool+BB
50.0848862	-5.8312903	0.91432374	-6.3777085	1.80E-10	1.19E-09	[Odoribacteraceae]	EasyDNA vs InnuPURE
8087.80815	4.56478376	0.72120492	6.32938523	2.46E-10	5.05E-09	Bifidobacteriaceae	FastDNA vs MagNAPure
188.732855	-4.5085906	0.73616033	-6.1244683	9.10E-10	5.36E-09	[Barnesiellaceae]	EasyDNA vs InnuPURE
8087.80815	4.52443467	0.7163253	6.31617319	2.68E-10	6.19E-09	Bifidobacteriaceae	FastDNA vs InnuPURE
195.592384	4.79061167	0.75894457	6.31220231	2.75E-10	6.19E-09	Coriobacteriaceae	FastDNA vs InnuPURE
261.894528	-3.9544048	0.64127777	-6.1664462	6.98E-10	3.35E-08	[Mogibacteriaceae]	QIAStool vs QIAStool+BB
3174.43554	2.51360524	0.41567177	6.04709158	1.47E-09	4.57E-08	Veillonellaceae	EasyDNA vs FastDNA
680.114734	-4.0974239	0.72004633	-5.6905004	1.27E-08	6.08E-08	Pasteurellaceae	EasyDNA vs QIAStool
52.7372451	-4.7893782	0.84104417	-5.6945621	1.24E-08	6.56E-08	Rikenellaceae	EasyDNA vs InnuPURE
680.114734	-4.5127186	0.79165921	-5.7003298	1.20E-08	8.67E-08	Pasteurellaceae	EasyDNA vs MagNAPure
55.8474481	4.94136357	0.85181641	5.80097251	6.59E-09	9.01E-08	Lactobacillaceae	FastDNA vs MagNAPure
195.592384	4.63413921	0.81137436	5.71146864	1.12E-08	1.15E-07	Coriobacteriaceae	FastDNA vs MagNAPure
118.367167	3.5108296	0.62265509	5.6384821	1.72E-08	1.58E-07	Streptococcaceae	EasyDNA vs PowerSoil.HMP
129.697728	-2.7013853	0.46425926	-5.8187	5.93E-09	1.90E-07	Erysipelotrichaceae	QIAStool vs QIAStool+BB
261.894528	3.6491912	0.65484847	5.57257343	2.51E-08	2.26E-07	[Mogibacteriaceae]	FastDNA vs QIAStool
3052.94009	-4.6400461	0.84608129	-5.4841611	4.15E-08	3.19E-07	Bacteroidaceae	EasyDNA vs PowerSoil.HMP
14192.9873	3.01986302	0.55817717	5.4102231	6.29E-08	5.04E-07	Ruminococcaceae	EasyDNA vs QIAStool+BB
12.8572515	4.68299174	0.87546116	5.34917137	8.84E-08	6.36E-07	Turicibacteraceae	FastDNA vs QIAStool
129.697728	2.47103163	0.46631338	5.29907944	1.16E-07	6.98E-07	Erysipelotrichaceae	FastDNA vs QIAStool
14192.9873	2.93847088	0.55833808	5.26288821	1.42E-07	9.14E-07	Ruminococcaceae	EasyDNA vs MagNAPure
14192.9873	2.89591167	0.55820633	5.18788757	2.13E-07	1.02E-06	Ruminococcaceae	EasyDNA vs InnuPURE
8087.80815	3.53845009	0.6544545	5.40671674	6.42E-08	1.06E-06	Bifidobacteriaceae	EasyDNA vs FastDNA
750.272969	-3.3818974	0.62676068	-5.3958354	6.82E-08	1.06E-06	Porphyromonadaceae	EasyDNA vs FastDNA
12.0964552	5.12847194	0.99982132	5.12938846	2.91E-07	1.28E-06	Actinomycetaceae	EasyDNA vs InnuPURE
188.732855	-4.5189874	0.86857022	-5.2027888	1.96E-07	1.29E-06	[Barnesiellaceae]	EasyDNA vs PowerSoil.HMP

128.300954	-3.9366676	0.7636346	-5.1551718	2.53E-07	1.47E-06	[Paraprevotellaceae]	EasyDNA vs MagNAPure
52.7372451	-4.3339548	0.84802512	-5.1106443	3.21E-07	1.69E-06	Rikenellaceae	EasyDNA vs MagNAPure
8087.80815	4.6840799	0.86240099	5.43144078	5.59E-08	1.87E-06	Bifidobacteriaceae	FastDNA vs PowerSoil.HMP
195.592384	5.52581549	1.03014075	5.36413637	8.13E-08	1.87E-06	Coriobacteriaceae	FastDNA vs PowerSoil.HMP
18839.0105	-4.1923769	0.82982709	-5.052109	4.37E-07	1.91E-06	Prevotellaceae	EasyDNA vs QIAStool
833.096381	-4.9687451	0.98221289	-5.0587252	4.22E-07	2.04E-06	Alcaligenaceae	EasyDNA vs MagNAPure
3052.94009	-3.5171962	0.69110702	-5.0892208	3.60E-07	2.21E-06	Bacteroidaceae	EasyDNA vs QIAStool+BB
188.732855	-3.7434477	0.73628464	-5.0842398	3.69E-07	2.21E-06	[Barnesiellaceae]	EasyDNA vs QIAStool+BB
12.0964552	4.43642641	0.88398443	5.0186703	5.20E-07	2.77E-06	Actinomycetaceae	EasyDNA vs QIAStool+BB
717.960343	2.58858006	0.5178777	4.99843893	5.78E-07	2.77E-06	Verrucomicrobiaceae	EasyDNA vs QIAStool+BB
55.8474481	3.13191814	0.62594371	5.00351402	5.63E-07	2.90E-06	Lactobacillaceae	FastDNA vs QIAStool
14192.9873	2.87901713	0.55820744	5.15761149	2.50E-07	3.10E-06	Ruminococcaceae	EasyDNA vs FastDNA
118.367167	2.38319975	0.48150613	4.94946913	7.44E-07	3.32E-06	Streptococcaceae	EasyDNA vs MagNAPure
129.697728	2.12324888	0.43267331	4.90727959	9.23E-07	3.69E-06	Erysipelotrichaceae	EasyDNA vs QIAStool
55.8474481	4.67527328	0.85115964	5.49282774	3.96E-08	3.80E-06	Lactobacillaceae	QIAStool+BB vs MagNAPure
118.367167	2.38330929	0.46888891	5.08288693	3.72E-07	3.84E-06	Streptococcaceae	EasyDNA vs FastDNA
128.300954	-3.3578675	0.69956976	-4.7999038	1.59E-06	5.86E-06	[Paraprevotellaceae]	EasyDNA vs QIAStool
52.7372451	-3.7686587	0.79504946	-4.7401563	2.14E-06	7.32E-06	Rikenellaceae	EasyDNA vs QIAStool
18839.0105	-4.359738	0.91400032	-4.7699524	1.84E-06	7.63E-06	Prevotellaceae	EasyDNA vs MagNAPure
3174.43554	-2.7465167	0.55002666	-4.9934247	5.93E-07	9.10E-06	Veillonellaceae	FastDNA vs PowerSoil.HMP
128.300954	-3.603787	0.76190786	-4.7299512	2.25E-06	9.16E-06	[Paraprevotellaceae]	EasyDNA vs InnuPURE
195.592384	3.86327715	0.72509531	5.32795768	9.93E-08	9.54E-06	Coriobacteriaceae	FastDNA vs QIAStool+BB
50.0848862	-4.1453968	0.88843557	-4.665951	3.07E-06	9.83E-06	[Odoribacteraceae]	EasyDNA vs QIAStool
50.0848862	-4.7429127	1.00547833	-4.717071	2.39E-06	1.38E-05	[Odoribacteraceae]	EasyDNA vs PowerSoil.HMP
50.0848862	-4.157899	0.9175784	-4.5313829	5.86E-06	2.56E-05	[Odoribacteraceae]	EasyDNA vs QIAStool+BB
129.697728	2.38013376	0.4664827	5.10229799	3.36E-07	3.22E-05	Erysipelotrichaceae	InnuPURE vs QIAStool
680.114734	-4.3059154	0.9546601	-4.5104173	6.47E-06	3.31E-05	Pasteurellaceae	EasyDNA vs PowerSoil.HMP
833.096381	-3.9165388	0.89595272	-4.3713677	1.23E-05	3.70E-05	Alcaligenaceae	EasyDNA vs QIAStool
833.096381	-4.2614855	0.98211348	-4.3390969	1.43E-05	5.42E-05	Alcaligenaceae	EasyDNA vs InnuPURE
188.732855	-3.231528	0.73880545	-4.3739905	1.22E-05	0.00010307	[Barnesiellaceae]	EasyDNA vs FastDNA
12.0964552	3.88737674	0.89260058	4.35511339	1.33E-05	0.00010307	Actinomycetaceae	EasyDNA vs FastDNA
55.8474481	-2.8658279	0.62501984	-4.585179	4.54E-06	0.00010886	Lactobacillaceae	QIAStool vs QIAStool+BB
680.114734	-3.3086423	0.79130163	-4.1812656	2.90E-05	0.00011596	Pasteurellaceae	EasyDNA vs QIAStool+BB
24.9775852	-3.3034167	0.79477138	-4.1564363	3.23E-05	0.00012499	S24-7	EasyDNA vs MagNAPure
164.683395	2.16861441	0.52508211	4.13004815	3.63E-05	0.00016321	Clostridiaceae	FastDNA vs QIAStool
3174.43554	1.83931522	0.41386092	4.44428346	8.82E-06	0.00016932	Veillonellaceae	QIAStool vs QIAStool+BB
52.7372451	-3.9821991	0.96797847	-4.1139335	3.89E-05	0.00017893	Rikenellaceae	EasyDNA vs PowerSoil.HMP
3174.43554	-1.9851163	0.45364194	-4.3759541	1.21E-05	0.00018135	Veillonellaceae	FastDNA vs InnuPURE
50.0848862	-3.8686456	0.92385007	-4.1875253	2.82E-05	0.00019428	[Odoribacteraceae]	EasyDNA vs FastDNA
12.8572515	4.81248863	1.14323741	4.20952688	2.56E-05	0.00020984	Turicibacteraceae	FastDNA vs MagNAPure
52.7372451	-3.2306201	0.84373606	-3.8289464	0.00012869	0.00047517	Rikenellaceae	EasyDNA vs QIAStool+BB
12.8572515	3.82479629	0.94129838	4.06331974	4.84E-05	0.00054427	Turicibacteraceae	FastDNA vs InnuPURE
2.90835369	5.801207	1.54291246	3.75990677	0.00016998	0.00058278	Flavobacteriaceae	EasyDNA vs QIAStool+BB
24.9775852	-2.6872704	0.73960427	-3.6333895	0.00027972	0.0007898	S24-7	EasyDNA vs QIAStool
20.5947041	-2.5767744	0.70109054	-3.6753803	0.0002375	0.00086092	Desulfovibrionaceae	EasyDNA vs MagNAPure
118.367167	1.7660869	0.47972483	3.68145821	0.0002319	0.00092762	Streptococcaceae	FastDNA vs QIAStool
20.5947041	-2.2850151	0.64393127	-3.5485389	0.00038737	0.001033	Desulfovibrionaceae	EasyDNA vs QIAStool
8087.80815	3.05449188	0.72131966	4.2345884	2.29E-05	0.00109906	Bifidobacteriaceae	QIAStool+BB vs MagNAPure
717.960343	2.05393444	0.51665248	3.97546612	7.02E-05	0.00112386	Verrucomicrobiaceae	QIAStool vs QIAStool+BB
717.960343	2.34984392	0.64833575	3.62442444	0.00028961	0.00121108	Verrucomicrobiaceae	EasyDNA vs PowerSoil.HMP
55.8474481	3.38899548	0.87536361	3.87152888	0.00010815	0.00124378	Lactobacillaceae	FastDNA vs PowerSoil.HMP
14192.9873	2.47838617	0.69245926	3.57910753	0.00034477	0.00132162	Ruminococcaceae	EasyDNA vs PowerSoil.HMP
833.096381	-4.1016476	1.16606691	-3.5175062	0.00043562	0.00154143	Alcaligenaceae	EasyDNA vs PowerSoil.HMP
3052.94009	-2.5171693	0.6913973	-3.6406987	0.0002719	0.00156067	Bacteroidaceae	EasyDNA vs FastDNA
55.8474481	-2.3828644	0.6553515	-3.6360097	0.00027689	0.00156067	Lactobacillaceae	EasyDNA vs FastDNA
2.90835369	5.46873449	1.5682075	3.48725183	0.00048801	0.00169597	Flavobacteriaceae	EasyDNA vs InnuPURE
24.9775852	-2.7362663	0.78754989	-3.4744037	0.00051199	0.00169597	S24-7	EasyDNA vs InnuPURE
833.096381	-3.3916737	0.98214712	-3.4533255	0.00055372	0.00177191	Alcaligenaceae	EasyDNA vs QIAStool+BB
50.0848862	-2.1811191	0.60399625	-3.6111467	0.00030485	0.00196263	[Odoribacteraceae]	FastDNA vs MagNAPure
164.683395	2.091138	0.58305007	3.5865496	0.00033508	0.00196263	Clostridiaceae	FastDNA vs MagNAPure
164.683395	-1.8905223	0.53660316	-3.5231293	0.00042648	0.0022035	Clostridiaceae	EasyDNA vs FastDNA
1.64901163	5.58253937	1.67724193	3.32840438	0.00087345	0.00220661	Corynebacteriaceae	EasyDNA vs QIAStool
680.114734	-2.7600494	0.79209328	-3.4845004	0.00049306	0.0023515	Pasteurellaceae	EasyDNA vs FastDNA
18839.0105	-3.0749852	0.91400099	-3.3643128	0.00076735	0.00239231	Prevotellaceae	EasyDNA vs InnuPURE
8087.80815	-3.0141428	0.71644091	-4.2071059	2.59E-05	0.00248315	Bifidobacteriaceae	InnuPURE vs QIAStool+BB

11.6781466	2.99916116	0.92454824	3.2439207	0.00117897	0.00282952	Staphylococcaceae	EasyDNA vs QIAStool
52.7372451	-2.5787944	0.72537335	-3.555127	0.0003778	0.00340017	Rikenellaceae	FastDNA vs InnuPURE
717.960343	2.18530557	0.56391756	3.87522171	0.00010653	0.00350549	Verrucomicrobiaceae	FastDNA vs QIAStool+BB
12.8572515	3.50252691	0.90541725	3.86841197	0.00010955	0.00350549	Turicibacteraceae	FastDNA vs QIAStool+BB
55.8474481	-2.1167741	0.65447232	-3.2343219	0.00121932	0.00365796	Lactobacillaceae	EasyDNA vs QIAStool+BB
128.300954	-2.9732328	0.91355445	-3.2545765	0.00113562	0.00373131	[Paraprevotellaceae]	EasyDNA vs PowerSoil.HMP
12.0964552	3.6784619	1.13856961	3.23077471	0.00123455	0.00378596	Actinomycetaceae	EasyDNA vs PowerSoil.HMP
33.4361572	2.81733334	0.86547361	3.25525042	0.00113292	0.00386527	Enterobacteriaceae	EasyDNA vs MagNAPure
1.82114912	5.23886708	1.65581045	3.16392923	0.00155655	0.00402037	Sphingobacteriaceae	EasyDNA vs QIAStool+BB
128.300954	-2.4114715	0.76277926	-3.1614277	0.00156998	0.00402037	[Paraprevotellaceae]	EasyDNA vs QIAStool+BB
33.4361572	2.54187917	0.80503432	3.15747925	0.0015914	0.00402037	Enterobacteriaceae	EasyDNA vs QIAStool+BB
3174.43554	-1.7292853	0.45259586	-3.8208156	0.00013301	0.00425635	Veillonellaceae	QIAStool+BB vs MagNAPure
12.0964552	2.92089424	0.9106931	3.20733102	0.00133973	0.0043169	Actinomycetaceae	EasyDNA vs MagNAPure
261.894528	2.6444657	0.67913295	3.89388516	9.87E-05	0.00473527	[Mogibacteriaceae]	InnuPURE vs QIAStool
18839.0105	-3.4408077	1.0966717	-3.1375002	0.00170395	0.00479902	Prevotellaceae	EasyDNA vs PowerSoil.HMP
2.73031934	-5.5948167	1.78991433	-3.1257455	0.00177355	0.00479902	Aerococcaceae	EasyDNA vs PowerSoil.HMP
55.8474481	3.46324375	0.85610471	4.04535067	5.22E-05	0.00501551	Lactobacillaceae	InnuPURE vs MagNAPure
717.960343	-2.0716363	0.56501412	-3.6665213	0.00024587	0.00590094	Verrucomicrobiaceae	QIAStool+BB vs MagNAPure
18839.0105	-2.7588902	0.91399679	-3.01849	0.00254038	0.00609691	Prevotellaceae	EasyDNA vs QIAStool+BB
3.42322903	4.0050664	1.33425048	3.00173503	0.00268446	0.0061359	Pseudomonadaceae	EasyDNA vs QIAStool+BB
750.272969	-2.1281593	0.63659993	-3.343009	0.00082875	0.00621564	Porphyromonadaceae	FastDNA vs InnuPURE
50.0848862	-1.9626447	0.59872204	-3.2780565	0.00104524	0.00671943	[Odoribacteraceae]	FastDNA vs InnuPURE
55.8474481	2.55849915	0.84174411	3.03952131	0.00236954	0.00723335	Lactobacillaceae	EasyDNA vs MagNAPure
3052.94009	-2.3734471	0.74564072	-3.1830975	0.00145709	0.00779712	Bacteroidaceae	FastDNA vs InnuPURE
680.114734	-2.6383223	0.83401698	-3.1633915	0.00155942	0.00779712	Pasteurellaceae	FastDNA vs InnuPURE
261.894528	3.13867473	0.8067979	3.89028617	0.00010013	0.00795635	[Mogibacteriaceae]	PowerSoil.HMP vs QIAStool
129.697728	2.20203213	0.58468421	3.76619048	0.00016576	0.00795635	Erysipelotrichaceae	PowerSoil.HMP vs QIAStool
2.73031934	-5.9403078	1.78353589	-3.3306354	0.00086648	0.00797162	Aerococcaceae	FastDNA vs PowerSoil.HMP
750.272969	-2.0086063	0.63715947	-3.1524389	0.00161913	0.00829803	Porphyromonadaceae	FastDNA vs MagNAPure
12.8572515	2.52115646	0.87157181	2.89265487	0.00382001	0.00873145	Turicibacteraceae	EasyDNA vs QIAStool
12.8249831	2.47583548	0.80177935	3.08792622	0.00201559	0.00907013	Peptostreptococcaceae	FastDNA vs InnuPURE
12.8572515	3.81865729	1.18165368	3.23162137	0.0012309	0.0094369	Turicibacteraceae	FastDNA vs PowerSoil.HMP
24.9775852	-2.6351797	0.91756811	-2.8719173	0.0040799	0.0104264	S24-7	EasyDNA vs PowerSoil.HMP
3052.94009	-2.2637531	0.74575245	-3.0355288	0.00240114	0.01086238	Bacteroidaceae	FastDNA vs MagNAPure
188.732855	-2.1015916	0.6991929	-3.0057393	0.00264936	0.01086238	[Barnesiellaceae]	FastDNA vs MagNAPure
128.300954	-2.2809055	0.7707057	-2.9595026	0.00308136	0.01148507	[Paraprevotellaceae]	FastDNA vs MagNAPure
717.960343	2.05912098	0.56396613	3.65114298	0.00026108	0.01253164	Verrucomicrobiaceae	InnuPURE vs QIAStool+BB
52.7372451	-2.1233709	0.7336022	-2.8944446	0.0037983	0.01297752	Rikenellaceae	FastDNA vs MagNAPure
2.90835369	4.49112126	1.61342038	2.78360267	0.00537588	0.0130153	Flavobacteriaceae	EasyDNA vs PowerSoil.HMP
3174.43554	1.11548884	0.41450174	2.69115597	0.00712049	0.01529563	Veillonellaceae	EasyDNA vs QIAStool+BB
11.6781466	2.68984272	1.00310889	2.6815062	0.00732916	0.01529563	Staphylococcaceae	EasyDNA vs QIAStool+BB
129.697728	-1.7885614	0.47769996	-3.7441104	0.00018103	0.01536327	Erysipelotrichaceae	QIAStool vs MagNAPure
118.367167	-1.7661964	0.4920531	-3.5894428	0.00033139	0.01536327	Streptococcaceae	QIAStool vs MagNAPure
50.0848862	-1.9043678	0.54541108	-3.4916192	0.0004801	0.01536327	[Odoribacteraceae]	QIAStool vs MagNAPure
12.8249831	-2.3955975	0.82460766	-2.9051361	0.00367093	0.01625699	Peptostreptococcaceae	EasyDNA vs FastDNA
833.096381	-2.9212505	1.04951845	-2.7834198	0.00537892	0.01696427	Alcaligenaceae	FastDNA vs MagNAPure
8087.80815	-3.173788	0.86249486	-3.6797762	0.00023344	0.01712544	Bifidobacteriaceae	PowerSoil.HMP vs QIAStool+BB
55.8474481	-3.1229052	0.87472716	-3.5701477	0.00035678	0.01712544	Lactobacillaceae	PowerSoil.HMP vs QIAStool+BB
11.6781466	2.79196033	1.02507165	2.72367334	0.00645603	0.01900943	Staphylococcaceae	EasyDNA vs InnuPURE
24.9775852	-2.0060866	0.78814538	-2.5453256	0.01091759	0.02183518	S24-7	EasyDNA vs QIAStool+BB
3.42322903	-4.3875174	1.50304307	-2.9190896	0.00351055	0.02298506	Pseudomonadaceae	FastDNA vs PowerSoil.HMP
164.683395	1.99810261	0.70063696	2.85183728	0.00434673	0.02298506	Clostridiaceae	FastDNA vs PowerSoil.HMP
717.960343	1.94656943	0.68516789	2.84101089	0.00449708	0.02298506	Verrucomicrobiaceae	FastDNA vs PowerSoil.HMP
1.64901163	4.49276703	1.71286008	2.62296208	0.0087169	0.025279	Corynebacteriaceae	EasyDNA vs MagNAPure
1.82114912	4.39593515	1.68839485	2.60361796	0.00922455	0.02547733	Sphingobacteriaceae	EasyDNA vs MagNAPure
24.9775852	-1.8300403	0.6984518	-2.6201383	0.00878941	0.02574042	S24-7	FastDNA vs MagNAPure
50.0848862	-1.8918656	0.59388004	-3.1856023	0.00144453	0.02773498	[Odoribacteraceae]	QIAStool+BB vs MagNAPure
1.64901163	4.22462547	1.68481228	2.50747548	0.0121597	0.02796731	Corynebacteriaceae	EasyDNA vs PowerSoil.HMP
1.82114912	4.10868763	1.67143497	2.45817976	0.01396433	0.03058852	Sphingobacteriaceae	EasyDNA vs PowerSoil.HMP
195.592384	1.736134	0.65109372	2.66648864	0.00766482	0.03168125	Coriobacteriaceae	EasyDNA vs FastDNA
3.42322903	4.86140167	1.48339223	3.27721931	0.00104835	0.03354717	Pseudomonadaceae	PowerSoil.HMP vs QIAStool+BB
3.42322903	3.53118218	1.35773544	2.6007881	0.00930099	0.03403901	Pseudomonadaceae	EasyDNA vs FastDNA
18839.0105	-2.3758961	0.91401379	-2.5994094	0.00933843	0.03403901	Prevotellaceae	EasyDNA vs FastDNA
52.7372451	-2.2105839	0.85684169	-2.579921	0.00988229	0.03403901	Rikenellaceae	EasyDNA vs FastDNA
750.272969	-1.5057613	0.58394147	-2.5786168	0.00991968	0.03571084	Porphyromonadaceae	FastDNA vs QIAStool

33.4361572	2.28011708	0.9191566	2.48066226	0.01311386	0.03584454	Enterobacteriaceae	FastDNA vs MagNAPure
12.8572515	-2.1618353	0.86641893	-2.4951386	0.0125908	0.04108575	Turicibacteraceae	EasyDNA vs FastDNA
20.5947041	-1.6911916	0.69469488	-2.434438	0.01491493	0.04160481	Desulfovibrionaceae	EasyDNA vs InnuPURE
3052.94009	-1.7009654	0.68492375	-2.4834376	0.01301211	0.0425851	Bacteroidaceae	FastDNA vs QIAStool
3.96627812	2.57800025	1.13190334	2.27757986	0.02275162	0.04368311	Comamonadaceae	EasyDNA vs QIAStool+BB
128.300954	-1.9480249	0.76899668	-2.5332033	0.01130254	0.04623766	[Paraprevotellaceae]	FastDNA vs InnuPURE
33.4361572	1.66163853	0.72289861	2.29857756	0.02152894	0.04635089	Enterobacteriaceae	EasyDNA vs QIAStool
1.29933147	3.68974709	1.61352576	2.28676057	0.0222098	0.04635089	Xanthomonadaceae	EasyDNA vs QIAStool
33.4361572	1.91151815	0.80544298	2.37325075	0.01763229	0.04672557	Enterobacteriaceae	EasyDNA vs InnuPURE
128.300954	-1.7021055	0.70735254	-2.4063043	0.01611483	0.0483445	[Paraprevotellaceae]	FastDNA vs QIAStool
3174.43554	-1.3981164	0.45362294	-3.0821114	0.00205538	0.0493291	Veillonellaceae	FastDNA vs QIAStool+BB
12.8572515	2.65065335	1.14044478	2.32422771	0.02011329	0.05302595	Turicibacteraceae	EasyDNA vs MagNAPure
52.7372451	-1.5580748	0.67048579	-2.3238	0.02013622	0.05576183	Rikenellaceae	FastDNA vs QIAStool
50.0848862	1.68589343	0.53952737	3.1247598	0.0017795	0.0569441	[Odoribacteraceae]	InnuPURE vs QIAStool
20.5947041	-1.4101301	0.61823455	-2.2808982	0.02255447	0.05779583	Desulfovibrionaceae	FastDNA vs MagNAPure
3174.43554	-1.2523153	0.41388181	-3.02578	0.00247993	0.05951824	Veillonellaceae	InnuPURE vs QIAStool
0.882185	4.0371131	1.78622755	2.26013371	0.02381295	0.06009936	Gemellaceae	EasyDNA vs InnuPURE
2.90835369	2.46128406	1.15648797	2.12824009	0.03331718	0.06663437	Flavobacteriaceae	EasyDNA vs QIAStool
14192.9873	1.24151496	0.55827928	2.2238242	0.02616027	0.06726928	Ruminococcaceae	FastDNA vs QIAStool
12.0964552	-3.2773268	1.09799283	-2.9848344	0.00283732	0.06809565	Actinomycetaceae	QIAStool vs MagNAPure
1.64901163	3.77525052	1.6511205	2.2864779	0.02222632	0.06890158	Corynebacteriaceae	EasyDNA vs FastDNA
1.64901163	3.58183657	1.63651122	2.18870272	0.02861845	0.06894446	Corynebacteriaceae	EasyDNA vs InnuPURE
164.683395	1.34909893	0.57279168	2.3553047	0.01850752	0.06940321	Clostridiaceae	FastDNA vs InnuPURE
0.882185	4.01816686	1.78597331	2.24984709	0.02445865	0.07095808	Gemellaceae	EasyDNA vs FastDNA
1.82114912	3.54033456	1.58145933	2.2386504	0.02517867	0.07095808	Sphingobacteriaceae	EasyDNA vs FastDNA
2.73031934	4.97244546	1.63772663	3.03618771	0.0023959	0.07666885	Aerococcaceae	PowerSoil.HMP vs QIAStool
1.29933147	3.36579674	1.65782097	2.03025344	0.04233078	0.07814914	Xanthomonadaceae	EasyDNA vs QIAStool+BB
3052.94009	-2.1228769	0.89019477	-2.384733	0.01709152	0.07862099	Bacteroidaceae	FastDNA vs PowerSoil.HMP
261.894528	1.477758	0.69172546	2.136336	0.03265204	0.07874903	[Mogibacteriaceae]	FastDNA vs MagNAPure
12.0964552	2.31084429	1.08368228	2.13240017	0.03297396	0.07913751	Actinomycetaceae	FastDNA vs QIAStool
680.114734	-1.7526692	0.83469225	-2.099779	0.03574829	0.08142665	Pasteurellaceae	FastDNA vs MagNAPure
128.300954	-1.6557621	0.76773858	-2.1566743	0.03103105	0.08364892	[Paraprevotellaceae]	EasyDNA vs FastDNA
11.6015749	-2.2504463	1.0243547	-2.1969405	0.0280247	0.09112658	Campylobacteraceae	FastDNA vs InnuPURE
55.8474481	1.47811982	0.67420109	2.19240201	0.02835049	0.09112658	Lactobacillaceae	FastDNA vs InnuPURE
0.882185	3.39994624	1.70295575	1.99649711	0.04587984	0.09593057	Gemellaceae	EasyDNA vs PowerSoil.HMP
20.5947041	-1.1183708	0.55185161	-2.0265789	0.04270549	0.09608736	Desulfovibrionaceae	FastDNA vs QIAStool
833.096381	-2.0474946	0.98295866	-2.0829916	0.03725199	0.09623431	Alcaligenaceae	EasyDNA vs FastDNA
1.19877439	3.49866232	1.81740826	1.92508331	0.05421894	0.09638922	Cryomorphaceae	EasyDNA vs QIAStool+BB
1.82114912	3.1528737	1.55024005	2.0337971	0.04197206	0.09671822	Sphingobacteriaceae	EasyDNA vs InnuPURE
18839.0105	-1.9838419	0.98946509	-2.004964	0.04496689	0.09703382	Prevotellaceae	FastDNA vs MagNAPure
2.73031934	-3.6259805	1.69469176	-2.1396106	0.03238625	0.09715874	Aerococcaceae	FastDNA vs InnuPURE
2.90835369	2.67380324	1.29756082	2.06063809	0.03933758	0.0975572	Flavobacteriaceae	EasyDNA vs FastDNA
11.6781466	2.16807721	1.05257629	2.05978154	0.03941943	0.0977461	Staphylococcaceae	EasyDNA vs MagNAPure
0.882185	3.61138826	1.76237664	2.04915804	0.04044666	0.0977461	Gemellaceae	EasyDNA vs MagNAPure
833.096381	-2.2139909	1.04942688	-2.1097143	0.03488297	0.09810835	Alcaligenaceae	FastDNA vs InnuPURE
3174.43554	-0.7238264	0.37173907	-1.9471356	0.05151847	0.09891547	Veillonellaceae	EasyDNA vs QIAStool
18839.0105	-1.8164808	0.91354221	-1.9883929	0.04676826	0.09903866	Prevotellaceae	FastDNA vs QIAStool
750.272969	-1.7309025	0.76559114	-2.2608707	0.02376727	0.09939039	Porphyromonadaceae	FastDNA vs PowerSoil.HMP
188.732855	-1.2492702	0.64121456	-1.9482873	0.0513806	0.09998639	[Barnesiellaceae]	FastDNA vs QIAStool
833.096381	-1.8690442	0.97057026	-1.9257176	0.05413965	0.09998639	Alcaligenaceae	FastDNA vs QIAStool
24.9775852	-1.213894	0.63403122	-1.9145651	0.05554799	0.09998639	S24-7	FastDNA vs QIAStool

Table S2B. Difference abundance of families - Pig fecal microbial community

baseMean	log2FoldChange	lfcSE	stat	pvalue	padj	family	compare
6226.33527	8.23337231	0.2901671	28.3745898	4.16E-177	2.96E-175	Turicibacteraceae	FastDNA vs QIAStool
6226.33527	-8.9600238	0.31794514	-28.181037	9.99E-175	7.99E-173	Turicibacteraceae	EasyDNA vs FastDNA
1329.00016	-7.3100807	0.26522532	-27.561776	3.20E-167	1.28E-165	Peptostreptococcaceae	EasyDNA vs FastDNA
1329.00016	6.29759341	0.23184427	27.1630318	1.78E-162	6.31E-161	Peptostreptococcaceae	FastDNA vs QIAStool
6226.33527	9.16361787	0.33891741	27.0379083	5.30E-161	3.44E-159	Turicibacteraceae	FastDNA vs MagNAPure
1329.00016	6.95674527	0.26345927	26.4053921	1.19E-153	6.18E-152	Peptostreptococcaceae	FastDNA vs InnuPURE
1329.00016	6.24786129	0.26803077	23.310239	3.49E-120	1.13E-118	Peptostreptococcaceae	FastDNA vs MagNAPure
6226.33527	8.37938709	0.35930749	23.3209362	2.72E-120	1.88E-118	Turicibacteraceae	FastDNA vs PowerSoil.HMP
14966.648	-7.3777612	0.32913136	-22.415856	2.76E-111	7.35E-110	Clostridiaceae	EasyDNA vs FastDNA
6226.33527	6.98112067	0.31642099	22.0627609	7.20E-108	1.87E-106	Turicibacteraceae	FastDNA vs InnuPURE
1329.00016	6.57818334	0.30059891	21.8835905	3.72E-106	1.28E-104	Peptostreptococcaceae	FastDNA vs PowerSoil.HMP
1329.00016	6.11331273	0.30715449	19.903055	3.83E-88	2.18E-86	Peptostreptococcaceae	FastDNA vs QIAStool+BB
14966.648	7.01386817	0.36120224	19.4181191	5.42E-84	1.25E-82	Clostridiaceae	FastDNA vs PowerSoil.HMP
14966.648	6.69216818	0.34940405	19.153093	9.12E-82	1.58E-80	Clostridiaceae	FastDNA vs InnuPURE
14966.648	6.16102466	0.34985453	17.6102467	2.06E-69	4.45E-68	Clostridiaceae	FastDNA vs MagNAPure
14966.648	5.44539513	0.318726	17.0848791	1.92E-65	4.55E-64	Clostridiaceae	FastDNA vs QIAStool
6226.33527	6.27791748	0.38345515	16.3719732	3.03E-60	8.64E-59	Turicibacteraceae	FastDNA vs QIAStool+BB
14966.648	5.80051888	0.42495209	13.6498186	2.02E-42	3.85E-41	Clostridiaceae	FastDNA vs QIAStool+BB
254.614729	-2.6451948	0.27545666	-9.6029437	7.77E-22	5.05E-20	S24-7	InnuPURE vs QIAStool
35.1041071	2.34749529	0.28446002	8.25246117	1.55E-16	1.18E-14	Peptococcaceae	EasyDNA vs QIAStool
576.775751	2.79069511	0.38546037	7.23990142	4.49E-13	8.98E-12	Streptococcaceae	EasyDNA vs FastDNA
408.703374	3.64804379	0.50165879	7.27196224	3.54E-13	3.51E-11	Lactobacillaceae	EasyDNA vs MagNAPure
254.614729	-1.8657731	0.26216238	-7.1168605	1.10E-12	4.20E-11	S24-7	EasyDNA vs QIAStool
6226.33527	2.88570039	0.40420787	7.13914937	9.39E-13	1.09E-10	Turicibacteraceae	QIAStool+BB vs MagNAPure
576.775751	2.21032995	0.32171271	6.87050862	6.40E-12	1.62E-10	Streptococcaceae	EasyDNA vs QIAStool
2576.15115	-2.3625033	0.3468527	-6.8112582	9.67E-12	1.72E-10	[Paraprevotellaceae]	FastDNA vs QIAStool
6226.33527	-2.6821064	0.38685912	-6.9330312	4.12E-12	2.06E-10	Turicibacteraceae	EasyDNA vs QIAStool+BB
576.775751	2.51571595	0.36417115	6.90805937	4.91E-12	2.43E-10	Streptococcaceae	EasyDNA vs MagNAPure
35.1041071	2.71384929	0.40707004	6.66678715	2.61E-11	8.63E-10	Peptococcaceae	EasyDNA vs MagNAPure
14966.648	-1.9323661	0.29731783	-6.4993278	8.07E-11	1.53E-09	Clostridiaceae	EasyDNA vs QIAStool
576.775751	-2.5501711	0.41120073	-6.201767	5.58E-10	7.26E-09	Streptococcaceae	FastDNA vs InnuPURE
6226.33527	2.1824972	0.34136207	6.39349657	1.62E-10	8.92E-09	Turicibacteraceae	InnuPURE vs MagNAPure
22.9687795	-4.1589171	0.64930852	-6.4051478	1.50E-10	9.46E-09	Helicobacteraceae	EasyDNA vs InnuPURE
254.614729	-1.8689828	0.30680193	-6.0918223	1.12E-09	1.59E-08	S24-7	FastDNA vs QIAStool
254.614729	-1.8826504	0.30472804	-6.1781333	6.49E-10	1.78E-08	S24-7	InnuPURE vs MagNAPure
1019.42191	-2.9556395	0.49110705	-6.0183202	1.76E-09	1.83E-08	Campylobacteraceae	FastDNA vs InnuPURE
6226.33527	-1.9789032	0.32055255	-6.1734127	6.68E-10	2.11E-08	Turicibacteraceae	EasyDNA vs InnuPURE
408.703374	3.99766567	0.64482803	6.19958424	5.66E-10	3.28E-08	Lactobacillaceae	QIAStool+BB vs MagNAPure
22.9687795	3.05927367	0.51002005	5.99834003	1.99E-09	6.48E-08	Helicobacteraceae	InnuPURE vs QIAStool
576.775751	2.27519194	0.39134713	5.81374372	6.11E-09	1.12E-07	Streptococcaceae	InnuPURE vs MagNAPure
62.3051996	-3.6585641	0.613405	-5.9643533	2.46E-09	1.87E-07	Enterobacteriaceae	EasyDNA vs PowerSoil.HMP
408.703374	2.43422116	0.43645264	5.57728586	2.44E-08	3.71E-07	Lactobacillaceae	EasyDNA vs QIAStool
576.775751	1.96980595	0.35224663	5.59212148	2.24E-08	4.86E-07	Streptococcaceae	InnuPURE vs QIAStool
2576.15115	-1.9272471	0.34922351	-5.5186637	3.42E-08	2.22E-06	[Paraprevotellaceae]	PowerSoil.HMP vs QIAStool
216.46708	6.27411432	1.18075226	5.31365855	1.07E-07	3.22E-06	Bacteroidaceae	PowerSoil.HMP vs QIAStool
62.3051996	2.92721007	0.55710414	5.25433196	1.49E-07	3.22E-06	Enterobacteriaceae	PowerSoil.HMP vs QIAStool
367.01539	-2.774783	0.54483681	-5.09287	3.53E-07	3.63E-06	Spirochaetaceae	FastDNA vs QIAStool
7308.6798	-1.2094339	0.23761089	-5.0899766	3.58E-07	3.63E-06	Prevotellaceae	FastDNA vs QIAStool
71.0660634	-2.1428817	0.41434301	-5.1717579	2.32E-07	3.77E-06	Alcaligenaceae	InnuPURE vs QIAStool
7308.6798	1.22465113	0.23872154	5.13004044	2.90E-07	4.63E-06	Prevotellaceae	EasyDNA vs FastDNA
62.3051996	-3.2981065	0.61505214	-5.3623202	8.22E-08	4.85E-06	Enterobacteriaceae	InnuPURE vs PowerSoil.HMP
2516.24433	1.09546943	0.21932759	4.99467231	5.89E-07	5.11E-06	Ruminococcaceae	FastDNA vs InnuPURE
6226.33527	-1.9554548	0.36446037	-5.3653428	8.08E-08	5.98E-06	Turicibacteraceae	QIAStool vs QIAStool+BB
35.1041071	1.49262726	0.29814049	5.00645612	5.54E-07	7.21E-06	Peptococcaceae	InnuPURE vs QIAStool
408.703374	2.70224502	0.53923011	5.01130217	5.41E-07	7.43E-06	Lactobacillaceae	InnuPURE vs MagNAPure
1019.42191	-2.2116186	0.45182578	-4.8948483	9.84E-07	8.73E-06	Campylobacteraceae	FastDNA vs QIAStool
88.4317884	3.06001619	0.6165416	4.96319502	6.93E-07	9.25E-06	Coriobacteriaceae	EasyDNA vs FastDNA
254.614729	-1.514942	0.30301081	-4.9996301	5.74E-07	9.33E-06	S24-7	PowerSoil.HMP vs QIAStool
62.3051996	3.74759308	0.72291645	5.18399201	2.17E-07	9.99E-06	Enterobacteriaceae	PowerSoil.HMP vs QIAStool+BB
7308.6798	-1.1800909	0.23943127	-4.928725	8.28E-07	1.08E-05	Prevotellaceae	PowerSoil.HMP vs QIAStool
35.1041071	1.93628224	0.38058991	5.0875816	3.63E-07	1.23E-05	Peptococcaceae	EasyDNA vs PowerSoil.HMP
22.9687795	-3.5480864	0.70506262	-5.0322997	4.85E-07	1.23E-05	Helicobacteraceae	EasyDNA vs PowerSoil.HMP
2516.24433	-1.1212344	0.22183279	-5.0544122	4.32E-07	1.27E-05	Ruminococcaceae	InnuPURE vs PowerSoil.HMP
7308.6798	1.19530817	0.24053344	4.96940538	6.72E-07	1.28E-05	Prevotellaceae	EasyDNA vs PowerSoil.HMP
6226.33527	-2.1014696	0.42138725	-4.9870271	6.13E-07	1.41E-05	Turicibacteraceae	PowerSoil.HMP vs QIAStool+BB
14966.648	-1.568473	0.33252325	-4.7168823	2.39E-06	2.59E-05	Clostridiaceae	PowerSoil.HMP vs QIAStool
254.614729	1.65814449	0.33449335	4.95718218	7.15E-07	2.65E-05	S24-7	QIAStool vs QIAStool+BB
2576.15115	-1.5902241	0.34177933	-4.6527802	3.27E-06	3.55E-05	[Paraprevotellaceae]	InnuPURE vs QIAStool

35.1041071	3.04129082	0.65712892	4.62814943	3.69E-06	4.22E-05	Peptococcaceae	EasyDNA vs FastDNA
367.01539	-2.2519712	0.48760528	-4.6184307	3.87E-06	4.47E-05	Spirochaetaceae	EasyDNA vs QIAStool
42.1528852	3.57440506	0.77612071	4.60547568	4.12E-06	4.47E-05	Bifidobacteriaceae	EasyDNA vs QIAStool
30.2465249	5.96837557	1.30932681	4.55835437	5.16E-06	4.79E-05	Verrucomicrobiaceae	PowerSoil.HMP vs QIAStool
71.0660634	-2.5632064	0.56711788	-4.5197066	6.19E-06	4.89E-05	Alcaligenaceae	FastDNA vs QIAStool
1019.42191	-2.3003618	0.49644019	-4.6337139	3.59E-06	6.20E-05	Campylobacteraceae	FastDNA vs PowerSoil.HMP
216.46708	6.67995824	1.44979616	4.60751549	4.08E-06	6.25E-05	Bacteroidaceae	PowerSoil.HMP vs QIAStool+BB
2516.24433	-0.98295	0.21745846	-4.5201736	6.18E-06	6.80E-05	Ruminococcaceae	InnuPURE vs MagNAPure
408.703374	-2.783843	0.59619079	-4.6693829	3.02E-06	7.45E-05	Lactobacillaceae	QIAStool vs QIAStool+BB
35.1041071	1.85898126	0.41668134	4.46139786	8.14E-06	7.46E-05	Peptococcaceae	InnuPURE vs MagNAPure
908.726526	-1.6630925	0.37476149	-4.4377359	9.09E-06	8.64E-05	Veillonellaceae	EasyDNA vs QIAStool
408.703374	2.48203872	0.54739067	4.53430954	5.78E-06	9.39E-05	Lactobacillaceae	FastDNA vs MagNAPure
42.1528852	4.04263564	0.88466376	4.56968603	4.88E-06	0.00010258	Bifidobacteriaceae	EasyDNA vs InnuPURE
810.892629	-2.4775883	0.57438549	-4.3134592	1.61E-05	0.00011411	Erysipelotrichaceae	FastDNA vs QIAStool
576.775751	1.69015115	0.37888619	4.4608413	8.16E-06	0.00012409	Streptococcaceae	EasyDNA vs PowerSoil.HMP
367.01539	-2.2701322	0.52308917	-4.339857	1.43E-05	0.00013239	Spirochaetaceae	InnuPURE vs QIAStool
29.6083625	1.69887968	0.38157423	4.45229143	8.50E-06	0.00013381	[Mogibacteriaceae]	EasyDNA vs InnuPURE
88.4317884	-2.803111	0.63669121	-4.4026225	1.07E-05	0.00013904	Coriobacteriaceae	FastDNA vs MagNAPure
6226.33527	1.25225163	0.2930255	4.27352444	1.92E-05	0.00015633	Turicibacteraceae	InnuPURE vs QIAStool
23.2927606	8.01906524	1.88146584	4.26213705	2.02E-05	0.00016452	Aerococcaceae	PowerSoil.HMP vs QIAStool
22.9687795	2.448443	0.58021396	4.21989674	2.44E-05	0.00017652	Helicobacteraceae	PowerSoil.HMP vs QIAStool
32.5255849	5.81457361	1.39301324	4.17409788	2.99E-05	0.00019446	Staphylococcaceae	PowerSoil.HMP vs QIAStool
367.01539	-2.2707797	0.54976209	-4.1304771	3.62E-05	0.00021392	Spirochaetaceae	PowerSoil.HMP vs QIAStool
71.0660634	-1.6777577	0.39962713	-4.1983078	2.69E-05	0.00022709	Alcaligenaceae	EasyDNA vs QIAStool
75.3501469	-3.3526702	0.78701599	-4.2599773	2.04E-05	0.0002576	Mycoplasmataceae	EasyDNA vs InnuPURE
22.9687795	-2.9648076	0.67504024	-4.3920458	1.12E-05	0.00027792	Helicobacteraceae	EasyDNA vs MagNAPure
17.2832685	-2.2084726	0.53750076	-4.1087804	3.98E-05	0.00028727	Desulfovibrionaceae	InnuPURE vs QIAStool
30.2465249	6.56938288	1.56924588	4.18633114	2.83E-05	0.00032602	Verrucomicrobiaceae	PowerSoil.HMP vs QIAStool+BB
15.8153467	4.63369833	1.14552937	4.04502795	5.23E-05	0.00033768	Moraxellaceae	FastDNA vs QIAStool
7308.6798	1.26879368	0.29728884	4.26788197	1.97E-05	0.00036507	Prevotellaceae	QIAStool vs QIAStool+BB
32.5255849	5.54202541	1.38556172	3.99984016	6.34E-05	0.00037503	Staphylococcaceae	FastDNA vs QIAStool
1329.00016	-1.0124872	0.24978175	-4.0534877	5.05E-05	0.00038349	Peptostreptococcaceae	EasyDNA vs QIAStool
1019.42191	1.93983364	0.47935342	4.04677127	5.19E-05	0.00040801	Campylobacteraceae	InnuPURE vs MagNAPure
7308.6798	1.28401094	0.29817511	4.3062311	1.66E-05	0.00041515	Prevotellaceae	EasyDNA vs QIAStool+BB
576.775751	-2.0306009	0.48938134	-4.1493222	3.33E-05	0.00047503	Streptococcaceae	FastDNA vs QIAStool+BB
15.4343002	1.53247414	0.38535741	3.97676056	6.99E-05	0.00048267	Corynebacteriaceae	EasyDNA vs QIAStool
31.9177507	-3.2697252	0.83566211	-3.912736	9.13E-05	0.0004984	Succinivibrionaceae	FastDNA vs QIAStool
5.3314072	4.35405821	1.11876983	3.89182661	9.95E-05	0.00053892	[Methanomassiliicoccaceae]	PowerSoil.HMP vs QIAStool
14966.648	-1.2467731	0.31964806	-3.9004556	9.60E-05	0.0005811	Clostridiaceae	InnuPURE vs QIAStool
2516.24433	-0.771523	0.19809811	-3.8946512	9.83E-05	0.0005811	Ruminococcaceae	InnuPURE vs QIAStool
216.46708	5.5853327	1.27809173	4.37005621	1.24E-05	0.00064592	Bacteroidaceae	PowerSoil.HMP vs MagNAPure
62.3051996	2.61404432	0.62207451	4.20214024	2.64E-05	0.00068745	Enterobacteriaceae	PowerSoil.HMP vs MagNAPure
2576.15115	-1.514051	0.37895822	-3.995298	6.46E-05	0.00069997	[Paraprevotellaceae]	FastDNA vs MagNAPure
2516.24433	0.80433271	0.20244058	3.97317915	7.09E-05	0.00074466	Ruminococcaceae	EasyDNA vs InnuPURE
22.9687795	-2.8972138	0.74038877	-3.913098	9.11E-05	0.0009112	Helicobacteraceae	EasyDNA vs FastDNA
42.1528852	3.64689481	0.9520937	3.83039483	0.00012794	0.0009504	Bifidobacteriaceae	FastDNA vs InnuPURE
42.1528852	3.17866423	0.85340488	3.72468483	0.00019556	0.00099177	Bifidobacteriaceae	FastDNA vs QIAStool
88.4317884	1.56076323	0.41573031	3.75426855	0.00017385	0.00110104	Coriobacteriaceae	EasyDNA vs QIAStool
88.4317884	1.78168807	0.46431042	3.83727784	0.00012441	0.00111965	Coriobacteriaceae	EasyDNA vs InnuPURE
35.1041071	-1.3708732	0.34666006	-3.9545174	7.67E-05	0.001135	Peptococcaceae	QIAStool vs QIAStool+BB
1019.42191	-1.6885426	0.44454014	-3.798403	0.00014563	0.00114685	Campylobacteraceae	EasyDNA vs InnuPURE
2576.15115	1.67665813	0.42907776	3.90758571	9.32E-05	0.00114975	[Paraprevotellaceae]	QIAStool vs QIAStool+BB
22.9687795	-2.9199313	0.73813187	-3.9558396	7.63E-05	0.00127111	Helicobacteraceae	EasyDNA vs QIAStool+BB
7.44045113	2.81616091	0.73648989	3.82376043	0.00013143	0.00138942	Anaeroplasmataceae	QIAStool vs QIAStool+BB
2576.15115	1.31919125	0.34980343	3.77123584	0.00016244	0.00144392	[Paraprevotellaceae]	EasyDNA vs FastDNA
14966.648	-1.5772423	0.40923926	-3.8540836	0.00011616	0.00145205	Clostridiaceae	EasyDNA vs QIAStool+BB
6226.33527	1.39826642	0.36161179	3.86676115	0.00011029	0.0016201	Turicibacteraceae	InnuPURE vs PowerSoil.HMP
20.723165	-5.185472	1.34638073	-3.8514158	0.00011744	0.0016201	Pseudomonadaceae	InnuPURE vs PowerSoil.HMP
71.0660634	-1.8593941	0.49253655	-3.7751394	0.00015992	0.0016201	Alcaligenaceae	InnuPURE vs PowerSoil.HMP
216.46708	-4.7736527	1.26699265	-3.7677036	0.00016476	0.0016201	Bacteroidaceae	InnuPURE vs PowerSoil.HMP
216.46708	-4.9350175	1.29105875	-3.8224577	0.00013213	0.00182337	Bacteroidaceae	FastDNA vs PowerSoil.HMP
810.892629	-2.373461	0.62905633	-3.7730501	0.00016126	0.00185453	Erysipelotrichaceae	FastDNA vs PowerSoil.HMP
1329.00016	-1.1967679	0.32086779	-3.7297852	0.00019164	0.00191643	Peptostreptococcaceae	EasyDNA vs QIAStool+BB
9.6383629	-2.1679125	0.6068059	-3.5726621	0.00035337	0.00206586	Fibrobacteraceae	EasyDNA vs QIAStool
15.4343002	1.84120962	0.50412395	3.65227395	0.00025993	0.00216607	Corynebacteriaceae	EasyDNA vs QIAStool+BB
71.0660634	-1.6471299	0.4610371	-3.5726625	0.00035337	0.00242942	Alcaligenaceae	InnuPURE vs MagNAPure
216.46708	-4.4629635	1.20206834	-3.7127369	0.00020503	0.00259705	Bacteroidaceae	EasyDNA vs PowerSoil.HMP
254.614729	-1.1032288	0.29278195	-3.7680902	0.0001645	0.002651	S24-7	EasyDNA vs MagNAPure
1329.00016	-1.0622194	0.28366823	-3.7445834	0.00018069	0.002651	Peptostreptococcaceae	EasyDNA vs MagNAPure
15.4343002	1.90293704	0.50943828	3.73536328	0.00018744	0.002651	Corynebacteriaceae	EasyDNA vs MagNAPure

71.0660634	-2.2797188	0.62584767	-3.6426097	0.00026989	0.00266032	Alcaligenaceae	FastDNA vs PowerSoil.HMP
14966.648	-1.2167365	0.33050128	-3.6814881	0.00023188	0.00286947	Clostridiaceae	EasyDNA vs MagNAPure
576.775751	1.44962715	0.4050545	3.57884468	0.00034512	0.00290884	Streptococcaceae	InnuPURE vs PowerSoil.HMP
86.5992792	2.99949869	0.85659517	3.5016526	0.00046238	0.00300548	Porphyromonadaceae	FastDNA vs InnuPURE
4.16403335	5.96645628	1.75160616	3.4062773	0.00065855	0.00311715	Carnobacteriaceae	FastDNA vs QIAStool
15.8153467	5.10720894	1.40712282	3.62954026	0.00028393	0.00323676	Moraxellaceae	FastDNA vs QIAStool+BB
15.4343002	1.94734349	0.53872155	3.61474956	0.00030064	0.00326407	Corynebacteriaceae	EasyDNA vs PowerSoil.HMP
15.4343002	1.5357668	0.43958903	3.4936422	0.00047648	0.00333535	Corynebacteriaceae	EasyDNA vs InnuPURE
15.8153467	4.25898236	1.24098327	3.43194181	0.00059928	0.00346248	Moraxellaceae	FastDNA vs InnuPURE
62.3051996	-2.2078122	0.62740538	-3.5189565	0.00043325	0.00346598	Enterobacteriaceae	EasyDNA vs FastDNA
75.3501469	-2.899597	0.86109053	-3.3673544	0.00075893	0.00394644	Mycoplasmataceae	FastDNA vs InnuPURE
2576.15115	-1.0433121	0.30923781	-3.3738179	0.00074133	0.00402438	[Paraprevotellaceae]	EasyDNA vs QIAStool
23.2927606	6.98533528	1.92158786	3.63518912	0.00027778	0.00426411	Aerococcaceae	PowerSoil.HMP vs MagNAPure
30.2465249	5.15082886	1.43392754	3.59211237	0.00032801	0.00426411	Verrucomicrobiaceae	PowerSoil.HMP vs MagNAPure
30.2465249	-4.7520422	1.38719877	-3.425639	0.00061336	0.00452349	Verrucomicrobiaceae	InnuPURE vs PowerSoil.HMP
408.703374	1.97172535	0.56084378	3.51564092	0.00043869	0.00456242	Lactobacillaceae	PowerSoil.HMP vs MagNAPure
35.1041071	-2.1864228	0.662999	-3.2977769	0.00097454	0.00460689	Peptococcaceae	FastDNA vs InnuPURE
29.6083625	-1.3718487	0.40734356	-3.3677929	0.00075772	0.00463054	[Mogibacteriaceae]	InnuPURE vs MagNAPure
254.614729	-1.1302528	0.33388242	-3.3851821	0.00071131	0.00466304	S24-7	InnuPURE vs PowerSoil.HMP
35.1041071	1.73722724	0.45243877	3.83969578	0.00012319	0.00476323	Peptococcaceae	QIAStool+BB vs MagNAPure
4.00044772	5.78336799	1.78215296	3.24515803	0.00117385	0.0050867	Cytophagaceae	FastDNA vs InnuPURE
1574.16355	1.17263609	0.35784849	3.27690659	0.00104951	0.00524755	Lachnospiraceae	PowerSoil.HMP vs QIAStool
71.0660634	-2.0674546	0.60167145	-3.4361853	0.00058997	0.00547827	Alcaligenaceae	FastDNA vs MagNAPure
29.6083625	1.43379183	0.42701614	3.35769937	0.00078594	0.00561386	[Mogibacteriaceae]	EasyDNA vs QIAStool+BB
23.2927606	-6.4898889	1.90765791	-3.4020192	0.0006689	0.00576926	Aerococcaceae	FastDNA vs PowerSoil.HMP
29.6083625	1.05593006	0.32473651	3.25165181	0.00114736	0.00581332	[Mogibacteriaceae]	EasyDNA vs QIAStool
908.726526	-1.4420041	0.41614103	-3.4651814	0.00052987	0.00582861	Veillonellaceae	EasyDNA vs MagNAPure
5.3314072	-4.298708	1.25525218	-3.4245772	0.00061576	0.00584969	[Methanomassiliicoccaceae]	EasyDNA vs PowerSoil.HMP
576.775751	1.75562174	0.47291523	3.71233916	0.00020535	0.00595522	Streptococcaceae	QIAStool+BB vs MagNAPure
32.5255849	4.24277218	1.31620267	3.22349458	0.00126637	0.00601524	Staphylococcaceae	EasyDNA vs QIAStool
88.4317884	1.85557658	0.56238288	3.29948979	0.00096861	0.0060538	Coriobacteriaceae	EasyDNA vs QIAStool+BB
254.614729	-1.1064385	0.33329617	-3.3196856	0.00090119	0.00703253	S24-7	FastDNA vs MagNAPure
7308.6798	-0.8575932	0.26003373	-3.2980077	0.00097373	0.00703253	Prevotellaceae	FastDNA vs MagNAPure
5.3314072	3.37108488	1.05056705	3.2088241	0.00133279	0.00721928	[Methanomassiliicoccaceae]	InnuPURE vs QIAStool
42.1528852	-3.1585672	0.98137257	-3.21852	0.00128854	0.00760239	Bifidobacteriaceae	InnuPURE vs PowerSoil.HMP
408.703374	1.67631844	0.50776167	3.3013883	0.00096208	0.0081242	Lactobacillaceae	EasyDNA vs PowerSoil.HMP
576.775751	-1.4502357	0.44128745	-3.2863743	0.00101486	0.00856339	Streptococcaceae	QIAStool vs QIAStool+BB
908.726526	1.66195963	0.50683851	3.27907132	0.00104149	0.00856339	Veillonellaceae	QIAStool vs QIAStool+BB
20.723165	4.83629545	1.47007325	3.28983298	0.00100247	0.00922271	Pseudomonadaceae	PowerSoil.HMP vs QIAStool+BB
408.703374	1.48842239	0.47936509	3.10498703	0.00190287	0.00951437	Lactobacillaceae	InnuPURE vs QIAStool
367.01539	2.08867133	0.64924642	3.2170702	0.00129507	0.00958351	Spirochaetaceae	QIAStool vs QIAStool+BB
4.00044772	4.90531979	1.59999884	3.06582709	0.00217069	0.00963243	Cytophagaceae	FastDNA vs QIAStool
23.2927606	-6.0628195	1.88263038	-3.2203982	0.00128013	0.00972896	Aerococcaceae	EasyDNA vs PowerSoil.HMP
15.8153467	4.05506684	1.27803736	3.172886	0.00150932	0.00981057	Moraxellaceae	FastDNA vs MagNAPure
88.4317884	-1.5247829	0.49109684	-3.1048518	0.00190374	0.01047059	Coriobacteriaceae	InnuPURE vs MagNAPure
5.3314072	-3.7627421	1.19345081	-3.1528255	0.00161698	0.0108779	[Methanomassiliicoccaceae]	QIAStool vs QIAStool+BB
42.1528852	2.69033662	0.88649264	3.03480989	0.00240687	0.01117477	Bifidobacteriaceae	PowerSoil.HMP vs QIAStool
23.2927606	6.07256394	1.92001451	3.16276981	0.00156276	0.01192028	Aerococcaceae	PowerSoil.HMP vs QIAStool+BB
408.703374	-2.0259403	0.64952423	-3.1191143	0.00181396	0.01192028	Lactobacillaceae	PowerSoil.HMP vs QIAStool+BB
810.892629	-1.9297672	0.62539388	-3.0856429	0.00203085	0.01200094	Erysipelotrichaceae	FastDNA vs MagNAPure
62.3051996	1.8473546	0.62901353	2.93690754	0.00331503	0.01326012	Enterobacteriaceae	FastDNA vs InnuPURE
29.6083625	1.83023832	0.58732018	3.11625309	0.00183165	0.01332109	[Mogibacteriaceae]	EasyDNA vs FastDNA
7308.6798	-0.8282503	0.26169677	-3.1649236	0.00155124	0.01344405	Prevotellaceae	PowerSoil.HMP vs MagNAPure
810.892629	-1.5061467	0.51416507	-2.9293057	0.0033972	0.01490319	Erysipelotrichaceae	EasyDNA vs QIAStool
75.3501469	-2.1087165	0.72280839	-2.9173935	0.0035297	0.01490319	Mycoplasmataceae	EasyDNA vs QIAStool
20.723165	4.16353174	1.34986951	3.08439573	0.00203966	0.01515177	Pseudomonadaceae	PowerSoil.HMP vs MagNAPure
86.5992792	-2.5986406	0.87070194	-2.9845352	0.0028401	0.01523324	Porphyromonadaceae	InnuPURE vs PowerSoil.HMP
30.7423537	-1.6327572	0.56577317	-2.8858865	0.00390313	0.01561251	Elusimicrobiaceae	EasyDNA vs QIAStool
29.6083625	1.32991944	0.43726084	3.04147851	0.00235419	0.01626534	[Mogibacteriaceae]	EasyDNA vs PowerSoil.HMP
22.9687795	1.79757037	0.62321785	2.8843371	0.00392239	0.01638173	Helicobacteraceae	FastDNA vs QIAStool
62.3051996	2.29684119	0.73468911	3.12627637	0.00177035	0.01681833	Enterobacteriaceae	FastDNA vs QIAStool+BB
908.726526	-1.1737315	0.4109824	-2.8559168	0.00429128	0.0169267	Veillonellaceae	FastDNA vs QIAStool
22.9687795	-1.8651642	0.54282237	-3.4360489	0.00059026	0.01771725	Helicobacteraceae	QIAStool vs MagNAPure
5.3314072	-3.6505736	1.06984005	-3.4122611	0.00064426	0.01771725	[Methanomassiliicoccaceae]	QIAStool vs MagNAPure
17.2832685	-2.4187504	0.86002934	-2.8124045	0.00491726	0.01837503	Desulfovibrionaceae	FastDNA vs QIAStool
1.66132016	4.416643	1.57030905	2.81259475	0.00491435	0.01867455	Leuconostocaceae	EasyDNA vs QIAStool
4.00044772	5.29050645	1.80899893	2.92454925	0.00344955	0.01868509	Cytophagaceae	FastDNA vs MagNAPure
5.8077209	-2.1252395	0.74134296	-2.8667427	0.0041472	0.01925486	Dethiosulfovibrionaceae	InnuPURE vs QIAStool
31.9177507	-1.3581394	0.47904029	-2.8351256	0.00458077	0.01984999	Succinivibrionaceae	InnuPURE vs QIAStool
9.6383629	-1.5817218	0.5627087	-2.810907	0.00494021	0.02006959	Fibrobacteraceae	InnuPURE vs QIAStool

1574.16355	-1.0713078	0.36311938	-2.9502908	0.00317475	0.02010675	Lachnospiraceae	EasyDNA vs PowerSoil.HMP
22.9687795	-1.8202879	0.62049856	-2.9335893	0.00335067	0.02066248	Helicobacteraceae	QIAStool vs QIAStool+BB
35.1041071	-2.0646688	0.68555435	-3.0116778	0.00259808	0.02115581	Peptococcaceae	FastDNA vs QIAStool+BB
1574.16355	-1.1146482	0.39092455	-2.8513128	0.00435391	0.02140673	Lachnospiraceae	InnuPURE vs PowerSoil.HMP
71.0660634	-1.3942701	0.48029311	-2.9029567	0.00369658	0.02161076	Alcaligenaceae	EasyDNA vs PowerSoil.HMP
31.9177507	2.50733503	0.858128	2.92186602	0.00347941	0.02319608	Succinivibrionaceae	EasyDNA vs FastDNA
31.9177507	-2.4656281	0.87371635	-2.8220006	0.00477251	0.02386254	Succinivibrionaceae	FastDNA vs MagNAPure
20.723165	-4.4393022	1.50402412	-2.9516164	0.00316115	0.02423551	Pseudomonadaceae	FastDNA vs PowerSoil.HMP
35.1041071	0.97662205	0.3461751	2.82117935	0.00478474	0.02483044	Peptococcaceae	EasyDNA vs QIAStool+BB
5.3314072	-3.7073919	1.31972034	-2.8092254	0.00496609	0.02483044	[Methanomassiliicoccaceae]	EasyDNA vs QIAStool+BB
88.4317884	1.46816717	0.5184466	2.83185802	0.00462784	0.02512256	Coriobacteriaceae	EasyDNA vs PowerSoil.HMP
35.1041071	0.85486803	0.29757508	2.87278099	0.00406876	0.02563319	Peptococcaceae	EasyDNA vs InnuPURE
35.1041071	1.08141422	0.39086961	2.76668788	0.00566289	0.02570083	Peptococcaceae	InnuPURE vs PowerSoil.HMP
23.2927606	-5.16892	1.89209931	-2.7318439	0.0062981	0.02654198	Aerococcaceae	InnuPURE vs PowerSoil.HMP
30.7423537	-1.9203408	0.72425206	-2.6514814	0.00801395	0.02844954	Elusimicrobiaceae	FastDNA vs QIAStool
1329.00016	-0.6591519	0.24790456	-2.6588937	0.00783977	0.0294383	Peptostreptococcaceae	InnuPURE vs QIAStool
7308.6798	-0.6240217	0.23586236	-2.6457027	0.00815214	0.0294383	Prevotellaceae	InnuPURE vs QIAStool
7.94956632	2.35515765	0.88703846	2.65507951	0.00792897	0.02945047	Christensenellaceae	FastDNA vs InnuPURE
5.3314072	-3.5952234	1.21352904	-2.9626183	0.00305035	0.03019842	[Methanomassiliicoccaceae]	EasyDNA vs MagNAPure
2576.15115	-1.0787948	0.38112544	-2.8305504	0.0046468	0.03020419	[Paraprevotellaceae]	PowerSoil.HMP vs MagNAPure
14966.648	-1.2133493	0.43533275	-2.7871767	0.00531695	0.03057245	Clostridiaceae	PowerSoil.HMP vs QIAStool+BB
4.2789796	-4.8902304	1.84214054	-2.6546457	0.00793918	0.03122742	Xanthomonadaceae	InnuPURE vs PowerSoil.HMP
75.3501469	2.20086404	0.8067117	2.72819154	0.00636826	0.03184131	Mycoplasmataceae	InnuPURE vs MagNAPure
408.703374	1.26821609	0.4885714	2.59576407	0.00943808	0.03190972	Lactobacillaceae	FastDNA vs QIAStool
62.3051996	1.47645817	0.57256579	2.57866992	0.00991815	0.03200858	Enterobacteriaceae	FastDNA vs QIAStool
5.3314072	-3.3157346	1.1972195	-2.7695294	0.00561373	0.03215138	[Methanomassiliicoccaceae]	EasyDNA vs InnuPURE
1019.42191	-1.6633229	0.59035526	-2.8174948	0.00483999	0.03448493	Campylobacteraceae	FastDNA vs QIAStool+BB
8.45632111	5.03537667	1.90368579	2.64506711	0.00816747	0.03539238	Rikenellaceae	PowerSoil.HMP vs QIAStool
2.95003546	4.7042716	1.82149714	2.58264012	0.00980475	0.03548387	Actinomycetaceae	EasyDNA vs QIAStool
1019.42191	1.26709688	0.45944542	2.75788338	0.00581769	0.0358012	Campylobacteraceae	EasyDNA vs FastDNA
1.25729577	3.9531204	1.56655913	2.52344155	0.01162124	0.03587426	Methanobacteriaceae	FastDNA vs QIAStool
9.6383629	-1.9706676	0.68484384	-2.8775431	0.00400785	0.03607067	Fibrobacteraceae	EasyDNA vs MagNAPure
7308.6798	0.63923894	0.23698127	2.69742385	0.00698783	0.03668609	Prevotellaceae	EasyDNA vs InnuPURE
7.44045113	2.10035711	0.79576123	2.6394313	0.00830453	0.03774784	Anaeroplasmataceae	EasyDNA vs QIAStool+BB
6226.33527	0.93024556	0.31721125	2.93257433	0.00336164	0.0381879	Turicibacteraceae	QIAStool vs MagNAPure
88.4317884	-1.3038581	0.44554611	-2.9264268	0.0034288	0.0381879	Coriobacteriaceae	QIAStool vs MagNAPure
32.5255849	-3.9854437	1.38548882	-2.8765614	0.00402034	0.0381879	Staphylococcaceae	QIAStool vs MagNAPure
254.614729	0.76254431	0.27049763	2.81904245	0.00481671	0.0381879	S24-7	QIAStool vs MagNAPure
1574.16355	-0.9682183	0.35443007	-2.7317613	0.00629968	0.0381879	Lachnospiraceae	QIAStool vs MagNAPure
14.253878	2.0500995	0.75196855	2.72631018	0.00640468	0.0381879	Pirellulaceae	QIAStool vs MagNAPure
1019.42191	1.19581281	0.43898336	2.72405041	0.00644867	0.0381879	Campylobacteraceae	QIAStool vs MagNAPure
367.01539	1.41359659	0.52364085	2.69955368	0.00694326	0.0381879	Spirochaetaceae	QIAStool vs MagNAPure
88.4317884	-1.499253	0.60509226	-2.4777262	0.01322226	0.03911584	Coriobacteriaceae	FastDNA vs QIAStool
42.1528852	-2.6816693	0.99400689	-2.6978378	0.00697914	0.03972744	Bifidobacteriaceae	QIAStool vs QIAStool+BB
254.614729	0.77942167	0.29736518	2.62109257	0.00876485	0.04149089	S24-7	EasyDNA vs InnuPURE
20.723165	3.34261958	1.28892921	2.59333062	0.00950513	0.04149089	Pseudomonadaceae	EasyDNA vs InnuPURE
5.22647741	4.53504399	1.75773924	2.58004366	0.00987878	0.04149089	Sphingobacteriaceae	EasyDNA vs InnuPURE
17.2832685	1.59191715	0.59981224	2.65402579	0.00795377	0.04204138	Desulfovibrionaceae	QIAStool vs QIAStool+BB
5.22647741	5.05202816	1.82265879	2.77179041	0.00557489	0.04331319	Sphingobacteriaceae	EasyDNA vs MagNAPure
13.1827693	-2.4080017	0.87080204	-2.7652688	0.00568759	0.04331319	RF16	EasyDNA vs MagNAPure
4.16403335	4.73703395	1.82302293	2.59845001	0.00936457	0.04347835	Carnobacteriaceae	FastDNA vs MagNAPure
2.29903269	-4.3063622	1.6450049	-2.6178416	0.00884879	0.04483386	Deferribacteraceae	EasyDNA vs PowerSoil.HMP
4.16403335	4.24556219	1.70906431	2.48414419	0.01298632	0.04486184	Carnobacteriaceae	EasyDNA vs QIAStool
6226.33527	-0.7266515	0.29467249	-2.4659632	0.01366454	0.04515238	Turicibacteraceae	EasyDNA vs QIAStool
810.892629	-1.5452431	0.62495608	-2.4725627	0.01341482	0.0465047	Erysipelotrichaceae	FastDNA vs InnuPURE
13.1827693	-2.0388951	0.78399797	-2.6006383	0.00930505	0.04652525	RF16	QIAStool vs MagNAPure
1019.42191	1.28455598	0.48482082	2.64954788	0.00805996	0.04656863	Campylobacteraceae	PowerSoil.HMP vs MagNAPure
2.05453403	3.06422851	1.22184327	2.50787363	0.01214601	0.04934316	[Cerasicoccaceae]	PowerSoil.HMP vs QIAStool
2.05453403	-3.5994508	1.41189289	-2.5493795	0.01079148	0.05096371	[Cerasicoccaceae]	EasyDNA vs PowerSoil.HMP
75.3501469	-2.0844308	0.82480975	-2.5271656	0.01149873	0.05096371	Mycoplasmataceae	EasyDNA vs PowerSoil.HMP
2576.15115	0.88393503	0.352154	2.51008088	0.01207035	0.05096371	[Paraprevotellaceae]	EasyDNA vs PowerSoil.HMP
8.45632111	-4.7100683	1.89200483	-2.4894589	0.01279377	0.05117508	Rikenellaceae	EasyDNA vs PowerSoil.HMP
1329.00016	-0.708884	0.28201899	-2.5136037	0.01195046	0.05213896	Peptostreptococcaceae	InnuPURE vs MagNAPure
17.2832685	-1.5355685	0.61355598	-2.5027357	0.01232375	0.05213896	Desulfovibrionaceae	InnuPURE vs MagNAPure
4.16403335	4.13609738	1.72422814	2.39881097	0.0164484	0.05345731	Carnobacteriaceae	FastDNA vs InnuPURE
7.44045113	2.72371757	0.77809085	3.50051356	0.00046436	0.05386606	Anaeroplasmataceae	InnuPURE vs QIAStool+BB
5.8077209	-2.048274	0.83138022	-2.463703	0.013751	0.05402178	Dethiosulfovibrionaceae	InnuPURE vs MagNAPure
15.4343002	1.75738715	0.67749264	2.59395756	0.00948782	0.05421611	Corynebacteriaceae	EasyDNA vs FastDNA
29.6083625	-1.5032074	0.60411383	-2.488285	0.01283608	0.05562303	[Mogibacteriaceae]	FastDNA vs MagNAPure
14.253878	2.21225682	0.86748571	2.55019397	0.0107663	0.05598476	Pirellulaceae	PowerSoil.HMP vs MagNAPure

810.892629	-1.4020193	0.57490323	-2.4387049	0.01474	0.05601199	Erysipelotrichaceae	EasyDNA vs PowerSoil.HMP
2.29903269	-3.8386124	1.56814864	-2.4478626	0.01437064	0.05658441	Deferribacteraceae	EasyDNA vs InnuPURE
908.726526	-1.0058792	0.41567683	-2.4198587	0.01552654	0.05753952	Veillonellaceae	EasyDNA vs InnuPURE
1019.42191	-0.9445218	0.40055924	-2.3580077	0.01837331	0.05818216	Campylobacteraceae	EasyDNA vs QIAStool
71.0660634	-1.1820059	0.44790155	-2.638986	0.00831544	0.05880205	Alcaligenaceae	EasyDNA vs MagNAPure
2.63481049	4.11471371	1.78335384	2.30728957	0.02103868	0.05943999	Cryomorphaceae	FastDNA vs QIAStool
14.253878	-2.3206117	1.01141826	-2.2944135	0.02176676	0.05943999	Pirellulaceae	FastDNA vs QIAStool
5.3314072	-4.2593782	1.62243256	-2.6253037	0.00865717	0.05973448	[Methanomassiliicoccaceae]	FastDNA vs PowerSoil.HMP
2576.15115	0.8484523	0.34193436	2.48133089	0.01308928	0.05999253	[Paraprevotellaceae]	QIAStool vs MagNAPure
576.775751	-1.100544	0.43076515	-2.5548584	0.0106231	0.06126359	Streptococcaceae	FastDNA vs PowerSoil.HMP
31.9177507	-2.2924488	0.90657913	-2.5286804	0.01144922	0.06126359	Succinivibrionaceae	FastDNA vs PowerSoil.HMP
15.8153467	3.36640237	1.3327887	2.52583353	0.01154242	0.06126359	Moraxellaceae	FastDNA vs PowerSoil.HMP
408.703374	1.21382263	0.49751615	2.4397653	0.01469681	0.0621788	Lactobacillaceae	QIAStool vs MagNAPure
8.45632111	4.57043088	1.84612969	2.47568246	0.01329818	0.06286412	Rikenellaceae	PowerSoil.HMP vs MagNAPure
15.8153467	-2.986528	1.18870072	-2.5124305	0.01199027	0.06394811	Moraxellaceae	EasyDNA vs FastDNA
7.44045113	-2.4644294	0.82396759	-2.99093	0.00278129	0.06452599	Anaeroplasmataceae	QIAStool+BB vs MagNAPure
254.614729	0.77621196	0.33732155	2.30110398	0.02138575	0.06541525	S24-7	FastDNA vs InnuPURE
1.25729577	3.29462871	1.41071562	2.3354308	0.01952093	0.06678212	Methanobacteriaceae	InnuPURE vs QIAStool
30.2465249	-3.1037748	1.32591991	-2.3408463	0.01924008	0.06907747	Verrucomicrobiaceae	EasyDNA vs PowerSoil.HMP
1329.00016	-0.7318973	0.31460229	-2.3264208	0.01999611	0.06907747	Peptostreptococcaceae	EasyDNA vs PowerSoil.HMP
7.94956632	1.80438583	0.77397907	2.33131089	0.01973697	0.0690794	Christensenellaceae	EasyDNA vs InnuPURE
1574.16355	-0.9102305	0.38780387	-2.3471412	0.01891808	0.0693663	Lachnospiraceae	InnuPURE vs MagNAPure
30.2465249	2.8646008	1.26008035	2.27334773	0.02300523	0.06993591	Verrucomicrobiaceae	EasyDNA vs QIAStool
7308.6798	-0.5854122	0.25988525	-2.2525795	0.02428568	0.07015862	Prevotellaceae	FastDNA vs InnuPURE
8.45632111	-4.399111	1.79521351	-2.4504667	0.01426711	0.07031649	Rikenellaceae	FastDNA vs PowerSoil.HMP
8.45632111	4.29348299	1.74750758	2.45691809	0.01401346	0.07049952	Rikenellaceae	PowerSoil.HMP vs QIAStool+BB
32.5255849	3.84761504	1.60906299	2.39121468	0.01679273	0.07049952	Staphylococcaceae	PowerSoil.HMP vs QIAStool+BB
2516.24433	0.64083569	0.26815705	2.38977754	0.01685858	0.07049952	Ruminococcaceae	PowerSoil.HMP vs QIAStool+BB
7308.6798	-0.916953	0.31545518	-2.9067617	0.00365191	0.07060364	Prevotellaceae	QIAStool+BB vs MagNAPure
1019.42191	-1.0332649	0.45044532	-2.2938743	0.02179772	0.07202725	Campylobacteraceae	EasyDNA vs PowerSoil.HMP
32.5255849	-3.3654336	1.44833673	-2.3236541	0.02014404	0.07428114	Staphylococcaceae	InnuPURE vs PowerSoil.HMP
5.22647741	4.41383425	1.81274855	2.434885	0.01489652	0.07448262	Sphingobacteriaceae	EasyDNA vs FastDNA
4.2789796	4.42090287	1.86386076	2.37190619	0.01769658	0.07448975	Xanthomonadaceae	PowerSoil.HMP vs MagNAPure
14966.648	-0.8528435	0.36244882	-2.3530039	0.01862244	0.07448975	Clostridiaceae	PowerSoil.HMP vs MagNAPure
42.1528852	2.1770014	0.86093835	2.52863796	0.01145061	0.07557401	Bifidobacteriaceae	EasyDNA vs MagNAPure
31.9177507	-1.9115858	0.86810273	-2.2020272	0.02766339	0.07571032	Succinivibrionaceae	FastDNA vs InnuPURE
71.0660634	1.18925034	0.49362875	2.40919991	0.01598754	0.07887186	Alcaligenaceae	QIAStool vs QIAStool+BB
32.5255849	3.0928854	1.44133799	2.14584325	0.03188549	0.08290226	Staphylococcaceae	FastDNA vs InnuPURE
254.614729	-0.7523977	0.32981434	-2.2812764	0.0225321	0.08369064	S24-7	PowerSoil.HMP vs MagNAPure
408.703374	1.16600506	0.49279223	2.36611902	0.01797566	0.08459136	Lactobacillaceae	EasyDNA vs FastDNA
88.4317884	-1.591849	0.67867494	-2.345525	0.0190003	0.0874014	Coriobacteriaceae	FastDNA vs PowerSoil.HMP
367.01539	-1.3611864	0.59325355	-2.294443	0.02176507	0.08842059	Spirochaetaceae	FastDNA vs MagNAPure
88.4317884	-1.211262	0.54246601	-2.232881	0.0255568	0.08859689	Coriobacteriaceae	PowerSoil.HMP vs MagNAPure
17.2832685	-1.4788953	0.66253337	-2.2321824	0.02560291	0.08865293	Desulfovibrionaceae	InnuPURE vs PowerSoil.HMP
8.45632111	-4.1252908	1.86593498	-2.2108438	0.02704666	0.08865293	Rikenellaceae	InnuPURE vs PowerSoil.HMP
408.703374	-1.5156269	0.63804615	-2.375419	0.01752904	0.09122372	Lactobacillaceae	FastDNA vs QIAStool+BB
7.94956632	2.22772913	0.93886337	2.37279374	0.01765412	0.09122372	Christensenellaceae	FastDNA vs QIAStool+BB
86.5992792	2.35593043	1.00429657	2.34585133	0.01898368	0.09122372	Porphyromonadaceae	FastDNA vs QIAStool+BB
2.60378858	4.2092934	1.79779891	2.34135941	0.01921366	0.09122372	Rhodocyclaceae	FastDNA vs QIAStool+BB
2516.24433	0.61507077	0.26609194	2.31149716	0.02080541	0.09122372	Ruminococcaceae	FastDNA vs QIAStool+BB
2.29903269	3.13747594	1.40445963	2.23393815	0.02548714	0.09345117	Deferribacteraceae	PowerSoil.HMP vs QIAStool
4.16403335	4.04801207	1.8168604	2.22802592	0.02587879	0.09345117	Carnobacteriaceae	PowerSoil.HMP vs QIAStool
62.3051996	-1.4507519	0.631896	-2.2958713	0.02168324	0.09350897	Enterobacteriaceae	FastDNA vs PowerSoil.HMP
29.6083625	-1.0028885	0.45984104	-2.180946	0.02918741	0.09485908	[Mogibacteriaceae]	PowerSoil.HMP vs MagNAPure
7.94956632	-1.8663171	0.86386794	-2.1604194	0.03074022	0.09545646	Christensenellaceae	InnuPURE vs PowerSoil.HMP
14.253878	-2.4827691	1.09644696	-2.2643768	0.02355095	0.09558915	Pirellulaceae	FastDNA vs PowerSoil.HMP
5.3314072	-3.2764048	1.58511548	-2.0669818	0.03873587	0.09591739	[Methanomassiliicoccaceae]	FastDNA vs InnuPURE
35.1041071	-0.9596602	0.42884473	-2.2377801	0.0252354	0.0967357	Peptococcaceae	PowerSoil.HMP vs QIAStool+BB
2.95003546	4.04538074	1.85721321	2.17819942	0.02939119	0.09745501	Actinomycetaceae	EasyDNA vs InnuPURE
5.22647741	3.87561089	1.79473927	2.15942836	0.03081695	0.09758701	Sphingobacteriaceae	EasyDNA vs PowerSoil.HMP
32.5255849	3.57506683	1.60327104	2.22985804	0.02575687	0.09766893	Staphylococcaceae	FastDNA vs QIAStool+BB
5.3314072	-3.6680621	1.66483055	-2.2032645	0.02757611	0.09766893	[Methanomassiliicoccaceae]	FastDNA vs QIAStool+BB
4.00044772	3.823021	1.74418415	2.19186775	0.02838906	0.09766893	Cytophagaceae	FastDNA vs QIAStool+BB
2.63481049	4.11626382	1.88669558	2.18173184	0.02912933	0.09766893	Cryomorphaceae	FastDNA vs QIAStool+BB
31.9177507	1.3344078	0.57886073	2.3052312	0.02115363	0.09783553	Succinivibrionaceae	QIAStool vs QIAStool+BB
2.95003546	4.1628328	1.90223045	2.18839563	0.0286408	0.09798167	Actinomycetaceae	PowerSoil.HMP vs QIAStool
30.2465249	-3.1673216	1.41854368	-2.2327981	0.02556226	0.09798868	Verrucomicrobiaceae	FastDNA vs PowerSoil.HMP
2576.15115	-0.7722792	0.37881863	-2.0386515	0.04148482	0.09805502	[Paraprevotellaceae]	FastDNA vs InnuPURE
20.723165	-2.6807638	1.24693501	-2.1498826	0.03156451	0.09811055	Pseudomonadaceae	InnuPURE vs QIAStool
7.94956632	-1.557632	0.72508445	-2.1482077	0.03169726	0.09811055	Christensenellaceae	InnuPURE vs QIAStool

7.46995934	-1.5690896	0.72723812	-2.1576009	0.03095888	0.09843754	RFP12	InnuPURE vs MagNAPure
22.9687795	1.19410946	0.55614757	2.14710901	0.0317846	0.09843754	Helicobacteraceae	InnuPURE vs MagNAPure
9.6383629	-1.384477	0.64643194	-2.1417212	0.03221592	0.09843754	Fibrobacteraceae	InnuPURE vs MagNAPure
9.6383629	-1.8427971	0.80591343	-2.2865943	0.02221951	0.09875339	Fibrobacteraceae	EasyDNA vs FastDNA
5.3314072	-3.5558936	1.59551674	-2.2286783	0.02583532	0.09878209	[Methanomassiliicoccaceae]	FastDNA vs MagNAPure
1574.16355	-0.86689	0.35975205	-2.4096875	0.01596619	0.0987908	Lachnospiraceae	EasyDNA vs MagNAPure
7308.6798	0.55606923	0.26154925	2.12605939	0.03349831	0.09882003	Prevotellaceae	InnuPURE vs PowerSoil.HMP
30.2465249	3.46560811	1.53285987	2.26087732	0.02376685	0.09902856	Verrucomicrobiaceae	EasyDNA vs QIAStool+BB
14966.648	0.71562954	0.32014118	2.23535608	0.02539398	0.09976207	Clostridiaceae	QIAStool vs MagNAPure
75.3501469	-1.6556432	0.80348783	-2.0605704	0.03934405	0.09978224	Mycoplasmataceae	FastDNA vs QIAStool
30.2465249	2.801054	1.35940485	2.06050023	0.03935074	0.09978224	Verrucomicrobiaceae	FastDNA vs QIAStool
8.41300461	2.92327159	1.38042248	2.11766444	0.0342035	0.09997945	Neisseriaceae	EasyDNA vs QIAStool

Table S2C. Difference abundance of families - Sewage microbial community

baseMean	log2FoldChange	lfcSE	stat	pvalue	padj	family	compare
649.223144	6.82534658	0.30988145	22.0256701	1.63E-107	3.12E-105	Bifidobacteriaceae	EasyDNA vs QIAStool
649.223144	5.53968468	0.33942717	16.3206873	7.03E-60	1.15E-57	Bifidobacteriaceae	EasyDNA vs PowerSoil.HMP
858.112951	4.52279196	0.28850532	15.6766329	2.19E-55	3.13E-53	Legionellaceae	EasyDNA vs FastDNA
649.223144	4.65718093	0.31803068	14.6438102	1.48E-48	1.99E-46	Bifidobacteriaceae	EasyDNA vs MagNAPure
283.625111	6.76475532	0.54891922	12.323772	6.75E-35	6.44E-33	Lactobacillaceae	EasyDNA vs QIAStool
649.223144	4.37138738	0.3572889	12.2348815	2.02E-34	3.06E-32	Bifidobacteriaceae	EasyDNA vs QIAStool+BB
458.885072	4.53073551	0.37259059	12.1600912	5.07E-34	3.23E-32	Carnobacteriaceae	EasyDNA vs QIAStool
649.223144	3.79123624	0.31179797	12.1592717	5.12E-34	3.66E-32	Bifidobacteriaceae	EasyDNA vs FastDNA
858.112951	3.26216457	0.29164006	11.1855845	4.80E-29	3.91E-27	Legionellaceae	EasyDNA vs PowerSoil.HMP
283.625111	6.53180922	0.59946607	10.8960449	1.20E-27	8.13E-26	Lactobacillaceae	EasyDNA vs MagNAPure
858.112951	3.3518407	0.31881432	10.5134571	7.49E-26	5.65E-24	Legionellaceae	EasyDNA vs QIAStool+BB
858.112951	-3.2399311	0.31273297	-10.360056	3.77E-25	4.18E-23	Legionellaceae	FastDNA vs MagNAPure
505.493872	3.75708256	0.37090716	10.129442	4.09E-24	1.95E-22	Erysipelotrichaceae	EasyDNA vs QIAStool
3668.6678	4.98922885	0.49392154	10.1012579	5.45E-24	2.45E-22	Lachnospiraceae	EasyDNA vs MagNAPure
458.885072	4.14006653	0.40630321	10.1895985	2.21E-24	3.16E-22	Carnobacteriaceae	FastDNA vs QIAStool
649.223144	3.76051584	0.391297	9.6103875	7.23E-22	1.18E-19	Bifidobacteriaceae	EasyDNA vs InnuPURE
103.008126	3.09894526	0.32848091	9.43417169	3.94E-21	1.51E-19	Enterococcaceae	EasyDNA vs QIAStool
3668.6678	4.06945931	0.44648024	9.11453394	7.90E-20	2.52E-18	Lachnospiraceae	EasyDNA vs QIAStool
458.885072	-3.9036097	0.42130142	-9.2655984	1.94E-20	2.70E-18	Carnobacteriaceae	QIAStool vs QIAStool+BB
505.493872	3.74637486	0.41138597	9.10671522	8.49E-20	2.87E-18	Erysipelotrichaceae	EasyDNA vs MagNAPure
649.223144	3.03411034	0.3389894	8.9504579	3.54E-19	2.53E-17	Bifidobacteriaceae	FastDNA vs QIAStool
49.8040986	4.33239316	0.49224425	8.80130772	1.35E-18	3.69E-17	Actinomycetaceae	EasyDNA vs QIAStool
4234.46977	2.87073548	0.32448599	8.84702452	8.99E-19	4.28E-17	Ruminococcaceae	EasyDNA vs FastDNA
858.112951	2.1585116	0.25552818	8.44725466	2.98E-17	7.12E-16	Legionellaceae	EasyDNA vs QIAStool
283.625111	5.01823943	0.58931798	8.51533406	1.66E-17	9.03E-16	Lactobacillaceae	EasyDNA vs PowerSoil.HMP
2750.67146	3.61050489	0.43024624	8.39171743	4.79E-17	1.29E-15	Bacteroidaceae	EasyDNA vs MagNAPure
4234.46977	2.65205125	0.32465158	8.16891528	3.11E-16	7.00E-15	Ruminococcaceae	EasyDNA vs MagNAPure
458.885072	4.13915825	0.49523734	8.35792836	6.38E-17	8.87E-15	Carnobacteriaceae	InnuPURE vs QIAStool
858.112951	-2.3642804	0.2878549	-8.213445	2.15E-16	1.02E-14	Legionellaceae	FastDNA vs QIAStool
719.094782	2.76017849	0.34134669	8.08614399	6.16E-16	2.51E-14	Streptococcaceae	EasyDNA vs PowerSoil.HMP
719.094782	2.39430118	0.30288969	7.90486188	2.68E-15	5.69E-14	Streptococcaceae	EasyDNA vs QIAStool
103.008126	2.8436344	0.36066035	7.88452173	3.16E-15	6.09E-14	Enterococcaceae	EasyDNA vs MagNAPure
103.008126	3.01172614	0.38076559	7.90965945	2.58E-15	8.41E-14	Enterococcaceae	EasyDNA vs PowerSoil.HMP
505.493872	3.18511617	0.40804762	7.80574613	5.92E-15	1.71E-13	Erysipelotrichaceae	EasyDNA vs FastDNA
3668.6678	3.84726205	0.49296566	7.80432056	5.98E-15	1.71E-13	Lachnospiraceae	EasyDNA vs FastDNA
33.4444509	4.99601637	0.68201466	7.32537975	2.38E-13	4.55E-12	Coriobacteriaceae	EasyDNA vs QIAStool
118.339928	-2.2031982	0.29992413	-7.3458517	2.04E-13	7.31E-12	[Chthoniobacteraceae]	FastDNA vs QIAStool
649.223144	3.06483074	0.41298549	7.42115838	1.16E-13	8.07E-12	Bifidobacteriaceae	InnuPURE vs QIAStool
49.8040986	3.68571012	0.5128394	7.18687007	6.63E-13	1.90E-11	Actinomycetaceae	FastDNA vs QIAStool
3668.6678	3.52725908	0.49371146	7.14437356	9.04E-13	2.15E-11	Lachnospiraceae	EasyDNA vs PowerSoil.HMP
505.493872	2.94927376	0.41296872	7.14163966	9.22E-13	2.15E-11	Erysipelotrichaceae	EasyDNA vs PowerSoil.HMP
186.231671	-2.2374803	0.31528142	-7.0967719	1.28E-12	8.88E-11	Rhizobiaceae	QIAStool vs QIAStool+BB
719.094782	2.55887653	0.36330665	7.04329673	1.88E-12	9.45E-11	Streptococcaceae	EasyDNA vs QIAStool+BB
858.112951	2.49333772	0.35533582	7.0168488	2.27E-12	1.85E-10	Legionellaceae	EasyDNA vs InnuPURE
3668.6678	3.35179711	0.49869049	6.72119718	1.80E-11	6.80E-10	Lachnospiraceae	EasyDNA vs QIAStool+BB
4234.46977	2.19156111	0.32972499	6.64663329	3.00E-11	9.06E-10	Ruminococcaceae	EasyDNA vs QIAStool+BB
993.963868	-2.9717971	0.45776516	-6.4919687	8.47E-11	1.47E-09	Opituaceae	EasyDNA vs QIAStool
505.493872	3.34829908	0.50439697	6.63822198	3.17E-11	1.73E-09	Erysipelotrichaceae	EasyDNA vs InnuPURE
49.8040986	3.93076665	0.601821	6.53145479	6.51E-11	3.02E-09	Actinomycetaceae	InnuPURE vs QIAStool
4234.46977	1.84495753	0.29167984	6.32528303	2.53E-10	4.02E-09	Ruminococcaceae	EasyDNA vs QIAStool
649.223144	-2.4539592	0.3810699	-6.4396564	1.20E-10	5.55E-09	Bifidobacteriaceae	QIAStool vs QIAStool+BB
505.493872	2.75232367	0.43583608	6.31504315	2.70E-10	6.80E-09	Erysipelotrichaceae	EasyDNA vs QIAStool+BB
458.885072	2.49945084	0.40038248	6.24265787	4.30E-10	7.26E-09	Carnobacteriaceae	EasyDNA vs MagNAPure
458.885072	2.52065316	0.40591353	6.20982789	5.30E-10	1.02E-08	Carnobacteriaceae	EasyDNA vs PowerSoil.HMP
47.9856545	-3.2556254	0.52513758	-6.1995666	5.66E-10	1.02E-08	Bacteriovoracaceae	EasyDNA vs PowerSoil.HMP
49.8040986	3.25085928	0.52651405	6.17430685	6.65E-10	1.02E-08	Actinomycetaceae	EasyDNA vs PowerSoil.HMP
33.4444509	4.67402796	0.75777778	6.16807211	6.91E-10	1.02E-08	Coriobacteriaceae	EasyDNA vs PowerSoil.HMP
858.112951	-1.9793037	0.31561673	-6.2712254	3.58E-10	2.50E-08	Legionellaceae	PowerSoil.HMP vs MagNAPure
622.632062	-2.1934305	0.35305951	-6.2126369	5.21E-10	2.50E-08	Flavobacteriaceae	PowerSoil.HMP vs MagNAPure
283.625111	3.63327256	0.59610996	6.09497039	1.09E-09	2.61E-08	Lactobacillaceae	FastDNA vs QIAStool
141.200238	-2.2467743	0.37457141	-5.9982536	1.99E-09	2.71E-08	Chitinophagaceae	EasyDNA vs PowerSoil.HMP
649.223144	-2.1681657	0.3447053	-6.2899111	3.18E-10	6.58E-08	Bifidobacteriaceae	QIAStool vs MagNAPure
616.023371	2.17009588	0.37325392	5.8139935	6.10E-09	9.15E-08	Veillonellaceae	EasyDNA vs MagNAPure

103.008126	2.0301104	0.34927259	5.81239536	6.16E-09	1.26E-07	Enterococcaceae	FastDNA vs QIAStool
858.112951	-2.0689798	0.34079652	-6.0710121	1.27E-09	1.31E-07	Legionellaceae	QIAStool+BB vs MagNAPure
141.200238	-2.3121306	0.39864712	-5.7999431	6.63E-09	1.43E-07	Chitinophagaceae	EasyDNA vs QIAStool+BB
2750.67146	2.58911904	0.42988976	6.02275112	1.71E-09	1.77E-07	Bacteroidaceae	QIAStool vs MagNAPure
579.211848	-1.7141861	0.29702596	-5.771166	7.87E-09	1.88E-07	Cryomorphaceae	EasyDNA vs FastDNA
622.632062	-1.8973044	0.31931391	-5.9418158	2.82E-09	1.94E-07	Flavobacteriaceae	QIAStool vs MagNAPure
283.625111	-3.671693	0.63002671	-5.8278371	5.62E-09	1.95E-07	Lactobacillaceae	QIAStool vs QIAStool+BB
118.339928	1.81003807	0.31730854	5.70434721	1.17E-08	2.39E-07	[Chthoniobacteraceae]	EasyDNA vs FastDNA
639.445335	-1.9434602	0.3470908	-5.5992847	2.15E-08	2.70E-07	Marinilabiaceae	EasyDNA vs PowerSoil.HMP
47.9856545	-2.7589818	0.49172905	-5.6107766	2.01E-08	2.96E-07	Bacteriovoraceae	EasyDNA vs QIAStool
50.9419349	-3.1533338	0.56681969	-5.5632045	2.65E-08	3.08E-07	Desulfobulbaceae	EasyDNA vs PowerSoil.HMP
4234.46977	1.80165225	0.3248057	5.54686163	2.91E-08	3.16E-07	Ruminococcaceae	EasyDNA vs PowerSoil.HMP
169.692077	-2.5020001	0.43818754	-5.7098842	1.13E-08	3.62E-07	Pirellulaceae	PowerSoil.HMP vs MagNAPure
283.625111	3.13148276	0.56384853	5.55376592	2.80E-08	5.00E-07	Lactobacillaceae	EasyDNA vs FastDNA
639.445335	-1.9819676	0.35653587	-5.558957	2.71E-08	5.12E-07	Marinilabiaceae	EasyDNA vs QIAStool+BB
47.9856545	-2.8371254	0.51608043	-5.4974481	3.85E-08	6.12E-07	Bacteriovoraceae	EasyDNA vs FastDNA
186.231671	-1.6588195	0.29104376	-5.6995534	1.20E-08	6.22E-07	Rhizobiaceae	QIAStool vs MagNAPure
49.8040986	2.6607802	0.48751569	5.45783502	4.82E-08	6.51E-07	Actinomycetaceae	EasyDNA vs MagNAPure
141.200238	-2.0144185	0.3687622	-5.4626491	4.69E-08	6.71E-07	Chitinophagaceae	EasyDNA vs FastDNA
33.4444509	3.7400853	0.68725107	5.44209458	5.27E-08	6.85E-07	Coriobacteriaceae	EasyDNA vs FastDNA
4234.46977	2.2738468	0.40301486	5.64209171	1.68E-08	6.85E-07	Ruminococcaceae	EasyDNA vs InnuPURE
50.9419349	-2.9788598	0.55810891	-5.3374166	9.43E-08	1.12E-06	Desulfobulbaceae	EasyDNA vs FastDNA
224.417038	-2.2900472	0.43302783	-5.2884528	1.23E-07	1.26E-06	Caulobacteraceae	EasyDNA vs PowerSoil.HMP
579.211848	-1.5784754	0.29952912	-5.2698562	1.37E-07	1.31E-06	Cryomorphaceae	EasyDNA vs PowerSoil.HMP
60.913589	1.80194128	0.33612104	5.36098928	8.28E-08	1.48E-06	SB-1	FastDNA vs QIAStool
50.9419349	-3.4975916	0.64107451	-5.455827	4.87E-08	1.59E-06	Desulfobulbaceae	EasyDNA vs InnuPURE
264.350163	1.67550628	0.29505857	5.67855486	1.36E-08	1.73E-06	Leptotrichiaceae	PowerSoil.HMP vs QIAStool
24.9235631	-2.9250852	0.56407379	-5.1856428	2.15E-07	1.95E-06	[Bryobacteraceae]	EasyDNA vs PowerSoil.HMP
2750.67146	2.31054863	0.43619881	5.29700806	1.18E-07	1.98E-06	Bacteroidaceae	EasyDNA vs QIAStool+BB
50.9419349	-2.9345835	0.56106287	-5.2304005	1.69E-07	2.08E-06	Desulfobulbaceae	EasyDNA vs MagNAPure
2750.67146	2.23082024	0.42928279	5.1966217	2.03E-07	2.23E-06	Bacteroidaceae	EasyDNA vs FastDNA
164.684355	2.43591952	0.46885051	5.19551427	2.04E-07	2.30E-06	Clostridiaceae	EasyDNA vs MagNAPure
616.023371	1.91438373	0.37167225	5.15073087	2.59E-07	2.65E-06	Veillonellaceae	EasyDNA vs FastDNA
579.211848	-1.5289396	0.29774329	-5.1350935	2.82E-07	2.93E-06	Cryomorphaceae	EasyDNA vs MagNAPure
264.350163	-1.5725743	0.29211855	-5.3833429	7.31E-08	3.03E-06	Leptotrichiaceae	QIAStool vs MagNAPure
47.9856545	-2.8959851	0.56014488	-5.1700644	2.34E-07	3.48E-06	Bacteriovoraceae	EasyDNA vs QIAStool+BB
283.625111	3.09306229	0.60001044	5.15501411	2.54E-07	3.48E-06	Lactobacillaceae	EasyDNA vs QIAStool+BB
104.663517	2.01013043	0.39723097	5.06035681	4.18E-07	4.04E-06	Rikenellaceae	EasyDNA vs MagNAPure
719.094782	2.18960446	0.41686817	5.2525105	1.50E-07	4.08E-06	Streptococcaceae	EasyDNA vs InnuPURE
49.8040986	-2.9975018	0.57075775	-5.2517935	1.51E-07	4.19E-06	Actinomycetaceae	QIAStool vs QIAStool+BB
28.0267224	3.26712548	0.65449599	4.99181896	5.98E-07	5.38E-06	Succinivibrionaceae	EasyDNA vs MagNAPure
3668.6678	3.086453	0.5976439	5.16436792	2.41E-07	5.62E-06	Lachnospiraceae	EasyDNA vs InnuPURE
22.1599751	3.76303745	0.74421723	5.05636971	4.27E-07	5.83E-06	Sanguibacteraceae	EasyDNA vs QIAStool
283.625111	3.40032645	0.64218587	5.29492571	1.19E-07	6.61E-06	Lactobacillaceae	FastDNA vs MagNAPure
186.231671	1.46311188	0.28954379	5.05316272	4.35E-07	6.90E-06	Rhizobiaceae	FastDNA vs QIAStool
131.739477	-1.6538134	0.33613801	-4.9200427	8.65E-07	7.30E-06	Hyphomicrobiaceae	EasyDNA vs MagNAPure
719.094782	1.64683881	0.33485216	4.91810712	8.74E-07	8.33E-06	Streptococcaceae	EasyDNA vs FastDNA
639.445335	-1.6956173	0.34566547	-4.9053709	9.33E-07	8.33E-06	Marinilabiaceae	EasyDNA vs FastDNA
7.94601326	4.25743862	0.86954011	4.8961958	9.77E-07	1.24E-05	Micrococcaceae	EasyDNA vs QIAStool
354.810526	-1.2016891	0.25020884	-4.8027444	1.57E-06	1.25E-05	Pseudomonadaceae	EasyDNA vs FastDNA
20.5932418	3.40730412	0.70969866	4.80105753	1.58E-06	1.25E-05	[Mogibacteriaceae]	EasyDNA vs FastDNA
47.9856545	-2.9519041	0.59210402	-4.9854486	6.18E-07	1.26E-05	Bacteriovoraceae	EasyDNA vs InnuPURE
33.4444509	3.30560527	0.69235087	4.77446541	1.80E-06	1.43E-05	Coriobacteriaceae	EasyDNA vs MagNAPure
31.145957	4.67590454	0.90414752	5.17161683	2.32E-07	1.47E-05	Aerococcaceae	PowerSoil.HMP vs QIAStool
1110.97909	-0.8552071	0.17647177	-4.8461411	1.26E-06	1.50E-05	Verrucomicrobiaceae	EasyDNA vs QIAStool
33.4444509	3.74329586	0.7734292	4.83986884	1.30E-06	1.63E-05	Coriobacteriaceae	EasyDNA vs QIAStool+BB
283.625111	3.43874693	0.67321308	5.10796215	3.26E-07	1.68E-05	Lactobacillaceae	QIAStool+BB vs MagNAPure
616.023371	2.26022292	0.4609869	4.90300899	9.44E-07	1.71E-05	Veillonellaceae	EasyDNA vs InnuPURE
92.6279814	2.25174145	0.47401585	4.75035053	2.03E-06	1.74E-05	Oxalobacteraceae	EasyDNA vs PowerSoil.HMP
858.112951	-2.0294542	0.37889716	-5.356214	8.50E-08	1.76E-05	Legionellaceae	FastDNA vs InnuPURE
44.0121162	-2.6519931	0.53414421	-4.9649384	6.87E-07	1.84E-05	Microbacteriaceae	QIAStool vs MagNAPure
458.885072	-2.0312847	0.410162	-4.9523961	7.33E-07	1.84E-05	Carnobacteriaceae	QIAStool vs MagNAPure
28.0267224	3.13232418	0.63285923	4.9494801	7.44E-07	1.84E-05	Succinivibrionaceae	QIAStool vs MagNAPure
104.663517	1.89306701	0.38356181	4.93549398	7.99E-07	1.84E-05	Rikenellaceae	QIAStool vs MagNAPure
283.625111	3.52238694	0.70298392	5.01062233	5.43E-07	1.89E-05	Lactobacillaceae	InnuPURE vs QIAStool

993.963868	2.43312399	0.49693261	4.89628565	9.77E-07	2.02E-05	Opitutaceae	QIAStool vs MagNAPure
224.417038	-2.0089553	0.43129404	-4.6579715	3.19E-06	2.40E-05	Caulobacteraceae	EasyDNA vs MagNAPure
2750.67146	2.5329193	0.52617828	4.81380436	1.48E-06	2.41E-05	Bacteroidaceae	EasyDNA vs InnuPURE
283.625111	3.24236838	0.67710254	4.78859282	1.68E-06	2.49E-05	Lactobacillaceae	EasyDNA vs InnuPURE
579.211848	-1.7395483	0.3646933	-4.7698939	1.84E-06	2.50E-05	Cryomorphaceae	EasyDNA vs InnuPURE
616.023371	1.78559065	0.37061859	4.81786587	1.45E-06	2.73E-05	Veillonellaceae	QIAStool vs MagNAPure
127.559639	-1.7610135	0.38087354	-4.6236173	3.77E-06	3.07E-05	Bradyrhizobiaceae	EasyDNA vs PowerSoil.HMP
354.810526	-1.1755292	0.25502204	-4.6095201	4.04E-06	3.13E-05	Pseudomonadaceae	EasyDNA vs PowerSoil.HMP
1110.97909	1.00356055	0.20789468	4.82725467	1.38E-06	3.21E-05	Verrucomicrobiaceae	QIAStool vs QIAStool+BB
458.885072	2.12998418	0.43685713	4.8756997	1.08E-06	3.63E-05	Carnobacteriaceae	FastDNA vs PowerSoil.HMP
103.008126	1.94289127	0.39867209	4.87340682	1.10E-06	3.63E-05	Enterococcaceae	FastDNA vs PowerSoil.HMP
649.223144	1.74844843	0.36606848	4.77628779	1.79E-06	3.63E-05	Bifidobacteriaceae	FastDNA vs PowerSoil.HMP
49.8040986	2.60417624	0.54562554	4.77282692	1.82E-06	3.63E-05	Actinomycetaceae	FastDNA vs PowerSoil.HMP
4830.98886	1.39637929	0.30634779	4.55815037	5.16E-06	3.82E-05	Moraxellaceae	EasyDNA vs PowerSoil.HMP
458.885072	2.10878186	0.43175866	4.8841681	1.04E-06	3.84E-05	Carnobacteriaceae	FastDNA vs MagNAPure
186.231671	1.44657273	0.29780928	4.8573796	1.19E-06	4.18E-05	Rhizobiaceae	PowerSoil.HMP vs QIAStool
458.885072	2.01008235	0.41555158	4.83714281	1.32E-06	4.18E-05	Carnobacteriaceae	PowerSoil.HMP vs QIAStool
127.559639	1.70915508	0.35669596	4.79163005	1.65E-06	4.20E-05	Bradyrhizobiaceae	PowerSoil.HMP vs QIAStool
858.112951	1.28286089	0.28337494	4.5270795	5.98E-06	4.25E-05	Legionellaceae	EasyDNA vs MagNAPure
113.494552	1.61419048	0.36198295	4.45929977	8.22E-06	5.55E-05	Alcaligenaceae	EasyDNA vs MagNAPure
2750.67146	1.91527189	0.42988963	4.45526421	8.38E-06	5.94E-05	Bacteroidaceae	EasyDNA vs PowerSoil.HMP
103.008126	1.77479953	0.37958812	4.67559296	2.93E-06	6.60E-05	Enterococcaceae	FastDNA vs MagNAPure
21.4381793	4.3110629	0.92258237	4.67282169	2.97E-06	6.60E-05	Turicibacteraceae	FastDNA vs MagNAPure
186.231671	-1.4894853	0.32863304	-4.5323662	5.83E-06	6.77E-05	Rhizobiaceae	EasyDNA vs QIAStool+BB
719.094782	1.45997524	0.33529292	4.35432771	1.33E-05	8.58E-05	Streptococcaceae	EasyDNA vs MagNAPure
24.9235631	-2.4240291	0.5584747	-4.3404456	1.42E-05	8.73E-05	[Bryobacteraceae]	EasyDNA vs MagNAPure
141.200238	-1.9858992	0.44217029	-4.4912543	7.08E-06	8.88E-05	Chitinophagaceae	EasyDNA vs InnuPURE
60.913589	-1.5556429	0.34130474	-4.5579295	5.17E-06	8.91E-05	SB-1	QIAStool vs MagNAPure
21.4381793	3.83550551	0.84989348	4.51292495	6.39E-06	9.14E-05	Turicibacteraceae	FastDNA vs QIAStool
47.9856545	-2.257859	0.52469926	-4.3031488	1.68E-05	9.88E-05	Bacteriovoracaceae	EasyDNA vs MagNAPure
129.664767	-1.7354308	0.40486896	-4.2864011	1.82E-05	0.00010214	Sphingomonadaceae	EasyDNA vs MagNAPure
186.231671	1.65979847	0.36004846	4.60993081	4.03E-06	0.00010414	Rhizobiaceae	InnuPURE vs QIAStool
7.94601326	4.39122021	0.9573039	4.5870702	4.50E-06	0.00010414	Micrococcaceae	InnuPURE vs QIAStool
103.008126	1.74349936	0.39463833	4.41796766	9.96E-06	0.00010746	Enterococcaceae	EasyDNA vs QIAStool+BB
24.9235631	-2.6423032	0.60049176	-4.4002323	1.08E-05	0.00010886	[Bryobacteraceae]	EasyDNA vs QIAStool+BB
639.445335	-1.4715753	0.34627638	-4.2497132	2.14E-05	0.00011558	Marinilabiaceae	EasyDNA vs MagNAPure
59.9667273	1.93226679	0.42523105	4.54403979	5.52E-06	0.00011681	Phyllobacteriaceae	PowerSoil.HMP vs QIAStool
21.4381793	4.27081182	0.9437759	4.52523934	6.03E-06	0.00011979	Turicibacteraceae	InnuPURE vs QIAStool
622.632062	-1.3757081	0.32541716	-4.2275217	2.36E-05	0.00012268	Flavobacteriaceae	EasyDNA vs MagNAPure
79.5100271	-1.7761899	0.42104528	-4.2185246	2.46E-05	0.00012295	Bdellovibrionaceae	EasyDNA vs MagNAPure
616.023371	1.72314588	0.3956667	4.35504397	1.33E-05	0.00012556	Veillonellaceae	EasyDNA vs QIAStool+BB
269.467613	3.83469555	0.91517191	4.19013688	2.79E-05	0.00013441	[Paraprevotellaceae]	EasyDNA vs MagNAPure
224.417038	-1.9574683	0.45287758	-4.3222901	1.54E-05	0.00013716	Caulobacteraceae	EasyDNA vs QIAStool+BB
622.632062	1.57003075	0.35300582	4.44760591	8.68E-06	0.00013893	Flavobacteriaceae	FastDNA vs PowerSoil.HMP
579.211848	-1.3487004	0.31530459	-4.2774525	1.89E-05	0.00015859	Cryomorphaceae	EasyDNA vs QIAStool+BB
17.3475351	-3.0168408	0.68543834	-4.4013307	1.08E-05	0.00017131	Isosphaeraceae	QIAStool vs MagNAPure
642.395629	-1.4644737	0.33909709	-4.3187444	1.57E-05	0.0001763	Rhodocyclaceae	EasyDNA vs QIAStool
21.4381793	4.36627385	0.95987712	4.54878417	5.40E-06	0.00018525	Turicibacteraceae	QIAStool+BB vs MagNAPure
736.495903	-3.1494341	0.72564066	-4.3402117	1.42E-05	0.00018979	Enterobacteriaceae	FastDNA vs PowerSoil.HMP
354.810526	-0.9826337	0.22992472	-4.273719	1.92E-05	0.00020399	Pseudomonadaceae	EasyDNA vs QIAStool
310.195483	-1.5002553	0.3530494	-4.2494202	2.14E-05	0.00021545	Aeromonadaceae	EasyDNA vs QIAStool
127.559639	-1.5335396	0.37748821	-4.0624835	4.86E-05	0.00022602	Bradyrhizobiaceae	EasyDNA vs MagNAPure
736.495903	3.2147007	0.72609436	4.42738697	9.54E-06	0.00022892	Enterobacteriaceae	PowerSoil.HMP vs MagNAPure
21.4381793	3.65889131	0.90779156	4.03054124	5.56E-05	0.00025042	Turicibacteraceae	EasyDNA vs MagNAPure
22.1599751	3.8088383	0.87850368	4.33559743	1.45E-05	0.00025257	Sanguibacteraceae	InnuPURE vs QIAStool
28.7248963	-2.4613767	0.5983952	-4.1132963	3.90E-05	0.0002592	Acetobacteraceae	EasyDNA vs PowerSoil.HMP
20.5932418	3.06459248	0.74584242	4.10890073	3.98E-05	0.0002592	[Mogibacteriaceae]	EasyDNA vs PowerSoil.HMP
21.4381793	-3.8907165	0.8934125	-4.3548937	1.33E-05	0.00026436	Turicibacteraceae	QIAStool vs QIAStool+BB
20.5932418	2.73509037	0.65453561	4.17867316	2.93E-05	0.00028002	[Mogibacteriaceae]	EasyDNA vs QIAStool
169.692077	-1.6566863	0.38767879	-4.2733476	1.93E-05	0.00028471	Pirellulaceae	QIAStool vs MagNAPure
19.2945448	-2.3378592	0.57446007	-4.0696636	4.71E-05	0.00029516	BA008	EasyDNA vs PowerSoil.HMP
59.9667273	-1.9028345	0.45384514	-4.1926957	2.76E-05	0.00031504	Phyllobacteriaceae	FastDNA vs PowerSoil.HMP
24.9235631	2.00013401	0.46676422	4.28510563	1.83E-05	0.00033138	[Bryobacteraceae]	PowerSoil.HMP vs QIAStool
118.159531	-1.6325049	0.41315264	-3.9513361	7.77E-05	0.00033844	Rhodobacteraceae	EasyDNA vs MagNAPure
31.145957	4.0364622	0.94315101	4.27976235	1.87E-05	0.00035922	Aerococcaceae	PowerSoil.HMP vs MagNAPure

127.559639	-1.4816812	0.35306201	-4.19666	2.71E-05	0.00036159	Bradyrhizobiaceae	QIAStool vs MagNAPure
118.339928	1.24831573	0.29795829	4.18956536	2.79E-05	0.00036159	[Chthoniobacteraceae]	QIAStool vs MagNAPure
59.9667273	-1.8333257	0.46049263	-3.981227	6.86E-05	0.0004139	Phyllobacteriaceae	EasyDNA vs PowerSoil.HMP
639.445335	1.40828004	0.33538657	4.19897568	2.68E-05	0.00042565	Marinilabiaceae	PowerSoil.HMP vs QIAStool
79.5100271	-1.6859989	0.41893896	-4.0244501	5.71E-05	0.00042982	Bdellovibrionaceae	EasyDNA vs FastDNA
24.9235631	-2.221681	0.55402879	-4.0100462	6.07E-05	0.00043405	[Bryobacteraceae]	EasyDNA vs FastDNA
118.339928	-1.589293	0.37966561	-4.1860337	2.84E-05	0.00043842	[Chthoniobacteraceae]	InnuPURE vs QIAStool
616.023371	-1.5298785	0.3690248	-4.1457335	3.39E-05	0.00044035	Veillonellaceae	FastDNA vs QIAStool
6.92597984	-3.2118216	0.79077444	-4.0616154	4.87E-05	0.00044325	[Chromatiaceae]	EasyDNA vs QIAStool
354.810526	-1.2624747	0.30675563	-4.1155713	3.86E-05	0.00044967	Pseudomonadaceae	EasyDNA vs InnuPURE
21.4381793	4.74636921	1.00321669	4.73115056	2.23E-06	0.00046213	Turicibacteraceae	InnuPURE vs MagNAPure
639.445335	-1.4467874	0.34517174	-4.1915002	2.77E-05	0.00048149	Marinilabiaceae	QIAStool vs QIAStool+BB
17.3475351	-3.0488699	0.72528896	-4.2036624	2.63E-05	0.00048587	Isosphaeraceae	FastDNA vs MagNAPure
91.6048239	-1.2060837	0.30202592	-3.9933119	6.52E-05	0.00053723	Halomonadaceae	EasyDNA vs QIAStool
50.9419349	-2.1166588	0.53101195	-3.986085	6.72E-05	0.00053723	Desulfobulbaceae	EasyDNA vs QIAStool
141.200238	-1.3720459	0.34431013	-3.9849128	6.75E-05	0.00053723	Chitinophagaceae	EasyDNA vs QIAStool
70.3986464	1.43164248	0.37367651	3.83123484	0.0001275	0.0005379	Desulfovibrionaceae	EasyDNA vs MagNAPure
103.008126	1.77265924	0.43241403	4.0994489	4.14E-05	0.00055112	Enterococcaceae	InnuPURE vs QIAStool
616.023371	-1.8757177	0.45889686	-4.0874494	4.36E-05	0.00055112	Veillonellaceae	InnuPURE vs QIAStool
11.2601759	3.51749155	0.88819118	3.9602865	7.49E-05	0.00057193	Peptococcaceae	EasyDNA vs QIAStool
310.195483	-1.5679178	0.3821482	-4.1029052	4.08E-05	0.00057573	Aeromonadaceae	PowerSoil.HMP vs QIAStool
50.9419349	-2.4153589	0.61074665	-3.9547641	7.66E-05	0.00060885	Desulfobulbaceae	EasyDNA vs QIAStool+BB
118.339928	1.41559349	0.34454956	4.10853378	3.98E-05	0.00061497	[Chthoniobacteraceae]	QIAStool vs QIAStool+BB
182.91656	-1.4289407	0.34903446	-4.0939816	4.24E-05	0.00067844	Planctomycetaceae	PowerSoil.HMP vs MagNAPure
458.885072	1.87232506	0.44582209	4.19971356	2.67E-05	0.00068818	Carnobacteriaceae	QIAStool+BB vs MagNAPure
593.953594	-0.8557847	0.22798878	-3.7536264	0.00017429	0.00071302	Cytophagaceae	EasyDNA vs MagNAPure
622.632062	1.27390462	0.3192542	3.99025176	6.60E-05	0.00073743	Flavobacteriaceae	FastDNA vs QIAStool
12.2684488	-2.4781182	0.62161852	-3.9865578	6.70E-05	0.00073743	Synergistaceae	FastDNA vs QIAStool
164.684355	1.63111762	0.42086767	3.87560684	0.00010636	0.00078133	Clostridiaceae	EasyDNA vs QIAStool
17.3475351	-2.7235825	0.73173203	-3.7221038	0.00019757	0.00078447	Isosphaeraceae	EasyDNA vs MagNAPure
12.0832988	2.86029733	0.72533982	3.94338935	8.03E-05	0.0008206	Microthrixaceae	FastDNA vs QIAStool
858.112951	-1.2606274	0.32020896	-3.9368897	8.25E-05	0.00082545	Legionellaceae	FastDNA vs PowerSoil.HMP
49.8040986	2.84923277	0.62918338	4.52846159	5.94E-06	0.00084963	Actinomycetaceae	InnuPURE vs PowerSoil.HMP
141.200238	-1.3748839	0.3732346	-3.6836989	0.00022987	0.00087939	Chitinophagaceae	EasyDNA vs MagNAPure
20.5932418	2.60659271	0.70858012	3.67861394	0.0002345	0.00087939	[Mogibacteriaceae]	EasyDNA vs MagNAPure
283.625111	3.28944084	0.74068442	4.44108277	8.95E-06	0.0009264	Lactobacillaceae	InnuPURE vs MagNAPure
21.4381793	3.18333391	0.83254469	3.82361927	0.00013151	0.00093029	Turicibacteraceae	EasyDNA vs QIAStool
129.664767	-1.5315971	0.40361056	-3.79474	0.0001478	0.00100644	Sphingomonadaceae	EasyDNA vs FastDNA
131.739477	-1.2097072	0.30846328	-3.921722	8.79E-05	0.00107054	Hyphomicrobiaceae	QIAStool vs MagNAPure
736.495903	-2.5211072	0.67768495	-3.7201758	0.00019908	0.00115895	Enterobacteriaceae	EasyDNA vs PowerSoil.HMP
105.619852	-1.7219182	0.47856033	-3.5981215	0.00032052	0.00116948	Nitrosomonadaceae	EasyDNA vs MagNAPure
118.159531	-1.6800207	0.44427482	-3.7814899	0.00015589	0.00117699	Rhodobacteraceae	EasyDNA vs QIAStool+BB
49.8040986	2.01409717	0.50832968	3.96218684	7.43E-05	0.00117765	Actinomycetaceae	FastDNA vs MagNAPure
616.023371	1.37539476	0.37422072	3.67535705	0.00023752	0.00133501	Veillonellaceae	EasyDNA vs PowerSoil.HMP
59.9667273	-1.7905328	0.4630503	-3.8668214	0.00011026	0.00153266	Phyllobacteriaceae	QIAStool vs QIAStool+BB
4421.89202	1.4148799	0.38994132	3.62844316	0.00028514	0.00154924	Campylobacteraceae	EasyDNA vs PowerSoil.HMP
7.76499835	3.00619582	0.78949885	3.80772665	0.00014025	0.00161288	Victivallaceae	QIAStool vs MagNAPure
858.112951	-1.103653	0.29099694	-3.7926618	0.00014904	0.00168722	Legionellaceae	PowerSoil.HMP vs QIAStool
118.339928	-1.1641985	0.30830147	-3.7761692	0.00015926	0.00168722	[Chthoniobacteraceae]	PowerSoil.HMP vs QIAStool
224.417038	1.54686811	0.40966719	3.77591408	0.00015942	0.00168722	Caulobacteraceae	PowerSoil.HMP vs QIAStool
31.145957	3.18626251	0.86955576	3.66424172	0.00024807	0.00169221	Aerococcaceae	EasyDNA vs QIAStool
264.350163	1.08538354	0.29166878	3.72128808	0.00019821	0.00188959	Leptotrichiaceae	FastDNA vs QIAStool
91.6048239	-1.1722866	0.3237836	-3.6205868	0.00029394	0.00191058	Halomonadaceae	EasyDNA vs FastDNA
7.94601326	3.25719295	0.87981198	3.70214661	0.00021378	0.00191069	Micrococcaceae	FastDNA vs QIAStool
15.0781299	2.03353532	0.58767751	3.46029119	0.00053959	0.00191697	[Odoribacteraceae]	EasyDNA vs MagNAPure
38.1375375	1.96739339	0.52466333	3.74982069	0.00017696	0.00192794	Procabacteriaceae	QIAStool vs MagNAPure
12.0832988	2.60554261	0.71947574	3.62144612	0.00029296	0.0019295	Microthrixaceae	EasyDNA vs QIAStool
269.467613	3.28592868	0.91384685	3.59571046	0.00032351	0.00201137	[Paraprevotellaceae]	EasyDNA vs FastDNA
59.9667273	-1.5618619	0.41967538	-3.7215952	0.00019797	0.00204897	Phyllobacteriaceae	QIAStool vs MagNAPure
92.6279814	1.60546842	0.44860355	3.57881343	0.00034516	0.00205114	Oxalobacteraceae	EasyDNA vs FastDNA
60.913589	-1.2906483	0.36164555	-3.5688212	0.00035859	0.00205114	SB-1	EasyDNA vs FastDNA
8.22875632	-3.0323745	0.85554848	-3.5443632	0.00039356	0.00206938	Solibacteraceae	EasyDNA vs PowerSoil.HMP
458.885072	2.1290759	0.52006231	4.09388616	4.24E-05	0.00207036	Carnobacteriaceae	InnuPURE vs PowerSoil.HMP
649.223144	1.77916883	0.43517387	4.08840914	4.34E-05	0.00207036	Bifidobacteriaceae	InnuPURE vs PowerSoil.HMP
118.159531	-1.4258087	0.38539195	-3.6996327	0.00021591	0.00212827	Rhodobacteraceae	QIAStool vs MagNAPure

3.24919787	3.6188219	1.00940315	3.58511056	0.00033694	0.00214516	Leuconostocaceae	EasyDNA vs QIAStool
33.4444509	2.98208837	0.80192515	3.71866173	0.00020028	0.00217639	Coriobacteriaceae	EasyDNA vs InnuPURE
1110.97909	-0.706397	0.19141459	-3.6904028	0.0002239	0.00218732	Verrucomicrobiaceae	PowerSoil.HMP vs QIAStool
858.112951	1.1933291	0.31822856	3.74991205	0.0001769	0.00223533	Legionellaceae	QIAStool vs QIAStool+BB
1181.58837	3.09655277	0.88052293	3.51672018	0.00043691	0.00240303	Prevotellaceae	EasyDNA vs FastDNA
28.7248963	-2.008188	0.59242969	-3.389749	0.00069957	0.00242158	Acetobacteraceae	EasyDNA vs MagNAPure
19.2945448	-1.9141579	0.56688273	-3.3766382	0.00073378	0.00246687	BA008	EasyDNA vs MagNAPure
92.6279814	1.52494408	0.4523832	3.37091228	0.0007492	0.00246687	Oxalobacteraceae	EasyDNA vs MagNAPure
169.692077	2.05994287	0.51874697	3.97099742	7.16E-05	0.00255871	Pirellulaceae	InnuPURE vs PowerSoil.HMP
354.810526	-0.98635	0.27628461	-3.5700506	0.00035691	0.00256637	Pseudomonadaceae	EasyDNA vs QIAStool+BB
129.664767	-1.3164872	0.3730486	-3.5289965	0.00041714	0.00257011	Sphingomonadaceae	EasyDNA vs QIAStool
4.87640195	3.65458882	1.09184921	3.34715525	0.00081645	0.00262432	Corynebacteriaceae	EasyDNA vs MagNAPure
22.1599751	-3.0574814	0.83143843	-3.6773395	0.00023568	0.00272995	Sanguibacteraceae	QIAStool vs QIAStool+BB
44.0121162	2.3469519	0.63926998	3.67130004	0.00024132	0.00279529	Microbacteriaceae	InnuPURE vs QIAStool
2.31382502	3.84858947	1.07433352	3.5823042	0.00034058	0.00286485	Nocardiaceae	FastDNA vs QIAStool
458.885072	2.10787358	0.51588	4.08597657	4.39E-05	0.00302853	Carnobacteriaceae	InnuPURE vs MagNAPure
17.3475351	-2.7947582	0.75672129	-3.6932464	0.00022141	0.00303647	Isosphaeraceae	PowerSoil.HMP vs MagNAPure
47.3908296	1.37466258	0.40163928	3.42262981	0.00062018	0.00328468	Christensenellaceae	EasyDNA vs FastDNA
169.692077	-1.7496979	0.46383396	-3.7722504	0.00016178	0.0033327	Pirellulaceae	QIAStool+BB vs MagNAPure
113.494552	1.2476785	0.34930574	3.57188087	0.00035443	0.00333483	Alcaligenaceae	QIAStool vs MagNAPure
113.494552	1.20497493	0.35380394	3.40577026	0.00065978	0.00336957	Alcaligenaceae	EasyDNA vs FastDNA
12.2684488	2.29895523	0.64642084	3.55643735	0.00037592	0.00338326	Synergistaceae	QIAStool vs MagNAPure
28.7248963	-1.9922099	0.56229317	-3.5430093	0.00039559	0.00351634	Acetobacteraceae	FastDNA vs PowerSoil.HMP
2750.67146	1.695233	0.46819787	3.6207619	0.00029374	0.00352484	Bacteroidaceae	PowerSoil.HMP vs MagNAPure
118.159531	-1.7715548	0.49520976	-3.5773826	0.00034705	0.00353559	Rhodobacteraceae	EasyDNA vs InnuPURE
92.6279814	-1.6411801	0.46414377	-3.5359305	0.00040634	0.00354978	Oxalobacteraceae	PowerSoil.HMP vs QIAStool
649.223144	1.28566191	0.36445274	3.52765052	0.00041927	0.00354978	Bifidobacteriaceae	PowerSoil.HMP vs QIAStool
44.0121162	-2.1137016	0.58933766	-3.5865715	0.00033505	0.00357391	Microbacteriaceae	PowerSoil.HMP vs MagNAPure
354.810526	-0.8207072	0.25232262	-3.2526104	0.0011435	0.00359006	Pseudomonadaceae	EasyDNA vs MagNAPure
12.0832988	-2.8162903	0.78971557	-3.5662084	0.00036218	0.00387258	Microthrixaceae	QIAStool vs QIAStool+BB
169.692077	1.41344996	0.42000085	3.36535026	0.00076447	0.003894	Pirellulaceae	EasyDNA vs PowerSoil.HMP
1110.97909	-0.8492203	0.23853992	-3.5600764	0.00037075	0.00396414	Verrucomicrobiaceae	InnuPURE vs QIAStool
118.159531	-1.5107086	0.41671211	-3.6253053	0.00028862	0.00400461	Rhodobacteraceae	FastDNA vs MagNAPure
639.445335	1.16043718	0.33390826	3.47531735	0.00051025	0.00403097	Marinilabiaceae	FastDNA vs QIAStool
993.963868	-1.7178466	0.49615794	-3.4622979	0.00053558	0.00403097	Opituaceae	FastDNA vs QIAStool
59.9667273	-1.4629208	0.45541119	-3.2123076	0.00131673	0.00403998	Phyllobacteriaceae	EasyDNA vs MagNAPure
7.4067071	-2.8464888	0.85314609	-3.3364612	0.00084852	0.00419119	Desulfobacteraceae	EasyDNA vs PowerSoil.HMP
719.094782	-1.3002032	0.36982603	-3.5157159	0.00043857	0.00421027	Streptococcaceae	PowerSoil.HMP vs MagNAPure
118.159531	-1.4733244	0.41877181	-3.5182035	0.00043448	0.00431376	Rhodobacteraceae	QIAStool vs QIAStool+BB
593.953594	-0.7654102	0.23067053	-3.3181968	0.00090601	0.0043435	Cytophagaceae	EasyDNA vs PowerSoil.HMP
59.9667273	-1.6915917	0.49533902	-3.4150181	0.00063778	0.00437747	Phyllobacteriaceae	EasyDNA vs QIAStool+BB
70.3986464	1.2061253	0.36301817	3.32249295	0.00089217	0.00439932	Desulfovibrionaceae	EasyDNA vs FastDNA
269.467613	3.16345108	0.93031323	3.40041501	0.00067284	0.00441732	[Paraprevotellaceae]	EasyDNA vs QIAStool+BB
1181.58837	2.792938	0.8806047	3.17161376	0.00151594	0.00454783	Prevotellaceae	EasyDNA vs MagNAPure
182.91656	-1.3835089	0.37942483	-3.6463319	0.00026601	0.00456651	Planctomycetaceae	QIAStool+BB vs MagNAPure
4421.89202	1.229723	0.38951484	3.15706329	0.00159367	0.00467707	Campylobacteraceae	EasyDNA vs MagNAPure
19.2945448	-2.0575467	0.61259008	-3.3587659	0.00078291	0.00476931	BA008	EasyDNA vs QIAStool+BB
92.6279814	1.70551314	0.50813645	3.35640779	0.00078962	0.00476931	Oxalobacteraceae	EasyDNA vs QIAStool+BB
15.0781299	2.00704774	0.61163899	3.28142545	0.00103284	0.00481008	[Odoribacteraceae]	EasyDNA vs PowerSoil.HMP
254.528648	-1.3406432	0.42677372	-3.1413445	0.00168174	0.00483053	Xanthomonadaceae	EasyDNA vs MagNAPure
129.664767	-1.4706229	0.44044394	-3.3389559	0.00084094	0.00488391	Sphingomonadaceae	EasyDNA vs QIAStool+BB
60.913589	1.45888204	0.41924116	3.4798159	0.00050176	0.00498174	SB-1	InnuPURE vs QIAStool
11.2601759	2.86575309	0.92393381	3.10168657	0.00192422	0.00541186	Peptococcaceae	EasyDNA vs MagNAPure
458.885072	-1.8935274	0.45074517	-4.2008822	2.66E-05	0.00550365	Carnobacteriaceae	PowerSoil.HMP vs QIAStool+BB
736.495903	-3.0921527	0.82941526	-3.7281117	0.00019292	0.00551751	Enterobacteriaceae	InnuPURE vs PowerSoil.HMP
129.664767	-1.3324584	0.41218724	-3.2326533	0.00122646	0.00555315	Sphingomonadaceae	EasyDNA vs PowerSoil.HMP
118.159531	-1.3509387	0.4195755	-3.2197749	0.00128291	0.00565175	Rhodobacteraceae	EasyDNA vs PowerSoil.HMP
11.2601759	3.04335101	0.94850206	3.20858662	0.00133389	0.00572169	Peptococcaceae	EasyDNA vs PowerSoil.HMP
616.023371	1.33864065	0.39319285	3.40453967	0.00066276	0.00607517	Veillonellaceae	QIAStool vs QIAStool+BB
103.008126	-1.3554459	0.40137289	-3.377024	0.00073275	0.00607517	Enterococcaceae	QIAStool vs QIAStool+BB
24.9235631	-1.717352	0.51140949	-3.3580762	0.00078487	0.00607517	[Bryobacteraceae]	QIAStool vs QIAStool+BB
60.913589	-1.3029449	0.38807828	-3.3574279	0.00078671	0.00607517	SB-1	QIAStool vs QIAStool+BB
22.5747955	1.6881564	0.55662937	3.0328195	0.0024228	0.00667507	Peptostreptococcaceae	EasyDNA vs MagNAPure
7.4067071	-2.6173944	0.80329827	-3.2583096	0.00112078	0.00668966	Desulfobacteraceae	EasyDNA vs QIAStool
12.2684488	2.11983061	0.66428024	3.19116918	0.00141698	0.00675428	Synergistaceae	EasyDNA vs FastDNA

104.663517	1.42534693	0.4244879	3.35780346	0.00078564	0.00685653	Rikenellaceae	PowerSoil.HMP vs MagNAPure
20.5932418	2.78844857	0.82723718	3.37079697	0.00074951	0.00718649	[Mogibacteriaceae]	EasyDNA vs InnuPURE
993.963868	-2.017559	0.60189132	-3.3520321	0.00080221	0.00743379	Opitutaceae	InnuPURE vs QIAStool
186.231671	-0.9108244	0.30550845	-2.9813394	0.0028699	0.00768192	Rhizobiaceae	EasyDNA vs MagNAPure
1110.97909	-0.5821481	0.19548775	-2.9779262	0.00290206	0.00768192	Verrucomicrobiaceae	EasyDNA vs MagNAPure
59.9667273	-1.5324296	0.44867974	-3.4154197	0.00063684	0.00785434	Phyllobacteriaceae	FastDNA vs MagNAPure
579.211848	-0.8687005	0.27121003	-3.2030546	0.00135978	0.00787025	Cryomorphaceae	EasyDNA vs QIAStool
103.008126	1.06883487	0.34143186	3.13044852	0.0017454	0.00805134	Enterococcaceae	EasyDNA vs FastDNA
118.159531	1.5648585	0.47276645	3.31000327	0.00093295	0.00810499	Rhodobacteraceae	InnuPURE vs QIAStool
8.22875632	2.50314686	0.76230268	3.28366528	0.00102467	0.00813328	Solibacteraceae	PowerSoil.HMP vs QIAStool
20.5932418	2.47178571	0.77677805	3.18210037	0.00146211	0.00817699	[Mogibacteriaceae]	EasyDNA vs QIAStool+BB
49.8040986	2.25915369	0.59809529	3.7772471	0.00015857	0.00820607	Actinomycetaceae	InnuPURE vs MagNAPure
103.008126	1.68544012	0.47255029	3.56668943	0.00036152	0.00858467	Enterococcaceae	InnuPURE vs PowerSoil.HMP
622.632062	1.49623869	0.42421902	3.52704296	0.00042023	0.00858467	Flavobacteriaceae	InnuPURE vs PowerSoil.HMP
6.92597984	-2.9939275	0.90591845	-3.3048533	0.00095026	0.00860514	[Chromatiaceae]	EasyDNA vs InnuPURE
7.94601326	2.36493345	0.80553543	2.93585279	0.00332632	0.00863564	Micrococcaceae	EasyDNA vs MagNAPure
28.0267224	2.247816	0.69396685	3.23908267	0.00119915	0.00959318	Succinivibrionaceae	PowerSoil.HMP vs MagNAPure
24.9235631	-1.4990779	0.45978042	-3.2604213	0.00111247	0.00959504	[Bryobacteraceae]	QIAStool vs MagNAPure
4421.89202	1.22665838	0.39306924	3.12071835	0.0018041	0.00972928	Campylobacteraceae	EasyDNA vs QIAStool+BB
736.495903	2.16738284	0.67437254	3.21392511	0.00130934	0.00978153	Enterobacteriaceae	PowerSoil.HMP vs QIAStool
19.2945448	1.55104302	0.48513414	3.19714258	0.00138796	0.00979285	BA008	PowerSoil.HMP vs QIAStool
269.467613	2.67877514	0.85846322	3.12043087	0.00180587	0.01014472	[Paraprevotellaceae]	EasyDNA vs QIAStool
254.528648	-1.2938909	0.42934953	-3.0136073	0.00258162	0.01078984	Xanthomonadaceae	EasyDNA vs PowerSoil.HMP
4421.89202	-1.2253129	0.38948425	-3.1459884	0.00165527	0.01106414	Campylobacteraceae	PowerSoil.HMP vs QIAStool
60.913589	-1.04435	0.3664465	-2.8499385	0.00437277	0.01113818	SB-1	EasyDNA vs MagNAPure
4234.46977	-1.025778	0.3242894	-3.163156	0.00156069	0.01115891	Ruminococcaceae	FastDNA vs QIAStool
17.3475351	-2.6763124	0.799256	-3.3485046	0.00081249	0.01146109	Isosphaeraceae	QIAStool+BB vs MagNAPure
622.632062	-1.2115734	0.36458966	-3.3231151	0.00089018	0.01146109	Flavobacteriaceae	QIAStool+BB vs MagNAPure
264.350163	0.85382433	0.2783669	3.0672624	0.00216029	0.01178902	Leptotrichiaceae	EasyDNA vs QIAStool
12.2684488	1.94066769	0.68710829	2.82439859	0.00473694	0.01184236	Synergistaceae	EasyDNA vs MagNAPure
993.963868	-1.503144	0.50502245	-2.9763905	0.00291663	0.01186219	Opitutaceae	EasyDNA vs PowerSoil.HMP
131.739477	-1.0249579	0.34517249	-2.969408	0.00298374	0.01186219	Hyphomicrobiaceae	EasyDNA vs PowerSoil.HMP
4.87640195	3.28815114	1.07742791	3.05185257	0.00227434	0.01206663	Corynebacteriaceae	EasyDNA vs QIAStool
31.145957	2.54682017	0.91162549	2.79371319	0.00521067	0.01278982	Aerococcaceae	EasyDNA vs MagNAPure
1110.97909	-0.7305016	0.22421632	-3.2580214	0.00112192	0.01283975	Verrucomicrobiaceae	QIAStool+BB vs MagNAPure
113.494552	1.40369726	0.44319404	3.16722954	0.00153899	0.0132029	Alcaligenaceae	EasyDNA vs InnuPURE
60.913589	1.09927041	0.35735416	3.07613715	0.00209701	0.01331604	SB-1	PowerSoil.HMP vs QIAStool
4.06716246	2.70023017	0.85880434	3.14417387	0.00166556	0.01379087	Elusimicrobiaceae	QIAStool vs MagNAPure
49.8040986	-1.671613	0.53708995	-3.1123519	0.00185603	0.0139388	Actinomycetaceae	QIAStool vs MagNAPure
224.417038	-1.2657762	0.40782434	-3.103729	0.00191098	0.0139388	Caulobacteraceae	QIAStool vs MagNAPure
182.91656	-0.9604176	0.30999369	-3.0981843	0.0019471	0.0139388	Planctomycetaceae	QIAStool vs MagNAPure
858.112951	-0.8756507	0.28271224	-3.0973216	0.00195278	0.0139388	Legionellaceae	QIAStool vs MagNAPure
4830.98886	0.90318919	0.30567964	2.954692	0.00312981	0.01398635	Moraxellaceae	EasyDNA vs FastDNA
127.559639	-1.1983362	0.38295763	-3.1291612	0.00175306	0.01402449	Bradyrhizobiaceae	FastDNA vs PowerSoil.HMP
4.81963649	-2.5920036	0.86507369	-2.9962807	0.00273295	0.01410792	auto67_4W	EasyDNA vs QIAStool
3.21882027	-2.9945197	1.02979292	-2.9078853	0.00363882	0.01412208	Geobacteraceae	EasyDNA vs PowerSoil.HMP
103.008126	1.32628602	0.42621318	3.11179028	0.00185957	0.01422317	Enterococcaceae	EasyDNA vs InnuPURE
269.467613	3.07998123	0.99275227	3.10246708	0.00191915	0.01422317	[Paraprevotellaceae]	EasyDNA vs InnuPURE
28.7248963	-2.1158352	0.68200314	-3.1023834	0.00191969	0.01422317	Acetobacteraceae	EasyDNA vs InnuPURE
12.0832988	2.60257042	0.83588608	3.11354678	0.00184853	0.01511447	Microthrixaceae	InnuPURE vs QIAStool
15.0781299	1.90584145	0.64043532	2.9758531	0.00292175	0.01521324	[Odoribacteraceae]	EasyDNA vs QIAStool+BB
7.94601326	-2.9156223	0.9471339	-3.0783634	0.00208141	0.01522715	Micrococcaceae	QIAStool vs QIAStool+BB
118.339928	-1.0389997	0.34156999	-3.0418353	0.00235141	0.01531368	[Chthoniobacteraceae]	FastDNA vs PowerSoil.HMP
4234.46977	-1.0690832	0.35423431	-3.0180115	0.00254439	0.01531368	Ruminococcaceae	FastDNA vs PowerSoil.HMP
719.094782	1.11333967	0.36942897	3.01367723	0.00258102	0.01531368	Streptococcaceae	FastDNA vs PowerSoil.HMP
169.692077	1.31061062	0.43885856	2.98640779	0.00282276	0.01531368	Pirellulaceae	FastDNA vs PowerSoil.HMP
283.625111	1.88675667	0.63288731	2.98118896	0.00287132	0.01531368	Lactobacillaceae	FastDNA vs PowerSoil.HMP
91.6048239	-0.9688326	0.3376377	-2.8694444	0.00411194	0.0153584	Halomonadaceae	EasyDNA vs PowerSoil.HMP
269.467613	2.62362002	0.91515855	2.86684753	0.00414583	0.0153584	[Paraprevotellaceae]	EasyDNA vs PowerSoil.HMP
169.692077	-1.0885501	0.39949647	-2.7248053	0.00643394	0.01551039	Pirellulaceae	EasyDNA vs MagNAPure
118.339928	0.85515562	0.31545607	2.71085484	0.006711	0.01589447	[Chthoniobacteraceae]	EasyDNA vs MagNAPure
4.27355164	-2.8506251	0.93680851	-3.0429112	0.00234301	0.01628395	Eubacteriaceae	QIAStool vs QIAStool+BB
21.5427213	2.62816676	0.89319412	2.94243625	0.00325641	0.01636774	Spirochaetaceae	EasyDNA vs QIAStool
59.9667273	1.56849566	0.51094835	3.0697734	0.00214221	0.01654264	Phyllobacteriaceae	InnuPURE vs QIAStool
118.339928	1.19613294	0.39335639	3.04083768	0.00235921	0.01671962	[Chthoniobacteraceae]	EasyDNA vs InnuPURE

224.417038	-1.3016651	0.44367212	-2.9338446	0.00334792	0.01673959	Caulobacteraceae	FastDNA vs PowerSoil.HMP
993.963868	-1.499755	0.51224519	-2.9278069	0.00341362	0.01679999	Opitutaceae	EasyDNA vs QIAStool+BB
131.739477	-1.0909518	0.37302612	-2.9245989	0.003449	0.01679999	Hyphomicrobiaceae	EasyDNA vs QIAStool+BB
21.5427213	2.53561288	0.94480086	2.68375379	0.00728007	0.01690336	Spirochaetaceae	EasyDNA vs MagNAPure
333.941631	1.06355661	0.39701881	2.67885697	0.00738739	0.01690336	Porphyromonadaceae	EasyDNA vs MagNAPure
28.7345598	-1.6199105	0.55646094	-2.9110947	0.00360165	0.01699528	Parachlamydiaceae	EasyDNA vs QIAStool+BB
21.4381793	2.87852569	0.94506216	3.04585858	0.00232017	0.01713356	Turicibacteraceae	PowerSoil.HMP vs MagNAPure
118.159531	-1.2291424	0.42307595	-2.9052522	0.00366957	0.01726858	Rhodobacteraceae	FastDNA vs PowerSoil.HMP
104.663517	1.26962895	0.43903252	2.89187904	0.00382945	0.01732243	Rikenellaceae	EasyDNA vs QIAStool+BB
21.5427213	2.85758764	0.99453991	2.87327597	0.00406239	0.01732243	Spirochaetaceae	EasyDNA vs QIAStool+BB
1181.58837	2.53355579	0.88411161	2.86565152	0.00416152	0.01732243	Prevotellaceae	EasyDNA vs QIAStool+BB
355.031722	1.13809798	0.39794486	2.85993889	0.00423723	0.01732243	[Weeksellaceae]	EasyDNA vs QIAStool+BB
113.494552	1.15165021	0.40396852	2.85084148	0.00436037	0.01732243	Alcaligenaceae	EasyDNA vs QIAStool+BB
28.7248963	-1.8378156	0.64637612	-2.8432603	0.00446546	0.01732243	Acetobacteraceae	EasyDNA vs QIAStool+BB
118.339928	1.02243338	0.35967603	2.84265084	0.00447401	0.01732243	[Chthoniobacteraceae]	EasyDNA vs QIAStool+BB
7.4067071	-2.533754	0.89397879	-2.834244	0.00459342	0.01734018	Desulfobacteraceae	EasyDNA vs QIAStool+BB
4.81963649	-2.5721622	0.89514869	-2.8734468	0.00406019	0.01759417	auto67_4W	EasyDNA vs FastDNA
38.1375375	-1.588819	0.53677015	-2.9599616	0.00307677	0.0176333	Procabacteriaceae	PowerSoil.HMP vs QIAStool
993.963868	-1.4686531	0.49715551	-2.9541121	0.0031357	0.0176333	Opitutaceae	PowerSoil.HMP vs QIAStool
377.356263	0.66481791	0.22547835	2.94847773	0.00319343	0.0176333	Sphingobacteriaceae	PowerSoil.HMP vs QIAStool
70.3986464	1.07418554	0.3564197	3.01382203	0.00257979	0.01780056	Desulfovibrionaceae	QIAStool vs MagNAPure
4830.98886	0.86879822	0.30910597	2.8106808	0.00494368	0.01820721	Moraxellaceae	EasyDNA vs QIAStool+BB
622.632062	1.20011256	0.39698082	3.02309962	0.002502	0.0183041	Flavobacteriaceae	InnuPURE vs QIAStool
22.1599751	2.34056949	0.78040891	2.99915784	0.00270727	0.01833074	Sanguibacteraceae	FastDNA vs QIAStool
579.211848	0.84548558	0.2831535	2.98596192	0.00282688	0.01833074	Cryomorphaceae	FastDNA vs QIAStool
310.195483	-1.1140138	0.37470056	-2.9730774	0.0029483	0.01833074	Aeromonadaceae	FastDNA vs QIAStool
21.5427213	2.67648334	0.94093314	2.84449896	0.00444813	0.01841406	Spirochaetaceae	EasyDNA vs FastDNA
3.24919787	2.81122058	0.98975752	2.84031241	0.00450694	0.01841406	Leuconostocaceae	EasyDNA vs FastDNA
118.159531	1.14424238	0.39230985	2.91668023	0.00353778	0.01872077	Rhodobacteraceae	PowerSoil.HMP vs QIAStool
1181.58837	2.37174212	0.82321907	2.88105827	0.00396342	0.01884447	Prevotellaceae	EasyDNA vs QIAStool
4830.98886	0.79076118	0.27480994	2.87748395	0.0040086	0.01884447	Moraxellaceae	EasyDNA vs QIAStool
4.27355164	2.54518735	0.8853998	2.87461931	0.00404515	0.01884447	Eubacteriaceae	EasyDNA vs QIAStool
113.494552	1.15379451	0.38639387	2.98605802	0.00282599	0.01937822	Alcaligenaceae	PowerSoil.HMP vs MagNAPure
92.6279814	1.64386434	0.5519235	2.97842786	0.00289731	0.01967758	Oxalobacteraceae	EasyDNA vs InnuPURE
18.7347412	1.74564306	0.58736763	2.97197698	0.00295889	0.01975774	[Barnesiellaceae]	QIAStool vs MagNAPure
24.9235631	-1.9126891	0.64531773	-2.9639494	0.00303718	0.01980243	[Bryobacteraceae]	EasyDNA vs InnuPURE
243.261282	-1.0090776	0.33828228	-2.9829456	0.00285489	0.01984147	GZKB119	InnuPURE vs QIAStool
12.0832988	1.9612567	0.7511158	2.61112429	0.00902451	0.02030515	Microthrixaceae	EasyDNA vs MagNAPure
2750.67146	1.28916277	0.43584769	2.95782861	0.00309814	0.02050676	Bacteroidaceae	QIAStool vs QIAStool+BB
243.261282	-0.7295655	0.25759568	-2.8322118	0.00462272	0.02102238	GZKB119	EasyDNA vs QIAStool
79.5100271	-1.191318	0.43277484	-2.7527432	0.00590982	0.0214067	Bdellovibrionaceae	EasyDNA vs PowerSoil.HMP
19.2945448	-1.5596526	0.56244073	-2.7730079	0.00555408	0.02193946	BA008	EasyDNA vs FastDNA
11.2601759	2.50254376	0.90615756	2.76170929	0.00574997	0.02193946	Peptococcaceae	EasyDNA vs FastDNA
593.953594	-0.62696	0.22739104	-2.7571889	0.00583007	0.02193946	Cytophagaceae	EasyDNA vs FastDNA
10.9576795	-2.0973714	0.76485713	-2.7421741	0.0061034	0.02194317	Acholeplasmataceae	EasyDNA vs QIAStool+BB
593.953594	-0.5824492	0.20721534	-2.8108403	0.00494123	0.02194826	Cytophagaceae	EasyDNA vs QIAStool
993.963868	1.47204213	0.50450751	2.91778038	0.00352533	0.02227365	Opitutaceae	QIAStool vs QIAStool+BB
10.9576795	-1.8987571	0.68029491	-2.7910793	0.00525326	0.02280392	Acholeplasmataceae	EasyDNA vs QIAStool
7.4067071	-2.284038	0.8363462	-2.7309719	0.00631478	0.02315421	Desulfobacteraceae	EasyDNA vs FastDNA
1110.97909	-0.5394896	0.18836632	-2.8640448	0.00418269	0.02338949	Verrucomicrobiaceae	FastDNA vs QIAStool
377.356263	0.62991124	0.22044344	2.85747327	0.00427029	0.02338949	Sphingobacteriaceae	FastDNA vs QIAStool
24.9235631	1.29672979	0.45420932	2.85491676	0.00430481	0.02338949	[Bryobacteraceae]	FastDNA vs QIAStool
31.145957	2.57510178	0.90456212	2.84679374	0.0044162	0.02338949	Aerococcaceae	FastDNA vs QIAStool
12.9936921	-1.8506044	0.72523789	-2.5517205	0.01071925	0.02372292	Hyphomonadaceae	EasyDNA vs MagNAPure
355.031722	0.99703315	0.36955369	2.69793854	0.00697703	0.02431227	[Weeksellaceae]	EasyDNA vs PowerSoil.HMP
264.350163	-0.8216819	0.30473801	-2.6963553	0.00701029	0.02431227	Leptotrichiaceae	EasyDNA vs PowerSoil.HMP
2750.67146	-1.2094344	0.42892542	-2.8196845	0.00480709	0.02455049	Bacteroidaceae	FastDNA vs QIAStool
283.625111	1.74651589	0.61997486	2.81707535	0.00484632	0.02461928	Lactobacillaceae	PowerSoil.HMP vs QIAStool
79.5100271	-1.2567499	0.46625307	-2.6954243	0.00702991	0.02468641	Bdellovibrionaceae	EasyDNA vs QIAStool+BB
3668.6678	1.63743174	0.54128315	3.02509274	0.00248557	0.02560136	Lachnospiraceae	QIAStool+BB vs MagNAPure
44.0121162	1.43218661	0.52197913	2.74376222	0.00607395	0.02578055	Microbacteriaceae	EasyDNA vs QIAStool
1756.14884	-0.8875915	0.35309787	-2.5137266	0.0119463	0.02592108	Neisseriaceae	EasyDNA vs MagNAPure
18.7347412	1.55956336	0.62150923	2.50931647	0.01209651	0.02592108	[Barnesiellaceae]	EasyDNA vs MagNAPure
2750.67146	-1.5115334	0.52589418	-2.8742159	0.00405032	0.02611934	Bacteroidaceae	InnuPURE vs QIAStool
4.27355164	2.76832398	0.97008355	2.85369643	0.00432138	0.02611934	Eubacteriaceae	InnuPURE vs QIAStool

17.3475351	2.27655242	0.79776644	2.85365781	0.00432191	0.02611934	Isosphaeraceae	InnuPURE vs QIAStool
38.1375375	-1.4159276	0.50824048	-2.7859402	0.00533727	0.02631827	Procabacteriaceae	FastDNA vs QIAStool
2.3093169	-2.8715958	1.08020362	-2.6583838	0.00785164	0.02694541	Myxococcaceae	EasyDNA vs QIAStool+BB
2.31382502	3.1273455	1.10587857	2.82792847	0.00468503	0.02713412	Nocardiaceae	InnuPURE vs QIAStool
254.528648	-1.1861816	0.44926195	-2.6402895	0.00828352	0.02779582	Xanthomonadaceae	EasyDNA vs QIAStool+BB
8.22875632	-2.3772037	0.9049355	-2.6269316	0.00861586	0.02828251	Solibacteraceae	EasyDNA vs QIAStool+BB
4.06716246	-2.6505323	0.9462686	-2.8010358	0.00509389	0.02832201	Elusimicrobiaceae	InnuPURE vs QIAStool
224.417038	-1.2142892	0.43070396	-2.8193127	0.00481266	0.02908521	Caulobacteraceae	QIAStool vs QIAStool+BB
8.97993102	-1.9720958	0.74525118	-2.6462163	0.00813978	0.0290997	Sediment-4	EasyDNA vs FastDNA
1.55959833	3.04990549	1.10011358	2.77235509	0.00556523	0.02914922	Dermatophilaceae	InnuPURE vs QIAStool
31.145957	2.73793794	0.98959174	2.76673484	0.00566208	0.02914922	Aerococcaceae	InnuPURE vs QIAStool
164.684355	2.00560381	0.59200326	3.38782561	0.00070449	0.0291659	Clostridiaceae	InnuPURE vs MagNAPure
21.4381793	2.4029683	0.87615781	2.74262042	0.00609511	0.02977227	Turicibacteraceae	PowerSoil.HMP vs QIAStool
22.1599751	-2.2288796	0.7867898	-2.8328781	0.0046131	0.02984097	Sanguibacteraceae	QIAStool vs MagNAPure
11.4855136	-1.8652956	0.66110538	-2.82148	0.00478026	0.02998528	Gemmatimonadaceae	QIAStool vs MagNAPure
7.76499835	2.08136474	0.84932062	2.4506231	0.01426092	0.03008163	Victivallaceae	EasyDNA vs MagNAPure
20.0435737	-1.4557469	0.56093764	-2.5952027	0.00945351	0.03037193	Saprospiraceae	EasyDNA vs QIAStool+BB
11.2601759	2.81611955	0.99962466	2.81717696	0.00484478	0.03037306	Peptococcaceae	EasyDNA vs InnuPURE
103.008126	1.51734838	0.45684108	3.32139215	0.0008957	0.03090151	Enterococcaceae	InnuPURE vs MagNAPure
639.445335	-0.9363952	0.33454193	-2.7990367	0.00512553	0.03114536	Marinilabiaceae	QIAStool vs MagNAPure
719.094782	-0.9343259	0.33484933	-2.7902876	0.00526612	0.03114536	Streptococcaceae	QIAStool vs MagNAPure
4.06716246	2.44686122	0.880977	2.77744052	0.00547889	0.03173189	Elusimicrobiaceae	QIAStool vs QIAStool+BB
858.112951	-1.2104768	0.37506448	-3.2273833	0.00124928	0.03314836	Legionellaceae	InnuPURE vs MagNAPure
1.55959833	3.53407623	1.09747746	3.22018115	0.0012811	0.03314836	Dermatophilaceae	InnuPURE vs MagNAPure
3.21882027	-2.7128444	1.06112987	-2.5565621	0.01057122	0.03325529	Geobacteraceae	EasyDNA vs QIAStool+BB
12.9936921	-1.8610075	0.71942811	-2.5867873	0.00968754	0.03378824	Hyphomonadaceae	EasyDNA vs FastDNA
49.8040986	1.33489132	0.52490846	2.54309356	0.01098758	0.03385969	Actinomycetaceae	EasyDNA vs QIAStool+BB
164.684355	1.20633872	0.46822066	2.57643208	0.00998258	0.03389918	Clostridiaceae	EasyDNA vs PowerSoil.HMP
2.61442095	-2.3742008	0.98904022	-2.4005099	0.01637225	0.0340039	Ellin515	EasyDNA vs MagNAPure
2750.67146	1.37968465	0.46764596	2.95027598	0.0031749	0.03433796	Bacteroidaceae	FastDNA vs MagNAPure
12.0832988	2.21601142	0.75663079	2.92878833	0.00340286	0.03433796	Microthrixaceae	FastDNA vs MagNAPure
2750.67146	1.02138586	0.38739727	2.63653347	0.00837579	0.03446271	Bacteroidaceae	EasyDNA vs QIAStool
2.90378317	-2.6542909	1.00834595	-2.6323217	0.00848035	0.03446271	Shewanellaceae	EasyDNA vs QIAStool
2.61442095	-2.5754714	0.93748593	-2.7472107	0.00601045	0.03456009	Ellin515	QIAStool vs MagNAPure
1.27584029	-3.0144798	1.10165276	-2.7363248	0.00621297	0.03475903	Methanobacteriaceae	QIAStool vs MagNAPure
4.18916425	2.38221387	0.99895023	2.38471727	0.01709225	0.03480958	Nocardioidaceae	EasyDNA vs MagNAPure
264.350163	-0.71875	0.30189642	-2.3807835	0.01727586	0.03480958	Leptotrichiaceae	EasyDNA vs MagNAPure
10.9576795	-2.197664	0.79886476	-2.7509837	0.00594166	0.03577349	Acholeplasmataceae	EasyDNA vs InnuPURE
1181.58837	2.64574182	0.96781367	2.73373057	0.00626213	0.03577349	Prevotellaceae	EasyDNA vs InnuPURE
59.9667273	-1.4695546	0.54001007	-2.7213466	0.00650165	0.03577349	Phyllobacteriaceae	EasyDNA vs InnuPURE
1.55959833	-2.9910789	1.10080263	-2.71718	0.00658408	0.03577349	Dermatophilaceae	EasyDNA vs InnuPURE
28.0267224	1.50756944	0.58964909	2.55672309	0.01056633	0.03597584	Succinivibrionaceae	EasyDNA vs FastDNA
616.023371	-0.9908895	0.37159274	-2.6666009	0.00766226	0.036041	Veillonellaceae	PowerSoil.HMP vs QIAStool
131.739477	-0.8564683	0.33629328	-2.5467899	0.01087189	0.03615534	Hyphomicrobiaceae	EasyDNA vs FastDNA
186.231671	0.74799507	0.28674278	2.60859252	0.00909154	0.03617677	Rhizobiaceae	EasyDNA vs QIAStool
118.339928	-0.9548824	0.33231545	-2.8734217	0.00406052	0.03636071	[Chthoniobacteraceae]	FastDNA vs MagNAPure
169.692077	-1.1913895	0.4193644	-2.8409409	0.00449807	0.03636071	Pirellulaceae	FastDNA vs MagNAPure
182.91656	-0.9532472	0.33627107	-2.8347584	0.00458604	0.03636071	Planctomycetaceae	FastDNA vs MagNAPure
91.6048239	-0.7763827	0.32992933	-2.3531788	0.01861368	0.03695363	Halomonadaceae	EasyDNA vs MagNAPure
22.1599751	1.97579153	0.77845518	2.53809287	0.01114584	0.03707698	Sanguibacteraceae	EasyDNA vs PowerSoil.HMP
28.7345598	-1.2703451	0.50442951	-2.5183798	0.01178961	0.03804106	Parachlamydiaceae	EasyDNA vs FastDNA
254.528648	-1.068488	0.42623526	-2.5068033	0.01218285	0.03804106	Xanthomonadaceae	EasyDNA vs FastDNA
6.92597984	-2.0759884	0.82940286	-2.5029916	0.01231485	0.03804106	[Chromatiaceae]	EasyDNA vs FastDNA
4421.89202	0.97253378	0.38938417	2.49762026	0.012503	0.03804106	Campylobacteraceae	EasyDNA vs FastDNA
993.963868	-1.2539505	0.50404268	-2.4877864	0.01285409	0.03829448	Opitutaceae	EasyDNA vs FastDNA
104.663517	1.15256553	0.42682061	2.70035119	0.00692663	0.03851207	Rikenellaceae	QIAStool vs QIAStool+BB
642.395629	-0.8728261	0.37490681	-2.3281148	0.01990601	0.03894653	Rhodocyclaceae	EasyDNA vs MagNAPure
6.92597984	-1.9641006	0.84680666	-2.3194203	0.02037226	0.03928935	[Chromatiaceae]	EasyDNA vs MagNAPure
3.5697885	-2.663582	0.99480591	-2.6774891	0.00741763	0.03965577	Beutenbergiaceae	QIAStool vs QIAStool+BB
8.22875632	-2.0849652	0.8464636	-2.4631481	0.0137723	0.04019264	Solibacteraceae	EasyDNA vs FastDNA
141.200238	0.87472846	0.33424873	2.61699863	0.00887067	0.04023482	Chitinophagaceae	PowerSoil.HMP vs QIAStool
4421.89202	1.04015603	0.38905723	2.67352962	0.00750576	0.04080907	Campylobacteraceae	QIAStool vs MagNAPure
4.27355164	-2.3695014	0.8889705	-2.6654444	0.00768866	0.04080907	Eubacteriaceae	QIAStool vs MagNAPure
28.0267224	1.91589962	0.72230348	2.65248565	0.00799015	0.04095456	Succinivibrionaceae	EasyDNA vs InnuPURE
639.445335	-1.1176738	0.42467418	-2.6318384	0.00849243	0.04095456	Marinilabiaceae	EasyDNA vs InnuPURE

224.417038	-1.3704338	0.52104201	-2.6301791	0.00853399	0.04095456	Caulobacteraceae	EasyDNA vs InnuPURE
104.663517	1.25209331	0.47716645	2.62401793	0.00868992	0.04095456	Rikenellaceae	EasyDNA vs InnuPURE
4830.98886	0.9966009	0.38038735	2.61996329	0.00879392	0.04095456	Moraxellaceae	EasyDNA vs InnuPURE
21.5427213	2.3806915	0.95400598	2.49546812	0.01257911	0.04100788	Spirochaetaceae	EasyDNA vs PowerSoil.HMP
4.18916425	2.4527686	0.95878924	2.5581937	0.01052175	0.04101334	Nocardioidaceae	EasyDNA vs QIAStool
3.21882027	-2.354502	1.02533773	-2.2963185	0.02165768	0.04118009	Geobacteraceae	EasyDNA vs MagNAPure
3668.6678	1.46196977	0.53675835	2.72370194	0.00645547	0.04131503	Lachnospiraceae	PowerSoil.HMP vs MagNAPure
9.46385082	-1.9959877	0.80350941	-2.4840875	0.01298839	0.04151191	Polyangiaceae	EasyDNA vs PowerSoil.HMP
28.7248963	-1.5390212	0.55585699	-2.7687358	0.00562742	0.04164294	Acetobacteraceae	FastDNA vs MagNAPure
355.031722	1.0232927	0.39395485	2.59748725	0.00939086	0.04173716	[Weeksellaceae]	FastDNA vs PowerSoil.HMP
7.76499835	2.85424593	0.91600206	3.11598202	0.00183333	0.04216669	Victivallaceae	InnuPURE vs MagNAPure
104.663517	1.13861777	0.41527505	2.74184006	0.00610961	0.04238541	Rikenellaceae	FastDNA vs MagNAPure
4421.89202	1.03709142	0.39261611	2.64148974	0.00825423	0.042494	Campylobacteraceae	QIAStool vs QIAStool+BB
622.632062	0.81772245	0.33204467	2.4626881	0.01378998	0.04266148	Flavobacteriaceae	EasyDNA vs PowerSoil.HMP
4.87640195	2.6847996	1.09159219	2.45952621	0.01391205	0.04266148	Corynebacteriaceae	EasyDNA vs PowerSoil.HMP
3.24919787	2.49848134	1.01818471	2.45385864	0.01413325	0.04266148	Leuconostocaceae	EasyDNA vs PowerSoil.HMP
28.7345598	-1.1664509	0.51264479	-2.275359	0.02288441	0.04290826	Parachlamydiaceae	EasyDNA vs MagNAPure
377.356263	-0.5824062	0.2219046	-2.6245793	0.00867561	0.04489627	Sphingobacteriaceae	QIAStool vs MagNAPure
2.31382502	-2.5964006	1.07577379	-2.4135191	0.0157993	0.045186	Nocardiaceae	EasyDNA vs FastDNA
719.094782	-1.0989013	0.39008341	-2.8170931	0.00484605	0.04537662	Streptococcaceae	QIAStool+BB vs MagNAPure
642.395629	-0.8990233	0.37439073	-2.4012969	0.01633708	0.04580788	Rhodocyclaceae	EasyDNA vs FastDNA
28.0267224	2.05253499	0.73852611	2.77923143	0.00544877	0.04586337	Succinivibrionaceae	QIAStool+BB vs MagNAPure
2750.67146	1.29995626	0.47394333	2.74285169	0.00609082	0.04586337	Bacteroidaceae	QIAStool+BB vs MagNAPure
355.031722	-1.1521059	0.42121139	-2.7352202	0.00623386	0.04586337	[Weeksellaceae]	QIAStool+BB vs MagNAPure
141.200238	-0.9400847	0.36121392	-2.6025706	0.00925278	0.04593342	Chitinophagaceae	QIAStool vs QIAStool+BB
164.684355	1.34712441	0.50064185	2.69079464	0.00712821	0.04654299	Clostridiaceae	FastDNA vs MagNAPure
59.9667273	-1.7611005	0.48922251	-3.5997946	0.00031847	0.04757822	Phyllobacteriaceae	FastDNA vs QIAStool+BB
118.159531	-1.5582244	0.44756329	-3.4815733	0.00049848	0.04757822	Rhodobacteraceae	FastDNA vs QIAStool+BB
858.112951	-1.1709513	0.34503626	-3.393705	0.00068954	0.04757822	Legionellaceae	FastDNA vs QIAStool+BB
44.0121162	-1.2198065	0.54714401	-2.2294067	0.02578686	0.04768803	Microbacteriaceae	EasyDNA vs MagNAPure
1756.14884	-1.401038	0.46013475	-3.0448428	0.00232802	0.04819001	Neisseriaceae	InnuPURE vs MagNAPure
1756.14884	-1.1234205	0.43466028	-2.5845943	0.00974937	0.04839863	Neisseriaceae	InnuPURE vs QIAStool
8.97993102	-1.6808038	0.75875734	-2.215206	0.02674593	0.04879324	Sediment-4	EasyDNA vs MagNAPure
3668.6678	1.90277585	0.63250342	3.00832499	0.00262692	0.04943387	Lachnospiraceae	InnuPURE vs MagNAPure
164.684355	1.08879511	0.4606952	2.36337412	0.01810938	0.04980079	Clostridiaceae	EasyDNA vs FastDNA
79.5100271	-0.9798214	0.380222	-2.5769718	0.00996701	0.05032123	Bdellovibrionaceae	QIAStool vs MagNAPure
186.231671	-0.7151168	0.30408292	-2.3517165	0.01868701	0.05041968	Rhizobiaceae	EasyDNA vs FastDNA
44.0121162	-1.4843431	0.56162354	-2.6429504	0.00821871	0.05068203	Microbacteriaceae	FastDNA vs MagNAPure
7.94601326	2.35096963	0.93228322	2.52173329	0.01167782	0.0511408	Micrococcaceae	PowerSoil.HMP vs QIAStool
333.941631	0.99974328	0.39058752	2.55958841	0.01047962	0.05164955	Porphyromonadaceae	QIAStool vs MagNAPure
19.2945448	-1.6538315	0.65473079	-2.5259718	0.01153788	0.05224094	BA008	EasyDNA vs InnuPURE
169.692077	1.21462904	0.47776311	2.5423249	0.01101178	0.05253157	Pirellulaceae	InnuPURE vs QIAStool
28.0267224	-1.7810983	0.703404	-2.5321129	0.01133775	0.05253157	Succinivibrionaceae	InnuPURE vs QIAStool
118.339928	0.77103841	0.32521728	2.37084086	0.01774767	0.05259764	[Chthoniobacteraceae]	EasyDNA vs PowerSoil.HMP
22.2584553	1.82334122	0.76805203	2.37398139	0.01759744	0.05314426	S24-7	EasyDNA vs QIAStool+BB
3.24919787	2.62550795	1.04618943	2.50959136	0.01208709	0.05324855	Leuconostocaceae	EasyDNA vs InnuPURE
243.261282	-0.6779435	0.28737692	-2.3590743	0.01832059	0.053326	GZKB119	EasyDNA vs PowerSoil.HMP
70.3986464	1.1324594	0.45348268	2.49724953	0.01251608	0.05368742	Desulfovibrionaceae	EasyDNA vs InnuPURE
7.94601326	1.90646899	0.81181724	2.34839678	0.01885442	0.05391704	Micrococcaceae	EasyDNA vs PowerSoil.HMP
129.664767	-1.213787	0.48816066	-2.4864499	0.01290247	0.05392572	Sphingomonadaceae	EasyDNA vs InnuPURE
12.0832988	2.17200442	0.8174185	2.65715106	0.00788041	0.05411217	Microthrixaceae	QIAStool+BB vs MagNAPure
28.0267224	1.75955604	0.67705141	2.59885144	0.00935362	0.05464486	Succinivibrionaceae	FastDNA vs MagNAPure
622.632062	-0.7523083	0.32535862	-2.3122433	0.02076428	0.05498688	Flavobacteriaceae	EasyDNA vs FastDNA
579.211848	0.70977486	0.28578343	2.48361094	0.01300578	0.05505781	Cryomorphaceae	PowerSoil.HMP vs QIAStool
11.2601759	2.2792279	0.96857952	2.35316549	0.01861435	0.05511307	Peptococcaceae	EasyDNA vs QIAStool+BB
22.9873604	-2.4720555	0.97855392	-2.5262333	0.01152929	0.05526107	Staphylococcaceae	QIAStool vs QIAStool+BB
310.195483	1.0118008	0.40293184	2.51109669	0.01203567	0.05576528	Aeromonadaceae	QIAStool vs QIAStool+BB
6.96503794	1.90349744	0.77276169	2.46323991	0.01376878	0.05640756	Piscirickettsiaceae	PowerSoil.HMP vs QIAStool
224.417038	-0.9883822	0.43171742	-2.2894192	0.02205501	0.05734302	Caulobacteraceae	EasyDNA vs FastDNA
2.31382502	-2.7483795	1.10394795	-2.4895916	0.012789	0.05734421	Nocardiaceae	QIAStool vs QIAStool+BB
182.91656	-0.6936928	0.32325492	-2.145962	0.03187601	0.05737304	Planctomycetaceae	EasyDNA vs MagNAPure
3.14112548	-2.3181345	1.08289035	-2.1406919	0.0322989	0.05737304	OM60	EasyDNA vs MagNAPure
186.231671	-0.9118034	0.37170792	-2.4530104	0.01416662	0.05772899	Rhizobiaceae	EasyDNA vs InnuPURE
5.49329045	-1.8319005	0.72941169	-2.5114767	0.01202272	0.05787682	Cyclobacteriaceae	QIAStool vs MagNAPure
8.97993102	-2.0613208	0.84378326	-2.4429505	0.01456773	0.05791562	Sediment-4	EasyDNA vs InnuPURE

12.2684488	-1.5691396	0.64325243	-2.4393838	0.01471233	0.05838958	Synergistaceae	PowerSoil.HMP vs QIAStool
104.663517	0.87151266	0.3830252	2.27534024	0.02288553	0.05843983	Rikenellaceae	EasyDNA vs FastDNA
127.559639	-0.9708623	0.3795923	-2.5576449	0.01053836	0.05848792	Bradyrhizobiaceae	FastDNA vs MagNAPure
11.4855136	1.65096216	0.68095654	2.42447508	0.01533054	0.05899933	Gemmatimonadaceae	PowerSoil.HMP vs QIAStool
4830.98886	0.64823777	0.30574015	2.12022453	0.03398711	0.0595878	Moraxellaceae	EasyDNA vs MagNAPure
4234.46977	0.80709372	0.3244551	2.48753594	0.01286315	0.06051525	Ruminococcaceae	QIAStool vs MagNAPure
6.96503794	1.86675561	0.74910667	2.49197568	0.01270347	0.06055322	Piscirickettsiaceae	FastDNA vs QIAStool
377.356263	-0.5435773	0.23659302	-2.2975206	0.02158909	0.06067279	Sphingobacteriaceae	EasyDNA vs PowerSoil.HMP
579.211848	0.87084781	0.35361388	2.46270821	0.01378921	0.06068046	Cryomorphaceae	InnuPURE vs QIAStool
50.9419349	1.38093288	0.56180151	2.45804409	0.0139696	0.06068046	Desulfobulbaceae	InnuPURE vs QIAStool
6.92597984	-1.9786325	0.86497437	-2.2875042	0.02216641	0.06123941	[Chromatiaceae]	EasyDNA vs PowerSoil.HMP
16.3205056	-1.7055823	0.7409508	-2.3018834	0.02134175	0.06197316	Methylophilaceae	EasyDNA vs QIAStool+BB
113.494552	-0.8384629	0.34080263	-2.4602596	0.01388365	0.06234019	Alcaligenaceae	FastDNA vs QIAStool
70.3986464	-0.8486684	0.34519175	-2.4585418	0.01395025	0.06234019	Desulfovibrionaceae	FastDNA vs QIAStool
355.031722	-1.0110411	0.39463636	-2.5619562	0.01040844	0.06245066	[Weeksellaceae]	PowerSoil.HMP vs MagNAPure
47.3908296	-0.9325292	0.38114769	-2.4466348	0.01441969	0.06248531	Christensenellaceae	FastDNA vs QIAStool
104.663517	-1.1350299	0.46606219	-2.4353614	0.01487692	0.06266339	Rikenellaceae	InnuPURE vs QIAStool
4.18916425	2.29762813	1.0113383	2.27186899	0.02309442	0.06273985	Nocardioidaceae	EasyDNA vs PowerSoil.HMP
28.0267224	-1.3727681	0.56486719	-2.4302494	0.01508844	0.06346019	Succinivibrionaceae	FastDNA vs QIAStool
264.350163	0.8838018	0.36532112	2.41924641	0.0155527	0.0635831	Leptotrichiaceae	InnuPURE vs QIAStool
113.494552	-1.0371853	0.43307898	-2.3949102	0.01662443	0.06456144	Alcaligenaceae	InnuPURE vs QIAStool
33.4444509	2.01392801	0.8425903	2.39016283	0.0168409	0.06456144	Coriobacteriaceae	InnuPURE vs QIAStool
3.5697885	2.44429648	1.0258444	2.38271659	0.01718542	0.06456144	Beutenbergiaceae	InnuPURE vs QIAStool
103.008126	1.10013504	0.42775185	2.57190011	0.01011421	0.06511021	Enterococcaceae	QIAStool+BB vs MagNAPure
649.223144	0.86594468	0.34641928	2.49970117	0.01242981	0.06570043	Bifidobacteriaceae	FastDNA vs MagNAPure
164.684355	1.13212163	0.49900063	2.26877795	0.02328183	0.06633126	Clostridiaceae	EasyDNA vs QIAStool+BB
28.7248963	1.22866154	0.51896204	2.36753646	0.01790696	0.06688775	Acetobacteraceae	PowerSoil.HMP vs QIAStool
186.231671	-0.6985777	0.3119467	-2.2394135	0.02512902	0.06714804	Rhizobiaceae	EasyDNA vs PowerSoil.HMP
3.21882027	-2.2403395	1.01657284	-2.203816	0.02753729	0.06722862	Geobacteraceae	EasyDNA vs FastDNA
243.261282	-0.6192531	0.28126021	-2.2017089	0.02768588	0.06722862	GZKB119	EasyDNA vs FastDNA
4.87640195	2.37711291	1.0817303	2.19750978	0.02798406	0.06722862	Corynebacteriaceae	EasyDNA vs FastDNA
377.356263	-0.5086707	0.23180568	-2.194384	0.02820781	0.06722862	Sphingobacteriaceae	EasyDNA vs FastDNA
4.27355164	2.11193339	0.88047945	2.39861747	0.0164571	0.067239	Eubacteriaceae	FastDNA vs QIAStool
264.350163	-0.7890883	0.32627137	-2.4185032	0.01558451	0.06769521	Leptotrichiaceae	QIAStool vs QIAStool+BB
17.3475351	-1.9832942	0.83527701	-2.3744149	0.01757679	0.06821467	Isosphaeraceae	EasyDNA vs InnuPURE
1181.58837	1.95342448	0.88069728	2.21804306	0.0265519	0.06980579	Prevotellaceae	EasyDNA vs PowerSoil.HMP
19.2945448	-1.2707305	0.53094712	-2.3933277	0.01669632	0.07032693	BA008	QIAStool vs QIAStool+BB
3.24919787	2.02439771	0.98885767	2.04720839	0.04063763	0.07033436	Leuconostocaceae	EasyDNA vs MagNAPure
70.3986464	0.99071862	0.39671413	2.49731114	0.01251391	0.07066679	Desulfovibrionaceae	PowerSoil.HMP vs MagNAPure
22.1599751	1.53415787	0.75524233	2.03134519	0.04221999	0.07214808	Sanguibacteraceae	EasyDNA vs MagNAPure
79.5100271	0.88963046	0.37786776	2.35434341	0.01855546	0.0724779	Bdellovibrionaceae	FastDNA vs QIAStool
5.49329045	1.69380301	0.72064323	2.3504044	0.01875302	0.0724779	Cyclobacteriaceae	FastDNA vs QIAStool
4.81963649	-2.3118932	0.98963119	-2.336116	0.0194852	0.07386249	auto67_4W	EasyDNA vs InnuPURE
2.55199853	-2.2827904	1.04339628	-2.187846	0.02868082	0.07420014	Holophagaceae	EasyDNA vs PowerSoil.HMP
182.91656	0.73524789	0.33701129	2.18167136	0.0291338	0.07420014	Planctomycetaceae	EasyDNA vs PowerSoil.HMP
22.9873604	2.14709978	0.91924374	2.33572414	0.01950562	0.07451148	Staphylococcaceae	EasyDNA vs QIAStool
642.395629	-0.8615187	0.3894487	-2.2121495	0.02695634	0.07537791	Rhodocyclaceae	EasyDNA vs QIAStool+BB
1.01494259	-2.6305311	1.10333506	-2.3841635	0.017118	0.07685828	Gemmataceae	QIAStool vs MagNAPure
310.195483	0.8934204	0.37551544	2.37918422	0.017351	0.07685828	Aeromonadaceae	QIAStool vs MagNAPure
254.528648	-0.9770092	0.41101457	-2.3770671	0.01745091	0.07685828	Xanthomonadaceae	QIAStool vs MagNAPure
4.06716246	-1.8067423	0.77930346	-2.3184066	0.02042724	0.07687091	Elusimicrobiaceae	FastDNA vs QIAStool
19.2945448	-1.1273417	0.47588079	-2.3689582	0.01783827	0.07692754	BA008	QIAStool vs MagNAPure
4.81963649	-2.1101927	0.9610752	-2.1956582	0.02811642	0.07719234	auto67_4W	EasyDNA vs QIAStool+BB
2.3093169	-2.3470892	1.01440126	-2.313768	0.02068045	0.07745031	Myxococcaceae	EasyDNA vs QIAStool
22.5747955	1.12730926	0.52924208	2.13004465	0.03316793	0.07775432	Peptostreptococcaceae	EasyDNA vs FastDNA
736.495903	2.47243917	0.73392038	3.36881115	0.00075493	0.07813539	Enterobacteriaceae	PowerSoil.HMP vs QIAStool+BB
22.9873604	1.9190924	0.96704297	1.98449547	0.04720063	0.07965107	Staphylococcaceae	EasyDNA vs MagNAPure
164.684355	1.22958079	0.50750114	2.42281388	0.01540081	0.07974548	Clostridiaceae	PowerSoil.HMP vs MagNAPure
4234.46977	0.850399	0.35438523	2.39964574	0.01641095	0.07974548	Ruminococcaceae	PowerSoil.HMP vs MagNAPure
141.200238	0.87189043	0.36402356	2.39514836	0.01661364	0.07974548	Chitinophagaceae	PowerSoil.HMP vs MagNAPure
649.223144	-0.8825037	0.37134215	-2.3765246	0.0174766	0.07989301	Bifidobacteriaceae	PowerSoil.HMP vs MagNAPure
377.356263	-0.4611657	0.23319393	-1.9776057	0.0479732	0.07995534	Sphingobacteriaceae	EasyDNA vs MagNAPure
13.1651756	-1.4992527	0.69891407	-2.1451174	0.03194346	0.08010438	Helicobacteraceae	EasyDNA vs PowerSoil.HMP
182.91656	1.18518163	0.41749716	2.83877771	0.00452867	0.08094996	Planctomycetaceae	InnuPURE vs PowerSoil.HMP
1.55959833	3.06644109	1.09689413	2.79556704	0.00518088	0.08231836	Dermatophilaceae	InnuPURE vs PowerSoil.HMP

131.739477	-0.797345	0.33199055	-2.4017101	0.01631864	0.08233494	Hyphomicrobiaceae	FastDNA vs MagNAPure
131.739477	-0.9313915	0.40909714	-2.2767002	0.02280414	0.08447896	Hyphomicrobiaceae	EasyDNA vs InnuPURE
13.1651756	-1.4567213	0.64093998	-2.272789	0.0230389	0.08462364	Helicobacteraceae	EasyDNA vs QIAStool
579.211848	-0.6602391	0.2839074	-2.3255439	0.0200429	0.08467101	Cryomorphaceae	QIAStool vs MagNAPure
92.6279814	-0.9949071	0.43808997	-2.2710109	0.02314632	0.08486984	Oxalobacteraceae	FastDNA vs QIAStool
29.0135086	-1.2323803	0.54478605	-2.2621363	0.02368898	0.08536974	Alteromonadaceae	EasyDNA vs QIAStool
243.261282	-0.6671317	0.31081555	-2.1463909	0.0318418	0.08585914	GZKB119	EasyDNA vs QIAStool+BB
38.1375375	1.08211604	0.55767402	1.94040963	0.05232993	0.08615293	Procabacteriaceae	EasyDNA vs MagNAPure
505.493872	-1.0047589	0.43784747	-2.2947692	0.02174636	0.08647001	Erysipelotrichaceae	QIAStool vs QIAStool+BB
4.18916425	-2.3294501	1.0153187	-2.2943043	0.02177303	0.08647001	Nocardioidaceae	QIAStool vs QIAStool+BB
11.4855136	-1.3573909	0.70420385	-1.927554	0.05391063	0.08768596	Gemmatimonadaceae	EasyDNA vs MagNAPure
5.49329045	-1.5087589	0.78541121	-1.9209795	0.05473429	0.08796582	Cyclobacteriaceae	EasyDNA vs MagNAPure
3.07801265	2.12104453	0.94429648	2.24616376	0.02469352	0.08827934	Caldilineaceae	FastDNA vs QIAStool
719.094782	0.74746237	0.33440796	2.23518117	0.02540546	0.08860928	Streptococcaceae	FastDNA vs QIAStool
254.528648	0.93025684	0.41369801	2.24863746	0.02453557	0.08902907	Xanthomonadaceae	PowerSoil.HMP vs QIAStool
20.0435737	-1.0503917	0.50791832	-2.0680327	0.03863694	0.08911424	Saprospiraceae	EasyDNA vs FastDNA
164.684355	1.30379789	0.53572935	2.43368761	0.01494589	0.0905545	Clostridiaceae	QIAStool+BB vs MagNAPure
141.200238	0.93724671	0.38880135	2.41060559	0.01592606	0.09113246	Chitinophagaceae	QIAStool+BB vs MagNAPure
377.356263	0.62154676	0.27788464	2.23670791	0.02530544	0.09256462	Sphingobacteriaceae	InnuPURE vs QIAStool
33.4444509	-1.6904111	0.74230936	-2.2772326	0.02277234	0.09427748	Coriobacteriaceae	QIAStool vs MagNAPure
105.619852	-1.021215	0.44996553	-2.2695406	0.02323547	0.09430867	Nitrosomonadaceae	QIAStool vs MagNAPure
22.1599751	1.78724592	0.80863491	2.21020129	0.0270912	0.09439858	Sanguibacteraceae	PowerSoil.HMP vs QIAStool
4.81963649	-1.8045008	0.81885869	-2.2036779	0.027547	0.09439858	auto67_4W	PowerSoil.HMP vs QIAStool
50.9419349	1.03667507	0.47253405	2.1938632	0.02824525	0.09439858	Desulfobulbaceae	PowerSoil.HMP vs QIAStool
8.22875632	-1.847976	0.8213212	-2.2500041	0.02444869	0.0943991	Solibacteraceae	QIAStool vs QIAStool+BB
29.0135086	-1.1098265	0.58897604	-1.8843321	0.05952006	0.09453186	Alteromonadaceae	EasyDNA vs MagNAPure
91.6048239	-0.7732133	0.36814481	-2.1002967	0.03570275	0.09458097	Halomonadaceae	EasyDNA vs QIAStool+BB
2.3093169	-2.2044833	1.06566745	-2.0686409	0.0385798	0.09503659	Myxococcaceae	EasyDNA vs PowerSoil.HMP
642.395629	-0.7773137	0.37669477	-2.0635108	0.03906412	0.09503659	Rhodocyclaceae	EasyDNA vs PowerSoil.HMP
118.159531	-1.6497585	0.49812382	-3.3119446	0.0009265	0.09589264	Rhodobacteraceae	FastDNA vs InnuPURE
49.8040986	-1.915968	0.59985447	-3.1940546	0.0014029	0.0967998	Actinomycetaceae	PowerSoil.HMP vs QIAStool+BB
31.145957	-2.1008028	0.92480483	-2.2716174	0.02310963	0.0973037	Aerococcaceae	FastDNA vs PowerSoil.HMP
15.0781299	1.10535598	0.54633858	2.02320688	0.04305182	0.0977208	[Odoribacteraceae]	EasyDNA vs FastDNA
44.0121162	-1.4733537	0.62305494	-2.3647251	0.01804346	0.09781453	Microbacteriaceae	QIAStool+BB vs MagNAPure
1756.14884	-1.2328734	0.46055978	-2.6769019	0.00743064	0.09793244	Neisseriaceae	InnuPURE vs PowerSoil.HMP
243.261282	-0.9574556	0.36120868	-2.6506993	0.00803253	0.09793244	GZKB119	InnuPURE vs PowerSoil.HMP
44.0121162	1.80866043	0.68432745	2.64297515	0.00821811	0.09793244	Microbacteriaceae	InnuPURE vs PowerSoil.HMP
2.61442095	2.04322522	0.93447232	2.18650159	0.02877893	0.0979854	Ellin515	FastDNA vs QIAStool
10.9576795	-1.3578423	0.72891946	-1.8628152	0.06248827	0.09809204	Acholeplasmataceae	EasyDNA vs MagNAPure
593.953594	-0.6205976	0.28170633	-2.2029949	0.0275951	0.09841013	Cytophagaceae	EasyDNA vs InnuPURE
28.7345598	-1.3134923	0.59690916	-2.2004895	0.02777218	0.09841013	Parachlamydiaceae	EasyDNA vs InnuPURE
22.2584553	1.27850913	0.68878738	1.85617385	0.06342878	0.09842397	S24-7	EasyDNA vs MagNAPure
44.0121162	1.16764997	0.53725954	2.17334431	0.02975441	0.09856449	Microbacteriaceae	FastDNA vs QIAStool
3.5697885	2.04025878	0.94204076	2.1657861	0.03032754	0.09856449	Beutenbergiaceae	FastDNA vs QIAStool
38.1375375	1.28724283	0.57931343	2.22201449	0.02628233	0.09873631	Procabacteriaceae	QIAStool vs QIAStool+BB
283.625111	1.51356979	0.66399228	2.27949907	0.02263742	0.09878145	Lactobacillaceae	PowerSoil.HMP vs MagNAPure
70.3986464	0.85713572	0.41436171	2.06856886	0.03858656	0.09879332	Desulfovibrionaceae	EasyDNA vs QIAStool+BB
4.87640195	2.25110021	1.09119838	2.06296146	0.03911629	0.09879332	Corynebacteriaceae	EasyDNA vs QIAStool+BB
1.41904526	2.24885942	1.09088645	2.06149725	0.03925563	0.09879332	Dehalobacteriaceae	EasyDNA vs QIAStool+BB
4.46443875	1.94043662	0.89629733	2.16494746	0.03039171	0.09896787	Xanthobacteraceae	PowerSoil.HMP vs QIAStool
28.7248963	-1.2327152	0.56097099	-2.1974669	0.02798712	0.09899147	Acetobacteraceae	EasyDNA vs QIAStool
38.1375375	-1.3782365	0.62635739	-2.2003995	0.02777856	0.09900564	Procabacteriaceae	InnuPURE vs QIAStool
7.94601326	2.49871504	0.90484448	2.76148565	0.0057539	0.09925485	Micrococcaceae	InnuPURE vs MagNAPure
13.1651756	-1.6658237	0.76085582	-2.1894078	0.02856721	0.09927107	Helicobacteraceae	InnuPURE vs QIAStool
224.417038	-1.0205732	0.44198438	-2.3090707	0.02093966	0.09943231	Caulobacteraceae	FastDNA vs MagNAPure
1756.14884	-0.8783179	0.38202572	-2.2991067	0.02149888	0.09943231	Neisseriaceae	FastDNA vs MagNAPure
49.8040986	1.32588889	0.5667839	2.33931996	0.01931888	0.09949224	Actinomycetaceae	QIAStool+BB vs MagNAPure
1756.14884	-0.7194269	0.35366435	-2.034208	0.04193062	0.09951465	Neisseriaceae	EasyDNA vs PowerSoil.HMP
10.9576795	-1.5130076	0.74448962	-2.0322749	0.04212583	0.09951465	Acholeplasmataceae	EasyDNA vs PowerSoil.HMP
1.41904526	2.33278776	1.06856001	2.18311347	0.02902745	0.09977434	Dehalobacteriaceae	EasyDNA vs InnuPURE
4.87640195	2.35164946	1.07956479	2.17833102	0.0293814	0.09977434	Corynebacteriaceae	EasyDNA vs InnuPURE
15.0781299	1.47004964	0.68024796	2.16104968	0.0306915	0.09985342	[Odoribacteraceae]	EasyDNA vs InnuPURE
60.913589	-0.9475891	0.43964149	-2.1553677	0.03113307	0.09985342	SB-1	EasyDNA vs InnuPURE
1.54406709	-2.3845863	1.10706543	-2.1539706	0.03124248	0.09985342	Mycobacteriaceae	EasyDNA vs InnuPURE