The Evolution of Genetic Bandwagoning

Idan S. Solon

University of Central Florida, Burnett School of Biomedical Sciences

Orlando, Florida, United States 32816

Correspondence: IdanSolon@gmail.com

Abstract

The concept of "genetic bandwagoning" is introduced. In genetic bandwagoning, a bandwagoning allele evaluates

the outlook for lineage fitness of the individual holding the allele. If its holder's lineage fitness figures to be low,

the bandwagoning allele induces its holder to relinquish some, or all, of the holder's own resources (e.g., food,

territory, mates), which are then used by individuals that happen to be nearby. While some nearby individuals may

not possess a copy of the bandwagoning allele, it is likely, if assortment is sufficiently positive, that numerous

individuals do and most of those that do have a better lineage fitness outlook than that of the individual that was

induced to forfeit the resources, since the inducement is contingent upon a poor lineage fitness outlook. Due to their

better lineage fitness outlook, the nearby individuals make better usage of the resources than the forfeiting

individuals would have and the ratio between the quantity of descendants nearby individuals produce with the

resources and the quantity of descendants that would have been produced by the forfeiting individuals tends to

increase with each generation due to multiple reasons examined in section 2. If assortment is sufficiently positive,

more copies of the allele are produced over the ensuing generations than if the resources had not been forfeited, as

the allele is selected by "jumping on the bandwagon" of these fit lineages that use the resources.

Two ways to bandwagon are introduced: resonation and reservation. In resonation, the bandwagoning allele

accomplishes the two steps of bandwagoning—1) evaluating its holder's lineage fitness outlook and 2) inducing a

holder with a low (i.e., poor) outlook to relinquish resources—sequentially. The necessity for reservation occurs

because a lineage's fitness outlook can depend upon aspects of quality which are difficult to evaluate during a

generation in which many individuals are able to avoid death or serious harm. In reservation, a bandwagoning allele

solves this problem of evaluating quality by inducing its holder to reserve from (i.e., withhold) some of its ability to

survive against one or more natural enemies, particularly predators or parasites: An individual that can remain

healthy in spite of reserving from its ability to survive predators or parasites is likely to have genes that enable its

descendants to survive when the predator or parasite threat is more severe; however, if the individual perishes or

becomes sick as a result of reservation, its lineage fitness outlook is determined to be low and the relinquishment of

resources occurs with this same perishing or sickness. That is, the individual's perishing or sickness due to

reservation serves simultaneously: 1) to identify that individual as having a low lineage fitness outlook and 2) as the

source of relinquishing the individual's resources on the basis of this low lineage fitness outlook.

Genetic bandwagoning constitutes a novel mechanism for the evolution of cooperation. Numerous paradoxes can be

explained by the actions of bandwagoning alleles that use a combination of resonation and reservation, including the

phenotypic cooccurrences of robust defenses against parasites and depressive symptoms; phenotypic cooccurrences

of robust defenses against parasites and photosynthetic costs; phenotypic cooccurrences of susceptibility to parasites

and androgenic benefits; the maintenance of honest signals when there are opportunities to cheat; the forfeiture of

reproductive opportunities by functionally capable individuals; and the differential nurturing of offspring.

Keywords:

handicap principle; honest signaling; cooperation; depression; social contraception

1) Introduction

Multiple scholars have discussed circumstances that might lead a lower-quality[1] individual to make an

uncoerced forfeiture of resources (e.g., food, territory, mating opportunities) in such a manner that they are not

directly bestowed to anyone but are instead left to be utilized by any nearby individual(s) and the individuals that

utilize them are often unaware of the forfeiting individual's identity or, indeed, that the resources were even

relinquished. (Therefore, such forfeitures are not explained by direct or indirect reciprocity: Nowak, 2012.)

For example, multiple scholars have hypothesized that if an individual is in moribund condition due to a

parasite infection or some other malady, it should commit suicide. That is, they have argued that there can be

natural selection for alleles that, under these circumstances, induce their holder to commit suicide, thereby

relinquishing the holder's resources to nearby individuals (e.g., Dawkins, 1976; Refardt et al., 2013). Such

scenarios are typically viewed in terms of Hamilton's (1964) rule, which holds that there can be natural selection for

a social act if $C < rB$, where C is the cost to the actor; B is the benefit to recipients of the act; and r is the relatedness

between the actor and the recipients. The cost of suicide is the reproduction that is foregone by the actor committing

it. Since a moribund actor has low reproductive outlook, its cost of suicide is low compared to the benefit nearby

individuals would receive from utilizing the actor's resources. The moribund actor can expect considerably lower

fitness than potential recipients would from the same resources. Consequently, there can be natural selection for a

trait that leads an actor to commit suicide in response to the actor's moribundity, even if relatedness between the

actor and those benefitting is low (Refardt et al., 2013).

In the present paper, I argue that the forfeiture of resources by a lower-quality individual to nearby

individuals has been naturally selected to occur: 1) in numerous circumstances besides the moribundity of the

lower-quality individual and 2) via numerous mechanisms besides suicide. Forfeiture of resources from a lower-

quality individual becomes naturally selected in more circumstances than are currently appreciated because most

analyses of the evolution of social behavior examine solely the costs and benefits that result from an act by the first

---

[1] The term "quality" is used often but inconsistently in evolutionary literature (Bergeron et al., 2011; Hill, 2011; Wilson & Nussey, 2010). An individual's "genetic quality" is sometimes distinguished from its "phenotypic quality"; when this occurs, the former typically refers to "good genes" and the latter to condition. The latter is typically interpreted to be an indication of the former and that is how the terms are used in the present paper. I assume, unless indicated otherwise, that an individual's "quality" in either sense covaries positively with the individual's outlook for lineage fitness, so that a higher-quality individual can, on average, expect to have a higher lineage fitness than a lower-quality individual.

generation following the act (Hunt et al., 2004). However, the descendants of a higher-quality individual can often

be expected to produce more offspring per individual than the descendants of a lower-quality individual. (Some

reasons for this expectation are explored in section 2.) If this expectation holds, then forfeiture of resources by a

lower-quality individual to higher-quality individuals is more likely to satisfy Hamilton's (1964) rule if the costs and

benefits are projected longer-term than if they are only calculated for a single generation. This point is developed

below.

Consider that a focus individual can expect to produce a quantity of A offspring but that, alternatively, it

can forfeit its resources to individuals that can expect to produce a total quantity of B offspring with the same

resources and that the relatedness between the focus individual and the individuals that would benefit is r.

According to Hamilton's (1964) rule, the focus individual should forfeit its resources if[2]:

$$A < rB \qquad (1)$$

Inequality (2) offers Hamilton's rule for the same scenario as inequality (1) but when looking two

generations ahead instead of one, where S represents the quantity of offspring that each of the focus individual's

offspring can be expected to produce, on average; V represents the quantity of offspring that each of the utilizing

individuals' offspring can be expected to produce, on average; and the other variables are the same as in inequality

(1):

$$AS < rBV \qquad (2)$$

Inequality (2) demonstrates that as the focus individual looks ahead two generations, the cost of forfeiting

the resources would be the grandoffspring it could obtain by using the resources (which is the product of A times S)

and the benefit would be the grandoffspring the utilizing individuals could obtain with the same resources (which is

the product of B times V). According to inequality (1), the focus individual that evaluates the costs and benefits

solely one generation ahead would only forfeit resources if $A < rB$; however, according to inequality (2), if the same

individual looks ahead two generations and anticipates that $S < V$, then it might forfeit resources even if $A \geq rB$.

Similarly, if the same individual looks three generations ahead, it becomes still more likely to forfeit resources than

---

[2] In a sexual population, there is a 50% likelihood that the focus individual shares an allele with its own offspring and there is the same 50% likelihood that the benefiting individuals share an allele with their own offspring; therefore, these effects cancel out.

if it looks two generations ahead, if it expects that its grandoffspring would produce fewer offspring per individual

than the grandoffspring of the individuals that would utilize the resources. And generally, if a focus individual looks

t+1 generations ahead, it becomes more likely to forfeit resources than if it had looked *t* generations ahead, as long

as it expects its descendants to produce fewer offspring per individual in generation *t* than the descendants of the

utilizing individuals produce in generation *t*.

In the present paper, I introduce the concept of genetic "bandwagoning." A bandwagoning allele[3] remains

latent when it is located within an individual with a sufficiently high lineage fitness outlook, but when it is located

within an individual with a low lineage fitness outlook, it induces that individual to act in a manner that leads that

individual to forfeit some or all of its resources, which are then utilized by whatever individuals are nearby. While

some nearby individuals may not possess a copy of the bandwagoning allele, numerous individuals do if assortment

is sufficiently positive and most nearby individuals have a higher lineage fitness outlook than that of the individual

that forfeited the resources, since the forfeiture was contingent upon a low lineage fitness outlook. If the outlooks

for lineage fitness of these individuals that use the resources are high enough compared to the lineage fitness outlook

of the forfeiting individual and if assortment is sufficiently high, more copies of the allele can be expected to be

produced over the ensuing generations if the resources are forfeited than if the resources are not forfeited. The allele

can be said to "jump on the bandwagon" of lineages that have a higher fitness outlook. If a person "jumps on the

bandwagon" of some entity (e.g., person, group, cause), it means that person has only decided to support that entity

because it has become successful, with the expectation of sharing in the entity's future success. A "bandwagoning

allele" fits this description. Regardless of whether or not the allele is located within an individual that has a high

lineage fitness outlook, it is acting to increase the likelihood that resources to which its holder has access are utilized

---

[3] Two clarifications about the usage of the term "bandwagoning allele" in the present paper: The singular "bandwagoning allele" 1) is used to represent any heritable element(s) (e.g., gene, epigenetic mark) responsible for bandwagoning within a single individual and 2) may represent a multiplicity of heritable elements within that individual. The term "bandwagoning allele" is used to represent the totality of bandwagoning elements operating within a single individual and "bandwagoning alleles" is used to represent the totality of bandwagoning elements operating on a scale larger than the individual.

by individuals with a high lineage fitness outlook: If it is located within an individual with a high lineage fitness

outlook, it is latent so that the individual can use its resources and if it is located within an individual with a low

lineage fitness outlook, it induces forfeiture of resources so that they can be utilized by nearby individuals that have

a higher lineage fitness outlook. Some of these forfeited resources are used by individuals that have a high lineage

fitness outlook and a copy of the bandwagoning allele and the copies of the bandwagoning allele increase in quantity

by being part of these lineages that have high fitness.

1.1) The concept of genetic bandwagoning illuminated via analogy

Consider an analogy to finance. A low-quality individual that is induced by a bandwagoning allele to

forfeit resources to higher-quality individuals to which it is related by r is analogous to a person who forgoes

investing his or her own capital in order to have it allocated by an investing manager who can earn a higher rate of

return on the capital but who also keeps all but a fraction, r, of the final sum. In this analogy, the rates of return

represent, respectively, the quantity of offspring the low-quality individual's lineage could produce per individual

per generation and the quantity of offspring the higher-quality individuals' lineages could, on average, produce per

individual per generation. The greater the difference in return rate, or in offspring produced per individual per

generation, and the longer this difference persists, the lower the r that is acceptable. The obvious difference between

these two cases is that in genetic bandwagoning, resources are forfeited to whatever individuals happen to be nearby,

whereas in investing, people are typically aware of the specific quantities of capital they are directing to particular

managers. However, in each case, the individual that gives up the assets (e.g., resources, capital) sacrifices some

part of the ultimate proceeds. In exchange for investing capital, managers typically would keep some fraction of the

final sum, leaving the person with the rest, which is analogous to how the descendants produced via forfeited

resources would not all possess the bandwagoning allele, as some of the forfeited resources would be spent by

individuals that do not share the bandwagoning allele. In both cases, the greater capabilities of those spending the

assets would, over time, make up for that fraction of the proceeds.

Another analogy may also be illuminating. Maynard Smith (1978) wrote that the natural selection of

alleles that increase mutation or recombination rates is best understood in terms of the hitchhiking effect, wherein an

allele is selected not for its influence upon an individual's phenotype, but because it is located next to alleles that

favorably influence an individual's phenotype. In Maynard Smith's thinking, a gene increasing the mutation rate

would exert mostly unfavorable effects upon the individuals in which it is located, but it would be selected if the

descendants of an individual that is favorably affected were to gain in quantity over the ensuing generations until

they more than make up for all of the individuals that were unfavorably affected. The mutator gene would be

carried to higher frequency by the descendants of the favorably-affected lineage, having "hitched a ride" with them

(Maynard Smith, 1978, pg. 113). A bandwagoning allele must also make up for a fitness loss it imposes upon some

of its holders. For the bandwagoning allele, this loss is incurred by the holders that forfeit resources, some of which

wind up being utilized by individuals that do not hold a copy of the bandwagoning allele. Like a hitchhiking allele,

a bandwagoning allele makes up for that loss by "hitching a ride"—not with individual mutations but with an entire

genotype as it gains frequency over the ensuing generations. An allele that jumps on the bandwagon of a genotype

in the manner introduced in the present paper would likely continue to be associated with a fitter genotype, even if

the genotype were subjected to recombination and sex, since, after recombination and sex, it would still be located

with approximately half of the alleles to which its fitness advantage was attributable and since the half with which it

would be newly located would likely also offer a fitness advantage, given sexual selection upon both males and

females (Amundsen & Forsgren, 2001; Clutton-Brock et al., 2006; Edward & Chapman, 2011, 2012, 2013) and

positive assortment by quality (Bos et al., 2009; Dakin & Montgomerie, 2014; Holveck et al., 2011; Holveck &

Riebel, 2010; Schultzhaus et al., 2017; Veen & Otto, 2015).

1.2) Outline of the remainder of the paper

Section 2 examines three factors—differential allocation, heritability of quality and fitness, and stochastic

dominance—that can often lead the lineage of a higher-quality individual to grow faster than that of a lower-quality

individual, which increases the likelihood of natural selection for genetic bandwagoning because a forfeiture of

resources from a lower-quality individual to higher-quality individuals is more likely to satisfy Hamilton's rule if the

ratio of a higher-quality individual's number of descendants can be expected to grow compared to a lower-quality

individual's number of descendants. Section 3 introduces two ways in which bandwagoning can occur: resonation

and reservation. In particular, a model combining resonation and reservation is introduced verbally in section 3.2.5

and algebraically in section 3.3. Section 4 offers guidelines for applying this model that combines resonation and

reservation to empirical phenomena and section 5 examines six empirical paradoxes that can be explained by this

model. The concluding section 6 includes interpretations of genetic bandwagoning as a mechanism for the evolution

of cooperation and as a phenomenon that exerts significant evolutionary influence.

2) Why the ratio of a higher-quality individual's number of descendants to a lower-quality individual's

number of descendants would grow over time

2.1) Differential allocation

Mothers commonly invest more resources in offspring that they produce with fitter and/or higher-quality

fathers (Burley, 1986; Harris & Uller, 2009; Ratikainen & Kokko, 2010; Sheldon, 2000; Stiver & Alonzo, 2009).

This has been reported to include investment in larger and more eggs, larger clutches, more androgens, and greater

parental care (Stiver & Alonzo, 2009). Fathers have also been found to invest more in offspring produced with

higher-quality mothers (Ratikainen & Kokko, 2010; Stiver & Alonzo, 2009). By investing more in offspring from

higher-quality mates, parents invest more in offspring that figure to "generate higher returns on investment" than

others (Harris & Uller, 2009; Sheldon, 2000).

A narrower set of analyses has reported what seems to be a reverse pattern, by which parents allocate more

resources to offspring with non-preferred parents (Gowaty et al., 2007; Ratikainen & Kokko, 2010; Stiver &

Alonzo, 2009). Findings fitting this pattern have usually been explained by invoking the reproductive compensation

hypothesis (Gowaty, 2003; Gowaty, 2008; Gowaty et al., 2007), which is intended to explain parental allocation

under the exclusive set of circumstances in which preference for progenitors is self-referential (Gowaty, 2008).

However, an analysis by Harris & Uller (2009) concluded that greater investment in offspring with higher-quality

mates can be expected to occur much more often than the reverse. Consequently, a lower-quality individual can

often anticipate that its offspring are likely to receive less resources from a parent that mates with multiple

individuals, which reduces the lower-quality individual's lineage fitness outlook compared to that of higher-quality

individuals.

2.2) Heritability of quality and fitness

2.2.1) Genic capture explains the lek paradox

192  If higher-quality and fitter individuals are more likely than other individuals to produce higher-quality and

193 fitter descendants, there is more likely to be natural selection for a bandwagoning allele that induces lower-quality

194 individuals to forfeit resources. Historically, theory has predicted the heritability of fitness to be minimally low—

195 that is, that variation in fitness between individuals would be far less attributable to genetic differences than to

196 chance—because any alleles substantially contributing to fitness would be expected to gain fixation (thereby

197 reducing genetic variance) far more rapidly than mutations would replenish variation at their loci (Borgia, 1979;

198 Taylor & Williams, 1982). However, this prediction is evidently contradicted by the appearance in many species

199 (e.g., birds of paradise, fish, frogs, insects) of "leks," which are groups in which males congregate to display signals

200 of their quality to females (Kirkpatrick, 1982; Kirkpatrick & Ryan, 1991). Even though the lekking males

201 contribute little to the females besides sperm, the females often exhibit a preference for the same male and incur

202 costs to mate with that male, which suggests their preference occurs to ensure that their offspring have better genes

203 (Kirkpatrick & Ryan, 1991; Rowe & Houle, 1996). This apparent contradiction between theory and empirical

204 observations is known as the lek paradox (Borgia, 1979; Kirkpatrick & Ryan, 1991; Kotiaho et al., 2001; Taylor &

205 Williams, 1982).

206  A well-received resolution to the lek paradox has emerged over the past two decades. It holds that: 1) An

207 individual's condition is influenced by alleles at many different loci—enough loci to allow a quantity of mutations

208 from one generation to the next that maintains substantial levels of additive genetic variance for condition

209 (Pomiankowski & Moller, 1995; Rowe & Houle, 1996) and 2) An individual's condition is a determinant of that

210 individual's ability to exhibit displays for mates (e.g., ornaments, dances, songs); therefore, 3) These mate-attracting

211 displays "capture" the individual's condition, so that a variance in display characteristics owes to a variance in

212 alleles that contribute to condition. In recent years, scholars have also pointed to epigenetic marks as a source of

213 heritable differences in condition, particularly since epigenetic marks tend to be subject to a higher mutation rate

214 than genes (Bonilla et al., 2016).

215 **2.2.2) Values of fitness heritability tend to underestimate the likelihood of natural selection for bandwagoning**

216  The heritability of fitness has been reported to be significant but low—approximately 10% (reviewed in

217 Burt, 1995 and in Merila & Sheldon, 2000). However, natural selection for genetic bandwagoning may be more

218 likely than values of fitness heritability would suggest. Heritability of fitness is often calculated as a parent-

offspring regression: the covariance between parent fitness and offspring fitness divided by the variance of parent

fitness. Multiple factors can exert an influence that leads the offspring of high-quality individuals to be fitter than

their parents and/or the offspring of low-quality individuals to be less fit than their parents, which can lead

covariance between parent fitness and offspring fitness (and, therefore, heritability of fitness) to be lower than

otherwise, while also making natural selection for genetic bandwagoning more likely since these factors contribute

to greater lineage fitness advantages for fitter individuals. These factors include differential allocation, which is

reviewed above in section 2.1. Additionally, sexual selection can have the effect of reducing reproduction (in some

cases to nil) by lower-quality individuals that would otherwise have the capacity to reproduce, since, in sexual

selection, females exercise their choice to mate with higher-quality individuals. The occurrence of either sexual

selection or genetic bandwagoning makes the other more likely to occur: If a female is likely to refuse to mate with

a low-quality individual, a bandwagoning allele is more likely to be selected to induce forfeiture of resources by that

low-quality individual prior to reproductive age, since it is less likely to reproduce anyway; likewise, a female is

more likely to refuse to mate with a low-quality individual if it is likely that bandwagoning alleles located within

that individual's offspring will induce them to forfeit their resources. These lower-quality individuals that do not

reproduce because of sexual selection or genetic bandwagoning are not included as parents in parent-offspring

regression calculations for heritability of fitness since they produce no offspring; however, they are included in these

calculations as offspring and it is likely that the reductions in their fitness that result from sexual selection and

genetic bandwagoning lead their fitness to be more dissimilar to that of their parents, which leads the covariance

between parent fitness and offspring fitness (and, therefore, heritability of fitness) to be lower than otherwise. In

accordance with this notion, sexual selection and genetic bandwagoning exert a greater influence upon the fitness of

lower-quality individuals than upon the condition of lower-quality individuals, which may explain why heritability

values for condition are often substantially higher than heritability values for fitness (Birkhead et al., 2006; Kotiaho

et al., 2001; Parker & Garant, 2004), even though the heritability of fitness is believed to occur through the

"capture" of heritable condition (Pomiankowski & Moller, 1995; Rowe & Houle, 1996). In sum, differential

allocation and sexual selection can increase the likelihood of natural selection for genetic bandwagoning, yet

differential allocation and sexual selection, as well as genetic bandwagoning, can reduce heritability values for

fitness and differential allocation and genetic bandwagoning can reduce heritability values for condition;

consequently, reported values of heritability for fitness and condition can underestimate the likelihood that there is

natural selection for genetic bandwagoning.

2.3) An increasing difference between two lineages' expected quantities of descendants

In section 2.2, it was noted that heritability values for both quality and fitness are often significantly

positive. When the heritability of fitness is positive, the expected quantity of descendants of a fitter individual's

lineage, ceteris paribus, increases from generation to generation in comparison to the expected quantity of

descendants of a less fit individual's lineage. This is explained as follows. Suppose that one individual produces s

offspring and another individual produces r offspring, with $s > r$, and suppose that the fitter individual's higher

fitness is, at least, partly due to its "good genes." This means that the fitter individual passes at least part of its

fitness advantage to its offspring. Suppose that this heritable fitness advantage is such that the offspring of the fit

individual are expected to have a mean fitness of a and the offspring of the less fit individual are expected to have a

mean fitness of b, with $a > b$. Due to germ-line mutation and recombination, different offspring from the same

parents inherit different mutations and genetic combinations, some of which have a positive effect upon fitness and

some of which have a negative effect. Consequently, there is variation in the fitnesses of the offspring of either

individual about these mean offspring fitness values of, respectively, a and b and this variation is heritable, since it

is due to mutations and genetic combinations. (There may also be environmental sources of variation in fitness, but

the point here is that the variation is, at least, partly heritable and the following algebra concerns this heritable

variation.) The heritable variation in the fitnesses of the fitter individual's offspring can be modeled as s random

choices from a distribution that has a mean of a and the heritable variation in the fitnesses of the less fit individual's

offspring can be modeled as r random choices from a distribution that has a mean of b. Since the fitter individual

produces more offspring, $s > r$, and since the mean fitness of the fitter individual's offspring exceeds the mean

fitness of the less fit individual's offspring, $a > b$. Generally, if s random numbers are chosen from a distribution

that has a mean of a and r random numbers are chosen from the same form of distribution (e.g., normal, poisson)

that has a mean of b and the same standard deviation, and $s > r$, then the difference between the expected i^{th} highest

number chosen after s choices, E[Xis], and the mean of the distribution from which those s choices are drawn, a,

exceeds the difference between the expected i^{th} highest number chosen after r choices, E[Xir], and the mean of the

distribution from which those r choices are drawn, b: E[Xis]-a > E[Xir]-b. Rearrangement provides: E[Xis]-E[Xir]

> *a-b*. In other words, for every offspring produced by the less fit individual, there is an offspring produced by the

fitter individual that has an expected fitness that is higher by more than *a-b*. Since these distributions represent the

heritable variation of fitness, this higher fitness expectation is due to heritable factors and the heritable fitness

advantage that is passed by the fitter individual's offspring to the fitter individual's grandoffspring exceeds *a-b*,

which is the heritable fitness advantage that is passed by the fitter individual to its offspring. This serves to illustrate

that if the heritability of fitness is positive, the difference between two lineages' expected quantities of descendants,

ceteris paribus, increases as the projection becomes increasingly long-term. This difference between the two

lineages' expected quantities of descendants would continuously increase from one generation to another in an

infinite population or continuously approach an asymptote from one generation to another in a finite population.

3) Two ways of bandwagoning: Resonation and Reservation

Section 3 introduces two ways in which bandwagoning occurs: resonation and reservation. In addition,

section 3 offers detail about how resonation and reservation accomplish the two steps of bandwagoning: 1)

evaluating the holder's lineage fitness outlook and 2) inducing a holder with a low lineage fitness outlook to forfeit

resources. Section 3 builds up to a particular combination of resonation and reservation which is presented and

explained verbally in section 3.25 and modeled algebraically in section 3.3.

3.1) Resonation

In resonation, the two aspects of bandwagoning are accomplished sequentially: A bandwagoning allele 1)

evaluates its holder's lineage fitness outlook and 2) if this outlook is low relative to that of other individuals, it

induces its holder to forfeit resources. The term "resonation" is chosen to represent this way of bandwagoning

because it describes how the bandwagoning allele responds to an evaluation of its holder's lineage fitness outlook

with a direct influence upon the holder's lineage fitness outlook that resonates with the evaluation: An already-low

lineage fitness outlook is the catalyst for the bandwagoning allele to directly make that lineage fitness outlook even

lower.

3.1.1) How a bandwagoning allele can evaluate its holder's lineage fitness outlook

Through various types of interactions with others, individuals receive information about their quality

relative to conspecifics. A bandwagoning allele within an individual would evaluate the lineage fitness outlook of

its holder by using information from these interactions directly or by using its holder's hormonal responses to these

interactions. Examples follow.

Parents evaluate the fitness outlooks of their offspring in order to determine how to allocate resources to

their offspring in a manner that maximizes their "return on investment" (Harris & Uller, 2009; Sheldon, 2000).

Consequently, the amount of resources and nurturing that an individual receives from its parents are indicators of

that individual's lineage fitness outlook compared to its siblings. Likewise, both the "good genes" (Andersson,

1982; Byers & Waits, 2006; Iwasa & Pomiankowski, 1994; Moller & Alatalo, 1999) and "sexy son" (Fisher, 1930;

Weatherhead & Robertson, 1981) hypotheses of sexual selection assume that individuals evaluate the fitness

outlooks of potential mates in order to determine whether the production of offspring with them would constitute a

good investment (Kokko et al., 2002). Therefore, the amount of positive attention an individual receives from

potential mates constitutes an indicator regarding how that individual's lineage fitness outlook compares to that of

conspecifics. Still another potential indicator of an individual's lineage fitness outlook involves direct competition,

such as a competitive bout. Performance in competitive bouts determines dominance, which is often heritable (Boag

& Alway, 1981; Dewsbury, 1990; Moore, 1990) and covaries with reproductive success (Andersson, 1994).

Dominance is also often reflective of other aspects of a lineage fitness outlook (e.g., immunocompetence: reviewed

in Georgiev et al., 2015), which supports the notion that a lineage fitness outlook determines dominance through an

influence upon performance in competitive bouts. Additionally, individuals can directly compare the resources

(e.g., territory) they have managed to obtain with those of other individuals. Individuals can also compare their

quality to others via the exchange of honest signals (Andersson, 1994)—for example, honest signals of levels of

stress incurred in development—which serves as another means of evaluating their lineage fitness outlook.

Many of the interactions mentioned above engender hormonal reactions that a bandwagoning allele can

potentially use in order to evaluate the lineage fitness outlook of its holder. For example, in many species,

individuals increase testosterone production in response to indications that they compare favorably to conspecifics,

whether this is indicated by interactions with parents (Gil et al., 1999; Stiver & Alonzo, 2009), potential mates

(Archer, 2006; Gleason et al., 2009; Oliveira, 2004), or rivals (Archer, 2006; Gleason et al., 2009; Mazur & Lamb,

1980; McCaul et al., 1992; Oliveira, 2004), which suggests a bandwagoning allele may be able to use the quantity

and/or timing of testosterone production as a cue in evaluating its holder's lineage fitness outlook.

3.1.2) How a bandwagoning allele can induce its holder to forfeit resources

If a bandwagoning allele evaluates its holder to have a low lineage fitness outlook, it then induces its holder

to forfeit resources. This can involve the suicide of the individual, which would result in a complete relinquishment

of the resources to which the individual has access and would have access in the future. However, even if an

individual's lineage fitness outlook is low compared to others, there is the possibility that it can change as a result

of, for example, the deaths of other individuals or a change in its environment. If an individual's lineage fitness

outlook is more likely to change, a bandwagoning allele is more likely to induce its holder to forfeit resources in a

manner that allows the holder to stay alive and/or allows the holder's offspring (if any) to stay alive while still

releasing some resources to nearby individuals that might have a more favorable lineage fitness outlook. One way

of accomplishing this is to impose upon each holder, irrespective of that holder's quality, an honest signal of that

holder's quality. The benefit from an honest signal of quality is correlated with the signaler's quality, so that honest

signalers of high quality are favorably influenced by the signal, but honest signals can cost low-quality signalers

food, territory, or mates. The food, territory, and mates are then utilized by nearby individuals. Thus, the

imposition of an honest signal of a holder's quality, irrespective of that holder's quality, is a way of inducing a

forfeiture of resources from lower-quality individuals while sparing higher-quality individuals, but also allowing

lower-quality individuals to remain alive in case any of their fitness outlooks change. Additionally, the inducement

of depression, self-handicapping, or a physical ailment upon a lower-quality individual can lead that individual to

forfeit resources it might otherwise utilize, but also leave the individual alive. Other ways in which an individual

with a low lineage fitness outlook can be induced to forfeit resources but remain alive include a cessation of the

functioning of the reproductive apparatus, a reduced interest in pursuing mates, and a diminishment in sexually-

selected characteristics. This point of allowing an individual with a low lineage fitness outlook to remain alive in

case this outlook changes also goes for that individual's lineage. Therefore, there may be natural selection for a

bandwagoning allele that allows an individual with a low lineage fitness outlook to produce some quantity of

offspring that is lower than the maximum quantity of offspring or grandoffspring that it could expect to support.

This would allow the individual's lineage to remain alive for another generation or two in case the lineage's fitness

outlook changes, yet it would also free up resources to be used by lineages with a higher fitness outlook.

Bandwagoning of this form might involve, for example, inducing an impairment in sperm competition or a

diminished allocation of food or nurturing to extant offspring.

3.2) Reservation

3.2.1) The problem: Difficulty encountered in evaluating a holder's lineage fitness outlook

In preceding sections, it has been explained that a bandwagoning allele evaluates the lineage fitness outlook

of the individual in which it is located and induces that individual to forfeit resources if that outlook is low. In

resonation, a bandwagoning allele accomplishes the two steps of bandwagoning—evaluation and forfeiture—

sequentially. However, there are heritable abilities that are relevant to an individual's lineage fitness outlook but

that are sometimes difficult to measure even for the individual itself (or for a bandwagoning allele within the

individual), which poses a challenge for the bandwagoning allele in its attempt to accomplish the first step of

bandwagoning—that of evaluating its holder's lineage fitness outlook. The dilemma is clarified with the following

discussion.

As a population coevolves with its natural enemies (e.g., predators, parasites, prey), there may be wide

intergenerational variance in the quantity of individuals in the population that are able to survive and reproduce. In

some generations, many individuals may be able to survive and reproduce. In other generations, a dearth of

available prey or a proliferation of well-adapted predators or parasites may mean that only a few individuals survive

to reproductive age. Individuals have a more favorable lineage fitness outlook if they have genes that can allow

survival when prey are not plentiful and/or when there is a prevalence of well-adapted predators or parasites, though

it can be difficult to determine which individuals have these "good genes" during a generation in which conditions

are unchallenging and numerous individuals can survive to reproductive age.

In particular, it can be difficult to determine which of numerous living individuals are better at surviving

predators and/or parasites, since it is frequently the case that an individual has either perished or has survived

without injury from predators or parasites. This binary manner of classifying an individual's success at surviving

predators and/or parasites differs from that of acquiring prey and/or territory, which is more given to representation

with a continuum, since an individual that occasionally fails at either prey or territory acquisition is more likely to

remain alive than an individual that fails against predators or parasites. As the life-dinner principle goes, a predator

that is unsuccessful loses its dinner, while a prey that is unsuccessful loses its life. In other words, an individual is

more likely to be able to determine, with honest signaling, whether the quantity of meals it has missed is more or

less than that of another individual than whether the quantity of predator encounters it has failed is more or less than

that of the other individual, since it is more likely that both individuals have survived all predator encounters without

injury than it is that both have never missed a meal.

Even if two individuals can survive to reproductive age in unchallenging conditions, one individual's

descendants may be less likely than the other individual's descendants to survive to reproductive age. One reason

this may occur is that the former individual's survival is more attributable to luck. Perhaps that individual would

have been captured by numerous predators that, instead, pursued other prey or would have been killed by several

parasite strains but was, instead, exposed to weaker ones, while the latter individual has genes and/or epigenetic

marks that allow it the speed, strength, mimicry or camouflage (or other factors) to survive many of the natural

enemies to which the former individual would have succumbed. Another possible reason is that, even if the two

individuals are exposed to largely the same natural enemies, the former individual may be able to survive by a lesser

margin than the latter individual. That is, some natural enemies may be nearly fast enough, strong enough, or

otherwise well-adapted enough to successfully predate or parasitize the former individual, but not so nearly the latter

individual because of genes and/or epigenetic marks held by the latter individual. Even though both individuals may

be able to survive to reproductive age, descendants of the predators and parasites that are better-adapted than their

predecessors are more likely to kill descendants of the former individual than those of the latter individual. Thus,

two reasons have been identified for why one individual may be able survive to reproductive age as well as another

individual, even though its descendants are less likely to survive and reproduce than descendants of the other: 1) the

former individual's luck of exposure to less well-adapted natural enemies and 2) the former individual's lower

margin of survival against the natural enemies that both faced. Hamilton's rule might be satisfied by a forfeiture of

resources from the former individual to the latter individual due to the higher lineage fitness outlook of the latter

individual, except that neither individual knows which has "better" genes for surviving the natural enemies that

afflict the population because both individuals are able to survive to reproductive age.

3.2.2) A solution to the problem of evaluating a holder's lineage fitness outlook: Reservation

A bandwagoning allele can solve this dilemma of evaluating its holder's lineage fitness outlook by

inducing its holder to act in a manner that detracts from its ability to survive predators and/or parasites. That is, the

bandwagoning allele can reserve (i.e., withhold) some of its holder's capacity to survive one or more of these natural

enemies. In section 3.2.1, it was argued that one surviving individual's descendants are less likely to survive to

reproductive age than those of another surviving individual if: 1) the former individual's survival was attributable to

a lack of exposure to more well-adapted predators or parasites in the population that would have killed it but the

latter individual's survival was less attributable to luck and could have survived more of the natural enemies that

would have killed the former individual or 2) there were natural enemies that were nearly well-adapted enough to

have killed the former individual but the latter individual was able to survive more securely. The reservation

induced by a bandwagoning allele can serve to reduce the likelihood of both of these two causes of the survival of a

less well-adapted individual. That is, a bandwagoning allele can reduce the likelihood of the first cause by

increasing its holder's exposure to natural enemies, which diminishes the likelihood that there are well-adapted

natural enemies to which the individual is not exposed. For example, it can induce its holder to position itself in

close proximity to predators, to make conspicuous displays or loud calls, or to eschew certain forms of mimicry or

camouflage, all of which would increase its holder's exposure to natural enemies. Furthermore, a bandwagoning

allele can reduce the likelihood of the second cause by inhibiting its holder's ability to survive the natural enemies to

which the holder is exposed, which leads to the perishing of individuals that, without reservation, would be able to

survive by a low margin. For example, it can induce its holder to do such things as exhibit weighty ornaments or

produce lower quantities of antibodies than otherwise, which would impair its holder's ability to survive the natural

enemies to which it is exposed.

If a holder can survive in spite of this reservation, it is more assured of being able to produce descendants

that will be able to survive to reproductive age during conditions that are more severe. Alternatively, a holder may

perish or become sick as a result of this reservation, in which case it would forfeit some or all of its resources to

nearby individuals. That is, if a holder perishes or becomes sick as a result of reservation, the two steps of

bandwagoning are accomplished simultaneously: The identification of the holder's poor lineage fitness outlook and

the forfeiture of that holder's resources both occur with the holder's perishing or sickness.

A reader may wonder about the purpose of inducing an individual to perish in order to forfeit resources to

individuals with "better genes" when that individual's descendants would forfeit their resources to the descendants

with "better genes" when they perish: If the resources are going to the descendants with "better genes" anyway,

what's the purpose of inducing reservation in order to forfeit the resources a generation or two beforehand? The

answer is that there is no guarantee that an individual with "better genes" will produce any surviving descendants, as

some of the population's predators or parasites could evolve to be so well-adapted that the entire population is killed

prior to reproductive age. However, the more descendants that are produced by individuals with "better genes," the

more likely it is that at least one will be able to survive to reproductive age. Likewise, the more descendants

produced by "better genes" individuals that hold the bandwagoning allele, the more likely the bandwagoning allele

gains fixation by being located within an individual capable of surviving a very well-adapted natural enemy

population.

Natural selection would act on multiple aspects of this reservation, including the natural enemies against

which the reservation occurs and the degree to which it occurs. A bandwagoning allele should not be expected to

evolve to induce reservation of its ability against all of its holder's natural enemies. If there are seldom generations

in which a particular natural enemy population evolves in such a manner that it kills most of a population's

individuals, it is unlikely that a bandwagoning allele in that population would evolve to induce its holders to reserve

from its ability against that natural enemy population, since lineages of holders that survive the reservation are

unlikely to have a fitness advantage in ensuing generations.

3.2.3) The possibility of adjusting the degree of reservation to the severity of the threat

A bandwagoning allele that induces its holder to reserve too little runs the risk of allowing its holder to

survive and reproduce, even though its genes may not be "good" enough to allow its offspring or grandoffspring to

survive against better-adapted natural enemies. Alternatively, a bandwagoning allele that induces too much

reservation runs the risk of inducing the death of every holder of a bandwagoning allele in the population, as even

the individuals with the "better genes" against the natural enemies would not be able to survive the reservation,

whereupon there would be no individuals with a bandwagoning allele left to utilize resources forfeited by the

perished individuals. However, what constitutes an appropriate degree of reservation suitable for the satisfaction of

Hamilton's rule may vary from generation to generation. As noted in section 3.2.1, in the absence of reservation,

the quantity of individuals in the population that perish prior to reproduction may vary widely from one generation

to another. During some generations, many individuals may be able to survive to reproductive age. During other

generations, there may be only a few individuals that can survive, owing to the proliferation of well-adapted

predator or parasite individuals. The more individuals that have perished during a generation and the greater alacrity

with which they have perished, the more severe the conditions that are indicated and, consequently, the less an

individual would have to be induced to reserve in order to demonstrate it has "good genes." If few individuals

remain alive, it is less likely that the average individual that would utilize the forfeited resources is substantially

more capable at evading the natural enemies than the forfeiting individual would be. Consequently, it is less likely

that Hamilton's rule would be satisfied by the forfeiture. Natural selection would, therefore, act upon a

bandwagoning allele to vary the degree of reservation it induces during a generation or a point in a generation, such

that it would reserve more if many individuals have been able to survive and it would reserve less if many

individuals have perished during the generation or if numerous individuals have rapidly perished within a short time

period.

A problem with this manner of calibrating the degree of reservation to the threat presented by a natural

enemy is that the threat presented by predators or parasites can escalate with little warning. A predator or parasite

population can evolve during what constitutes a generation for, respectively, a prey population or a host population.

Indeed, a parasite epidemic is often apparent only in retrospect. Therefore, a bandwagoning allele that induces all of

its holders to engage in reservation at the same time and to the same degree runs the risk of leading to the death of

all of its holders if the predator or parasite threat rapidly intensifies. If this were to occur, there would be no

bandwagoning allele holders left to utilize the resources forfeited by the perished individuals. Therefore, it is likely

that there would be natural selection for a bandwagoning allele that hedges against the possibility that conditions

rapidly deteriorate by varying the timing and degree of the reservation it induces its holders to undertake.

3.2.4) Varying the timing and degree of reservation

A bandwagoning allele that hedges in this manner may vary the timing and degree of the reservation it

induces depending upon certain characteristics of its holder, such as, for example, whether the holder is male or

female or whether the holder is a first-born or later-born individual. A hedge based upon the timing of reservation

may involve a bandwagoning allele that induces holders with one characteristic to reserve during a particular span in

its life history and holders with another characteristic to reserve during another span in its life history. A hedge

based upon the degree of reservation may involve a bandwagoning allele that induces holders with one characteristic

to reserve more over the totality of its life than holders with another characteristic. In a model offered introduced in

section 3.2.5, a bandwagoning allele varies the degree of reservation based upon indications about the holder's

quality.

3.2.5) Hedging the degree of reservation based upon a tentative evaluation of a holder's lineage fitness

outlook

Prior sections have included discussion of the premise that at any particular time during a generation, an

individual (and a bandwagoning allele within the individual) has partial but incomplete information about that

individual's lineage fitness outlook. It was explained in section 3.2.1 that an individual (and a bandwagoning allele

within the individual) is more likely to know how its ability to procure food or territory compares with that of other

individuals than how its ability to survive predators or parasites compares with that of other individuals.

Reservation has been presented as a way that can be used by a bandwagoning allele to identify its holder's abilities

in areas that are difficult to evaluate when conditions are unchallenging (e.g., its holder's ability to survive against

predators or parasites) in order to garner greater certainty about the holder's lineage fitness outlook. Discussion in

section 3.2.4 focused upon the role of varying the timing and degree of reservation based on the holder's particular

characteristics. The present section 3.2.5 considers a bandwagoning allele that varies the degree of reservation it

induces depending upon the information about the holder's lineage fitness outlook that it has already received.

Specifically, holders that have received information indicative of a favorable lineage fitness outlook are induced to

reserve more over the course of their lives and individuals that have received information indicative of a less

favorable lineage fitness outlook are induced to reserve less over the course of their lives. This information

regarding the holder's lineage fitness outlook can come in the forms reviewed in section 3.1.1. That is, individuals

that have, for example, been preferred more by parents and mates, acquired more food and territory, and performed

better in competitive bouts are induced to reserve more against a natural enemy. In accordance with the discussion

in section 3.2.1, it is less likely that individuals would receive information about their ability to survive against

parasites or predators; however, any individuals that are sickened by parasites or wounded by predators but remain

alive are induced to reserve less against a natural enemy. Natural selection would act upon a bandwagoning allele to

vary reservation in this manner for the following reasons.

If the information available suggests that a holder's lineage fitness outlook is high—that is, if a holder fares

well in comparison to other individuals in, for example, food, territory, and mate acquisition—a bandwagoning

allele would induce its holder to reserve against one or more natural enemies in order to determine whether the

holder has high ability against these natural enemies, in addition to the other abilities that have led to the favorable

outlook; if so, its outlook can be evaluated to be high with greater certainty. However, the more poorly a holder

compares to other individuals in food, territory, and mate acquisition, the more likely it is that Hamilton's rule

would be satisfied by the holder's forfeiture of resources if the bandwagoning allele's information about its holder's

lineage fitness outlook were more complete—if, for example, the holder's abilities against predators and parasites

were known in comparison to other individuals. This is because the lower a holder's abilities to acquire food,

territory, and mates compared to others, the more likely it is that there are numerous individuals that have both better

abilities to acquire food, territory, and mates and better abilities against predators and parasites and, therefore, the

more likely it is that Hamilton's rule would be satisfied if the holder were to forfeit its resources because there

would be enough individuals utilizing the resources that have a more favorable lineage fitness outlook and a copy of

the bandwagoning allele. In fact, if the tentative evaluation is that the individual's lineage fitness outlook is very

low, the best chance that Hamilton's rule would not be satisfied by a total forfeiture of that individual's reproduction

is likely to be if severe conditions suddenly ensue, characterized by a newly well-adapted natural enemy population,

such that many individuals that formerly had a more favorable lineage fitness outlook and were induced to engage in

reservation were to perish due to the reservation. However, if the holder with a low lineage fitness outlook were to

have reserved as much as other individuals and these severe conditions were to have occurred, the holder would

have likely been among the perished due to its reservation. In other words, a holder with a very low lineage fitness

outlook will probably satisfy Hamilton's rule by forfeiting resources even if they can survive in spite of reservation,

unless conditions are severe enough to kill most (or all) reserving individuals, but if they were to reserve, they would

also probably perish. That is, reservation is probably pointless for individuals with a very low lineage fitness

outlook, but if they do not undertake reservation, they have a chance of surviving the severe conditions that would

warrant their reproduction (that is, conditions that would mean their forfeitures would not satisfy Hamilton's rule).

Therefore, the lower a holder's lineage fitness outlook, the less the bandwagoning allele induces that holder to

reserve. Instead, the bandwagoning allele induces a holder with a low lineage fitness outlook to forfeit some

resources but also allows the holder to remain alive in case the holder's lineage fitness outlook changes. This is in

accordance with the observation in section 3.1.2: If an individual's lineage fitness outlook is more likely to change,

there is more likely to be natural selection for a bandwagoning allele that induces its holder to forfeit resources in a

manner that allows the holder to stay alive and/or allows the holder's offspring (if any) to stay alive while still

releasing some resources to nearby individuals that might have a more favorable lineage fitness outlook.

In sum, for a bandwagoning allele located within a holder that has had interactions that suggest a high

lineage fitness outlook, the holder reserves and, if it survives, it reproduces; however, for a bandwagoning allele

located within a holder that has had interactions that suggest a low lineage fitness outlook, it does not induce the

holder to reserve as much and instead induces the holder to forfeit some resources but allows the holder to remain

alive in case its lineage fitness outlook changes. Figure 1 is a flowchart of the actions that a bandwagoning allele

would take if it were to engage in bandwagoning of the form introduced in the present section 3.2.5.

Figure 1. Flowchart that represents how a bandwagoning allele acts according to the model introduced in section 3.

A bandwagoning allele that acts in this manner can be said to combine resonation and reservation: If its

holder is suggested to have a high lineage fitness outlook by the information that has already accrued to the allele,

the allele induces reservation; if its holder is suggested to have a low lineage fitness outlook by the information that

has already accrued to the allele, the allele induces its holder to refrain from reproduction. In the interest of offering

further clarity, a bandwagoning allele that combines resonation and reservation in this manner introduced in the

present section is algebraically modeled in section 3.3, which serves to illustrate why a bandwagoning allele that

combines resonation and reservation in this manner would be selected in favor of a bandwagoning allele that solely

practices resonation.

3.3) Model

The model in this section serves to demonstrate why there would be natural selection for one type of

bandwagoning allele, one which does both resonation and reservation in the manner described in section 3.2.5, at the

expense of another type of bandwagoning allele, one which only does resonation. Therefore, in this model, there are

two bandwagoning alleles in the host population: an R_R allele which does both resonation and reservation in the

manner described in section 3.2.5 and an R allele which does only resonation.

Each individual in the host population is closer than any other individual to one unit of resources. This

resource unit is intended to represent food, territory, or mates that an individual could use in order to produce

offspring or, alternatively, that an individual could forfeit, in which case it is used for reproduction by other

individuals. If the individual forfeits its unit, it is divided for reproduction by other individuals according to the

assortment of the population. In modeling the assortment of a population, it is common to suppose, without

specifying how the assortment arises in three-dimensional space, that an individual interacts with a same-type

individual with probability r and with a random member of the population (including same-type individuals) with

probability 1-r (Allen & Nowak, 2015; Cooney et al., 2016; Grafen, 1979; Okasha & Marens, 2016; Van Cleve &

Akcay, 2014). Analogously, it is supposed in the present model that if an individual does not use its resource unit,

an r fraction of that unit is used by individuals with the same allele and a 1-r fraction is divided evenly among the

entire population (including individuals with the same allele).

Individuals in the host population differ in two dimensions of heritable ability: A_R ability, which refers to an

individual's ability to resist parasites, and A_O ability, which refers to an individual's ability in all other aspects (e.g.,

obtaining food, territory, and mates and avoiding predators). Each individual in the population has either high or

low A_R ability and high or low A_O ability. An allele within a host individual knows whether the host individual has

high or low A_O ability but does not know whether the host has high or low A_R ability unless, or until, the host does

reservation. An individual with high A_R ability survives in spite of reservation, but an individual with low A_R ability

cannot; consequently, reservation can be used to identify individuals with high A_R ability because an individual that

survives in spite of reservation is guaranteed to have high A_R ability.

Additionally, the parasite threat in the population during generation t is either T_M or T_S. A T_M threat means

that half of the individuals in the population have high A_R ability and half have low A_R ability, while a T_S threat

means that there are no individuals in the population with high A_R ability. Since a holder with high A_R ability can

survive in spite of reservation but a holder with low A_R ability cannot, this means that if the threat is T_M, half of the

holders that reserve can survive in spite of reservation, but if the threat is T_S, no holders can survive reservation.

This feature of a T_S threat will help illustrate the reason a bandwagoning allele varies the degree of reservation it

induces.

Suppose that if an individual reproduces in generation t, the quantity of descendants of the same type that it

can expect to have by some future generation $t+x$ (where $x > 1$) for every unit of resources it uses to reproduce in

generation t varies according to the individual's heritable ability, as described in Table 1. It is assumed that both

high A_O ability and high A_R ability lead to greater quantities of descendants by a future generation $t+x$ than,

respectively, low A_O ability and low A_R ability; therefore, $C > D > F$ and $C > E > F$.

Heritable ability	Descendants by generation $t+x$ per unit used in generation t
High A_O ability and high A_R ability:	C
High A_O ability and low A_R ability:	D
Low A_O ability and high A_R ability:	E
Low A_O ability and low A_R ability:	F

Table 1. The descendants by generation t+x that individuals with each combination of ability can expect per unit of resources

used in generation t.

The difference between the two types of bandwagoning alleles in the host population is that an R allele

induces resonation only: If a holder of an R allele has high A_O ability, it reproduces; if it has low A_O ability, it does

not reproduce. An R_R allele induces both resonation and reservation: If a holder of an R_R allele has high A_O

ability, it reserves and only if it survives in spite of reservation does it reproduce; if a holder has low A_O ability, it

does not reserve and it does not reproduce unless some of the individuals with high A_O ability have perished.

At generation t, there are $4A$ holders of the R_R allele and $4B$ holders of the R allele. Half of the holders of

each allele have high A_O quality and half have low A_O quality. With probability G, the parasite threat in generation t

is T_M and with probability $1\text{-}G$, the parasite threat in generation t is T_S. If the threat is T_M, half of each allele's

holders of both high A_O ability and low A_O ability have high A_R ability and half have low A_R ability. If the threat is

T_S, there are no holders of either allele with high A_R ability, as all holders of each allele have low A_R ability.

3.3.1) The quantities of descendants expected by generation $t+x$ if the threat in generation t is T_M

When the threat is T_M, there are A R_R holders and B R holders each with: 1) high A_O ability and high A_R

ability; 2) high A_O ability and low A_R ability; 3) low A_O ability and high A_R ability; and 4) low A_O ability and low A_R

ability. When the threat is T_M, the only R_R holders to reproduce are those with high A_O ability and high A_R ability.

This is because when the threat is T_M, the A R_R holders with high A_O ability but low A_R ability perish due to

reservation and all $2A$ R_R holders with low A_O ability refrain from reproducing due to resonation. Since three-

fourths of the $4A$ R_R holders do not use their resource units for reproduction, the total resource units forfeited by

R_R holders is $3A$. By contrast, R holders reproduce if they have high A_O ability, whether they have high or low A_R

ability. The one-half of the $4B$ R holders with low A_O ability forfeit their resources due to resonation, so the total

resource units forfeited by R holders is $2B$.

Consequently, if the threat during generation t is T_M, R_R holders can expect a quantity of descendants by

generation $t+x$ that is given by (3), which is calculated as the quantity of descendants the reproducing R_R holders

can expect per resource unit times the resource units they use. When the threat is T_M, the only R_R holders to

reproduce are those with high A_O ability and high A_R ability. According to Table 1, these holders can expect C

descendants per resource unit. The resource units they use are their own resource units, A; plus r times the resource

units forfeited by other R_R holders, $3A$; plus $(1\text{-}r)$ times the total resource units forfeited by both R_R and R

holders, $3A+2B$, times the proportion of reproducing individuals that are R_R holders, $A/(A+2B)$.

$C[A+r3A+(1\text{-}r)(3A+2B)A/(A+2B)]$ (3)

If the threat during generation t is T_M, R holders can expect a quantity of descendants by generation $t+x$ that

is given by (4). The reproducing R holders with high A_O ability and high A_R ability can expect C descendants per

resource unit and the reproducing R holders with high A_O ability and low A_R ability can expect D descendants per

resource unit. The resource units available for R holders are their own resource units, $2B$; plus r times the resource

units forfeited by other R holders, $2B$; plus $(1-r)$ times the total resource units forfeited by both R_R and R holders,

$3A+2B$, times the proportion of reproducing individuals that are R holders, $2B/(A+2B)$. Since the quantity of R

holders with high A_O ability and high A_R ability is equal to the quantity of R holders with high A_O ability and high A_R

ability, half of the resource units available for R holders are multiplied by C and half are multiplied by D.

$(C+D)[B+rB+(1-r)(3A+2B)B/(A+2B)]$ (4)

3.3.2) The quantities of descendants expected by generation $t+x$ if the threat in generation t is T_S

If the threat is T_S, there are no holders of either allele that have high A_R ability (that is, there are no holders

of either allele that have the ability to survive reservation). Instead, there are $2A$ R_R holders and $2B$ R holders that

have high A_O ability and low A_R ability and $2A$ R_R holders and $2B$ R holders that have low A_O ability and low A_R

ability.

The R_R holders with high A_O ability and low A_R ability perish due to reservation. For the R_R allele, the

perishing of individuals with high A_O ability is the criterion for allowing reproduction by holders with low A_O

ability. If the threat is T_S, all of the R_R holders with high A_O ability perish; consequently, R_R holders with low A_O

ability reproduce. Since one-half of the $4A$ R_R holders forfeit their resource units, the total resource units forfeited

by R_R holders is $2A$. Alternatively, R holders reproduce if they have high A_O ability and there is no reproduction

by R holders with low A_O ability. The one-half of the $4B$ R holders with low A_O ability forfeit their resources due to

resonation, so the total resource units forfeited by R holders is $2B$.

If the threat during generation t is T_S, R_R holders can expect a quantity of descendants by generation $t+x$

that is given by (5). Since the reproducing R_R holders have low A_O ability and low A_R ability, they can expect F

descendants per resource unit. The resource units they use are their own resource units, $2A$; plus r times the

resource units forfeited by other R_R holders, $2A$; plus $(1-r)$ times the total resource units forfeited by both R_R and

R holders, $2A+2B$, times the proportion of reproducing individuals that are R_R holders, $2A/(2A+2B)$.

$F[2A+r2A+(1-r)(2A+2B)2A/(2A+2B)]$ (5)

If the threat during generation t is T_S, R holders can expect a quantity of descendants by generation $t+x$ that

is given by (6). Since the reproducing R holders have high A_O ability and low A_R ability, they can expect D

descendants per resource unit. The resource units they use are their own resource units, $2B$; plus, r times the

resource units forfeited by other R holders, $2B$; plus $(1-r)$ times the total resource units forfeited by both R_R and R

holders, $2A+2B$, times the proportion of reproducing individuals that are R holders, $2B/(2A+2B)$.

$D[2B+r2B+(1-r)(2A+2B)2B/(2A+2B)]$ (6)

3.3.3) The quantities of descendants expected by generation $t+x$

The expected quantity of descendants for R_R holders by generation $t+x$ (7) is the expected quantity of

descendants for R_R holders by generation $t+x$ if the threat in generation t is T_M, which is given by (3), times the

probability that the threat is T_M, G, plus the expected quantity of descendants for R_R holders by generation $t+x$ if

the threat in generation t is T_S, which is given by (5), times the probability that the threat is T_S, $(1-G)$.

$GC[A+r3A+(1-r)(3A+2B)A/(A+2B)]+(1-G)F[2A+r2A+(1-r)(2A+2B)2A/(2A+2B)]$ (7)

The expected quantity of descendants for R holders by generation $t+x$ (8) is the expected quantity of

descendants for R holders by generation $t+x$ if the threat in generation t is T_M, which is given by (4), times the

probability that the threat is T_M, G, plus the expected quantity of descendants for R holders by generation $t+x$ if the

threat in generation t is T_S, which is given by (6), times the probability that the threat is T_S, $(1-G)$.

$G(C+D)[B+rB+(1-r)(3A+2B)B/(A+2B)]+(1-G)D[2B+r2B+(1-r)(2A+2B)2B/(2A+2B)]$ (8)

The ratio of (7) to (8) represents the ratio of the expected quantity of descendants for R_R holders by

generation $t+x$ to the expected quantity of descendants for R holders by generation $t+x$. The R_R allele gains in

frequency between generation t and generation $t+x$ if the ratio of (7) to (8) exceeds the ratio of the quantity of R_R

holders at generation t to the quantity of R holders at generation t. The quantity of R_R holders at generation t is $4A$

and the quantity of R holders at generation t is $4B$, so the ratio of the quantity of R_R holders at generation t to the

quantity of R holders at generation t is $A{:}B$. This ratio of $A{:}B$ is equivalent to each of: a) the ratio of the first term

inside the brackets in the first half of (7) to its counterpart of (8); b) the ratio of the third term inside the brackets in

the first half of (7) to its counterpart of (8); and c) the ratio of the entire value inside the brackets in the second half

of (7) to its counterpart of (8). Since $D > 0$, this means that the ratio of GC times the first term inside the brackets in

the first half of (7) to $G(C+D)$ times its counterpart of (8) is exceeded by $A{:}B$. Additionally, this means that the ratio

of GC times the third term inside the brackets in the first half of (7) to $G(C+D)$ times its counterpart of (8) is

exceeded by $A{:}B$. Additionally, since $D > F$, the ratio of $(1{-}G)F$ times the entire value inside the brackets in the

second half of (7) to $(1{-}G)D$ times its counterpart of (8) is exceeded by $A{:}B$. That is, each of these terms in (7), in

comparison to their counterparts in (8), would, on their own, contribute to a loss of frequency for the R_R allele

between generation t and generation $t{+}x$. The only way the R_R allele can gain frequency between generation t and

generation $t{+}x$ is if this loss is made up by the difference between GC times the second term inside the brackets in

the first half of (7), $r3A$, minus $G(C+D)$ times the second term inside the brackets in the first half of (8), rB. In other

words, what is responsible for any expected increase in frequency for the R_R allele between generation t and

generation $t{+}x$ is that R_R holders in the aggregate forfeit a higher percentage of the resources to which they are

close ($3A$ by R_R holders versus $2B$ by R holders) and, of these resources, $r3A$ are used by R_R holders with high A_O

ability and high A_R ability, whereas only rB are used by R holders with high A_O ability and high A_R ability (since rB

are also used by R holders with high A_O ability and low A_R ability); consequently, the fraction of resources close to

R_R holders that are used by R_R holders with high A_O ability and high A_R ability is triple the fraction of resources

close to R holders that are used by R holders with high A_O ability and high A_R ability.

The second halves of (7) and (8) give the likelihood that the parasite threat during generation t is T_S times

the expected quantities of descendants by generation $t{+}x$ of, respectively, R_R holders and R holders if the parasite

threat during generation t is T_S. Since the ratio of the second half of (7) to its counterpart of (8) is exceeded by $A{:}B$,

a higher likelihood that the parasite threat during generation t is T_S means a lower frequency of the R_R allele at

generation $t{+}x$. This makes sense: When reservation occurs during a generation in which there is a threat that is

sufficiently considerable that no individual in the population is able to survive in spite of reservation, it has a

negative effect upon the frequency of the bandwagoning allele that practices it because there are no holders of the

allele left to utilize the resources forfeited by the perished individuals.

This model makes the simplifying assumptions that an individual either reproduces to capacity or

completely forfeits reproduction and that an individual can delay reproduction (that is, an individual with high A_O

ability can delay reproduction until it has survived reservation and an individual with low A_O ability can delay

reproduction until it determines whether individuals with high A_R ability have survived reservation or perished

because of it) and still reproduce to full capacity. In a more realistic scenario, an individual that delays reproduction

may forego reproductive opportunities and may, therefore, reproduce less than it could have, once the

bandwagoning allele that it holds has determined that it should reproduce.

3.4) Parallels between the model and established theory

A model for the natural selection of a bandwagoning allele that practices both resonation and reservation is

represented verbally in section 3.2.5 and algebraically in section 3.3. For the purposes of conferring both clarity and

credibility to this model, it is worthwhile to note parallels between it and both the handicap principle (Getty, 2006;

Grafen, 1990; Szamado, 2011; Zahavi, 1975) and bet-hedging theory (Gillespie, 1973, 1974; Schreiber, 2015; Seger

& Brockmann, 1987; Slatkin, 1974; Starrfelt & Kokko, 2012).

Zahavi (1975) introduced the handicap principle to explain exhibitions that apparently impair their

exhibitor, such as loud songs, conspicuous displays, and colorful plumage. Zahavi argued that individuals engage in

these costly exhibitions (i.e., handicaps) in order to demonstrate their superior genes to potential mates. He reasoned

that individuals with superior genes tend to be in better condition than others and are, therefore, capable of surviving

a larger handicap; consequently, individuals that maintain a larger handicap do so to demonstrate their superior

genes. While Zahavi's original examples focused on individuals that impair their ability to evade predators for the

purpose of signaling their quality to potential mates, the handicap principle has since been developed further (Getty,

2006; Grafen, 1990; Szamado, 2011) and has been applied across the animal and plant kingdoms to explain

ostensibly avoidable costs incurred by individuals that impair their abilities against parasites, prey, and conspecifics

(Folstad & Karter, 1992; Johnstone, 1995; Zahavi & Zahavi, 1997). Later applications of the handicap principle

have involved the demonstration of ability that may or may not owe to "good genes," which is a departure from

Zahavi's original formulation that emphasized heritable ability, though it is a general feature of the handicap

principle that costs are incurred by an individual in order to demonstrate that individual's ability.

The model of bandwagoning described in sections 3.2.5 and 3.3 shares numerous properties with the

handicap principle. Both bandwagoning theory and Zahavi's original formulation of the handicap principle maintain

that natural selection has promoted alleles that help identify whether their holder has "good genes" by burdening that

holder. Both maintain that this identification of good genes occurs through the satisfaction of a criterion of

differential costliness: The burden is prohibitively costly for individuals without good genes and not as costly (and

possibly not at all costly) for individuals with superior genes. Both maintain that this burden is incurred

conditionally and is, therefore, not necessarily incurred by the parent to the same extent that it is incurred by the

offspring. Specifically, both predict that the extent of the burden incurred varies in this respect: Individuals that are

better at procuring food, territory, mates, and other areas incur a greater burden. However, the two models make

this prediction for different reasons. The handicap principle holds that individuals that are better in these areas are

of higher quality and can, therefore, afford to incur a greater burden. Bandwagoning theory holds that individuals

that are inferior in these areas would, even if all of their abilities were fully known, likely have a lower lineage

fitness outlook, and would, therefore, be rendered by bandwagoning alleles to forfeit resources through resonation

unless the natural enemy threat is severe enough that numerous individuals with a higher lineage fitness outlook

perish, but they would only survive a severe natural enemy threat if they were to not undertake the degree of

reservation undertaken by individuals with a higher lineage fitness outlook.

749   A major difference between the handicap principle and bandwagoning theory is that the handicap principle

holds that an individual incurs the burden in order to demonstrate (i.e., signal) the individual's good genes to others

(particularly mates), while bandwagoning theory holds that the individual incurs the burden in order to identify

whether the individual has good genes for the bandwagoning allele that induces the burden. Consequently,

bandwagoning theory is better at accounting for situations in which it does not appear that an honest signal could be

enforced because of the opportunity for selfish dishonesty. In particular, this includes situations in which the

intended recipient of the signal cannot directly perceive the mechanism that makes the signal costly to individuals

with inferior genes. For example, Folstad & Karter (1992) argued that testosterone 1) reduces immunocompetence

and 2) contributes to the development of ornaments and display behaviors; therefore, individuals can demonstrate

good genes for resisting parasites by increasing testosterone production and, therefore, signaling to potential mates

via their ornaments and display behaviors that they can resist parasites in spite of the associated immunity deficits.

However, since potential mates that view these signals cannot verify, via direct observation, the antibody production

within the body of the individual that exhibits these signals, there may have been mutations over evolutionary time

scales that would have allowed an individual to maintain immunity while also producing attractive ornaments and

displays. If there is evidence of the potential for these mutations, their absence in natural populations is not

explained by the handicap principle, but it is explained by bandwagoning theory. Bandwagoning theory would hold

that the production of these ornaments and display behaviors does, indeed, constitute an honest signal of both the

individual's quality (as indicated by information available to the bandwagoning allele about the abilities that have

been identified) and the immunity burden incurred by the individual, but that this honesty is enforced by a

bandwagoning allele which is practicing resonation by inducing low-quality holders to signal suboptimally and is

practicing reservation by inducing high-quality holders to incur a burden versus parasites. Bandwagoning theory

would, therefore, hold that there was natural selection for a bandwagoning allele (or alleles) responsible for this dual

effect of testosterone. (This concept is visited further in section 5.)

Additionally, there are signals that: 1) are communicated honestly between individuals with conflicting

interests but 2) do not appear to involve any burden associated with communicating a more favorable signal. In

order to account for the honesty with which such signals are communicated, scholars have advanced the index

hypothesis, which states that a signal reporting a characteristic of an individual's condition is reliable if the signal is

tied (i.e., indexed) to the characteristic in such a manner that a dishonest signal cannot be made (Maynard Smith &

Harper, 1995). Biernaskie et al. (2014) argued that the honesty with which index signals are communicated also

owes to the criterion for the preservation of signal honesty on which the handicap principle is based: Dishonest

signals would be too costly to fake. It follows that if there is any evidence that a signal that is communicated

honestly with no cost could have evolved to have been communicated dishonestly with no additional cost, it is not

well explained by the handicap principle, but it is attributable to bandwagoning in the form of resonation: Honest

signals of quality are less favorably transmitted by individuals of lower quality, so that by signaling honestly, lower-

quality individuals wind up foregoing food, mates, and territory compared to higher-quality individuals that signal

honestly.

The handicap principle relies upon (and bandwagoning theory allows) the assumption that it is possible, at

times, for individuals to identify, prior to incurring a burden upon their ability to resist a natural enemy, that their

ability to resist the natural enemy is better or worse than that of another individual. For bandwagoning theory, this

can occur if: 1) one individual or the other has become sick or wounded by the particular natural enemy and 2) this

sickness or wounding is visible or is signaled honestly. (Signaling honesty may, in fact, be imposed by a

bandwagoning allele and this notion is discussed further in section 5.4.) Both the handicap principle and

bandwagoning theory hold that an individual that identifies that its ability to resist a natural enemy is worse than that

of another individual would, ceteris paribus, incur a more modest burden upon its ability to resist that natural enemy

than that other individual. However, bandwagoning theory can better explain situations in which an individual

incurs more of a burden to its ability to resist a natural enemy than another individual even when it appears that it

could not have determined that its ability to resist the natural enemy is better than that of the other individual. For

example, the handicap principle and the model of bandwagoning offered in sections 3.25 and 3.3 both predict that an

individual that compares favorably to others in obtaining food and territory would incur a greater burden upon its

ability to resist parasites. If there is evidence that indicates that individuals that are better at obtaining food and

territory cannot expect to be better at resisting parasites, the greater burdens against parasites that they incur is not

well explained by the handicap principle and is, instead, better explained by bandwagoning theory.

801   A recent result (Schreiber, 2015) that has developed out of bet-hedging literature[4] holds that long-term

natural selection favors a genotype with, ceteris paribus, 1) a higher expected number of offspring produced by

individuals with the genotype across space and time; 2) a lower net variance of offspring produced by individuals

with the genotype across space and time; and 3) a lower correlation of offspring produced between two randomly

selected individuals with the genotype from the same generation. The bandwagoning model presented in sections

3.25 and 3.3 has features that satisfy each of these three properties. By engaging in resonation in order to ensure

that lower-quality holders forfeit resources, a bandwagoning allele yields resources to lineages that have, ceteris

paribus, a higher expected number of offspring (property 1 above), some of which would hold copies of the

bandwagoning allele. A lineage that is not capable of surviving a severe environment has, ceteris paribus, a higher

net variance in offspring produced than a lineage that can survive in a severe environment; therefore, by engaging in

reservation in order to ensure that holders forfeit resources if they do not have a genotype that enables survival in a

severe environment, a bandwagoning allele yields resources to lineages that have, ceteris paribus, a lower net

variance in offspring produced (property 2 above), some of which would hold copies of the bandwagoning allele.

Likewise, by not inducing as much reservation in some holders as in others for the purpose of ensuring that some

holders stay alive if conditions are severe, a bandwagoning allele lowers the correlation in offspring number

between two holders of the allele during the same generation (property 3 above).

---

[4] While Schreiber's (2015) result is developed out of bet-hedging literature, a bandwagoning allele that acts according to the model in sections 3.25 and 3.3 is not necessarily bet-hedging because these acts do not necessarily result in a reduction of arithmetic mean fitness for the holders of the allele.

4) Guidelines for applying bandwagoning theory to empirical phenomena

In theory, it is possible that in populations of some species, there is natural selection for a bandwagoning

allele to induce resonation but not reservation. This is more likely if the bandwagoning allele is selected in a

population that tends not to be subjected to substantial intergenerational variance of the quantity of individuals that

survive to reproductive age as a result of, in particular, predators or parasites. And just as it is theoretically possible

for there to be natural selection for a bandwagoning allele that does not induce reservation at all or induces

reservation from ability against one natural enemy, it is also theoretically possible for there to be natural selection

for a bandwagoning allele that induces reservation against multiple natural enemies in separate ways. However, the

model in sections 3.25 and 3.3 was offered due to its ability to account uniquely for numerous paradoxical empirical

phenomena (examples of which are offered in sections 5.1-5.6). The fundamental elements of this model are a

response by a bandwagoning allele to an indication that its holder has a high lineage fitness outlook with

reservation—a burden imposed to its holder's ability to survive a natural enemy—and a response by a

bandwagoning allele to an indication that its holder has a low lineage fitness outlook with resonation---a cost

imposed against its holder's ability to utilize resources. It was explained in section 3 that as a result of the

possibility that individuals with a high lineage fitness outlook would perish, individuals with a low lineage fitness

outlook may engage in some forfeitures of resources, but are less likely to completely forfeit all of the resources to

which they currently have access and all of the resources to which they will have access in the future. In lieu of an

absolute forfeiture of resources, partial forfeitures can occur numerous times through an individual's life, in

response to indications that the individual's lineage fitness outlook is low (but potentially subject to change).

Additionally, given the present paper's emphasis upon analyses longer than a single generation, it should be noted

that there may be a partial forfeiture not just by an individual but also by that individual's lineage. This would

involve an individual with a low lineage fitness outlook that, for example, produces some non-zero quantity of

offspring but does not produce as many offspring as it could or does not nurture its offspring as much as it could.

As the severity presented by the natural enemy increases, it becomes increasingly likely that an individual

previously judged to have a low lineage fitness outlook no longer has a low lineage fitness outlook because

individuals previously judged to have a higher lineage fitness outlook have perished; therefore, as the severity

presented by the natural enemy increases, the degree to which an individual previously judged to have a low lineage

fitness outlook utilizes its resources can also be expected to increase.

What follows in section 5 is an analysis of six paradoxical phenomena. It is argued in section 5 that these

phenomena represent: 1) an individual's response to indications that its lineage fitness outlook is low by restricting

its ability to utilize resources (which would suggest resonation) and with an increase in ability against a natural

enemy (which would be suggestive of a relaxation of reservation) and 2) an individual's response to indications of a

high lineage fitness outlook with a reversal of a restriction against resource utilization (which would suggest a

relaxation of resonation) or with a decrease in ability against a natural enemy (which would be suggestive of

reservation). Additionally, it is argued that there would have been mutations over evolutionary time scales that

would have allowed an individual to be spared of the costs associated with the restriction and/or the burden that

would not exceed the costs of the restriction and/or burden. Their absence is indicative of natural selection for a

coupling of the restriction in an individual's ability to use resources and that individual's ability against the natural

enemy, such that individuals tentatively evaluated to have a low lineage fitness outlook restrict their ability to use

resources but can resist fully against the natural enemy and individuals tentatively evaluated to have a high lineage

fitness outlook do not restrict their ability to use resources but cannot resist fully against the natural enemy. In this

paper, the term "bandwagoning allele" represents whatever heritable elements are responsible for this pattern and

these elements may involve genes or epigenetic marks that control some genes involved in resource utilization and

natural enemy resistance, but may also involve mutations to some of these genes that are involved in resource

utilization and natural enemy resistance. There is natural selection for the "bandwagoning allele" if there is natural

selection for whatever heritable elements are responsible for this pattern. Table 2 lists, for each of the six

phenomena examined in section 5, the restriction incurred by individuals with a low lineage fitness outlook in

utilizing resources and the burden incurred by individuals with a high lineage fitness outlook in resisting a natural

enemy.

Section	Restriction to resource utilization	Burden against natural enemy
5.1	Costs of depression.	Susceptibility to parasites.
5.2	Photosynthetic cost.	Susceptibility to parasites.
5.3	Diminished sexually-selected characteristics.	Susceptibility to parasites.

5.4	Less advantageous signaling.	Susceptibility to parasites.
5.5	Reproductive inhibition.	Susceptibility to parasites.
5.6	Costs from diminished nurturing.	Susceptibility to parasites.

Table 2.

Additionally, there is a focus in section 5 upon burdens incurred against parasites as examples of

reservation; however, this should not be interpreted to necessarily mean that reservation occurs more commonly

against parasites than predators. Burdens incurred that make survival against predators more difficult, such as

behaviors, ornaments or displays, are highly visible to conspecifics; therefore, these burdens are accounted for by

well-established theory, such as the handicap principle discussed above. They are also accounted for by

bandwagoning theory, but it is typically not necessary to invoke bandwagoning theory in order to explain burdens

incurred against predators. The focus in the section below upon burdens that make survival against parasites more

difficult occurs because these burdens are not accounted for well by the handicap principle, since they involve

burdens to immunity that are not directly observable by conspecifics. Some scholars claim that signals relied upon

by conspecifics are indexed to these burdens, but without an ability to directly observe these burdens, there is also

the potential for the evolution of a dishonest signal. If there is an opportunity for a dishonest signal to evolve that

would allow individuals to signal that they are bearing the burdens without actually incurring them, then the

handicap principle does not account for these burdens that are incurred, but bandwagoning theory might. Therefore,

there is an emphasis in the sections below upon burdens incurred that make survival against parasites more difficult.

5) Empirical applications

5.1) The coupling of robustness against parasites and depressive symptoms

Depression is a response to chronic and uncontrollable stress that is characterized by sadness, self-loathing,

and anhedonia; impaired cognition, concentration, and memory; changes in psychomotor, sleeping, and eating

patterns; and diminished fertility and libido (Anders et al., 2013; Austin et al., 2001; Sun et al., 2013). Among

humans, depression is both prevalent and costly: It increases the mortality rate for all major disease-related causes

of death (Mykletun et al., 2007) and is a significant risk factor for suicide (Hagen, 2003), which claims

approximately one million people worldwide each year (Hawton & van Heeringen, 2009). Individuals in many

species are susceptible to learned helplessness (Seligman, 1976), which is provoked by stress, is characterized by

many of the symptoms associated with depression (e.g., anhedonia, psychomotor retardation, reduced self-care, and

diminished appetite and libido), and has been induced, via application of uncontrollable stress, in many species,

including insects, cats, dogs, rats, birds, fish, and primates (Franklin et al., 2012; Sapolsky, 1998; Weiss, J. et al.,

1982). Though there are obstacles in determining the intent, some models of learned helplessness have included

suicide and self-mutilation (Crawley et al., 1985; Preti, 2011; Sapolsky, 1998; Seligman, 1976). These depressive

symptoms are considered maladaptive and are reversed in animals by the same therapies that are used to treat

depression in humans (Russo et al., 2012; Weiss, J. et al., 1982).

Some scholars have argued that there was natural selection for depression (e.g., Andrews & Thomson,

2009; Raison & Miller, 2013; Watson & Andrews, 2002), citing the commonness of depressive responses among

humans of all ages—including ages at which natural selection is strong (Medawar, 1952)—and the prevalence of

depression risk alleles (Anders et al., 2013; Raison & Miller, 2013). Many evolutionary explanations for depression

have been offered, yet none have been embraced by a scholarly consensus (Hagen, 2003; Kinney & Tanaka, 2009;

Nettle, 2004; Raison & Miller, 2013; Varga, 2012). Of particular relevance to genetic bandwagoning theory, a

recent set of hypotheses have been advanced, holding that depression was selected in conjunction with the

prevention of infection by parasites and pathogens (Anders et al., 2013; Kinney & Tanaka, 2009; Raison & Miller,

2013). These scholars have pointed to evidence for a mutually causal relationship between immune activation and

depression, noting that immune activation leads to depression, whereas depressive symptoms are associated with

greater resistance against infections. According to these hypotheses, the depressive mood is adaptive for fighting

infection since it encourages energy preservation and it also leads to solitude, which protects the afflicted individual

from further exposure to infection and also protects kin from exposure to the afflicted individual. Under these

views, the immunity activation and depression that occur in response to a wide range of stressors (including

psychosocial stressors) are adaptive, since over evolutionary time scales, most stressors related to hunting, escaping

predators, or fighting conspecifics in dominance hierarchies, and infection risk was increased by wounds associated

with these activities. These hypotheses account well for the immune activation among depressives, but do not

account as well for the self-loathing and its associated costs. Depression is a response to a wide variety of stress

sources and while some causes of depression might have suggested an increased infection risk, other common ones

(e.g., the death of kin or the loss of a mate without combat) may not have imposed enough of an infection risk to

make depressive mood beneficial to the individual or to kin, especially since kin can be adversely affected by an

individual's depression (Ramchandani et al., 2005; Shen et al., 2016). Indeed, depression is caused by numerous

non-infectious diseases, even if the individual is unaware that he or she is diseased (Dowlati et al., 2010; Irving &

Lloyd-Williams, 2010; Parker & Brotchie, 2017; Pryce & Fontana, 2017; Yirmiya et al., 1999). Additionally,

depression is a costly way of exerting a preference for solitude and it would seem that mutations could have

occurred over evolutionary time scales that would have instilled in a sick individual a preference for solitude without

requiring that individual to incur the costs of self-loathing, given that some well-adjusted individuals exhibit a

preference for solitude (Berger, 1995).

The self-loathing and immunity activation features of depression are well-explained by genetic

bandwagoning theory. Since chronic stress is an indication of lower quality and a lower lineage fitness outlook[5], the

reproductive costs incurred by the chronically stressed (e.g., diminished fertility and libido; suicide) are re-

interpreted as resonation by individuals with a lower lineage fitness outlook, which allows their resources to be

utilized by nearby individuals. The augmented defenses versus parasites that have been associated with depression

are also re-interpreted: Individuals of higher quality that do not undertake these defenses are engaging in reservation

against these natural enemies.

Genetic bandwagoning theory also accounts better for the finding that depression tends to result not from a

single incidence of stress but from a prolonged (i.e., chronic) bout with stress and that the likelihood of developing

depression is higher if an individual has a history of it (Brilman & Ormel, 2001; Ghaziuddin et al., 1990; Kendler et

al., 2000; Solomon et al., 2000). Evolutionary explanations that maintain that depression is a response to infection

risk fare poorly in explaining the increasing intensification of depressive symptoms, since a history of chronic stress

that has not yet resulted in an infection would be suggestive of a reduced marginal risk of infection from further

incidents of stress. However, in the context of genetic bandwagoning theory, a history of chronic stress is more

suggestive than a single stress bout that the stress-afflicted individual's lower quality is irreparable; therefore, a

---

[5] Some stressors, such as stressors relating to physical exercise, are not indicative of lower quality or a lower lineage fitness outlook, but these stressors also do not typically contribute to the likelihood of depression (Schuch et al., 2016).

bandwagoning allele is more likely to induce its holder to completely forfeit its resources if the holder has been

subjected to a history of chronic stress.

5.2) The coupling of robustness against parasites and photosynthetic costs

Plants of many species respond to a wide variety of stressors by increasing anthocyanin production, which

makes their leaves redder. Examples of sources of stress that lead to an increase in anthocyanin production include

strong light, UV-B radiation, temperature extremes, drought, nutrient deficiencies, bacterial and fungal infections,

wounding, herbivory, herbicides, and various pollutants (Gould et al., 2009). This production of anthocyanins

imposes a photosynthetic cost, as anthocyanins absorb visible light without being photosynthetic (Gould et al., 2009;

Karageorgou et al., 2008; Manetas, 2006).

The reddening of plant leaves[6] is biochemically well-explained but is an evolutionary paradox that has,

over the past century, invited many hypotheses, none of which are thought to adequately explain the variation in

spatial and temporal patterns of anthocyanin production (Cooney et al., 2012; Gould et al., 2009; Kovinich et al.,

2014; Manetas, 2006; Menzies et al., 2016). Of interest to genetic bandwagoning theory, anthocyanins are produced

in a common biosynthetic pathway with defensive compounds against parasites (e.g., phenols, tannins: Schaefer &

Rolshausen, 2006), so that when plants produce anthocyanins, they also augment their defenses versus parasites.

Therefore, when plants that are subjected to stress produce anthocyanins, they incur a photosynthetic cost, but also

enhance their defenses against parasites. One reason for the paradox is that the tradeoff between photosynthesis and

defense against parasites does not appear to have been imposed by physical constraints. These plants evidently

could have evolved to produce anthocyanins and defense compounds against parasites in separate pathways without

making additional energy expenditures or incurring other costs. This would have allowed a plant to experience the

parasite defense without the photosynthetic cost from the anthocyanins. The absence of plants with this ability (or

the pervasiveness of plants without it) is well-explained by genetic bandwagoning theory: Plants that are afflicted

by a stress source, ceteris paribus, have a lineage fitness outlook that is lower than plants that are able to resist the

same source without becoming stressed. By producing anthocyanins, plants with a lower lineage fitness outlook

incur a photosynthetic cost, which is consistent with resonation, but they also enhance their defenses against

parasites. The individuals that do not produce additional anthocyanins (that is, individuals with a higher lineage

____

[6] Plants produce anthocyanins in other contexts besides stress response that likely are unrelated to genetic
bandwagoning theory.

fitness outlook) do not enhance their defenses against parasites and are, in fact, engaging in reservation by

underperforming their maximum resistance against parasites. The interpretation in terms of genetic bandwagoning

theory is that, over evolutionary time scales, many plants could defend against parasites without incurring a

photosynthetic cost, which allowed individuals with a high lineage fitness outlook to defend to greater capacity

against parasites and which allowed individuals with a low lineage fitness outlook to avoid a photosynthetic cost.

However, their absence today, and the common coupling of anthocyanin production and production of defense

compounds, is evidence of natural selection for genetic bandwagoning alleles that have imposed this tradeoff

between photosynthesis and defense against parasites.

5.3) The coupling of susceptibility against parasites and androgenic benefits

Individuals of lower quality tend to utilize androgens (e.g., testosterone) less than individuals of higher

quality (Folstad & Karter, 1992; Johnstone, 1995; Muehlenbein & Bribiescas, 2005). Since androgens are utilized

in strength, aggression, and mate-attracting ornaments, the lesser utilization of androgens by lower-quality

individuals leads them to be disadvantaged in competition for food, territory, and mates (Folstad & Karter, 1992;

Gil, 2008; Groothuis et al., 2005). An explanation commonly offered for the lesser utilization of androgens by

lower-quality individuals is that the utilization of androgens steers energetic resources into reproduction and away

from immunity, which those with weaker immunity cannot afford (Folstad & Karter, 1992; Groothuis et al., 2005;

Habig & Archie, 2015; Muehlenbein & Bribiescas, 2005). This explanation typically invokes the handicap

principle. The point raised in section 3.4 is pertinent: The handicap principle relies upon an assumption that an

individual can determine prior to incurring a burden (i.e., a handicap) that the individual has superior capability for

overcoming the burden in comparison to other individuals. Individuals that perform better in competitive bouts or at

attracting mates raise androgen levels (Archer, 2006; Oliveira, 2004), even though it is questionable that individuals

can know they are more capable of resisting parasites than others on the basis of their ability to fight conspecifics or

attract mates.

Moreover, there is evidence that a tradeoff between androgens and immunity is not imposed by energy

constraints. In a meta-analysis, Roberts et al. (2004) found no effect of androgens upon direct measures of

immunity, but they did find androgens to increase ectoparasite abundance in several studies. In a larger, more recent

meta-analysis, Habig & Archie (2015) replicated the finding that dominant males are consistently more susceptible

to a breadth of parasites, but found that result "puzzling," since they could find no significant differences between

the immune responses of dominant and subordinate males. They suggested that dominant males may suppress

antibody production in response to antigens, which is an aspect of the less energetically costly Th-2 mediated

immunity and which is contrary to how individuals reducing immunity to save energy would be predicted to operate

(Lee, 2006). Consequently, Habig & Archie (2015) concluded that their results did not support the hypothesis that

dominant individuals sacrifice immune function to engage in greater reproductive effort. Relatedly, Kankova et al.

(2014) found that when Japanese quail offspring were subjected to moderate caloric restriction during development,

the offspring with high egg testosterone content retained their growth advantage and did not suffer an immunity

disadvantage, which, they concluded, contradicted the hypothesized energetic tradeoff between androgen-mediated

growth and immunity.

Bandwagoning theory offers the explanation that by utilizing androgens in lesser quantity, lower-quality

individuals that have a lower lineage fitness outlook are engaging in resonation via the forfeiture of resources that

occurs by possessing lower strength, aggression, and mate-attracting ornaments, and that by utilizing androgens in

greater quantity, higher-quality individuals that have a better lineage fitness outlook are engaging in reservation via

the lesser ability to resist parasites. The evidence that there could have been mutations over evolutionary time scales

that would have allowed an individual the strength, aggression, and ornaments associated with androgens without

incurring a cost against parasites and the absence of these mutations, or the prevalence of androgens exerting this

dual effect, is evidence of natural selection for bandwagoning alleles that mediate this dual effect.

5.4) The maintenance of honest signaling when a selfishly dishonest signal would be uncostly

It was argued in section 3.4 that the handicap principle would not account well for evidence that an

uncostly signal is communicated honestly despite opportunities to communicate it dishonestly. Additionally, it was

argued in section 3.1.2 that honest signaling is well explained by bandwagoning in the form of resonation: If

individuals signal honesty, irrespective of quality, lower-quality individuals are disadvantaged because they signal

less favorably; consequently, mating opportunities and other resources that are determined by signaling are more

likely to be obtained by higher-quality individuals than if honest signaling were not to occur. Honest signaling is,

therefore, a means by which a lower-quality individual relinquishes resources to higher-quality individuals.

The index hypothesis has been advanced to account for the honesty of some uncostly signals (Husak &

Swallow, 2011). However, there is evidence that signals that were thought to be indices are not true indices because

they are subject to selection (Emlen et al., 2012; Simons et al., 2015). Thus, there is the paradoxical occurrence of

honest but apparently uncostly signals.

For example, carotenoid ornaments reliably reflect quality, even though carotenoids are evidently not

limiting (Hill, 2006, 2014; McGraw & Blount, 2009; Simons et al., 2014) and there are mutations that would allow

an individual to cheat with a dishonest signal that would cost the cheater less than the cost imposed by honesty

(Simons et al., 2015). Additionally, zebra finches use song as an honest signal of quality, even though it is

metabolically cheap (Ritschard et al., 2010; Zollinger et al., 2011); stressed birds incur song deficits even after a

song allotment that is adequate for exact copying in other birds (Brumm et al., 2009); and some nutritionally

stressed populations of wild zebra finches have more complex songs than populations with greater food availability

(Zann & Cash, 2008), which indicates that zebra finches could have evolved the ability to learn complex song

despite undergoing nutritional stress.

There is a role for testosterone in mediating the honesty of these signals: In many species, an individual's

testosterone levels are reduced by stressors that are suggestive of that individual's lower quality and lower lineage

fitness outlook, such as nutritional, developmental, and psychological stressors (Dong et al., 2004; Lynn et al., 2015;

Nelson et al., 1989). Chronic stress is more likely than acute stress to lead to a decrease in testosterone production

(Chichinadze & Chichinadze, 2008), which is consistent with the notion that an individual's testosterone levels

correlate with its lineage fitness outlook, since a chronic stressor is a more reliable indicator than an acute stressor

that the stressed individual's lineage fitness outlook is low. Additionally, there is evidence that the expression of

carotenoid ornaments is dependent upon testosterone and that testosterone mediates the relationship between

individual quality and carotenoid ornament color by being produced more in higher-quality individuals and

engendering more prominent carotenoid ornament production (McGraw & Blount, 2009; Weaver et al., 2017).

There is also evidence that the structure of zebra finch song, which is important in attracting mates

(Woodgate et al., 2012), is an indicator of the testosterone levels that zebra finch has experienced in its history

(Fedurek et al., 2016; Ritschard et al., 2011) and, therefore, is an indicator of the degree to which that zebra finch's

lineage fitness outlook. In addition to augmenting signals, testosterone leads to greater susceptibility to parasites,

which was discussed in section 5.3, and there is evidence that this also occurs in songbirds, including zebra finches

(Alonso-Alvarez et al., 2007). In sum, testosterone is ramped up within individuals with a higher lineage fitness

outlook, which leads these individuals to signal favorably to mates and conspecifics but to be more susceptible to

parasites, while individuals with a lower lineage fitness outlook signal unfavorably but are less susceptible to

parasites. In accordance with the above discussion, there are likely mutations that would have occurred over

evolutionary time scales that would have allowed an individual to signal favorably and to resist parasites without

additional susceptibility. This could have involved, for example, a way to signal favorably, irrespective of

testosterone levels, or it could have involved a way of increasing testosterone without incurring the susceptibility to

parasites. That these mutations are not prevalent is an indication of natural selection for genetic bandwagoning

alleles that lead individuals with a low lineage fitness outlook to engage in resonation by signaling unfavorably and

that lead individuals with a high lineage fitness outlook to engage in reservation by becoming more susceptible to

parasites.

5.5) The forfeiture of reproduction by lower-quality individuals

In many species, subordinate individuals cooperatively care for the offspring of dominant individuals

without having their own offspring, which is considered a paradox (Clutton-Brock et al., 2010; Johnstone, 2000;

Saltzman et al., 2009). In some species, a group's subordinates are completely bereft of reproductive ability;

however, their fertility and libido quickly return if the dominant group members depart the group (Sapolsky, 2001).

The states of dominance and subordinance are commonly attributed to differential quality: Dominance consistently

reflects certain aspects of quality, especially immunocompetence (Cavigelli & Chaudhry, 2012; Georgiev et al.,

2015). Subordinates' reproductive inhibition (which has been called "socially induced suppression," "physiological

suppression," and "social contraception") has been documented in mammals, birds, fish, and invertebrates (Saltzman

et al., 2009) and sometimes occurs even though subordinates show comparable body condition to dominants and

overlap dominants substantially in age (Harrison et al., 2013; Saltzman et al., 2009).

Despite well-established theory regarding how individuals in a group should be expected to divide

reproduction (Vehrencamp, 1983), multiple large reviews have determined that theory is not matched well by

experimental and observation results (Field, J. et al., 1998; Nonacs & Hager, 2011). Recently, a comprehensive

review by Nonacs & Hager (2011) found that there is little evidence that the behavior of members in cooperatively

breeding groups is governed by an evaluation of patterns of relatedness or competitive ability in order to set

reproductive skew. The review's authors wrote (pg. 295) that "some individuals appear to behave suboptimally and

lose fitness by either joining disadvantageous groups or not leaving them."

Genetic bandwagoning theory offers the explanation that the forfeiture of reproduction by lower-quality

individuals looks maladaptive when analyzed after one generation, but that an analysis over a longer time period

would demonstrate its adaptiveness. Indeed, Nonacs & Hager (2011) noted that high heritability is suggested in

species in which reproductive skew is unresponsive to internal group characteristics such as relatedness, which

indicates an emphasis upon a longer-term analysis.

The reproductive inhibition of subordinates coincides with dominants' greater susceptibility to parasites,

which was discussed in section 5.3. The reproductive inhibition by subordinates is an example of a forfeiture of

resources by individuals with a lower lineage fitness outlook and the greater susceptibility to parasites by dominants

is an example of a reservation of ability against a natural enemy by individuals with a higher lineage fitness outlook.

A recurring point of emphasis in section 5 is that both phenomena are considered paradoxes, but are well-explained

by bandwagoning theory: The reproductive inhibition of subordinates is an example of resonation and the greater

susceptibility to parasites by dominants is an example of reservation.

5.6) Nurturing deficits toward lower-quality offspring

In numerous rodents and primates, higher levels of maternal nurturing (e.g., licking and grooming) lead to

more growth (Denenberg & Karas, 1959; Field, T., 1998; Field, T. et al., 1986; Schanberg, 1995), greater cognitive

function and memory (Field, T., 1998; Korosi & Baram, 2009; Zhang & Meaney, 2010), higher alertness (Field, T.,

1998; Field, T. et al., 1986), greater behavioral maturity (Field, T. et al., 1986), and lower mortality risk (Denenberg

& Karas, 1959; Field, T. et al., 1986), while lower levels of maternal nurturing lead to depression (Franklin et al.,

2010; Franklin et al., 2011; Weiss, I. et al., 2011), anxiety (reviewed in Korosi & Baram, 2009; Weiss, I. et al.,

2011), and greater tendencies to resign to stressful sources (Nestler, 2012). Despite the costs incurred by offspring

that are nurtured less than others, mothers that are chronically stressed exhibit diminished nurturing across the

aggregate of their offspring in comparison to mothers that are not chronically stressed (Bosch et al., 2007; Heiming

et al., 2011; Mansuy et al., 2013; Zhang & Meaney, 2010). A mother's nurturing practices tend to remain consistent

through her lifetime and her offspring are likely to impart the same levels of nurturing to their offspring that they

received from their mother—within-lineage consistency of maternal nurturing which is mediated epigenetically

(Drury et al., 2016; Mansuy et al., 2013; Meaney, 2001; Zhang & Meaney, 2010).

1104   The prevalent evolutionary explanation for diminished nurturing by stressed mothers is that it acts as a

signal to offspring regarding the level of vigilance required in the environment, particularly versus predators

(Champagne et al., 2003; Drury et al., 2016; Pittet et al., 2012). According to this view, the additional anxiety and

increased stress response associated with diminished maternal nurturing better enables the offspring to avoid danger,

especially predators. However, evidence that has accumulated against this explanation includes: 1) Offspring from

the same mother (and evidently the same formative environment) are induced via maternal effects to incur costs that

are determined by their father's quality, even in the absence of their father's postpartum presence (Dietz et al., 2011;

Dietz & Nestler, 2012; Mashoodh et al., 2012); 2) Exposure to predator odor during gestation has been found to

stimulate postpartum maternal affection (Mashoodh et al., 2009; McLeod et al., 2007)—the opposite of what would

be expected if reduced nurturing were intended to increase vigilance toward predators; and 3) Offspring of preferred

males, which tend to receive more nurturing, exhibit more risk-averse behavior in a predator-avoidance test (Curley

et al., 2011).

1116   Work by Sapolsky (1998) and Suomi (1997) also suggests that reduced maternal nurturing exerts a negative

influence upon offspring. They independently documented what they considered inefficient stress responses in,

respectively, baboons and rhesus monkeys. These animals apparently do not distinguish in their reactions to

threatening and innocuous situations: Their resting glucocorticoid levels are pathologically higher than the average

but their glucocorticoid response to a real stressor is smaller and slower (Sapolsky, 1998). This condition, which

Sapolsky (1998) and Suomi (1997) each considered maladaptive, is both a cause and an effect of reduced maternal

nurturing.

1123   Genetic bandwagoning theory offers another evolutionary explanation for differential maternal nurturing.

A chronically stressed individual has, ceteris paribus, a lower lineage fitness outlook. Therefore, a bandwagoning

allele would respond to chronic stress experienced by its holder by inducing its holder to nurture its offspring less

than other individuals nurture their offspring, which disadvantages its offspring and leads these offspring to use less

resources for their own reproduction in comparison to the offspring of other individuals. This is a way of inducing

lineages with a lower lineage fitness outlook to use less resources while not eliminating the lineages, in case of a

change to their lineage outlooks.

Moreover, there is evidence that in addition to the resonation occurring from differential nurturing, there is

also reservation by nurtured individuals. In numerous species, diminished maternal nurturing is associated with

increases in inflammation (Chen et al., 2011; Figueiredo et al., 2016; Gracia-Rubio et al., 2016; Miller et al., 2011),

which are associated with resistance against parasites (Anders et al., 2013; Raison & Miller, 2013). Since it has not

been argued that individuals incur costs from diminished nurturing in order to save energy directed into resistance

against parasites, it appears that, over evolutionary time scales, the resistance against parasites could have occurred

without incurring the costs from diminished nurturing. The scarcity in natural populations of this coupling in the

same individual is a paradox that is explained by bandwagoning theory: Resonation is exemplified by the costs

from diminished nurturing and reservation is exemplified by the greater susceptibility to parasites by nurtured

individuals.

6) Conclusions

In this paper, it is argued that there can be natural selection for an allele that induces its holder to forfeit

resources if that holder's outlook for lineage fitness is sufficiently low in comparison to the individuals nearby

and/or if other parameters (e.g., values for heritability and relatedness) are satisfactorily high. When this occurs

sequentially—that is, when an allele responds to an indication that its holder has a low lineage fitness outlook by

inducing that holder to forfeit resources—it is called resonation. However, if a holder's lineage fitness outlook

depends upon heritable dimensions that are difficult to evaluate under non-severe conditions (e.g., abilities against

parasites or predators), an allele induces that holder to reserve from (i.e., withhold) its ability to survive one or more

natural enemies, which leads that holder to become sick or perish if its ability to survive the natural enemy against

which it reserves is sufficiently low that its lineage fitness outlook is low, and this sickness or perishing by a holder

with a low lineage fitness outlook leads that holder to be less capable of procuring, attracting, or utilizing resources,

which constitutes a forfeiture of resources to nearby individuals. This is called reservation.

The focus in this paper is upon a model in which an allele uses a particular combination of resonation and

reservation: The allele evaluates its holder's lineage fitness outlook on the basis of information it has available, such

as information about the holder's ability to obtain food, territory, and mates, and whether the holder has been

sickened by parasites or wounded by predators (however, in accordance with the discussion in this paper, it is less

likely that the allele has information about the holder's ability to survive against parasites or predators). The allele

then induces its holder to a degree of reservation against a natural enemy that is positively correlated to that

evaluation of its holder's lineage fitness outlook and a degree of resource forfeiture that is negatively correlated to

that evaluation of its holder's lineage fitness outlook: The higher that holder's lineage fitness outlook is evaluated,

the less the allele induces it to forfeit resources and the more the allele induces it to reserve against a natural enemy

and the lower that holder's lineage fitness outlook is evaluated, the more the allele induces it to forfeit resources and

the less the allele induces it to reserve against a natural enemy. Holders that are tentatively evaluated to have a high

lineage fitness outlook are induced to reserve from their ability against the natural enemy in order to determine

which of these holders have "good genes" for surviving the natural enemy and to induce forfeitures by individuals

that do not have "good genes" for surviving the natural enemy. Holders that are tentatively evaluated to have a low

lineage fitness outlook are also induced to forfeit resources, but are induced to reserve less against the natural enemy

for the following reason: If the natural enemy threat is of typical severity, Hamilton's rule is likely satisfied by the

forfeiture of resources by individuals that are tentatively evaluated to have a low lineage fitness outlook, even if

these individuals are good at surviving natural enemies. The best chance that Hamilton's rule would not be satisfied

by the forfeiture of individuals that are tentatively evaluated to have a low lineage fitness outlook is if the natural

enemy threat becomes unexpectedly severe, in which case numerous individuals that had a high lineage fitness

outlook would perish due to their reservation, but if individuals with a low lineage fitness outlook were to reserve as

much, they, too, would perish.

The six empirical phenomena in section 5 are presented as examples of well-documented but heretofore

paradoxical phenomena, which are uniquely accounted for by this model of bandwagoning. Each case involves: 1) a

response to an indication that an individual's lineage fitness outlook is low with a restriction of that individual's

ability to utilize resources (which indicates resonation) and an increase in the robustness of that individual's

defenses against natural enemies (which indicates a relaxation of reservation) and 2) a response to an indication that

an individual's lineage fitness outlook is high with a relaxation of that restriction against resource utilization (which

indicates a relaxation of resonation) and a decrease in the robustness of that individual's defenses against natural

enemies (which indicates reservation). In each case, it appears that mutations could have occurred over evolutionary

time scales that would have allowed the individual to avoid the restriction and/or the burden against natural enemies

at a cost lower than the cost associated with the restriction and/or the burden. The paradoxical absence of such

mutations is indicative of natural selection for a bandwagoning allele that would induce these responses. Each of

these six cases lends: a) support for the notion that the parameters (e.g., relatedness, heritability of fitness and

condition) required for the natural selection of a bandwagoning allele do tend to occur in natural populations and b)

relevance to genetic bandwagoning as an evolutionary theory that can account for empirical phenomena.

Sections 5.1, 5.2 and 5.3 examine phenomena for which the restriction and the burden are reported as

aspects of the same phenomena and individuals are assumed (by, at least, some scholars) to be constrained to choose

one or the other. That is, individuals are assumed to have to choose, respectively, the costs of depression or

susceptibility to parasites; the costs of anthocyanin production or susceptibility to parasites; and the absence of

sexually-selected characteristics or susceptibility to parasites. In each case, they are assumed to be unable to have

both. Sections 5.4, 5.5 and 5.6 examine phenomena for which the restriction and the burden are not reported

together. That is, robustness (or the absence of susceptibility) against parasites is assumed to merely coincide with,

respectively, disadvantageously honest signaling, reproductive inhibition, and diminished nurturing. These cases for

which tradeoffs are not even suspected lend credibility to the argument that the phenomena that fit this pattern that is

consistent with the model of genetic bandwagoning introduced in section 3 (3.25 and 3.3) are not attributable to

tradeoffs imposed by physical constraints, but are rather attributable to bandwagoning alleles.

Thus, there are the following major theories presented in this paper:

1) There can be natural selection for genetic bandwagoning in the form of resonation, in which an allele evaluates its

holder's lineage fitness outlook and acts, in any of a large number of ways, to induce that holder to forfeit resources

if that outlook is low.

2) There can be natural selection for genetic bandwagoning in the form of reservation, in which an allele cannot

normally evaluate some facets of its holder's lineage fitness outlook and, therefore, induces its holder to reserve

from its ability to evade predators, resist parasites, capture prey, or otherwise survive, which allows the allele to

identify its holder's lineage fitness outlook on the basis of the degree to which it can overcome this reservation and

which leads a holder with an unfavorable lineage fitness outlook to forfeit resources due to death, injury, or sickness

from the reservation.

3) There can be natural selection for genetic bandwagoning in the form of a combination of resonation and

reservation, which is verbally outlined in section 3.2.5 and algebraically demonstrated in section 3, in which

individuals that receive information that suggests they have a high lineage fitness outlook reserve from their ability

to compete or survive against one or more natural enemies, while individuals that receive information that suggests

they have a low lineage fitness outlook forfeit resources to nearby individuals but reserve less (if at all) from their

ability against natural enemies and remain alive insofar as it is likely that their lineage fitness outlook can change.

4-9) The six empirical phenomena analyzed in sections 5.1 to 5.6 are not well-explained by contemporary

evolutionary theory and are best explained as products of natural selection for genetic bandwagoning that uses the

combination of resonation and reservation that is referenced in theory 3 above in this summary.

6.1) Challenging the theories presented in this paper

Successful challenges to the theories presented might involve a demonstration that, for example: a) the

parameters required for the natural selection of bandwagoning do not usually occur in nature and/or b) it is not

necessary for an allele within an individual to induce reservation against a natural enemy in order to evaluate that

individual's ability against that natural enemy. Any argument that demonstrates that any of the six phenomena

considered in section 5 does not require an explanation by bandwagoning theory would subtract some relevance

from the theory and reduce support for the notion that the parameters required for the natural selection of

bandwagoning tend to occur in nature.

6.2) Potential for future research

The foregoing examples do not constitute an exhaustive set of phenomena that can be accounted for by

resonation and/or reservation. Productive avenues for further utilization of the concepts of resonation and

reservation in evolutionary explanations may include, for example, the phenomena by which 1) stressed individuals

tend to produce male offspring (Cameron, 2004), even though theory holds that less competitive individuals should

generally prefer to produce female offspring (Trivers & Willard, 1973) and 2) individuals that are defeated in

competitive bouts tend, ceteris paribus, to perform more poorly in subsequent bouts (Dugatkin, 1997; Hsu et al.,

2006; Rutte et al., 2006).

6.2) A bandwagoning allele does not necessarily act against the interests of alleles at other loci

By inducing resonation and/or reservation, a bandwagoning allele should not necessarily be interpreted to

be acting against the interests of alleles at other loci, even if it acts to induce its holder to forfeit resources. Insofar

as a bandwagoning allele's frequency can be expected to increase over ensuing generations if the individual in

which it is located forfeits resources to individuals with a more favorable lineage fitness outlook, the frequencies of

alleles at other loci within the same individual can also be expected to increase over the ensuing generations as a

result of the same forfeiture of resources—via shifting the resources to fitter lineages, some of which would hold a

copy of the same alleles. That is, an effect of a bandwagoning allele upon alleles at other loci would be to induce

the forfeiture resources from the holders of those alleles that have the lowest lineage fitness outlooks to the holders

of those alleles that have the highest lineage fitness outlooks, allowing those alleles to, ceteris paribus, benefit from

having the resources utilized by holders with the highest lineage fitness outlooks. The term "ceteris paribus" is used

above because if an allele at another loci has a negative influence upon fitness, it would likely be disproportionately

represented within the genotypes of the forfeiting individuals, thereby being disadvantaged by the bandwagoning

allele, and if an allele at another loci has a positive influence upon fitness, it would likely be disproportionately

represented within the genotypes of the utilizing individuals, thereby being benefited by the bandwagoning allele.

6.3) Genetic bandwagoning as a mechanism for the evolution of cooperation

The evolution of cooperation is a perennial question (Hamilton, 1964; Trivers, 1971; Nowak, 2006, 2012;

West et al., 2007; West et al., 2011). Cooperation is defined (West et al., 2007) as "a behaviour which provides a

benefit to another individual (recipient), and which is selected for because of its beneficial effect on the recipient."

The resource forfeitures by low-quality individuals that result from genetic bandwagoning constitute cooperation

that affords indirect genetic benefits (West et al., 2007; West et al., 2011) to the forfeiting individual. In

comprehensive reviews of the evolution of cooperation, kin discrimination, greenbeard discrimination, group

selection, limited dispersal, and spatial selection have been offered as ways in which there can be evolution of

cooperation for indirect benefits (Nowak, 2006, 2012; West et al., 2007; West et al., 2011). These are all ways of

referring to a state in which cooperation evolves primarily because of a high degree of positive assortment in the

population. In other words, cooperation that owes to these explanations satisfies Hamilton's rule primarily via high

relatedness between the cooperating and benefiting individuals rather than a high benefit-to-cost ratio associated

with the cooperation. Genetic bandwagoning is distinct from these explanations because cooperation that owes to

genetic bandwagoning can satisfy Hamilton's rule even if relatedness is low, since it satisfies Hamilton's rule

(which is C < rB) primarily via a high benefit-to-cost ratio (high B/C), which occurs due to the better abilities of

"good genes" individuals (and ultimately, their lineages) to turn the resources into fitness over ensuing generations.

Figure 2 depicts various explanations for cooperation on a Cartesian curve whose coordinates are the combinations

of the minimum values of relatedness and benefit-to-cost ratio that allow Hamilton's rule to be satisfied by

cooperation.

1268

1269

1270   *Figure 2. The minimum values for B/C for a given r value that are required for the satisfaction of Hamilton's rule. The curve*

1271   *represented is B/C=1/r. (Curve is not drawn to scale.)*

1272

At one end of the Cartesian curve in Figure 2 are kin discrimination, greenbeard discrimination, group

selection, limited dispersal, and spatial selection, which are explanations for the evolution of cooperation due to

indirect genetic benefits that are typically reliant upon high relatedness ($r \gg 0$) and that can accommodate a low

benefit-to-cost ratio ($B/C \approx 1$). At the other end is genetic bandwagoning, which relies upon a high benefit-to-cost

ratio ($B/C \gg 1$) and can accommodate relatedness not much higher than 0 ($r \approx 0$).

6.4) The evolutionary influences exerted by a bandwagoning allele

1279   By inducing resonation and/or reservation, a bandwagoning allele exerts an evolutionary influence upon

1280 other genes in the population. Through both resonation and reservation, the bandwagoning allele acts to direct

1281 resources from genotypes with a long-term fitness disadvantage to genotypes with a long-term fitness advantage and

1282 the bandwagoning allele becomes selected as its copies ride on the bandwagon of the genotypes that offer this long-

1283 term fitness advantage. In directing resources to genotypes with a long-term fitness advantage, the bandwagoning

1284 allele accelerates the fixation of the genes at other loci on these genotypes that are responsible for this long-term

1285 fitness advantage.

1286   By inducing reservation against natural enemies, a bandwagoning allele exerts a reverse effect upon these

1287 natural enemy populations. Just as reservation of ability against predators or parasites makes it more difficult for

1288 reserving individuals to survive against these natural enemies, it also makes the challenge of predating or

1289 parasitizing easier for, respectively, the predators or parasites against which the reservation occurs. Just as

1290 reservation is differentially costly—that is, costlier to individuals that do not have high ability against these natural

1291 enemies—to reserving individuals, reservation is also differentially beneficial to individuals in these natural enemy

1292 populations—that is, it is more beneficial to low-ability predators or parasites. Predator individuals or parasite

1293 individuals that might have no success at predating or parasitizing an individual that does not reserve might be able

1294 to successfully predate or parasitize a reserving individual. Thus, a bandwagoning allele that induces reservation

1295 from ability that is used against a natural enemy population can act to attenuate the intensity of natural selection

1296 upon that natural enemy population.

1297   In sum, a bandwagoning allele acts to increase the intensity of natural selection upon the population in

1298 which the bandwagoning allele is located. A bandwagoning allele that induces reservation from ability used against

1299 a natural enemy population can also act to attenuate the intensity of natural selection upon that natural enemy

1300 population.

1301 **Acknowledgments**

1302 This research did not receive any specific grant from funding agencies in the public, commercial, or non-for-profit

1303 sectors.

1304 **Declarations of Interest**

No competing interests.

References

Allen, B., Nowak, M., 2015. Games among relatives revisited. *J. Theor. Biol.* (378), 103-116.

Alonso-Alvarez, C., Bertrand, S., Faivre, B., Chastel, O., Sorci, G., 2007. Testosterone and oxidative stress: the
oxidation handicap hypothesis. *Proc. R. Soc. B: Biol. Sci.* 274(1611), 819-825.

Amundsen, T., Forsgren, E., 2001. Male mate choice selects for female coloration in a fish. *Proc. Natl. Acad. Sci.*
U.S.A. 98(23), 13155-13160.

Anders, S., Tanaka, M., Kinney, D., 2013. Depression as an evolutionary strategy for defense against infection.
Brain Behav. Immun. (31), 9-22.

Andersson, M., 1982. Sexual selection, natural selection and quality advertisement. *Biol. J. Linn. Soc.* 17(4), 375-
393.

Andersson, M., 1994. *Sexual Selection.* Princeton: Princeton University Press.

Andrews, P., Thomson, J., Jr., 2009. The bright side of being blue: depression as an adaptation for analyzing
complex problems. *Psychol. Rev.* 116(3), 620-654.

Archer, J., 2006. Testosterone and human aggression: an evaluation of the challenge hypothesis. *Neurosci.*
Biobehav. Rev. 30(3), 319-345.

Austin, M., Mitchell, P., Goodwin, G., 2001. Cognitive deficits in depression: possible implications for functional
neuropathology. *Br. J. Psychiatry* 178(3), 200-206.

Berger, J., 1995. Individual differences in preference for solitude. *J. Res. Pers.* 29(1), 85-108.

Bergeron, P., Baeta, R., Pelletier, F., Reale, D., Garant, P., 2011. Individual quality: tautology or biological reality?
J. Anim. Ecol. 80(2), 361-364.

Biernaskie, J., Grafen, A., Perry, J., 2014. The evolution of index signals to avoid the cost of dishonesty. *Proc. R.*
Soc. B: Biol. Sci. 281(1790), 20140876.

Birkhead, T., Pellatt, E., Matthews, I., Castillo-Juarez, H., 2006. Genic capture and the genetic basis of sexually
selected traits in the zebra finch. *Evolution* 60(11), 2389-2398.

Birnbaum, M., 2005. A comparison of five models that predict violations of first-order stochastic dominance in risky
decision making. *J. Risk Uncertain.* 31(3), 263-287.

Boag, D., Alway, J., 1981. Heritability of dominance status among Japanese quail: a preliminary report. *Can. J.*
Zool. 59(3), 441-444.

Bonilla, M., Zeh, J., Zeh, D., 2016. An epigenetic resolution of the lek paradox. *BioEssays* 38(4), 355-366.

Borgia, G., 1979. Sexual selection and the evolution of mating systems. In Blum, M., Blum, N. (eds.), *Sexual*
Selection and Reproductive Competition in Insects, pgs. 19-80. New York: Academic Press.

Bos, D., Williams, R., Gopurenko, D., Bulut, Z., Dewoody, J., 2009. Condition-dependent mate choice and a
reproductive disadvantage for MHC-divergent male tiger salamanders. *Mol. Ecol.* 18(15), 3307-3315.

Bosch, O., Musch, W., Bredewold, R., Slattery, D., Neumann, I., 2007. Prenatal stress increases HPA axis activity
and impairs maternal care in lactating female offspring: implications for postpartum mood disorder.
Psychoneuroendocrinology 32(3), 267-278.

Brilman, E., Ormel, J., 2001. Life events, difficulties and onset of depressive episodes in later life. *Psychol. Med.*
31(5), 859-869.

Brumm, H., Zollinger, S., Slater, P., 2009. Developmental stress affects song learning but not song complexity and
vocal amplitude in zebra finches. *Behav. Ecol. Sociobiol.* 63(9), 1387-1395.

Burley, N., 1986. Sexual selection for aesthetic traits in species with biparental care. *Am. Nat.* 127(4), 415-445.

Burt, A., 1995. The evolution of fitness. *Evolution* (49), 1-8.

Byers, J., Waits, L., 2006. Good genes sexual selection in nature. *Proc. Natl. Acad. Sci. U.S.A.* 103(44), 16343-
16345.

Cameron, E., 2004. Facultative adjustment of mammalian sex ratios in support of the Trivers-Willard hypothesis:
evidence for a mechanism. *Proc. R. Soc. B: Biol. Sci.* 271(1549), 1723-1728.

Cavigelli, S., Chaudhry, H., 2012. Social status, glucocorticoids, immune function, and health: Can animal studies

help us understand human socioeconomic-status-related health disparities? *Horm. Behav.* 62(3), 295-313.

Champagne, F., Francis, D., Mar, A., Meaney, M., 2003. Variations in maternal care in the rat as a mediating

influence for the effects of environment on development. *Physiol. Behav.* (79), 359-371.

Chen, E., Miller, G., Kobor, M., Cole, S., 2011. Maternal warmth buffers the effects of low early-life socioeconomic

status on pro-inflammatory signaling in adulthood. *Mol. Psychiatry* 16(7), 729-737.

Chichinadze, K., Chichinadze, N., 2008. Stress-induced increase of testosterone: contributions of social status and

sympathetic reactivity. *Physiol. Behav.* 94(4), 595-603.

Clutton-Brock, T., Hodge, S., Spong, G., Russell, A., Jordan, N., Bennett, N., Sharpe, L., Manser, M., 2006.

Intrasexual competition and sexual selection in cooperative mammals. *Nature* (444), 1065-1068.

Clutton-Brock, T., Hodge, S., Flower, T., Spong, G., Young, A., 2010. Adaptive suppression of subordinate

reproduction in cooperative mammals. *Am. Nat.* 176(5), 664-673.

Cooney, D., Allen, B., Veller, C., 2016. Assortment and the evolution of cooperation in a Moran process with

exponential fitness. *J. Theor. Biol.* (409), 38-46.

Cooney, L., van Klink, J., Hughes, N., Perry, N., Schaefer, H., Menzies, I., Gould, K., 2012. Red leaf margins

indicate increased polygodial content and function as visual signals to reduce herbivory in Pseudowintera

colorata. *New Phytol. 19*4(2), 488-497.

Crawley, J., Sutton, M., Pickar, D., 1985. Animal models of self-destructive behavior and suicide. *Psychiatr. Clin.*

North Am. 8(2), 299-310.

Curley, J., Mashoodh, R., Champagne, F., 2011. Epigenetics and the origins of paternal effects. *Horm. Behav.* 59(3),

306-314.

Dakin, R., Montgomerie, R., 2014. Condition-dependent mate assessment and choice by peahens: implications for

sexual selection. *Behav. Ecol.* 25(5), 1097-1104.

Dawkins, R., 1976. *The Selfish Gene.* Oxford: Oxford University Press.

Denenberg, V., Karas, G., 1959. Effects of differential infantile handling upon weight gain and mortality in the rat
and mouse. *Science* 130(3376), 629-630.

Dertwinkel-Kalt, M., Koster, M., 2015. Violations of first-order stochastic dominance as salience effects. *J. Behav.*
Exp. Econ. (59), 42-46.

Dewsbury, D., 1990. Fathers and sons: genetic factors and social dominance in deer mice, Peromyscus maniculatus.
Anim. Behav. 39(2), 284-289.

Dietz, D., LaPlant, Q., Watts, E., Hodes, G., Russo, S., Feng, J., Oosting, R., Vialou, V., Nestler, E., 2011. Paternal
transmission of stress-induced pathologies. *Biol. Psychiatry* 70(5), 408-414.

Dietz, D., Nestler, E., 2012. From father to offspring: paternal transmission of depressive-like behaviors.
Neuropsychopharmacology 37(1), 311-312.

Dong, Q., Salva, A., Sottas, C., Niu, E., Holmes, M., Hardy, M., 2004. Rapid glucocorticoid mediation of
suppressed testosterone biosynthesis in male mice subjected to immobilization stress. *J. Androl.* 25(6), 973-
981.

Dowlati, Y., Herrmann, N., Swardfager, W., Liu, H., Sham, L., Reim, E., Lanctot, K., 2010. A meta-analysis of
cytokines in major depression. *Biol. Psychiatry* (67), 446-457.

Drury, S., Sanchez, M., Gonzalez, A., 2016. When mothering goes awry: challenges and opportunities for utilizing
evidence across rodent, nonhuman primate and human studies to better define the biological consequences
of negative early caregiving. *Horm. Behav.* (77), 182-192.

Dugatkin, L., 1997. Winner and loser effects and the structure of dominance hierarchies. *Behav. Ecol.* 8(6), 583-587.

Edward, D., Chapman, T., 2011. The evolution and significance of male mate choice. *Trends Ecol. Evol.* 26(12),
647-654.

Edward, D., Chapman, T., 2012. Measuring the fitness benefits of male mate choice in Drosophila melanogaster.
Evolution 66(8), 2646-2653.

Edward, D., Chapman, T., 2013. Variation in male mate choice in Drosophila melanogaster. *Plos One* 8(2), e56299.

Emlen, D., Warren, I., Johns, A., Dworkin, I., Lavine, L., 2012. A mechanism of extreme growth and reliable

signaling in sexually selected ornaments and weapons. *Science* 337(6096), 860-864.

Fedurek, P., Slocombe, K., Enigk, D., Thompson, M., Wrangham, R., Muller, M., 2016. The relationship between

testosterone and long-distance calling in wild male chimpanzees. *Behav. Ecol. Sociobiol.* 70(5), 659-672.

Field, J., Solis, C., Queller, D., Strassmann, J., 1998. Social and genetic structure of paper wasp cofoundress

associations: tests of reproductive skew models. *Am. Nat.* 151(6), 545-563.

Field, T., 1998. Touch therapy effects on development. *Int. J. Behav. Dev.* 22(4), 779-797.

Field, T., Schanberg, S., Scafidi, F., Bauer, C., Vega-Lahr, N., Garcia, R., Nystrom, J., Kuhn, C., 1986.

Tactile/kinesthetic stimulation effects on preterm neonates. *Pediatrics* 77(5), 654-658.

Figueiredo, I., Frota, P., da Cunha, D., da Silva Raposo, R., Canuto, K., de Andrade, G., Sousa, N., Moore, S.,

Anstead, G., Alvarez-Leite, J., Guerrant, R., Oria, R., 2016. Prolonged maternal separation induces

undernutrition and systematic inflammation with disrupted hippocampal development in mice. *Nutrition*

32(9), 1019-1027.

Fisher, R., 1930. *The Genetical Theory of Natural Selection.* Oxford: The Clarendon Press.

Folstad, I., Karter, A., 1992. Parasites, bright males, and the immunocompetence handicap. *Am. Nat.* 139(3), 603-

622.

Franklin, T., Linder, N., Russig, H., Thony, B., Mansuy, I., 2011. Influence of early stress on social abilities and

serotonergic functions across generations in mice. *Plos One* 6(7), e21842.

Franklin, T., Russig, H., Weiss, I., Graff, J., Linder, N., Michalon, A., Vizi, S., Mansuy, I., 2010. Epigenetic

transmission of the impact of early stress across generations. *Biol. Psychiatry* 68(5), 408-415.

Franklin, T., Saab, B., Mansuy, I., 2012. Neural mechanisms of stress resilience and vulnerability. *Neuron* 75(5),

747-761.

Georgiev, A., Muehlenbein, M., Prall, S., Thompson, M., Maestripieri, D., 2015. Male quality, dominance rank, and

mating success in free-ranging rhesus macaques. *Behav. Ecol.* 26(3), 763-772.

Getty, T., 2006. Sexually selected signals are not similar to sports handicaps. *Trends Ecol. Evol.* 21(2), 83-88.

Ghaziuddin, M., Ghaziuddin, N., Stein, G., 1990. Life events and the recurrence of depression. *Can. J. Psychiatry*
1426   35(3), 239-242.

1427   Gil, D., 2008. Hormones in avian eggs: physiology, ecology and behavior. In Brockmann, H., Roper, T., Naguib,
1428   M., Wynne-Edwards, K., Barnard, C., Mitani, J. (eds.), *Advances in the Study of Behavior, Volume 38*, pgs.
1429   337-398. San Diego, CA: Academic Press.

1430   Gil, D., Graves, J., Hazon, N., Wells, A., 1999. Male attractiveness and differential testosterone investment in zebra
1431   finch eggs. *Science* 286(5437), 126-128.

1432   Gillespie, J., 1973. Natural selection with varying selection coefficients—a haploid model. *Genet. Res.* (21), 115-
1433   120.

1434   Gillespie, J., 1974. Natural selection for within-generation variance in offspring number. *Genetics* 76(3), 601-606.

1435   Gleason, E., Fuxjager, M., Oyegbile, T., Marler, C., 2009. Testosterone release and social context: When it occurs
1436   and why. *Front. Neuroendocrinol.* 30(4), 460-469.

1437   Gould, K., Davies, K., Winefield, C., 2009. *Anthocyanins: Biosynthesis, Functions, and Applications.* New York:
1438   Springer-Verlag.

1439   Gowaty, P., 2003. Power asymmetries between the sexes, mate preferences, and components of fitness. In Travis, C.
1440   (ed.), *Women, Evolution, and Rape*, pgs. 61-86. Cambridge, MA: MIT Press.

1441   Gowaty, P., 2008. Reproductive compensation. *J. Evol. Biol.* 21(5), 1189-1200.

1442   Gowaty, P., Anderson, W., Bluhm, C., Drickamer, L., Kim, Y., Moore, A., 2007. The hypothesis of reproductive
1443   compensation and its assumptions about mate preferences and offspring viability. *Proc. Natl. Acad. Sci.*
1444   *U.S.A.* 104(38), 15023-15027.

1445   Gracia-Rubio, I., Moscoso-Castro, M., Pozo, O., Marcos, J., Nadal, R., Valverde, O., 2016. Maternal separation
1446   induces neuroinflammation and long-lasting emotional alterations in mice. *Prog. Neuropsychopharmacol.*
1447   *Biol. Psychiatry* (65), 104-117.

Grafen, A., 1979. The hawk-dove game played between relatives. *Anim. Behav.* (27), 905-907.

Grafen, A., 1990. Biological signals as handicaps. *J. Theor. Biol.* 144(4), 517-546.

Groothuis, T., Muller, W., von Engelhardt, N., Carere, C., Eising, C., 2005. Maternal hormones as a tool to adjust
offspring phenotype in avian species. *Neurosci. Biobehav. Rev.* 29(2), 329-352.

Habig, B., Archie, E., 2015. Social status, immune response and parasitism in males: a meta-analysis. *Philos. Trans.*
Royal Soc. B 370(1669), 20140109.

Hagen, E., 2003. The bargaining model of depression. In Hammerstein, P. (ed.), *Genetic and Cultural Evolution of*
Cooperation, pgs. 97-123. Cambridge: The MIT Press.

Hamilton, W., 1964. The genetical evolution of social behavior. I. *J. Theor. Biol.* 7(1), 1-16.

Hanoch, G., Levy, H., 1969. The efficiency analysis of choices involving risk. *Rev. Econ. Stud.* 36(3), 335-346.

Harris, W., Uller, T., 2009. Reproductive investment when mate quality varies: differential allocation versus
reproductive compensation. *Philos. Trans. Royal Soc. B* 364(1520), 1039-1048.

Harrison, X., York, J., Cram, D., Hares, M., Young, A., 2013. Complete reproductive skew within white-browed
sparrow weaver groups despite outbreeding opportunities for subordinates of both sexes. *Behav. Ecol.*
Sociobiol. 67(12), 1915-1929.

Hawton, K., van Heeringen, K., 2009. Suicide. *Lancet* 373(9672), 1372-1381.

Heiming, R., Bodden, C., Jansen, F., Lewejohann, L., Kaiser, S., Lesch, K., Palme, R., Sachser, N., 2011. Living in
a dangerous world decreases maternal care: a study in serotonin transporter knockout mice. *Horm. Behav.*
60(4), 397-407.

Hill, G., 2006. Environmental regulation of ornamental coloration. In Hill, G., McGraw, K. (eds.), *Bird Coloration,*
Volume 1, pgs. 507-560. Cambridge, MA: Harvard University Press.

Hill, G., 2011. Condition-dependent traits as signals of the functionality of vital cellular processes. *Ecol. Lett.* 14(7),
625-634.

Hill, G., 2014. Cellular respiration: the nexus of stress, condition, and ornamentation. *Integr. Comp. Biol.* 54(4), 645-657.

Holveck, M., Geberzahn, N., Riebel, K., 2011. An experimental test of condition-dependent male and female mate choice in zebra finches. *Plos One* 6(8), e23974.

Holveck, M., Riebel, K., 2010. Low-quality females prefer low-quality males when choosing a mate. *Proc. R. Soc. B: Biol. Sci.* (277), 153-160.

Hsu, Y., Earley, R., Wolf, L., 2006. Modulation of aggressive behaviour by fighting experience: mechanisms and contest outcomes. *Biol. Rev.* 81(1), 33-74.

Hunt, J., Bussiere, L., Jennions, M., Brooks, R., 2004. What is genetic quality? *Trends Ecol. Evol. 19*(6), 329-333.

Husak, J., Swallow, J., 2011. Compensatory traits and the evolution of male ornaments. *Behaviour* 148(1), 1-29.

Irving, G., Lloyd-Williams, W., 2010. Depression in advanced cancer. *Eur. J. Oncol. Nurs.* 14(5), 395-399.

Iwasa, Y., Pomiankowski, A., 1994. The evolution of mate preferences for multiple sexual ornaments. *Evolution* 48(3), 853-867.

Jarrow, R., 1986. The relationship between arbitrage and first order stochastic dominance. *J. Finance* 41(4), 915-921.

Johnstone, R., 1995. Sexual selection, honest advertisement and the handicap principle: reviewing the evidence. *Biol. Rev.* 70(1), 1-65.

Johnstone, R., 2000. Models of reproductive skew: A review and synthesis. *Ethology* 106(1), 5-26.

Kankova, Z., Zeman, M., Okuliarova, M., 2014. Selection for high egg testosterone and immune response of young Japanese quail under mild food restriction. *Avian Biol. Res.* 7(1), 25-32.

Karageorgou, P., Buschmann, C., Manetas, Y., 2008. Red leaf color as a warning signal against insect herbivory: honest or mimetic? *Flora* 203(8), 648-652.

Kendler, K., Thornton, L., Gardner, C., 2000. Stressful life events and previous episodes in the etiology of major depression in women: an evaluation of the "kindling" hypothesis. *Am. J. Psychiatry* 157(8), 1243-1251.

Kinney, D., Tanaka, M., 2009. An evolutionary hypothesis of depression and its symptoms, adaptive value, and risk
factors. *J. Nerv. Ment. Dis. 19*7(8), 561-567.

Kirkpatrick, M., 1982. Sexual selection and the evolution of female choice. *Evolution* 36(1), 1-12.

Kirkpatrick, M., Ryan, M., 1991. The evolution of mating preferences and the paradox of the lek. *Nature* (350), 33-
38.

Kokko, H., Brooks, R., McNamara, J., Houston, A., 2002. The sexual selection continuum. *Proc. R. Soc. B: Biol.*
Sci. 269(1498), 1331-1340.

Korosi, A., Baram, T., 2009. The pathways from mother's love to baby's future. *Front. Behav. Neurosci.* 3(27).
DOI: 10.3389/neuro.08.027.2009

Kotiaho, J., Simmons, L., Tomkins, J., 2001. Towards a resolution of the lek paradox. *Nature* (410), 684-686.

Kovinich, N., Kayanja, G., Chanoca, A., Riedl, K., Otegui, M., Grotewold, E., 2014. Not all anthocyanins are born
equal: distinct patterns induced by stress in Arabidopsis. *Planta* 240(5), 931-940.

Lee, K., 2006. Linking immune defenses and life history at the levels of the individual and the species. *Integr.*
Comp. Biol. 46(6), 1000-1015.

Lynn, S., Perfito, N., Guardado, D., Bentley, G., 2015. Food, stress, and circulating testosterone: cue integration by
the testes, not the brain, in male zebra finches (Taeniopygia guttata). *Gen. Comp. Endocrinol.* (215), 1-9.

Manetas, Y., 2006. Why some leaves are anthocyanic and why most anthocyanic leaves are red? *Flora* 201(3), 163-
177.

Mansuy, I., Mashoodh, R., Champagne, F., 2013. Transgenerational inheritance in mammals. In Sweatt, J., Meaney,
M., Nestler, E., Akbarian, S. (eds.), *Epigenetic Regulation in the Nervous System*: *Basic Mechanisms and*
Clinical Impact, pgs. 323-338. London: Academic Press.

Mashoodh, R., Franks, B., Curley, J., Champagne, F., 2012. Paternal social enrichment effects on maternal behavior
and offspring growth. *Proc. Natl. Acad. Sci. U.S.A.* (109), 17232-17238.

Mashoodh, R., Sinal, C., Perrot-Sinal, T., 2009. Predation threat exerts specific effects on rat maternal behaviour

and anxiety-related behaviour of male and female offspring. *Physiol. Behav.* 96(4-5), 693-702.

Maynard Smith, J., 1978. *The Evolution of Sex.* Cambridge: Cambridge University Press.

Maynard Smith, J., Harper, D., 1995. Animal signals: models and terminology. *J. Theor Biol.* 177(3), 305-311.

Mazur, A., Lamb, T., 1980. Testosterone, status, and mood in human males. *Horm. Behav.* 14(3), 236-246.

McCaul, K., Gladue, B., Joppa, M., 1992. Winning, losing, mood, and testosterone. *Horm. Behav.* 26(4), 486-504.

McGraw, K., Blount, J., 2009. Control and function of carotenoid coloration in birds: selected case studies. In

Landrum, J. (ed.), *Carotenoids: Physical, Chemical, and Biological Functions and Properties*, pgs. 487-

510, Boca Raton, FL: CRC Press.

McLeod, J., Sinal, C., Perrot-Sinal, T., 2007. Evidence for non-genomic transmission of ecological information via

maternal behavior in female rats. *Genes Brain Behav.* 6(1), 19-29.

Meaney, M., 2001. Maternal care, gene expression, and the transmission of individual differences in stress reactivity

across generations. *Neuroscience* (24), 1161-1192.

Medawar, P., 1952. *An Unsolved Problem in Biology*. London: H.K. Lewis.

Menzies, I., Youard, L., Lord, J., Carpenter, K., van Klink, J., Perry, N., Schaefer, H., Gould, K., 2016. Leaf colour

polymorphisms: a balance between plant defence and photosynthesis. *J. Ecol.* 104(1), 104-113.

Merila, J., Sheldon, B., 2000. Lifetime reproductive success and heritability in nature. *Am. Nat.* 155(3), 301-310.

Miller, G., Chen, E., Parker, K., 2011. Psychological stress in childhood and susceptibility to the chronic diseases of

aging: moving toward a model of behavioral and biological mechanisms. *Psychol. Bull.* 137(6), 959-997.

Moller, A., Alatalo, R., 1999. Good-genes effects in sexual selection. *Proc. R. Soc. B: Biol. Sci.* 266(1414), 85-91.

Moore, A., 1990. The inheritance of social dominance, mating behaviour and attractiveness to mates in male

Nauphoeta cinerea. *Anim. Behav.* 39(2), 388-397.

Muehlenbein, M., Bribiescas, R., 2005. Testosterone-mediated immune functions and male life histories. *Am. J.*
Hum. Biol. 17(5), 527-558.

Mykletun, A., Bjerkeset, O., Dewey, M., Prince, M., Overland, S., Stewart, R., 2007. Anxiety, depression, and
cause-specific mortality: the HUNT study. *Psychosom. Med.* 69(4), 323-331.

Nelson, R., Frank, D., Bennett, S., Carter, C., 1989. Simulated drought influences reproduction in male prairie voles.
Physiol. Behav. 46(5), 849-852.

Nestler, E., 2012. Epigenetics: stress makes its molecular mark. *Nature* 490(7419), 171-172.

Nettle, D., 2004. Evolutionary origins of depression: a review and reformulation. *J. Affect. Disord.* 81(2), 91-102.

Nonacs, P., Hager, R., 2011. The past, present and future of reproductive skew theory and experiments. *Biol. Rev.*
86(2), 271-298.

Nowak, M., 2006. Five rules for the evolution of cooperation. *Science* 314(5805), 1560-1563.

Nowak, M., 2012. Evolving cooperation. *J. Theor. Biol.* (299), 1-8.

Okasha, S., Martens, J., 2016. Hamilton's rule, inclusive fitness maximization, and the goal of individual behaviour
in symmetric two-player games. *J. Evol. Biol.* 29(3), 473-482.

Oliveira, R., 2004. Social modulation of androgens in vertebrates: mechanisms and function. In Slater, P.,
Rosenblatt, J., Snowdon, C. (eds.), *Advances in the Study of Behavior, Volume 34*, pgs. 165-239. San
Diego: Elsevier Academic Press.

Parker, G., Brotchie, H., 2017. Pancreatic cancer and depression: a narrative review. *J. Nerv. Ment. Dis.* 205(6),
487-490.

Parker, T., Garant, D., 2004. Quantitative genetics of sexually dimorphic traits and capture of genetic variance by a
sexually-selected condition-dependent ornament in red junglefowl (Gallus gallus). *J. Evol. Biol.* 17(6),
1277-1285.

Pittet, F., Coignard, M., Houdelier, C., Richard-Yris, M., Lumineau, S., 2012. Age affects the expression of
maternal care and subsequent behavioural development of offspring in a precocial bird. *Plos One* 7(5),
e36835.

Pomiankowski, A., Moller, A., 1995. A resolution of the lek paradox. *Proc. R. Soc. B: Biol. Sci.* 260(1357), 21-29.

Preti, A., 2011. Animal model and neurobiology of suicide. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 35(4),
818-830.

Pryce, C., Fontana, A., 2017. Depression in autoimmune diseases. *Curr. Top. Behav. Neurosci.* (31), 139-154.

Raison, C., Miller, A., 2013. The evolutionary significance of depression in pathogen host defense (PATHOS-D).
Mol. Psychiatry 18(1), 15-37.

Ramchandani, P., Stein, A., Evans, J., O'Connor, T., 2005. Paternal depression in the postnatal period and child
development: a prospective population study. *Lancet* 365(9478), 2201-2205.

Ratikainen, I., Kokko, H., 2010. Differential allocation and compensation: Who deserves the silver spoon? *Behav.*
Ecol. 21(1), 195-200.

Refardt, D., Bergmiller, T., Kummerli, R., 2013. Altruism can evolve when relatedness is low: evidence from
bacteria committing suicide upon phage infection. *Proc. R. Soc. B: Biol. Sci.* 280(1759), 20123035.

Ritschard, M., Laucht, S., Dale, J., Brumm, H., 2011. Enhanced testosterone levels affect singing motivation but not
song amplitude in Bengalese finches. *Physiol. Behav.* 102(1), 30-35.

Ritschard, M., Riebel, K., Brumm, H., 2010. Female zebra finches prefer high-amplitude song. *Anim. Behav.* 79(4),
877-883.

Roberts, M., Buchanan, K., Evans, M., 2004. Testing the immunocompetence handicap hypothesis: a review of the
evidence. *Anim. Behav.* 68(2), 227-239.

Rowe, L., Houle, D., 1996. The lek paradox and the capture of genetic variance by condition dependent traits. *Proc.*
R. Soc. B: Biol. Sci. 263(1375), 1415-1421.

Russo, S., Murrough, J., Han, M., Charney, D., Nestler, E., 2012. Neurobiology of resilience. *Nat. Neurosci.* 15(11),
1475-1484.

Rutte, C., Taborsky, M., Brinkhof, M., 2006. What sets the odds of winning and losing? *Trends Ecol. Evol.* 21(1),
16-21.

Saltzman, W., Digby, L., Abbott, D., 2009. Reproductive skew in female common marmosets: What can proximate
mechanisms tell us about ultimate causes? *Proc. R. Soc. B: Biol. Sci.* 276(1656), 389-399.

Sapolsky, R., 1998. *Why Zebras Don't Get Ulcers. 2nd edition.* New York: W.H. Freeman and Company.

Sapolsky, R., 2001. Physiological and pathophysiological implications of social stress in mammals. In McEwen, B.
(ed.), *Handbook of Physiology: Section 7, The Endocrine System; Volume IV: Coping With The*
Environment, pgs. 517-532. Oxford: Oxford University Press.

Schaefer, H., Rolshausen, G., 2006. Plants on red alert: Do insects pay attention? *BioEssays* 28(1), 65-71.

Schanberg, S., 1995. The genetic basis for touch effects. In Field, T. (ed.), *Touch in Early Development*, pgs. 67-80.
Mahwah, NJ: Lawrence Erlbaum Associates.

Schreiber, S., 2015. Unifying within- and between-generation bet-hedging theories: an ode to J.H. Gillespie. *Am.*
Nat. 186(6), 792-796.

Schuch, F., Vancampfort, D., Richards, J., Rosenbaum, S., Ward, P., Stubbs, B., 2016. Exercise as a treatment for
depression: a meta-analysis adjusting for publication bias. *J. Psychiatr. Res.* (77), 42-51.

Schultzhaus, J., Nixon, J., Duran, J., Carney, G., 2017. Diet alters Drosophila melanogaster mate preference and
attractiveness. *Anim. Behav.* (123), 317-327.

Seger, J., Brockmann, H., 1987. What is bet-hedging? In Harvey, P., Partridge, L. (eds.), *Oxford Surveys In*
Evolutionary Biology, Volume 4, pgs. 182-211. Oxford: Oxford University Press.

Seligman, M., 1976. *Helplessness: On Depression, Development, and Death.* San Francisco: W.H. Freeman and
Company.

Sheldon, B., 2000. Differential allocation: tests, mechanisms and implications. *Trends Ecol. Evol.* 15(10), 397-402.

Shen, H., Magnusson, C., Rai, D., 2016. Associations of parental depression with child school performance at age

16 years in Sweden. *JAMA Psychiatry* 73(3), 239-246.

Simons, M., Groothuis, T., Verhulst, S., 2015. An appraisal of how the vitamin A-redox hypothesis can maintain

honesty of carotenoid-dependent signals. *Ecol. Evol.* 5(1), 224-228.

Simons, M., Maia, R., Leenknegt, B., Verhulst, S., 2014. Carotenoid-dependent signals and the evolution of plasma

carotenoid levels in birds. *Am. Nat.* 184(6), 741-751.

Slatkin, M., 1974. Hedging one's evolutionary bets. *Nature* (250), 704-705.

Solomon, D., Keller, M., Leon, A., Mueller, T., Lavori, P., Shea, M., Coryell, W., Warshaw, M., Turvey, C., Maser,

J., Endicott, J., 2000. Multiple recurrences of major depressive disorder. *Am. J. Psychiatry* 157(2), 229-233.

Starrfelt, J., Kokko, H., 2012. Bet-hedging—a triple trade-off between means, variances and correlations. *Biol. Rev.*

87(3), 742-755.

Stiver, K., Alonzo, S., 2009. Parental and mating effort: Is there necessarily a trade-off? *Ethology* 115(12), 1101-

1126.

Sun, H., Kennedy, P., Nestler, E., 2013. Epigenetics of the depressed brain: role of histone acetylation and

methylation. *Neuropsychopharmacology* 38(1), 124-137.

Suomi, S., 1997. Early determinants of behaviour: evidence from primate studies. *Br. Med. Bull.* 53(1), 170-184.

Szamado, S., 2011. The cost of honesty and the fallacy of the handicap principle. *Anim. Behav.* 81(1), 3-10.

Taylor, P., Williams, G., 1982. The lek paradox is not resolved. *Theor. Popul. Biol.* 22(3), 392-409.

Trivers, R., 1971. The evolution of reciprocal altruism. *Q. Rev. Biol.* 46(1), 35-57.

Trivers, R., Willard, D., 1973. Natural selection of parental ability to vary the sex ratio of offspring. *Science*

179(4068), 90-92.

Van Cleve, J., Akcay, E., 2014. Pathways to social evolution: reciprocity, relatedness, and synergy. *Evolution* 68(8),

2245-2258.

Varga, S., 2012. Evolutionary psychiatry and depression: testing two hypotheses. *Med. Health Care Philos.* 15(1), 41-52.

Veen, T., Otto, S., 2015. Liking the good guys: amplifying local adaptation via the evolution of condition-dependent mate choice. *J. Evol. Biol.* 28(10), 1804-1815.

Vehrencamp, S., 1983. A model for the evolution of despotic versus egalitarian societies. *Anim. Behav.* (31), 667-682.

Watson, P., Andrews, P., 2002. Toward a revised evolutionary adaptationist analysis of depression: the social navigation hypothesis. *J. Affect. Disord.* 72(1), 1-14.

Weatherhead, P., Robertson, R., 1981. In defense of the "sexy son" hypothesis. *Am. Nat.* 117(3), 349-356.

Weaver, R., Koch, R., Hill, G., 2017. What maintains signal honesty in animal colour displays used in mate choice? *Philos. Trans. Royal Soc. B* 372(1724), 20160343.

Weiss, I., Franklin, T., Vizi, S., Mansuy, I., 2011. Inheritable effect of unpredictable maternal separation on behavioral responses in mice. *Front. Behav. Neurosci.* 5(3). DOI: 10.3389/fnbeh.2011.00003

Weiss, J., Bailey, W., Goodman, P., Hoffman, L., Ambrose, M., Salman, S., Charry, J., 1982. A model for neurochemical study of depression. In Spiegelstein, M., Levy, A. (eds.), *Behavioral Models and the Analysis of Drug Action*, pgs. 195-223. Amsterdam: Elsevier Scientific Publishing Company.

West, S., Griffin, A., Gardner, A., 2007. Social semantics: altruism, cooperation, mutualism, strong reciprocity and group selection. *J. Evol. Biol.* 20(2), 415-432.

West, S., El Mouden, C., Gardner, A., 2011. Sixteen common misconceptions about the evolution of cooperation in humans. *Evol. Hum. Behav.* 32(4), 231-262.

Wilson, A., Nussey, D., 2010. What is individual quality? An evolutionary perspective. *Trends Ecol. Evol.* 25(4), 207-214.

Woodgate, J., Mariette, M., Bennett, A., Griffith, S., Buchanan, K., 2012. Male song structure predicts reproductive success in a wild zebra finch population. *Anim. Behav.* 83(3), 773-781.

Yirmiya, R., Weidenfeld, J., Pollak, Y., Morag, M., Morag, A., Avitsur, R., Barak, O., Reichenberg, A., Cohen, E.,

Shavit, Y., Ovadia, H., 1999. Cytokines, "depression due to a general medical condition," and

antidepressant drugs. In Dantzer, R., Wollman, E., Yirmiya, R. (eds.), *Cytokines, Stress, and Depression.*

Advances in Experimental Medicine and Biology (461), 283-316.

Zahavi, A., 1975. Mate selection—a selection for a handicap. *J. Theor. Biol.* 53(1), 205-214.

Zahavi, A., Zahavi, A., 1997. *The Handicap Principle: A Missing Piece of Darwin's Puzzle.* Oxford: Oxford

University Press.

Zann, R., Cash, E., 2008. Developmental stress impairs song complexity but not learning accuracy in non-

domesticated zebra finches (Taeniopygia guttata). *Behav. Ecol. Sociobiol.* 62(3), 391-400.

Zhang, T., Meaney, M., 2010. Epigenetics and the environmental regulation of the genome and its function. *Annu.*

Rev. Psychol. (61), 439-466.

Zollinger, S., Goller, F., Brumm, H., 2011. Metabolic and respiratory costs of increasing song amplitude in zebra

finches. *Plos One* 6(9), e23198.

1669