Understanding How Microbiomes Influence the Systems they Inhabit: Insight from Ecosystem Ecology

Authors: Hall, E.K.[1*], Bernhardt, E.S[2], Bier, R.L.[2], Bradford, M.A.[3], Boot, C.M.[1], Cotner, J.B.[4], del Giorgio, P.A.[5], Evans, S.E.[6], Graham E. B[2,7,8], Jones, S.E.[9], Lennon, J.T.[10], Locey, K.[10] Nemergut, D.[2], Osborne, B.[11], Rocca, J.D.[1,2], Schimel J.S.[12], Waldrop, M.P.[13], Wallenstein, M.W.[1]

Article Type: Perspectives
Words in Abstract: 272
Words in Main Text: 4,431
2 Figures, 0 Tables, 51 References

Running Title: How microbiomes influence ecosystems
Keywords: microbiome, microbial ecology, ecosystem ecology, emergent properties, community aggregated traits

Affiliations: [1]Colorado State University, Fort Collins CO (ed.hall@colostate.edu, claudia.boot@colostate.edu, matt.wallenstein@colostate.edu) [2]Duke University, Durham, N.C. (ebernhar@duke.edu, ebgraham2@colorado.edu, diana.nemergut@duke.edu) [3]Yale University, New Haven, CT (mark.bradford.yale@gmail.com) [4]University of Minnesota, Saint Paul, MN (cotne002@umn.edu) [5]Université du Québec à Montréal, Montréal, CA (del_giorgio.paul@uqam.ca) [6]Michigan State University, Hickory Corners, MI (evanssar@gmail.com) [7]Institute of Arctic and Alpine Research, University of Colorado at Boulder [8] Pacific Northwest National Laboratory, Richland, WA, USA [9]Notre Dame University, South Bend, IN (sjones20@nd.edu) [10]Indiana University, Bloomington, IN (lennonj@indiana.edu, ken@weecology.org) [11]Brown University, Providence, Rhode Island (brooke_osborne@brown.edu) [12]University of California, Santa Barbara, Santa Barbara, CA (josh.schimel@lifesci.ucsb.edu) [13]U.S. Geological Survey, Menlo Park, CA (mwaldrop@usgs.gov) Biological Sciences Division,

* **corresponding author**: Dr. Ed Hall, Natural Resource Ecology Laboratory, Department of Ecosystem Science and Sustainability, Campus Delivery 1499, Colorado State University, Fort Collins, CO 80523, ed.hall@colostate.edu, 970-491-2162

Contribution: All listed authors have contributed to the conceptualization, writing, and preparation of the current manuscript.

Abstract

The well-documented significance of microorganisms to the function of virtually all

ecosystems has led to the assumption that more information on microbiomes will

improve our ability to understand and predict system-level processes. Notably, the

importance of the microbiome has become increasingly evident in the environmental

sciences and in particular ecosystem ecology. However, translating the ever-

increasing wealth of information on environmental microbiomes to advance

ecosystem science is proving exceptionally challenging. One reason for this

challenge is that correlations between microbiomes and the ecosystem processes

they influence are often reported without the underlying causal mechanisms. This

limits the predictive power of each correlation to the time and place at which it was

identified. In this paper, we assess the assumptions and approaches currently used

to establish links between environmental microbiomes and the ecosystems they

influence, propose a framework to more effectively harness our understanding of

microbiomes to advance ecosystem science, and identify key challenges and

solutions required to apply the proposed framework. Specifically, we suggest

identifying each microbial process that contributes to the ecosystem process of

interest *a priori.* We then suggest linking information on microbial community

membership through microbial community properties (such as biomass elemental

ratios) to the microbial processes that drive each ecosystem process (e.g. N -

mineralization). A key challenge in this framework will be identifying which microbial

community properties can be determined from the constituents of the community

(community aggregated traits, CATs) and which properties are unable to be

predicted from a list of their constituent taxa (emergent properties, EPs). We view

this directed approach as a promising pathway to advance our understanding of how

microbiomes influence the systems they inhabit.

Current approaches in linking microbial characteristics and ecosystem processes

Recently there has been a broad call, including the National Microbiome Initiative

led by the executive branch of the United States Federal Government, for a

coordinated effort to evaluate the role of microorganisms in all environments[1,2].

Coordinating efforts to explore microbiomes and their functioning across such a

broad range of systems is exciting and ambitious and holds the potential to transform

societies approach many of the most important challenges we currently face[3].

However, advances in this direction require an assessment of our progress to date

and an attempt to identify the most promising paths forward.

In most ecosystems, many processes are carried out primarily by microorganisms,

and virtually all processes are influenced by microorganisms. Therefore it is common

to apply high-resolution analytical techniques to better describe microbial

communities, assuming that greater resolution of the community, (including its

associated transcripts, proteins, and metabolic products) should lead to better

predictions of ecosystem processes. However, such justifications assume that

microbial metrics (e.g. 16S rRNA gene libraries, metagenomes, enzymatic activities)

will improve our ability to understand, model, and predict ecosystem processes. This

assumption may not necessarily be valid, because microbial information may not

provide additional explanatory power for understanding ecosystem process rates

beyond what can be predicted by environmental factors alone[4,5]. In addition, when

correlations between microbial information and an ecosystem process are identified

the underlying causal associations may remain difficult to resolve and may not be

causal at all—e.g. the microbiome and the ecosystem may be responding in concert

to a common underlying driver; limiting the predictive ability of each result across

additional systems.

Two recent meta-analyses suggest that the current research approach at the

intersection of ecosystem and microbial ecology has the potential to be better

focused to more effectively achieve the anticipated insights into how microbiomes

influence ecosystems[6,7]. The first meta-analysis evaluated studies that related the

relative abundance of protein encoding gene copy or transcript abundance to

associated biogeochemical processes[6]. Of 416 identified studies that attempted to

address the correlation between the relative abundance of a protein-encoding gene

(or transcript) and an ecosystem process, only 56 measured both genes or transcript

copy number and the corresponding process. Within these 56 studies, 14% of the

observations showed a significant negative correlation between gene copy number

and process rate, 38% had a significant positive relationship, and 48% had no

significant relationship. Thus, the effect size for the relationship between gene copy

number and process rate had an approximately normal distribution with a mean near

zero[6]. The second meta-analysis evaluated links between microbial community

composition and ecosystem processes in response to an experimentally-induced

disturbance in the environment (e.g. drought, warming, nutrient addition)[7]. Whereas

40% of published papers reported concomitant changes in microbial community

structure and ecosystem function, only about a third of those cases (only 12% of

total studies) attempted to identify a statistical relationship between community

composition and an ecosystem process. Interestingly, many of the studies that did

not measure both community composition and a corresponding ecosystem process

still framed their study in the microbial structure-ecosystem function framework[7].

These meta-analyses illustrate that links between microbial characteristics and

ecosystem processes are often assumed to be present but are rarely tested. When

linkages are explicitly tested, connections between microbial structure and

ecosystem processes are more often than not weak or non-existent[6]. These findings

suggest that our current approach to linking microbial metrics to ecosystem function

should be refocused with more attention paid to empirically identifying explicit

linkages between microbial characteristics and the ecosystem processes that they

influence.

Linking microbial characteristics and ecosystem processes

A key challenge in linking microbial information to an ecosystem process is that

conceptual research frameworks often do not align directly with the available

methods or the data they generate. For example, environmental factors act on the

physiology of individual organisms, which alters their competitive ability, relative

abundance, collective physiology, and ultimately their contribution to ecosystem

processes (Figure 1a). However, designing an observational study or experiment

from this conceptual framework (Figure 1a) assumes that environmental metrics can

be empirically linked to measureable microbial characteristics across multiple

categories of ecological organization (i.e. individuals, populations, and communities)

at the appropriate temporal and spatial scales. Yet in nature, relationships between

environmental variables and microbial characteristics are dynamic and non-linear[8],

simultaneously affected by a plethora of biotic and abiotic variables[9,10] and

decoupled in both time and space[7]. Each of these aspects of microbial-environment

interactions obscures the relationships among microbial characteristics collected at

each level of ecological organization and the ecosystem processes they affect

(Figure 2b). To address this challenge we propose a framework that explicitly

identifies the ecosystem process of interest and how it relates to microbial

characteristics. The proposed framework illustrates the relationship among different

categories of microbial characteristics and conceptually defines their contribution to

ecosystem processes.

Identifying the Ecosystem Process The first step to understand how microorganisms

influence an ecosystem process is to define each of its sub-processes, the set of

constituent reactions that combine to dominate the net flux of the ecosystem process

of interest. Ecosystem processes are defined as a change in a pool size or a flux

from one pool to another (e.g. NH_4^+ to NO_3^-, or dissolved organic matter

mineralization to CO_2). Few, if any of these processes are carried out by a single

physiological pathway or a single organism. Rather, ecosystem processes are

aggregate processes consisting of complementary or antagonistic sub-processes

carried out by a breadth of phylogenetically diverse microorganisms[11]. For example,

net ecosystem productivity (NEP) is the balance between C-fixation and C-

mineralization. Each sub-process of NEP can be further partitioned into a series of

metabolic pathways (e.g. heterotrophic fermentation and aerobic respiration or

chemoautotrophic nitrification and photoautotrophic C-fixation). Partitioning each

ecosystem process in this hierarchical manner can continue until the sub-process

maps directly to specific microbial metabolic pathways (e.g. acetoclastic

methanogenesis). Subsequently each of these metabolic pathways can be

categorized as either broad or narrow[12]. Broad processes are phylogenetically

common (i.e. widely distributed among taxa), whereas narrow processes are

phylogenetically conserved (i.e. limited to a specific subset of taxa). For example,

denitrification is broad, while both methanogenesis and methanotrophy are narrow

(with some notable exceptions[13]).

The second step is to identify the controls or constraints on each constituent sub-

process. For example, kinetics of a given metabolic pathway in a model organism

may help understand the rate limiting steps of a narrow process, but insights from

model organisms are much less likely to be useful for a broad process where

phenotypic variation among phylogenetically diverse organisms should be much

greater. Defining the ecosystem process, its critical sub-processes, and the known

phylogenetic distribution of the metabolic pathways that drive those sub-processes in

this manner creates an explicit conceptual pathway that directly links the ecosystem

process to the microorganisms that influence them. Once this conceptual pathway

has been identified a concerted empirical approach can be applied to investigate

how the microbiome influences the ecosystem process of interest.

Understanding the relationship between categories of microbial

characteristics

Categories of microbial characteristics At present, researchers are measuring a wide

variety of characteristics of microbial communities (e.g. sequence or relative

abundance of genes, transcripts or proteins, enzyme expression, and process rates).

Much of that work does not clearly articulate how these measurements differ in their

specificity, precision, or linkage among each other or how they inform the microbial

contribution to ecosystem processes. We propose that by categorizing microbial

characteristics into three distinct categories, 1) microbial processes, 2) microbial

community properties, and 3) microbial community membership (Figure 2), we can

frame how different metrics interact with each other and how they can elucidate the

microbial contribution to an ecosystem process.

Microbial Processes Microbial processes are the collective physiology of the

microbial community that drives changes in pools and fluxes of important elements

or compounds[14] (Figure 2). They are the level of microbial information that can most

readily be incorporated into ecosystem-level models because many of these

processes are the sub-processes that contribute to an overall pool or flux. Examples

include nitrogen fixation, denitrification, nitrification, phosphorus uptake and

immobilization, primary production, respiration, and carbon use efficiency. The rates

of many microbial processes can be estimated through physiological assays, and

while they do not open the "black box" of the microbial community, they do directly

quantify the microbial contribution to the transformation of substrates moving through

the box.

However, physiological assays to estimate microbial processes are often

logistically challenging, require experimental manipulations that inevitably deviate

from the *in situ* conditions, and often depend on the environment in which they were

measured. For example, the relationship between microbial process rates and

temperature vary across geographical temperature gradients[15] (enzyme activity) and

among seasons[16] (phosphorus use efficiency, PUE). Thus observations of the effect

of temperature on either enzyme activity or PUE are time and place dependent.

Therefore, without an underlying physiological mechanism, to accurately quantify the

microbial process the relationship between temperature and community physiology

must be measured through a direct assay at each location and at each time.

Because of these limitations, a microbial community property that can be measured

in situ or collected and preserved in the field for later analysis in the laboratory has

several advantages over using bioassays to directly measure microbial processes. A

community property may include characteristics of community biomass such as

elemental ratios (biomass C:N or C:P ratios) that estimate potential to mineralize or

immobilize limiting nutrients, or the relative abundance of genes that encode for an

known physiology or physiological response (such as the relative abundance of cold

shock proteins to estimate cold tolerance)[17]. We refer to these *in situ* measurements

that allow estimation of microbial processes as microbial community properties

(hereafter community properties, Figure 2). Community properties represent an

integrated characteristic of the extant microbial community that has the potential to

estimate the microbial process of interest.

Microbial Community Properties

Microbial community properties can be separated into two categories, emergent

properties (EPs) such as biofilm thickness, which cannot be determined from the

properties of their constituent populations[18], and community aggregated traits

(CATs) such as nitrification potential, which can be estimated from community

membership or at least characteristics such as relative gene abundance (e.g.

AmoA), of those members[17].

The potential importance of EPs to influence ecosystem processes was

demonstrated in series of experiments conducted in flow-through flumes that

mimicked development and metabolism of stream biofilms[19]. Both transient storage

(i.e. an increase in residence time of the water and its solutes near the biofilm

relative to the flow around it) and the biofilm community's ability to use arabinose

relative to glucose increased as the microbial biofilm density increased and porosity

decreased. Microbial biofilm thickness and density are both EPs that affected the

important ecosystem processes of hydrological transient storage and substrate use

specificity[19]. In this case biofilm thickness was affected by physical factors (i.e. flow)

but biofilm thickness may also be influenced by other environmental characteristics

such as P availability[20].

While EPs are powerful metrics for understanding ecosystem processes (Figure 2,

Letter F or H) they cannot, by definition, be estimated from a list of constituent taxa

or characteristics of those taxa (Figure 2, Letter E) and thus must remain as an

intermediary between environmental drivers such as flow or P availability (Figure 2,

Letter C) and ecosystem processes.

Unlike EP's CATs can be estimated from characteristics of their constituents and

provide one pathway to link microbial community membership to the community

properties that drive ecosystem processes[17] (Figure 2, Letter E and F). Microbial

community biomass stoichiometry (e.g. biomass C:N or C:P) is one example of a

putative CAT that has been shown to be a useful predictor of nutrient immobilization

or mineralization during litter decomposition[21], and can predict both respiration and

N-mineralization better than microbial biomass alone[22]. In a study of soil microcosms

amended with organic carbon and reactive N, the relationship between the resource

C:N and microbial biomass C:N was able to better predict whether C would be

respired or immobilized relative to microbial biomass alone[22]. Biomass stoichiometry

has been shown to differ among phylogenetically different organisms. For example,

at the broadest level microbial biomass C:N differs between fungi and bacteria[24], and

has been shown to be variable among a wide range of taxa grown on the same

media[25,26]. Thus community biomass stoichiometry has the potential to be empirically

deconstructed into the biomass stoichiometry of its constituent taxa[27], linking

community membership and a community property (e.g. biomass C:N) with the

power to estimate an important microbial process (e.g. N-mineralization).

Microbial Community Membership

Although analysis of community membership by sequencing phylogenetic marker

genes provides organism identity, the direct coupling of microbial phylogeny to its

physiology and ecology is often weak[28] (Figure 2, Letter G). For example, most

organic carbon molecules can be metabolized by a phylogenetically diverse suite of

organisms and denitrification is also a phylogenetically broad process. The result is

that with the exception of some specialists (e.g. nitrifiers), we can infer very little

about the function of microbial communities from a list of their constituent taxa.

Whereas it is clear that microbial populations are non-randomly distributed in space

and time[29] and some microbial traits appear to be conserved at coarse taxonomic

scales[30,31,32] the underlying physiological mechanisms for phylogenetic sorting

across environmental gradients is often unknown. This prevents an explicit link

between the organism's relative abundance and their role in the collective

community physiology that influences ecosystems. The paucity of associated

physiological data that accompanies phylogenetic sequence data in most studies

limits the ecological insight from phylogenetic analyses and constrains our ability to

attribute microbial processes to community membership even of well-defined

consortia.

In addition to the paucity of ecological inference provided by an analysis of

community membership, community analysis using current methods has two

exceptional challenges that need to be addressed in order to gain insight from

community membership to drivers of ecosystem processes. First, bulk extraction of

DNA from environmental samples, often the first step in analysis of microbial

communities, may or may not represent the extant and active microbial community at

the time of sampling. In microbial ecology, unlike in plant or animal ecology, the

number, biomass, and identity of different populations cannot be assessed with

confidence[33,34]. When DNA is extracted from the environment the presence or

relative abundance of a given sequence is not necessarily proportionate to the

absolute abundance or biomass of that organism within the community. One reason

for this is that the extracted nucleic acid may not have been contained within a

microbial cell at the time of extraction. This idea has been well established[35] and the

proportion of extracellular DNA is known to vary among ecosystems[36] (e.g. there

may be more DNA in sediment than the water column). It is clear that a portion of

sequences derived from any environmental sample are not from intact cells[37,38,39].

Even when specific phylotypes can be empirically linked to intact cells (e.g. using *in*

situ hybridization), the viability and metabolism of that cell typically remains

unknown. Because of this, the presence of a sequence within a "community" does

not indicate the associated organism is participating in the microbial process of

interest, and if active, it does not indicate that that the organism's contribution is

proportionate to the relative abundance of its DNA sequence. While these facts are

readily acknowledged, they can be very challenging to address, and therefore are

often overlooked or ignored as a problematic but non-addressable constraint,

something of an inconvenient truth of microbial ecology.

The second major challenge in linking relative abundance of microbial populations

with microbial processes is that the diversity, growth rate, and metabolic complexity

of environmental microorganisms are orders of magnitude greater than all other

organisms. The result is that when scaling from individuals through populations to

microbial communities, the obfuscating factors described in Figure 1b introduce and

ultimately accumulate more uncertainty at higher levels of ecological organization,

relative to the same analyses applied to macroorganism-ecosystem linkages. This

uncertainty is further confounded because community composition and processes

are routinely measured at different spatial scales[40]. Translating microbial

measurements to an ecosystem flux typically requires linking measurements from

microbial physiology (10^{-12} m), to microbial process measurements (10^{-1} or 10^{-2} m),

to ecosystem process measurements (m to 10^3 m). This enormous scale (15 orders

of magnitude) over which to interpolate data, raises challenges analogous to

mechanistically analyzing the global carbon cycle, i.e. linking experiments on grams

of carbon in soils (e.g. g C per g dry weight) to global fluxes of Petagrams (10^{15} g).

However, unlike global C cycle research, microbial-to-ecosystem research often

addresses these in a descriptive manner using correlative approaches, without the

rigor and quantitative modeling approaches typically applied to global

biogeochemical cycles[41]. Each of these challenges must be effectively addressed in

order to rigorously incorporate the growing wealth of information on microbiomes to

system level processes.

A Way Forward at the Intersection of Microbial and Ecosystem Science

The conceptual diagram presented here (Figure 2) provides a road map for

organizing and linking the diverse suite of microbial characteristics that are

commonly measured. Ecosystem ecology has traditionally been confined to the

interactions depicted within the horizontal arrow, moving from environmental

parameters to ecosystem processes (Letter A, Figure 2). As the role of the

microbiome has come to the forefront of environmental sciences it is clear that

microbial ecology has a great deal to contribute. However, microbial ecology has

traditionally been confined to interactions depicted within the vertical arrow, moving

from individuals (microbial community membership) to process rates (microbial

processes) of populations or more recently communities (Figure 2). The excitement

to integrate microbial metrics into ecosystem science has led to a range of novel

approaches of linking characteristics on the microbiome to ecosystem processes.

Direct connections between microbial community membership and ecosystem

processes (Figure 2, Letter J) or community properties and ecosystem processes

(Figure 2, Letter I) are almost exclusively correlative in nature. Whereas many of

these results are intriguing, the relationships discovered across these pathways, in

the absence of the defining physiological mechanism, are often restricted to the time

and place they are identified. Because of this, moving from correlative,

phenomenological approaches to a causative and mechanistic understanding is a

challenging but necessary step for microbiome science.

Because microbial processes can be estimated by community properties (e.g. N

mineralization and biomass C:N), understanding the drivers of community properties

is a way to more explicitly link environmental microorganisms with the ecosystem

processes they control. We propose that identifying which community properties best

describe microbial processes (Figure 2, Letter F), then identifying whether or not the

community properties that best describe each process are a CAT (Letter E, Figure 2)

or an EP (Figure 2 Letter C) provides clear pathway to understand whether

environmental drivers or microbial drivers dominate ecosystem processes. Currently,

many microbial community properties are EPs (i.e. cannot be predicted from their

constituent members or their characteristics). Understanding when community

properties can be predicted by membership is an important and open research

direction. Distinguishing which community properties represent aggregated traits

(CATs) and which are actually EPs may be an essential link in advancing our ability

to apply microbial information to ecosystem science. This is not a trivial task,

however a suite of existing methods already provides the ability to directly pursue

this challenge.

Applying the proposed framework

Understanding the principal drivers of community properties will require a series of

complementary approaches applied in concert, including; stable isotope probing of

cultured isolates and mixed communities, single cell methods that can assay cells in

the same physiological state they occur in in the environment, sorting of complex

communities into subsets of populations or consortia for further investigation, and

physiological assays of isolates grown in culture. Studies that use either labeled

substrates or single cell techniques (or both) have been successful in linking

community composition and process rates. One example, stable isotope probing

(SIP, in which an isotopically labeled element (or elements) from a defined substrate

can be tracked into microbial biomass) is a method used to identify which organisms

may be participating in an ecosystem process of interest[42]. For example, a study of

sulfate reduction in a Scottish peatland revealed that a single species of

Desulfosporosinus was most likely responsible for the totality of sulfate reduction

within the peatland even though it only comprised 0.0006% of the retrieved

sequences[43]. In this case the *Desulfosporosinus* species represented the only

known sulfate reducer within the community and thus the kinetics of this organism

seemingly defined the kinetics of sulfate reduction in the entire system. Whereas this

is a single example of using confirmative ecophysiology to link categories of

microbial information (pathway G in Figure 2), there is a suite of culture-free

techniques (such as Raman microspectroscopy (MS), NanoSIMS, or X-ray

microanalysis, XRMA) that complement sequence-based microbiome analysis by

reporting on the physiological and compositional characteristics of individual cells *in*

situ[27, 44,45]. All three methods (Raman MS, NanoSIMS and XRMA) can be coupled

with phylogentic labels (*in situ* hybridization) that can visualize identification of

phylotypes simultaneously with macromolecular (Raman MS), isotopic (NanoSIMS),

and elemental composition (XRMA). In addition, both Raman MS and NanoSIMS

can trace substrates that are labeled with stable isotopes into cells, providing the

ability to identify populations that are participating in specific metabolic pathways

within complex communities. These powerful approaches applied in concert with

sequence analysis have the potential to empirically link the categories of microbial

information defined here (Figure 2). These methods also have the greatest potential

to begin to unravel, which community properties are EPs, and which are CATs.

In addition to direct visualization of individuals cells from mixed populations or

consortia using single cell approaches, there are abundant examples of

immunocapture (e.g. bromodeosyuridine, BrdU)[46] or other labeling and cell sorting

approaches (eg. fluorescent in situ hybridization(FISH) coupled with flow cytometry

cell sorting, FACS)[47] that provide powerful tools to constrain the complexity of

microbial communities and link community membership to microbial characteristics

that influence ecosystem processes. Labeling and sorting techniques allow the cells

that incorporate a labeled substrate or can be targeted with a stain or fluorescent

reporter to be separated from the broader community and then assayed for

membership or for biomass composition. For example, a study of a North Atlantic

bacterial community labeled the actively growing component of the community using

BrdU and then separated those populations from the rest of the community using an

immune capture technique[46]. Similarly, cells can be labeled with phylogenetic probes

(e.g. FISH) that fluorescent at different wavelengths and can be separated from the

general community using a flow cytometry to only select those cells that reported for

the phylogenetic label[47].

In addition to these approaches for assaying natural communities, there is a need

for broader representation of cultured taxa that better represent physiologies that are

similar to those phyla found in the environment. For example, microbial community

carbon use efficiency (the amount of carbon allocated to biomass production relative

to carbon respired or stored) is a central parameter in many ecosystem level carbon

cycling models. Differences in CUE among phyla depend largely on the relative

plasticity of growth rate, carbon storage capacity, and maintenance respiration. *E.*

coli, the "poster bacterium" for physiological assays is unique both in the plasticity of

its growth rate and its capacity to store energy as organic carbon and thus not

necessarily informative to develop a better understanding of CUE[48]. *Streptococcus*

Bovis has proven to as a more appropriate organism to study energy cycling in

bacteria[48]. Physiological studies of isolates from a broader distribution of

representative phyla are key to advancing our understanding of environmental

microbiomes. However, it is unlikely that information about specific phenotypes

estimated in isolation from pure-culture studies can be directly used to estimate

community properties because of the plasticity of organismal physiology and

because of the prevalence of competitive interactions when isolates are grown in co-

culture with even one other organism[49]. Therefore studies of isolates grown in

culture would provide more powerful information if they reported the plasticity of a

given phenotype, rather than only the phenotype under a single set of environmental

conditions. For example, a recent study of 24 freshwater bacterial isolates showed

differences in phenotypic plasticity in the biomass stoichiometry of the taxa studied[26].

Some of the taxa had highly variable biomass C:P ratios whereas others

demonstrated a high degree of homeostasis in their C:P ratios. This allowed the

authors to hypothesize how communities or consortia with homeostatic phenotypes

would respond to environmental drivers compared to communities or consortia

composed of populations with more plastic phenotypes.

Whereas linking microbial membership to system level processes is an exceptional

challenge the tools and approaches to address this challenge already exist. These

existing and established microbiological methods both relatively novel (NanoSIMS)

and foundational in the field (physiological assays of isolates) can be applied in

concert to begin to parse the exceptional complexity of environmental microbiomes.

Designing microbiome research to maximize insights into system-level

processes

The meta-analyses discussed above clearly illustrate that a more directed

approach to microbiome research is necessary. We suggest that rather than looking

for linkages among microbial community membership and system-level processes in

every study, research efforts would benefit from strategically targeting the linkages

and processes for which an *a priori* understanding of microbial physiology should

allow us to improve our understanding of the ecosystem process. These cases may

be identified first by noting patterns in which environmental factors explain little of the

variability in an ecosystem process, or where system-level responses deviate

significantly from rates that are predicted from environmental factors alone. These

deviations could include spatial or temporal heterogeneity in an ecosystem response

where the environmental characteristics do not have the same level of heterogeneity.

For example, the discovery of novel microbial metabolic pathways (i.e. annamox)

has helped explain otherwise puzzling chemical transformations, such as the

oxidation of ammonium under anoxic conditions[50]. Such advances are most likely to

be cases where the microbial process of interest can be directly linked to a

phylogenetically constrained group (i.e. a narrow process) and where the system-

level behavior of the process reflects the organismal-level physiological controls,

such as N-sensitivity of methane monooxygenase[51].

The framework presented here provides one approach to formalize inquiry across

microbiome science and encourage empirical linkages between the presence of

organisms in a system and the processes that characterize that system. Whereas we

draw examples from environmental microbiomes and the ecosystems they inhabit,

the framework presented here should also benefit the analysis of microbiomes

associated with other systems such as host organisms and those of engineered

environments. We assert that this framework provides an important and

straightforward starting point as the global research community aims to undergo one

of the most exciting concerted efforts in the microbial sciences to date.

Acknowledgements

This work is a product of the Next Generation of Ecosystem Indicators Working

Group, supported by the USGS John Wesley Powell Center for Synthesis and

Analysis. Preparation of this manuscript was supported by NSF DEB IOS #1456959

awarded to EKH. Chuck Pepe-Ranney provided valuable feedback on a previous

version of this manuscript.

Figures

Figure 1 Diagram of microbial-ecosystem linkages A) how linkages are commonly conceptualized across levels of ecological organization and B) the series of environmental filters that create challenges when attempting to link metrics from one level of ecological organization to the other.

Figure 2 Reframing how we study microbial-ecosystem linkages. Shown is the intersection between microbial (vertical) and ecosystem (horizontal) ecology with each of the three categories of microbial information (microbial processes, emergent community properties, and microbial community membership) as defined in the text. We argue for an increased focus on studies that elucidate pathways E, F and H. In addition we note that pathways G, J and I are less likely to effectively incorporate microbial information into ecosystem science. The delta symbol in each category indicates an emphasis on how changes within a category may lead to a change in a connected category.

References

1. Alivisatos A. P., M. J. Blaser, E. L. Brodie, M. Chun, J. L. Dangl, T. J. Donohue, P. C. Dorrestein, J A Gilbert, J. L. Green, J. K. Jansson, R. Knight, M. E. Maxon, M. J. McFall-Ngai, J. F. Miller, K. S. Pollard, E. G. Ruby, S. A. Taha, Unified Microbiome Initiative Consortium 2015: A unified initiative to harness Earth's microbiomes. Science 350, 503-504. DOI: 0.1126/science.aac84809. Angilletta, M. J. Jr. (2009). Thermal Adaptation. A Theoretical and Empirical Synthesis. Oxford: Oxford University Press, New York, New York

2. Dubilier N., McFall-Mgai, M. and Zhao L. (2015): Create a global microbiome effort. Nature 529 (631-634).DOI: 10.1038/526631a

3. Blaser, M.J., Cardon, Z. G., Cho, M.K., Dangl, J.L, Donohue, T.J.,Green, J.L., Knight, R., Maxon, M.E., Northen, T.R., Pollard, K.S. and Brodie, E.L. 2016 Toward a Predictive Understanding of Earth's Microbiomes to Address 21st Century Challenges 7(3) e00714-16

4. Graham E. B., Wieder, W.R., Leff, J.R., Weintraub, S. W., Townsend, A.R., Cleveland, C.C., Philippot L., Nemergut D.R. 2014 Do we need to understand microbial communities to predict ecosystem function? A comparison of statistical models of nitrogen cycling processes Soil Biology and Biochemistry 68, 279–282

5. Graham E. B. et al. 2016 Microbes as engines of ecosystem function: when does community structure enhance predictions of ecosystem processes? Frontiers in Microbiology 7: doi: DOI=10.3389/fmicb.2016.0021

6. Rocca J.D., Hall E.K., Lennon J.T., Evans S.E., Waldrop M.P., Cotner J.B., Nemergut D.R., Graham E.B., Wallenstein M.D. (2015). ISME J. 9, 1693–1699

7. Bier R.L., E.S. Bernhardt, CM Boot, EB Graham, EK Hall, JT Lennon, D Nemergut, B Osborne, C Ruiz-González, JP Schimel, MP Waldrop, MD. Wallenstein How are we forging conceptual, analytical, and mechanistic links between microbial community structure and ecosystem process? (in review at FEMS Microbiol.)

8. Felip M., Pace M.L., Cole, J.J. (1996). Regulation of planktonic bacterial growth rates: The effects of temperature and resources. Microb Ecol. 31(1):15-28.

9. Hochachka P.W., Somero G.N. (2002). Biochemical adaptation: Mechanism and process in physiological evolution. Oxford University Press, New York

10. Angilletta, M. J. Jr. (2009). Thermal Adaptation. A Theoretical and Empirical Synthesis. Oxford: Oxford University Press, New York, New York

11. Schimel, J.P., J. Bennett, and N. Fierer. 2005. Microbial community composition and soil N cycling: is there really a connection? In: Biological diversity and function in soils. Bardgett, R.D., D.W. Hopkins, and M.B. Usher (Eds.) Cambridge University Press. pp. 171-188

12. Schimel, J.P. (1995). Ecosystem consequences of microbial diversity and community structure in Arctic and Alpine Biodiversity: Patterns, Causes and Ecosystem Consequences Ecological Studies Volume 113, pp 239-254

13. Lenhart K., M. Bunge, S. Ratering, T. R. Neu, I. Schüttmann, M. Greule, C. Kammann, S. Schnell, C. Müller, H. Zorn & F. Keppler. (2012). Evidence for methane production by saprotrophic fungi Nature Comm. 3:1046

14. Wallenstein, M. D., and E. K. Hall. (2012). A trait-based framework for predicting when and where microbial adaptation to climate change will affect ecosystem functioning. Biogeochem.: 109: 1-3, 35-47

15. German D.P., Marcelo, K.R.B., Stone, M.M and Allison, S.D. (2012). The michaelis-menten kinetics of soil extracellular enzymes in response to temperature: a cross-latitudinal study Global Change Biology 18,1468-1479

16. Hall, E. K., A. R. Dzialowski, S. M. Stoxen, and J. B. Cotner. (2009). The effect of temperature on the coupling between phosphorus and growth in lacustrine bacterioplankton communities. Limnol. and Oceanogr. 54:880-889.

17. Fierer, N., A. Barberán, D. Laughlin. (2014). Seeing the forest for the genes: Using metagenomics to infer the aggregated traits of microbial communities. Frontiers in Microbiology. 5:614

18. Konopka, A. (2009) What is microbial community ecology ISME J., 3:11, 1223-12230

19. Battin, T., L. A. Kaplan, L. Newbold, X. Cheng, and C. Hansen. (2003). Effects of current velocity on the nascent architecture of stream microbial biofilms. Appl. and Env. Microbiol. 69:5443-5452.

20. Pepe-Ranney, C.P. and Hall, E.K. 2015 Carbon subsidies affect planktonic niche partitioning and recruitment of bacteria to marine biofilms http://dx.doi.org/10.3389/fmicb.2015.00703 Frontiers in Aquatic Microbiology

21. Manzoni, S., R. B. Jackson, J. A. Trofymow, and A. Porporato. (2008). The global stoichiometry of litter nitrogen mineralization. Science 321:684-686.

22. Buchkowski, R.W., Schmitz, O.J., Bradford, M.A. (2015) Microbial stoichiometry overrides biomass as a regulator of soil carbon and nitrogen cycling. Ecology, 96, 1139-1149.

23. Strickland M.S., and Rousk J. 2010 Considering fungal:bacterial dominance in soils- methods, controls and ecosystem implications Soil Biology and Biochemistry 42: 1385-1396

24. Keiblinger, K.M.† Hall, E.K†, Szukics, U., Hämmerle, I., Ellersdorfer, G., Sterflinger, K., Wanek, W., Richter, A., Jandl R., and Zechmeister -Boltenstern, S. 2010 The effect of resource quantity and resource stoichiometry on microbial carbon use efficiency FEMS Microbiology Ecology 73(3): 430-440

25. Mouginot, C., Kawamura, R., Matulich, K.L., Berlemont, R., Allison, S.D., Amend, A.S. and Martiny, A.C. (2014) Elemental stoichiometry of fungi and bacteria strains from grassland leaf litter Soil Biol. & Biochem. 76, 278-285

26. Godwin, C.M. and J.B. Cotner. 2015 Stoichiometric Flexibility in Diverse Aquatic Heterotrophic Bacteria Is Coupled to Differences in Cellular Phosphorus Quotas. Name: Frontiers in Microbiology 6 (2015): doi:10.3389/fmicb

27. Hall, E.K., Maixner F., Franklin O., Daims, H., Richter, A. and Battin, T. (2011) Linking microbial and ecosystem ecology using ecological stoichiometry: A synthesis of conceptual and empirical approaches Ecosystems 14, 261-273

28. Newton, R. J, S. E. Jones, A. Eiler, K. D McMahon, and S. Bertilsson. (2011). A guide to the natural history of freshwater lake bacteria. Microbiol. and Molec. Biol. Rev. 75:1, doi:10.1128/MMBR.00028

29. Martiny, J.B.H., B. Bohannan, J. Brown, R. Colwell, J. Fuhrman, J. Green, M.C. Horner-Devine, M. Kane, J. Krumins, C. Kuske, P. Morin, S. Naeem, L. Ovreas, A.-L. Reysenbach, V. Smith, J. Staley. 2006. "Microbial biogeography: Putting microorganisms on the map." Nature Reviews Microbiology 4: 102-112

30. Martiny JBH, Jones SE, Lennon JT, Martiny AC (2015) Microbiomes in light of traits: a phylogenetic perspective. Science 350: aac9323 (pdf)

31. Lennon JT, Aanderud ZA, Lehmkuhl BK, Schoolmaster DR (2012) Mapping the niche space of soil microorganisms using taxonomy and traits. Ecology 93: 1867-1879 (pdf)

32. Treseder KK, Lennon JT (2015) Fungal traits that drive ecosystem dynamics. Microbiology and Molecular Biology Reviews 79: 243-262 (pdf)

33. Tilman D, PB Reich, J Knops, D Wedin, T Mielke, C Lehman. (2001). Diversity and productivity in a long-term grassland experiment Science, 294:5543, 843-845

34. Reich P. B., David Tilman, Forest Isbell, Kevin Mueller, Sarah E. Hobbie, Dan F. B. Flynn, and Nico Eisenhauer. (2012). Impacts of biodiversity loss escalate through time as redundancy rades Science: 336:6081, 589-592.

35. Klein, D. A., 2011 Bulk Extraction-Based Microbial Ecology: Three Critical Questions Microbe Sept

36. Corinaldesi C., R. Danovaro, and A. Dell'Anno (2005) Simultaneous Recovery of Extracellular and Intracellular DNA Suitable for Molecular Studies from Marine Sediments Appl. and Environ. Microbiol., 71:1, 46–50

37. Dell'Anno A. and Roberto Danovaro (2005) Extracellular DNA Plays a Key Role in Deep-Sea Ecosystem Functioning Science 309 (5744), 2179

38. Torti, A., Leverb, M.A., Jørgensen, B.B. 2015 Origin, dynamics, and implications of extracellular DNA pools in marine sediments Marine Genomics doi:10.1016/j.margen.2015.08.007

39. Carini P., Patrick J. Marsden, Jonathan W. Leff, Emily E. Morgan , Michael S.Strickland, Noah Fierer 2016 Relic DNA is abundant in soil and obscures estimates of soil microbial diversity http://dx.doi.org/10.1101/043372

40. Clark, J. S., Bell, D. M., Hersh, M. H., Kwit, M. C., Moran, E., Salk, C., Stine, A., Valle, D. and Zhu, K. (2011) Individual-scale variation, species-scale differences: inference needed to understand diversity. Ecol. Lett., 14, 1273–1287

41. Thornton P.E., Doney S.C., Linsay K., Moore J.K., Mahowald N., Randerson J.T., Fung I., Lamarque J.F., Feddema J.J., Lee, Y.H. (2009) Carbon-nitrogen interactions regulate climate-carbon cycle feedbacks: results from an atmosphere-ocean general circulation model Biogeosciences, 6, 2099-2120

42. Neufeld J.D., Vohra J., Dumont M.G., Lueders T., Manefield M., Friedrich M.W., Murrell J.C. 2007 DNA stable-isotope probing. Nat Protoc. 2:4, 860-866

43. Pester, M., Bittner, N., Deevong, P., Wagner, M., and Loy, A. (2010). A 'rare biosphere' microorganism drives sulfate reduction in a peatland. ISME J. 4:1591–1602.

44. Wagner, M. (2009). Single cell ecophysiology of microbes as revealed by raman microspectroscopy or secondary ion mass spectrometry imaging. Annu Rev Microbiol 63:411-429.

45. Behrens, S., Kappler, A., Obst M. (2012). Linking environmental processes to the in situ functioning of microorganisms by high-resolution secondary ion mass spectrometry (NanoSIMS) and scanning transmission X-ray microscopy (STXM) Env. Microbiol. 14:11, 2851-69

46. Galand P.E., L. Alonso-Sáez, S. Bertilsson, C. Lovejoy, and E. O. Casamayor (2013) Contrasting activity patterns determined by BrdU incorporation in bacterial ribotypes from the Arctic Ocean in winter Front Microbiol. 2013; 4: 118

47. Czechowska, K., Johnson, D.R., van der Meer, J.R. (2008) Use of flow cytometric methods for single-cell analysis in environmental microbiology Curr. Opin. In Microbiol. 11:3, 205-212

48. Russell, J.B. (2007). The energy spilling reactions of bacteria and other organisms. Journal of Molecular Microbiology and Biotechnology 13, 1-11

49. Foster, K.R. and Bell, T. 2012 Competition, not cooperation, dominates interations among culturable microbial species Curr. Biol. 22, 1845-1850

50. Mulder A., A. A. Van De Graaf, L. A. Robertson, and J. G. Kuenen. (1995). Anaerobic ammonium oxidation discovered in a denitrifying fluidized bed reactor, FEMS Microbiology Ecology, 16:3, 177–184,
51. Paul LE Bodelier, Interactions between nitrogenous fertilizers and methane cycling in wetland and upland soils, Current Opinion in Environmental Sustainability, Volume 3, Issue 5, October 2011, Pages 379-388, ISSN 1877-3435, http://dx.doi.org/10.1016/j.cosust.2011.06.002.