Understanding How Microbiomes Influence The Systems They Inhabit: moving from a correlative to a causal research framework

Authors: Hall, E.K.[1*], Bernhardt, E.S[2], Bier, R.L.[2], Bradford, M.A.[3], Boot, C.M.[1], Cotner, J.B.[4], del Giorgio, P.A.[5], Evans, S.E.[6], Graham E. B [2,7,8], Jones, S.E.[9], Lennon, J.T.[10], Locey, K.J.[10] Nemergut, D.[2], Osborne, B.B.[11], Rocca, J.D.[1,2], Schimel J.S.[12], Waldrop, M.P.[13], Wallenstein, M.W.[1]

Article Type: Perspectives
Words in Abstract: 271
Words in Main Text: 5,167
2 Figures, 0 Tables, 53 References

Running Title: How microbiomes influence ecosystems
Keywords: microbiome, microbial ecology, ecosystem ecology, emergent properties, community aggregated traits

Affiliations: [1]Colorado State University, Fort Collins CO (ed.hall@colostate.edu, claudia.boot@colostate.edu, matt.wallenstein@colostate.edu) [2]Duke University, Durham, N.C. (ebernhar@duke.edu, ebgraham2@colorado.edu, jennifer.rocca@duke.edu) [3]Yale University, New Haven, CT, Dept. of Forestry and Environmental Studies (mark.bradford@yale.edu), [4]University of Minnesota, Saint Paul, MN (cotne002@umn.edu) [5]Université du Québec à Montréal, Montréal, CA (del_giorgio.paul@uqam.ca) [6]Michigan State University, Hickory Corners, MI (evanssar@gmail.com) [7]Institute of Arctic and Alpine Research, University of Colorado at Boulder [8]Pacific Northwest National Laboratory, Richland, WA, USA [9]Notre Dame University, South Bend, IN (sjones20@nd.edu) [10]Indiana University, Bloomington, IN (lennonj@indiana.edu, ken@weecology.org) [11]Brown University, Providence, RI (brooke_osborne@brown.edu) [12]University of California, Santa Barbara, Santa Barbara, CA (josh.schimel@lifesci.ucsb.edu) [13]U.S. Geological Survey, Menlo Park, CA (mwaldrop@usgs.gov) Biological Sciences Division,

* **Corresponding Author**: Dr. Ed Hall, Department of Ecosystem Science and Sustainability, Natural Resource Ecology Laboratory Campus Delivery 1499, Colorado State University, Fort Collins, CO 80523, ed.hall@colostate.edu, 970-491-2162

Contribution: All listed authors have contributed to the conceptualization, writing, and preparation of the current manuscript.

Abstract

Translating the ever-increasing wealth of information on microbiomes (environment,

host, or built environment) to advance the understanding of system-level processes

is proving to be an exceptional research challenge. One reason for this challenge is

that relationships between characteristics of microbiomes and the system-level

processes they influence are often evaluated in the absence of a robust conceptual

framework and reported without elucidating the underlying causal mechanisms. The

reliance on correlative approaches limits the potential to expand the inference of a

single relationship to additional systems and advance the field. In this perspective

piece we propose that research focused on how microbiomes influence the systems

they inhabit should work within a common framework and target known microbial

processes that contribute to the system-level process of interest. Here we identify

three distinct categories of microbiome characteristics (microbial processes,

microbial community properties, and microbial membership) and propose a

framework to empirically link each of these categories to each other and the broader

system level processes they affect. We posit that it is particularly important to

distinguish microbial community properties that can be predicted from constituent

taxa (community aggregated traits, CATs) from those properties that are currently

unable to be predicted from constituent taxa (emergent properties, EPs). We discuss

how a series of existing methods in microbial ecology can be applied to more

explicitly elucidate properties within each of these categories and connect these

three categories of microbial characteristics with each other. We view this proposed

framework, gleaned from a breadth of research on environmental microbiomes and

ecosystem processes, as a promising pathway with the potential to advance

microbiome science across a broad range of disciplines.

Current Approaches Linking Microbial Characteristics and Ecosystem Processes

In all ecosystems, virtually all processes are influenced by microorganisms and many processes are carried out primarily/exclusively by microorganisms. This has sometimes led to the assumption that a better description of the microbiome (including its associated transcripts, proteins, and metabolic products) should lead to a better understanding and predictions of system level processes. However, such justifications assume that measureable characteristics of the microbiome (e.g. 16S rRNA gene libraries, metagenomes, enzymatic activities) can inform our ability to better understand and predict system-level processes. Unfortunately, additional information about the microbiome does not always provide a clearer understanding of ecosystem processes beyond what can be predicted by environmental factors alone[1,2].

Two recent meta-analyses[3,4] suggest that research at the intersection of ecosystem science and microbial ecology often rely on assumed correlations between microbiome characteristics and ecosystem processes and less frequently test to see if those correlations are present. The first, an examination of 415 studies, found little evidence that protein-encoding genes (sometimes referred to as "functional genes") or gene transcripts correlate with associated biogeochemical processes[3]. Although all studies attempted (or presumed) to link microbial genes or transcripts with function, only 14% measured both the copy number of genes or transcripts and the corresponding process (n = 59 studies, comprising 224 individual effects as most studies had measured multiple gene-function relationships). Of the 224 effects where both characteristics were measured, only 38% exhibited a positive relationship between molecular and process measurements that many assume to

exist. The effect size for the relationship between gene copy number and process

rate had an approximately normal distribution with a mean near zero[3]. This result

was consistent whether functional gene or transcript abundance was used as the

response variable. The second study, compiled a separate dataset of 148 studies

that examined microbial membership and ecosystem processes in response to

experimental manipulations[4]. Whereas 40% of included studies reported

concomitant changes in microbial membership and an ecosystem process, only one

third of those cases reported the relationship between microbial membership and an

ecosystem process. Interestingly, of the 53 studies that posed a hypothesis or

objective about links between microbial membership and ecosystem processes,

more than half (53%) did not test for a statistical structure-function link[4].

Microbiomes are the engines that power system-level processes[5]. However the

meta-analyses described above illustrate that the current approach to study the links

between microbiome characteristics and ecosystem processes are not well

formulated and relationships between microbiome characteristics and system level

processes are rarely tested. When linkages are explicitly tested, significant

correlations between microbiome characteristics and ecosystem processes are

sometimes present, but more frequently not present [3,4]. One reason for the ambiguity

between microbiome characteristics and system level processes is that many studies

are conducted in the absence of a conceptual framework that illustrates how different

measurable microbial characteristics relate to one another and to the system level

process of interest.

Microbial characteristics can range in resolution from cellular abundance to the

entire genetic potential of a species-rich community (i.e., a metagenome). Because

all measureable microbial characteristics do not exist within the same plane of detail,

a conceptual structure is required to clarify how various microbial characteristics are

related to each other and to the processes they affect. Identifying correlations

between a microbial characteristic and an ecosystem process in the absence a

broader conceptual underpinning creates the potential for correlations to be

mediated by a third (or more) unaccounted variable(s). The absence of an underlying

mechanism also limits the potential for each result to be applied to additional

systems or to expand to broader spatial and temporal scales, for additional testing,

replication, and confirmation.

Challenges in Linking Microbial Characteristics and Ecosystem Processes

A key challenge in linking microbial information to a system-level process is that

conceptual research frameworks often do not effectively align with the methods

being applied or the data those methods generate. For example, environmental

factors act on the physiology of individual organisms, which alters their competitive

ability, abundance, collective physiology, and ultimately their contribution to an

ecosystem process. However, designing an observational study or experiment from

this framework assumes that environmental and microbial characteristics are

measurable across multiple categories of ecological organization (i.e., individuals,

populations, and communities) at the temporal and spatial scales at which they

influence microorganisms (Figure 1a). In addition, the relationships between

environmental variables and microbial characteristics can be decoupled in both time

and space[4], and are often non-linear[6]. Recent immigration, phenotypic plasticity,

disequilibrium between the environment and the extant microbiome at the time of

sampling, functional redundancy, and dormancy can all mask the relationship

between measurable microbial characteristics and the processes microorganisms

influence (Figure 1b). [4,7,8,9] As micrometer scale characteristics of microbiomes (10^{-6}

m) are scaled to the level of ecosystems (m to km), we assume that our conceptual

understanding is also scalable. However, each of the aforementioned confounding

factors aggregate over multiple orders of magnitude often masking the very

relationships we seek to elucidate (Figure 1b). To formalize how measureable

microbiome characteristics are linked with system-level processes we have

conceptually defined the intersection of microbial and ecosystem ecology and

identified three categories of microbial characteristics that illustrate the relationship

among each category of microbial characteristics and how they may contribute to an

ecosystem process (Figure 2).

Mapping Ecosystem Processes to Microbial Characteristics

Ecosystem processes are defined as a qualitative change in a pool or a flux from

one pool to another (e.g. NH_4^+ to NO_3^-, or dissolved organic matter to CO_2). The first

step to understand how the microbiome influences an ecosystem process, is to

define the ecosystem process of interest and each sub-process that contributes to it

(i.e., the set of constituent reactions that combine to determine net flux). Few, if any

ecosystem processes involve a single metabolic pathway, or are carried out by a

single organism (e.g., a notable exception being the recent discovery of commamox

which can independently carry out nitrification, the conversion of NH_4^+ to NO_3^-)[10,11].

Rather, ecosystem processes are composites of complementary or antagonistic sub-

processes, carried out by phylogenetically and metabolically diverse

microorganisms[12]. For example, net ecosystem productivity (NEP) is the balance

between antagonistic processes of C-fixation and C-mineralization. Each sub-

process of NEP can be further partitioned into a series of metabolic pathways (e.g.,

chemoautotrophic nitrification and photoautotrophic C-fixation or heterotrophic

fermentation and aerobic respiration). Partitioning each ecosystem process in this

hierarchical manner can continue until the sub-processes maps directly to specific

microbial metabolic pathways (e.g., acetoclastic methanogenesis). Subsequently

each of these metabolic pathways can be categorized as either phylogenetically

broad or narrow[13]. Broad processes are phylogenetically common (i.e., widely

distributed among taxa), whereas narrow processes are phylogenetically conserved

(i.e., limited to a specific subset of taxa). For example, denitrification and

photosynthesis are phylogenetically broad processes, while both methanogenesis

and methanotrophy are phylogenetically narrow processes (with at least one notable

exception[14]).

The second step is to identify, the controls or constraints on each constituent sub-

process. For example, the kinetics of a single metabolic pathway in a model

organism may help us understand the rate limiting steps of a narrow process, but

insights from model organisms are much less likely to capture the full spectrum of

responses of a broad process where phenotypic variation among phylogenetically

diverse organisms is likely to be much greater[15,16]. Defining the ecosystem process,

its critical sub-processes, and the known phylogenetic distribution of the metabolic

pathways that drive those sub-processes creates an explicit conceptual pathway that

links the ecosystem process to the microorganisms that contribute to it. Once the

ecosystem process has been conceptually partitioned into its component parts and

their primary controls, a concerted approach can be applied to investigate how

characteristics of the microbiome influence the ecosystem process of interest within

the complexity of a natural environment.

Categories of Microbial Characteristics

We propose that attempts to elucidate the microbial contribution to system-level

processes needs to explicitly identify three distinct categories of microbial

characteristics: 1) microbial processes, 2) microbial community properties, and 3)

microbial membership (Figure 2). The contribution of the microbiome to ecosystem

processes is exerted through aggregate community properties that are shaped by

both microbial membership and environmental factors. This proposed framework

allows the researcher to clearly identify how different measurements used to

characterize a microbiome interact with each other and identify the potential of each

characteristic to elucidate the microbial contribution to the system level process.

These categories are hierarchically connected (Fig. 2), but they represent distinct

degrees of aggregation that are not simply an additive function of the previous

category. Furthermore, each category is potentially subject to different modes of

regulation, and, each category has different putative linkages to system-level

processes (Fig. 2). All measureable characteristics of microbial communities (e.g.,

abundance of cells, sequence of genes, transcripts, or proteins; enzyme expression

or activity) can be placed within one of the above categories, but most studies rarely

articulate how these measurements differ in their specificity (i.e., the level of

phylogenetic resolution at which they are applied), precision (i.e., the ability of the

method to repeatedly describe the characteristic of interest), or context (i.e., how a

characteristic relates to other characteristics or the ecosystem within which they

were measured). This conceptual structure that orientates each microbial category

within a broader context creates the opportunity to improve the design of

observational and experimental studies in micobiome research.

Microbial Processes - Microbial processes are the collective metabolism of the

microbiome that contribute to changes in pools and fluxes of elements or compounds

(i.e., Figure 2, Letter K). This is the level of microbial information that can most

readily be incorporated into system-level models because many microbial processes

represent the key sub-processes that contribute to a particular ecosystem process

(e.g., methanogenesis + methanotrophy ≈ methane efflux). Commonly measured

microbial processes in ecosystem science include nitrogen fixation, denitrification,

nitrification, phosphorus uptake and immobilization, carbon fixation, and organic

carbon mineralization. The rates of many microbial processes can be approximated

through physiological assays (e.g., biological oxygen demand to estimate microbial

community respiration), and while they do not open the "black box" of the microbial

community, they do directly quantify the microbial contribution (or at least the

potential contribution) to changes in resources moving through the box. Microbial

processes can be distinguished from other microbial characteristics because they

are all rates (i.e., have time in the denominator) and require a bioassay to estimate.

Assays used to estimate microbial processes are often logistically challenging,

require manipulations that inevitably deviate from the *in situ* conditions, and often

depend on the environment from which the microbiome was sampled. For example,

the relationship between temperature and microbial processes such as enzyme

activity and phosphorus use efficiency (PUE) vary across latitudinal gradients[17] and

among seasons[18]. Thus, observations of the effect of temperature on either enzyme

activity or PUE depend on where (e.g., at what latitude) and when (e.g., during which

season) they were measured. In the absence of an understanding of the underlying

physiological mechanism (e.g., the physiological change that allows a community to

perform differently at different temperatures), the relationship between and

environmental driver (e.g., temperature) and a microbial process must be measured

through a direct assay at each location and at each time. This limits the inference

possible from relying only on measurements of microbial processes alone.

Microbial Community Properties - Microbial community properties include a broad

set of microbial characteristics such as community biomass or biomass elemental

ratios (e.g., biomass C:N or C:P ratios) and the majority of phylogenetically

undifferentiated aggregate sequence based measurements (e.g., gene abundance,

metagenomes, transcriptomes). Microbial community properties (Figure 2) represent

an integrated characteristic of the microbiome that has the potential to predict or at

least constrain the estimates of microbial processes. For example, microbial

community biomass C:N (a community property) indicates a microbiome's potential

to mineralize or immobilize N.[19] Community biomass stoichiometry has been shown

to be a useful predictor of nutrient immobilization or mineralization during litter

decomposition[19], and in soils can predict both respiration and N-mineralization better

than microbial biomass alone[20]. The power of biomass elemental ratios to explain

nutrient cycling has also been shown in freshwater[21] and marine ecosystems[22]

including the seminal paper that demonstrated the similarities between the

stoichiometry of marine algal biomass and that of the dissolved fraction of nutrients

in the ocean[23].

Microbial community properties can be separated into two categories, emergent

properties (EPs) and community aggregated traits (CATs). Emergent properties have

been used to refer to a variety of phenomena in ecology, however here we use

emergent properties as it has been defined by Salt (1979)[24]: "An emergent property

of an ecological unit is one which is wholly unpredictable from observation of the

components of that unit", which is consistent with its contemporary use in microbial

ecology.[25] For example, the potential importance of emergent properties to influence

ecosystem processes has been demonstrated in a series of experimental flow-

through flumes that mimicked development and metabolism of stream biofilms[26].

Transient storage (i.e., an increase in residence time of the water and its solutes

near the biofilm relative to the flow around it) increased as the microbial biofilm

density increased[26]. Microbial biofilm formation is an EP[27] that affects the important

ecosystem process of hydrological transient storage[26]. Another example of an

emergent property is the relative abundance of a certain traits within a microbiome.

Trait based approaches have a rich history in ecology and have been increasingly

applied to address questions in multiple areas of microbial ecology.[28] For example,

specific functions (i.e. uptake of an individual organic substrate) are associated with

traits which can involve multiple genes, among different taxa, all capable of

performing the function albeit with differences in the underlying physiology

and efficiency. The distribution and expression of these functional gene variants

generates a trait structure among microbiomes, which determines the overall

performance of the microbiome for that given function (i.e. uptake of a given organic

substrate), but which cannot be predicted simply from the presence of the taxa that

carry the genes conferring that trait.[29] While characterization of EPs may improve the

understanding of microbial processes (Figure 2, Letter G) they cannot, in principle,

be estimated or predicted on the basis of the constituent taxa (i.e. membership)

alone (Figure 2, Letter F), and thus must remain as an intermediary between

environmental drivers (Figure 2, Letter C) and microbial processes (Figure 2, Letter

G).

Unlike EPs, CATs can potentially be estimated from characteristics of their

constituents and provide a pathway to link microbial community membership to the

community properties that drive microbial processes (Figure 2, Letter E)[30]. For

example, CATs may include commonly measured community properties such as

functional gene abundance as estimated from qPCR (e.g., pmoA which encodes a

subunit of the enzyme involved in methane oxidation, can be used to estimate

potential for methanotrophy and as a phylogenetic marker for methanotrophs)[31]. A

recent perspective article discussed the role of CATs in microbial ecology and noted

a series of additional CATs (e.g., maximum growth rate, dormancy, osmoregulation)

that could be inferred from metagenomic data of the extant community[30].

Understanding when, and which, community properties that shape microbial

processes can be predicted by membership is a critical research question, and an

important step in understanding how the microbiome contributes to system level

processes. Whether or not a community property is an EP or a CAT is an exciting

area of research and provides an important framework to advance research at the

microbial-ecosystem nexus. New approaches, like studying higher-level interactions

in ecological communities could help understand how a microbiome's constituents

interact to from emergent properties.[32] This is not a trivial task, yet a suite of existing

methods, discussed below, already provides the ability to directly pursue this

challenge.

Microbial Community Membership - Although the now commonplace analysis of

community membership by sequencing phylogenetic marker genes (e.g., regions of

the 16S, 18S, or ITS genes) or suites of phylogenetically conserved protein

sequences identifies constituent microbial taxa, the direct coupling of microbial

phylogeny to physiology and ecology remains elusive (Figure 2, Letter H).[33,34,35] In

general the paucity of associated physiological data or information on population

phenotypes that accompany phylogenetic sequence data limits the system-level

inference that is possible from analyses of community membership. This constrains

our ability to attribute microbial processes to community membership of even

relatively simple environmental consortia. Whereas it is clear that microbial

populations are not randomly distributed in space and time[35], and that some

microbial traits are conserved at coarse taxonomic scales [28,36,37], the physiological

mechanisms underlying non-random distributions of microbial taxa across

environmental gradients is often unknown. The limited understanding of the

metabolism of most bacterial phyla limits an explicit understanding between the

organism's abundance and its role in the microbial process that contributes to an

ecosystem process.

A Path Forward

We suggest that a challenging but necessary step for microbiome science is to

move away from identifying correlative relationships between characteristics of the

microbiome and system level processes, and towards identifying more causative and

mechanistic relationships. The conceptual diagram (Figure 2) is a road map to

organize and link the diverse suite of measureable microbial characteristics that are

currently available to researchers. Figure 2 does not represent how these

components necessarily interact in the environment; rather it is a map that identifies

potential links between measureable microbial and system-level characteristics that

can help structure our exploration of how microorganisms influence the systems they

inhabit. Ecosystem ecology has traditionally been confined to interactions between

environmental parameters and ecosystem processes (depicted within the horizontal

arrow, Figure 2). Similarly, microbial ecology (depicted within the vertical arrow,

Figure 2) has historically focused on phylogenetically undefined aspects of microbial

communities (e.g., bacterial abundance) and microbial processes (e.g., bacterial

production) or on the physiology of microbial isolates (e.g., sulfate reducing bacteria)

or the collective physiology of highly reduced communities with known membership

(e.g., waste water treatment microbiome). The routine inclusion of sequence-based

approaches in studies of environmental microorganisms has lead to an increasingly

detailed description of the world's microbiomes and an increasing interest in how

constituents of those communities interact to influence the system as a whole.

The drive to include microbial characteristics into system-level science has led to a

range of approaches for linking characteristics of the microbiome to ecosystem

processes. Direct connections between microbial membership and ecosystem

processes (Figure 2, Letter I), or community properties and ecosystem processes

(Figure 2, Letter J), have proven consistently difficult to establish[3,4]. We propose 1)

identifying which microbial processes are likely to contribute to ecosystem-level

pools and fluxes *a priori* (Figure 2, Letter K), 2) determining which microbial

community properties best describe and predict these microbial processes (Figure 2,

Letter G), and 3) identifying whether the community properties that best describe

each process are a CAT or an EP (Community Properties, Figure 2). If the

community property is a CAT then exploring the link between microbial membership

and community properties may lead to further understanding and perhaps an

enhanced predictive power (Figure 2, Letter E). However, if the community property

is an EP elucidating the microbial membership that contributes to the EP is unlikely

to improve understanding of the drivers of that community property (Figure 2, Letter

F) and understanding how environmental drivers structure the EP will be more

insightful. Formalizing microbiome research into a structured, conceptual framework

will allow the research community to better focus on potential links between

microbiome characteristics and system-level processes that are most likely to be

detected empirically. This approach will also allow researchers working in different

systems to test the same pathways among defined microbiome characteristics and

thus increase the possibility of understanding the casual mechanism (or absence of

causality) for observed correlations. Equally as important, we suggest that

attempting to link community membership (Figure 2, Letter I) or community

properties (Figure 2, Letter J) directly to ecosystem processes is by definition

correlative and therefore a less powerful approach to integrating microbiome

characteristics into system-level science. Thus future research endeavors will be

most powerful if they focus on elucidating connections through the complete path of

microbial ecology (Figure 2, blue arrow, Letters E, F, and G) and not direct

connections between microbial membership or community properties and ecosystem

processes (Figure 2, Letter I and J).

Applying and Testing the Proposed Framework

Applying and testing the proposed framework will depend on the ability to more

robustly evaluate each category of microbial characteristics and to directly measure

the arrows that connect each category (Figure 2). Both labeling/sorting approaches

and phenotypic description of isolates provide an opportunity to better understand

how microbial membership contributes to community properties (Figure 2, Letter E or

F). Labeling and cell sorting approaches (e.g., fluorescent in situ hybridization (FISH)

coupled with flow cytometry cell sorting[38], or immunocapture such as with

bromodeosyuridine, BrdU)[39] provide powerful tools to constrain the complexity of the

microbiome and directly test hypotheses that link membership to community

properties or microbial processes. Labeling and sorting techniques allow the cells

that can be targeted with a stain or other label to be separated from the broader

community and then assayed for membership or phenotypes such as activity or

biomass composition. For example, a study of an Arctic Ocean bacterial community

labeled the actively growing component of the community using BrdU and then

separated those populations from the rest of the community using an immunocapture

technique to better understand the portion of the microbiome that was driving

community dynamics[39]. By simplifying the community, researchers were able to link

membership to secondary production (a microbial process, Figure 2, Letter H) and

begin to better understand which members of the community were contributing to

secondary production.

In addition, physiological studies of isolates from a broader distribution of

representative phyla are key to advancing our understanding of how membership

contributes to community properties (Figure 2, Letter E). Finding isolates that are

representative of important community properties has the potential to better

understand phenotypic plasticity and how constituent populations do (CATs) or do

not (EPs) contribute to a community property[15]. For example, work on the marine

bacterioplankton SAR11 has led to an increased understanding of how this

ubiquitous member of the marine microbiome interacts with elemental cycles in the

open ocean[40]. Similarly, a rich body of work on multiple isolates of the comparably

ubiquitous photoautotroph *Prochlorococcus* has advanced our understanding of the

ecology and physiology of one of the most abundant phototrophs on the planet[41].

Detailed studies of isolates of common environmental OTUs have clearly

demonstrated immense variation within a given OTU (i.e., "microdiversity") that in

part explains the challenge of linking membership to a community property[16]. For

example, work on *Prochlorococcus has led to* a better understanding of how

ecotypes within a single taxonomic unit (OTU) can lead to specialization in

temperature and substrate affinity[41]. OTUs that form a substantial portion of the

microbiome's sequence abundance provide potential candidates for further

investigation of possible phenotypic plasticity and or microdiversity[16]. For example, a

single phylotype of the class *Spartobacteria* within the phyla *Verrucomicrobia* was

found to be present in a broad range of soil ecosystems and comprised as much as

31% of all 16S sequences returned from prairie soils[42], making it an excellent

candidate for targeted isolation and physiological studies. Whereas it is challenging

to isolate and culture many microorganisms from the environment, existing

approaches to isolation have been led to successful isolation of both abundant and

rare members of environmental microbiomes. A recent study isolated members of an

apple orchard soil microbiome where most isolates were from the least abundant

members of the community[43]. Previous studies have had success isolating members

of the pelagic marine microbiome by using filter-sterilized seawater with a dilution to

extinction approach[44]. Thus there is a potential to target both abundant[40,41] and rare[43]

members of diverse microbiomes to learn more about their influence on community

level properties. Studies of environmental isolates are essential in building a broader

understanding of how community membership does or does not contribute to

community properties (H2, Letter E and F).

The most commonly measured microbial characteristics can be associated with

one of the three categories of microbial characteristics defined here. However, the

key to moving from a correlative and descriptive approach to a causative and

mechanistic approach comes in measuring the arrows represented by letters in

Figure 2. There is a suite of powerful methods already being employed in microbial

ecology that can actively measure many of the arrows illustrated in Figure 2. These

include: stable isotope probing of mixed communities[45], single cell methods that can

assay cells in the physiological state they occur in in the environment[46], and labeling

individual cells with stable isotopes for single cell analyses[46]. Studies that use stable

isotope probing or any form of tracking stable isotopes into a population have been

successful in linking microbial membership to microbial processes (Figure 2, Letter

H). For example, a study of sulfate reduction in a Scottish peatland using SIP

revealed that a single species of *Desulfosporosinus* was most likely responsible for

the totality of sulfate reduction within the peatland even though it only comprised

0.006% of the retrieved 16S rRNA gene sequences[47]. In this case, the

Desulfosporosinus species represented the only known sulfate reducer within the

community and thus the kinetics of this organism seemingly defined the kinetics of

sulfate reduction for the entire system. Whereas this is a single example of using SIP

to link microbial membership to microbial processes (Figure 2, Letter H), there is a

suite of culture-free techniques (such as Raman microspectroscopy (MS),

NanoSIMS, or X-ray microanalysis, XRMA) that complement sequence-based

microbiome analyses by reporting on the physiological and phenotypic

characteristics of individual cells *in situ* [46,48,49]. For example, Raman MS has the

ability to elucidate the macromolecular composition of uncultured individual cells *in*

situ[46]. Incorporation of stable isotopes into a cell's macromolecules can be visualized

as a shift in the Raman spectra. This provides information not only on which cells are

incorporating the substrate but what macromolecular pool those substrates are being

allocated to. Similarly, NanoSims allows for visualization of isotopes within a cell,

and while it cannot be used to identify which macromolecule pool an isotopic label

went into, it can visualize which cells are actively incorporating isotopically labeled

substrates. Both Raman and NanoSims can be coupled with a range of *in situ*

hybridization techniques (e.g., fluorescent in situ hybridization, FISH) to identify

which populations are contributing to community properties (Figure 2, Letter E) or

microbial processes (Figure 2, Letter H). For example, the study of a microbial

consortia from the Sippewissett Salt Marsh on the coast of Massachusetts, USA

used a combination of FISH and NanoSIMs to confirm a syntrophic association

between a population of autotrophic purple sulfur bacteria and heterotrophic sulfate

reducing bacteria (SRB)[50]. Whereas several pieces of evidence pointed to a

syntrophic association, the authors confirmed the association by first using FISH to

visualize the physical association of each population. Further analysis with

NanoSIMS after incubation with ^{34}S enriched sulfate and ^{14}C enriched bicarbonate

confirmed the presence of carbon fixation by the purple photosynthetic bacteria with

sulfide as the electron donor was coupled to the reduction of sulfate by the SRB.

Using existing methods of confirmatory ecophysiology allows for direct

measurements of the arrows connecting membership with microbial processes in a

stable microbial consortia (in this case both carbon fixation and sulfate reduction,

Figure 2, Letter H). These approaches applied in concert with sequence-based

analyses have the potential to empirically link the categories of microbial information

defined here (Figure 2), moving microbiome science from a descriptive and

correlative approach to a mechanistic and causative approach.

These culture free approaches also create the potential to begin to determine

which community properties are EPs, and which are CATs. For example, microbial

community biomass stoichiometry (e.g., biomass C:N or C:P) cannot currently be

predicted (or even constrained) from a list of it's constituent taxa (Figure 2, Letter F).

However, microbial biomass stoichiometry is a community property with power to

predict the microbial contribution to nutrient cycling (Figure 2, Letter G).[19-22]

Independently, the biomass stoichiometry of microbial isolates grown on the same

media has been shown to differ among different taxa suggesting a relationship

between an organisms' identity and the elemental composition of its biomass. [51,52]

However, there is an abundance of evidence that suggests that the biomass

stoichiometry of many isolates is a function of the stoichiometry of the media they

were cultured on[52]. Electron dispersal spectroscopy (EDS) has the power to

measure the C:N:P of individual bacterial cells growing *in situ* (i.e. not in culture)[49].

The potential to couple EDS analysis with a phylogenetic label presents the

opportunity to assay mixed microbial communities and assess the link between

phylogenetic identity and biomass stoichiometry under natural conditions[53]. Thus,

community biomass stoichiometry can potentially be deconstructed into the biomass

stoichiometry of its constituent taxa[53]. This approach would provide a direct link

between community membership and a community property (e.g., biomass C:N,

Figure 2, Letter E), that influences an important microbial process (i.e. nutrient

recycling).

Designing microbiome research

It is critical that we recognize the influence of the taxonomic and functional

composition of the microbiome is exerted through multiple pathways, some that are

direct and can be readily identified, some that are indirect, and mediated by complex

interactions at the community level. We must also recognize that the influence of

microbiomes will vary depending on the system-level process in question, because

analysis of microbial characteristics may simply not improve the environmentally-

based prediction of certain processes (Figure 2, Letter A), whereas other system-

level processes may indeed benefit from the inclusion of microbial characteristics

including membership. For the latter, the challenge then is to determine which

microbial category is the most relevant predictor of the system level process of

interest: microbial processes, community properties, or microbial membership.

Establishing the links between these microbial dimensions (Figure 2, Letters E, F,

and G) further contributes to our understanding of the mechanistic underpinnings

that affect system level processes and thus will have greater explanatory power in a

broader range of systems.

The framework presented here provides one approach to formalize inquiry across

microbiome science and encourages empirical linkages between the presence of

organisms in a system and the processes that characterize that system. Whereas we

draw examples from environmental microbiomes and the ecosystems they inhabit,

this structured approach has the potential to benefit the analysis of microbiomes

associated with other systems such as host organisms and those of the built

environment. As important as establishing causal links among microbial

membership, community properties, microbial processes, and ecosystem processes,

is determining when these links are unlikely to be present. Research that

indiscriminately seeks to identify correlations, which does not recognize the

hierarchy of effects, and that places all metrics on an equal plane are susceptible to

confirmation bias and will continue to yield conflicting and ambiguous results that not

only fail to provide new insight into ecosystem processes, but also blurs the

connections that do exist. We suggest that rather than looking for linkages among

microbiome membership and system-level processes in every study, research efforts

would benefit from strategically targeting the linkages and processes for which an *a*

priori understanding of microbial physiology should allow us to improve our

understanding of the ecosystem process.

Acknowledgements

This work is a product of the Next Generation of Ecosystem Indicators Working Group, supported by the USGS John Wesley Powell Center for Synthesis and Analysis. Preparation of this manuscript was supported by NSF DEB IOS #1456959 awarded to EKH. Chuck Pepe-Ranney and Ariane Peralta provided valuable feedback on previous versions of this manuscript. This paper is dedicated to Diana Nemergut, an integral part of our working group who passed away during the preparation of this manuscript. She is one of a kind.

Figures

Figure 1 Diagram of microbial-ecosystem linkages A) how linkages are commonly conceptualized across levels of ecological organization and B) the series of ecological phenomena that create challenges when attempting to link metrics from one level of ecological organization to the other.

Figure 2 Shown is a conceptual map of the intersection between microbial (vertical) and ecosystem (horizontal) ecology with each of the three categories of microbial characteristics (microbial processes, community properties, and microbial membership) as defined in the text. We argue for an increased focus on studies that elucidate pathways E, F, and G. In addition, we note that pathways I and J are less likely to effectively incorporate microbiome characteristics into system-level science. The delta symbol in each category indicates an emphasis on how changes within a category may lead to a change in a connected category. The dotted arrow for letter F denotes that many emergent properties cannot currently be linked to membership and is an important area for active research.

References

1. Graham E. B., Wieder, W.R., Leff, J.R., Weintraub, S. W., Townsend, A.R., Cleveland, C.C., Philippot L., Nemergut D.R. (2014) Do we need to understand microbial communities to predict ecosystem function? A comparison of statistical models of nitrogen cycling processes Soil Biology and Biochemistry 68, 279–282

2. Graham E. B. et al. (2016) Microbes as engines of ecosystem function: when does community structure enhance predictions of ecosystem processes? Frontiers in Microbiology 7: 214 doi:10.3389/fmicb.2016.0021

3. Rocca J.D., Hall E.K., Lennon J.T., Evans S.E., Waldrop M.P., Cotner J.B., Nemergut D.R., Graham E.B., Wallenstein M.D. (2015) Relationships between protein-encoding gene abundance and corresponding process are commonly assumed yet rarely observed ISME J. 9: 1693–1699

4. Bier R.L., E.S. Bernhardt, CM Boot, EB Graham, EK Hall, JT Lennon, D Nemergut, B Osborne, C Ruiz-González, JP Schimel, MP Waldrop, MD. Wallenstein (2015) How are we forging conceptual, analytical, and mechanistic links between microbial community structure and ecosystem process? FEMS microbiology ecology FEMS microbiology ecology 91(10): fiv113

5. Falkowski, P.G., Fenchel, T., and DeLong, E.F. (2008) The microbial engines that drive earth's biogeochemical cycles Science 320: 1034-1038

6. Felip M., Pace M.L., Cole, J.J. (1996) Regulation of planktonic bacterial growth rates: The effects of temperature and resources. Microb. Ecol. 31(1): 15-28.

7. Lennon, J. T. and S. E. Jones (2011) Microbial seed banks: the ecological and evolutionary implications of dormancy. Nature Microbiology Reviews 9: 119-130.

8. Comte, J., Fauteux, L. and del Giorgio, P.A. (2013) Links between Metabolic Plasticity and Functional Redundancy in Freshwater Bacterioplankton Communities. Frontiers in Microbiology 4: 112. doi: 10.3389/fmicb.2013.00112

9. Adams, H. E., Crump, B.C. and Kling, G.W. (2014) Metacommunity Dynamics of Bacteria in an Arctic Lake: The Impact of Species Sorting and Mass Effects on Bacterial Production and Biogeography. Frontiers in Microbiology 5 (2014): 82. 10.3389/fmicb.2014.00082

10. Daims, H., Lebedeva, E.V., Pjevac, P., Han, P., Herbold, C., Albertsen, M., Jehmlich, N., Palatinszky, M., Vierhellig, J., Bulaev, A., Kirkegaard, R. H., von Bergen M., Rattei, T., Bendinger, B., Nielsen, P.H., Wagner, M. (2015) Complete nitrification by Nitrospira bacteria Nature 528: 504-509 doi: 10.1038/nature16461

11. van Kessel, Jaartje A.H.J., Speth, D.R., Albertsen, M., Nielsen, P.H., Op den Camp, H.J.M., Kartal, B.,Jetten, M.S.M., Lueker, S. (2015) Complete nitrification by a single microorganism 528: 555-559 doi: 10.1038/nature16549

12. Schimel, J.P., J. Bennett, and N. Fierer. (2005) Microbial community composition and soil N cycling: is there really a connection? In: Biological diversity and function in soils. Bardgett, R.D., D.W. Hopkins, and M.B. Usher (Eds.) Cambridge University Press. pp. 171-188

13. Schimel, J.P. (1995) Ecosystem consequences of microbial diversity and community structure in Arctic and Alpine Biodiversity: Patterns, Causes and Ecosystem Consequences Ecological Studies Volume 113, pp 239-254

14. Lenhart K., M. Bunge, S. Ratering, T. R. Neu, I. Schüttmann, M. Greule, C. Kammann, S. Schnell, C. Müller, H. Zorn & F. Keppler. (2012) Evidence for methane production by saprotrophic fungi Nature Comm. 3:1046

15. Hagstroem, A., Azam, F., Berg, C., Zweifel U.L. (2017) Isolates as models to study bacterial ecophysiology and biogeochemistry Aquat. Microb. Ecol. 80:15-27

16. Larkin, A.A, and Martiny, A. (2017) Microdiversity shapes the traits, niche space and biogeography of microbial taxa Environ. Microbiol. Rep. 9(2): 55-70 doi: 10.111/1758-2229

17. German D.P., Marcelo, K.R.B., Stone, M.M and Allison, S.D. (2012) The Michaelis-Menten kinetics of soil extracellular enzymes in response to temperature: a cross-latitudinal study Global Change Biology 18: 1468-1479

18. Hall, E. K., A. R. Dzialowski, S. M. Stoxen, and J. B. Cotner. (2009) The effect of temperature on the coupling between phosphorus and growth in lacustrine bacterioplankton communities. Limnol. and Oceanogr. 54: 880-889.

19. Manzoni, S., R. B. Jackson, J. A. Trofymow, and A. Porporato. (2008) The global stoichiometry of litter nitrogen mineralization. Science 321: 684-686.

20. Buchkowski, R.W., Schmitz, O.J., Bradford, M.A. (2015) Microbial stoichiometry overrides biomass as a regulator of soil carbon and nitrogen cycling. Ecology 96: 1139-1149.

21. Elser, J.J., Chrzanowski, T.H., Sterner, R.W., Schampel, J.H., Foster D.K. (1995) Elemental ratios and the uptake and release of nutrients by phytoplankton and bacteria in three lakes of the canadian shield Microbial Ecology 29: 145-162

22. Goldman, J.C., Caron, D.A., and Dennett, M.R. (1987) Regulation of gross growth efficiency and ammonium regeneration in bacteria by substrate C:N ratio Limnology and Oceanography 32 (6): 1239-1252

23. Redfield, A.C (1958) The biological control of chemical factors in the environment American Scientist 46 (3): 205-221

24. Salt, G.W. (1979) A comment on the use of the term emergent properties American Naturalist 113: 145-148

25. Konopka, A. (2009) What is microbial community ecology? ISME J., 3 (11): 1223-12230

26. Battin, T., L. A. Kaplan, L. Newbold, X. Cheng, and C. Hansen. (2003) Effects of current velocity on the nascent architecture of stream microbial biofilms. Appl. and Env. Microbiol. 69: 5443-5452.

27. Flemming, H-C., Wingender, J., Szewzyk, U., Steinberg, P., Rice, S.A., Kjelleberg, S. (2016) Biofilms: an emergent form of bacterial life Nature Reviews Microbiology 14: 563-575

28. Martiny JBH, Jones SE, Lennon JT, Martiny AC (2015) Microbiomes in light of traits: a phylogenetic perspective. Science 350: doi: 10.1126/science.aac9323

29. Ruiz-González, C., Niño-García, J. P., Lapierre, J. F., & del Giorgio, P. A. (2015) The quality of organic matter shapes the functional biogeography of bacterioplankton across boreal freshwater ecosystems. Global Ecology and Biogeography 24(12): 1487-1498.

30. Fierer, N., A. Barberán, Laughlin, D. (2014) Seeing the forest for the genes: Using metagenomics to infer the aggregated traits of microbial communities. Frontiers in Microbiology 5: 614

31. Judd, C.R., Koyama, A., Simmons, M.P., Brewer, P., and von Fischer, J.C. (2016) Co-variation in methanotroph community composition and activity in three temperate grassland soils Soil Biol. and Biochem. 95:78-86

32. Grilli, J., Barabas G., Michalska-Smith, M.J., and Allesina, S. (2017) Higher-order interactions stabilize dynamics in competitive network models Nature. 2017 548: 210–213

33. Newton, R. J, S. E. Jones, A. Eiler, K. D McMahon, Bertilsson, S. (2011) A guide to the natural history of freshwater lake bacteria. Microbiol. and Molec. Biol. Rev. 75:1, doi:10.1128/MMBR.00028

34. Hug, L.A., Baker, B.J., Anantharaman, K., Brown, C.T., Probst, A.J., Castelle, C.J., Butterfield, C.N., Hernsdorf, A.W., Amano, Y., Ise, K., Suzuki, Y., Dudek, N., Relman, D.A., Finstad, K.M., Amundson, R., Thomas, B.C., Banfield, J.F. (2016) 1: 16048 doi: 10.1038/nmicrobiol.2016.48

35. Martiny, J.B.H., B. Bohannan, J. Brown, R. Colwell, J. Fuhrman, J. Green, M.C. Horner-Devine, M. Kane, J. Krumins, C. Kuske, P. Morin, S. Naeem, L. Ovreas, A.-L. Reysenbach, V. Smith, J. Staley. (2006) Microbial biogeography: Putting microorganisms on the map. Nature Reviews Microbiology 4: 102-112

36. Lennon JT, Aanderud ZA, Lehmkuhl BK, Schoolmaster DR (2012) Mapping the niche space of soil microorganisms using taxonomy and traits. Ecology 93: 1867-1879

37. Treseder KK, Lennon JT (2015) Fungal traits that drive ecosystem dynamics. Microbiology and Molecular Biology Reviews 79: 243-262

38. Czechowska, K., Johnson, D.R., van der Meer, J.R. (2008) Use of flow cytometric methods for single-cell analysis in environmental microbiology Curr. Opin. In Microbiol. 11:3, 205-212

39. Galand P.E., L. Alonso-Sáez, S. Bertilsson, C. Lovejoy, and E. O. Casamayor (2013) Contrasting activity patterns determined by BrdU incorporation in bacterial ribotypes from the Arctic Ocean in winter Front Microbiol. (2013); 4: 118

40. Giovannoni S.J. 2017 SAR11 Bacteria: The most abundant plankton in the oceans. Annual Review Marine Science 9: 231-255

41. Biller, S.J., Berube, P.M., Lindell, D., Chisholm, S.W. (2015) *Procholorcoccus:* The structure and function of collective diversity Nature Reviews Microbiology 13:13-27

42. Brewer, T.E., Handley, K.M., Carini, P., Gilbert, J.A., Fierer, N. (2016) Genome reduction in an abundant and ubiquitous soil bacterium 'Candidatus Udaeobacter copiosus' Nature Microbiology 2: 16198

43. Shade, A., Hogan, C.S., Klimowicz, A.K., Linske, M., McManus, P.S., and Handelsman, J. (2012) Culturing captures members of the soil rare bioshphere Environmental Microbiology 14(9): 2247-2252

44. Schut, F., De Vries, E.J., Gottschal, J.C., Robertson, B.R., Harder, W., Prins, R.A., Button, D.K. (1993) Isolation of typical marine bacteria by dilution culture: Growth, maintenance and characteristics of isolates under laboratory conditions Applied and Environmental Microbiology 59(7): 2150-2160

45. Neufeld J.D., Vohra J, Dumont M.G., Lueders T., Manefield M., Friedrich M.W., Murrell J.C. (2007) DNA stable-isotope probing. Nat Protoc. 2:4, 860-866

46. Wagner, M. (2009) Single cell ecophysiology of microbes as revealed by Raman microspectroscopy or secondary ion mass spectrometry imaging. Annu Rev Microbiol 63:411-429.

47. Pester, M., Bittner, N., Deevong, P., Wagner, M., and Loy, A. (2010) A 'rare biosphere' microorganism drives sulfate reduction in a peatland. ISME J. 4:1591–1602.

48. Behrens, S., Kappler, A., Obst M. (2012) Linking environmental processes to the in situ functioning of microorganisms by high-resolution secondary ion mass spectrometry (NanoSIMS) and scanning transmission X-ray microscopy (STXM) Env. Microbiol. 14:11, 2851-69

49. Norland S, Fagerbakke K, Heldel M. (1995) Light element analysis of individual bacteria by X-ray microanalysis. Appl. Environ. Microbiol. 61:1357–62.

50. Wilbanks, E.G., Jaekel, U., Salman, V., Humphrey, P.T., Eisen, J.A., Faciotti, M.T., Buckley, D.H., Zinder, S.H., Druschel, G.K., Fike, D.A., Orphan,V.J., (2014) Microscale sulfur cyclingin the photophrophic pink berry consortia of the Sippewissett Salt Marsh Environ. Micro 16: 3398-3415

51. Mouginot, C., Kawamura, R., Matulich, K.L., Berlemont, R., Allison, S.D., Amend, A.S. and Martiny, A.C. (2014) Elemental stoichiometry of fungi and bacteria strains from grassland leaf litter Soil Biol. & Biochem. 76, 278-285

52. Godwin, C.M. and Cotner, J.B. (2015) Stoichiometric Flexibility in Diverse Aquatic Heterotrophic Bacteria Is Coupled to Differences in Cellular Phosphorus Quotas. Frontiers in Microbiology 6: doi:10.3389/fmicb

53. Hall, E.K., Maixner F., Franklin O., Daims, H., Richter, A. and Battin, T. (2011) Linking microbial and ecosystem ecology using ecological stoichiometry: A synthesis of conceptual and empirical approaches Ecosystems 14: 261-273