Understanding How Microbiomes Influence The Systems They Inhabit: Moving From A Correlative To A Causal Research Framework

Authors: Hall, E.K.[1*], Bernhardt, E.S[2], Bier, R.L.[2], Bradford, M.A.[3], Boot, C.M.[1], Cotner, J.B.[4], del Giorgio, P.A.[5], Evans, S.E.[6], Graham E. B [2,7,8], Jones, S.E.[9], Lennon, J.T.[10], Locey, K.J.[10] Nemergut, D.[2], Osborne, B.B.[11], Rocca, J.D.[1,2], Schimel J.S.[12], Waldrop, M.P.[13], Wallenstein, M.W.[1]

Article Type: Perspectives
Words in Abstract: 262
Words in Main Text: 4,238
2 Figures, 0 Tables, 45 References

Running Title: How microbiomes influence ecosystems
Keywords: microbiome, microbial ecology, ecosystem ecology, emergent properties, community aggregated traits

Affiliations: [1]Colorado State University, Fort Collins CO (ed.hall@colostate.edu, claudia.boot@colostate.edu, matt.wallenstein@colostate.edu) [2]Duke University, Durham, N.C. (ebernhar@duke.edu, ebgraham2@colorado.edu, jennifer.rocca@duke.edu) [3]Yale University, New Haven, CT, Dept. of Forestry and Environmental Studies (mark.bradford@yale.edu), [4]University of Minnesota, Saint Paul, MN (cotne002@umn.edu) [5]Université du Québec à Montréal, Montréal, CA (del_giorgio.paul@uqam.ca) [6]Michigan State University, Hickory Corners, MI (evanssar@gmail.com) [7]Institute of Arctic and Alpine Research, University of Colorado at Boulder [8]Pacific Northwest National Laboratory, Richland, WA, USA [9]University of Notre Dame, South Bend, IN (sjones20@nd.edu) [10]Indiana University, Bloomington, IN (lennonj@indiana.edu, ken@weecology.org) [11]Brown University, Providence, RI (brooke_osborne@brown.edu) [12]University of California, Santa Barbara, Santa Barbara, CA (josh.schimel@lifesci.ucsb.edu) [13]U.S. Geological Survey, Menlo Park, CA (mwaldrop@usgs.gov)

* **Corresponding Author**: Dr. Ed Hall, Department of Ecosystem Science and Sustainability, Natural Resource Ecology Laboratory Campus Delivery 1499, Colorado State University, Fort Collins, CO 80523, ed.hall@colostate.edu, 970-491-2162

Contribution: All listed authors have contributed to the conceptualization, writing, and preparation of the current manuscript.

Abstract

Translating the ever-increasing wealth of information on microbiomes (environment,

host, or built environment) to advance the understanding of system-level processes

is proving to be an exceptional research challenge. One reason for this challenge is

that relationships between characteristics of microbiomes and the system-level

processes they influence are often evaluated in the absence of a robust conceptual

framework and reported without elucidating the underlying causal mechanisms. The

reliance on correlative approaches limits the potential to expand the inference of a

single relationship to additional systems and advance the field. We propose that

research focused on how microbiomes influence the systems they inhabit should

work within a common framework and target known microbial processes that

contribute to the system-level processes of interest. Here we identify three distinct

categories of microbiome characteristics (microbial processes, microbial community

properties, and microbial membership) and propose a framework to empirically link

each of these categories to each other and the broader system level processes they

affect. We posit that it is particularly important to distinguish microbial community

properties that can be predicted from constituent taxa (community aggregated traits)

from and those properties that are currently unable to be predicted from constituent

taxa (emergent properties). Existing methods in microbial ecology can be applied to

more explicitly elucidate properties within each of these categories and connect

these three categories of microbial characteristics with each other. We view this

proposed framework, gleaned from a breadth of research on environmental

microbiomes and ecosystem processes, as a promising pathway with the potential to

advance discovery and understanding across a broad range of microbiome science.

Current Approaches Linking Microbial Characteristics and Ecosystem Processes

Virtually all ecosystem processes are influenced by microorganisms, and many processes are carried out exclusively by microorganisms. This has sometimes led to the assumption that a better description of the microbiome (including its associated transcripts, proteins, and metabolic products) should lead to a better understanding and predictions of system level processes. However, such justifications assume that measurable characteristics of the microbiome (e.g. 16S rRNA gene libraries, metagenomes, enzymatic activities) can inform our ability to better understand and predict system-level processes. Unfortunately, additional information about the microbiome does not always provide a clearer understanding of ecosystem processes beyond what can be predicted by environmental factors alone[1,2].

Two recent meta-analyses[3,4] suggest that research at the intersection of ecosystem science and microbial ecology often relies on assumed relationships between microbiome characteristics and ecosystem processes and often do not test to see if those relationships are present. The first, an examination of 415 studies, found little evidence that protein-encoding genes (sometimes referred to as "functional genes") or gene transcripts correlate with associated biogeochemical processes[3]. Although all studies attempted (or presumed) to link microbial genes or transcripts with function, only 14% measured both the abundance of genes or transcripts and the corresponding process. When the relationship between microbial characteristic and ecosystem process was measured and tested only 38% exhibited a positive correlation[3]. This result was consistent whether functional gene or transcript abundance was used as the response variable.

The second study compiled a separate dataset of 148 studies that

examined microbial membership and ecosystem processes in response to

experimental manipulations[4]. Whereas 40% of included studies reported

concomitant changes in microbial membership and an ecosystem process, only one

third of those cases reported the relationship between microbial membership and an

ecosystem process. Interestingly, of the 53 studies that posed a hypothesis

about links between microbial membership and ecosystem processes, more than

half (53%) did not report testing for a statistical link between membership and

processes[4].

Microbiomes are the engines that power system-level processes[5]. However the

meta-analyses described above illustrate that the current approach to study the links

between microbiome characteristics and ecosystem processes are not well

formulated and relationships between microbiome characteristics and system level

processes are implied yet rarely tested. When linkages are tested, significant

correlations between microbiome characteristics and ecosystem processes are

sometimes present, but more frequently not[3,4]. One reason for the ambiguity

between microbiome characteristics and system level processes is that many studies

are conducted in the absence of a conceptual framework that illustrates how different

measurable microbial characteristics relate to one another and to the system level

process of interest.

Challenges in Linking Microbial Characteristics and Ecosystem Processes

Current conceptual research frameworks that attempt to link microbial information

to a system-level process often do not effectively align with the methods being

applied or the data those methods generate. For example, environmental factors act

on the physiology of individual organisms, which alters their competitive ability,

abundance, and ultimately their contribution to an ecosystem process. However,

designing an observational study or experiment from this framework assumes that

environmental and microbial characteristics are measurable across multiple

categories of ecological organization (i.e., individuals, populations, and communities)

at the temporal and spatial scales at which they influence system level processes

(Figure 1a). In addition, the relationships between environmental variables and

microbial characteristics can be decoupled in both time and space[4], and are often

non-linear[6]. Recent immigration, phenotypic plasticity, disequilibrium between the

environment and the extant microbiome at the time of sampling, functional

redundancy, and dormancy can all mask the relationship between measurable

microbial characteristics and the processes microorganisms influence (Figure

1b).[4,7,8,9] As micrometer scale characteristics of microbiomes (10^{-6} m) are scaled to

the level of ecosystems (m to km), we assume that our conceptual understanding is

also scalable. However, each of the aforementioned confounding factors aggregate

over multiple orders of magnitude often masking the very relationships we seek to

elucidate (Figure 1b). To formalize how measurable microbiome characteristics are

linked with system-level processes we have conceptually defined the intersection of

microbial and ecosystem ecology and identified three categories of microbial

characteristics to illustrate how they interact with each other and may contribute to

an ecosystem process (Figure 2).

Mapping Ecosystem Processes to Microbial Characteristics

Defining the ecosystem process, its critical sub-processes, and the known

phylogenetic distribution of the metabolic pathways that drive those sub-processes

creates an explicit conceptual pathway that links the ecosystem process to the

microorganisms that contribute to it. Ecosystem processes are defined as qualitative

changes in a pool, or a flux from one pool to another (e.g., NH_4^+ to NO_3^-, or dissolved

organic matter to CO_2). The first step to understand how a microbiome influences an

ecosystem process is to define the ecosystem process of interest and each sub-

process that contributes to it (i.e., the set of constituent reactions that combine to

determine a net flux). Ecosystem processes are composites of complementary or

antagonistic sub-processes, carried out by phylogenetically and metabolically

diverse microorganisms[10]. For example, net ecosystem productivity (NEP) is the

balance between antagonistic processes of C-fixation and C-mineralization. Each

sub-process of NEP can be further partitioned into a series of metabolic pathways

(e.g., chemoautotrophic nitrification and photoautotrophic C-fixation or heterotrophic

fermentation and aerobic respiration). Partitioning each ecosystem process in this

hierarchical manner can continue until the sub-processes maps directly to specific

microbial metabolic pathways (e.g., acetoclastic methanogenesis). Subsequently,

each of these metabolic pathways can be categorized as either phylogenetically

broad or narrow[11]. Broad processes are phylogenetically common (i.e., widely

distributed among taxa), whereas narrow processes are phylogenetically conserved

(i.e., limited to a specific subset of taxa). For example, denitrification and

photosynthesis are phylogenetically broad processes, while both methanogenesis

and methanotrophy are currently thought to be phylogenetically narrow processes

(with at least one notable exception[12]).

The second step is to identify the controls or constraints on each constituent sub-

process. For example, the kinetics of a single metabolic pathway in a model

organism may help us understand the rate limiting steps of a narrow process, but

insights from model organisms are much less likely to capture the full spectrum of

responses of a broad process where phenotypic variation among phylogenetically

diverse organisms is likely to be much greater[13,14]. Once the ecosystem process has

been conceptually partitioned into its component parts and their primary controls, a

concerted approach can be applied to investigate how characteristics of the

microbiome influence the ecosystem process of interest within the complexity of a

natural environment.

Categories of Microbial Characteristics

The contribution of the microbiome to ecosystem processes is exerted through

aggregate community properties that are shaped by both microbial membership and

environmental factors. Most current studies rarely articulate how measurements of

microbial characteristics differ in their specificity (i.e., the level of phylogenetic

resolution), precision (i.e., the ability of the method to repeatedly describe the

characteristic of interest), or context (i.e., how a characteristic relates to other

microbial characteristics or the ecosystem). To aid in designing experiments that link

microorganisms to the processes they influence we propose three distinct categories

of microbial characteristics: 1) microbial processes, 2) microbial community

properties, and 3) microbial membership (Figure 2). This proposed framework allows

the researcher to clearly identify how different measurements used to characterize a

microbiome interact with each other, and to identify the potential of each

characteristic to elucidate the microbial contribution to the system level process. All

measurable characteristics of microbial communities (e.g., abundance of cells,

sequence of genes, transcripts, or proteins; enzyme expression or activity) can be

placed within one of the above categories. This conceptual structure that orientates

each microbial category within a broader context creates the opportunity to improve

the design of observational and experimental studies in microbiome research.

Microbial Processes - Microbial processes are the collective metabolisms of the

microbiome that contribute to changes in pools and fluxes of elements or compounds

(i.e., Figure 2, Letter K). This is the category of microbial information that can most

readily be incorporated into system-level models because many microbial processes

represent the key sub-processes that contribute to a particular ecosystem process

(e.g., methanogenesis + methanotrophy ≈ methane efflux). Commonly measured

microbial processes in ecosystem science include nitrogen fixation, denitrification,

nitrification, phosphorus uptake and immobilization, carbon fixation, and organic

carbon mineralization. The rates of many microbial processes can be approximated

through physiological assays (e.g., biological oxygen demand to estimate microbial

respiration), and while they do not open the "black box" of the microbial community,

they do directly quantify the microbial contribution (or potential contribution) to

changes in resources moving through the box. Microbial processes can be

distinguished from other microbial characteristics because they are all rates (i.e.,

have time in the denominator) and require a bioassay to estimate.

Assays used to estimate microbial processes are often logistically challenging,

require manipulations that inevitably deviate from *in situ* conditions, and often

depend on the environment from which the microbiome was sampled. For example,

the relationship between temperature and microbial processes such as enzyme

activity and phosphorus use efficiency vary across latitudinal gradients[15] and among

seasons[16], respectively. Thus, observations of the effect of temperature on either

enzyme activity or phosphorus use efficiency depend on where (e.g., at what

latitude) and when (e.g., during which season) they were measured. In the absence

of an understanding of the underlying physiological mechanism (e.g., the

physiological change that allows a community to perform differently at different

temperatures), the relationship between and environmental driver (e.g., temperature)

and a microbial process must be measured through a direct assay at each location

and at each time. This limits the inference possible from relying on measurements of

microbial processes alone to understand the microbial contribution to an ecosystem

process.

Microbial Community Properties - Microbial community properties include a broad

set of microbial characteristics such as community biomass or biomass elemental

ratios (e.g., biomass C:N or C:P ratios) and the majority of phylogenetically

undifferentiated aggregate sequence based measurements (e.g., gene abundance,

metagenomes, transcriptomes). Microbial community properties (Figure 2) represent

an integrated characteristic of the microbiome that has the potential to predict or at

least constrain the estimates of microbial processes. For example, microbial

community biomass C:N (a community property) has been shown to indicate a

microbiome's potential to mineralize or immobilize N in terrestrial[17], freshwater[18] and

marine[19] ecosystems.

Microbial community properties can be separated into two categories, emergent

properties and community aggregated traits. It is generally agreed that emergent

properties refer to a quality of the whole that is unique and distinguishable from the

additive properties of its constituents. Whereas in some cases, emergent properties

may be predicted from their constituent parts, for example, prediction of the physical

properties of carbon polymers is possible based on the atomic organization of

carbon atoms within each structure[20], in microbial ecology our understanding of the

constituent parts and their interactions are more often than not insufficient to predict

emergent properties. Here we use emergent properties as it has been defined

previously in ecology[21]: "An emergent property of an ecological unit is one which is

wholly unpredictable from observation of the components of that unit", which is also

consistent with its contemporary use in microbial ecology[22].

Emergent properties of microbiomes influence important ecosystem processes. For

example, a series of experiments in flow-through flumes mimicked development and

metabolism of stream biofilms[23]. Transient storage (i.e., an increase in residence

time of the water and its solutes near the biofilm relative to the flow around it)

increased as the microbial biofilm density increased.[23] In this case biofilm density

was an emergent property of the microbiome and transient storage was the microbial

process it mediated. Another example of an emergent property is the distribution of

traits that influences key microbial processes. Trait based approaches have a rich

history in ecology and have been increasingly applied to address questions in

multiple areas of microbial ecology.[24] For example, uptake of an organic substrate

can often involve the expression of multiple genes, differing among individual

organisms, all capable of performing uptake of the organic substrate, albeit with

differences in the underlying efficiency. The distribution of the expression of these

functional gene variants generates a trait distribution that is an emergent property of

the microbiome. That emergent trait distribution determines the overall performance

of the microbiome for that function (i.e., the uptake of the organic substrate), but it

cannot be predicted from the presence of the organisms conferring that trait using

current methods.[25] While characterization of emergent properties may improve the

understanding of microbial processes (Figure 2, Letter G) currently they most often

cannot be estimated or predicted on the basis of the constituent taxa (i.e.

membership) alone (Figure 2, Letter F), and thus must remain as an intermediary

between environmental drivers (Figure 2, Letter C) and microbial processes (Figure

2, Letter G).

Unlike emergent properties, community aggregated traits can potentially be

estimated from characteristics of their constituents and provide a pathway to link

microbial community membership to the community properties that drive microbial

processes (Figure 2, Letter E).[26] For example, community aggregated traits may

include commonly measured community properties such as functional gene

abundance as estimated from qPCR (e.g., *pmo*A which encodes a subunit of the

enzyme involved in methane oxidation, can be used to estimate potential for

methanotrophy and as a phylogenetic marker for methanotrophs)[27]. A recent

perspective article that discussed the role of community aggregated traits in

microbial ecology noted a series of additional putative community aggregated traits

(e.g., maximum growth rate, dormancy, osmoregulation) that could be inferred from

metagenomic data of the extant community.[26]

Understanding which community properties can be predicted by membership is a

critical research question and an important step in understanding how the

microbiome contributes to system level processes. Whether or not a community

property is likely to be an emergent property or a community aggregated trait is an

exciting area of research and provides an important framework to advance research

at the microbial-ecosystem nexus. New approaches, like studying higher-level

interactions in ecological communities could help understand how a microbiome's

constituents interact to from emergent properties.[28] This is not a trivial task, yet a

suite of existing methods, discussed below, provides the ability to directly pursue this

challenge.

Microbial Community Membership - Although the now commonplace analysis of

community membership by sequencing phylogenetic markers (e.g., ITS regions or

regions of the16S rRNA and 18S rRNA genes) or suites of phylogenetically

conserved protein sequences identifies constituent microbial taxa, the direct coupling

of microbial phylogeny to physiology and ecology remains elusive (Figure 2, Letter

H).[29,30,31] In general the paucity of associated physiological data or information on

population phenotypes that accompany phylogenetic analyses limits the system-level

inference that is possible from analyses of community membership. Even when the

physiology of an organism is known, it appears many metabolic pathways are

phylogenetically broad, and that any given microbiome will contain a diverse set of

microorganisms with the genes that encode many of the same common microbial

metabolic pathways, often referred to as "functional redundancy"[32]. There also

appears to be no consistent phylogenetic resolution at which specific microbial

metabolic pathways are constrained[32]. This constrains our ability to attribute

microbial processes to community membership of even relatively simple

environmental consortia. Whereas it is clear that microbial populations are not

randomly distributed in space and time[31], and that some microbial traits are

conserved at coarse taxonomic scales, [24,33,34] the physiological mechanisms

underlying non-random distributions of microbial taxa across environmental

gradients are often unknown. This limited understanding of the metabolism of most

bacterial phyla is one thing that currently prevents linking a microorganism's

abundance in an environment to its role in a related microbial process.

A Path Forward

We suggest that a challenging but necessary step for microbiome science is to

move away from identifying correlative relationships between characteristics of the

microbiome and system level processes, and towards identifying more causative and

mechanistic relationships. The conceptual diagram (Figure 2) is a road map to

organize and link the diverse suite of measurable microbial characteristics that are

currently available to researchers. Figure 2 does not represent how these

components necessarily interact in the environment; rather it is a map that identifies

potential links between measurable microbial and system-level characteristics that

can help structure our exploration of how microorganisms influence the systems they

inhabit. Ecosystem ecology has traditionally been confined to interactions between

environmental parameters and ecosystem processes (depicted within the horizontal

arrow, Figure 2). Similarly, microbial ecology (depicted within the vertical arrow,

Figure 2) has historically focused on phylogenetically undefined aspects of microbial

communities (e.g., bacterial abundance) and microbial processes (e.g., bacterial

production), or the physiology of microbial isolates (e.g., sulfate reducing bacteria) or

the collective physiology of highly reduced communities with known membership

(e.g., waste water treatment microbiome). The routine inclusion of sequence-based

approaches in studies of environmental microorganisms has led to an increasingly

detailed description of the world's microbiomes and an increasing interest in how

constituents of those microbiomes interact to influence the system as a whole.

Currently, researchers use a range of approaches for attempting to link

characteristics of the microbiome to ecosystem processes. Direct connections

between microbial membership and ecosystem processes (Figure 2, Letter I), or

community properties and ecosystem processes (Figure 2, Letter J), have proven

difficult to establish.[3,4] We propose 1) identifying which microbial processes are likely

to contribute to ecosystem-level pools and fluxes *a priori* (Figure 2, Letter K), 2)

determining which microbial community properties best describe and predict these

microbial processes (Figure 2, Letter G), and 3) identifying whether the community

properties that best describe each process are a community aggregated trait or a

emergent property (Community Properties, Figure 2). If the community property is

likely to be a community aggregated trait then exploring the link between microbial

membership and community properties may lead to further understanding and

perhaps an enhanced predictive power (Figure 2, Letter E). However, if the

community property is likely an emergent property elucidating the microbial

membership that contributes to the emergent property is, given the current

understanding, unlikely to improve understanding of the drivers of that community

property (Figure 2, Letter F). Formalizing microbiome research into a structured,

conceptual framework should help the research community better focus on potential

links between microbiome characteristics and system-level processes that are most

likely to be detected empirically. This approach will also allow researchers working in

different systems to test the same pathways among defined microbiome

characteristics and thus increase the possibility of understanding the causal

mechanism (or absence of causality) for observed correlations. Thus, future

research endeavors will be most powerful if they focus on elucidating connections

through the complete path of microbial ecology (Figure 2, blue arrow, Letters E, F,

and G) and not direct connections between microbial membership or community

properties and ecosystem processes (Figure 2, Letters I and J).

Applying and Testing the Proposed Framework

Applying and testing the proposed framework will depend on the ability to more

robustly characterize each category of microbial characteristics and to directly

measure the arrows that connect each category (Figure 2). Both labeling/sorting

approaches and phenotypic description of isolates provide an opportunity to better

understand how microbial membership contributes to community properties (Figure

2, Letter E or F). Labeling and cell sorting approaches (e.g., fluorescent in situ

hybridization (FISH) coupled with flow cytometry cell sorting,[35] or immunocapture,

such as with bromodeoxyuridine, BrdU)[36] provide powerful tools to constrain the

complexity of the microbiome and directly test hypotheses that link membership to

community properties or microbial processes. For example, a study of an Arctic

Ocean bacterial community labeled the actively growing component of the

community using BrdU and then separated those populations from the rest of the

community using an immunocapture technique to better understand the portion of

the microbiome that was contributing to secondary production.[36] In addition,

physiological studies of isolates that are representative of important community

properties have the potential to advance understanding of the role of phenotypic

plasticity in structuring how constituent populations do or do not contribute to

community properties (Figure 2, Letter E)[13]. Detailed studies of isolates of common

environmental OTUs have demonstrated immense variation within a given OTU (i.e.,

"microdiversity") that in part explains the challenge of linking membership to a

community property[14]. For example, work on *Prochlorococcus* has led to a better

understanding of how ecotypes within a single taxonomic unit (OTU) can lead to

specialization in temperature, and substrate affinity[37]. OTUs that form a substantial

portion of the microbiome's sequence abundance provide potential candidates for

further investigation of possible phenotypic plasticity and or microdiversity[14]. For

example, a single phylotype of the class *Spartobacteria* within the phyla

Verrucomicrobia was found to be present in a broad range of soil ecosystems and

comprised as much as 31% of all 16S rRNA gene sequences returned from prairie

soils[38], making it an excellent candidate for targeted isolation and physiological

studies. Studies of environmental isolates are essential in building a broader

understanding of how community membership does or does not contribute to

community properties (Figure 2, Letters E and F).

In addition to a better description of each category of microbial characteristics, an

important step in moving from a correlative and descriptive approach to a causative

and mechanistic approach comes in measuring the arrows represented by letters in

Figure 2. There is a suite of powerful methods already being employed in microbial

ecology that can actively measure many of the arrows illustrated in Figure 2,

including: stable isotope probing of mixed communities[39], single cell methods that

can assay cells in the physiological state they occur in in the environment, and

labeling individual cells with stable isotopes for single cell analyses[40]. Studies that

use stable isotope probing or any form of tracking isotopically labeled substrates into

a population have been successful in linking microbial membership to microbial

processes (Figure 2, Letter H). For example, a study of a Scottish peatland used SIP

to reveal that a single species of *Desulfosporosinus* was most likely responsible for

the totality of sulfate reduction within the peatland even though it only comprised

0.006% of the retrieved 16S rRNA gene sequences[41]. In addition to this example of

using SIP to link microbial membership and microbial processes (Figure 2, Letter H),

there is a suite of cultivation independent techniques (such as Raman

microspectroscopy (MS), NanoSIMS, or energy dispersive spectroscopy, EDS) that

complement sequence-based microbiome analyses by reporting on the physiological

and phenotypic characteristics of individual cells *in situ*. [40,42,43] Both Raman MS and

NanoSims can be coupled with a range of *in situ* hybridization techniques (e.g.,

fluorescent in situ hybridization, FISH) to identify which populations are contributing

to community properties (Figure 2, Letter E) or microbial processes (Figure 2, Letter

H). For example, a study of a microbial consortia from the Sippewissett Salt Marsh

on the coast of Massachusetts, USA used a combination of FISH and NanoSIMs to

confirm a syntrophic association between a population of autotrophic purple sulfur

bacteria and heterotrophic sulfate reducing bacteria (SRB)[44]. These existing

methods of confirmatory ecophysiology allowed for direct measurements of the

arrows connecting membership with microbial processes (in this case both carbon

fixation and sulfate reduction, Figure 2, Letter H) in a stable microbial consortium.

These cultivation independent approaches also create the potential to begin to

determine which community properties are emergent properties, and which are

community aggregated traits. For example, microbial community biomass

stoichiometry (e.g., biomass C:N or C:P) cannot currently be predicted (or even

constrained) from a list of its constituent taxa (Figure 2, Letter F). However, microbial

biomass stoichiometry is a community property with power to predict the microbial

contribution to nutrient cycling (Figure 2, Letter G).[17-19] Energy dispersive

spectroscopy (EDS) has the power to measure the C:N:P of individual bacterial cells

growing *in situ* (i.e. not in culture)[43]. The potential to couple EDS analysis with a

phylogenetic label presents the opportunity to assay mixed microbial communities

and assess the link between phylogenetic identity and biomass stoichiometry under

natural conditions[45]. This approach would provide a direct link between community

membership and a community property (e.g., biomass C:N, Figure 2, Letter E), that

influences an important microbial process (i.e., nutrient recycling). These

approaches applied in concert with sequence-based analyses have the potential to

empirically link the categories of microbial information defined here (Figure 2) with

the processes they influence, moving microbiome science from a descriptive and

correlative approach to a mechanistic and causative approach.

Designing microbiome research

The framework presented here provides one approach to formalize inquiry across

microbiome science and encourages empirical linkages between the presence of

organisms in a system and the processes that characterize that system. Whereas we draw examples from environmental microbiomes and the ecosystems they inhabit, this structured approach has the potential to benefit the analysis of microbiomes associated with other systems such as host organisms and those of the built environment. As important as establishing causal links among microbial membership, community properties, microbial processes, and ecosystem processes, is determining when these links are unlikely to be present. Research that indiscriminately seeks to identify correlations, which places all microbial characteristics on an equal plane and does not explicitly recognize the relationships between microbial characteristics, are likely to continue to yield conflicting and ambiguous results that not only fail to provide new insight into ecosystem processes, but also blur the connections that do exist. We suggest that rather than looking for linkages among microbiome membership and system-level processes in every study, research efforts would benefit from strategically targeting the linkages and processes for which an *a priori* understanding of microbial physiology should allow us to improve our understanding of the ecosystem process.

Acknowledgements

This work is a product of the Next Generation of Ecosystem Indicators Working

Group, supported by the USGS John Wesley Powell Center for Synthesis and

Analysis. Preparation of this manuscript was supported by NSF DEB IOS #1456959

awarded to EKH. Chuck Pepe-Ranney and Ariane Peralta provided valuable

feedback on previous versions of this manuscript. This paper is dedicated to Diana

Nemergut, an integral part of our working group who passed away during the

preparation of this manuscript. She is one of a kind.

Figures

Figure 1 Diagram of microbial-ecosystem linkages A) how linkages are commonly conceptualized across levels of ecological organization and B) the series of ecological phenomena that create challenges when attempting to link metrics from one level of ecological organization to the other.

Figure 2 A conceptual map of the intersection between microbial (vertical) and ecosystem (horizontal) ecology illustrating each of the three categories of microbial characteristics (microbial processes, community properties, and microbial membership) as defined in the text. We argue for an increased focus on studies that elucidate pathways E, F, and G. In addition, we note that pathways I and J are less likely to effectively incorporate microbiome characteristics into system-level science. The delta symbol in each category indicates an emphasis on how changes within a category may lead to a change in a connected category. The dotted arrow for letter F denotes that many emergent properties cannot currently be linked to membership and is an important area for active research. CAT = community aggregated trait, EP=emergent property.

References

1.Graham E. B., Wieder, W.R., Leff, J.R., Weintraub, S. W., Townsend, A.R., Cleveland, C.C., Philippot L., Nemergut D.R. (2014) Do we need to understand microbial communities to predict ecosystem function? A comparison of statistical models of nitrogen cycling processes Soil Biology and Biochemistry 68, 279–282

2.Graham E. B. et al. (2016) Microbes as engines of ecosystem function: when does community structure enhance predictions of ecosystem processes? Frontiers in Microbiology 7: 214 doi:10.3389/fmicb.2016.0021

3.Rocca J.D., Hall E.K., Lennon J.T., Evans S.E., Waldrop M.P., Cotner J.B., Nemergut D.R., Graham E.B., Wallenstein M.D. (2015) Relationships between protein-encoding gene abundance and corresponding process are commonly assumed yet rarely observed ISME J. 9: 1693–1699

4.Bier R.L., E.S. Bernhardt, CM Boot, EB Graham, EK Hall, JT Lennon, D Nemergut, B Osborne, C Ruiz-González, JP Schimel, MP Waldrop, MD. Wallenstein (2015) How are we forging conceptual, analytical, and mechanistic links between microbial community structure and ecosystem process? FEMS microbiology ecology FEMS microbiology ecology 91(10): fiv113

5. Falkowski, P.G., Fenchel, T., and DeLong, E.F. (2008) The microbial engines that drive earth's biogeochemical cycles Science 320: 1034-1038

6.Felip M., Pace M.L., Cole, J.J. (1996) Regulation of planktonic bacterial growth rates: The effects of temperature and resources. Microb. Ecol. 31(1): 15-28.

7. Lennon, J. T. and S. E. Jones (2011) Microbial seed banks: the ecological and evolutionary implications of dormancy. Nature Microbiology Reviews 9: 119-130.

8. Comte, J., Fauteux, L. and del Giorgio, P.A. (2013) Links between Metabolic Plasticity and Functional Redundancy in Freshwater Bacterioplankton Communities. Frontiers in Microbiology 4: 112. doi: 10.3389/fmicb.2013.00112

9. Adams, H. E., Crump, B.C. and Kling, G.W. (2014) Metacommunity Dynamics of Bacteria in an Arctic Lake: The Impact of Species Sorting and Mass Effects on Bacterial Production and Biogeography. Frontiers in Microbiology 5 (2014): 82. 10.3389/fmicb.2014.00082

10. Schimel, J.P., J. Bennett, and N. Fierer. (2005) Microbial community composition and soil N cycling: is there really a connection? In: Biological diversity and function in soils. Bardgett, R.D., D.W. Hopkins, and M.B. Usher (Eds.) Cambridge University Press. pp. 171-188

11. Schimel, J.P. (1995) Ecosystem consequences of microbial diversity and community structure in Arctic and Alpine Biodiversity: Patterns, Causes and Ecosystem Consequences Ecological Studies Volume 113, pp 239-254

12. Lenhart K., M. Bunge, S. Ratering, T. R. Neu, I. Schüttmann, M. Greule, C. Kammann, S. Schnell, C. Müller, H. Zorn & F. Keppler. (2012) Evidence for methane production by saprotrophic fungi Nature Comm. 3:1046

13. Hagstroem, A., Azam, F., Berg, C., Zweifel U.L. (2017) Isolates as models to study bacterial ecophysiology and biogeochemistry Aquat. Microb. Ecol. 80:15-27

14. Larkin, A.A, and Martiny, A. (2017) Microdiversity shapes the traits, niche space and biogeography of microbial taxa Environ. Microbiol. Rep. 9(2): 55-70 doi: 10.111/1758-2229

15. German D.P., Marcelo, K.R.B., Stone, M.M and Allison, S.D. (2012) The Michaelis-Menten kinetics of soil extracellular enzymes in response to temperature: a cross-latitudinal study Global Change Biology 18: 1468-1479

16. Hall, E. K., A. R. Dzialowski, S. M. Stoxen, and J. B. Cotner. (2009) The effect of temperature on the coupling between phosphorus and growth in lacustrine bacterioplankton communities. Limnol. and Oceanogr. 54: 880-889.

17. Manzoni, S., R. B. Jackson, J. A. Trofymow, and A. Porporato. (2008) The global stoichiometry of litter nitrogen mineralization. Science 321: 684-686.

18. Elser, J.J., Chrzanowski, T.H., Sterner, R.W., Schampel, J.H., Foster D.K. (1995) Elemental ratios and the uptake and release of nutrients by phytoplankton and bacteria in three lakes of the canadian shield Microbial Ecology 29: 145-162

19. Goldman, J.C., Caron, D.A., and Dennett, M.R. (1987) Regulation of gross growth efficiency and ammonium regeneration in bacteria by substrate C:N ratio Limnology and Oceanography 32 (6): 1239-1252

20. Jensen, B.D., Wise, K.E. and Odegard G.M. 2015 Simulation of the Elastic and Ultimate Tensile Properties of Diamond, Graphene, Carbon Nanotubes, and Amorphous Carbon Using a Revised ReaxFF Parametrization *J. Phys. Chem. A*, *119* (37)**:** 9710–9721

21. Salt, G.W. (1979) A comment on the use of the term emergent properties American Naturalist 113: 145-148

22. Konopka, A. (2009) What is microbial community ecology? ISME J., 3 (11): 1223-12230

23. Battin, T., L. A. Kaplan, L. Newbold, X. Cheng, and C. Hansen. (2003) Effects of current velocity on the nascent architecture of stream microbial biofilms. Appl. and Env. Microbiol. 69: 5443-5452.

24. Martiny JBH, Jones SE, Lennon JT, Martiny AC (2015) Microbiomes in light of traits: a phylogenetic perspective. Science 350: doi: 10.1126/science.aac9323

25. Ruiz-González, C., Niño-García, J. P., Lapierre, J. F., & del Giorgio, P. A. (2015) The quality of organic matter shapes the functional biogeography of bacterioplankton across boreal freshwater ecosystems. Global Ecology and Biogeography *24*(12): 1487-1498.

26. Fierer, N., A. Barberán, Laughlin, D. (2014) Seeing the forest for the genes: Using metagenomics to infer the aggregated traits of microbial communities. Frontiers in Microbiology 5: 614

27. Judd, C.R., Koyama, A., Simmons, M.P., Brewer, P., and von Fischer, J.C. (2016) Co-variation in methanotroph community composition and activity in three temperate grassland soils Soil Biol. and Biochem. 95:78-86

28. Grilli, J., Barabas G., Michalska-Smith, M.J., and Allesina, S. (2017) Higher-order interactions stabilize dynamics in competitive network models Nature. 2017 548: 210–213

29. Newton, R. J, S. E. Jones, A. Eiler, K. D McMahon, Bertilsson, S. (2011) A guide to the natural history of freshwater lake bacteria. Microbiol. and Molec. Biol. Rev. 75:1, doi:10.1128/MMBR.00028

30. Hug, L.A., Baker, B.J., Anantharaman, K., Brown, C.T., Probst, A.J., Castelle, C.J., Butterfield, C.N., Hernsdorf, A.W., Amano, Y., Ise, K., Suzuki, Y., Dudek, N., Relman, D.A., Finstad, K.M., Amundson, R., Thomas, B.C., Banfield, J.F. (2016) 1: 16048 doi: 10.1038/nmicrobiol.2016.48

31. Martiny, J.B.H., B. Bohannan, J. Brown, R. Colwell, J. Fuhrman, J. Green, M.C. Horner-Devine, M. Kane, J. Krumins, C. Kuske, P. Morin, S. Naeem, L. Ovreas, A.-L. Reysenbach, V. Smith, J. Staley. (2006) Microbial biogeography: Putting microorganisms on the map. Nature Reviews Microbiology 4: 102-112

32. Louca, S., Polz, M.F., Mazel, F., Albright, M.B.N., Huber, J. A., O'Connor, M.I., Ackermann, M. Hahn, A.S., Srivastava, D.S., Crowe, S.A., Doebeli, M. and Parfrey, L. W. 2018 Function and functional redundancy in microbial systems https://doi.org/10.1038/s41559-018-0519-1

33. Lennon JT, Aanderud ZA, Lehmkuhl BK, Schoolmaster DR (2012) Mapping the niche space of soil microorganisms using taxonomy and traits. Ecology 93: 1867-1879

34. Treseder KK, Lennon JT (2015) Fungal traits that drive ecosystem dynamics. Microbiology and Molecular Biology Reviews 79: 243-262

35. Czechowska, K., Johnson, D.R., van der Meer, J.R. (2008) Use of flow cytometric methods for single-cell analysis in environmental microbiology Curr. Opin. In Microbiol. 11:3, 205-212

36. Galand P.E., L. Alonso-Sáez, S. Bertilsson, C. Lovejoy, and E. O. Casamayor (2013) Contrasting activity patterns determined by BrdU incorporation in bacterial ribotypes from the Arctic Ocean in winter Front Microbiol. (2013); 4: 118

37. Biller, S.J., Berube, P.M., Lindell, D., Chisholm, S.W. (2015) *Procholorcoccus:* The structure and function of collective diversity Nature Reviews Microbiology 13:13-27

38. Brewer, T.E., Handley, K.M., Carini, P., Gilbert, J.A., Fierer, N. (2016) Genome reduction in an abundant and ubiquitous soil bacterium 'Candidatus Udaeobacter copiosus' Nature Microbiology 2: 16198

39. Neufeld J.D., Vohra J., Dumont M.G., Lueders T., Manefield M., Friedrich M.W., Murrell J.C. (2007) DNA stable-isotope probing. Nat Protoc. 2:4, 860-866

40. Wagner, M. (2009) Single cell ecophysiology of microbes as revealed by Raman microspectroscopy or secondary ion mass spectrometry imaging. Annu Rev Microbiol 63:411-429.

41. Pester, M., Bittner, N., Deevong, P., Wagner, M., and Loy, A. (2010) A 'rare biosphere' microorganism drives sulfate reduction in a peatland. ISME J. 4:1591–1602.

42. Behrens, S., Kappler, A., Obst M. (2012) Linking environmental processes to the in situ functioning of microorganisms by high-resolution secondary ion mass spectrometry (NanoSIMS) and scanning transmission X-ray microscopy (STXM) Env. Microbiol. 14(11): 2851-69

43. Norland S, Fagerbakke K, Heldel M. (1995) Light element analysis of individual bacteria by X-ray microanalysis. Appl. Environ. Microbiol. 61:1357–62.

44. Wilbanks, E.G., Jaekel, U., Salman, V., Humphrey, P.T., Eisen, J.A., Faciotti, M.T., Buckley, D.H., Zinder, S.H., Druschel, G.K., Fike, D.A., Orphan,V.J., (2014) Microscale sulfur cyclingin the photophrophic pink berry consortia of the Sippewissett Salt Marsh Environ. Micro 16: 3398-3415

45. Hall, E.K., Maixner F., Franklin O., Daims, H., Richter, A. and Battin, T. (2011) Linking microbial and ecosystem ecology using ecological stoichiometry: A synthesis of conceptual and empirical approaches Ecosystems 14: 261-273