Prefrontal neurons expand their representation of space by increase in dimensionality and decrease in noise correlation

Mohammad-Reza A Dehaqani [1], Abdol-Hossein Vahabie [1], Mohammadbagher Parsa[2], Behard Noudoost [3], Alireza Soltani [4]

[1] School of Cognitive Sciences, Institute for Research in Fundamental Sciences, Tehran, Iran.

[2] Department of Computer Science, Montana State University, Bozeman MT

[3] Department of Cell Biology and Neuroscience, Montana State University, Bozeman MT

[4] Department of Psychological and Brain Sciences, Dartmouth College, NH 03755

Correspondence: A.S., Department of Psychological and Brain Sciences, HB 6207, Dartmouth College, Hanover, NH, 03755 (Email: soltani@dartmouth.edu) or B.N., Department of Cell Biology and Neuroscience, Montana State University, Bozeman MT, 59717 (Email: bnoudoost@montana.edu)

July 22, 2016

Manuscript information: Total 41 pages (including references, figures and figure captions, and Supplementary figures and methods), 4 figures and 5 Supplementary figures; Abstract 150 words; Introduction 680 words; Results 2490 words; Discussion 700 words

Abstract

Encoding of information by populations of neurons is critical for neural processes, but underlying mechanisms are poorly understood. High-dimensional neural representations and changes in inter-neuronal correlations are two candidates for enhancing population-level information. To examine the contribution of both mechanisms, we applied a combination of decoding and encoding methods to simultaneously recorded neural data. We found that increases in the information content of prefrontal neurons prior to saccadic selection of visual targets were greater for the population of neurons than for individual neurons. This differential increase occurred primarily for regions of space weakly encoded by single-cell responses (encoding expansion), relied on high-dimensional neural representations, and was accompanied by selective reductions in noise correlation. Our results show that population-level information is enhanced both by location-specific increases in dimensionality and decreases in noise correlation, enabling prefrontal ensembles to recruit non-informative individual neurons to encode a larger part of space.

Introduction

The discovery of neurons that are highly selective to external stimuli (Hubel and Wiesel 1959; Hubel and Wiesel 1968; Quiroga et al. 2005) or highly informative about the animal's performance (Newsome, Britten, and Movshon 1989) raised the long-lasting question of how the information content of neural ensembles exceeds that of selective/informative individual neurons. Recently, the combination of improved techniques for large-scale recording from neuronal ensembles and novel computational methods has led to a surge of interest in understanding the contribution of population-level representation to cognitive functions such as working memory, visual attention, decision making, and categorization (Hung et al. 2005; Cohen and Maunsell 2010; Zhang et al. 2011; Rigotti et al. 2013; Mante et al. 2013; Cunningham and Yu 2014). On the one hand, a large body of theoretical work (Shadlen and Newsome 1998; Abbott and Dayan 1999; Panzeri et al. 1999; Nirenberg and Latham 2003; Pola et al. 2003; Averbeck, Latham, and Pouget 2006; Pillow et al. 2008) and electrophysiological studies (Zohary, Shadlen, and Newsome 1994; Cohen and Newsome 2009; Gutnisky and Dragoi 2008; Mitchell, Sundberg, and Reynolds 2009; Cohen and Maunsell 2009; Adibi et al. 2014) have revealed the contribution of inter-neuronal correlations to the coding capacity of neural populations. For example, it has been shown that noise correlation can affect signal-to-noise ratio of the population response (Zohary, Shadlen, and Newsome 1994) and modulate sensory coding more than the firing rate of individual neurons (Cohen and Maunsell 2009; Mitchell, Sundberg, and Reynolds 2009). On the other hand, others have argued that high-dimensional neural representations are more valuable for cognitive functions than representations based on highly specialized neurons (Mante et al. 2013; Rigotti et al. 2013; Mazzucato, Fontanini, and La Camera 2016; Fusi, Miller, and Rigotti 2016).

Although high-dimensional representations and changes in noise correlation are the leading candidates to account for population-level enhancement, the mechanisms by which these factors contribute to the information content of the population have not been examined together. This gap in our knowledge is primarily due to the distinct nature of analyses and methods used to investigate each mechanism. High-dimensional neural representations have been studied with decoding methods such as support vector machines (Tremblay et al. 2015; Hung et al. 2005; Rigotti et al. 2013), whereas changes in noise and signal correlations have been explored using correlation techniques (Cohen and Newsome 2009; Mitchell, Sundberg, and Reynolds 2009; Gutnisky and Dragoi 2008). Studies of these mechanisms have also tended to use different types of neural data (high-dimensional: non-simultaneously recorded neural activity; correlations: simultaneously recorded neural activity) and different experimental paradigms (high-dimensional: working memory, categorization; correlations: attention, adaptation paradigm).

Here we examined the contributions of both high-dimensional neural representations and noise correlation to spatial representation of visual stimuli and saccadic preparation by applying a combination of coding and encoding methods to simultaneously recorded neural activity. The recording was performed in the Frontal Eye Field (FEF), a prefrontal area implicated in controlling eye movements and attention (Bruce and Goldberg 1985; Bruce et al. 1985; Clark et al. 2014), during an oculomotor delayed response task. This data allowed us to study neural representations from visual encoding to saccadic target selection. We found that prior to saccadic selection of visual targets, the information content of the population of neurons increased more than that of individual neurons. More specifically, although individual neurons maintain their limited spatial sensitivity, the ability of the population to encode spatial locations far from the neurons' preferred locations improved prior to saccades, a phenomenon which we refer to as

encoding expansion. In other words, locations poorly encoded by individual neurons (non-preferred locations) could be discriminated much better at the population level. We found that this enhanced population-level information relied on high-dimensional neural representations for non-preferred locations and was accompanied by reductions in noise correlation for the same locations. Overall, we show for the first time that by differently impacting preferred and non-preferred locations of visual space, both high-dimensional representations and changes in the correlated activity contribute to the enhanced information present in the population code.

Results

Using 16-channel linear array electrodes (V-probe, Plexon Inc.), we simultaneously recorded from a population of neurons in the FEF of two monkeys during an oculomotor delayed response (ODR) task (188 single neurons, 1165 pairs of neurons; see Materials and methods). The ODR task (Fig. 1A) enabled us to trace the transformation of visual input into motor action (saccade), and has been classically used to determine the contribution of FEF neurons to each of these processes (Bruce and Goldberg 1985; Bruce et al. 1985; Clark et al. 2014). Our goal was to determine how visual space was encoded differently by a given ensemble of neurons and the participating individual neurons. Therefore, we analyzed the simultaneously recorded neural data to measure spatial information encoded at the single-cell and population levels, as well as information embedded in the correlated activity of pairs of neurons.

Presaccadic expansion of the population code. The average activity of an example recorded neuron during the ODR task revealed the typical spatial tuning/selectivity of FEF neurons (Fig.1B). This neuron responded differently to stimuli presented at eight possible angles during both visual encoding (Enc, one way Anova, F = 18.51, $p < 10^{-3}$) and just before the saccade to

the remembered location (Sac, one way Anova $F = 48.72$, $p < 10^{-3}$). However, the neural

response was more graded during saccade preparation than visual encoding. Similarly, the

average normalized activity across all recorded neurons showed stronger spatial tuning during

saccade preparation compared to visual encoding such that the tuning was narrower for the

former epoch (tuning dispersion: Sac = 2.06±0.04, Enc = 2.25±0.04, $p < 10^{-3}$; see Materials and

methods). We found a similar relationship between the response during saccade preparation and

visual encoding even when we restricted the analysis to the population of neurons that showed

mixed selectivity ($n = 65$, Fig. S1).

We next quantified the information content of individual neurons using the mutual information

between the response and stimulus location (see Materials and methods). We refer to this mutual

information as 'spatial discriminability' because it quantifies the ability of single neuron to

discriminate between various target locations. For the example neuron, the spatial

discriminability was greater during saccade preparation than visual encoding (spatial

discriminability: Enc = 0.09±0.01, Sac = 0.18±0.02, $p < 10^{-3}$). This difference was also observed

across all neurons such that the spatial discriminability was greater during saccade preparation

than visual encoding for the entire population of neurons as well as selective neurons (Enc_{all} =

0.15±0.01, Sac_{all} = 0.23±0.02, $p < 10^{-3}$; $n = 188$, $Enc_{selective}$ = 0.21±0.02, $Sac_{selective}$ = 0.27±0.02,

$p = 0.007$; Fig. 1D). This enhanced representation of saccade targets is consistent with previous

studies (Bruce and Goldberg 1985) and the known role of FEF in oculomotor control.

In addition to mutual information, which reflects the spatial discriminability of single neurons,

we employed a support vector machine (SVM) to quantify the ability of the population of FEF

neurons and individual FEF neurons to represent the target location. The decoding accuracy

measures the ability of an external observer (the classifier) to determine the location of the target

solely based on the simultaneous activity of recorded single and multi-units. The SVM classifier was designed to classify eight target angles either when all the units were considered individually (i.e. separately running the SVM for each unit, $SVM_{individual}$) or when the simultaneous activity of the ensemble of units recorded in one session was taken into account (i.e. running the SVM for each session, $SVM_{ensemble}$; see Materials and methods). The time course of the SVM performance applied to 36 individual units recorded within a single session of the experiment showed similar accuracy during saccade preparation and visual encoding (Fig. 1E). However, when the activity of the population of simultaneously recorded units was taken into account, the decoding accuracy during saccade preparation improved dramatically (Fig. 1F). This improvement was most pronounced for locations in the opposite hemifield. For example, the population decoding accuracy for the 180-degrees away location was about 5 times larger than the decoding accuracy of individual units (decoding accuracy$_{ensemble}$= 0.6±0.09, considering the chance level of 0.125). In contrast, the decoding accuracy for individual units was 0.34±0.18 for the preferred location and dropped to 0.23±0.22 for a location 180 degrees away. Therefore, recordings during this example session exhibited a population-level enhancement of neural encoding during saccade preparation ('presaccadic expansion') which occurred primarily for parts of space weakly encoded by the single-cell response.

The presaccadic expansion of the neural code was consistently observed across all recording sessions. For each unit or session, we determined the preferred location based on the decoding accuracy of the SVM taking the whole visual, memory and presaccadic activity into account. Overall, we found that preferred locations showed almost no presaccadic change in decoding accuracy at the level of individual units (Δdecoding accuracy$_{individual}$= 0.0074±0.0135, p = 0. 0.722, n = 481; Fig.2A). At the same time, nonpreferred locations (>90 degrees away from the

preferred location) exhibited a modest change (Δdecoding accuracy$_{individual}$ = 0.0313±0.0042, p < 10^{-3}). In contrast, when the activity of the population of simultaneously recorded units was taken into account, the decoding accuracy during the presaccadic period improved dramatically for both preferred and nonpreferred locations. This presaccadic enhancement was observed across all eight locations, as indicated by the greater overall decoding accuracy in SVM$_{ensemble}$ during saccade preparation than visual encoding (Δdecoding accuracy$_{ensemble}$ = 0.09±0.03, p = 0.008; Fig. 2A). Importantly, this presaccadic population benefit was not equally strong across all target locations: the benefit was small for the neurons' preferred location (Δdecoding accuracy$_{pref}$ = 0.02±0.03; p = 0.711), and much larger for locations far from the preferred locations (Δdecoding accuracy$_{nonpref}$ = 0.12±0.03, p = 0.008; Fig. 2B).

Overall, these results reveal an expansion of the population code prior to saccadic target selection. Such an expansion may increase the robustness or precision of the population code by incorporating a larger ensemble of FEF neurons. Note that this presaccadic enhancement was mainly observed at the population level and not the individual units (Fig. 2B), implying that the expansion of the neural code relies on high-dimensional neural representations and also depends on changes in the inter-neuronal relationships. Thus, we next examined the contributions of these factors to presaccadic expansion.

Presaccadic expansion relies on high-dimensional neural representations. We found that during saccade preparation, the ability of the population to represent target locations expanded far beyond the capability of single neurons: locations which were poorly encoded by single neurons could be accurately discriminated using an ensemble-level population read-out. Recent studies have indicated that high-dimensional neural representations are beneficial for information encoding (Mante et al. 2013; Rigotti et al. 2013). Therefore, we next examined whether the

encoding expansion during saccade preparation was associated with a greater dimensionality in the population.

To do so, we sorted the simultaneously recorded units in each session based on their mutual information (spatial discriminability), and made a set of neural ensembles by subsequently adding new units (with decreasing mutual information). Using this approach, we first computed the overall decoding accuracy as a function of the number of units (NU) in different epochs of the experiment (Fig. S2). We found that about 4 units are enough to explain 75% of the ultimate classifier performance during both saccade preparation and visual encoding ($NU_{Enc} = 4.44 \pm 0.84$, $NU_{Sac} = 3.82 \pm 0.67$, errors based on bootstrapping; see Materials and methods).

The encoding expansion during saccade preparation occurred because the SVM performance increased more strongly for nonpreferred than preferred locations. Therefore, we next computed the dimensionality of neural representation separately for preferred and nonpreferred (>90 deg) locations (Fig. 2C). We found that 6.54 ± 2.21 units were necessary to achieve 75% of the whole-ensemble level of performance for nonpreferred locations, compared to only 2.22 ± 0.45 units for preferred locations, indicating that the population response had a much larger dimensionality for representing nonpreferred compared to preferred locations (bootstrap confidence interval, $p < 10^{-3}$; Fig. 2C). Note that the exact value of the dimensionality strongly depends on the threshold (here 75%) such that a larger dimensionality is obtained with a larger value of the threshold. Nevertheless, our result indicates that about three times more neurons are needed to encode nonpreferred locations than preferred locations.

We next examined the direct relationship between encoding expansion and the difference in dimensionality between nonpreferred and preferred locations by defining an expansion index for

each session of the experiment (see Materials and methods). There was a positive correlation

between the encoding expansion index and the difference in dimensionality between

nonpreferred and preferred locations (Spearman correlation, $r = 0.62$, $p = 0.014$; Fig. 2D). This

correlation demonstrates that a larger difference in dimensionality between nonpreferred and

preferred locations is accompanied by larger increase in the population-level, differential

encoding of nonpreferred than preferred locations during saccade preparation (i.e. presaccadic

expansion). Therefore, these results suggest that high-dimensional neural representations are

crucial for the observed presaccadic expansion.

Changes in inter-neuronal correlations contribute to encoding expansion during saccade

preparation. To address the contribution of correlated neural activity to the enhanced spatial

representation prior to saccades, we examined the effects of inter-neuronal correlations at the

level of pairs of neurons (see Materials and methods). The correlated activity of a pair of neurons

could be decomposed into two components: 1) signal correlation (SC), which in this task reflects

the degree to which two neurons exhibit similar response to a given target; and 2) noise

correlation (NC), measuring shared variability unrelated to the target location (Cohen and Kohn

2011). By transforming the problem of population level analysis to the level of pairs of neurons

and stimuli (~ 32400 neuron-stimulus pairs), we were able to precisely study the contribution of

noise and signal correlations to encoding expansion during saccade preparation.

We found a higher SC during saccade preparation than visual encoding ($SC_{Enc} = 0.0164 \pm 0.0005$,

$SC_{Sac} = 0.0447 \pm 0.0008$, $p < 10^{-3}$). This greater correlation in the presaccadic spatial tuning of

FEF neurons was consistent with the enhanced response (Fig. 1C) and spatial discriminability

(Fig. 2A) observed during this epoch within the same population. In contrast to SC, the NC was

reduced in the presaccadic period ($NC_{Enc} = 0.046 \pm 0.0007$, $NC_{Sac} = 0.039 \pm 0.0007$, $p < 10^{-3}$; Fig.

3A). Interestingly, the presaccadic reduction in NC was prominent for pairs of nonpreferred locations (>90 deg) compared to preferred locations ($\Delta NC_{nonpref} = -0.0152\pm0.0029$, $\Delta NC_{pref} = -0.0007\pm0.0025$, $p < 10^{-3}$; Fig. 3A), however, this differential change was not observed for increases in signal correlation ($\Delta SC_{pref} = 0.0139\pm0.0021$, $\Delta SC_{nonpref} = 0.0104\pm0.0012$, $p = 0.376$). The differential decreases in noise correlation (from visual encoding to saccade preparation) for preferred and nonpreferred locations dovetail with the presaccadic expansion of population code measured by the SVM.

Because signal and noise correlations are defined based on the activity of pairs of neurons and not the whole populations, computing their contribution to the observed encoding expansion of the population code requires measuring the information content of pairs of neurons (as a proxy for the population). Therefore, we next measured the information content of a given pair of neurons by the ability of this pair to discriminate a pair of locations ('joint' discriminability) using the linear discriminant analysis (LDA) technique (see Materials and methods and Figure S3A). We found that joint discriminability was enhanced during saccade preparation compared to visual encoding (Enc = 0.1572 ± 0.0009, Sac = 0.2274 ± 0.0013, $p < 10^{-3}$). Moreover, like the SVM results (Fig. 2A), nonpreferred locations showed greater presaccadic improvements in discriminability than preferred locations (Δjoint discriminability$_{nofpref} = 0.0281\pm0.0027$, Δjoint discriminability$_{pref} = 0.0105\pm0.0031$, $p < 10^{-3}$; Fig. 3B).

Therefore, we next tested whether the differential improvement in joint discriminability for nonpreferred than preferred locations reflects the encoding expansion measured by the SVM. To do so we computed the correlation between the encoding expansion index and a similar index based on joint discriminability (after subtracting a baseline related to the information content of individual cells; see Materials and methods). A significant correlation between encoding

expansion index based on the SVM and joint discriminability demonstrated that joint discriminability reflects encoding expansion (Spearman correlation, $r = 0.53$ $p = 0.025$). Overall, the analyses reported above based on pairwise measures confirmed the expanded presaccadic spatial encoding, which was originally revealed by the decoding analysis.

Unlike encoding expansion based on the SVM, joint discriminability measures the information content of a pair of neurons on a trial-by-trial basis and therefore, can be used to look for behavioral correlates of changes in the information content between visual encoding and saccade preparation. By binning the change in joint discriminability from visual encoding to saccade preparation, we found that greater changes in joint discriminability were associated with faster reaction times (Fig. 3C). More specifically, there was ~5% difference between the average RT when the joint discriminability was negative and positive (235.32 ± 0.39 vs. 224.60 ± 0.68 msec, $p < 10^{-3}$), showing that changes in joint discriminability had a significant impact on the behavior.

Having established that changes in joint discriminability reflect encoding expansion, we next examined the relationship between signal correlation, noise correlation, and joint discriminability for the same pairs of neurons during visual encoding and saccade preparation. This analysis allowed us to study the contribution of signal and noise correlations to encoding expansion. To do so we first partitioned our dataset according to the sign of signal and noise correlations in each pair. Interestingly, we found that indeed when two neurons had a common source of signal (a positive SC), they were three times more likely to have a common source of noise (a positive NC; Fig. S4). Moreover, pairs with a positive signal and noise correlations comprised about 45% of the dataset. Importantly, presaccadic expansion based on joint discriminability was at least two times greater in these pairs than in those pairs with one or more negative correlation values (Fig. S4).

By examining the relationship between signal correlation and joint discriminability during visual encoding for pairs of neuron with positive signal and noise correlations, we found that a higher signal correlation was associated with greater discriminability (Fig. 4A). However, by separating the population based on whether the noise correlation decreased or increased during the presaccadic period (comprising 76% and 24% of population, respectively), we found that for a given level of SC the joint discriminability depended upon the change in noise correlation such that a decrease in noise correlation improved joint discriminability. This result indicated that a change in noise correlation was sufficient to improve encoding expansion during saccade preparation (also see Figure S5A).

As a further control, we verified that the isolated discriminability (which excludes the contribution of NC yet incorporates each unit's spatial sensitivity) stayed the same between neurons for which NC decreased during saccadic preparation and those for which NC increased (Fig. S5B). Finally, we examined the presaccadic change in joint discriminability as a function of changes in SC and NC and found that the change in discriminability depended on the change in NC, independent of the change in SC (Fig. 4B). Overall, these results indicated that decorrelation (i.e. a reduction in noise correlation) contributed to the presaccadic encoding expansion.

Discussion

The brain's processing capacity is limited by physical constraints including intrinsic noise in neural firing, the number of neurons, and their connectivity. Therefore, changes in neural encoding which improve the ability of a fixed population of neurons to represent information are of particular interest to neuroscientists, because such changes enable the brain to dynamically allocate limited neural resources based on task demands and to maximize neural computations

(Mansouri 2006; Asaad, Rainer, and Miller 1998). Here, we examined neural representations of saccade targets in the FEF and found improved spatial encoding during saccade preparation, especially at the population level. When moving from sensory encoding to saccade preparation, our results also showed an enhanced population representation of parts of space. This novel phenomenon, which we refer to as the presaccadic encoding expansion, was not present in the single-cell responses and relied on high-dimensional neural representations.

Our results dovetail with recent findings showing the importance of high-dimensional neural representations for different cognitive functions and demonstrate that neural populations can encode information not evident in the firing rate of individual neurons (Hung et al. 2005; Cohen and Maunsell 2010; Zhang et al. 2011; Rigotti et al. 2013). Interestingly, we found increased dimensionality for encoding the nonpreferred than the preferred target locations, corresponding with the larger population-level encoding benefit for nonpreferred locations. This boost indicates that a high-dimensional representation is particularly useful for improving population encoding when signal improvement cannot occur through single-cell mechanisms (e.g. gain modulation), which is compatible with a recent finding that reduced dimensionality accompanied poorer performance (Rigotti et al. 2013). In addition, we found that encoding expansion was accompanied by reduction in noise correlation for nonpreferred locations only, indicating the importance of inter-neuronal interactions between neurons selective to different locations.

Here we used two very different measures (decoding accuracy based on the SVM and joint discriminability based on the LDA) to assess the information content of the population response. Importantly, both measures captured changes in the representation of targets between stimulus encoding and saccade preparation. However, only joint discriminability allowed measurement of the information content of pairs of neurons for which signal and noise correlations could be

computed as well. Numerous studies have linked reductions in noise correlation to enhanced cognitive functions such as attention (Cohen and Maunsell 2009; Mitchell, Sundberg, and Reynolds 2009; Tremblay et al. 2015). Here, we used the LDA to isolate the effect of changes in inter-neuronal correlations and found that a presaccadic reduction in noise correlation enhances joint discriminability. Results based on both the SVM and LDA reveal a presaccadic expansion of the population code, with population-level benefits at the nonpreferred locations attributable both to the dimensionality of the population representation and changes in noise correlation between neurons. Therefore, we were able to quantify the contributions of both high-dimensional neural representation and reduction in noise correlations within the same study.

Interestingly, a recent study has suggested that the amount of pair-wise correlations could put an upper bound on the dimensionality of neural ensembles (Mazzucato, Fontanini, and La Camera 2016). Another study measured the effect of noise correlation on population encoding in prefrontal cortex and found that removing noise correlation increased decoding accuracy for the cue and attention location but not for the saccade (Tremblay et al. 2015). Together, these results suggest that reduction in noise correlation and increase in dimensionality are complementary processes involved in enhancing information.

We find that during saccade preparation, the FEF's representation of space undergoes expansion at the level of the population code, showing another example of how a population of neuron can be more informative than individual neurons. A recent study has shown that the FEF neurons are able to encode targets of eye movements far from their classic receptive field (Zirnsak et al. 2014). Our results illustrate how population-level changes can further expand the neural code beyond what individual neural can achieve. This encoding expansion relies on both high-dimensional neural representations and reductions in noise correlation between neurons. Overall,

our results demonstrate the ability of prefrontal neural ensembles to recruit non-informative individual neurons to encode a larger part of space, and to exploit high-dimensional neural representations and changes in inter-neuronal relationships for improving the encoding capacity of the whole population.

Material and methods

Experimental paradigm and recording. Two monkeys (macaca mulatta) were trained to perform an oculomotor delayed response (ODR) task. Each trial started with the monkey fixating on the fixation point, followed by the presentation of a visual target (black circle) in one of 16 locations defined with two possible eccentricities (7 and 14 visual degrees) and eight different angles (0, 45, 90, 135, 180, 225, 270, and 315 degrees). The target remained on screen for one second, followed by 1 second of delay after which the monkey could saccade to the remembered location. The monkey was rewarded with a drop of juice if the saccade fell within 5 degrees of visual angle from the target location. The neural recording was done using a V-probe linear array electrode, allowing us to simultaneously record from 16 channels. Using a strict criterion for spike sorting, we identified 188 single-unit (1165 pairs of neurons) and 320 clusters of multi-unit activity (total 508) during the 20 recording sessions (with an average of 784 trials per session). All experimental procedures were in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals, the Society for Neuroscience Guidelines and Policies, and the Montana State Animal Care and Use Committee.

Data analysis. The baseline mutual information (MI) was computed using neural firing rates during non-overlapping 150 msec windows during the fixation interval for each pair of locations.

$$MI = -\sum_{r} P[r] \log_2 P[r] + \sum_{s,r} P[s] P[r \mid s] \log_2 P[r \mid s] \quad \text{(Eq.1)}$$

where *s* is the set of stimuli, *r* is the set of neural response (mean firing rate), *P(r|s)* is the conditional probability of neural response *r* given stimulus *s*, and *p(s)* is a prior probability of stimulus *s*. We refer to this mutual information as 'spatial discriminability' because it quantifies the ability of single neuron to discriminate between various target locations.

Neural selectivity across different epochs was then quantified by comparing the MI for each pair of locations during sensory encoding (50-200 msec time window aligned to the target onset), memory (1500-1650 msec time window aligned to the target onset), and saccade preparation (-120-30 msec time window aligned to the saccade onset), to the baseline MI (using two-sided Wilcoxon ranksum and *p* < .05 as the significant test).

To extract decoding information, we trained a SVM classifier with a linear kernel using the population and single neural responses to 70% of the stimuli. Decoding accuracy was computed by the performance of the classifier on the remaining 30% of the stimuli. We repeated the calculations 500 times and in each repetition, the stimuli were randomly partitioned into training and test sets. In order to compute the dimensionality of neural representation, we first sorted the simultaneously recorded units in each session based on their mutual information (spatial discriminability) and subsequently constructed a set of neural ensembles by subsequently adding new units (with decreasing mutual information). We next computed the overall decoding accuracy as a function of the number of units (NU) in different epochs of the experiment. The dimensionality was defined as the number of units needed to exceed 75% of maximum decoding accuracy. To measure the standard error of the dimensionality, we used a bootstrapping method

that sampled the session's ensembles with replacement (1000 times). More details are provided in the Supplementary methods.

The preferred location is defined based on the maximum of the average response (neural tuning curve) or the average decoding accuracy (decoding tuning curves) of locations in visual encoding, memory, and saccade preparation intervals for individual units and the ensemble. For the dimensionality analysis we used an average of the three intervals across all dimensions. We computed the tuning dispersion of spatial tuning by fitting a Gaussian probability density function and using the standard deviation of the fitted function.

Moreover, in order to estimate the information content of a pair of neurons, we applied Fisher linear discriminant analysis (LDA) on each pair of neurons (2-dimensional neural space) and extracted the optimal weight vector w_{opt} as:

$$w_{opt} = C^{-1}R \qquad \text{(Eq.2)}$$

where R is the difference between the mean vectors of a given two conditions, and C^{-1} is the inverse of the summation of the covariance matrixes of two conditions in the 2-dimensional neural space constructed by a pair of neurons. The joint discriminability was computed by a projection of the 2 locations represented in the 2-dimensional neural space on to the weight vector. The joint discriminability is the mutual information between the projections of the neural response and target location. The geometric mean of MI values in single cells of a given pair was considered as the single-cell version of the joint discriminability, which we refer to as 'isolated' discriminability and was used as a control for measuring encoding expansion (see below).

Different encoding expansion indices were computed using the performance based on the SVM or discriminability as follows:

$$(X_{nonpref}^{Sac} - X_{nonpref}^{Enc}) - (X_{pref}^{Sac} - X_{pref}^{Enc}) \qquad \text{(Eq.3)}$$

where X can be $SVM_{individual}$, $SVM_{ensemble}$, $discriminability_{isolated}$, or $discriminability_{joint}$. To compute the final measure the encoding expansion index using the SVM analysis or the LDA, we subtracted single-unit expansion index ($X = SVM_{individual}$ or $discriminability_{isolated}$) from the corresponding ensemble expansion index ($X = SVM_{ensemble}$ or $discriminability_{joint}$).

Statistical analysis were performed using the Wilcoxon's signed-rank test (for paired comparisons) or ranksum (for unpaired comparisons), unless otherwise specified. All *p* values are reported up to three digits and values below 0.001 are all reported as *p < 10^{-3}*.

Acknowledgments

We thank Dr. Kelsey Clark for her helpful comments and edits on the manuscript. The work was supported by MSU startup fund, Whitehall 2014-5-18, NSF BCS143221, and NEI R01EY026924 to BN and Dartmouth startup fund to AS.

References

Abbott LF, and P Dayan. 1999. "The Effect of Correlated Variability on the Accuracy of a Population Code." *Neural Computation* 11 (1):1–101. doi:10.1162/089976699300016827.

Adibi M, McDonald JS, Clifford CWG, and E Arabzadeh. 2014. "Population Decoding in Rat Barrel Cortex: Optimizing the Linear Readout of Correlated Population Responses." *PLoS Computational Biology* 10 (1): e1003415. doi:10.1371/journal.pcbi.1003415.

Asaad WF, Rainer G, and EK Miller. 1998. "Neural Activity in the Primate Prefrontal Cortex during Associative Learning." *Neuron* 21 (6): 1399–1407.

Asaad WF, Santhanam N, McClellan S, and DJ Freedman. 2013. "High-Performance Execution of Psychophysical Tasks with Complex Visual Stimuli in MATLAB." *Journal of Neurophysiology* 109 (1): 249–60.

Averbeck BB, Latham PE, and A Pouget. 2006. "Neural Correlations, Population Coding and Computation." *Nature Rev Neuroscience* 7 (5): 358–66. doi:10.1038/nrn1888.

Bruce CJ, and ME Goldberg. 1985. "Primate Frontal Eye Fields. I. Single Neurons Discharging before Saccades." *Journal of Neurophysiology* 53 (3): 603–35.

Bruce CJ, Goldberg ME, Bushnell MC, and GB Stanton. 1985. "Primate Frontal Eye Fields. II. Physiological and Anatomical Correlates of Electrically Evoked Eye Movements." *Journal of Neurophysiology* 54 (3): 714–34.

Clark KL, Noudoost B, Schafer RJ, and T Moore. 2014. "Neuronal Mechanisms of Attentional Control: Frontal Cortex." *The Oxford Handbook of Attention*, no. August. Oxford University Press Oxford: 1–18. doi:10.1093/oxfordhb/9780199675111.013.010.

Cohen MR, and JHR Maunsell. 2010. "A Neuronal Population Measure of Attention Predicts Behavioral Performance on Individual Trials." *Journal of Neuroscience* 30 (45): 15241–53.

doi:10.1523/JNEUROSCI.2171-10.2010.

Cohen MR, and A Kohn. 2011. "Measuring and Interpreting Neuronal Correlations." *Nature Neuroscience* 14 (7): 811–19. doi:10.1038/nn.2842.

Cohen MR, and JHR Maunsell. 2009. "Attention Improves Performance Primarily by Reducing Interneuronal Correlations." *Nature Neuroscience* 12 (12): 1594–1600. doi:10.1038/nn.2439.

Cohen MR, and WT Newsome. 2009. "Contest-Dependent Changes in Functional Circuitry in MT." *Neuron* 60 (1): 162–73. doi:10.1016/j.neuron.2008.08.007.

Cortes C, and V Vapnik. 1995. "Support-Vector Networks." *Machine Learning* 20 (3): 273–97. doi:10.1007/BF00994018.

Cunningham JP, and BM Yu. 2014. "Dimensionality Reduction for Large-Scale Neural Recordings." *Nature Neuroscience* 17 (11): 1500–1509. doi:10.1038/nn.3776.

Duda RO, Hart PE, and DG Stork. 2012. *Pattern Classification*. John Wiley & Sons.

Fusi S, Miller EK, and M Rigotti. 2016. "Why Neurons Mix: High Dimensionality for Higher Cognition." *Current Opinion in Neurobiology* 37: 66–74. doi:10.1016/j.conb.2016.01.010.

Gutnisky DA, and V Dragoi. 2008. "Adaptive Coding of Visual Information in Neural Populations." *Nature* 452 (7184): 220–24. doi:10.1038/nature06563.

Hubel DH, and TN Wiesel. 1959. "Receptive Fields of Single Neurones in the Cat's Striate Cortex." *Journal of Physiology* 148 (3): 574–91. doi:10.1113/jphysiol.2009.174151.

Hubel DH, and TN Wiesel. 1968. "Receptive Fields and Functional Architecture of Monkey Striate Cortex." *Journal of Physiology (London)* 195 (1): 215–43.

Hung CP, Kreiman G, Poggio P, and JJ DiCarlo. 2005. "Fast Readout of Object Identify from Macaque Inferior Temporal Cortex." *Science* 310 (5749): 863-6.

doi:10.1126/science.1117593.

Mansouri FA. 2006. "Prefrontal Cell Activities Related to Monkeys' Success and Failure in Adapting to Rule Changes in a Wisconsin Card Sorting Test Analog." *Journal of Neuroscience* 26 (10): 2745–56. doi:10.1523/JNEUROSCI.5238-05.2006.

Mante V, Sussillo D, Shenoy KV, and WT Newsome. 2013. "Context-Dependent Computation by Recurrent Dynamics in Prefrontal Cortex." *Nature* 503 (7474): 78–84. doi:10.1038/nature12742.

Mazzucato L, Fontanini A, and G La Camera. 2016. "Stimuli Reduce the Dimensionality of Cortical Activity." *Frontiers in Systems Neuroscience* 10 (11). doi: 10.3389/fnsys.2016.00011

Mitchell JF, Sundberg KA, and JH Reynolds. 2009. "Spatial Attention Decorrelates Intrinsic Activity Fluctuations in Macaque Area V4." *Neuron* 63 (6)879–88. doi:10.1016/j.neuron.2009.09.013.

Newsome WT, Britten KH, and JA Movshon. 1989. "Neuronal Correlates of a Perceptual Decision." *Nature* 342: 52–54. :doi:10.1038/341052a0.

Nirenberg S, and PE Latham. 2003. "Decoding Neuronal Spike Trains: How Important Are Correlations?" *Proceedings of the National Academy of Sciences of the United States of America* 100 (12): 7348–53. doi:10.1073/pnas.1131895100.

Pachitariu M, Lyamzin DR, Sahani M, and NA Lesica. 2015. "State-Dependent Population Coding in Primary Auditory Cortex." *Journal of Neuroscience* 35 (5): 2058–73. doi:10.1523/JNEUROSCI.3318-14.2015.

Panzeri S, Schultz SR, Treves A, and ET Rolls. 1999. "Correlations and the Encoding of Information in the Nervous System." *Proceedings of the Royal Society of London B:*

Biological Sciences 266 (1423): 1001–12. doi:10.1098/rspb.1999.0736.

Pillow JW, Shlens J, Paninski L, Sher A, Litke AM, Chichilnisky EJ, and EP Simoncelli. 2008. "Spatio-Temporal Correlations and Visual Signalling in a Complete Neuronal Population." *Nature* 454 (7207): 995–99. doi:10.1038/nature07140.

Pola G, Thiele A, Hoffmann KP, and S Panzeri. 2003. "An Exact Method to Quantify the Information Transmitted by Different Mechanisms of Correlational Coding." *Network-Computation in Neural Systems* 14 (1): 35–60. doi:10.1088/0954-898X/14/1/303.

Quiroga RQ, Reddy L, Kreiman G, Koch C, and I Fried. 2005. "Invariant Visual Representation by Single Neurons in the Human Brain." *Nature* 435 (7045): 1102–7. doi:10.1038/nature03687.

Rigotti M, Barak O, Warden MR, Wang XJ, Daw ND, Miller EK, and S Fusi. 2013. "The Importance of Mixed Selectivity in Complex Cognitive Tasks." *Nature* 497 (7451): 1–6. doi:10.1038/nature12160.

Shadlen MN, and WT Newsome. 1998. "The Variable Discharge of Cortical Neurons: Implications for Connectivity, Computation, and Information Coding." *The Journal of Neuroscience* 18 (10): 3870–96. doi:0270-6474/98/183870-27$05.00/0.

Shannon CE. 1948. "A Mathematical Theory of Communication." *The Bell System Technical Journal* 27: 379–423, 623–56. doi:10.1145/584091.584093.

Shoham S, Fellows MR, and RA Normann. 2003. "Robust, Automatic Spike Sorting Using Mixtures of Multivariate T-Distributions." *Journal of Neuroscience Methods* 127 (2): 111–22. doi:10.1016/S0165-0270(03)00120-1.

Suykens JAK, and J Vandewalle. 1999. "Least Squares Support Vector Machine Classifiers." *Neural Processing Letters* 9 (3): 293–300.

Tremblay S, Pieper F, Sachs A, and J Martinez-Trujillo. 2015. "Attentional Filtering of Visual Information by Neuronal Ensembles in the Primate Lateral Prefrontal Cortex." *Neuron* 85 (1): 202–15. doi: 10.1016/j.neuron.2014.11.021

Zhang Y, Meyers EM, Bichot NP, Serre S, Poggio T, and R Desimone. 2011. "Object Decoding with Attention in Inferior Temporal Cortex." *Proceedings of the National Academy of Sciences of the United States of America* 108(21):8850-5. doi: 10.1073/pnas.1100999108.

Zirnsak M, Steinmetz NA, Noudoost B, Xu KZ, and Moore T. 2014. "Visual Space Is Compressed in Prefrontal Cortex before Eye Movements." *Nature* 507 (7493): 504–7. doi:10.1038/nature13149.

Zohary E, Shadlen MN, and WT Newsome. 1994. "Correlated Neuronal Discharge Rate and Its Implications for Psychophysical Performance." *Nature* 370 (6485): 140–43. doi:10.1038/370140a0.

Figures and figure legends

Figure 1. Spatial encoding of a population of FEF neurons expands beyond the ability of individual neurons prior to saccade. **(A)** Behavioral paradigm. Each trial of the oculomotor delayed response task started with fixation on a central spot for 1000 msec followed by the presentation of a visual target (black circle) for 1000 msec in one of the 16 possible locations (at 8 angles and 2 eccentricities). Disappearance of the target was followed by a 1000 msec delay, after which the central spot disappeared and the monkey made a saccade to the remembered target location. **(B)** The mean response of an example FEF neuron to the target presented at 8

different angles (averaged over two eccentricities), indicated by color in the inset. The neural response is aligned to the onset of the target (left panels) or saccade (right panel). (**C**) Normalized response of 188 FEF neurons to targets presented at different angles relative to the neuron's preferred location during visual encoding (Enc, green) and saccade preparation (Sac, blue) epochs. The error bars represent s.e.m. There was only a slight improvement in spatial tuning of individual neurons from visual encoding to saccade preparation. (**D**) Comparison of the spatial discriminability during visual encoding and saccade preparation. FEF neurons exhibiting mixed selectivity are highlighted in red (n=65). Histogram on the diagonal shows the difference in spatial discriminability. (**E-F**) Plotted is the decoding accuracy for different target locations based on the SVM classifier, when the activity of individual units (E) or the ensemble of units (F) recorded in an example session is used. The time course of decoding accuracy showed similar accuracy during saccade preparation and visual encoding for individual units (E) but larger accuracy by the ensemble of neurons during saccade preparation than visual encoding for nonpreferred locations. The left (right) panels shows the results when the activity is aligned to target (saccade) onset.

Figure 2. Presaccadic expansion relies on high-dimensional neural representation of space. (**A**) Plotted is the decoding accuracy (Dec. accuracy) of the SVM classifier for various locations (relative to preferred) for individual (circles) and ensemble (stars) of units during visual encoding (Enc) and saccade preparation (Sac). There was a remarkable difference for decoding accuracy of the ensemble of units for nonpreferred locations. (**B**) The change in decoding accuracy between visual encoding and saccade preparation for individual units and the ensemble of units as a function of target locations. Larger improvements are visible for nonpreferred locations. (**C**) The decoding accuracy of the ensemble of units plotted as a function of the number of units separately for the preferred and nonpreferred target locations during saccade preparation. (**D**) The difference in dimensionality for preferred and nonpreferred locations predicted encoding expansion. Plotted is the encoding expansion index during a session as a function of the difference in dimensionality between nonpreferred and preferred locations during Sac. The dashed line illustrates least-squares line fitted to points and the error bars represent s.e.m.

Figure 3. Noise correlation and information content of pairs of neurons (joint discriminability) differentially change for preferred and nonpreferred locations, and the latter modulates behavior. (**A**) Average noise correlation across all locations and separately for preferred and non-preferred locations during visual encoding (Enc, green) and saccade preparation (Sac, blue). There was an overall reduction in noise correlation during Sac, but this effect was driven by the reduction in noise correlation at nonpreferred locations. (**B**) Plotted is the joint discriminability separately measured for preferred and nonpreferred locations during visual encoding and saccade preparation. Nonpreferred locations showed a larger improvement in joint discriminability from visual encoding to saccade preparation. (**C**) Faster reaction times corresponded with greater improvement in joint discriminability from visual encoding to saccade preparation. The error bars represent s.e.m.

Figure 4. Changes in noise correlation contributes to changes in joint discriminability and therefore, presaccadic expansion, however, this effect is revealed by controlling for the change in signal correlation. (**A**) Joint discriminability depended upon both signal and noise correlation. Plotted is joint discriminability as a function of signal correlation computed separately for neuron-location pairs during visual encoding (Enc, green) and during saccade preparation. The latter is plotted for pairs of neurons that showed a large (> 0.1) reduction (blue) or increase (red) in noise correlation from visual encoding to saccade preparation. The shaded area represents s.e.m. For a given value of signal correlation, joint discriminability of neurons that became decorrelated was larger than those which became more correlated during saccade preparation. (**B**) Change in joint discriminability was explained by a reduction in noise correlation when the change in signal correlation was taken into account. The change in joint discriminability (separately z-scored across the same change of signal correlation, each row) is plotted as a function of the changes in signal and noise correlation.

Supplementary figures and figure legends

Figure S1. Normalized response of FEF neurons with mixed selectivity to targets presented at different angles relative to their preferred location (0 degree) during visual encoding (Enc, green) and saccade preparation (Sac, blue). The error bars represent s.e.m. Overall, the response profile of neurons with mixed selectivity was very similar to the average of all recorded neurons.

Figure S2. Dimensionality analysis based on the decoding accuracy of the SVM classifier. Plotted is the decoding accuracy of the ensemble of neurons as a function of the number of neurons used for the SVM classifier, separately for visual encoding (Enc, green) and saccade preparation (Sac, blue) epochs. The error bars represent s.e.m.

Figure S3. (A) Schematic of the linear discriminant analysis (LDA) used for computing joint discriminability. Ellipses represent the variability of neural population response in two conditions (green and blue). The LDA weight vector for the pair of neurons (solid line in the left panel) is the optimum weight vector for population decoding, and is equal to a linear combination of the two neurons that maximally separates neural responses in the two conditions. The dashed line corresponds to the best criterion to discriminate responses to the two conditions (i.e. the LDA line). **(B)** Schematic of the computations of signal and noise correlations. The signal correlation was computed after shuffling the order of repeated trials within each condition (right panel). The noise correlation was obtained by subtracting the signal correlation from the total correlation (i.e. correlation coefficient without shuffling). **(C)** Effects of changes in inter-neuronal correlations on the discriminability of paired neural responses. Plotted is an idealized response of two neurons with positive signal and noise correlations to the target at two different locations (green and blue), and ellipses represent the variability of the neural response. A decrease in the noise correlation between the two neurons (decorrelated case) produces a reduction in the overlap of the distributions of the neural responses to the two stimuli, corresponding to improved discrimination between two locations. In contrast, an increase in correlation between the two neurons (correlated case) creates more overlap between the neural responses to the two stimuli, indicating a drop in discriminability.

Figure S4. Neuron pairs with positive signal and noise correlations comprised about half of the population, and exhibited a change in joint discriminability twice other pairs of neurons. Plotted is the change in joint discriminability from visual encoding to saccade preparation for various groups of pairs, sorted based on the sign of the signal and noise correlations for the neuron-condition pairs. The number of neuron-condition pairs in each group is shown above each bar.

Figure S5. Although the change in the information content of pairs of neurons depends on the change in noise correlation, the information content of individual neurons does not depend on change in noise correlation between visual encoding and saccade preparation. (**A**) Plotted is the change in joint discriminability for pairs of neurons that do not show a negligible change in SC between visual encoding and saccade preparation (ΔSC < 0.05, *n* = 6419), illustrating the direct contribution of NC changes to joint discriminability. Improved joint discriminability during saccade preparation compared to visual encoding is most prominent when NC is reduced in saccade preparation relative to visual encoding in the same pair. (**B**) Plotted is isolated discriminability as a function of signal correlation computed separately for neuron-location pairs during visual encoding (Enc, green) and during saccade preparation. The latter is plotted for pairs of neurons that showed a large (> 0.1) reduction (blue) or increase (red) in noise correlation from visual encoding to saccade preparation. The shaded area represents s.e.m. This figure provides a control of Figure 4A, and illustrates that the information content of individual neurons in the same pairs of neurons is the same for visual encoding and saccade preparation, and for pairs which become more correlated and decorrelated during saccade preparation.

Supplementary materials and methods

General and surgical procedures. two male rhesus monkeys (Maccaca mulatta) were used in this experiment. All experimental procedures were in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals, the Society for Neuroscience Guidelines and Policies, and the Montana State Animal Care and Use Committee. Each animal was surgically implanted with a head post and recording chamber. Surgery was conducted using aseptic techniques under general anesthesia (isoflurane), and analgesics were provided during postsurgical recovery. Eye position monitoring was performed via optical tracking with a high frequency camera (the eyeLink system; 2000 Hz). Eye monitoring, stimulus presentation, data acquisition, and behavioral monitoring were controlled by the Monkey Logic system (Asaad et al. 2013). Visual targets (black circles with 1 visual angle radius) presented in one of 16 locations defined with 2 possible radii (7 and 14 visual angles) and 8 different angles (0, 45, 90, 135, 180, 225, 270, and 315 degree). Stimuli are presented on an LED-lit monitor (ASUS VG248QE: 24in, resolution 1920x1080, refresh rate 144 Hz) positioned 28.5 cm in front of the animal's eyes.

Single-neuron recording in FEF. Single-neuron recordings in awake monkeys were made through a surgically implanted cubic titanium chamber (30×30 mm). The sixteen-site, linear V-probe electrode (125 μm distance of sites) was lowered into the cortex using a hydraulic microdrive (Narashige) to record the extracellular activity of single cells. During each experiment, a recording site in the FEF was first localized by the ability to evoke fixed-vector, saccadic eye movements with stimulation at currents of<50μA (Bruce et al. 1985).

Spike sorting. To isolate single cells' activity, we used multivariate t-distribution decomposition and Expectation-Maximization for clustering (Shoham, Fellows, and Normann 2003). After that,

to find the well isolated clusters in spike sorting, we trained a binary classifier for each pair of clusters in one session of recording. For each cluster, we assigned the minimum performance of binary classifiers which separated that cluster from other clusters as an index to measure the isolation power of cluster. For each classifier, the number of samples was equalized and 10-fold cross validation was applied on equalized samples.

Individual and population neural response. At each time period we quantified the firing rate of individual single or multi-unit clusters. The population activity included all of the simultaneously recorded units in a single recording session. So the population representation of location used for the SVM analysis is a point in R^N space, where N is the number of simultaneously recorded units (single and multi) in one session of recording. For neuron-condition pairs, 2 locations were represented in R^2 neural space.

Neural response tuning curves. In order to compute the neural response relative to each neuron's preferred location, for each neuron the 8 stimulus locations were circularly shifted such that the zero-degree location indicated the maximum response, computed by averaging the neural response across the three task periods (visual, memory, and saccade).

Mutual information (MI). We used mutual information as a measure to quantify the ability of neural responses to represent the target location (Shannon 1948). Mutual information quantifies how well an ideal observer can discriminate between target locations based on the neural response. We computed mutual information using the neural firing rate during non-overlapping 150 msec windows within different intervals:

$$MI = -\sum_{r} P[r]\log_2 P[r] + \sum_{s,r} P[s]P[r\,|\,s]\log_2 P[r\,|\,s]$$

Where s is the set of Stimulus and r is the set of neural responses, i.e. mean firing rates. P(r|s) is the conditional probability of observing a neural response r given an stimulus s. p(s) is the prior probability of stimulus s.

The support vector machine (SVM) classifier. We trained an SVM classifier with a linear kernel (Cortes and Vapnik 1995) using the population and individual neural responses on a randomly selected 70% of trials as a training set. Then we measured the classification accuracy of the trained classifier on the remaining 30% of trials. Training was done using a least square method for finding the separating hyper-plane (Suykens and Vandewalle 1999). To calculate the standard error of classification accuracy in one session, we repeated the calculations 500 times with different sets of training and test trials. In each repetition, the data were randomly partitioned into training and test sets. We trained and tested the SVM with stimuli represented either in 1 dimension (for 'individual' decoding accuracy), or n dimensions, where n is the number of units in a given recording session (for 'ensemble' decoding accuracy).

Decoding tuning curves. To compute decoding tuning curves (Fig. 1E) we extracted the classification accuracy of condition C (C is one angle) by using the classifier confusion matrix. The eight possible target locations were circularly shifted such that zero location indicted the location with maximum performance, computed by averaging classification performance over the three task epochs (visual encoding, working memory, and saccade preparation).

Signal and noise correlation. The signal and noise correlation between each pair of simultaneously recorded units was computed for each pair of target locations (a neuron-condition pair). The correlation coefficient across trials captured the total correlation for each neuron-condition pair. The signal correlation was the correlation coefficient computed after shuffling the order of repeated trials for each target location (Fig. S3B). We repeated the shuffling and

calculation of the correlation coefficient 500 times, and used the mean of these *r*-values as the

measure of the signal correlation for that neuron-condition pair. The noise correlation was

defined by subtracting the signal correlation from the total correlation (Pachitariu et al. 2015;

Tremblay et al. 2015).

In Figure 3B, we computed the signal and noise correlations for pairs of conditions for preferred

and nonpreferred target locations based on the SVM classifier tuning curves. First we calculated

the SVM performance and decoding tuning curve for each session; the target location with

maximal decoding accuracy was considered the 'preferred' location, and the location with the

lowest decoding accuracy was considered the 'nonpreferred' location. Using these locations, we

computed the noise and signal correlations for all pairs of neurons recorded during that session.

Fisher linear discriminant analysis (LDA) and joint discriminability. In each neuron-

condition pair, 2 locations were represented in a 2 dimensional neural space, based on the

simultaneously recorded activity of two neurons. We used Fisher linear discriminant analysis

(LDA) to find the optimum vector for the linear discrimination between the two responses

(Duda, Hart, and Stork 2012). For a population of 2 neurons, let R_s (s = 1, 2) as a 2×N matrix (N

is number of trials) and 2×2 matrix C_s denote the neuronal response covariance matrix for

stimulus s. Let the 2×1 matrix $\bar{R}_s$ be the average population response to stimulus s. We

calculated the optimal weight vector w$_{opt}$ that yields maximum discrimination between two target

locations as

$$w_{opt} = C^{-1}R$$

Where R is the difference of mean vectors of two conditions ($R = \bar{R}_1 + \bar{R}_2$) and C^{-1} is the inverse

of summation of covariance matrixes of the two conditions ($C = C_1 + C_2$). After finding w_{opt}, we

projected the population response of each trial onto the optimal weight vector w_{opt}. This projection is a mapping from two dimensions to scalar. So we could estimate the distribution of projected neural response in one dimension and compute the joint discriminability as the mutual information between the projections of the neural response the target location.

Dimensionality analysis. We sorted the simultaneously recorded units in each session based on their MI, and made a set of neural ensembles by subsequently adding new units (with decreasing MI). Using this approach, we computed the decoding accuracy as a function of the number of units (NU). Then we fitted $y = \alpha(1 - e^{(\frac{-n}{\tau})})$ function, where y is SVM performance and n is number of units, using least-squares and computed dimension as a number of units needed to exceed 75% of maximum decoding accuracy. To measure the standard error of the dimensionality, we used a bootstrapping method that sampled the session's ensembles with replacement (1000 times).

Supplementary references

Asaad W F, Santhanam N, McClellan S, and DJ Freedman. 2013. "High-Performance Execution of Psychophysical Tasks with Complex Visual Stimuli in MATLAB." *Journal of Neurophysiology* 109 (1): 249–60.

Bruce CJ, Goldberg ME, Bushnell MC, and GB Stanton. 1985. "Primate Frontal Eye Fields. II. Physiological and Anatomical Correlates of Electrically Evoked Eye Movements." *Journal of Neurophysiology* 54 (3): 714–34.

Cortes C, and V Vapnik. 1995. "Support-Vector Networks." *Machine Learning* 20 (3): 273–97.

Duda RO, Hart PE, and DG Stork. 2012. *Pattern Classification*. John Wiley & Sons.

Pachitariu M, Lyamzin DR, Sahani M, and NA Lesica. 2015. "State-Dependent Population

Coding in Primary Auditory Cortex." *Journal of Neuroscience* 35 (5): 2058–73.

Shannon CE. 1948. "A Mathematical Theory of Communication." *The Bell System Technical Journal* 27: 379–423, 623–56.

Shoham S, Fellows MR, and RA Normann. 2003. "Robust, Automatic Spike Sorting Using Mixtures of Multivariate T-Distributions." *Journal of Neuroscience Methods* 127 (2): 111–22.

Suykens JAK, and J Vandewalle. 1999. "Least Squares Support Vector Machine Classifiers." *Neural Processing Letters* 9 (3): 293–300.

Tremblay S, Pieper F, Sachs A, and J Martinez-Trujillo. 2015. "Attentional Filtering of Visual Information by Neuronal Ensembles in the Primate Lateral Prefrontal Cortex." *Neuron* 85 (1): 202–15.