Evolution of passwords for cost-free, honest signalling between symbionts and hosts

Marco Archetti[1], J. Benjamin Miller[1], Douglas W. Yu[1,2,*]

[1] School of Biological Sciences, University of East Anglia, Norwich Research Park, Norwich, Norfolk NR47TJ UK

[2] State Key Laboratory of Genetic Resources and Evolution, Kunming Institute of Zoology, Chinese Academy of Sciences, Kunming, Yunnan 650223, China

* Corresponding author: tel: +44 7510308272 and +86 18608717369; fax: +441603592250; dougwyu@gmail.com & douglas.yu@uea.ac.uk; Twitter: @insectsoup

Abstract: 210 words

Main text: 5462 words

References: 31

Figures: 5

Tables: 0

Text boxes: 0

Appendix: 1

Running title: Password signalling in mutualisms

Abstract

Honest communication between potential partners with conflicting interests is generally thought to require costly signals. Costly signalling can explain partner choice when it is possible to link a strategic cost to an individual's quality, like in mate choice. However, in mutualisms, it is usually impossible to link a cost to the likelihood that a potential partner will behave cooperatively in the future. In fact, signals like Nod factors in rhizobial bacteria, which form symbioses with leguminous plants, are evidence of cost-free, honest signals in situations of potential conflict. How can such a signalling system evolve? We use a population-genetics model to show that a cost-free, honest signal can evolve when the receiver is under *soft selection*, which is when high juvenile mortality does not lead to a corresponding reduction in fitness, a common occurrence in many species. Under soft selection, senders evolve increasingly complex messages of identity, a system akin to a password or a lock and key. Thus, a symbiont can signal that it shares a coevolutionary history with a potential host, and if that history is mutualistic, then the host can believe that the symbiont is mutualistic. Password signalling might also explain the evolution of complex species-recognition signals in mate choice and in the acquisition of defensive symbionts.

Key words: evolutionary game theory, legume, horizontal transmission, mate choice, microbiome, mutualism, Nod factor, partner choice, partner fidelity feedback, rhizobia, signalling, defensive symbiosis, vertical transmission

"...Say now Shibboleth: and he said Sibboleth: for he could not frame to pronounce it right. Then they took him, and slew him...." Judges 12:6, King James Version

Introduction

A 'mechanism-design' problem in the rhizobia-legume mutualism

Let us think of communication between symbionts and hosts as a signalling problem. While both mutualistic and parasitic partners have an incentive to enter a host, parasites decrease a host's fitness. Hence interests are not aligned, similar to what is found in mate choice, where males of both high and low quality have an incentive to mate, and females have an incentive to choose high-quality males. The host faces a 'mechanism-design' problem: how to design a signalling system in which a mutualistic symbiont can uniquely identify itself as a mutualist to

a host. In other words, a symbiont must be able to signal its 'mutualistic identity' (i.e. that the

signaller will be reliably 'nice' to the host).

The difficulty with using the classic mechanism of costly signals (Grafen 1990; Maynard Smith

and Harper 2003) is that it is not readily apparent how a mutualistic nature can be correlated

with the strategic cost that is required for the maintenance of honest signalling in situations of

non-aligned interests. Costly signalling is arguably possible in a few cases like big,

symmetrical flowers, in which a signal of vigour *per se* can honestly signal that the flower is

likely carrying high amounts of rewards. But in most mutualisms, the mere demonstration that

a symbiont is vigorous does not also demonstrate that the symbiont will also *behave*

mutualistically in the future (Edwards and Yu 2007). Thus, a strategic cost does not seem to

provide a good explanation for honest signalling in mutualisms.

Nonetheless, evolution is exceedingly clever and seems to have solved this mechanism-design

problem for Nod factors in legume-rhizobia symbioses (Oldroyd 2013). When the root of a

leguminous plant perceives Nod factor from rhizobial bacteria, the root initiates signalling

events that results in the formation of an 'infection thread' to allow the bacteria to colonize the

host plant. All sorts of bacteria would benefit from gaining entry to a root, so there is a strong

temptation for pathogens to counterfeit Nod factor. Nonetheless, only rhizobial bacteria appear

to make Nod factors that are successfully recognised by the plant to initiate signalling and

infection events.

There are three general theoretical classes of honest signalling. We can rule out the '**costly,**

honest' signal explanation of Nod factor, since there is no obvious *strategic* cost (additional to

the mere cost of producing the necessary molecules [Maynard Smith and Harper, 2003]) to

Nod factor. That is, Nod factor is not a bundle of ammonium molecules, serving as evidence

that the bacterium is capable of fixing nitrogen. We can also exclude '**cheap talk**' signalling

(Crawford and Sobel 1982), which requires shared (partially aligned) interests between

signaller and receiver individuals, such as occurs between kin. But in horizontally transmitted

mutualisms, host and symbiont are different species and disperse separately, erasing shared

interests. The third class, '**verifiable information**,' requires that the signal be true on its face.

In biology, verifiable-information signals exist within the concept of the *index* (Maynard Smith

and Harper 2003). For instance, claw marks high up on a tree trunk are a self-evident, and thus

believable, signal of a tiger's large size. For Nod factor to be an index, it would need to be a

unique by-product of the same biochemical pathway that leads to the quality being sought

(nitrogen fixation), so that the mere presence of Nod factor would indicate a mutualistic

symbiont, but this is not the case. Nod factor can, in principle, be synthesized by non-nitrogen-

fixing rhizobial bacteria, which would appear to rule out verifiable-information signals as well,

but we will show how Nod factor can be included in this class.

Cost-free, honest signals: passwords and Nod factors

To start, we propose that Nod factors be thought of as '**password signals**,' cost-free messages

of arbitrarily high complexity that can honestly convey identity. The idea that Nod factors serve

as passwords arises naturally from the many descriptions of Nod factors as chitin-based chains

adorned with multiple "decorations" that vary across rhizobial species (**Fig. 1**) and the portrayal

of Nod factors and Nod-factor receptors acting in a 'lock-and-key' manner (Parniske and

Downie 2003), such that different Nod factors are accepted by different plant species (Perret

et al. 2000*a*), leading to a high degree of species-specificity in rhizobia-hostplant associations.

Nod factors are famously variable in structure, with a lipochitooligosaccharide (LCO)

backbone of three to five N-acetylglucosamine residues to which multiple different

"decorations" can be added, e.g. addition of an acetyl, methyl, sulphate, or sugar moiety (**Fig.**

1). The set of Nod factor decorations varies across rhizobial species and biovars, as does the

length of the LCO backbone and the acyl (fatty-acid) chain added to the non-reducing terminus.

These degrees of freedom give rhizobia the capacity to produce a multitude of Nod-factor

variants (Miller and Oldroyd 2011). For instance, if each of 7 positions on the LCO backbone

can have one of three possible decorations (including no decoration), there are $3^7 = 2,187$

possible variants, not counting length variation in the LCO backbone and acyl chains. The

genetic architectures of Nod factors and Nod-factor-receptor genes both appear to favour rapid

diversification. Known Nod-factor receptor genes are found in tandem arrays, and this might

allow rapid evolution of these genes via recombination (Parniske and Downie 2003).

Similarly, the diversity of *nod* genes allows rhizobia to add multiple and different decorations

to the LCO backbone and to simultaneously produce multiple different Nod factor molecules

(Miller and Oldroyd 2011).

	Acyl	**R₁**	**R₂**	**R₃**	**R₄**	**R₅**	**R₆**	**n**
Nod factors	C16, C18, C20 (varying degrees of saturation; between 0 and 4 double bonds)	H, S, Ac, Fuc, MeFuc, AcFuc, AcMeFuc, SMeFuc	H, Ara	H, OH, Ac, Fuc	H, Ac, Cb	H, OH, Cb	H, Me	-1, 0, 1, 2, 3
Myc factors	C16:0, C16:1, C16:2, C18:0, C18:1 (N.B. non-LCO Myc factors have also been described; see Genre *et al.* 2013)	H, S	H	OH	H	OH	H	1, 2

Fig. 1. Structures of Nod and Myc factors. **A**. The major Nod factor produced by *Sinorhizobium meliloti* consists of four N-acetylglucosamine residues (black) and a C16:2 acyl group (green). This lipochitooligosaccharide (LCO) backbone is further decorated with sulphate (red) and acetyl (blue) groups. **B**. Generalised structure of Nod and Myc factors. The table shows some of the major decorations and length variants that have been characterised to date. **Ac** acetyl, **Ara** arabinosyl, **Cb** carbamoyl, **Fuc** fucosyl, **H** hydrogen, **Me** methyl, **OH** hydroxyl, **S** sulphate, **AcFuc** acetylated fucose, **MeFuc** methylfucose, **AcMeFuc** acetylated methylfucose, **SMeFuc** sulphated methylfucose. Figure adapted from Perret et al. (2000), Wais et al. (2002), and Miller and Oldroyd (2011).

However, what are the conditions under which passwords evolve via natural selection? Let us say that a bacterial lineage has co-evolved with a plant and that during this time, the symbiosis has progressed from an ancestral state of close association at the root surface (e.g. the plant

root secretes carbohydrate compounds into its immediate surroundings and captures

ammonium by diffusion and/or active transport) to the more intimate state of endosymbiosis,

via the formation of an infection thread and root nodule. The benefit of evolving endosymbiosis

is increased efficiency of nitrogen fixation in an oxygen-free environment, and the exclusion

of free-riders is a possible additional benefit. *Nod*-gene duplication and mutation allowed

rhizobia to evolve Nod factors of greater and greater complexity (more decorations added to

the LCO chain). Nod factor does not have to be costly, except in the trivial sense that it needs

some energy to be synthesized, and indeed Nod factor does not seem to have a *strategic* cost.

However, Nod factor does need to be uniquely recognizable, which could explain why it is

complex.

During the evolution of Nod factor, we assume that the host plant also had ancestral

physiological mechanisms for shedding or withdrawing resources from worthless roots that

have failed to take up fixed nitrogen (Partner Fidelity Feedback PFF, Weyl et al. 2010) or had

even evolved punishment (Host Sanctions HS, Kiers et al. 2003). Either way, rhizobial lineages

with the correct Nod factor have been subjected to selection for nitrogen fixation and against

pathogenicity. The usefulness of passwords is that they can truthfully signal that a bacterial

lineage shares a co-evolutionary history with a host lineage, and if that history is mutualistic,

then the plant is selected to engage in symbiosis with bearers of the password. Thus, passwords

signal a particular evolutionary history (an 'identity'), and since that history is written into the

bacterial symbiont's genome, the genome enforces a particular behaviour.

This is what a person does with a password on a bank website: he credibly signals to the bank

that they share a specific history of transactions, and if those transactions have been acceptable

to the bank, there is a good chance that the person will continue to act acceptably, so the bank

should allow continued transactions. (Passwords are an alternative to repeated games. By

definition, players of repeated games build up interaction histories with other players and thus

need to recognize individuals in order to apply the correct history to each player, but with

passwords, it is possible for a member of a host lineage to recognize a member of a specific

symbiont lineage, even if that pair of individuals has never met.)

For a password to be a reliable signal of identity, the password needs to be complex. Otherwise,

a password that identifies one lineage could easily be evolved *de novo* by other lineages. Thus,

our challenge becomes one of explaining why there is directional selection for signal

complexity. We also need to explain 'strictness' in receivers, where strict means that the

receiver rejects passwords with (too many) errors. It is the combination of a sufficiently

complex signal and a sufficiently strict receiver that renders it essentially impossible for

parasites to evolve a working password *de novo*.

Passwords can evolve under soft selection

In the context of mutualistic interactions, the process we envisage leading to the evolution of

cost-free but still honest signalling is the following. Consider a population of hosts harbouring

only mutualistic symbionts. Mutations in the symbionts lead to a slightly different (simpler or

more complex) password or to a parasitic phenotype. We assume that double mutations are

negligible, meaning that a symbiont will not simultaneously evolve a new password and a

newly parasitic phenotype. We also let mutations in the hosts lead them to accept a slightly

different (simpler or more complex) password. Hosts accepting the original password will

therefore accept both mutualistic and the new mutant parasitic partners, but hosts with

mutations that cause them to accept a different password will reject parasitic symbionts, which

still use the original password under the assumption that double mutations are negligible.

Mutant *hosts* will have higher fitness than the resident hosts if the benefit of avoiding parasites

is higher than the cost of the difficulty of finding the rarer partners carrying the right mutant

password (i.e. some mutant hosts fail to find a symbiont partner). If this is the case, mutant

symbionts with the slightly different (simpler or more complex) passwords will also have

higher fitness, because mutualistic symbionts with the original password suffer some of the

cost of PFF or HS triggered by the co-colonising parasitic mutants (because the host expends

energy to trigger PFF or HS and because of the opportunity cost of mutualists having lost out

on colonisation opportunities to parasites). Then, we must only explain why the mutant

symbionts with the more complex password have higher fitness than the mutants with the

simpler password. We will show that, *provided that the hosts undergo soft selection*, the hosts

that mutated to accept the more complex passwords have a selective advantage, and as a result,

the signallers that mutated to slightly more complex passwords will also have a selective

advantage. Repeating this scenario over time leads to increasingly complex passwords that are

honest and cost-free.

The critical, non-intuitive step in this scenario is the contribution of *soft selection* (Buchholz

1922; Wallace 1981; Klekowski 1988) (**Fig. 2**). Soft selection occurs when juveniles are

produced in excess of available carrying capacity for adults. For instance, a plant produces

many more seeds than there are patches in the environment to support adults, and as a result,

juvenile populations are unavoidably bottlenecked by this exogenous mechanism, or to put it

another way, when a parent loses some fecundity, its fitness remains virtually unaffected, since

some offspring would have died no matter what. Since all offspring contain *de novo* mutations,

which, if non-silent, are more likely to be deleterious than beneficial, soft-selected lineages

evolve genes that are hypersensitive to mutation (i.e. antirobust); that is, soft-selected genes

evolve to suffer large losses in function after mutation, causing the offspring that carry more

mutations to be much more likely to die, which reduces competition with their more fortunate

siblings that carry fewer mutations. Parental fitness benefits more from this filtering out of

mutated offspring than is lost to reduced fecundity, since most offspring are destined to die

anyway in a soft-selection scenario.

In what follows, we use a population-genetics model to show that soft selection leads to the

evolution of more complex passwords. We stick with the scenario in which the signaller is a

bacterium and the receiver is a host plant, but the model applies in general to partner choice.

Figure 2. Soft selection leads
to antirobustness. Alleles x_1
and x_2 have the same fitness.
However, x_1's mutants x_3
happen to have lower fitness
than do x_2's mutants x_4 ($z_3 > z_4$,
so $(1-\mu z_3) < (1-\mu z_4)$), and
hence, x_1 is less robust than x_2.
If there is a juvenile selection
stage, x_3 mutants are more
likely to be eliminated before
the adult stage; if juvenile
selection does not cause a loss
of fecundity (because only a
small fraction of the progeny

goes on to the adult stage M anyway), the adult progeny of x_1 will have a selective advantage

over the adult progeny of x_2, and x_1 will increase in frequency (Image reproduced with some

changes from Archetti 2009).

Model

Consider a population of nitrogen-fixing rhizobial bacteria producing a password composed of separate elements *a, b, c,* etc. and allow each of these elements to arise or disappear in a stepwise fashion by a mutation that can add or remove one element (a mutation is denoted by adding or removing a letter from the password). For simplicity of notation, the order of the elements does not matter; that is, passwords of increasing complexity are denoted by *a, ab, abc,…,* etc., and *ab=ba, abc=acb,* etc.. Note also that any element of a password can mutate (e.g. *abc → tbc → tbu*), but we ignore these scenarios here because our focus is on the origin of complexity *per se*, which we are representing by a longer password. In other words, we focus on explaining how to get from passwords of low to high complexity (*a → abcde*), and we ignore the diversification of passwords (*abcde → uewix*). Our way of denoting more complex passwords mimics the idea that Nod factor has evolved increased complexity by the proliferation of additional *nod* genes that have added more "decorations" to the basic LCO backbone (Miller and Oldroyd 2011).

In each generation, a fraction μ_P of the population produces a simpler password, and a fraction μ_P produces a more complex password. For instance, in a population of bacteria sending password *ab*, some mutants will send the password *a*, and some will send the password *abc*. Bacteria can also evolve to be parasitic by failing to export fixed nitrogen to the host plant. Hence, the population will also have a fraction μ_D of parasitic *ab** bacteria. Assuming that double mutations are rare, we can ignore *a** and *abc** mutants. Finally, we denote plants that accept a given password by capital letters; each plant type only accepts its own corresponding bacterium type (AB accepts only *ab* and *ab**; A accepts only *a*; ABC accepts only *abc*). This assumption of strictness in the plants is here made only for explanatory convenience, and we explicitly model the evolution of strictness in the next section. We start with a population fixed on AB plants and *ab* bacteria, with a low frequency of mutants ABC, A, *abc, a,* and *ab**.

Signaller (Bacteria)

Selection coefficients s_i against the different bacterial types *i* are as follows (See **Appendix** for model details):

- $s_{ab}>0$ because AB plants can be colonized by *ab* and *ab** bacteria. *ab* bacteria suffer some fitness loss due to *ab**'s cheating, as a consequence of the costs of host response against *ab** (direct costs of nodule senescence via HS or PFF in mixed nodules, indirect costs of

senescence due to plant expenditure of energy, and opportunity costs of *ab* bacteria having

suffered from competition with *ab** bacteria for limited colonization opportunities).

- $s_{ab*}>0$ for the same reasons as above. If the plant reacts indiscriminately against *ab* or *ab**

then $s_{ab*}<s_{ab}$ since *ab** do not pay the cost of nitrogen fixation. If HS or PFF is

preferentially targeted toward *ab**-inhabited nodules, then $s_{ab*}>s_{ab}$.

- $s_{abc}=0$ because there is no costly host response by ABC plants, since there are no *abc**

bacteria. Strictly $s_{abc} = 0$ when *abc* and ABC have zero additional difficulty in finding each

other, relative to *ab* and AB finding each other. The assumption is reasonably upheld under

soft selection because ABC plants make many ABC seedlings, which sample the bacterial

population, and only some need to pick up *abc* bacteria. Note that this is an origin scenario

for *abc* passwords, so a successful colonization of ABC plants only has to happen once in

history for *abc* bacteria to start to be selected.

- $s_a=0$ for the same reason that $s_{abc}=0$. There is no costly host response by A plants since

there are no *a** bacteria in the population, due to our assumption of no double mutations.

Therefore it is trivial to conclude that simple and complex passwords (*a* and *abc*) have the same

fitness and that their fitnesses are greater than that of *ab* and *ab**, because under our assumption

of no double mutants, only *ab* evolves cheating mutants *ab**, which elicit nodule senescence

in plants and compete for nodules and which therefore reduce fitness for *ab* as well. So either

a simpler password *a* or a more complex password *abc* will evolve from *ab*. Which one of the

two prevails is determined by their respective hosts. If ABC plants increase in the population,

then *abc* bacteria will have an advantage over *a*, and selection is for more complex passwords.

So we have to explain why ABC plants evolve, rather than A plants. In other words, so far we

have easily proven the well-known fact that selection induced by parasites induces diversity in

a population. We are left to explain why this diversity leads to increased signal complexity.

Our solution in the next section will invoke soft selection.

Before going on, it is possible to outline a simpler but perhaps less general scenario for the

evolution of password signalling. It might be the case that $s_a>0$ (i.e. *a* suffers a fitness cost) if

there are *a** bacteria in the population, which elicit PFF response in plants, and this response

harms *a*, as in the argument for *ab* having $s_{ab}>0$. This could occur if *a* passwords are simple

enough that they can be evolved *de novo* by parasitic bacteria. In this case, it seems obvious

that *abc* will go to evolutionary fixation, as long as the benefit of being a pathogen is not greater

than the cost of PFF. Then, ABC plants will increase simply because they accept only *abc*

bacteria, whereas A and AB plants accept some pathogenic bacteria (*a** and *ab**). Note that

this scenario still requires soft selection, because we assume that there is no loss of fitness for

ABC in finding the (initially) rare partner type *abc*.

In the following, let us retain the more conservative assumption that $s_a=0$ (i.e. parasites induce

diversity in the host, but this diversity is not biased towards higher or lower complexity). The

next part of our explanation is to show why higher complexity (*abc*) has a selective advantage

over lower complexity (*a*).

Receivers (Plants)

Let us consider a locus with four alleles i = ABC, A, ABC*, A*, with frequencies x_i, coding for

the receiver's recognition system. We only need to model two types (A and ABC) and their

nonsense or missense mutants (A*, ABC*):

•   Allele ABC always accepts password *abc*

•   Allele ABC* is a parasitic mutant of allele ABC

•   Allele A always accepts password *a*

•   Allele A* is a parasitic mutant of allele A

Our key assumption is that a mutant of A (A*) is more likely to still accept its password *a* than

a mutant of allele ABC (ABC*) will accept its password *abc*. In other words, we posit that

allele A is more *robust* to mutation and that allele ABC is *antirobust* to mutations (sensitive).

Note that robustness is unrelated to viability. A and ABC have exactly the same fitness (since

abc and *a* bacteria have the same effect on the plant's fitness); it is their mutants (A* and ABC*)

that have different fitnesses.

The mechanism behind our assumption could simply be that an ABC receiver system is

necessarily made up of more numerous or complex molecules that interact with each other,

since password *abc* is physically more complex than password *a*. Thus, even a mutation of

'small effect' in one component of ABC might prevent the *abc* signal molecule from fitting

properly in the other component(s) of ABC and thus prevent the different receptor molecules

from interacting correctly with each other to trigger a signalling cascade. A complexly

interacting receptor is also more likely to be a strict receiver, since even a small change in the

Nod factor would cause it fit differently in one or more of the receptor's molecules and thus

interfere with the interaction of those molecules.

We are thus looking for evidence that the Nod-factor receptor complex is composed of multiple

molecules that need to interact with each other in order to trigger a proper signalling cascade.

As it happens, there is considerable empirical support for this model (Oldroyd 2013). The Nod-
factor receptor complex is made up of two separately produced receptor-like kinases (NFR1
and NFR5, using *Lotus japonicus* nomenclature), each of which carries extracellular LysM
motifs that bind to the *N*-acetylglucosamine backbone of the Nod-factor backbone
(Broghammer et al. 2012). The two proteins heterodimerise *in vivo* (Madsen et al. 2011), and
mutation in either of the two receptor-like kinases prevents rhizobial infection (Radutoiu et al.
2003). Importantly, NFR5 is a non-functional kinase (this is known because it lacks essential
protein subdomains), and thus, NFR5 can act only via its interaction with NFR1 (Madsen et al.
2011). The activated NFR1/5 complex then appears to activate a third receptor-like kinase
(SYMRK), which is necessary for downstream propagation of the infection pathway (Radutoiu
et al. 2003). Finally, Morieri et al. (2013) have shown that the removal of just one acetyl group
decoration from the *Sinorhizobium meliloti* Nod factor is enough to prevent calcium influx in
its host, *Medicago truncatula*, and this failed influx prevents the initiation of an infection
thread. Morieri *et al*. propose a model in which *only the correct Nod factor is able to bring*
about "cooperative interactions" between receptor proteins "such that the resulting
interaction alters the kinase activity or specificity of the receptor complex", triggering the
calcium influx that is needed for successful infection-thread initiation. In short, the Nod factor
receptor is clearly a machine of many interdependent parts, and thus, of many points of failure.

The evidence that an A receptor would be more robust to mutation is sparser, because all known
Nod-factor receptors (and Nod factors) are complex (Perret et al. 2000*b*; Madsen et al. 2011;
Miller and Oldroyd 2011; Broghammer et al. 2012). However, the Myc factors produced by
arbuscular mycorrhizal (AM) fungi are structurally simpler (Maillet et al. 2011), and multiple,
distantly related plant species will enter into symbiosis with the same AM fungal genotype,
despite the reasonable expectation that the Myc receptor complexes from different plant
lineages have mutated during plant diversification. However, because Myc-factor receptors are
not yet characterised, it is not yet possible to rule out the alternative hypothesis that individual
plant species produce multiple, species-specific Myc receptors.

Now, the evolution of password complexity requires that ABC increase in frequency over A.
The question therefore is: why should ABC increase in frequency, given that alleles ABC and
A are neutral?

If anything, it seems that allele A should increase, since it is more robust to mutation and will
therefore have a higher rate of back mutations from allele A[*], whereas allele ABC[*] is less likely
to survive and will provide fewer back mutants to ABC (Wagner et al. 1997; Hermisson et al.

2002; de Visser et al. 2003). We can see this effect in **Fig. 3**, where we assume the standard

hard-selection scenario in which juvenile mortality reduces adult fitness ($z_{A*}=z_{ABC*}=0$). The

frequency (x_A) of the robust allele A increases, provided that the selection coefficient s_{ABC*}

against the defective mutant ABC* is high enough (approximately higher than the mutation

rate). If selection is too weak compared to the mutation rate, the differential amount of back

mutations (to ABC and A from AB) is negligible, and the two alleles A and ABC maintain the

same frequencies. If selection is strong enough, however, with no soft selection, the robust

allele A increases in frequency over ABC because it receives more back mutations from A*

than ABC receives from ABC*.

However, with soft selection, antirobust alleles increase in frequency over robust alleles (**Figs.**

3, 4, Appendix 1). In fact, if the selection coefficients are high enough, the antirobust allele

ABC can go to fixation. Strong selection is necessary, but not high mutation rates. Contrary to

the evolution of robustness, the magnitude of the mutation rate does not make any relevant

difference to the evolution of anti-robustness because the driving force is not the rate of back

mutation, but the soft selection process that eliminates the non-functional ABC mutants (Otto

and Hastings 1998). Lower rates of mutation to non-functional ABC mutants do enable

complex passwords to evolve for lower levels of soft selection (**Fig. 4A**).

The fraction of complex passwords at equilibrium increases as the ratio between the mutation

rates for passwords and the mutation rate to defective alleles decreases (**Fig. 4B**), because if

the mutation rate to defective alleles is too high, the advantage derived from soft selection is

offset by partial loss of fecundity. A similar effect is observed for z_i/s_i (**Fig. 4D**), because when

juvenile selection is stronger, defective alleles are more easily eliminated from the offspring

(hence, the original antirobust allele is more likely to increase in frequency). The influence of

s_{ABC*}/s_{A*} is more complex, as it depends on the value of z_{ABC*} (**Fig. 4C**). Note that soft selection

can lead to an increase in the complex (antirobust) allele even if complexity itself has a small

cost (that is if $s_{ABC} > 0$; **Figure 5.**)

Figure 3. Equilibrium frequencies of the four receptor alleles as a function of selection, with

hard selection (left, $z_i=0$) or with soft selection (right, $z_i=s_i$); $s_{A*}/s_{ABC*}=1/10$; $s_A=s_{ABC}=0$. $\mu_S=10^{-}$

^{4}, $\mu_P=10^{-7}$. s_{A*} and s_{ABC*} are selection coefficients against dysfunctional A and ABC alleles,

which arise with probability μ_S. The x-axis in the top two figures is logarithmic, and in the

bottom two figures is linear.

Figure 4. Equilibrium frequency of the complex, antirobust receiver system (x_{ABC}) as a

function of juvenile selection (z_{ABC*}); $s_A = s_{ABC} = 0$. (A) for different values of μ_S; $s_{ABC*}/s_{A*} = 10$;

$s_i/z_i = 1$; $\mu_P = \mu_S/50$. (B) for different values of μ_S/μ_P; $s_{ABC*}/s_{A*} = 10$; $s_i/z_i = 1$; $\mu_S = 10^{-2}$. (C) for

different values of s_{ABC*}/s_{A*}; $s_i/z_i =1$; $\mu_S=10^{-2}$; $\mu_S/\mu_P=50$; (D) for different values of s_i/z_i;

$s_{ABC*}/s_{A*}=10$; $\mu_S=10^{-2}$; $\mu_P=\mu_S/50$.

Figure 5. Equilibrium frequency of the complex, antirobust receiver system (x_{ABC}) as a function of juvenile selection (z_{ABC*}); $s_A=0$, for different values of s_{ABC}; $\mu_S=1\times10^{-4}$, $\mu_P=1\times10^{-6}$ $s_{ABC*}/s_{A*}=10$, $s_{ABC*}/z_{ABC*} =s_{A*}/z_{A*} =1$; (A): with juvenile selection equal to adult selection on ABC ($s_{ABC}=z_{ABC}$); (B) with only juvenile selection on ABC ($s_{ABC}=0$); (C): with juvenile selection equal to 1/10 of adult selection on ABC ($s_{ABC}= z_{ABC}/10$). Antirobustness/complexity evolves even if there is strong selection against the antirobust/complex allele (ABC) in the

394 juveniles (B); if there is selection on the adults too, however, antirobustness/complexity

395 evolves only with weak selection against the antirobust/complex allele (ABC).

Discussion

Game theory recognizes three classes of signalling models: costly-honest signals, cheap talk,

and verifiable information. Passwords can be thought of as a variant of the verifiable-

information class. By virtue of their complexity and thus the low likelihood that they can evolve

de novo, they serve as signals of identity that are self-evidently true so long as the receiver can

recognise the password. In the context of horizontally transmitted mutualisms, passwords can

evolve to reliably signal a shared coevolutionary history, and a coevolutionary history between

mutualistic lineages strongly implies that the individual sending the password is itself a

mutualist or a recent descendent of a mutualist. There is a superficial similarity of password

signalling with green-beard signals (Jansen and van Baalen 2006), which allow kin to identify

each other, but green beards are a within-species mechanism, and passwords can act between

species. Green-beard signals can also be simple, due to their linkage with cooperation loci.

As a result, password signalling can allow hosts to engage in successful Partner Choice. As it

happens, it has been shown experimentally that legume plants are able to associate

preferentially with 'more mutualistic' (nitrogen-fixing) rhizobial bacteria (Heath and Tiffin

2009; Gubry-Rangin et al. 2010; Sachs et al. 2010). Importantly, these studies used mutualistic

and parasitic rhizobial bacteria that had been isolated from the same soil as the host plant, and

we predict that the rejected parasitic bacterial lineages used in these experiments may have

been producing Nod factors that had diverged from the mutualistic lineages that were accepted.

Our proposed scenario for the evolution of password signalling derives from the observation

that plants are subject to a non-trivial degree of soft selection, since plants generally make

many more juveniles than can possibly grow into adults. We then posit that mutations in the

genes for complex-signal receptors (ABC) are inherently more likely to result in non-functional

receptors than are mutations in the genes for simple-signal receptors (A), because we know

that complex Nod-factor receptors are constructed from multiple, interdependent parts. Thus,

only fully functional ABC-receptors are likely to be represented in adult plants, because ABC-

mutant juveniles will have died due to an inability to recruit rhizobia. In contrast, when the

genes for A-type receptors mutate, the receptors are more likely to retain some function

because they are simple, and some of these lower fitness mutants will thus be represented in

the adult stage. Competition between ABC and A adults will then favour ABC, and thus,

complex rhizobial passwords (*abc*) will also be favoured, and the system will evolve toward

complex signals of identity. After enough rounds, Nod factor will have evolved to a high

enough degree of complexity that it will be essentially impossible for a bacterium to evolve a

working Nod factor *de novo*.

We also recall our first, and simpler, scenario for the evolution of complex signalling

passwords, which relies on the possibility that simple Nod factors can evolve *de novo* in non-

mutualistic bacteria (a*). In this situation, bacteria that evolve more complex passwords (*abc*)

are favoured over those that evolve simpler passwords (*a*), as the former will find themselves

in parasite-free hostplants, at least until *abc* bacteria evolve parasitic behaviour *abc**. This

scenario also relies on soft selection, in that many ABC juvenile plants will die before finding

a suitable *abc* partner, but as long as there are lots of ABC juveniles, some will be successful,

and these will form the next generation.

Limits to complexity in passwords. - In either scenario, we expect a natural upper limit to the

complexity of passwords because there will be physical limits on the reliable functioning of

complex-signal receptors ($s_{ABC} > 0$), and receptors that evolve beyond these limits will likely

fail to perceive any symbionts, which reduces the effect of soft selection (Fig. 5). Thus, the

evolution of complexity in passwords cannot escape indefinitely from the evolution of parasitic

genotypes within rhizobial lineages. There must also be mechanisms to senesce nodules that

have been colonised by parasites (Kiers et al. 2003; Weyl et al. 2010). And of course, such

mechanisms were necessary to proliferate the mutualistic genotypes of rhizobia in the first

place, or there would have been no mutualistic lineages for the plant to recognise.

Once a combination of password signalling and selective nodule senescence has evolved,

mutualistic strains of rhizobia should grow to dominate soils. As a result, it is possible to

imagine situations where some hosts will evolve to relax the strictness of association or evolve

to accept multiple passwords. As one example, some leguminous tree species, and the non-

legume Cannabaceae plant genus *Parasponia*, are early-successional species that colonise low-

nutrient soils, and they can be colonised by multiple rhizobial genera, including strains from

different continents (Behm et al. 2014). Under such conditions, we expect that the risk of being

colonised by non-productive or even pathogenic bacteria is outweighed by the benefit being

able to fix nitrogen. It will be interesting to see if these species have evolved multiple Nod

factor receptors, or if their receptors are less strict (which should make them more robust to

mutation). Indeed, in *Parasponia andersonii*, it appears that the latter might be true, because

this species uses the same receptor for both Myc and Nod factors (Op den Camp et al. 2011).

Myc factors, which consist of simple, almost entirely undecorated LCOs (Maillet et al. 2011),

provide an interesting counterexample to complex Nod factors. Why have AM fungi not

evolved complex passwords? Part of the answer is likely due to the fact that plants are

colonized by multiple AM fungal species, and by doing so, plants make the fungi compete for

plant carbon, thereby reducing the carbon cost of AM-provided phosphorous (Argüello et al.

2016). A plant that evolved a more complex receptor would reduce its diversity of fungal

partners and thus reduce the number of competing fungal suppliers. It is also possible that each

AM fungal genotype benefits from colonizing multiple plant species, if plants vary temporally

in the photosynthate that they are able to transfer to their fungal partners. An individual AM

fungus that evolved a more complex Myc factor recognised only by the rare plant genotype

that had also evolved a matching receptor would not be able to create to create such networks.

Password signalling in mate recognition and defensive-symbiont acquisition. - Soft selection

occurs in practically all vascular plants, and also in many cryptogamic plants and in animals

(Buchholz 1922; Wallace 1981; Klekowski 1988; Archetti 2009). Although detailed treatments

are outside the scope of this paper, we hypothesize that password signalling can evolve in other

recognition systems. For example, a post-mating recognition system in gametes could require

a complex or a simple signal to differentiate conspecifics from heterospecifics. If the complex

recognition system is more antirobust to mutation, then gametes that have suffered mutation

will die unmated. However, those that survive will only have accepted conspecifics. In contrast,

gametes that accept a simple signal might be robust to mutation and thus accept heterospecifics,

producing hybrids. Under the twin assumptions that soft selection is acting (most juveniles die

before achieving adulthood) and that hybrids have lower fitness, there will be selection for a

mate-recognition system that requires a complex signal. Another possible class of password

signals are polymorphic toxin systems (PTS), which comprise complex, multi-domain

molecules that exhibit high levels of allelic diversity. Hillman and Goodrich-Blair (2016) have

proposed that eukaryotic hosts can directly identify suitable defensive symbionts to acquire by

sensing the PTS produced by those symbionts, and they review evidence that hosts produce

PTS-receptors that are specific to particular symbiont lineages.

Acknowledgments

We thank István Scheuring and Gergely Boza for illuminating discussions. DWY was

supported by the National Natural Science Foundation of China (41661144002), the Ministry

of Science and Technology of China (2012FY110800), the University of East Anglia, and the

State Key Laboratory of Genetic Resources and Evolution at the Kunming Institute of Zoology

(GREKF13-13, GREKF14-13, and GREKF16-09). JBM is supported by a fellowship from the

University of East Anglia and the Eastern Academic Research Consortium.

References

Archetti, M. 2009. Survival of the steepest: hypersensitivity to mutations as an adaptation to soft
selection. Journal Of Evolutionary Biology 22:740–750.

Argüello, A., M. J. O'Brien, M. G. A. Heijden, A. Wiemken, B. Schmid, and P. A. Niklaus. 2016.
Options of partners improve carbon for phosphorus trade in the arbuscular mycorrhizal
mutualism. (H. Maherali, ed.)Ecology Letters.

Behm, J. E., R. Geurts, and E. T. Kiers. 2014. *Parasponia*: a novel system for studying mutualism
stability. Trends in Plant Science 19:757–763.

Broghammer, A., L. Krusell, M. Blaise, J. Sauer, J. T. Sullivan, N. Maolanon, M. Vinther, et al. 2012.
Legume receptors perceive the rhizobial lipochitin oligosaccharide signal molecules by direct
binding. Proceedings of the National Academy of Sciences USA 109:13859–13864.

Buchholz, J. T. 1922. Developmental selection in vascular plants. Botanical Gazette 73:249–286.

Crawford, V., and J. Sobel. 1982. Strategic information transmission. Econometrica 50:1431–1451.

de Visser, J. A. G. M., J. Hermisson, G. P. Wagner, L. Ancel Meyers, H. Bagheri-Chaichian, J. L.
Blanchard, L. Chao, et al. 2003. Perspective: Evolution and detection of genetic robustness.
Evolution 57:1959–1972.

Edwards, D. P., and D. W. Yu. 2007. The roles of sensory traps in the origin, maintenance, and
breakdown of mutualism. Behavioral Ecology And Sociobiology 61:1321–1327.

Grafen, A. 1990. Biological signals as handicaps. Journal of Theoretical Biology 144:517–546.

Gubry-Rangin, C., M. Garcia, and G. Bena. 2010. Partner choice in *Medicago truncatula-*
Sinorhizobium symbiosis. Proceedings of the Royal Society B: Biological Sciences 277:1947–
1951.

Heath, K. D., and P. Tiffin. 2009. Stabilizing mechanisms in a legume-rhizobium mutualism.
Evolution 63:652–662.

Hermisson, J., O. Redner, H. Wagner, and E. Baake. 2002. Mutation–selection balance: ancestry,
load, and maximum principle. Theoretical population biology 62:9–46.

Hillman, K., and H. Goodrich-Blair. 2016. Are you my symbiont? Microbial polymorphic toxins and
antimicrobial compounds as honest signals of beneficial symbiotic defensive traits. Current
Opinion In Microbiology 31:184–190.

Jansen, V. A. A., and M. van Baalen. 2006. Altruism through beard chromodynamics. Nature
440:663–666.

Kiers, E. T., R. A. Rousseau, S. A. West, and R. F. Denison. 2003. Host sanctions and the legume-

rhizobium mutualism. Nature 425:78–81.

Klekowski, E. J. 1988. Mutation, developmental selection, and plant evolution. Columbia University
Press.

Madsen, E. B., M. Antolín Llovera, C. Grossmann, J. Ye, S. Vieweg, A. Broghammer, L. Krusell, et
al. 2011. Autophosphorylation is essential for the *in vivo* function of the *Lotus japonicus* Nod
factor receptor 1 and receptor-mediated signalling in cooperation with Nod factor receptor 5. The
Plant Journal 65:404–417.

Maillet, F., V. Poinsot, O. André, V. Puech-Pagès, A. Haouy, M. Gueunier, L. Cromer, et al. 2011.
Fungal lipochitooligosaccharide symbiotic signals in arbuscular mycorrhiza. Nature 469:58–63.

Maynard Smith, J., and D. Harper. 2003. Animal Signals. Oxford University Press.

Miller, J. B., and G. E. D. Oldroyd. 2011. The role of diffusible signals in the establishment of
rhizobial and mycorrhizal symbioses. Pages 1–30 *in* S. Perotto and F. Baluška, eds. Signaling and
Communication in Plant Symbiosis, Signaling and Communication in Plants (Vol. 11). Springer
Berlin Heidelberg, Berlin, Heidelberg.

Morieri, G., E. A. Martinez, A. Jarynowski, H. Driguez, R. Morris, G. E. D. Oldroyd, and J. A.
Downie. 2013. Host-specific Nod-factors associated with *Medicago truncatula* nodule infection
differentially induce calcium influx and calcium spiking in root hairs. New Phytologist 200:656–
662.

Oldroyd, G. E. D. 2013. Speak, friend, and enter: signalling systems that promote beneficial
symbiotic associations in plants. Nature Reviews Microbiology 11:252–263.

Op den Camp, R., A. Streng, S. De Mita, Q. Cao, E. Polone, W. Liu, J. S. S. Ammiraju, et al. 2011.
LysM-type mycorrhizal receptor recruited for rhizobium symbiosis in nonlegume *Parasponia*.
Science 331:909–912.

Otto, S. P., and I. M. Hastings. 1998. Mutation and selection within the individual. Pages 507–524 *in*
R. C. Woodruff and J. N. Thompson, eds. Mutation and Evolution, Contemporary Issues in
Genetics and Evolution (Vol. 7). Springer Netherlands, Dordrecht.

Parniske, M., and J. A. Downie. 2003. Plant biology: Locks, keys and symbioses. Nature 425:569–
570.

Perret, X., C. Staehelin, and W. J. Broughton. 2000*a*. Molecular basis of symbiotic promiscuity.
Microbiology and Molecular Biology Reviews 64:180–201.

Perret, X., C. Staehelin, and W. J. Broughton. 2000*b*. Molecular basis of symbiotic promiscuity.
Microbiology and Molecular Biology Reviews 64:180–201.

Radutoiu, S., L. H. Madsen, E. B. Madsen, H. H. Felle, Y. Umehara, M. Grønlund, S. Sato, et al.
2003. Plant recognition of symbiotic bacteria requires two LysM receptor-like kinases. Nature
425:585–592.

Sachs, J. L., J. E. Russell, Y. E. Lii, K. C. Black, G. Lopez, and A. S. Patil. 2010. Host control over
infection and proliferation of a cheater symbiont. Journal Of Evolutionary Biology 23:1919–1927.

Wagner, G. P., G. Booth, and H. Bagheri-Chaichian. 1997. A population genetic theory of
canalization. Evolution 51:329.

Wallace, B. 1981. Basic Population Genetics. Columbia University Press.

Weyl, E. G., M. E. Frederickson, D. W. Yu, and N. E. Pierce. 2010. Economic contract theory tests
models of mutualism. Proceedings of the National Academy of Sciences USA 107:15712–
15716.

Appendix

Plants

We assume that alleles A and ABC can mutate to each other with the same probability μ_P and that each can mutate to dysfunctional alleles (respectively $A*$ and $ABC*$) with probability μ_S. We assume that all alleles have the same total mutation rate, hence $A*$ and $ABC*$ have other mutants (A^0 and ABC^0) with frequency μ_P that have zero fitness:

$$ABC^0 \leftarrow\mu_P\rightarrow ABC* \leftarrow\mu_S\rightarrow ABC \leftarrow\mu_P\rightarrow A \leftarrow\mu_S\rightarrow A* \leftarrow\mu_P\rightarrow A^0$$

Individuals with allele i have fitness $1-s_i$ in the adult phase and $1-z_i$ in the juvenile phase. We can assume with no loss of generality that $s_{ABC}=s_A=z_{ABC}=z_A=0$.

The recurrence equations for this system are (see **Table A1**):

$$T \cdot x_{ABC}' = x_{ABC}(1-s_{ABC})(1-\mu_S-\mu_P)(1-z_{ABC})/\alpha_P + x_A(1-s_A)\mu_P(1-z_{ABC})/\beta_P + x_{ABC*}(1-s_{ABC*})\mu_S(1-z_{ABC})/\chi_P$$

$$T \cdot x_A' = x_{ABC}(1-s_{ABC})\mu_P(1-z_A)/\alpha_P + x_A(1-s_A)(1-\mu_S-\mu_P)(1-z_A)/\beta_P + x_{A*}(1-s_{A*})\mu_S(1-z_A)/\delta_P$$

$$T \cdot x_{ABC*}' = x_{ABC}(1-s_{ABC})\mu_S(1-z_{ABC*})/\alpha_P + x_{ABC*}(1-s_{ABC*})(1-\mu_S-\mu_P)(1-z_{ABC*})/\chi_P$$

$$T \cdot x_{A*}' = x_A(1-s_A)\mu_S(1-z_{A*})/\beta_P + x_{A*}(1-s_{A*})(1-\mu_S-\mu_P)(1-z_{A*})/\delta_P$$

where T is a normalizing factor obtained by summing the right-hand side of the four above equations; α_P, β_P, χ_P, δ_P are normalizing factors of the offspring frequencies (see Table 1) ;

$\phi_P=M/N$ is the degree of soft selection, where N is the number of offspring before soft selection (the same for all alleles) and M ($<N$) is the maximum number of individuals that can go on to the adult phase after soft selection. N_i' is the number of offspring of an individual with allele i after soft selection:

$$N_{ABC}' = N[(1-\mu_S-\mu_P)(1-z_{ABC})+\mu_P(1-z_A)+\mu_S(1-z_{ABC*})]$$

$$N_A' = N[(1-\mu_S-\mu_P)(1-z_A)+\mu_P(1-z_{ABC})+\mu_S(1-z_{A*})]$$

$$N_{ABC*}' = N[(1-\mu_S-\mu_P)(1-z_{ABC*})+\mu_S(1-z_{ABC})+\mu_P(0)]$$

$$N_{A*}' = N[(1-\mu_S-\mu_P)(1-z_{A*})+\mu_S(1-z_A)+\mu_P(0)]$$

and, since $s_{ABC}=s_A=z_{ABC}=z_A=0$:

$$N_{ABC}' = N(1-\mu_S z_{ABC*})$$

$$N_A' = N(1-\mu_S z_{A*})$$

$$N_{ABC*}' = N[1-\mu_P-z_{ABC*}(1-\mu_P-\mu_S)]$$

$$N_{A*}' = N[1-\mu_P-z_{A*}(1-\mu_P-\mu_S)]$$

602 **Table A1.** Offspring and fitness for the four plant types.

	fitness	x_1	x_2	x_3	x_4
			offspring		
x_{ABC}	$1-s_{ABC}$	$(1-\mu_S-\mu_P)(1-z_{ABC})/\alpha_P$	$\mu_P(1-z_A)/\alpha_P$	$\mu_S(1-z_{ABC*})/\alpha_P$	0
x_A	$1-s_A$	$\mu_P(1-z_{ABC})/\beta_P$	$(1-\mu_S-\mu_P)(1-z_A)/\beta_P$	0	$\mu_S(1-z_{A*})/\beta_P$
x_{ABC*}	$1-s_{ABC*}$	$\mu_S(1-z_{ABC})/\chi_P$	0	$(1-\mu_S-\mu_P)(1-z_{ABC*})/\chi_P$	0
x_{A*}	$1-s_{A*}$	0	$\mu_S(1-z_A)/\delta_P$	0	$(1-\mu_S-\mu_P)(1-z_{A*})/\delta_P$

$\alpha_P = 1-(1-\phi_P)[\mu_S z_{ABC*}]$

$\beta_P = 1-(1-\phi_P)[\mu_S z_{A*}]$

$\chi_P = 1-(1-\phi_P)[\mu_P + z_{ABC*}(1-\mu_P-\mu_S)]$

$\delta_P = 1-(1-\phi_P)[\mu_P + z_{A*}(1-\mu_P-\mu_S)]$

The effect of soft selection on offspring frequencies is that frequencies are normalized (because

we are assuming no loss of viability) after juvenile selection by dividing them by the total

frequencies of the surviving offspring (α_P, β_P, χ_P, δ_P as appropriate, see **Table 1**). Individuals

with these normalized frequencies go on to the adult phase, where another round of (hard)

selection occurs. Selection in the juvenile phase has no effect on fecundity if $z_i < 1-\phi$.

The equilibrium frequencies of the four alleles can be found by specifying the parameters

μ_S, μ_P, s_i and z_i for the system above and calculating the leading eigenvector. The situation in

which allele 1 is less robust than allele 2, is given by $s_{ABC*} > s_{A*} > 0$ and $z_{ABC*} > z_{A*} > 0$.

Bacteria

We assume that alleles *a* and *abc* can mutate to *ab* only among the alleles that can enter plants

but that the total mutation rate is the same for all alleles. Hence, *a* and *abc* also have other

mutants with frequency μ_P that produce either a password that has no match in the plant

population or a defective bacterium (and therefore zero fitness):

$abc^0 \leftarrow \mu_P \rightarrow abc* \leftarrow \mu_D \rightarrow abc \leftarrow \mu_P \rightarrow a \leftarrow \mu_D \rightarrow a* \leftarrow \mu_P \rightarrow a^0$

The recurrence equations for this system are (see **Table A2**):

$T \cdot y_{ab}' = y_{ab}(1-s_{ab})(1-2\mu_P-\mu_D)(1-z_{ab})/\alpha_B + y_a(1-s_a)\mu(1-z_{ab})/\beta_B + y_{abc}(1-s_{abc})\mu(1-z_{ab})/\chi_B + y_{ab*}(1-$

$s_{ab*})\mu(1-z_{ab})/\delta_B$

$T \cdot y_a' = y_{ab}(1-s_{ab})\,\mu(1-z_a)/\alpha_B + y_a(1-s_a)(1-2\mu_P-\mu_D)(1-z_a)/\beta_B$

$T \cdot y_{abc}' = y_{ab}(1-s_{ab})\,\mu(1-z_{abc})/\alpha_B + y_{abc}(1-s_{abc})(1-2\mu_P-\mu_D)(1-z_{abc})/\chi_B$

$T \cdot y_{ab*}' = y_{ab}(1-s_{ab})\,\mu(1-z_{ab*})/\alpha_B + y_{ab*}(1-s_{ab*})(1-2\mu_P-\mu_D)(1-z_{ab*})/\delta_B$

Where α_B, β_B, χ_B, δ_B are normalizing factors of the offspring frequencies (see Table 2) and ϕ_B

$=m/n$ is the degree of soft selection, where n is the number of offspring before soft selection

(the same for all alleles) and m ($<n$) is the maximum number of individuals that can go on to

the adult phase after soft selection. n_i' is the number of offspring of an individual with allele i

after soft selection:

$n_{ab}' = n[(1-2\mu_P-\mu_D)(1-z_{ab})+\mu_P(1-z_a)+\mu_P(1-z_{abc})+ \mu_D(1-z_{ab*})]$

$n_a' = n[(1-2\mu_P-\mu_D)(1-z_a)+ \mu_P(1-z_{ab})]$

$n_{abc}' = n[(1-2\mu_P-\mu_D)(1-z_{abc})+ \mu_P(1-z_{ab})]$

$n_{ab*}' = n[(1-2\mu_P-\mu_D)(1-z_{ab*})+ \mu_D(1-z_{ab})]$

Table A2. The offspring and fitness of the four types of bacteria

		offspring			
freq.	fitness	y_1	y_2	y_3	y_4
y_{ab}	$1-s_{ab}$	$(1-2\mu_P-\mu_D)(1-z_{ab})/\alpha_B$	$\mu_P(1-z_a)/\alpha_B$	$\mu_P(1-z_{abc})/\alpha_B$	$\mu_D(1-z_{ab*})/\alpha_B$
y_a	$1-s_a$	$\mu_P(1-z_{ab})/\beta_B$	$(1-2\mu_P-\mu_D)(1-z_a)/\beta_B$	0	–
y_{abc}	$1-s_{abc}$	$\mu_P(1-z_{ab})/\chi_B$	0	$(1-2\mu_P-\mu_D)(1-z_{abc})/\chi_B$	0
y_{ab*}	$1-s_{ab*}$	$\mu_D(1-z_{ab})/\delta_B$	0	0	$(1-2\mu_P-\mu_D)(1-z_{ab*})/\delta_B$

637     $\alpha_B = 1-(1-\phi_B)[(1-2\mu_P-\mu_D)(1-z_{abc})+\mu_P(1-z_a)+\mu_P(1-z_{abc})+ \mu_D(1-z_{ab*})]$

638     $\beta_B = 1-(1-\phi_B)[(1-2\mu_P-\mu_D)(1-z_a)+ \mu_P(1-z_{ab})]$

639     $\chi_B = 1-(1-\phi_B)[(1-2\mu_P-\mu_D)(1-z_{abc})+ \mu_P(1-z_{ab})]$

640     $\delta_B = 1-(1-\phi_B)[(1-2\mu_P-\mu_D)(1-z_{ab*})+ \mu_D(1-z_{ab})]$