Drosophila Mutants that Are Motile but Respond Poorly to All Stimuli Tested: Response by Various Organisms and a Proposal

Lar L. Vang and Julius Adler*

Departments of Biochemistry and Genetics, University of Wisconsin-Madison, Madison, Wisconsin 53706, USA

*e-mail: adler@biochem.wisc.edu

ABSTRACT

Adult Drosophila melanogaster fruit flies were placed into one end of a tube near to repellents (benzaldehyde and heat) and away from the other end containing attractants (light and a favored temperature). They escaped from the repellents and went to the attractants. Five motile mutants that failed to do that were isolated. They did not respond to any external attractants tested or external repellents tested. In addition, they did not respond well to internal sensory stimuli like hunger, thirst, and sleep. The mutants, although motile, failed to respond to stimuli at both 34°C and at room temperature. Some of the mutants have been mapped. It is proposed that the information from the different sensory receptors comes together at an intermediate, called "inbetween" (Inbet), that brings about a behavioral response. The Boss is defined here.

I. INTRODUCTION

Organisms are constantly exposed to a variety of external attractants and external repellents as well as to a variety of internal sensory stimuli. How organisms respond to these to bring about behavior is a basic question of life.

One approach for discovering how this works is the isolation and study of mutants that fail here. In this report we show that Drosophila flies can be mutated in such a way that, although still motile, they no longer respond well to any sensory stimulus tested. This includes various external attractants and various external repellents as well as internal sensory functions like hunger, thirst, and sleep. The isolation of the mutants that failed to respond was carried out at a high temperature (34°C), but they also failed to respond at room temperature. An account of some of this work has appeared (Lar Vang and Julius Adler, 2016). A preliminary report of some of the results has been presented (Adler, 2011; Vang, Alexei Medvedev, and Adler, 2012).

It is proposed that information from all the different sensory receptors comes together in the central brain at a newly found intermediate called "Inbetween" (the products of the inbet genes), which sends information to bring about a behavioral response. See Figure 1.

Figure 1. From sensing to response by way of the central brain. Mutations described in this report are located in the central brain.

Work by others has shown that in insects the central brain is a part of behaviors such as courtship (Haria Pavlou and Stephen Goodwin, 2013), audition (Jan Clemens et al., 2015), vision (Peter Weir and Michael Dickinson, 2015), and smell and taste (Leslie Voshall and Reinhard Stocker, 2007). The central brain includes the central complex, which is a system of neuropils consisting of the protocerebral bridge, the fan-shaped body, the ellipsoid body, and noduli (Ulrike Hanesch, Karl-Friedrich Fischbach, and Martin Heisenberg, 1989; J.M. Young and Douglas Armstrong, 2010; Tanya Wolf, Nirmala Iyer, and Gerald Rubin, 2015).

II. RESULTS

   A. RESPONSES TO EXTERNAL STIMULI

      1. RESPONSE TO STIMULI USED TOGETHER

In a 34°C dark room flies were started near two repellents (0.1M benzaldehyde and 37°C) at one end of a tube, away from two attractants (light at 1000 lux and 27°C) at the other end (Figure 2). The parent responded by going away from the repellents and to

Figure 2. Apparatus for isolating and testing mutants in a 34°C room. At the left end were repulsive 0.1M benzaldehyde and repulsive 37°C (due to a hot plate at 150°C). At the right end were attractive light (1000 lux) and attractive 27°C (due to ice water). The middle was close to 34°C.

the attractants (Figure 3A). Mutants that were not motile were rejected, only the motile

Figure 3. Response to stimuli used together. Repellents (0.1M benzaldehyde and high temperature (37°C) were at the left end, attractants (light, 1000 lux, and a favored temperature (27°C) at the right end. **A**, Parental response (n=7). **B**, Mutant 2 (n=8). Flies were tested in a 34°C room with 10 to 20 flies used per trial. Data are mean±SEM.

mutants were studied. This consisted of five mutants, named 1 to 5. Figure 3B shows that such a mutant failed to respond when the four stimuli were present. Each of the five mutants failed to respond when the four stimuli were present (Vang and Adler, 2016).

2. RESPONSE TO INDIVIDUAL STIMULI

A single stimulus was presented to flies that were derived from ones that had already experienced the four stimuli. For example, the parent went to light only (Figure 4A) while a mutant did not (Figure 4B). Each of the five mutants failed to

Figure 4. Response to light alone. Light (1000 lux) was placed at the right end as in Fig. 2. *A*, Parental response (n=4). *B*, Mutant 1 response (n=5). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

respond to light only (Vang and Adler, 2016).

For heat alone, the parent was repelled (Figure 5A) but the mutant was not repelled (Figure 5B). That was the case for Mutants 1 and 2 (Vang and Adler, 2016). (The other mutants were not tested for this.)

Figure 5. Response to heat gradient alone. The heat source was placed at the left end as in Figure 2. *A*, Parental response (n=4). *B*, Mutant 1 response (n=5). Flies were tested at 34°C with 10 to 20 flies per trial. The warm side measured 37°C and the cool side 27°C. Data are mean±SEM.

A similar result was found for benzaldehyde alone: the parent was repelled by benzaldehyde while Mutants 1 and 2 were not repelled. See Vang and Adler, 2016, for the figures. (The other mutants were not tested for this.)

Thus the mutants were defective not only for the four stimuli used together but also for each stimulus used alone.

3. RESPONSE TO OTHER EXTERNAL STIMULI

These mutants were in addition tested with stimuli that were not among those four used to obtain the mutants:

The mutants were tested for response to the attractant sucrose after starvation (Robert Edgecomb, Cara Harth, and Anne Schneiderman, 1994) for 17 to 20 hours. Compared to the wild-type, both Mutants 1 and 2 consumed less sucrose, about 20% as much as the wild-type. See Vang and Adler, 2016, for the figures. (The other mutants were not tested for this.)

In the case of the repellent quinine, flies were started in a 0.1M quinine half and then they had the opportunity to go into a non-quinine half (see Vang, Medvedev, and Adler, 2012, for details of the method). The parent went into the non-quinine half but Mutant 1 and Mutant 2 did not. See Vang and Adler, 2016, for the figures. (The other mutants were not tested for this.)

To test response to gravity, these flies were placed into a vertical tube and pounded down, then at every minute the flies in each third of the tube were counted (see Vang, Medvedev, and Adler, 2012, for details of the method). The parent responded by climbing up while Mutants 1 and 2 climbed up 10% as well after subtraction of movement without any added stimuli. See Vang and Adler, 2016, for the figures. (The other mutants were not tested for this.)

Thus these mutants, isolated by use of the four stimuli, were defective even for stimuli that were not present during their isolation.

4. MOVEMENT WITHOUT ANY ADDED STIMULI

In the absence of any stimulus added by the experimenters, the parent (Figure 6A) and the mutant (Figure 6B) moved similarly, indicating that motility alone is about the

Figure 6. Response without added stimuli. *A*, Parental response (n=4). *B*, Mutant 1 response (n=6). Flies were tested at 34°C with 10 to 20 flies used per trial. Data are mean±SEM.

same in parent and mutant. This was found also for Mutants 2, 3, 4 and 5 (Vang and Adler, 2016). Aside from our seeing the flies, these results tell that the mutants are motile.

5. EFFECT OF INCUBATION TEMPERATURE

All the work reported above was carried out in a 34°C room in order to allow, if necessary, isolation and study of conditional mutants, i.e. mutants defective at 34°C but not defective at room temperature. We measured response to light (1000 lux) at room temperature (21 to 23°C). The parent responded to light but all five of the mutants failed to respond to light or responded only 10% as well as the parent, just as they did at 34°C (Vang and Adler, 2016). Thus the mutations are not conditional.

Presumably these mutants are defective to all stimuli at room temperature, not just to light. Figures below show defects at room temperature for hunger, thirst, and sleep. Then how could the mutants survive and grow at room temperature? It must be that the mechanism studied here is not an essential one: flies live and reproduce without it.

B. RESPONSES TO INTERNAL STIMULI

1. HUNGER

Here we focus on hunger (Edgecomb *et al.,* 1994; Melche *et al.,* 2007; Fujikawa *et al.,* 2009; Farhadian *et al.,* 2012; Hong *et al*., 2012; Itskov and Ribeiro, 2012). To measure hunger we used an apparatus (Fig. 7), inspired and modified from

Figure 7. Apparatus for measuring hunger and for measuring thirst. For details see (Vang and Adler, 2016). Tube 1 is called "origin". Flies were tested at room temperature (21-23°C) for up to 40 hours.

an earlier design (Brown *et al.*, 1960), that we described (Vang and Adler, 2016).

Briefly, in a dark room at 21-23°C male flies – parent or mutants - were transferred into one end (tube 1) of a 5 x 140 cm apparatus containing throughout its length a 5 cm wide strip of wet paper to satisfy thirst but containing no food. Starvation for food began once the flies were put in. Every 10 hours the location of the flies was measured with light on for a few seconds.

At 20 hours the parent had largely left the origin (tube 1) and had begun to accumulate at the end (tube 4) (Figure 8A, solid bars), while the mutant had moved

Figure 8. Movement of flies at 20 hours in search for food. Solid: water but no food (no sucrose). Open: water and food (0.1M sucrose). *A*, Parental response with water only (n=5) and with water + sucrose (n=9). *B*, Mutant 2 response with water only (n=5) and with water + sucrose (n=4). Data are mean±SEM. See (Vang and Adler, 2016) for Mutant 1; the other mutants were not tested for this. Flies were tested at room temperature (21-23°C) with 40 to 60 flies used per trial.

towards the end very little (Figure 8B, solid bars). This is interpreted to mean that the parent is searching for food while the mutant is defective in searching for food.

When food (0.1M sucrose) was added throughout the tube along with the wet strip of paper, the parent moved less far (rather than accumulating at the end) (Figure 8A open bars), while the mutant remained mostly where placed (Figure 8B, open bars). Since sucrose inhibited the movement of the parent, it is supposed that movement without sucrose is due largely to hunger. From these results we conclude that the mutants are defective in hunger.

2. THIRST

To study thirst, flies were deprived of water. The procedure is the same as for hunger except that water was omitted and solid sucrose was layered throughout (Vang and Adler, 2016). Mutants 1 and 2 were tested, the other mutants not (Vang and Adler, 2016).

By 30 hours the parent had moved out, presumably to search for water since addition of water inhibited this (Vang and Adler, 2016). The mutant moved out less well than the parent (Vang and Adler, 2016), so we conclude that the mutants are defective in thirst.

3. SLEEP-WAKE

The parent and mutants isolated here were studied for sleep and wake according to the procedure of Cory Pfeiffenberger *et al*., 2010. The parent was different from the mutants (Figure 9). The parent showed greatest activity at the start and end of the day but

Figure 9. Circadian response. Individual flies are placed into a tube (5 x 20 mm) with an infrared light beam intersecting at the middle of the tube. Mutant 1 (n=24), Mutant 2 (n=24), and parental response (n=24) are recorded over a 24 hour period at

22° C. Data are mean±SEM. There are smaller differences between the parent and the three other mutants (Vang and Adler, 2016).

not in the middle of the day. Mutant 2 showed high activity throughout the day. Mutant 1 was less active than the parent at the start of the day.

C. MAPPING OF THE MUTANTS

We found that Mutant 1 maps in a small gap between 12E3 and 12E5 on the X-chromosome (Vang and Adler, 2016) which we call "*inbetween A*" (*inbetA*). We found that Mutant 2 maps in the *CG1791* gene, a part of the fibrinogen gene of the X-chromosome, or next to it (Vang and Adler, 2016), which we call "*inbetween B*" (*inbetB*).

III. DISCUSSION

Here we describe the isolation and some properties of *Drosophila* mutants that are motile but yet they each fail in response to all external attractants and repellents tested (Figures 3-5) and also they are deficient in response to internal stimuli tested (Figures 8 and 9). Thus, although the mutants are motile, they have:

decreased responsiveness to light
decreased responsiveness to heat and to favorable temperature
decreased responsiveness to repulsive chemicals (like benzaldehyde)
decreased responsiveness to sweet tastants (like sucrose)
decreased responsiveness to bitter tastants (like quinine)
decreased responsiveness to gravity
decreased responsiveness to hunger
decreased responsiveness to thirst
abnormality in some sleep

Because all of these different behaviors are defective in the mutants, it seems reasonable to say that there is a single place that is responsible, rather than a defect in each of the many different sensory receptors. One possibility for this place is the interaction between sensory receptors and processing of sensory stimuli (Vang and Adler, 2016). Another possibility is that this place is in the central brain (Figure 1) between sensing and response, where all the sensory information comes together at a newly discovered place we call "Inbetween", the proteins of the *inbetA* and *inbetB* genes.

The mechanism and function of these Inbetween proteins need to be determined further.

An analogy can be made between *Drosophila* mutants missing the part of the central brain studied here and *Escherichia coli* mutants missing their chemotaxis mechanism (John Armstrong, Adler, and Margaret Dahl, 1967; Adler, 1969; John S. Parkinson, 1976).

A further analogy can be made with humans. Since the patient's fall off his bicycle at the age of 10, he had severe epileptic seizures. William Scoville performed experimental surgery to remove the brain's temporal lobes at the patient's age of 27. After the operation the seizures reduced drastically but the patient now suffered from amnesia until death at 84. Brenda Milner, a student of Scoville's, studied the patient: he failed to remember all stimuli (Scoville and Milner, 1957). The patient's brain after death was examined by histological sectioning, which revealed affected parts in the medial temporal lobes and to a small degree in the orbitofrontal cortex (Jacobo Annese and coauthors, 2014).

It would appear that all three organisms – bacteria, flies, and people – may have a related mechanism for the path from sensory stimuli to behavioral responses.

In each of these cases, when the behavior was absent the organism was still motile and it still reproduced: thus this behavior is not essential to life, though of course the defective organism must be severely handicapped.

The path from sensory stimuli to behavioral responses in vertebrates and flies has been compared by Nicholas Strausfeld and Frank Hirth (2013); they reported that the basal ganglia of mammals and the central complex of insects have multiple similarities and they suggest deep homology between them, see Figure 10. Origin of the cerebral cortex has

After Strausfeld and Hirth, Science 340:157 (2013)

Figure 10. Comparison of the basal ganglia of vertebrates and the central complex of insects according to modification of Fig. 2 of Strausfeld and Hirth (2013). The green arrows lead to the response.

been studied by use of an ancient ancestor by Dedlev Arendt and coauthors (2010) and Heather Marlow and Arendt (2014).

Starting in the 1870's it became apparent to some psychologists that there is a part of the mammalian brain, the frontal cortex, that is master of the whole brain; see reviews up to 1970 by the neurophysiologist Aleksandr Luria, who himself modernized this concept and studied syndromes resulting from deficiencies of the frontal cortex (Luria, 1973, 1980). This part of the brain became known as the "central executive" through the research of the psychologist Alan Baddeley (1966) or as the "executive brain" through the research of the neuropsychologist Elkhonon Goldberg (2001), a student of Luria's. It is now known as "executive control" or "executive function" as well as "prefrontal cortex". For reviews see Goldberg and Dmitri Bougakov (2007), Joaquin Fuster (2008), Sam Gilbert and Paul Burgess (2008), Alfredo Ardilla (2008), and Hyun Chung, Lisa Weyandt, and Anthony Swentosky (2014).

Helen Barbas reported (2000, 2013), as described in Figure 11, that the orbitofrontal

Figure 11. The orbitofrontal cortex receives information from sensory cortices and also from the amygdala's recording of past events and emotions and also (not shown here) from the thalamus, as described in Fig. 3 of Barbas (2000). For more detail, see Figs. 41.14 and 41.31 of Barbas (2013).

cortex of humans and other primates receives information from the sensory cortices (which contain visual, auditory, somatosensory, gustatory, and olfactory data as it is received) and also from the amygdala (which contains data about memory and emotion). Barbas suggested that "the orbitofrontal cortex is thus capable of sampling the entire external and internal environment and may act as an environmental integrator" (2000). This is presented in more detail by Barbas in 2013: "The extensive connections suggest that the posterior part of the orbitofrontal cortex has a panoramic view of the entire sensory world, including our internal environment of motives and

drives" (Fig. 41.16 and p.23 of 2013). After receiving all this information, the prefrontal cortex decides what action should be brought about (Barbas, 2013). Then the prefrontal cortex informs the premotor cortex (Barbas, 2013), whose function is to produce action. The prefrontal cortex has been reported in rodents (Verity Brown and Eric Bowman, 2002; Harry Uylings, Henk Groenwegen, and Bryan Kolb, 2003).

It is suggested here that the excitatory pathways and the inhibitory pathways of the prefrontal cortex (Figures 41.14 and 41.31 of Barbas, 2013) are together to be called "The Boss". The Boss is in charge of the prefrontal cortex, The Boss controls the prefrontal cortex. The Boss is an idea: there is at this time no evidence for it. The Boss is a chemical (or chemicals) that occurs in the organism. It is proposed that The Boss is to be found throughout biology, in animals, plants, and microorganisms, and that The Boss directs each organism (Adler, 2011, "My life with nature", Figure 13 and page 60; Adler, 2016, "A search for The Boss: The thing inside each organism that is in charge"). The idea presented here is that every organism has something in control of it. This differs from earlier work.

The control of genes by The Boss could be studied by finding mutants that are in it. Such mutants would most likely be missing the response to all stimuli at a higher temperature but would be normal at room temperature. In the present report Boss mutants did not show up: the mutants described failed at 34°C but they also failed at room temperature. However, only five mutants were studied here; possibly if a much larger number of mutants were isolated there would be some among them that are defective at 34°C but normally responsive at room temperature.

Decision making has been studied in animals, plants, and microorganisms. Decision making (also called "choice behavior" and "cross-modal object recognition") occurs in primates and rodents, as described by Jacob Cloke, Derek Jacklin, and Boyer Winters (2015). For a review of decision making in mice, *Drosophila*, and *Caenorhabditis elegans* see Nilay Yapici, Manuel Zimmer, and Ana Domingos (2014); about *Drosophila* they say, "Nutrient sensing that guides decision-making has also been studied in *Drosophila* and found to be regulated at the central nervous system level…These neurons are located in the lateral protocerebrum". In *Drosophila* integration of conflicting sensory information (repellent CO_2 plus attractant vinegar) requires a higher brain circuit identified as the mushroom body, according to Laurence Lewis, Gerald Rubin, Ilona Kadow, and coworkers (2015). *Drosophila* mutants were isolated by Adler and Vang (2016) that fail in making a decision (repellent methyl eugenol plus attractant light) at 34°C but the mutants respond normally to attractant alone and repellent alone at 34°C and they succeed in making a decision at room temperature, so it seems that The Boss could be in control of decision making. This was proposed by Adler in 2011 ("My life with nature", Fig. 13, page 60).

Compare decision making in *Drosophila* (Adler and Vang, 2016) with decision making in *Salmonella* or *E. coli* (Nora Tsang, Robert Macnab, and Daniel Koshland, Jr 1973; Adler and Wung-Wai Tso, 1974; Ian Strauss *et al.,* 1995; Yevgeniy Kalinin *et al*., 2010). In both flies and bacteria the organism is of course repelled when repellent exceeds attractant. Mutants in decision making in *Drosophila* have been isolated (Figure 3, Adler and Vang, 2016) but it remains to isolate mutants in decision making in bacteria. Such mutants would allow

comparison of the mechanism of decision making in animals and in bacteria. That search might tell about The Boss in bacteria.

The Boss would be in control of motivation. "Primary motives are thought to include hunger, thirst, sex, avoidance of pain, and perhaps aggression and fear" according to Charles Cofer and Herbert Petri (2017). "Motivation is a theoretical construct used to explain behavior. It gives the reason for people's action, desires, and needs" says Wikipedia (2017). For additional reports about motivation see Vincent Dethier (1976), and Paul Kleinginna, Jr and Anne Kleinginna (1981).

Motivation directs behavior, so by being in control of motivation, The Boss would become in control of behavior. Examples of behavior are eating, drinking, smelling, tasting, seeing, thermosensing, humidity sensing, and gravity sensing, as reviewed by Vang, Medvedev, and Adler (2012) and references cited there.

Thus The Boss would control decision making, motivation, and behavior. Additional functions of The Boss are control of synthesis and function of DNA, RNA, and proteins and control of ATP level, see Figs. 10 and 11 of Adler (2016).

What is The Boss? Of course it must be a gene or genes because it is inherited. The genes for The Boss would seem to be made of DNA, but not necessarily since RNA may have been present in organisms before there was any DNA according to ideas of Carl Woese (1968), David Baltimore (1980), Walter Gilbert (1986), Raymond Gesteland and John Atkins (1993), and Gerald Joyce (2002). RNA viruses and RNA-dependent RNA polymerases must have played a vital role early in evolution, when RNA was the primary genetic material, says Paul Ahlquist (2002). Some cells lacking viruses use RNA instead of DNA in roles that possibly occurred early, according to Lakshminarayan Iyer, Eugene Koonin, and Iyer Aravind (2003). See also Andrew Fire and Craig Mello ((1998) and see RNA-dependent RNA polymerase (Wikipedia, 2017). Since The Boss may already have occurred in the earliest organisms, The Boss genes might well be RNA that does not go through any DNA at all.

IV. OVERVIEW

Much of the above Discussion is put together in Figure 12. There are sensory inputs, both

Figure 12. Where decision making, final pathway, and behavioral response act in organisms in general. Decision making is led by executive function, which is led by The Boss.

for external and internal stimuli. Then the organism chooses what to do about these by means of decision making. That result is then sent to the final pathway, which dictates a behavioral response. Decision making is directed by executive function which in turn is controlled by The Boss.

V. METHODS

Details of methods used here are found in in the previous paper (Vang and Adler, 2016): A. Isolation of mutants. B. How to study response to external stimuli. C. How to study response to internal stimuli.

ACKNOWLEDGEMENTS

Julius Adler is grateful to the Camille and Henry Dreyfus Foundation for six years of grants in support of 31 undergraduate research students named in Adler and Vang, 2016, and in Vang and Adler, 2016. Lar Vang is currently associate research specialist in the Adler laboratory. Robert A. Kreber, a research specialist in Barry Ganetzky's laboratory, has helped us in studies of the genetics of our mutants. We are grateful to Erin Gonzales and Jerry Yin for showing us how to use the *Drosophila* activity monitory system. Julius Adler thanks Barry Ganetzky for teaching him about fruit flies. Thanks to Millard Susman for criticism. We are thankful to Laura Vanderploeg for the art work.

BIBLIOGRAPHY

Adler J (2011) My life with nature. Annu Rev Biochem. 80:42–70

Adler J (2016) A search for The Boss: The thing inside each organism that is in charge. Anatomy Physiol Biochem Int J. Vol. 1, July 2016.

Adler J, Tso W-W (1974) "Decision"-making in bacteria: chemotactic response of Escherichia coli to conflicting stimuli. Science 184:1292-1294.

Adler J, Vang LL (2016) Decision making By *Drosophila* flies. bioRxiv, doi: https://doi.org/10.1101/045666

Ahlquist P (2002) RNA-dependent RNA polymerases, viruses, and RNA silencing. Science 296:1270-1273.

Annese J, Shenker-Ahmed NM, Bartsch H *et al.* (2014) Postmortem examination of patient H.M.'s brain based on histological sectioning and digit 3D reconstruction. Nature Comm 5:1-23.

Ardila A. (2008) On the evolutionary origins of executive functions. Brain Cogn 68:92-99.

Armstrong JB, Adler J, Dahl MM (1967) Nonchemotactic mutants of *Escherichia coli*. J Bacteriol 93:390-398.

Baddeley A (1966) Exploring the central executive. Quar J Expl Psychol 49A(1):5-28.

Baltimore D (1980) Evolution of RNA viruses. Ann N Y Acad Sci 354:492-497.

Barbas H (2000) Connections underlying the synthesis of cognition, memory, and emotion in primate prefrontal cortices. Brain Res Bull 52:319-330.

Barbas H (2013) Frontal cortex. Chapter 41, pp. 1289-1334. *Neuroscience in the 21st Century*, ed. Donald W. Pfaff, SpringerLink.

Brown VJ, Bowman EM (2002) Rodent models of prefrontal cortical function. Trends Neurosci 25:340-343.

Browne LB, Evans DR (1960) Locomotor activity of the blowfly as a function of feeding and starvation. J Insect Physiol 4:27–37.

Chung HJ, Weyandt LL, Swentosky A (2014) The physiology of executive functioning. pp. 13-27, in Goldstein S, Naglieri JA, *Handbook of Executive Functioning*, SpringerLink, Springer Science.

Clemens J, Giradin CC, Coen P, Guan X-J, Dickson BJ, Murthy M (2015) Connecting neural codes with behavior in the auditory system of *Drosophila*. Neuron 87:1332-1343.

Cofer CN, Petri HL (2017) Motivation behavior. Encyclopedia Britannica. 1-30, 1-18, 1-8.

Dethier VG (1976) *The Hungry Fly*, Chapter 10, The fly and the concept of motivation. Harvard University Press.

Edgecomb RS, Harth CE, Schneiderman AM (1994) Regulation of feeding behavior in adult *Drosophila melanogaster* varies with feeding regime and nutritional state. J Exp Biol 197:215-35.

Farhadian SF, Suárez-Fariñas M, Cho CE, Pellegrino M, Vosshall LB (2012) Post-fasting olfactory, transcriptional, and feeding responses in *Drosophila.* Physiol Behav 105:544–53.

Fire A, Xu SQ, Montgomery MK, Kostas SA, Driver SE, Mello CC (1998) Potent and specific genetic interference by double-stranded RNA in *Caenorhabditis elegance*. Nature 391:806-811.

Fujikawa K, Takahashi A, Nishimura A, Itoh M, Takano-Shimizu T, Ozaki M (2009) Characteristics of genes up-regulated and down-regulated after 24 h starvation in the head of *Drosophila*. Gene 446:11–17.

Fuster JM. IV. Executive function. In: *The Prefrontal Cortex*, Fourth Edition. London: Academic Press, 2008. pp. 178-193.

Gallistel CR (1975) Motivation as central organizing process: The psychophysical approach to its functional and neurophysiological analysis. Nebraska Symposium on Motivation, 1974, University of Nebraska Press, 22:185-225.

Gestiland RF, Atkins JF, editors (1993) *The RNA World, the Nature of Modern RNA Suggests a Prebiotic RNA World*. Cold Spring Harb Lab Press.

Gilbert W (1986) Origin of life, the RNA world. Nature 319:618.

Gilbert SJ, Burgess PW (2008) Executive function. Curr Biol 18(3):R110-R114.

Goldberg E (2001) *The Executive Brain, Frontal Lobes and the Civilized Mind*. Oxford University Press, USA.

Goldberg EM, Bougakov DH (2007) Goals, executive control, and action. Chapter 12 In: Baars BJ and Gage NM (Eds) *Cognition, Brain, and Consciousness: Introduction to Cognitive Neuroscience*, Elsevier, pp. 343-368.

Hanesch U, Fischbach K-F, Heisenberg M (1989) Neuronal architecture of the central complex in *Drosophila melanogaster*. Cell Tissue Res 257:343-366.

Hong S-H, Lee K-S, Kwak S-J, Kim A-K, Bai H, Jung M-S, Kwon OY, Song W-J, Tatar M, and Yu K (2012) Minibrain/Dyrk1a regulates food intake through the Sir2-FOXO-sNPF/NPY pathway in *Drosophila* and mammals. PLoS Genet 8:e1002857. doi: 10.1371/annotation/8c2c8644-1beb-4410-8bef-c388b7256738.

Itskov PM, Ribeiro C (2013) The dilemmas of the gourmet fly: the molecular and neuronal mechanisms of feeding and nutrient decision making in *Drosophila*. Front Neurosci 7:1–7. doi: 10.3389/fnins.2013.00012.

Iyer LM, Koonin EV, Aravind L (2003) Evolutionary connection between the catalytic subunits of DNA-dependent RNA polymerases and eukaryotic RNA-dependent RNA polymerases and the origin of RNA polymerases. BMC Struct Biol 3:1.

Joyce GF (2002) The antiquity of RNA-based evolution. Nature 418:214-221.

Kalinin Y, Neumann S, Sourjik V, Wu, M. (2010) Responses of *Escherichia coli* bacteria to two opposing chemoattractant gradients depend on the chemoreceptor ratio. J Bacteriol 192:1796-1800.

Kleinginna PR Jr, Kleinginna AM (1981) A categorized list of motivational definitions, with a suggestion for a consensual definition. Motiv Emot 5:263-291.

Lewis LPC, Sigu KP, Aso Y, Friedrich AB, Bulteel AJB, Rubin GM, Kadow ICG (2015) A higher brain circuit for immediate integration of conflicting sensory information in *Drosophila*. Curr Biol 25:2202-2214.

Luria AR (1973) *The Working Brain, an Introduction to Neuropsychology*, translated by Basil Haigh, Penguin Books Ltd., pp. 79-99, 187-225.

Luria AR (1980) *Higher Cortical Functions in Man*, second American edition, translated by Basil Haigh, pp. 246-365, Basic Books Inc.

Marlow H, Arent D (2014) Evolution: Ctenophore genomes and the origin of neurons. Curr Biol 24:R757-R761.

Melche C, Bader R, Pankratz MJ (2007) Amino acids, taste cirquits, and feeding behavior in *Drosophila*: towards understanding the psychology of feeding in flies and man. J Endocrinol 192:467-462.

Parker MO, Brock AS, Walton RT, Brennan CH (2013) The role of zebrafish (*Danio verio*) in dissecting the genetics and neural circuits of executive function. Front Neural Circuits 7:63-81.

Parkinson JS (1976) cheA, cheB, and cheC genes of *Escherichia coli* and their role in chemotaxis. J Bacteriol 126:758-770.

Pavlou HJ, Goodwin SF (2013) Courtship behavior in *Drosophila melanogaster*: towards a 'courtship connectome'. Curr Opin Neorobiol 23:76-83.

Pfeiffenberger C, Lear BC, Keagan KP, Allada R (2010) Processing circadian data collected from the *Drosophila* activity monitoring (DAM) system. Cold Spring Harb Protoc 2010:1242–1250.

Scoville WB, Milner B (1957) Loss of recent memory after bilateral hippocampal lesions. J Neurol Neurosurg Psychiat 20:11-21.

Strausfeld NJ, Hirth F (2013) Deep homology of arthropod central complex and vertebrate basal ganglia. Science 340:157-161.

Strauss I, Frymier PD, Hahn CM, Ford RM (1995) Analysis of bacterial migration: II. Studies with multiple attractant gradients. Bioeng Food Nat Prod 41:402-414.

Tomer R, Denes AS, Tessmar-Raible,K, Arendt D (2010) Profiling by image registration reveals common origin of annelid mushroom bodies and vertebrate pallium. Cell 142:800-809.

Tsang N, Macnab R, Koshland DE Jr (1973) Common mechanism for repellents and attractants in bacterial chemotaxis. Science 181:60-63.

Uylings HBM, Groenewegen HJ, Kolb B (2003) Do rats have a prefrontal cortex? Behav Brain Res 146:3-17.

Vang LL, Medvedev AV, Adler J (2012) Simple ways to measure behavioral responses of *Drosophila* to stimuli and use of these methods to characterize a novel mutant. PLoS ONE 7:e37495. doi: http://dx.doi.org/10.1371/journal.pone.0037495.

Vang LL, Adler J (2016) *Drosophila* mutants that are motile but respond poorly to all stimuli tested: These may have a defect in interaction with stimuli or in executive function. bioRxiv. doi: http://dx.doi.org/10.1101/045062. April 13, 2016, see also March 21, 2016.

Vosshall LB, Stocker RF (2007) Molecular architecture of smell and taste in *Drosophila*. Annu Rev Neurosci 30:505-533.

Weir PT, Dickinson MH (2015) Functional divisions for visual processing in the central brain of flying *Drosophila*. Proc Natl Acad Sci USA 112:E5523-E5532.

Wikipedia (2017) Executive functions.

Wikipedia (2017) Motivation.

Wikipedia (2017) RNA-dependent RNA polymerse

Woese C (1968) *The Genetic Code*. Harper & Row, pp.179-195.

Wolff T, Iyer NA, Rubin GM (2015) Neuroarchitecture and anatomy of the *Drosophila* central complex: A GAL4-based dissection of protocerebral bridge neurons and circuits. J Comp Neurol 523:997-1037.

Young JM, Armstrong JD (2010) Structure of the adult central complex in *Drosophila*: Organization of distinct neuronal subsets. J Comp Neurol 518:1500-1524.