Cohort Specific Effects of Cereal-bar Supplementation in Overweight Patients

With or Without Type 2 Diabetes Mellitus

Short Title: Cohort specific effects of diet supplementation

Christian L. Lauber [1*], Chieh Jason Chou[1], Anirikh Chakrabarti[1], Jay Siddharth[1],

Anne Chalut-Carpentier[2], Zoltan Pataky[2], Alain Golay[2¶], Scott Parkinson[1¶]

[1]Nestlé Institute of Health Sciences S.A., Lausanne, Switzerland
12

13   [2] Service of Therapeutic Education for Chronic Diseases, WHO Collaborating

14   Centre, University Hospitals of Geneva and University of Geneva, Geneva,

15   Switzerland

21   ¶ Senior authors

23   * Corresponding author. Nestlé Institute of Health Sciences, SA. École

24   Polytechnique Fédérale de Lausanne, Bâtiment H, 1015, Lausanne, Switzerland.

25   Email: christianlynn.lauber@rd.nestle.com.

Abstract

The importance of gut microbes to metabolic health is becoming more evident and nutrition-based therapies to alter the composition of bacterial communities to manage metabolic disease are an attractive avenue to ameliorate some effects of Western diets. While the composition of gut microbial communities can vary significantly across disease states, it is not well known if these communities have common responses to nutritional interventions. To better understand diet-bacterial community interactions, we collected biological parameters and fecal samples of overweight non-diabetic (OND) and diabetic (OD) individuals before and after daily supplementation of 2.8 g β-glucan on their habitual diet for 30 days. Fecal bacterial communities in an age-matched cohort were measured by sequencing partial 16S rRNA genes and imputed metagenomic content. Unexpectedly, we observed disconnected responses of biological measurements and the bacterial community. Based on average effect size, biological measurements were greater in the OND group while effects on the bacterial community were greatest on the OD cohort, and we suspect these observations are due to the significantly lower alpha diversity in the OD cohort. Our data indicate that responses to cereal-bar supplementation are cohort specific and this should be considered when manipulating the microbiome via diet supplementation.

Introduction

As a result of a globalization of the economy, obesity and diabetes are becoming major issues not just for the developed but the developing world. The

global adoption of a Western lifestyle includes dietary modification to diets low

in fiber and high energy density. Deaths accounted for by non-communicable

diseases have now overtaken deaths due to communicable diseases virtually

world wide [1]. This statistic highlights the dilemma for providing high quality

foods and avoiding over nutrition for both the developed and developing world.

The gut bacterial community serves as one conduit by which the

consequences of nutrition and dietary choices integrate into human health, and

therefore must be considered in the context of any nutritional intervention.

Recent work shows abundances of dominant taxa in the distal gut vary widely

across individuals and habitual food choices [2-6]. As Firmicutes and

Bacteroidetes are by far the most prevalent taxa in the human gastro-intestinal

tract [7], their presence and response to diet can have a significant impact on

human health. For instance, members of the *Fecalbacterium* (a firmicute)

produce short-chain fatty acids such as butyrate [8] that serve as an energy

source for colonic epithelia. *Bacteroides and Ruminococcaceae* (both

Bacteroidetes) degrade polysaccharides found in foods containing fiber [9, 10].

Conversely, abundant taxa such as the Proteobacteria are typically associated

with dysbiosis observed in persons with inflammatory bowel disorders [11, 12]

or in the guts of those consuming a high fat or typical Western style diet [2].

Despite the inter- and intra-individual variability in the abundance of bacterial

taxa, these and other studies demonstrate the importance of the bacterial

communities to integrate diet and host metabolism. Dietary supplements aiming

to leverage the benefits of microbial function therefore may become practical

solutions to Western lifestyle associated diseases if their efficacy and the

molecular mechanisms underlying the bacteria-diet interactions could be

demonstrated.

Diet supplemenation is one approach proposed to minimize the effects of

Western life styles by potentially altering the composition and metabolism of the

gut bacterial community [13, 14]. However it is not widely known if a diet

supplement has the same effect on microbiomes in subjects of related but

distinct disease states. For instance, there are numerous studies that document

fiber induced alterations in the microbiomes of monotonic cohorts [15-19].

Hooda et al. [20] demonstrated the significant effect of corn derived fibers on

abundances of genera of belonging to Firmicutes, Bacteroidetes and

Verrucomicrobia in generally healthy subjects, while Benus et al. [21] further

demonstrated that pea fiber and fructo-oligosaccharides supplementation was

associated with increased abundances of butyrate producing bacteria in healthy

subjects. Though these studies have given us great insight into the effect of fiber

on the microbiome, each used unique diet amendments in cohorts that consisted

almost exclusively of healthy, overweight, obese, or subjects with diagnosed

metabolic disease. Given that bacterial community composition and diversity

vary significantly across disease states, it is reasonable to assume that effects in

one cohort may not be applicable to others. This lack of knowledge may be

important when identifying patients whose microbiomes may be more or less

responsive to diet supplementation and thus resistant to improvement of host

metabolic markers potentially mediated by gut bacteria. Thus quantifying the

responsiveness of the microbiome in similar but distinctive cohorts may help to

define the limits of diet supplementation via changes in the microbiome.

Overweight diabetic (OD) and overweight non-diabetics (OND) represent

two metabolic states that can be distinguished by the degree of insulin

resistance. However, it is not known whether the two populations exhibit cohort

specific effects to diet supplementation on the bacterial communities and host

metabolism. To answer this question, we conducted a diet supplementation

study in OD and OND subjects with a cereal-bar containing 1.4 g β-glucan. We

investigated how the bacterial communities responded to consumption of two

cereal-bars per day and determined whether there were unique features of the

response in OND vs OD subjects. Specifically we predicted that cereal-bar

supplementation would increase abundances of Bacteroidetes and

Verrucomicrobia while decreasing Firmicutes and Proteobacteria in both

cohorts with concomitant improvement in host metabolic markers.

Results

Biological Parameters

We recruited 46 overweight and obese individuals with and without

diabetes to assess the effect of a β-glucan containing cereal-bar supplementation

on biological parameters and the distal gut bacterial community. The subject

characteristics are summarized in S1 Table. Subjects went through a diet

normalization phase followed by consumption of 2 cereal-bars per day for 30

days to deliver 2.8g β-glucan fiber to the habitual diet. Twenty-six individuals

were excluded for compliance issues, lack of sequencing success, or fell outside

the age-range, leaving 10 individuals in each group for analysis (S5, S6). Non-

parametric testing indicated no significant difference in any biological

measurement between the cohorts at the pre-supplementation time point.

However, LDL and triglyceride concentrations were statistically different

between the OND and OD post supplementation (Table 1, S2 Table), though

these differences were not significant improvements from their pre-

supplementation values. The remaining biological parameters were unaffected

by consumption of the cereal-bar. Lastly, Kruskal-Wallis tests showed there were

no differences within each cohort between pre- and post-supplementation time

points (Table1).

Table 1. Biological parameters of age-matched cohorts at enrollment, pre-

supplementation, and post-supplementation. The mean and standard

deviation (in parentheses) are shown. Except for *Age and *Height, the rest of

parameters were included in biological effect size calculation. Superscript "a"

indicates significant difference between cohorts post-supplementation.

	OND Mean			OD Mean		
	Enrollment	Pre-supplementation	Post-supplementation	Enrollment	Pre-supplementation	Post-supplementation
*Age	48 (3.5)	48 (3.5)	48 (3.5)	47 (4.3)	47 (4.3)	47 (4.3)
*Height (cm)	167 (10.3)	167 (10.3)	167 (10.3)	171 (10)	171 (10)	171 (10)
Weight (kg)	95 (14.7)	95 (14.1)	94 (14.4)	98 (17)	96 (17)	96 (16)
BMI (kg m^2)	34 (5.2)	34 (5.2)	34 (5.3)	34 (4.66)	33 (4.5)	33 (4.4)
Waist Circumference (cm)	104 (11.4)	105 (11.3)	106 (11.3)	113 (10)	113 (10)	113 (12)
Hip Circumfrenece (cm)	117 (8.1)	117 (8.3)	116 (8.2)	116 (12)	114 (11)	115 (12)
Diastolic BP (mmHg)	95 (16)	87 (5.1)	89 (13)	87 (9.86)	85 (9.63)	88 (14)
Systolic BP (mmHg)	131 (15)	124 (13)	131 (21)	130 (17)	129 (22)	131 (17)
Cardiac Frequency (beats min^{-1})	67 (11)	68 (12)	72 (16)	79 (24)	78 (13)	77 (14)
Total Cholesterol (mmol L^{-1})	5 (0.74)	4.9 (0.87)	5.2 (0.69)	4.9 (1.3)	4.6 (0.82)	4.7 (0.49)
Triglyceride (mmol L^{-1})	1.1 (0.42)	1.2 (0.51)	1.2 (0.43)	4.1 (5.8)	3.6 (4.3)	2.9 (2.1)
HDL cholesterol (mmol L^{-1})	1 (0.23)	1 (0.3)	1.1 (0.25)	0.9 (0.24)	0.8 (0.24)	0.8 (0.16)
LDL (mmol L^{-1})	3.5 (0.61)	3.4 (0.68)	3.6 (0.65)[a]	2.8 (0.74)	2.7 (0.7)	2.7 (0.67)[a]
Glucose (mmol L^{-1})	5 (0.68)	5 (0.63)	5.1 (0.46)[a]	6.4 (2)	6.5 (2.2)	6.3 (1.5)[a]
Insulin (uIU ml^{-1})	17 (7.1)	19 (8.1)	19 (7.6)	20 (13)	17 (10)	18 (12)
FFA (mmol L^{-1})	0.5 (0.13)	0.5 (0.2)	0.6 (0.23)	0.6 (0.25)	0.5 (0.19)	0.5 (0.19)
HOMA_IR	3.9 (1.8)	4.4 (2.1)	4.3 (1.6)	6.2 (5.8)	5.1 (3.7)	5.5 (5)
LBP (ug ml^{-1})	28 (6)	27 (3.8)	29 (4.8)	31 (8.7)	27 (7.4)	29 (7.7)

Taxon abundance and KEGG functions pre- and post-supplementation

Bacterial communities in each cohort were dominated by Firmicutes and

Bacteroidetes, followed by the Actinobacteria, Proteobacteria and

Verrucomicrobia (Fig 1A-1D, S3 Table). Abundances at the pre-supplementation

time point were, however, not statistically different between OND and OD groups

(Fig 1A-1D). Likewise the abundances of KEGG functions from predicted

metagenomes were not different between the cohorts pre-supplementation (S4

Table). However, we did observe a significant difference in alpha diversity

between OND and OD groups at enrollment and before the intervention (Fig 1E,

S3 Table). The OND cohorts had significantly greater community richness

compared to the OD group (Fig 1E, S3 Table).

Fig 1. **Proportional abundances of dominant taxa and Shannon diversity at**

enrollment (EN), pre-supplementation (Pre) and post-supplementation

(Post). Bar graphs for proportional abundances of dominant taxa are indicated

as (A) Firmicutes, (B) Bacteroidetes, (C) Proteobacteria, and (D)

Verrucomicrobia. (E) Shannon diversity. Filled bars represent the OND cohort,

open bars represent the OD group. Error bars depict the standard deviation for

the mean of 10 subjects. p-values for significant differences are shown.

Post-supplementation, we observed a decline in Firmicutes (Fig 1A) with

simultaneous increase in Bacteroidetes (Fig 1B) abundances in the OD cohort.

However, only the Firmicutes were statistically different between the cohorts

even though the Bacteroidetes increased by 12% over the pre-supplementation

abundance in the OD group (S3 Table). The remaining taxon abundances were

not significantly impacted by the supplementation but we did observe that the

Proteobacteria (Fig 1C) consistently decreased, while the Verrucomicrobia (Fig

1D) consistently increased in abundance after supplementation. The abundance

of individual KEGG functions were not significantly altered within or between

cohorts post-supplementation (S3 Table). Alpha diversity (Shannon Index)

remained significantly higher in the OND cohort at the last time point (Fig 1E).

Changes in effect size pre- and post-supplementation

173   We next examined the response to the β-glucan cereal-bar

supplementation by comparing average effect size across the biological

parameters, phyla, and predicted metagenomic content within each cohort (Fig

2, S2- S4 Tables). Firstly, effect size was not significantly different after

supplementation in the OND cohort when compared to pre-supplementation

abundances (p > 0.3 in all cases, Fig 2A-2C). Mean effect size was largest for the

phyla (mean 0.26, range 0.04-0.62, Fig 2B), followed by the biological (mean

0.20, range 0.04-0.42, Fig 2A) and predicted metagenome content (mean 0.11,

range 0.00-0.37, Fig 2C). In contrast, β-glucan supplementation had a significant

effect in the OD cohort for all data categories (Fig 2A-C). Effects for the phyla and

predicted metagenome content were significantly greater after supplementation

while (p < 0.03, Fig 2B) while physiology effect size was significantly greater pre-

supplementation for the OD (p = 0.04, Fig 2A, S3 Table).

Fig 2. **Effect size of cereal-bar supplementation on measured parameters.**

Box and whisker plots showing the mean, the minimum and maximum values for

(A) biological, (B) phyla, and (C) predicted metagenome. Kruskal-Wallis

comparisons with p-values are shown. EN-Pre = effect size for enrollment and

pre-supplementation time points; Pre-Post = effect size for pre-supplementation

and post-supplementation time points. Data can be found in S2-S4 Tables.

Finally, we compared the average effect size between cohorts using the

Kruskal-Wallis test to determine which cohort was more responsive to the

cereal-bar supplementation. Comparing the two treated groups illustrated

significant differences in average effect size on the biological parameters, phyla,

and predicted metagenomic content ($p < 0.02$, Fig 2A-C). Average effect size on

biological parameters was higher in the OND compared to OD (0.19 and 0.09,

respectively, Fig 2A, S2 Table), whereas average effects on the phyla and

predicted metagenomic content were both higher in the OD cohorts (Fig 2B and

2C). Differences in effect sizes were most apparent for the predicted

metagenomic content as there was a 4 fold greater effect in the OD versus the

OND group (0.49 vs 0.11, respectively, $p = 4.3 \times 10^{-9}$, Fig 2C).

Discussion

We tested a cereal-bar supplementation containing β-glucan for 30 days

in OD and OND subjects with the goal of quantifying the effect in two similar

disease cohorts. We chose a cereal-bar with the goal of increasing fiber

consumption and minimizing side effects of sudden increase of fiber intake.

However, we found that consumption of 2 cereal-bars per day did not

significantly affect blood glucose in either cohort nor did we observe

improvement in any of the other individual biological parameters. The

ineffectiveness of the cereal-bar supplementation could result from several

possibilities. For instance, the amount of fiber delivered in this study was

relatively small even though it is near the FDA recommended amount for

cholesterol reduction (CFR 21 101.81, [22]). Martinez et al., [17] has showed

improvement in blood glucose with 60 g of additional oat fiber over four weeks

in healthy individuals while DeAngelis et al. [15] showed a similar result with

pasta containing β-glucan fiber in health individuals. In addition our subjects

were asked not to alter their normal dietary habit and the habitual fiber

consumption was not controlled, possibly confounding the effect of our cereal-

bar supplementation. We must also consider the possibility that our

observations are related to the medications taken by the OD subjects as we

asked the subjects not to change their drug treatment during the study.

Regardless of the reasons behind our observations, we found no significant effect

of cereal-bar supplementation on biological parameters in our cohorts.

As we had predicted, the abundance of Bacteroidetes responded

positively to the cereal-bar supplementation in the OD cohort while there was a

simultaneous decrease in the abundance of Firmicutes, suggesting small

additions of β-glucan in cereal-bars can alter taxonomic abundances in gut

bacterial communities. These observations are in line with studies highlighting

the saccharolytic nature of the Bacteroidetes in the human gut [23].

Interestingly, we did not observe any significant change in Bacteroidetes or

Firmicutes abundances in the OND participants. One possible explanation is that

the specific strains of Bacteroidetes and Firmicutes in the OND group are

somehow resistant to change with such a small infusion of nutrients. As

community analysis using 16S rRNA genes does not allow for strain level

resolution, an alternative approach may be necessary in order to identify specific

members of the microbial community that are resistant or susceptible to cereal-

bar supplementations. However, it is also possible that abundances of

Bacteroidetes and Firmicutes in the OND cohort are at levels where small diet

modifications can not overcome the inherent functional state of the bacterial

communities requiring more intense interventions to shift the abundance of

these taxa. As these two taxa represented more than 93% of the sequences in

this study, the possibility exists that consuming fiber containing cereal-bars may

be insufficient to garner consistent and significant changes of dominant taxa

across all patient cohorts.

Changes in abundance of Proteobacteria and Verrucomicrobia matched

our *a priori* predictions in both cohorts. The decline of Proteobacteria is

concomitant with studies indicating these bacteria are negatively associated with

diets high in fiber and positively associated with high fat diets [24, 25]. The

reduced abundance of the Proteobacteria has been linked to declines in

inflammatory markers and general host inflammation [26]. The reduction of

these organisms in human gut microbial communities is proposed to be

beneficial. Conversely, increases in Verrucomicrobia are associated with healthy

gut microbial communities [27] and diets high in fiber. The changes observed in

both cohorts suggest that small supplements of fiber delivered via cereal-bar can

alter abundances of these taxa. Though abundances of Proteobacteria and

Verrucomicrobia did not change significantly, our data nevertheless indicates

these taxa are responsive to diet modification across similar disease cohorts

possibly reflecting a general life history strategy of these organisms.

The alternative hypothesis that 2.8g β-glucan per day would have a

significant impact on average effect size in both cohorts was not supported.

Instead we observed significantly larger effects on the phyla and predicted

metagenomic content of the OD cohort compared to the OND individuals and

postulate that the lower species richness in the OD group is responsible for these

observations. Greater mean effect sizes in the OD group indicate these gut

communities were more susceptible to our cereal-bar supplementation than

were the OND group and coincide with recent report documenting richness as

factor in determining microbial responses to fiber supplementations [18]. Our

data build upon this observation and further suggest significantly changing

microbiomes with β-glucan containing cereal-bars may be limited to disease

cohorts harboring microbial communities that are relatively species poor as

these ecosystems have lost the functional flexibility needed to cope with a

changing environment. Unfortunately our experimental design did not allow us

to identify robust causal relationships between microbial communities and low

intensity diet supplementation (e.g. we used a single dosing level with small

cohort size). Nonetheless, our data suggest that small diet modifications

delivered by a cereal-bar have the potential to disproportionally affect low

diversity microbial communities and thus may be useful in studies addressing

ecological questions *in situ*, such as community stability, without altering host

physiology.

Given the significant effect on the OD bacterial communities, we would

have expected there to be a parallel effect on the biological parameters of these

participants as gut microbes and host phenotypes are clearly linked [7, 28, 29].

However, even though effect sizes were small in both cohorts, the physiology of

the OND cohort was more affected by the cereal-bar supplementation. The

intrinsic metabolic state of the host could explain this observation. In our

subjects it may be possible that insulin resistance, pancreatic beta cell functions

and even drug treatment dominate the biological set point of the diabetics and

small shifts in diet or bacterial communities had limited influence on the

metabolic outcomes. Unlike the diabetics, these same intrinsic factors in non-

diabetics may not be of equal intensity leaving a window for relatively small

dietary supplementations to have some degree of influence on the host. Low

dose fiber supplementation in healthy individuals reported by Martinez et al.

[17] would support this observation. Altogether, the results of this study suggest

bacterial community diversity in different cohorts may play a role in the

response of microbes to dietary supplementation aimed to manage host

metabolism.

Materials and Methods

Study population

A total of 46 overweight subjects with Body Mass Index (BMI) between 20

and 30 kg/m^2 were recruited in the Service of Therapeutic education for Chronic

Diseases of the University Hospitals of Geneva. After the inclusion and according

to the results of the initial screening, patients were assigned to either group 1 –

with type 2 diabetes mellitus (OD, n=21) or group 2 – without type 2 diabetes

mellitus (OND, n=25). We defined type 2 diabetes mellitus as fasting plasma

glucose > 7.0 mmol/l and/or HbA1c >7% and/or the presence of any glucose-

lowering treatment. OND subjects were matched for age, gender, BMI and ethnic

background of the OD subjects. Exclusion criteria were based on use of drugs

altering intestinal permeability (nonsteroidal anti-inflammatory drugs,

corticoids) or intestinal digestion and absorption (Orlistat, Colestipol,

anticoagulants, α-glucosidase inhibitors) and antibiotics administered in the 4

weeks preceding inclusion; previous abdominal surgery, gastro-intestinal

diseases interfering with intestinal absorption, cancer, bulimia, pregnancy;

parenteral nutrition or other ongoing dietary intervention, and diarrhea (>2

stools/day) within 7 days before enrolment. From this initial population, 26

were excluded for compliance with the supplementation protocol, they failed to

contribute samples or the sequencing effort was insufficient at any visit. The

resulting subset had a mean age of 42 yrs and 51 yrs for OND and OD,

respectively. Ten individuals for each cohort with overlapping age range were

selected for microbial analysis. Patient data is summarized in S1 Table. The study

is registered at ClinicalTrials.gov under the study number NCT02694172

(https://clinicaltrials.gov/ct2/results?term=NCT02694172).

Study design and intervention

A case controlled, single center prospective clinical trial design was used.

The study consisted of 4 visits and 3 periods. The 3 periods were i. recruiting

period (10 to30 days), ii. Diet-normalization period (14 days), and iii.

intervention period (30 days). After their recruitment, subjects received

instructions by a nutritionist for dietary normalization. After the dietary

normalization period, subjects received instructions for taking Nutren Balance

cereal-bars per day (one between breakfast and lunch and the other one

between lunch and dinner), rich in viscous soluble fiber β-glucan. Each cereal-

bar contained 65 Kcal. The total carbohydrate is 10.3 g that includes 3.8 g of

sugars and 2.5 g of fructose. The total protein is 1.9 g per bar. There is 1.8 g of

total fat of which saturated fat is 0.7 g, monounsaturated fat is 0.7 g and

polyunsaturated fat is 0.3 g. Total dietary fiber is 4.4 g of which 1.4 g is β-glucan.

Sodium content is 33 mg in each bar. The cereal-bar was well tolerated by all

patients without clinically significant side effects. Subjects submitted blood and

fecal samples at the enrollment, pre- and post- cereal-bar supplementation to

monitor host physiology and bacterial community composition. The study

protocol (06.42NRC) was approved by the Geneva ethical committee.

Participants were informed about the aims of the study and gave their written

consent.

Fecal Sample Collection, DNA extraction, PCR and sequencing

Stools were collected in sterile plastic 50 mL containers and frozen at -

80°C until processing. Frozen samples were partially thawed and from which

0.25 g of fecal matter was placed in the lysis tube and extracted according to the

manufacturers instructions. DNA was frozen at -20°C until its use in PCR

reactions to generate barcoded amplicons for sequencing on the MiSeq platform

[30]. Briefly, individual samples were amplified in triplicate, pooled then the

PCR products were quantified using PicoGreen dsDNS reagent. Equal amounts of

amplicon from each sample were then combined and sequenced on the MiSeq

platform. Sequencing was performed at the Nestlé Institute of Health Sciences

Functional Genomics Core facility.

Sequence analysis

Sequence data were quality filtered and demultiplexed in Qiime 1.8 [31]

using the default settings for the split_libraries_fastq.py command followed by

closed reference OTU picking at 97% sequence similarity against the gg_13_5

release (http://greengenes.secondgenome.com/downloads/database/13_5)

with the parallel_pick_otus_uclust_ref.py command. We additionally filtered out

Cyanobacteria to avoid chloroplast sequences and filtered out low abundance

OTUs according to Bokulich et al. 2013 [32] using the

filter_otus_from_otu_table.py command. Samples were then rarefied to 50000

sequences per sample (single_rarefaction.py) from which the relative abundance

of taxa (classified to the phylum level using summarize_taxa.py) was calculated

and used for downstream analysis. An estimate of bacterial richness was

performed using the Shannon index (alpha_diversity.py) on the rarefied data.

Predicted metagenomic content was also calculated using PiCRUST [33] and

summarized to KEGG level 2 for statistical evaluation. All bacterial community

data was expressed as the proportional abundance in each sample.

Data analysis

From the initial cohort we chose an age-matched sub set of patients to

assess changes in host metabolic measurements, taxonomic abundance, and

KEGG functions in response to the β-glucan cereal-bar supplementation. Using

Kruskal-Wallis tests implemented in Spotfire® (Göteborg, Sweden), we

compared group means within (e.g. pre- vs post-supplementation in each cohort)

and between cohorts (e.g. OND vs OD pre- supplementation and post-

supplementation) with α = 0.05 for all individual measurements (e.g. blood

glucose, taxon abundance, and individual KEGG functions, etc.) to identify

changes in these measurements. Bonferroni multiple comparison correction was

applied to the p-values for the taxonomic abundances and KEGG functions. We

next evaluated the magnitude of response of a data category (biological, phyla,

and predicted metagenome; S2-S4 Tables) by testing average effect size between

cohorts before and after cereal-bar supplementation [34]. The absolute values of

the effect size, calculated by Eq.1 where Es= effect size, m = mean, and σ =

standard deviation, were compared using Kruskal-Wallis tests to determine of

there were statistical differences in response.

Eq. 1

$$\mathbf{Es} = \frac{(m_1 - m_2)}{\sqrt{\frac{\sigma_1^2 + \sigma_2^2}{2}}}$$

Acknowledgements

405 We would like to thank Bernard Berger of the Nestlé Research Center Lausanne

406 for his assistance, Patrick Descombes and Deborah Moine of the NIHS Functional

407 Genomics core for their technical help with sequencing the 16S rRNA amplicons.

Supporting Information

S1 Table. Mean and standard deviation of biological parameters of 46

subjects at recruitment.

S2 Table. Means, standard deviations, and effect sizes for biological

parameters for the age-matched cohort. *Age and *height are not included in

the mean effect size calculation. Superscript "a" indicates significant difference

between cohorts post-supplementation.

S3 Table. Means, standard deviations, and effect sizes for taxa and Shannon

Diversity for the age-matched cohort.

S4 Table. Means, standard deviations, and effect sizes for the KEGG

functions predicted by PiCRUSt for the age-matched cohort.

S5. Clinical Protocol

S6. TREND Flowchart

S7. TREND Checklist

References

1.   World Health Organization, 2003.

2.   Wu GD, Chen J, Hoffmann C, Bittinger K, Chen Y-Y, Keilbaugh SA, et al.

Linking Long-Term Dietary Patterns with Gut Microbial Enterotypes.

Science. 2011; 334:105-8. doi: 10.1126/science.1208344. PMID:3368382.

3.   Matijašić B, Obermajer T, Lipoglavšek L, Grabnar I, Avguštin G, Rogelj I.

Association of dietary type with fecal microbiota in vegetarians and

omnivores in Slovenia. European Journal of Nutrition. 2014; 53:1051-64.

doi: 10.1007/s00394-013-0607-6. PMID: 24173964

4.   Kabeerdoss J, Shobana Devi R, Regina Mary R, Ramakrishna BS. Faecal

microbiota composition in vegetarians: comparison with omnivores in a

cohort of young women in southern India. British Journal of Nutrition.

2012; 108:953-7. doi:10.1017/S0007114511006362. PMID: 22182464

5.   Glick-Bauer M, Yeh M-C. The Health Advantage of a Vegan Diet: Exploring

the Gut Microbiota Connection. Nutrients. 2014; 6:4822-38 doi:

10.3390/nu6114822. PMID: 25365383

6.   De Filippo C, Cavalieri D, Di Paola M, Ramazzotti M, Poullet JB, Massart S,

et al. Impact of diet in shaping gut microbiota revealed by a comparative

study in children from Europe and rural Africa. Proceedings of the

National Academy of Sciences of the United States of America. 2010;

107:14691-6. doi: 10.1073/pnas.1005963107. PMID: 20679230.

7.   Ley RE, Hamady M, Lozupone C, Turnbaugh P, Ramey RR, Bircher JS, et al.

Evolution of mammals and their gut microbes. Science. 2008. 320:1647-

51. doi: 10.1126/science.1155725. PMID:2649005.

8.   Machiels K, Joossens M, Sabino J, De Preter V, Arijs I, Eeckhaut V, et al. A

decrease of the butyrate-producing species Roseburia hominis and

Faecalibacterium prausnitzii defines dysbiosis in patients with ulcerative

colitis. Gut. 2014; 63:1275-83. doi: 10.1136/gutjnl-2013-304833. PMID:

24021287

9.   Flint HJ, Scott KP, Duncan SH, Louis P, Forano E. Microbial degradation of

complex carbohydrates in the gut. Gut Microbes. 2012; 3:289-306. doi:

10.4161/gmic.19897. PMID: 22572875

10.   Salyers AA, Vercellotti JR, West SE, Wilkins TD. Fermentation of mucin

and plant polysaccharides by strains of Bacteroides from the human

colon. Applied and Environmental Microbiology. 1977; 33:319-22. PMID:

848954

11.   Sartor RB, Mazmanian SK. Intestinal Microbes in Inflammatory Bowel

Diseases. American Journal of Gastroenterology. Suppl. 2012; 1:15-21

doi:10.1038/ajgsup.2012.4. PMID: 19107650

12.   Gevers D, Kugathasan S, Denson LA, Vázquez-Baeza Y, Van Treuren W,

Ren B, et al. The treatment-naïve microbiome in new-onset Crohn's

disease. Cell Host & Microbe. 2014; 15:382-92.

doi:10.1016/j.chom.2014.02.005. PMID: 4059512

13.   Flint HJ, Duncan SH, Scott KP, Louis P. Interactions and competition

within the microbial community of the human colon: links between diet

and health. Environmental Microbiology. 2007; 9:1101-11. doi:

10.1111/j.1462-2920.2007.01281.x. PMID: 17472627

14.   Louis P, Scott KP, Duncan SH, Flint HJ. Understanding the effects of diet on

bacterial metabolism in the large intestine. Journal of Applied

Microbiology. 2007; 1197-208. doi: 10.1111/j.1365-2672.2007.03322.x.

PMID: 17448155

15.   De Angelis M, Montemurno E, Vannini L, Cosola C, Cavallo N, Gozzi G, et al.

The role of whole-grain barley on human fecal microbiota and

metabolome. Applied and Environmental Microbiology. 2015; doi:

10.1128/aem.02507-15. PMID: 26386056

16.   Dewulf EM, Cani PD, Claus SP, Fuentes S, Puylaert PGB, Neyrinck AM, et al.

Insight into the prebiotic concept: lessons from an exploratory, double

blind intervention study with inulin-type fructans in obese women. Gut.

2013; 62:1112-21. doi: 10.1136/gutjnl-2012-303304. PMID:3711491.

17.   Martinez I, Lattimer JM, Hubach KL, Case JA, Yang J, Weber CG, et al. Gut

microbiome composition is linked to whole grain-induced immunological

improvements. The ISME Journal. 2013; 7:269-80.

doi:10.1038/ismej.2012.104. PMID: 23038174

18.   Tap J, Furet J-P, Bensaada M, Philippe C, Roth H, Rabot S, et al. Gut

microbiota richness promotes its stability upon increased dietary fibre

intake in healthy adults. Environmental Microbiology. 2015; doi:

10.1111/1462-2920.13006. PMID: 26235304

19.   Walker AW, Ince J, Duncan SH, Webster LM, Holtrop G, Ze X, et al.

Dominant and diet-responsive groups of bacteria within the human

colonic microbiota. The ISME Journal. 2011; 5:220-30.

doi:10.1038/ismej.2010.118. PMID: 20686513

20.   Hooda S, Boler BMV, Serao MCR, Brulc JM, Staeger MA, Boileau TW, et al.

454 Pyrosequencing Reveals a Shift in Fecal Microbiota of Healthy Adult

Men Consuming Polydextrose or Soluble Corn Fiber. The Journal of

Nutrition. 2012; 142:1259-65. doi: 10.3945/jn.112.158766. PMID:

22649263

21.   Benus RFJ, van der Werf TS, Welling GW, Judd PA, Taylor MA, Harmsen

HJM, et al. Association between Faecalibacterium prausnitzii and dietary

fibre in colonic fermentation in healthy human subjects. British Journal of

Nutrition. 2010; 104:693-700. doi:10.1017/S0007114510001030.

PMID: 20346190.

22.   US Food and Drug Administration. TITLE 21—Food and drugs Chapter 1,

subchapter B –Food for human consumption. 2015.

w.accessdata.fda.gov/scripts/cdrh/cfdocs/cfcfr/cfrsearch.cfm?fr=101.81

23.   Martens EC, Chiang HC, Gordon JI. Mucosal Glycan Foraging Enhances

Fitness and Transmission of a Saccharolytic Human Gut Bacterial

Symbiont. Cell host & microbe. 2008; 4(5):447-57. doi:

10.1016/j.chom.2008.09.007. PMID: 18996345

24.   Hildebrandt MA, Hoffmann C, Sherrill–Mix SA, Keilbaugh SA, Hamady M,

Chen YY, et al. High-Fat Diet Determines the Composition of the Murine

Gut Microbiome Independently of Obesity. Gastroenterology. 2009;

137:1716-24.e1-2. doi: 10.1053/j.gastro.2009.08.042. PMID: 19706296

25.   Shin N-R, Whon TW, Bae J-W. Proteobacteria: microbial signature of

dysbiosis in gut microbiota. Trends in Biotechnology. 2015; 33:496-503.

doi: doi.org/10.1016/j.tibtech.2015.06.011. PMID: 26210164

26.   Carvalho Frederic A, Koren O, Goodrich Julia K, Johansson Malin EV,

Nalbantoglu I, Aitken Jesse D, et al. Transient Inability to Manage

Proteobacteria Promotes Chronic Gut Inflammation in TLR5-Deficient

Mice. Cell Host & Microbe. 2012; 12:139-52.

doi:10.1016/j.chom.2012.07.004. PMID: 22863420

27.   Everard A, Belzer C, Geurts L, Ouwerkerk JP, Druart C, Bindels LB, et al.

Cross-talk between Akkermansia muciniphila and intestinal epithelium

controls diet-induced obesity. Proceedings of the National Academy of

Sciences. 2013; 110:9066-71. doi: 10.1073/pnas.1219451110. PMID:

23671105

28.   Larsen N, Vogensen FK, van den Berg FWJ, Nielsen DS, Andreasen AS,

Pedersen BK, et al. Gut Microbiota in Human Adults with Type 2 Diabetes

Differs from Non-Diabetic Adults. PLoS ONE. 2010; 5:e9085. doi:

10.1371/journal.pone.0009085. PMID: 20140211

29.   Turnbaugh PJ, Ley RE, Mahowald MA, Magrini V, Mardis ER, Gordon JI. An

obesity-associated gut microbiome with increased capacity for energy

harvest. Nature. 2006; 444:1027-131. doi:10.1038/nature05414. PMID:

17183312

30.   Caporaso JG, Lauber CL, Walters WA, Berg-Lyons D, Huntley J, Fierer N, et

al. Ultra-high-throughput microbial community analysis on the Illumina

HiSeq and MiSeq platforms. The ISME Journal 2012; 6:1621-4.

doi:10.1038/ismej.2012.8. PMID: 22402401

31.   Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello

EK, et al. QIIME allows analysis of high-throughput community

sequencing data. Nat Meth. 2010;7:335-6. doi:10.1038/nmeth.f.303.

PMID: 20383131

32.   Bokulich NA, Subramanian S, Faith JJ, Gevers D, Gordon GI, Knight R, et al.

554   Quality-filtering vastly improves diversity estimates from Illumina

555   amplicon sequencing. Nature Methods. 2013; 10:57-59.

556   doi:10.1038/nmeth.2276. PMID: 23202435

557   33.   Langille MGI, Zaneveld J, Caporaso JG, McDonald D, Knights D, Reyes JA, et

558   al. Predictive functional profiling of microbial communities using 16S

559   rRNA marker gene sequences. Nature Biotechnology. 2013; 814-21.

560   doi:10.1038/nbt.2676. PMID: 23975157

561   34.   Sullivan GM, Feinn R. Using Effect Size—or Why the P Value Is Not

562   Enough. Journal of Graduate Medical Education. 2012; 4(3):279-82.

563   doi: 10.4300/JGME-D-12-00156.1. PMID: 23997866

	OND Mean			OD Mean		
	Enrollment	Pre-supplementation	Post-supplementation	Enrollment	Pre-supplementation	Post-supplementation
*Age	48 (3.5)	48 (3.5)	48 (3.5)	47 (4.3)	47 (4.3)	47 (4.3)
*Height (cm)	167 (10.3)	167 (10.3)	167 (10.3)	171 (10)	171 (10)	171 (10)
Weight (kg)	95 (14.7)	95 (14.1)	94 (14.4)	98 (17)	96 (17)	96 (16)
BMI (kg m^2)	34 (5.2)	34 (5.2)	34 (5.3)	34 (4.66)	33 (4.5)	33 (4.4)
Waist Circumference (cm)	104 (11.4)	105 (11.3)	106 (11.3)	113 (10)	113 (10)	113 (12)
Hip Circumfrenece (cm)	117 (8.1)	117 (8.3)	116 (8.2)	116 (12)	114 (11)	115 (12)
Diastolic BP (mmHg)	95 (16)	87 (5.1)	89 (13)	87 (9.86)	85 (9.63)	88 (14)
Systolic BP (mmHg)	131 (15)	124 (13)	131 (21)	130 (17)	129 (22)	131 (17)
Cardiac Frequency (beats min^{-1})	67 (11)	68 (12)	72 (16)	79 (24)	78 (13)	77 (14)
Total Cholesterol (mmol L^{-1})	5 (0.74)	4.9 (0.87)	5.2 (0.69)	4.9 (1.3)	4.6 (0.82)	4.7 (0.49)
Triglyceride (mmol L^{-1})	1.1 (0.42)	1.2 (0.51)	1.2 (0.43)	4.1 (5.8)	3.6 (4.3)	2.9 (2.1)
HDL cholesterol (mmol L^{-1})	1 (0.23)	1 (0.3)	1.1 (0.25)	0.9 (0.24)	0.8 (0.24)	0.8 (0.16)
LDL (mmol L^{-1})	3.5 (0.61)	3.4 (0.68)	3.6 (0.65)[a]	2.8 (0.74)	2.7 (0.7)	2.7 (0.67)[a]
Glucose (mmol L^{-1})	5 (0.68)	5 (0.63)	5.1 (0.46)[a]	6.4 (2)	6.5 (2.2)	6.3 (1.5)[a]
Insulin (uIU ml^{-1})	17 (7.1)	19 (8.1)	19 (7.6)	20 (13)	17 (10)	18 (12)
FFA (mmol L^{-1})	0.5 (0.13)	0.5 (0.2)	0.6 (0.23)	0.6 (0.25)	0.5 (0.19)	0.5 (0.19)
HOMA_IR	3.9 (1.8)	4.4 (2.1)	4.3 (1.6)	6.2 (5.3)	5.1 (3.7)	5.5 (5)
LBP (ug ml^{-1})	28 (6)	27 (3.8)	29 (4.8)	31 (8.7)	27 (7.4)	29 (7.7)