Temporal Stability and Molecular Persistence of the Bone Marrow Plasma Cell Antibody Repertoire

Gabriel C. Wu[1], Nai-Kong V. Cheung[3], George Georgiou[2,4,5,6] , Edward M. Marcotte[1,2,6*] and Gregory C. Ippolito[2,*]

1. Center for Systems and Synthetic Biology University of Texas at Austin, Austin, TX, USA
2. Department of Molecular Biosciences, University of Texas at Austin, Austin, Texas, USA
3. Department of Pediatrics, Memorial Sloan-Kettering Cancer Center, New York, New York, USA
4. Department of Biomedical Engineering, University of Texas at Austin, Austin, Texas, USA
5. Department of Chemical Engineering, University of Texas at Austin, Austin, Texas, USA
6. Institute for Cell and Molecular Biology, University of Texas at Austin, Austin, TX, USA

*To whom correspondence should be addressed: Edward M. Marcotte, marcotte@icmb.utexas.edu; Gregory C. Ippolito, gci@mail.utexas.edu.

ABSTRACT

Plasma cells in human bone marrow (BM PCs) are thought to be intrinsically
long-lived and to be responsible for sustaining lifelong immunity through the constitutive
secretion of antibody—but the underlying basis for this serological memory remains
controversial. Here, we analyzed the molecular persistence of serological immunity by
an examination of BM PC immunoglobulin heavy-chain (IGH) transcripts derived from
serial bone marrow specimens obtained during a span of several years. Using high-
throughput sequence analysis of the same individual for 6.5 years, we show that the BM
PC repertoire is remarkably stable over time. We find that the bias in IGH V, D, and J
individual gene usage and also the combinatorial V–D, V–J, D–J, and V-D-J usage
across time to be nearly static. When compared to a second donor with time points 2
years apart, these overall patterns are preserved, and surprisingly, we find high
correlation of gene usage between the two donors. Lastly, we report the persistence of
numerous BM PC clonal clusters (~2%) identifiable across 6.5 years at all time points
assayed, supporting a model of serological memory based, at least in part, upon intrinsic
longevity of human PCs. We anticipate that this longitudinal study will facilitate the ability
to differentiate between healthy and diseased antibody repertoire states, by serving as a
point of comparison with future deep-sequencing studies involving immune intervention.

INTRODUCTION

The human bone marrow (BM) is a specialized immune compartment that is responsible for both the initial generation of newly-formed B cells and also the maintenance of terminally differentiated, antibody-secreting plasma cells (PCs). The BM, and the PCs it harbors, is a major site of antibody production and is the major source of all classes and subclasses of human immunoglobulins (Ig) detectable in the serum[1,2]. Ig-secreting BM PCs are generally believed to be "long-lived" and maintained for the lifespan of the organism[3]. In this regard, it has been well established in longitudinal serological studies that antiviral serum antibodies can be remarkably stable, with half-lives ranging from 50 years (e.g. varicella-zoster virus) to 200 years for other viruses (e.g. measles and mumps); however, in contrast, antibody responses to non-replicating antigens (e.g. tetanus and diphtheria bacterial toxins) rapidly decay with much shorter half-lives of only 10-20 years[4]. Not only does this suggest that antigen-specific mechanisms play a substantial role in the establishment and/or maintenance of serological memory, but raises the question of whether the differential stability of antibody responses might reflect differential intrinsic longevity of PCs themselves. This has been previously proposed in the context of vaccinations and infections[4,5], and is also supported by observations of differential stability of autoantibody titers when using B-cell depleting therapies to treat autoimmune diseases[6,7].

The basis underlying lifelong serological memory (antibody responses) remains controversial[3,8,9]. Data supporting a model for intrinsic longevity in PC survival (and hence longevity in serum antibody maintenance) has been posited for the laboratory mouse[10,11], but data for human PCs are absent. Based upon the murine models, it has been assumed that human BM PCs are similarly long-lived and that these long-lived PCs are the major source of serum antibodies; however, only recently have antigen-specific BM PCs been ascertained for their contribution to the pool of serum antibodies in humans[5,12]. Despite these notable advances, the availability of corresponding molecular data (namely, sequence data of BM PC Ig transcripts) and of information regarding PC dynamics *in vivo* are scarce.

Three studies have generated BM PC data using next-generation sequencing techniques, but none of which have examined the temporal changes that occur in the immune repertoire over time[5,13,14]. Here, building upon our prior experiences with the

comprehensive analysis of human cellular and serological antibody repertoires[15–19], we
present the first longitudinal study of serially acquired human BM PCs assayed by next-
generation deep sequencing. To directly measure the temporal dynamics of BM PCs—
and to indirectly gain insight into long-lived serological memory—we sequenced the
recombined VHDJH region, which encodes the variable region (V domain) of IGH heavy
chains of BM PCs from the same individual over seven timepoints encompassing a total
of 6.5 years and from a second individual with two timepoints over 2.3 years. The
temporal resolution and duration of sampling provides a method to interrogate the *in vivo*
temporal dynamics of BM PCs in a previously uncharacterized way. We provide detailed
temporal information on the individual genes (IGH V, D, and J), gene combinations (V-D,
V-J, D-J, V-D-J), and persistent CDR-H3 clonotypes. The second individual provides
support that our observations are not unique. Overall, our results (i) underscore the
temporal stability of the IGH V region repertoire according to multiple metrics (temporally
stable IGH molecular phenotypes), and (ii) provide unequivocal sequence-based
evidence for the persistence of PC cellular clonotypes spanning 6.5 years.

RESULTS

Serial Bone Marrow Biopsies Followed by Next-Generation Sequencing
Demonstrate Temporal Dynamics of the Immune Repertoire.

To investigate the temporal dynamics of the IGH antibody gene repertoire of
bone marrow plasma cells (BM PCs), we sampled, sorted, and performed high-
throughput sequencing (Fig. 1a). Serial bone marrow biopsies were obtained from two
adolescents (10–17 years of age) as part of routine evaluations for non-immuno-
hematological disease. BM PCs were isolated using fluorescence-activated cell sorting
(FACS). BM PCs were obtained by sorting for CD38++ CD138+ cells within the
mononuclear light-scatter gate (Fig. 1b). Additionally, the cells were uniformly positive
for the TNF-receptor superfamily member CD27 (Fig. 1b, inset). Importantly, we avoided
gating of the pan-B cell marker CD19 since previous characterizations of human BM
PCs show heterogeneous expression of CD19[20,21]. Therefore, our method captured all
recently described BM PC subpopulations[5,12] with an overall CD19$^{+/-}$ CD27$^+$ CD38^{++}

CD138⁺ phenotype. Subsequently, transcripts were amplified from BM PCs expressing

IgM, IgG, and IgA using RT-PCR followed by high-throughput sequencing.

In total, 51,200 BM PC were sorted, which generated 503,415 reads after quality-

threshold filtering (see Methods and Supplementary Table S1). These data were

distributed across seven timepoints spanning 6.5 years (Fig. 1c). A biological replicate, a

second frozen ampule derived from the same bone marrow aspiration, was also

collected from each donor and analyzed. Multiple sampling from the same donor allows

us to accurately identify the active heavy chain genes that compose this donor's

antibody repertoire. Specifically, we identify 38 IGHV genes, 21 IGHD genes, and 6

IGHJ genes (4,788 combinations).

IGH V, D, and J Frequency Are Highly Stable Over 6.5 Years.

To determine the stability of individual gene usage, we assessed the frequency

of each IGH V, D, and J gene across time (Fig. 2). Surprisingly, we saw stable behavior

of these genes, with the most frequently used genes (e.g. IGHV4-34) showing

consistently high expression while less frequently used genes (e.g. IGHV3-72) showed

consistently low expression. This observation was quantified using the Mann-Kendall

Test, which evaluates trends in time series data. We found that 89% of IGHV genes,

95% of IGHD genes and 100% IGHJ genes showed no statistically significant trends

(Mann-Kendall test, $p > 0.05$), indicating that the IGHV (Fig. 2a), IGHD (Fig. 2b), and

IGHJ (Fig. 2c) genes were time stable.

Next, we analyzed population behavior of gene usage. Averaging across all

timepoints, we observed a highly skewed distribution of individual gene frequencies,

consistent with previous single timepoint observations. Only 6 IGHV genes (16%)

accounted for greater than 50% of total IGHV gene usage by frequency (Fig. 2d). IGHD

genes that have previously been shown to have biased usage IGHD2-2, IGHD3-3, and

IGHD3-22[22] together accounted for 33% of total IGHD usage (Fig. 2b). In addition,

known biases in IGHJ usage[23] are recapitulated as IGHJ4, IGHJ6, and IGHJ5 account

for 86% of total IGHJ usage. Furthermore, our analysis demonstrated that IGH V, D, and

J gene usage were not significantly different from a log-normal distribution (Anderson-

Darling, H=0, $p > 0.05$).

IGH V-D, D-J, V-J, and V-D-J Combinations Are Stable Over Time.

Given the temporal stability of individual genes, we hypothesized that differential

intrinsic longevity might be found in gene combinations. Surprisingly, our analysis

suggests that gene combinations, like their individual component genes, are time stable

as well. We found that 92% V-J (Fig. 4), 97% V-D (Supplementary Fig. 1a), 95% D-J

(Supplementary Fig. 1b), and 97% V-D-J (Supplementary Fig. 1c) do not show

significant trends (Mann-Kendall, H=0, p>0.05).

To better understand the nature of gene combinations, we analyzed preferential

gene pairing biases by comparing the expected versus observed frequency of pairwise

gene combinations. The observed frequency of each gene combination was found to be

correlated to its expected frequency (Spearman r): V-D (0.74), V-J (0.87), D-J (0.93),

and V-D-J (0.65) (Fig. 5a-d). This high level of correlation and lack of significant outliers

suggests that there is minimal gene pairing linkage and that the gene pairing is a

random process.

Persistent Clonotypes Are Stable Over Time.

To understand how each of these individual genes and gene combinations

together might indicate the existence of long lived PCs, we analyzed the behavior of the

CDR-H3, the highest resolution possible for a single identifier of an antibody producing

cell. To eliminate errors and ambiguities, we clustered CDR-H3s into clonotypes based

on previously established criteria (see Methods). On average, we found that 16% of

clonotypes are shared between adjacent timepoints (Fig. 6a, top). Interestingly, 23

clonotypes persisted across all timepoints (Fig. 6b). We found that 100% of these

persistent clonotypes were time stable (Fig. 6a, bottom, Mann-Kendall test, h=0, p>0.05)

and 78% (18/23) were of the IgA isotype. In addition, characteristics of the complete

CDR-H3 population, specifically CDR-H3 lengths (Supplementary Fig. 2) and hydropathy

index (Supplementary Fig. 3), are unchanged over time. The overall total distribution of

CDR-H3 lengths are consistent with previously reported single timepoint values[27]. Also,

higher expressing CDR-H3s tended to be neither hydrophobic nor hydrophilic

(Supplementary Fig. 3) and no significant trends between hydrophobicity and expression

level were found.

A Second Donor Corroborates The Observations From The First Donor.

To verify our longitudinal observations of stability and random gene choices from

Donor 1, we analyzed a second donor across two years (Fig. 7). We identified 38 IGHV

genes, 22 IGHD genes, and 6 IGHJ genes (5,016 combinations, 6,763 cells, 48,525

reads) (Supplementary Table 1 and Supplementary Fig. 4). Donor 1 and Donor 2 show

highly correlated IGHV gene usage (r=0.82). Thus, the trends observed in Donor 1 were

also observed in Donor 2. Specifically, individual IGHV, IGHD, and IGHJ gene usages

were time stable (Supplementary Fig. 4), as were the gene combinations

(Supplementary Fig. 5). Consistent with Donor 1, Donor 2 showed no preferential

pairing in gene combinations (Supplementary Fig. 6). These results are highly consistent

with the trends observed in Donor 1, and together, they indicate that BM PC antibody

gene and gene combination usage show surprisingly minimal variation between

individuals and across time.

DISCUSSION

Next-generation sequencing has enabled unprecedented ability to explore the

details of the human immune repertoire[28,29]. Whereas previous studies have been able

to describe some aspects of immune repertoire at a single point in time, our results

harness the power of next-generation sequencing to elucidate the temporal dynamics of

BM PCs over 6.5 years. Importantly, our data provide molecular resolution of antibody

identity in the form of clonotypes, which is not possible with classic techniques like

enzyme-linked immunosorbent assay (ELISA).

In this study, we show that the immune system is naturally polarized in both gene

choice and gene combination usage and that the polarization is maintained over time.

We also found that the bias is not primarily a result of gene linkage, suggesting there are

additional genomic or extrinsic factors that contribute to polarization. For example, there

may be an Eμ like enhancer in humans that results in higher than expected IGHJ4

usage[30]. Alternatively, the long arms race between the human immune system and the

antigens it faces throughout evolutionary history may have established a preferential

gene choice long ago. So higher expressing genes are likely broad-spectrum antibodies

that have been useful in fighting particular classes of disease and continue to do so

today. For example, IGHV1-69 is implicated in many viral diseases like influenza and

HIV. IGHV4-34 is associated with a range of autoimmune disorders, like cold agglutinin

disease and systemic lupus erythematosus (SLE).

Taken together, our results indicate that the BM PC antibody gene and gene

combination usage is highly stable over long periods of time. While short term studies

over days, weeks or at most a year show that antigen stimulus can result in dramatic

immune repertoire changes over weeks[31] our study extends over half a decade and is

the longest longitudinal study to date. Given the known resistance we have for a large

and diverse array of pathogenic antigens, it is surprising that the overall effect of these

changes is not reflected in gene usage. This suggests that (i) either the existing immune

gene combinations are established early on or that the mechanism of gene choice

results in consistent bias over long periods of time, (ii) the effective size and variability of

the immune repertoire is restricted, and (iii) antigen challenges in the form of

vaccinations and illness do not have lasting changes at the gene usage level. Therefore,

the immune repertoire must defend against a wide array of pathogens with only relatively

small changes in immune diversity. On the other hand, intrinsic immunological memory

is a well-established concept and BM PCs are thought to persist for long periods of time.

An outstanding question remains of how intrinsic longevity is established and

maintained. Although our results are unable to verify that any particular PC member

persists, we can conclude that the clonotype, which defines binding specificity at the

molecular level, does persist for at least 6.5 years. Our results suggest clonotype

persistence contributes to the mechanism underlying long term immunological memory.

Moreover, we observe steady usage of gene and gene combinations throughout

our experiment. Our observations suggest that there are large resident pools of plasma

cells of the same identity, from which we can sample continuously with no loss of relative

expression levels. We conclude that a large population of clonally-related PCs is

responsible for persistent and stable gene expression of a given antibody, and that any

given PC clone does not contribute greatly to the overall expression of that antibody.

In conclusion, we have used high-throughput, next-generation sequencing to

definitively identify long-term persistent BM PC clonotypes, which has implications in

clinical intervention studies, vaccines, and immunotherapy. Future next-generation

sequencing studies can provide an even more detailed picture of the B cell immune

repertoire as these studies could include advances in VH:VL native-pair sequencing

(paired BCR-seq[19]), the analysis of correlations between BM PC repertoires and serum

immunoglobulin species (Ig-seq[17,18]), and an examination of the connectivity of B cells at
various developmental stages (e.g., clonal relationships between circulating memory B
cells and sessile BM PCs) . Our study provides a foundation upon which these further
studies can be built.
252
253   **MATERIALS & METHODS**
254
255   **Bone Marrow Specimens**
256
Serially acquired human bone marrow specimens were collected from two donors by
aspiration from the ileac crest, and mononuclear cells were enriched by Ficoll hypaque
centrifugation. The two adolescent–teenage donors were originally diagnosed with
neuroblastoma but had been asymptomatic and disease-free for many years according
to routine bone marrow histology. A complete description of the donors' past medical
history and ages at the time of the multiple time point collections is included in
Supplementary Table 1. All procedures were performed per a standard operating
procedure at the Memorial Sloan-Kettering Cancer Center and collected according to a
longstanding protocol approved by the MSKCC Institutional Review Board. Aspirates
were withdrawn from four sites and combined (total of 8–10 mL from 4 sites, 2–2.5 mL
per site) drawn from the following: anterior right iliac crest, anterior left iliac crest,
posterior right iliac crest, and posterior left iliac crest. The same attending physicians
performed these procedures and usually biopsied through the same surgical site each
time. De-identified specimens were shipped overnight on dry ice to the University of
Texas at Austin.
272
273   **Flow Cytometry and Isolation of Plasma Cells (PC)**
274
BM samples were quick-thawed in a 37 °C H_2O bath and slowly diluted into RPMI-1640
complete medium containing DNaseI (Sigma D 4513; 20 U/mL), pelleted, washed and
re-suspended in 2 mL FACS buffer (Dulbecco's PBS + 0.5% BSA Fraction V). Cell
viability was determined using Trypan Blue exclusion and on average was approximately
90% per specimen. After a one-hour recovery at room temperature, BM cells were
stained for 30 minutes at room temperature using empirically-determined optimal
titrations of monoclonal antibodies: CD38-FITC (HIT2), CD138-PE (B-B4), CD27-APC

(M-T271), and CD19-v450 (HIB19). $CD19^{+/-}CD38^{++}CD138^{+}$ cells in human BM were

collected as plasma cells (PC). PCs were observed to be heterogeneous for expression

of the CD19 B-lineage marker; therefore, CD19-gating was avoided. $CD38^{++}CD138^{+}$

PCs were additionally gated by light scatter properties (FSC v. SSC) to exclude debris,

apoptotic cells, and remnant granulocytes. All cell sorts were performed on a FACSAria

flow cytometer. Cells were sorted directly into TRI Reagent for RNA preservation.

RT-PCR, High Throughput Sequencing of IGH V, D, and J Genes

All methods and reagents were as previously described[15]. Variable genes (recombined

VHDJH region, which encodes the V domain) of IGH isotypes IgM, IgG, and IgA were

amplified from oligo-dT cDNA and sequenced at high-throughput using the Roche 454

GS FLX technology using titanium long-read chemistry.

Data Processing, Analysis, and Visualization

All sequence data have been deposited to NCBI SRA under BioProject number

PRJNA310043. IGHV, IGHD, IGHJ, and CDR-H3 regions for each read was quality

filtered, processed and annotated using the VDJFasta utility described previously[32].

Reference IGHV, IGHD, and IGHJ genes from the international ImMunoGeneTics

(IMGT) database were used. Mann-Kendall Tests were performed in Matlab, against the

null hypothesis of no trend (alpha=0.05). Spearman r non-parametric correlation analysis

was performed in python using the scipy library. To perform clonal clustering analysis,

CDR-H3 sequences were clustered to form clonotypes, as established previously[33,34],

using CD-HIT[35] with the following criteria: (i) Same IGHV and IGHJ, (ii) minimum length

of 5, and (iii) 85% similarity threshold on the amino acid level.

Circular visualization plots were created with Circos software v0.67-7[36] where genes

were sorted by expression within each timepoint and connected to adjacent timepoints

via colored lines showing their expression levels. All other data visualization was

performed using Python and matplotlib.

Acknowledgements

This work was supported by NSF Graduate Research Fellowship DGE-1110007 (to
GCW), HDTRA1-12-C-0105 from DTRA (GG, GCI, EMM), and grants from the NIH,
NSF, and Welch Foundation (F-1515) to EMM.
319

320 **Authorship contributions**

322 GG and NKC conceived the study. GCW and GCI designed and performed experiments,
323 analyzed data, prepared figures, and wrote the manuscript, under the supervision of
324 EMM. All authors reviewed the manuscript.
325

326 **Conflict of Interest**

328 The authors declare no conflict of interest.
329

REFERENCES

1.   McMillan, R. *et al.* Immunoglobulin synthesis by human lymphoid tissues: normal
bone marrow as a major site of IgG production. *J. Immunol.* **109,** 1386–94 (1972).

2.   Benner, R., Hijmans, W. & Haaijman, J. J. The bone marrow: the major source of
serum immunoglobulins, but still a neglected site of antibody formation. *Clin. Exp.*
Immunol. **46,** 1–8 (1981).

3.   Radbruch, A. *et al.* Competence and competition: the challenge of becoming a
long-lived plasma cell. *Nat. Rev. Immunol.* **6,** 741–50 (2006).

4.   Amanna, I. J., Carlson, N. E. & Slifka, M. K. Duration of humoral immunity to
common viral and vaccine antigens. *N. Engl. J. Med.* **357,** 1903–15 (2007).

5.   Halliley, J. L. *et al.* Long-Lived Plasma Cells Are Contained within the
CD19−CD38hiCD138+ Subset in Human Bone Marrow. *Immunity* **43,** 132–45
(2015).

6.   Cambridge, G. *et al.* Serologic changes following B lymphocyte depletion therapy
for rheumatoid arthritis. *Arthritis Rheum.* **48,** 2146–54 (2003).

7.   Cambridge, G. *et al.* B cell depletion therapy in systemic lupus erythematosus:
effect on autoantibody and antimicrobial antibody profiles. *Arthritis Rheum.* **54,**
3612–22 (2006).

8.   Amanna, I. J. & Slifka, M. K. Contributions of humoral and cellular immunity to
vaccine-induced protection in humans. *Virology* **411,** 206–15 (2011).

9.   Zinkernagel, R. On plasma cell longevity or brevity. *Expert Rev. Vaccines* **13,**
821–3 (2014).

10.   Manz, R. A., Thiel, A. & Radbruch, A. Lifetime of plasma cells in the bone marrow.
Nature **388,** 133–4 (1997).

11.   Slifka, M. K., Antia, R., Whitmire, J. K. & Ahmed, R. Humoral immunity due to
long-lived plasma cells. *Immunity* **8,** 363–72 (1998).

12.   Mei, H. E. *et al.* A unique population of IgG-expressing plasma cells lacking CD19
is enriched in human bone marrow. *Blood* **125,** 1739–48 (2015).

13.   Cowan, G. *et al.* Massive parallel IGHV gene sequencing reveals a germinal
center pathway in origins of human multiple myeloma. *Oncotarget* **6,** 13229–40
(2015).

14.   Tschumper, R. C. *et al.* Comprehensive assessment of potential multiple
myeloma immunoglobulin heavy chain V-D-J intraclonal variation using massively
parallel pyrosequencing. *Oncotarget* **3,** 502–13 (2012).

15.   Ippolito, G. C. *et al.* Antibody repertoires in humanized NOD-scid-IL2Rγ(null) mice
and human B cells reveals human-like diversification and tolerance checkpoints in
the mouse. *PLoS One* **7,** e35497 (2012).

16.   DeKosky, B. J. *et al.* High-throughput sequencing of the paired human
immunoglobulin heavy and light chain repertoire. *Nat. Biotechnol.* **31,** 166–9
(2013).

17.   Wine, Y. *et al.* Molecular deconvolution of the monoclonal antibodies that
comprise the polyclonal serum response. *Proc. Natl. Acad. Sci. U. S. A.* **110,**
2993–8 (2013).

18.   Lavinder, J. J. *et al.* Identification and characterization of the constituent human
serum antibodies elicited by vaccination. *Proc. Natl. Acad. Sci. U. S. A.* **111,**
2259–64 (2014).

19.   DeKosky, B. J. *et al.* In-depth determination and analysis of the human paired
heavy- and light-chain antibody repertoire. *Nat. Med.* **21,** 86–91 (2014).

20.   Kaminski, D. A., Wei, C., Qian, Y., Rosenberg, A. F. & Sanz, I. Advances in
Human B Cell Phenotypic Profiling. *Front. Immunol.* **3,** 302 (2012).

21.   Flores-Montero, J. *et al.* Immunophenotype of normal vs. myeloma plasma cells:
Toward antibody panel specifications for MRD detection in multiple myeloma.
Cytometry B. Clin. Cytom. (2015). doi:10.1002/cyto.b.21265

22.   Larimore, K., McCormick, M. W., Robins, H. S. & Greenberg, P. D. Shaping of
human germline IgH repertoires revealed by deep sequencing. *J. Immunol.* **189,**
3221–30 (2012).

23.   Wasserman, R. *et al.* The pattern of joining (JH) gene usage in the human IgH
chain is established predominantly at the B precursor cell stage. *J. Immunol.* **149,**
511–6 (1992).

24.   Theofilopoulos, A. N. & Bona, A. N. *The Molecular Pathology of Autoimmune*
Diseases. (CRC Press, 2002).

25.   Mortuza, F. Y. Immunoglobulin heavy-chain gene rearrangement in adult acute
lymphoblastic leukemia reveals preferential usage of JH-proximal variable gene
segments. *Blood* **97,** 2716–2726 (2001).

26.   Berman, J. E. *et al.* VH gene usage in humans: biased usage of the VH6 gene in
immature B lymphoid cells. *Eur. J. Immunol.* **21,** 1311–4 (1991).

27.   Rao, S. P. *et al.* Biased VH gene usage in early lineage human B cells: evidence
for preferential Ig gene rearrangement in the absence of selection. *J. Immunol.*
163, 2732–40 (1999).

28.   Georgiou, G. *et al.* The promise and challenge of high-throughput sequencing of
the antibody repertoire. *Nat. Biotechnol.* **32,** 158–68 (2014).

29.   Wine, Y., Horton, A. P., Ippolito, G. C. & Georgiou, G. Serology in the 21st
century: the molecular-level analysis of the serum antibody repertoire. *Curr. Opin.*
Immunol. **35,** 89–97 (2015).

30.   Perlot, T., Alt, F. W., Bassing, C. H., Suh, H. & Pinaud, E. Elucidation of IgH
intronic enhancer functions via germ-line deletion. *Proc. Natl. Acad. Sci.* **102,**
14362–14367 (2005).

31.   Laserson, U. *et al.* High-resolution antibody dynamics of vaccine-induced immune
responses. *Proc. Natl. Acad. Sci. U. S. A.* **111,** 4928–33 (2014).

32.   Glanville, J. *et al.* Precise determination of the diversity of a combinatorial
antibody library gives insight into the human immunoglobulin repertoire. *Proc.*
Natl. Acad. Sci. U. S. A. **106,** 20216–21 (2009).

33.   Wang, C. *et al.* B-cell repertoire responses to varicella-zoster vaccination in
human identical twins. *Proc. Natl. Acad. Sci. U. S. A.* **112,** 500–5 (2015).

34.   Tipton, C. M. *et al.* Diversity, cellular origin and autoreactivity of antibody-
secreting cell population expansions in acute systemic lupus erythematosus. *Nat.*
Immunol. **16,** 755–65 (2015).

35.   Li, W. & Godzik, A. Cd-hit: a fast program for clustering and comparing large sets
of protein or nucleotide sequences. *Bioinformatics* **22,** 1658–9 (2006).

36.   Krzywinski, M. *et al.* Circos: an information aesthetic for comparative genomics.
Genome Res. **19,** 1639–45 (2009).

 FIGURE LEGENDS

Figure 1. Serial sampling, isolation by FACS, and NGS of BM PCs reveal temporal
dynamics of antibody repertoire. (a) Overview of antibody repertoire characterization
method. Serial sampling of human bone marrow plasma cells (BM PCs) over 6.5 years
(left). Analysis of individual genes, gene combinations, and CDR-H3s (center) show
temporally stable expression of persistent entities (right). (b) Representative
fluorescence-activated cell sorting (FACS) gates of BM PCs (CD138+, CD38++) isolated
from bone marrow mononuclear cells (BMMCs). (c) Sample collection timeline and
summary of cell counts, sequencing reads, and unique CDR-H3s.

Figure 2. **Individual gene use frequencies for Donor 1 are temporally stable.** (a-c)
IGHV (a), IGHD (b), and IGHJ (c) gene usage frequency over time. Plots are sorted by
decreasing mean frequency. Only gene calls that appear in all timepoints are shown. (d)
Mean frequency of IGHV gene use. Error bars are standard deviation.

Figure 3. Usage frequencies of gene combinations are temporally stable. IGH V-J
usage frequencies for Donor 1 are shown. Plots are sorted by decreasing mean
frequency. Only gene calls that appear in all timepoints are shown. See Supplementary
Figure 1(a-c) for usage frequencies of IGH V-D, D-J, and V-D-J.

Figure 3. Gene combinations do not preferentially associate and are instead
randomly assorted. (a-d) Spearman's rank correlation of expected versus observed
IGH V-D (a), V-J (b), D-J (c), and V-D-J (d) gene combination frequencies. Expected (by
random association) frequencies are calculated as products of the frequencies of the
individual component genes. Diagonal lines in red indicate no difference between the
expected and observed frequencies.

Figure 5. Usage frequencies of persistent clonotypes are temporally stable. (a)
Circos plot of shared clonotypes between adjacent timepoints (top). Circos plot of the
persistent clonotypes across all timepoints (bottom). Each band in the outermost
perimeter represents the clonotypes found in a given timepoint, sorted by decreasing
expression. The inner curved lines indicate the same clonotype shared by two
timepoints. Green indicates high expression; purple, low expression; with lighter colors

indicating intermediate expression. (b) Gene usage frequency over time of the 23
persistent clonotypes (see Methods) found in all timepoints. Plots are sorted by
decreasing mean frequency. Gene names (for IGHV and IGHJ), representative amino
acid sequences, and isotype are above each plot.

Figure 6. Gene and gene combination use frequencies correlate between Donor 1
and Donor 2. (a) Spearman's rank correlation of individual gene frequencies between
the two donors: IGHV (top), IGHD (center), and IGHJ (bottom). (b-e) Spearman's rank
correlation of combination gene frequencies between the two donors: V-D (b), V-J (c), D-
J (d), and V-D-J (e). (a-e) Red lines indicate least squares regression.

Serial Collection of Human BM PCs   **Multiple resolution analysis of IgH repertoire**   **Global repertoire stability**

FACS Sort for Plasma Cells   **High Throughput Sequencing of Ig Heavy Chain**

490   **Figure 1**
491

507 **Figure 2**

Figure 3

Figure 4

Figure 5

602 **Figure 6**
603