Exploring mechanisms that affect coral cooperation: symbiont transmission mode, cell density and community composition

Carly D Kenkel[a,1] and Line K Bay[b,2]

[a]Department of Biological Sciences, University of Southern California, 3616 Trousdale Parkway, Los Angeles, CA 90089, USA

[b]Australian Institute of Marine Science, PMB No 3, Townsville MC, Queensland 4810, Australia

[1]Corresponding author, email: ckenkel@usc.edu; phone: 213-821-1705; fax: 213-740-8123

[2]Email: L.Bay@aims.gov.au

RUNNING TITLE: coral-algal cooperation mechanisms

KEYWORDS: Symbiosis, transmission mode, reef corals, bleaching, carbon translocation

ABSTRACT

The coral symbiosis is the linchpin of the reef ecosystem, yet the mechanisms that

promote and maintain cooperation between hosts and symbionts have not been fully resolved.

We used a phylogenetically controlled design to investigate the role of vertical symbiont

transmission, an evolutionary mechanism in which symbionts are inherited directly from

parents, predicted to enhance cooperation and holobiont fitness. Six species of coral, three

vertical transmitters and their closest horizontally transmitting relatives, which exhibit

environmental acquisition of symbionts, were fragmented and subjected to a two-week thermal

stress experiment. Symbiont cell density, photosynthetic function and translocation of

photosynthetically fixed carbon between symbionts and hosts were quantified to assess

changes in physiological performance and cooperation. All species exhibited similar decreases

in symbiont cell density and net photosynthesis in response to elevated temperature, consistent

with the onset of bleaching. Yet baseline cooperation, i.e. translocation of photosynthate, in

ambient conditions and the reduction in cooperation in response to elevated temperature

differed among species. Although *Porites lobata* and *Galaxea acrhelia* did exhibit the highest

levels of baseline cooperation, we did not observe universally higher levels of cooperation in

vertically transmitting species. *Post hoc* sequencing of the *Symbiodinium* ITS-2 locus was used

to investigate the potential role of differences in symbiont community composition.

Interestingly, reductions in cooperation at the onset of bleaching tended to be associated with

increased symbiont community diversity among coral species. The theoretical benefits of

evolving vertical transmission are based on the underlying assumption that the host-symbiont

relationship becomes genetically uniform, thereby reducing competition among symbionts.

Taken together, our results suggest that it may not be vertical transmission *per se* that

influences host-symbiont cooperation, but genetic uniformity of the symbiont community,

although additional work is needed to test this hypothesis.

INTRODUCTION

Cooperation between species has played a fundamental role in the evolution and

diversification of life (Friesen & Jones 2013; Kiers & West 2015; Maynard Smith &

Szathmary 1995; Moran 2006). In the case of reef-building corals, the intracellular symbiosis

between dinoflagellates in the genus *Symbiodinium* and a calcifying Cnidarian host forms the

basis of one of the most biodiverse and productive ecosystems on the planet (Hatcher 1988;

Knowlton et al. 2010). The process of host calcification, which builds the three dimensional

structure of the reef, is largely powered by symbiont primary productivity (Roth 2014).

However, climate change and other anthropogenic processes threaten reefs because of the

sensitivity of the coral-dinoflagellate symbiosis to environmental stress (Hoegh-Guldberg et

al. 2007; Hughes et al. 2003), indicating that host-symbiont cooperation is not stable over

ecological timescales.

Recent work has suggested that a transition to parasitism may precipitate the

breakdown of the host-symbiont relationship (Baker et al. 2018), but this is not a unique

feature of the coral symbiosis. Across taxa, mutualisms are better defined as a spectrum that

ranges from negative parasitic interactions to mutually beneficial symbioses, both within the

context of a focal inter-species interaction and when comparing relationships across taxa

(Doebeli & Knowlton 1998; Lesser et al. 2013; Nowak et al. 1994; Sachs et al. 2011). Less

well-resolved, particularly for the coral-*Symbiodinium* symbiosis, are the mechanisms that

promote and maintain positive interactions between partners (Lesser et al. 2013; Sachs &

Wilcox 2006).

One major factor predicted to influence levels of cooperation is the mode of symbiont

transmission (Anderson & May 1982; Ebert & Bull 2003). In corals as in other symbioses,

two transmission modes predominate: symbionts can be acquired horizontally from the local

environment, usually during a defined larval stage, or vertically from parents, typically

through the maternal germ line (reviewed in (Bright & Bulgheresi 2010). Virulence theory

predicts that horizontal transmission allows symbionts to adopt selfish strategies, potentially

harmful to the host (Bull 1994). A transition from horizontal to vertical transmission is

predicted to align the reproductive interests of partners (via partner-fidelity feedback *sensu*

(Sachs et al. 2004) and optimize resource sharing to maximize holobiont (the combination of

host and symbiont) fitness (Ebert 2013; Frank 1994; Herre et al. 1999).

Experimental manipulations of transmission mode in other systems have provided

empirical support for a reduction in pathogen virulence under enforced vertical transmission

scenarios (Bull et al. 1991; Dusi et al. 2015; Sachs & Wilcox 2006; Stewart et al. 2005).

Bacteriophages forced into vertical transmission evolved lower virulence and lost the

capacity to transmit horizontally (Bull et al. 1991). Similarly, *Symbiodinium microadriaticum*

under an experimentally enforced horizontal transmission regime proliferated faster within

their *Cassiopea* jellyfish hosts while reducing host reproduction and growth (Sachs & Wilcox

2006). However, studies attempting to quantify the evolutionary consequences of natural

shifts in transmission mode remain rare, with Herre's demonstration of a negative

relationship between vertical transmission and virulence in nematodes that parasitize fig

wasps a notable exception (Herre 1993).

Reef-building corals are a potential system in which to study naturally occurring

transitions in transmission mode in a mutualistic symbiosis. Corals (Cnidaria: Anthozoa:

Scleractinia) are colonial animals that harbour intracellular populations of dinoflagellate

algae in the genus *Symbiodinium*. This symbiosis is considered obligate as the breakdown of

the relationship between host animals and their intracellular *Symbiodinium* algae, commonly

known as coral bleaching, has major fitness consequences for both partners, and can be lethal

(reviewed in (Brown 1997). This inter-species partnership is ancient (evolved ~ 250 MYA

(Stanley & Swart 1995), prolific (600+ coral species worldwide (Daly et al. 2007), and

constitutes the foundation of one of the most bio-diverse ecosystems on the planet. The

majority of coral species (~85%) acquire their *Symbiodinium* horizontally from the local

environment in each generation (Harrison & Wallace 1990). However, vertical transmission

has independently evolved at least four times, such that both transmission strategies can be

exhibited by different coral species within the same genus (Baird et al. 2009b; Hartmann et

al. 2017).

We compared physiological components of cooperation and fitness proxies between

horizontal and vertical transmitters in a phylogenetically controlled design using three pairs

of related coral species exhibiting different strategies: (1) *Galaxea acrhelia* (vertical

transmitter, VT) and *G. astreata* (horizontal transmitter, HT); (2) *Porites lobata* (VT) *and*

Goniopora columna (HT); (3) *Montipora aequituberculata* (VT) *and Acropora millepora*

(HT). Species comparisons were drawn from the same or sister genera and replicate

comparisons from more distantly related clades (Hartmann et al. 2017). Species also

represented a diversity of reproductive modes (e.g. broadcast spawner, brooder), sexual

systems (e.g. hermaphroditic, gonochoric), and morphologies (e.g. massive, branching), and

host different subclades of *Symbiodinium* (Tonk et al. 2013), such that mode of symbiont

transmission was the only consistent difference between pairs (Baird et al. 2009b; Franklin et

al. 2012; Kerr et al. 2011). We quantified changes in symbiont cell density, photosynthetic

function and translocation of photosynthetically fixed carbon between symbionts and hosts.

We defined cooperation as the proportion of photosynthetically fixed carbon translocated to

the host, while the degree of symbiont parasitism was calculated as the difference in the

proportion of photosynthetically fixed carbon translocated to hosts between control and heat-

treated samples at the end of the experiment, *sensu* Baker et al. (2018). Bleaching, or the

reduction in symbiont density in response to sustained thermal stress was used as a proxy for

holobiont fitness. While we observed differences in host-symbiont cooperation, both at a

baseline level and during the onset of bleaching, vertically transmitting species did not

exhibit universally elevated levels of cooperation. Additional *post hoc* analysis of

Symbiodinium ITS-2 diversity among coral species was therefore used to investigate whether

symbiont community composition could better explain physiological trait patterns. Symbiont

community composition did not explain a significant portion of the variation in physiological

components of cooperation or fitness proxies; however, diversity tended to be associated with

the degree of symbiont parasitism at the onset of bleaching, suggesting that reduced genetic

diversity of symbionts, rather than vertical transmission *per se*, may influence host-symbiont

cooperation.

MATERIALS and METHODS

Coral Collection and Acclimation

Fragments from sixty unique coral colonies, ~ 20 cm in diameter, were collected from

reefs on the Central GBR from the 8-22 April 2015 under the Great Barrier Reef Marine Park

Authority permits G12/35236.1 and G14/37318.1, prioritizing collection of focal species

pairs by transmission mode from the same reef environment, including location and depth.

Ten corals of each species, *Galaxea acrhelia* and *G. astreata* were collected from Davies

Reef (18°49.816', 147°37.888', 11 April 2015), ten corals each of *Acropora millepora* and

Montipora aequituberculata were collected from Pelorus Island (18°33.358', 146°30.276', 18

April 2015) and ten corals each of *Goniopora columna* and *Porites lobata* were collected

from Pandora Reef (18°48.778', 146°25.593', 21 April 2015) from depths of <10m across all

reefs, and from the same depth within each site. Corals were transported to the Australian

Institute of Marine Science Sea Simulator facility and placed in shaded outdoor holding tanks

with 0.2 µM filtered flow-through seawater (FSW, 27°C, 150 µmol quanta m^{-2} s^{-1}). Each

individual coral colony was further cut into 6 replicates using a diamond blade band saw and

these fragments were mounted on aragonite plugs using either super-glue or marine epoxy.

On 1 May 2015, fragments were moved into indoor experimental aquaria, which consisted of

12 50-L treatment tanks fitted with 3.5-watt Turbelle nanostream 6015 pumps (Tunze,

Germany) with flow-through filtered seawater (FSW, ~25 liters/hour) at 27°C with Hydra 52

lights (Aqua Illumination, C2 Development, Inc., USA) on a 12:12 light/dark cycle

mimicking natural irradiance patterns, peaking at 130-160 μmol quanta m^{-2} s^{-1} at "midday",

which actually occurred at 10:30 GMT+10. Tanks, coral racks and plugs were cleaned daily.

Beginning on 7 May, at 20 mins before dark (16:30 GMT +10) every day, corals were fed

Artemia naupli at a density of 1-1.5 naupli/5ml and rotifers at a density of 1-3/ml.

Experimental Conditions and Physiological Trait Measurements

Beginning on 28 May 2015, effective quantum yield of *Symbiodinium* photosystem II

(EQY) was measured daily for all experimental fragments using a pulse amplitude modulated

fluorometer (diving-PAM, Walz) fitted with a plastic fibre optic cable (Fig. S1).

Measurements were made using factory settings with a measuring intensity of 12 and a gain

of 5 and taken at peak light intensity (between 09:30 – 1130 GMT +10). EQY values were

used to guide the timing of the final sampling time point (Fig. S2), where a decline reflects an

impact on the photosynthetic condition of the *Symbiodinium* (Ralph et al. 2005) because we

aimed to target the onset of the coral bleaching response rather than the end-point.

On 31 May 2015, temperatures in the heat treatment tanks were increased at a rate of

1°C per day until temperatures reached 31°C (day 4, Fig S3). Sample time points occurred on

day 2 (29°C, 1 Jun), day 4 (31°C, 3 Jun) and day 17 (31°C, 16 Jun, Fig S3). On each

sampling day, one replicate fragment of each colony (n=10) and species (n=6) from each

temperature treatment (n=2; n=120 total per sampling day) were used to measure net

photosynthesis following the two-point method originally described and validated for *A.*

millepora in Strahl et al. (2015). It is important to note that this method was not validated for

additional species or under the experimental conditions used in the present study, nor were

species specific photosynthesis-irradiance curves quantified to determine an appropriate

saturating irradiance. Briefly, corals were incubated in enclosed 600-ml acrylic chambers at

their respective treatment temperatures and light levels for 1.5 h. Chambers were placed onto

custom built tables with rotating magnets, which served to power stir bars within each

chamber to facilitate water mixing. For each run, four chambers without corals were used as

blanks to account for potential changes in oxygen content due to the metabolic activity of

other microorganisms in the seawater. For net photosynthesis measures, the O_2 concentration

of the seawater in each chamber was measured at the end of the run using a hand-held

dissolved oxygen meter (HQ30d, equipped with LDO101 IntelliCAL oxygen probe, Hach,

USA). Values from blank chambers were subtracted from measures made in coral chambers

and the subsequent rate of net photosynthesis was related to coral surface area, calculated in

$\mu g\ O_2/cm^2/min$.

To measure the fraction of autotrophically derived carbon translocated to host

animals, five colony fragments of each species from each treatment (n=30 control, n=30 heat)

were placed into 18-L of FSW with a 5-W aquarium pump for circulation and ^{14}C-

bicarbonate (specific activity: 56 mCi/mmol) was added to a final concentration of 0.28

μCi/ml. Corals were incubated for 5 hours in experimental tanks which experienced the

normal experimental irradiance profile from 09:00 – 1400 GMT +10, rinsed with flow-

through FSW for one hour to remove remaining unfixed ^{14}C then snap frozen in liquid

nitrogen.

Tissue was removed from snap frozen coral skeletons using an air gun and

homogenized for 60 s using a Pro250 homogenizer (Perth Scientific Equipment, AUS). A

300-µl aliquot of the tissue homogenate was fixed with 5% formalin in FSW and used to

quantify *Symbiodinium* cell density. The average cell number was obtained from four

replicate haemocytometer counts of a 1-mm^3 area and cell density was related to host protein

content (as assessed below) and expressed as cells/mg host protein. Although surface area is

common used as the normalization factor for *Symbiodinium* cell density, it has been

recognized that areal abundance does not account for differences in host biomass (reviewed

in (Cunning & Baker 2014). We found that normalization to soluble host protein more

accurately reflected this difference in biomass, given that tissue thickness is significantly

greater in *Goniopora columna* and not adequately accounted for by skeletal surface area

normalization alone (Fig. S4). We determined the degree of bleaching, or the reduction in

symbiont density in response to sustained thermal stress, by calculating the difference in

symbiont cell densities between heat-treated and paired control samples following the full 17

days of experimental treatment and used this value as a proxy of holobiont fitness.

An additional 1-ml aliquot of holobiont homogenate was frozen at -20°C. The

remaining homogenate was centrifuged for 2 min at 3500 rcf to separate host and symbiont

fractions and 2-ml of the host tissue slurry was frozen at -20°C. Total soluble protein was

quantified for host tissue samples in duplicate using a colorimetric assay (Bio-Rad Protein

Assay Kit II, USA) following the manufacturer's instructions. Coral skeletons were rinsed

with 5% bleach then dried at room temperature. Skeletal surface area was quantified using

the single wax dipping method (Veal et al. 2010) and skeletal volume (used to standardize

respirometry chamber volumes) was determined by calculating water displacement in a

graduated cylinder.

Sample radioactivity was determined using a liquid scintillation counter (Tri-Carb

2810TR v2.12, Perkin Elmer, USA). Triplicate host and duplicate holobiont 300 µl tissue

homogenate aliquots were mixed with 3.5 ml Ultima Gold XR liquid scintillation cocktail

(Perkin Elmer, USA). Samples were temperature and light adapted for 1 hour and then

counted for 1.5 min using the default parameters. Counts per minute were converted to

disintegrations per minute (DPM) using a standard curve derived from a ^{14}C Ultima Gold

Quench Standards Assay (Perkin Elmer, USA). Technical replicates were averaged. Host

DPM values were divided by holobiont DPM values to yield the fraction of autotrophically

derived carbon shared between partners, which we defined as our metric of host-symbiont

cooperation. The degree of symbiont parasitism was calculated as the difference in this

proportion of photosynthetically fixed carbon translocated to hosts between heat-treated and

paired control samples following the full 17 days of experimental treatment, *sensu* Baker et

al. (2018).

To identify major *Symbiodinium* clades hosted by focal species, DNA was extracted

from symbiont fractions of the tissue homogenate for each replicate colony of each species

(n=60, all control tank fragments) using Wayne's method (Wilson et al. 2002). A restriction

digest of the LSU region of *Symbiodinium* rRNA (Baker & Rowan 1997; Palstra 2000)

consistently revealed single bands indicating the dominance of a single *Symbiodinium* clade

for four of the six species (*A. millepora*, *M. aequituberculata*, *G. columna* and *P. lobata*)

whereas communities in *G. astreata* and *G. acrhelia* appeared more variable (Fig. S5).

Therefore, DNA was pooled in equal proportions by species (n=6 samples) to identify

general species-specific communities using amplicon sequencing of the ITS2 region of

Symbiodinium rRNA. Additional DNA samples for each of the *G. astreata* and *G. acrhelia*

colony replicates (n=20 samples) were also submitted for sequencing at the Genome

Sequencing and Analysis Facility at the University of Texas at Austin. Given the high

diversity subsequently observed in pooled samples of *P. lobat*a and *M. aequituberculata,*

additional colony replicate samples of each (n=3 and n=5, respectively) were later submitted

for sequencing at Oregon State University's Center for Genome Research and Biocomputing

to compare symbiont community composition from individual samples.

Amplicon Sequencing Analysis

For sample libraries prepared and sequenced at UT Austin's GSAF, the ITS2 primers

of Pochon et al. (Pochon et al. 2001) were used. For colony replicate samples sequenced at

OSU's CGRB (individual *P. lobata* and *M. aequituberculata* samples), the ITS2 primers of

LaJeunesse (LaJeunesse 2002) were used to prepare libraries. All sequencing libraries were

subsequently analyzed together. Prior to analysis, raw read data was filtered to remove reads

which contained Illumina sequencing adapters or did not begin with the correct ITS2

amplicon primer sequence using the BBMAP package ver. 37.75

(http://sourceforge.net/projects/bbmap/). The DADA2 pipeline (Callahan 2016) implemented

in R (Team 2017) was then used to infer sequence variants. Read data was analyzed

according to the following tutorial (https://benjjneb.github.io/dada2/tutorial.html) with the

following modifications: ITS2 primers were trimmed from the beginning of each read and

forward and reverse reads were truncated at 210bp and 160bp respectively to remove low

quality bases at the end of reads. Additional paired end reads were discarded if they exhibited

more than one expected error or when a quality score of 2 or less was encountered. In

addition, when inferring sequencing variants using the *dada*() command, the BAND_SIZE

flag was set to 32 as is recommended for ITS data

(https://benjjneb.github.io/dada2/tutorial.html). The distribution of raw read data per sample

and reads lost during quality filtering and processing steps can be found in Table S1. Post-

clustering curation of identified amplicon sequence variants (ASVs) was accomplished with

the LULU algorithm, using the default parameters (Frøslev et al. 2017). The MCMC.OTU

package (Green et al. 2014) was then used to remove sample outliers with low counts overall

(z-score <-2.5) and remaining ASVs of abundance less than 0.001 (Quigley et al. 2014) prior

to statistical analysis. In total, 12 ASVs were identified across samples that satisfied these

criteria. These high confidence sequence variants were taxonomically classified through a

blast search against the GeoSymbio ITS2 database

https://sites.google.com/site/geosymbio/downloads (Franklin et al. 2012), and the best match

was recorded. In cases where variants matched equally well to multiple references, all top hits

were reported (Table S2). Resequencing of the individual *P. lobata* and *M. aequituberculata*

samples indicated that the *P. lobata* pooled sample was contaminated at some stage of the

sequencing process, as individual samples did not match the host pool (Fig. S6). In addition,

two *Galaxea* samples (Gacr1 and Gast4) appeared mislabeled, based on the pattern of

symbiont community differences among species (Fig S6) and we removed these samples

prior to further analysis. To account for differences between individual samples and species

pools, we calculated the normalized mean abundance of each variant across individual

samples within a species and replaced the pooled sequence sample with this *in silico* pooled

value, when available.

Statistical Analyses

All statistical analyses were performed in R, version 3.4.2 (Team 2017). A series of

linear mixed models were used to determine the effect of species, temperature treatment and

sampling day on physiological metrics. We used a conservative approach to evaluate the

effect of transmission mode on physiological metrics. Rather than modelling fixed effects of

transmission mode directly, we modelled the fixed effects of species (levels: Amil, Maeq,

Gcol, Plob, Gast, Gacr), sampling day (levels: day 2, day 4, day 17), temperature treatment

(levels: heat, control) and all possible interactions on *Symbiodinium* cell density, net

photosynthesis, and host-symbiont resource sharing using the *lme* command of the *nlme*

package (Pinheiro et al. 2013), including source coral colony identity nested within reef of

origin as a scalar random factor. All models were assessed for normality of residuals and

homoscedasticity. Symbiont densities required a log-transformation to meet normality

assumptions. To assess goodness-of-fit, we used the function *rsquared.glmm* to calculate the

conditional R^2 value for each of our mixed models (Johnson 2014). Significance of fixed

factors within models was evaluated using Wald tests and Tukey's post-hoc tests were used

to evaluate significance among levels within factors and interactions when warranted. In

these models, a significant effect of transmission mode would have been detected first as a

significant difference among species, but significant differences in the hypothesized direction

between each pair of vertical and horizontal transmitters in the subsequent Tukey's tests were

also required to satisfy reporting an effect of transmission mode overall.

To determine the impact of specific *Symbiodinium* community characteristics on

responses among species, the DESeq package (Anders & Huber 2010) was used to construct

a series of generalized linear models to evaluate differences in in the abundance of each ASV

by species with respect to aspects of host-symbiont cooperation: mean initial symbiont

density (Day 2, heat samples); the mean fraction of carbon shared by symbionts under

ambient conditions (Day 2, 4, 17 control samples), the mean difference in the fraction of

carbon shared by symbionts during bleaching (Day17, heat vs. control samples) and the mean

difference in symbiont density during bleaching (Day17, heat vs. control samples). Models

were run for 30 iterations and for models that did not converge, p-values were converted to

NAs, the standard notation for missing data, prior to applying a multiple test correction

(Benjamini & Hochberg 1995).

Predictive relationships between *Symbiodinium* cell density, the degree of bleaching,

cooperation, and symbiont parasitism, in addition to *Symbiodinium* community diversity

(quantified using the inverse Simpson index, (Magurran 2004)) and composition (the

proportion of Clade D) were explored at the species level using a series of linear models.

When a significant difference was detected among species for physiological trait

comparisons, the model was re-run within each species and a multiple test correction using

the method of Benjamini & Hochberg (1995) was applied. Analyses of symbiont community

composition were run both using the full six species averages and excluding the species for

which individual sample sequencing replicates were not available (*A. millepora* and *G.*

columna). As models did not substantially differ, we report results for the full six species.

RESULTS

Physiological metrics of thermal tolerance and cooperation

Significant fixed effects of species, temperature treatment, sampling time and the

time*treatment interaction were detected for the *Symbiodinium* cell density model

(R^2_{GLMM}=0.53, Table S3). The fixed difference among species was driven by the low

symbiont density on average in *P. lobata*, which differed significantly from densities in *A.*

millepora, *M. aequituberculata*, *G. astreata* and *G. acrhelia* (Tukey's HSD < 0.05, Fig. 1).

No differences were detected in post-hoc tests between focal species pairs by symbiont

transmission mode. Across species, cell densities did not differ between control and heat-

treated corals on days 2 and 4, but were reduced in heat treated corals on day 17, by 5.4 x 10^5

cells/mg host protein on average (Tukey's HSD < 0.001, Fig. 1).

The model for changes in the rate of net photosynthesis identified significant effects

of species, sampling time and the temperature treatment*time interaction (R^2_{GLMM}=0.53,

Table S4). Average net O_2 production rate was highest in *M. aequituberculata,* differing

significantly from *A. millepora*, *G. columna*, *G. acrhelia,* and *P. lobata* Tukey's HSD < 0.05,

Fig. 1), and lowest in *G. columna*, significantly more so in comparison to *M.*

aequituberculata, P. lobata, G. astreata and G. acrhelia (Tukey's HSD < 0.001, Fig. 1). Net

O_2 production rate was also lower on average in *A. millepora* than in *M. aequituberculata* and

G. astreata (Tukey's HSD < 0.01, Fig. 1). Considering differences between focal species

pairs by symbiont transmission mode, the rate of net photosynthesis was significantly higher

in the vertically transmitting *M. aequituberculata* and *P. lobata* than in their horizontally

transmitting counterparts, *A. millepora* and *G. columna*. However, it is important to note that

absolute rates of oxygen production among focal species are much lower than previously

reported values for corals (e.g. (Anthony et al. 2008). Corals were acclimated to a common

light environment, but data on species-specific photosynthesis-irradiance curves are lacking.

Consequently, the differences among species may be influenced by variation in species-

specific photobiology.

Figure 1. *Symbiodinium* cell density, expressed as cells per mg of host protein, rate of net
photosynthesis, expressed as mg O_2 / cm^2 / min and the proportion of photosynthetically
fixed carbon translocated to the host, calculated as the ratio of total radioactivity in DPM in
host to holobiont tissue fractions of focal coral species (Amil: *Acropora millepora*, Maqe:
Montipora aequituberculata, Gcol: *Goniopora columna*, Plob: *Porites lobata*, Gast:
Galaxea astreata, Gach: *Galaxea acrhelia*) under control (27°C, blue circles) and elevated
(31°C, red triangles) temperature following 2, 4 and 17 days of treatment.

Over the course of the experiment, net photosynthesis rates were elevated in heat-

treated corals on days 2 and 4 relative to their respective controls (day2: 0.09 µg O_2/cm^2/min;

day 4: 0.06 µg O_2/cm^2/min; Tukey's HSD < 0.01, Fig. 1) and reduced in heat-treated corals

on day 17, by 0.1 µg O_2/cm^2/min (Tukey's HSD < 0.001, Fig. 1).

The proportion of carbon photosynthetically fixed by symbionts then translocated to

hosts was significantly different among species, temperature treatment, sampling time, the

treatment*time interaction and the species*treatment interaction (R^2_{GLMM}=0.73, Table S5).

Carbon sharing in ambient conditions was significantly lower in *G. columna* in comparison to

all other species (Tukey's HSD < 0.05, Fig. 1) and highest in the vertically transmitting *G.*

acrhelia and *P. lobata*, significantly more so than in *M. aequituberculata* and their

horizontally transmitting counterparts, *G. astreata* and *G. columna* (Tukey's HSD < 0.01,

Fig. 1). Over time, no significant differences were detected in carbon sharing between control

and heat-treated corals on days 2 and 4, but proportional translocation was significantly lower

in heat-treated corals on day 17 (Tukey's HSD < 0.001, Fig. 1). Relative to controls, carbon

sharing decreased slightly under heat treatment on average in all species save *G. columna*,

though the only significant difference was observed in *G. acrhelia* (heat vs. control Tukey's

HSD < 0.01, Fig. 1). Although the species*treatment*time interaction was not significant in

the final model, this was likely driven by the decrease in heat-treated corals on day 17, which

is most evident in *A. millepora*, *M. aequituberculata, G. astreata* and *G. acrhelia*.

Relationships between symbiont density, degree of bleaching and carbon translocation

Symbiont cell density did not explain a significant proportion of the variance in

carbon translocation under ambient conditions (control corals on days 2, 4 and 17) due to

differences among species ($F_{1,5}$ = 25.57, P < 0.001, Fig. 2). Within species, carbon

translocation increased with increasing symbiont density in *G. acrhelia* (R^2=0.22) but this

relationship became non-significant after applying a multiple test correction.

Figure 2. The proportion of photosynthetically fixed carbon translocated to the host in ambient conditions (27°C) as a function of *Symbiodinium* cell density by species (Amil: *Acropora millepora*, Maqe: *Montipora aequituberculata*, Gcol: *Goniopora columna*, Plob: *Porites lobata*, Gast: *Galaxea astreata*, Gach: *Galaxea acrhelia*).

Symbiont cell density in heat-treated corals on day 2 predicted a small portion of the variance in bleaching intensity on day 17 (calculated as the difference in cell density between paired control and heat-treated coral fragments). Greater bleaching was associated with higher initial symbiont cell densities (β = -0.34, R^2 = 0.06, P = 0.04, Fig. 3A) and this relationship did not differ significantly among species. A similar, but non-significant trend was also observed between bleaching on day 17 and symbiont cell density on day 4 (β = -0.24, R^2 = 0.04, P = 0.07).

No significant relationship was detected between the degree of bleaching and the degree of parasitism (calculated as the difference in carbon translocation between heat and control treated samples of paired corals by source colony) at the end of the experiment (Fig. 3B). Nor was the degree of parasitism explained by differences in symbiont cell density of heat-treated corals on day 2, but relationships differed among species ($F_{1,5}$ = 6.88, P < 0.001). In *P. lobata*, a higher symbiont density on day 2 was associated with lower parasitism, or an increase in proportional carbon translocation in heat-treated corals on day 17, and this relationship remained marginally significant even after applying a multiple test correction (R^2 = 0.89, P = 0.06, Fig 3C).

Figure 3. Relationships between changes in symbiont cell density and the proportion of photosynthetically fixed carbon translocated to hosts. (A) Bleaching intensity, calculated as the change in symbiont cell density between control (27°C) and heat-treated (31°C) samples on day 17 as a function of symbiont cell density in heat-treated corals on day 2. (B) The degree of symbiont parasitism, calculated as the difference in the proportion of photosynthetically fixed carbon translocated to hosts between control and heat-treated samples on day 17. (C) The degree of symbiont parasitism as a function of symbiont cell density in heat-treated corals on day 2. Regression line shown for Plob only. Amil: *Acropora millepora*, Maqe: *Montipora aequituberculata*, Gcol: *Goniopora columna*, Plob: *Porites lobata*, Gast: *Galaxea astreata*, Gach: *Galaxea acrhelia*.

Figure 4. Normalized percent abundance of sequence variants by pooled sample. The 'is' label indicates samples for which sequence variant abundances across individually sequenced samples were pooled *in silico*. The best BLAST match against the GeoSymbio ITS2 database (Franklin et al. 2012) for each sequence variant is also reported.

The role of Symbiodinium *community composition*

Across all individual and pooled samples, 12 ASVs of sufficient representation

(>0.1% abundance, *sensu* (Quigley et al. 2014) were identified and consisted of clade C and

D-type *Symbiodinium* (Table S2, Fig. S6). *Symbiodinium* community composition varied

among species (Fig. 4) but no relationships were detected between the abundance of

individual ASVs and symbiont density, degree of bleaching or carbon translocation. Nor did

we detect any relationship between *Symbiodinium* community diversity overall and symbiont

density on day 2, mean carbon translocation under ambient conditions or the degree of

bleaching at the end of the experiment (Fig. 5A,B,D). A marginally significant negative

relationship was detected between community diversity and the degree of parasitism at the

end of the experiment, where a more diverse community was associated with a greater degree

of symbiont parasitism ($R^2 = 0.56$, P = 0.054, Fig 5C). We did not detect any relationships

between the percent of Clade D in the symbiont community and symbiont density on day 2,

mean carbon translocation under ambient conditions, or the degree of parasitism or bleaching

at the end of the experiment (Fig. 5E-H), but this was likely due to the fact that the abundance

of D was very low in all species except *G. acrhelia*. *G. acrhelia,* however*,* did exhibit the

highest carbon translocation rates under ambient conditions and the greatest transition

towards parasitism on day 17 (Fig. 5F,G).

Figure 5. Relationships between changes in symbiont cell density or the proportion of photosynthetically
fixed carbon translocated to hosts as a function of symbiont community diversity, expressed as the inverse
Simpson index (A-D) or the normalized proportion of Clade D *Symbiodinium* (E-H) across species. For the
dependent variables: (A,E) mean symbiont cell density in heat-treated corals (31°C) on day 2; (B,F) mean
proportion of photosynthetically fixed carbon translocated to hosts under ambient conditions (27°C); (C,G)
The degree of symbiont parasitism, calculated as the difference in the proportion of photosynthetically fixed
carbon translocated to hosts between control and heat-treated corals on day 17.

DISCUSSION

Understanding variation in the degree of cooperation between corals and their

Symbiodinium will be critical for assessing survival potential among species and populations

in the face of increasing environmental change (Lesser et al. 2013). As in other mutualisms

(Ebert 2013; Frank 1994; Herre et al. 1999), vertical transmission has been proposed as an

evolutionary mechanism for enhancing holobiont fitness in the Cnidarian-algal symbiosis

(Putnam et al. 2012). However, we did not observe universally consistent differences in

cooperation between vertical and horizontally transmitting species. The vertically

transmitting *P. lobata* and *G. acrhelia* exhibited the highest levels of carbon translocation in

ambient conditions, which we interpret as symbiont-host cooperation, significantly more so

than their respective horizontally-transmitting counterparts, *G. columna* and *G. astreata*; but

cooperation was not different between *M. aequituberculata* and *A. millepora*, and tended to

be higher in the latter horizontally transmitting species. However, species-specific

photosynthesis-irradiance curves were not measured and the potential for interactions

between photophysiology and baseline rates of carbon translocation must also be explored.

We also expected that the degree of breakdown in the host-symbiont relationship under heat

stress, which we interpret as a transition towards parasitism *sensu* (Baker et al. 2018), would

be comparatively intensified in horizontally transmitting species, but again, this was not the

case. *P. lobata* did exhibit the least change in carbon translocation in spite of showing the

same signs of bleaching as other species, but there was no difference in comparison to *G.*

columna which also largely maintained its baseline translocation under elevated temperature.

In addition, the greatest relative declines in the proportion of carbon translocated from

symbionts to hosts actually occurred in the other two vertically transmitting species, *G.*

acrhelia and *M. aequituberculata*.

Although we did not find significant support for the role of vertical transmission we

still observed significant differences in cooperation among our six focal species, both

baseline differences under ambient conditions and in the degree of transition towards

parasitism under elevated temperature stress. We therefore conducted a series of *post hoc*

analyses to explore other putative drivers of differential cooperation and thermal tolerance:

differences in *Symbiodinium* cell density and/or in symbiont community composition.

The density of symbiont cells was recently proposed as a major driver underpinning

the degree of cooperation between coral hosts and symbionts and the functional response of

the coral symbiosis to environmental stressors (Cunning & Baker 2014). Studies in other

species have shown that corals with greater initial *Symbiodinium* cell densities, as quantified

by the symbiont to host cell ratio, are subsequently associated with greater bleaching severity

in response to elevated thermal stress (Cunning & Baker 2013; Silverstein et al. 2014). This

association has been hypothesized to result from the proportional increase in oxidative

cellular stress: more symbionts yield more reactive oxygen species when the photosynthetic

machinery is overloaded (Cunning & Baker 2014), though the recent work of Baker et al.

(Baker et al. 2018), adds another potential explanation. They showed that under non-limiting

nutrient conditions, *Symbiodinium* cell division in *Orbicella faveolata* actually increased in

response to sub-bleaching temperature exposure, but that the metabolic costs were borne by

the coral hosts (Baker et al. 2018), supporting the prediction that a transition to parasitism

precedes unsustainable proliferation of the symbiont community which ultimately results in

bleaching (Wooldridge 2009).

In examining relationships between symbiont cell densities, the intensity of the

bleaching response at the end of our 17-day temperature exposure and carbon translocation

rates, we did observe a negative relationship between initial symbiont cell density on days 2

and 4 and subsequent bleaching response on day 17, which did not significantly differ among

our six focal species. However, initial symbiont cell densities did not predict the degree of the

subsequent transition to parasitism. In fact, the majority of species showed a trend in which

greater initial cell densities were associated with a greater maintenance of cooperation under

bleaching stress. We also found no relationship between the degree of bleaching and the

degree of parasitism on day 17, nor did symbiont cell density explain variation in cooperation

among species in ambient conditions. For some species, cooperation tended to decrease with

increasing density of symbionts, whereas in others it increased. Taken together, these

observations support the association between initial symbiont cell density and subsequent

bleaching intensity, but disagree with the proposed role of an alteration of host-symbiont

cooperation in mediating the bleaching response.

The discrepancy in results among studies may be due to the importance of nutrient

enrichment for observing a parasitic increase in symbiont communities, or in the difference in

study duration and sampling design. Corals were exposed to a constant light environment,

which likely did not provide a saturating irradiance across species. Consequently, alteration

of carbon translocation as a function of species-specific photophysiology may explain

baseline differences in ambient conditions, and future work should aim to test this hypothesis.

In addition, while corals received supplemental feeding throughout the duration of this

experiment, which likely introduced organic nitrogen, we did not explicitly manipulate

inorganic nitrogen levels. It is also possible that variation among hosts in their ability to limit

their symbionts' nitrogen supply may have influenced the observed variation in the degree of

parasitism (Cunning et al. 2017; Wooldridge 2009), though additional studies are needed to

test this. In addition, Baker et al. (Baker et al. 2018) exposed corals to a +5°C temperature

ramp over 8 days, sampling once 24 hours after the final temperature was reached, whereas

the present study increased temperatures by +4°C over 5 days and maintained that difference

for an additional 12 days, sampling at three time points, comparatively earlier and later. We

did not observe an initial increase in symbiont cell density or decrease in carbon translocation

on days 2 and 4 under elevated temperature, the experimental time-frame most analogous to

that of Baker et al. (Baker et al. 2018). It is possible that these dynamics occurred during a

time frame in which we did not sample; however, our final results argue against this

explanation. On day 17, we observed symptoms of bleaching that did not differ across

species: symbiont cell densities and rates of net photosynthesis were uniformly decreased.

However, the transition to parasitism was not uniform, as some species exhibited significant

differences in carbon translocation in response to heat stress whereas others did not. We

therefore conclude that while symbiont density alone may be a reasonable predictor of the

potential for observing a bleaching response under elevated temperature, it does not predict

cooperative dynamics that likely also influence holobiont fitness.

We therefore also explored the role of symbiont community diversity on bleaching

stress and cooperation. Predictions regarding the cooperative and fitness benefits of evolving

vertical transmission are based on the assumption that the host-symbiont relationship

becomes exclusive: symbiont population sizes are substantially reduced, resulting in genetic

uniformity, more rapid co-evolution of partner traits and reduction in intra-symbiont

community competition (Herre et al. 1999; Maynard Smith & Szathmary 1995). In this case,

it may not be vertical transmission *per se* that influences host-symbiont cooperation, but the

relative diversity of the symbiont community. A prior meta-analysis found that symbiont

specificity was correlated with transmission mode, with horizontally transmitting species

being more likely to interact with generalist symbionts (Fabina et al. 2012). However, the

relationship between transmission mode and overall community diversity was not explored.

Other more recent studies have also shown the potential for ontogenetic shifts in

Symbiodinium community composition of putative vertical transmitters, potentially indicating

the capacity for mixed-mode or cryptic horizontal transmission (Byler et al. 2013; Reich et al.

2017). Our results show that symbiont diversity does not partition by transmission mode.

While communities in the vertically transmitting *M. aequituberculata* and *P. lobata* were

largely uniform, consisting predominantly of C35 and C15-type sequence variants,

respectively, *Symbiodinium* community diversity was more than twice as high in *G. acrhelia*

in comparison to *G. astreata* (1/D = 3.6 vs. 1.7). In addition, community composition in the

horizontally transmitting *G. columna* was also highly uniform, second only to that of *P.*

lobata (1/D = 1.08 and 1.00, respectively).

In exploring the relationship between symbiont diversity at the ITS2 locus and metrics

of host-symbiont cooperation and bleaching independent of transmission mode, we did not

find any formally significant correlations, likely due to the fact that our symbiont community

analysis was limited to the level of differences among the six species, greatly reducing our

statistical power. However, there was a weak negative relationship between community

diversity and the degree of parasitism under thermal stress. Species with the most genetically

uniform symbiont communities, *P. lobata* and *G. columna,* maintained the highest levels of

cooperation in spite of showing signs of bleaching. Yet there was no relationship between

community diversity and baseline differences in cooperation among species under ambient

conditions, as high rates of translocation were observed in species with both the lowest (*P.*

lobata) and highest community diversity (*A. millepora, G. acrhelia*), though again, species-

specific interactions between photophysiology and carbon translocation remain to be

explored.

The presence of particular symbiont types has also been shown to influence holobiont

fitness and carbon translocation. For example, conspecific corals hosting clade D

Symbiodinium exhibit greater thermal tolerance than those hosting C1 or C2-types

(Berkelmans & van Oppen 2006; Jones et al. 2008), however they generally grow more

slowly under non-stressful conditions (Jones & Berkelmans 2010; Little et al. 2004) and

receive less photosynthetically fixed carbon from their symbionts (Cantin et al. 2009). We

found no significant relationships between the proportional abundance of Clade D-type

symbionts and metrics of host-symbiont cooperation or bleaching. Most species had no, or a

low proportion of Clade D, but the species with the greatest proportion of Clade D symbionts

(*G. acrhelia*) exhibited both the highest carbon translocation under ambient conditions and

the greatest transition to parasitism under elevated temperature stress. Similar to our

observations regarding symbiont cell density, these results support prior observations that

Symbiodinium community composition alone is not sufficient to explain variation in

holobiont performance (Abrego et al. 2008; Baird et al. 2009a; Kenkel et al. 2013). However,

we reiterate that our analyses are limited in comparing only averages among species. There

was some variation observed in dominant symbiont types among individual coral colonies

within species (Fig. S6) and a priority for future study should be to investigate whether these

conclusions hold when considering intraspecific variation in symbiont community

composition in addition to these broader interspecific differences.

Quantifying cooperation between symbiotic partners in terms of biologically realistic

costs and benefits remains an outstanding question for many symbioses (Herre et al. 1999).

The transfer of photosynthetically fixed carbon has long been known as a major cooperative

benefit to the coral host as up to 95% of a coral's energy requirements can be met through

this mechanism (Muscatine 1990); however, reciprocal products shared by hosts with their

symbionts remain largely unknown (Yellowlees et al. 2008). Similarly, heterotrophic feeding

can offset the need for symbiont-derived carbon in some species and in these cases other

symbiont-derived metabolites may be more critical for host fitness (Grottoli et al. 2006).

Substantial variation in both intra- and inter-specific bleaching thresholds (Marshall & Baird

2000), suggests that levels of cooperation between host and symbiont may also vary.

Significant work has gone into investigating coral bleaching over the past three decades, yet

fundamental questions remain unresolved (Edmunds & Gates 2003). Ultimately, a greater

understanding of both fine-scale interactions between coral hosts and symbionts and the

evolutionary and ecological mechanisms that maintain and strengthen cooperation will be

essential for managing these ecosystems (Davy et al. 2012; Lesser et al. 2013).

CONCLUSIONS

This study investigated whether corals employing vertical symbiont transmission also exhibit

enhanced cooperation and holobiont fitness. Contrary to theoretical predictions, we did not

find significant support for the role of vertical transmission in spite of significant differences

in cooperation among our six focal species. In a *post hoc* analysis of other drivers, we found

that a greater initial symbiont cell density was associated with a greater bleaching intensity,

but this association did not appear to result from an alteration of host-symbiont cooperation.

Rather, the reduction in cooperation across species at the onset of bleaching was marginally

associated with symbiont community diversity. The theoretical benefits of evolving vertical

transmission are based on the underlying assumption that the host-symbiont relationship

becomes genetically uniform, thereby reducing competition among symbionts. Taken

together, our results suggest that it may not be vertical transmission *per se* that influences

host-symbiont cooperation, but genetic uniformity of the symbiont community, though future

work is needed to directly test this hypothesis.

ACKNOWLEDGEMENTS

The authors gratefully acknowledge the efforts of A Bouriat in maintaining aquaria

and processing coral samples. M Salmon, G Milton, A Severati, C Humphrey implemented

the experimental aquaria design. Coral collection was accomplished with the help of S

Noonan, V Mocellin, A Severati and M Nayfa. P Muir provided advice on coral taxonomic

identification. Comments from J Caley, B Schaffelke and multiple anonymous reviewers

greatly improved this manuscript.

REFERENCES

Abrego D, Ulstrup KE, Willis BL, and van Oppen MJH. 2008. Species-specific interactions between
algal endosymbionts and coral hosts define their bleaching response to heat and light stress.
Proceedings of the Royal Society B-Biological Sciences 275:2273-2282.
10.1098/rspb.2008.0180
Anders S, and Huber W. 2010. Differential expression analysis for sequence count data. *Genome*
Biology 11:R106.
Anderson RM, and May RM. 1982. Coevolution of hosts and parasites. *Parasitology* 85:411-426.
Anthony KRN, Kline DI, Diaz-Pulido G, Dove S, and Hoegh-Guldberg O. 2008. Ocean acidification
causes bleaching and productivity loss in coral reef builders. *Proceedings of the National*
Academy of Sciences of the United States of America 105:17442-17446.
10.1073/pnas.0804478105
Baird AH, Bhagooli R, Ralph PJ, and Takahashi S. 2009a. Coral bleaching: the role of the host. *Trends*
in Ecology & Evolution 24:16-20. 10.1016/j.tree.2008.09.005
Baird AH, Guest JR, and Willis BL. 2009b. Systematic and Biogeographical Patterns in the
Reproductive Biology of Scleractinian Corals. *Annual Review of Ecology Evolution and*
Systematics 40:551-571. 10.1146/annurev.ecolsys.110308.120220
Baker AC, and Rowan R. 1997. Diversity of symbiotic dinoflagellates (zooxanthellae) in scleractinian
corals fo the Caribbean and Eastern Pacific. Proceedings of the 8th International Coral Reef
Symposium. p 1301-1306.
Baker DM, Freeman CJ, Wong JCY, Fogel ML, and Knowlton N. 2018. Climate change promotes
parisitism in a coral symbiosis. *The ISME Journal*. https://doi.org/10.1038/s41396-018-
0046-8
Benjamini Y, and Hochberg Y. 1995. Controlling the false discovery rate - a practical and powerful
approach to multiple testing. *Journal of the Royal Statistical Society Series B-Methodological*
57:289-300.
Berkelmans R, and van Oppen MJH. 2006. The role of zooxanthellae in the thermal tolerance of
corals: a 'nugget of hope' for coral reefs in an era of climate change. *Proceedings of the*
Royal Society B-Biological Sciences 273:2305-2312. 10.1098/rspb.2006.3567
Bright M, and Bulgheresi S. 2010. A complex journey: transmission of microbial symbionts. *Nature*
Reviews Microbiology 8:218-230.
Brown BE. 1997. Coral bleaching: causes and consequences. *Coral Reefs* 16:S129-S138.
Bull JJ. 1994. Perspective: virulence. *Evolution* 48:1423-1437.
Bull JJ, Molineux IJ, and Rice WR. 1991. Selection of benevolence in a host-parasite system. *Evolution*
45:857-882.
Byler KA, Carmi-Veal M, Fine M, and Goulet TL. 2013. Multiple symbiont acquisition strategies as an
adaptive mechanism in the coral Stylophora pistillata. *PLoS ONE* 8:e59596.
Callahan BJ, McMurdie, Paul J, Rosen, Michael J, Han, Andrew W, Johnson, Amy Jo A, Holmes, Susan
P. 2016. DADA2: High-resolution sample inference from Illumina amplicon data. *Nature*
Methods 13:581-583.
Cantin NE, van Oppen MJH, Willis BL, Mieog JC, and Negri AP. 2009. Juvenile corals can acquire more
carbon from high-performance algal symbionts. *Coral Reefs* 28:405-414. 10.1007/s00338-
009-0478-8
Cunning R, and Baker AC. 2013. Excess of algal symbionts increase the susceptibility of reef corals to
bleaching. *Nature Climate Change* 3:259-262.
Cunning R, and Baker AC. 2014. Not just who, but how many: the importance of partner abundance
in reef coral symbioses. *Frontiers in Microbiology* 5. 10.3389/fmicb.2014.00400
Cunning R, Muller EB, Gates RD, and Nisbet RM. 2017. A dynamic bioenergetic model for coral-
Symbiodinium symbioses and coral bleaching as an alternate stable state. *Journal of*
Theoretical Biology 431:49-62. 10.1016/j.jtbi.2017.08.003

Daly M, Brugler MR, Cartwright P, Collines AG, Dawson MN, Fautin DG, France SC, McFadden CS, Opresko DM, Rodriguez E, Romano SL, and Stake JL. 2007. The phylum Cnidaria: A review of phylogenetic patterns and diversity 300 years after Linnaeus. In: Zhang ZQ, and Shear WA, eds. *Linnaeus Tercentenary: Progress in Invertebrate Taxonomy*: Zootaxa, 127-182.

Davy SK, Allemand D, and Weis VM. 2012. Cell biology of Cnidarian-dinoflagellate symbiosis. *Microbiology and Molecular Biology Reviews* 76:229-261.

Doebeli M, and Knowlton N. 1998. The evolution of interspecific mutualisms. *Proceedings of the National Academy of Sciences of the United States of America* 95:8676-8680.

Dusi E, Gougat-Barbera C, Berendonk TU, and Kaltz O. 2015. Long-term selection experiment produces breakdown of horizontal transmissibility in parasite with mixd transmission mode. *Evolution* 69:1069-1076.

Ebert D. 2013. The epidemiology and evolution of symbionts with mixed-mode transmission. *Annual Review of Ecology, Evolution, and Systematics* 44:7.1-7.21.

Ebert D, and Bull JJ. 2003. Challenging the trade-off model for the evolution of virulence: is virulence management feasible? *Trends in Microbiology* 11:15-20.

Edmunds PJ, and Gates RD. 2003. Has coral bleaching delayed our understanding of fundamental aspects of coral-dinoflagellate symbioses? *BioScience* 53:976-980.

Fabina NS, Putnam HM, Franklin EC, Stat M, and Gates RD. 2012. Transmission mode predicts specificity and interaction patterns in coral-Symbiodinium networks. *PLoS ONE* 7:e44970. doi10.1371/journal.pone.0044970

Frank SA. 1994. Genetics of mutualism: the evolution of altruism between species. *Journal of Theoretical Biology* 170:393-400.

Franklin EC, Stat M, Pochon X, Putnam HM, and Gates RD. 2012. GeoSymbio: a hybrid, cloud-based web application of global geospatial bioinformatics and ecoinformatics for Symbiodinium-host symbioses. *Molecular Ecology Resources* 12:369-373.

Friesen ML, and Jones EI. 2013. Modelling the evolution of mutualistic symbioses. In: van Helden J, Toussaint A, and Thieffry D, eds. *Bacterial Molecular Networks: Methods and Protocols*: Springer Science+Business Media, 481-499.

Frøslev TG, Kjøller R, Bruun HH, Ejrnæs R, Brunbjerg AK, Pietroni C, and Hansen AJ. 2017. Algorithm for post-clustering curation of DNA amplicon data yields reliable biodiversity estimates. *Nature Communications* 8:1188. 10.1038/s41467-017-01312-x

Green EA, Davies SW, Matz MV, and Medina M. 2014. Quantifying cryptic Symbiodinium diversity with Orbicella faveolata and Orbicella franksi at the Flower Garden Banks, Gulf of Mexico. *PeerJ* 2:e386. 10.7717/peerj.386

Grottoli AG, Rodrigues LJ, and Palardy JE. 2006. Heterotrophic plasticity and resilience in bleached corals. *Nature* 440:1186-1189. 10.1038/nature04565

Harrison PL, and Wallace CC. 1990. Reproduction, dispersal and recruitment of scleractinian corals. In: Dubinsky Z, ed. *Ecosystems of the World: Coral Reefs*. New York: Elsevier, 133-207.

Hartmann AC, Baird AH, Knowlton N, and Huang D. 2017. The paradox of environmental symbiont acquisition in obligate mutualisms. *Current Biology* 27:3711-3716.

Hatcher BG. 1988. Coral reef primary productivity: A beggar's banquet. *Trends in Ecology and Evolution* 3:106-111.

Herre EA. 1993. Population structure and the evolution of virulence in nematode parasites of fig wasps. *Science* 259:1442-1445.

Herre EA, Knowlton N, Mueller UG, and Rehner SA. 1999. The evolution of mutualisms: exploring the paths between conflict and cooperation. *Trends in Ecology & Evolution* 14.

Hoegh-Guldberg O, Mumby PJ, Hooten AJ, Steneck RS, Greenfield P, Gomez E, Harvell CD, Sale PF, Edwards AJ, Caldeira K, Knowlton N, Eakin CM, Iglesias-Prieto R, Muthiga N, Bradbury RH, Dubi A, and Hatziolos ME. 2007. Coral reefs under rapid climate change and ocean acidification. *Science* 318:1737-1742. 10.1126/science.1152509

Hughes TP, Baird AH, Bellwood DR, Card M, Connolly SR, Folke C, Grosberg R, Hoegh-Guldberg O, Jackson JBC, Kleypas J, Lough JM, Marshall P, Nystrom M, Palumbi SR, Pandolfi JM, Rosen B, and Roughgarden J. 2003. Climate change, human impacts, and the resilience of coral reefs. *Science* 301:929-933.

Johnson PCD. 2014. Extension of Nakagawa & Schielzeth's R2GLMM to random slopes models. *Methods in Ecology and Evolution* 5:944-946.

Jones A, and Berkelmans R. 2010. Potential Costs of Acclimatization to a Warmer Climate: Growth of a Reef Coral with Heat Tolerant vs. Sensitive Symbiont Types. *PLoS ONE* 5. 10.1371/journal.pone.0010437

Jones A, Berkelmans R, van Oppen MJH, Mieog JC, and Sinclair W. 2008. A community change in the algal endosymbionts of a scleractinian coral following a natural bleaching event: field evidence of acclimatization. *Proceedings of the Royal Society B-Biological Sciences* 275:1359-1365.

Kenkel C, Goodbody-Gringley G, Caillaud D, Davies SW, Bartels E, and Matz M. 2013. Evidence for a host role in thermotolerance divergence between populations of the mustard hill coral (Porites astreoides) from different reef environments. *Molecular Ecology* 22:4335-4348.

Kerr AM, Baird AH, and Hughes TP. 2011. Correlated evolution of sex and reproductive mode in corals (Anthozoa: Scleractinia). *Proceedings of the Royal Society B-Biological Sciences* 278:75-81.

Kiers TE, and West SA. 2015. Evolving new organisms via symbiosis. *Science* 348:392-394.

Knowlton N, Brainard RE, Fisher R, Moews M, Plaisance L, and Caley MJ. 2010. Coral Reef Biodiversity. In: McIntyre A, ed. *Life in the World's Oceans: Diversity, Distribution, and Abundance*. Oxford: Blackwell Publishing Ltd., 65-78.

LaJeunesse TC. 2002. Diversity and community structure of symbiotic dinoflagellates from Caribbean coral reefs. *Marine Biology* 141:387-400.

Lesser MP, Stat M, and Gates RD. 2013. The endosymbiotic dinoflagellates (Symbiodinium sp.) of corals are parasites and mutualists. *Coral Reefs* 32:603-611.

Little AF, van Oppen MJH, and Willis BL. 2004. Flexibility in algal endosymbioses shapes growth in reef corals. *Science* 304:1492-1494.

Magurran AE. 2004. *Measuring biological diversity*. Oxford, UK: Blackwell Science Ltd.

Marshall PA, and Baird AH. 2000. Bleaching of corals on the Great Barrier Reef: differential susceptibilities among taxa. *Coral Reefs* 19:155-163.

Maynard Smith J, and Szathmary E. 1995. *The Major Transitions in Evolution*. Oxford: Freeman.

Moran NA. 2006. Symbiosis. *Current Biology* 16:R866-R871.

Muscatine L. 1990. The role of symbiotic algae in carbon and energy flux in reef corals. In: Dubinsky, ed. *Ecosystems of the world 25: Coral reefs*. New York: Elsiever, 75-87.

Nowak MA, Bonhoeffer S, and May RM. 1994. Spatial games and the maintenance of cooperation. *Proceedings of the National Academy of Sciences of the United States of America* 91:4877-4881.

Palstra F. 2000. Host-endosymbiont specificity in Acropora corals of the Indo-Pacific? James Cook University.

Pinheiro J, Bates D, DebRoy S, Sarkar D, and Team RDC. 2013. nlme: Linear and nonlinear mixed effects models. R package version 3.1-113 ed.

Pochon X, Pawlowski J, Zaninetti L, and Rowan R. 2001. High genetic diversity and relative specificity among Symbiodinium-like endosymbiotic dinoflagellates in soritid foraminiferans. *Marine Biology* 139:1069-1078.

Putnam HM, Stat M, Pochon X, and Gates RD. 2012. Endosymbiotic flexibility associates with environmental sensitivity in scleractinian corals. *Proceedings of the Royal Society B: Biological Sciences* 279:4352-4361.

Quigley KM, Davies SW, Kenkel CD, Willis BL, Matz MV, and Bay LK. 2014. Deep-Sequencing Method for Quantifying Background Abundances of Symbiodinium Types: Exploring the Rare

Symbiodinium Biosphere in Reef-Building Corals. *PLoS ONE* 9:e94297.
doi:10.1371/journal.pone.0094297

Ralph PJ, Larkum AWD, and Kuhl M. 2005. Temporal patterns in effective quantum yield of individual
zooxanthellae expelled during bleaching. *Journal of Experimental Marine Biology and*
Ecology 316:17-28.

Reich HG, Robertson DL, and Goodbody-Gringley G. 2017. Do the shuffle: Changes in Symbiodinium
consortia throughout juvenile coral development. *PLoS ONE* 12:e0171768.
10.1371/journal.pone.0171768

Roth MS. 2014. The engine of the reef: photobiology of the coral-algal symbiosis. *Frontiers in*
Microbiology 5:422. 10.3389/fmicb.2014.00422

Sachs JL, Essenberg CJ, and Turcotte MM. 2011. New paradigms for the evolution of beneficial
infections. *Trends in Ecology & Evolution* 26:202-209.

Sachs JL, Mueller UG, Wilcox TP, and Bull JJ. 2004. The evolution of cooperation. *Quarterly Review of*
Biology 79:135-160.

Sachs JL, and Wilcox TP. 2006. A shift to parasitism in the jellyfish symbiont Symbiodinium
microadriaticum. *Proceedings of the Royal Society B-Biological Sciences* 273:425-429.

Silverstein RN, Cunning R, and Baker AC. 2014. Change in algal symbiont communities after
bleaching, not prior heat exposure, increases heat tolerance of reef corals. *Global Change*
Biology 21:236-249.

Stanley GD, and Swart PK. 1995. Evolution of the coral-zooxanthellae symbiosis during the Triassic: a
geochemical approach. *Paleobiology* 21:179-199.

Stewart AD, Logsdon JM, and Kelley SE. 2005. An empirical study of the evolution of virulece under
both horizontal and vertical transmission. *Evolution* 59:730-739.

Strahl J, Stolz I, Uthicke S, Vogen N, Noonan SHC, and Fabricius KE. 2015. Physiological and ecological
performance differs in four coral taxa at a volcanic carbon dioxide seep. *Comparative*
Biochemistry and Physiology Part A: Molecular & Integrative Physiology 184:179-186.

Team RC. 2017. R: A language and environment for statistical computing. Vienna: R Foundation for
Statistical Computing.

Tonk L, Bongaerts P, Sampayo EM, and Hoegh-Guldberg O. 2013. SymbioGBR: a web-based database
of Symbiodinium associated with cnidarian hosts on the Great Barrier Reef. *BMC Ecology* 13.

Veal CJ, Carmi M, Fine M, and Hoegh-Guldberg O. 2010. Increasing the accuracy of surface area
estimation using single wax dipping of coral fragments. *Coral Reefs* 29:893-897.

Wilson K, Li Y, Whan V, Lehnert S, Byrne K, Moore S, Pongsomboon S, Tassanakajon A, Rosenberg G,
Ballment E, Fayazi Z, Swan J, Kenway M, and Benzie J. 2002. Genetic mapping of the black
tiger shrimp Penaeus monodon with amplified fragment length polymorphism. *Aquaculture*
204:297-309.

Wooldridge SA. 2009. Water quality and coral bleaching thresholds: formalising the linkage for the
inshore reefs of the Great Barrier Reef, Australia. *Marine Pollution Bulletin* 58:745-751.

Yellowlees D, Rees T, and Leggat W. 2008. Metabolic interactions between algal symbionts and
invertebrate hosts. *Plant, Cell & Environment* 31:679-694.

817　**Supplementary Material: Exploring mechanisms that affect coral cooperation: symbiont**

818　**transmission mode, cell density and community composition**

820　Carly D. Kenkel and Line K. Bay

837   Table S1. Raw read counts and reads lost during filtering and clustering with DADA2.

Sample	Raw	Input_DADA2	filtered	denoised	merged	tabled	nonchim
AmilAll	35820	35061	31194	31194	28315	28315	24013
Gach1	74818	74054	67294	67294	66374	66301	62896
Gach10	72354	71449	60322	60322	59915	59451	50573
Gach2	74662	74012	63885	63885	62752	62250	50278
Gach3	35176	34552	25626	25626	25323	25140	22602
Gach4	93883	93234	80667	80667	79625	79064	63253
Gach5	30755	30541	26595	26595	26375	26167	22976
Gach6	40510	39898	34705	34705	34374	34019	31309
Gach7	4472	4332	2643	2643	2611	2598	2528
Gach8	33640	33267	29026	29026	28780	28443	26381
Gach9	40512	40026	34308	34308	34026	33542	31669
GachAll	25286	24821	21705	21705	21399	21246	17958
Gast1	1061	942	582	582	581	581	571
Gast10	33882	33390	30573	30573	30231	30212	29162
Gast2	21197	20863	18934	18934	18696	18696	17876
Gast3	20029	19875	18266	18266	18003	18003	17431
Gast4	16471	16248	13906	13906	13741	13558	12598
Gast5	11943	11730	10196	10196	10172	10172	9182
Gast6	45291	44813	40014	40014	39595	39574	38225
Gast7	33652	33320	29961	29961	29883	29883	26478
Gast8	23892	23539	21491	21491	21312	21287	20885
Gast9	734	621	374	374	374	374	337
GastAll	31819	31304	27836	27836	27425	27367	23894
GcolAll	10196	9940	7972	7972	7839	7839	7661
Maqe10	77568	69196	44998	44998	44705	44698	42679
Maqe6	57788	46772	31275	31275	31160	31114	30225
Maqe6b	51170	27633	17137	17137	17074	17047	16564

Maqe7	29361	23378	16091	16091	15970	15970	15338
Maqe7b	17131	16676	11653	11653	11567	11567	11491
Maqe8	72073	43668	29519	29519	29346	29346	27746
Maqe8b	12976	12418	8388	8388	8314	8314	8214
Maqe9	20282	19533	13318	13318	13226	13226	13131
Maqe9b	59314	31986	19854	19854	19715	19715	19002
MaqeAll	14445	14232	11940	11940	11754	11754	10322
Plob6	16697	5454	3630	3630	725	584	561
Plob6b	28036	3924	2461	2461	1354	1294	1184
Plob7	1469	454	263	263	44	30	3
Plob7b	17267	1798	1135	1135	197	109	109
Plob8b	36191	19086	12600	12600	11986	11940	11837
PlobAll	31619	30869	27527	27527	26407	26341	21754

854   Table S2. Best BLASTn matches to the GeoSymbio ITS2 database (Franklin et al. 2012) for curated

855   amplicon sequence variants (ASVs) detected across samples. Value in % column indicates mean

856   percent abundance across all samples in which the sequence variant, or sub-variants (identified

857   through LULU curation), were identified. Match bit-scores are reported and hits separated by '/' in

858   GeoSymbio column indicate equally high scoring blast matches.

ASV	%	Score	GeoSymbio BLAST
Sq1	31.6	523	C1
Sq2	18.8	501	D1
Sq3	10.6	508	C35(type2)/C26.b1
Sq4	7.0	501	D1a
Sq5	11.4	508	C35a(type2)/C26a/C26
Sq6	6.6	512	C35(type2)/C26.b1
Sq7	3.8	501	D6
Sq8	4.9	523	C3/Cspc
Sq10	2.5	512	C35a(type2)/C26a/C26
Sq18	1.6	520	C15
Sq23	0.8	523	C15
Sq40	0.3	523	C15a

860   Table S3. Wald test statistics for individual factors and interaction terms for the *Symbiodinium* cell

861   density model. Significant terms are shown in bold.

862		numDF	denDF	F-value	p-value
863	-----------------	------	------	-------------	----------
864	(Intercept)	1	270	1.989512e+05	0.0000000
865	**Species**	**5**	**52**	**9.116880e+00**	**0.0000028**
866	**Temp**	**1**	**270**	**1.260908e+01**	**0.0004528**
867	**Day**	**2**	**270**	**6.089511e+00**	**0.0025898**
868	Species:Temp	5	270	1.104898e+00	0.3580558
869	Species:Day	10	270	1.452625e+00	0.1573259
870	**Temp:Day**	**2**	**270**	**1.388694e+01**	**0.0000018**
871	Species:Temp:Day	10	270	4.876186e-01	0.8975041

872   Table S4. Wald test statistics for individual factors and interaction terms for the net photosynthesis

873   model. Significant terms are shown in bold.

	numDF	denDF	F-value	p-value
-----------------	------	------	-------------	----------
(Intercept)	1	270	1648.4748360	0.0000000
Species	**5**	**52**	**40.2840568**	**0.0000000**
Temp	1	270	3.4308423	0.0650808
Day	**2**	**270**	**5.7878681**	**0.0034574**
Species:Temp	5	270	1.2021088	0.3084131
Species:Day	10	270	0.8998245	0.5338268
Temp:Day	**2**	**270**	**36.4032169**	**0.0000000**
Species:Temp:Day	10	270	1.0755283	0.3486423

885   Table S5. Wald test statistics for individual factors and interaction terms for the fraction of DPM

886   shared model. Significant terms are shown in bold.

	numDF	denDF	F-value	p-value
-----------------	------	------	-------------	----------
(Intercept)	1	120	5585.5585773	0.0000000
Species	**5**	**22**	**18.7172944**	**0.0000003**
Temp	**1**	**120**	**18.0748728**	**0.0000423**
Day	**2**	120	**24.4071621**	**0.0000000**
Species:Temp	**5**	**120**	**3.4767095**	**0.0057086**
Species:Day	10	120	0.6096878	0.8031208
Temp:Day	**2**	**120**	**14.5121654**	**0.0000023**
Species:Temp:Day	10	120	0.8893596	0.5453031

899   Figure S1. Daily measures of effective quantum yield (mean EQY±SEM, note that range of error bars
900   is extremely small, in most cases barely exceeding points) of *Symbiodinium* photosystem II by
901   temperature treatment. Bars show the difference in mean EQY between treatments through time
902   (n=720 per bar).

Figure S2. Daily measures of effective quantum yield (EQY±SEM) of *Symbiodinium* photosystem II
by temperature treatment and species.

Figure S3. Temperature profile of experimental treatment tanks (±SD) by date. Bars indicate time of
sampling.

Figure S4. *Symbiodinium* cell density, expressed as cells per cm^2 of coral surface area of focal coral
species (Amil: *Acropora millepora*, Maeq: *Montipora aequituberculata*, Gcol: *Goniopora columna*,
Plob: *Porites lobata*, Gast: *Galaxea astreata*, Gacr: *Galaxea acrhelia*) under control (27°C, blue
circles) and elevated (31°C, red triangles) temperature following 2, 4 and 17 days of treatment.

Figure S5. Taq1 digest of *Symbiodinium* LSU rRNA. Samples are grouped by species pairs. Standard
shows expected banding pattern for *Symbiodinium* genotypes in clades C and D.

Fig S6. Normalized proportional abundance of sequence variants by samples sequenced at UT
Austin's GSAF (All and individual Gast/Gacr samples) and OSU's CGRB (individual Maeq and Plob
samples). Gacr1 and Gast4 were excluded from subsequent analyses as they appear to be mis-labeled
and for analyses comparing differences among species, the 'All' samples for Gacr, Gast, Maeq and
Plob were replaced by *in silico* averages calculated using the individual sample replicates shown here.

REFERENCES

Franklin, E. C., M. Stat, X. Pochon, H. M. Putnam, and R. D. Gates. 2012. GeoSymbio: a hybrid, cloud-
based web application of global geospatial bioinformatics and ecoinformatics for
Symbiodinium-host symbioses. Molecular Ecology Resources 12:369-373.
Veal, C. J., M. Carmi, M. Fine, and O. Hoegh-Guldberg. 2010. Increasing the accuracy of surface area
estimation using single wax dipping of coral fragments. Coral Reefs 29:893-897.