Dendrites of DG granule cells contribute to pattern separation by controlling sparsity

Spyridon Chavlis[1,2], Panagiotis C. Petrantonakis[1], Panayiota Poirazi[1]

[1]Institute of Molecular Biology and Biotechnology (IMBB), Foundation for Research and Technology – Hellas (FORTH), Heraklion, Crete

[2]Department of Biology, School of Sciences and Engineering, University of Crete, Heraklion, Crete

Abstract

The hippocampus plays a key role in pattern separation, the process of transforming similar incoming information to highly dissimilar, non-overlapping representations. Sparse firing granule cells (GCs) in the dentate gyrus (DG) have been proposed to undertake this computation, but little is known about which of their properties influence pattern separation. Dendritic atrophy has been reported in diseases associated with pattern separation deficits, suggesting a possible role for dendrites in this phenomenon. To investigate whether and how the dendrites of GCs contribute to pattern separation, we build a simplified, biologically relevant, computational model of the DG. Our model suggests that the presence of GC dendrites is associated with high pattern separation efficiency while their atrophy leads to increased excitability and performance impairments. These impairments can be rescued by restoring GC sparsity to control levels through various manipulations. We predict that dendrites contribute to pattern separation as a mechanism for controlling sparsity.

Introduction

The hippocampus is known to be involved in memory formation, storage and consolidation (Squire et al., 2004), but its specific functionalities remain a mystery. One such functionality is the ability to rapidly store non-overlapping representations of similar inputs and thereafter, retrieve them given a partial or noisy stimulus. Theoretical models refer to those processes as pattern separation and pattern completion, respectively (Marr, 1971; Treves and Rolls, 1994; Yassa and Stark, 2011; Santoro, 2013). The Dentate Gyrus (DG), in particular, has been proposed to implement pattern separation by sparsifying and orthogonalizing its input, coming mainly from the Entorhinal Cortex (EC), and thereafter, projecting this information to the CA3 area via the mossy fibers (Treves and Rolls, 1994). DG has been hypothesized to separate two distinct but overlapping patterns through the activation of different Granule Cells (GCs), through the expression of different firing rates in identical neuronal populations (Deng et al., 2010) or a combination of the two. While several studies have investigated pattern separation both in rodents (Leutgeb et al., 2004, 2005, 2007) and humans (Kirwan and Stark, 2007; Bakker et al., 2008; Lacy et al., 2011; Motley and Kirwan, 2012), the role of dendrites in this phenomenon remains unknown.

The DG is the first subregion of the hippocampus that receives incoming information from other brain areas. DG principal neurons, the Granule Cells (GCs), receive input from excitatory afferents coming from EC layer II cells and project to the CA3 subregion. In addition, they receive input from other DG excitatory cells, the Mossy Cells (MCs), and various interneurons (Sik et al., 1997) with Basket Cells (BCs) and Hilar Perforant Path associated (HIPP) cells being the most important. MCs form an inhibitory circuit as their axons contact the BCs. The net effect of MC excitation to both GCs and BCs is considered to be inhibitory (Jinde et al., 2012).

Experimental studies have shown that only a small population of GCs, ~5%, are active in a single context (Marrone et al., 2011; Satvat et al., 2011; Danielson et al., 2016), a

phenomenon termed sparse coding (Olshausen and Field, 2004). It has been proposed that

sparse coding in GCs enhances pattern separation by recruiting different subgroups of GCs

to encode similar incoming stimuli (Treves et al., 2008; Petrantonakis and Poirazi, 2014,

2015). Computational models (Santhakumar et al., 2005; Yim et al., 2015) and experimental

studies (Nitz and McNaughton, 2004; Jinde et al., 2012) have proposed that inhibition

controls GC activity which, in turn, mediates pattern separation (Myers and Scharfman,

2009, 2011; Ikrar et al., 2013; Faghihi and Moustafa, 2015).

The ability to perform pattern separation is critical for normal brain functioning and its

impairment is associated with cognitive decline. Diseases such as schizophrenia (Das et al.,

2014) and Alzheimer's Disease (AD) (Ally et al., 2013), where cognitive decline is evident,

are both characterized by pattern separation deficiencies. Interestingly, these conditions are

also characterized by alterations in the anatomical properties of GC dendrites, such as a

decrease in the total dendritic length (Einstein et al., 1994) and spine loss (Jain et al., 2012).

Dendritic growth on the other hand has been associated with pattern separation

enhancements. Specifically, voluntary running was recently shown to enhance pattern

separation and this enhancement was attributed to an increase in the neurogenesis rate that

was accompanied by increased GC dendrite outgrowth in active compared to sedentary

animals (Bolz et al., 2015). These findings suggest that the dendrites of GCs are likely to

play a key role in pattern separation mediated by the DG.

To investigate this possibility, we implemented a morphologically simple, yet biologically

relevant, scaled-down spiking neural network model of the DG. The model consists of four

types of cells (MCs, BCs, HIPP and GCs) modeled as simplified integrate-and-fire neurons.

The GC model alone was extended to incorporate dendrites. The electrophysiological

properties of all cell types were calibrated according to a range of experimental data. An

advantage of using such a simplified approach lies in the small number of parameters that

make it possible to characterize their role in the behavior of the model. Despite its simplicity,

the model exhibits realistic pattern separation under several conditions and explains how

inhibition to GCs provided directly from BCs and indirectly via the inhibitory circuitry through

MCs impact this task, as suggested by a number of recent studies (Myers and Scharfman,

2011; Jinde et al., 2012), thus, supporting its biological relevance. We use the model to

investigate whether and how GC dendrites may contribute to pattern separation.

Results

The Dentate Gyrus model

The DG network model consists of four different neuronal types, namely GCs, BCs, MCs, and HIPP cells. MCs, BCs and HIPP cells are modeled as adaptive exponential I&F (aEIF) point neurons (Brette and Gerstner, 2005). GCs consist of a leaky integrate-and-fire somatic compartment connected to a variable number of active dendritic compartments (3, 6 or 12) whose morphology relies loosely on anatomical data (see Materials and Methods). I&F models were selected primarily because of their correspondence with experimental parameters (e.g. the C_m, R_m, R_{in}) that facilitated constraining with experimental data. Since dentate GCs are known to have 10-15 dendrites (Claiborne et al., 1990), we consider the GC model with 12 dendrites as the control. A schematic illustration of the DG network is shown in Figure 1A.

All computational neuron models were validated against experimental data with respect to their activity and basic electrophysiological properties (Lübke et al., 1998; Bartos et al., 2001; Krueppel et al., 2011). The spiking profiles of the four neuronal model types are depicted in Figure 2 while the respective I-V and I-F curves compared against experimental data are shown in Figure 2-figure supplement 1 and Figure 2-figure supplement 2. The control (12-dendrite) GC model was validated against the experimental data of (Krueppel et al., 2011) with respect to its dendritic input-output function, which was found to be slightly above linearity (see Figure 2-figure supplement 3(A)). Table 1 lists the model and experimental values of basic intrinsic properties for each of these cell types. Overall, the electrophysiological properties of the computational models of neurons included in the DG network are in fair alignment with experimental findings. It should be noted that spiking profiles are not taken into account when estimating pattern separation in the DG network model. A GC model neuron was considered active even if it produced a single spike. Thus, we chose to fit average values rather than temporal profiles of the model neurons.

Pattern separation in the DG model

Following validation of the individual neuron models, we tested the DG network's ability to

implement pattern separation when presented with pairs of inputs characterized by various

degrees of similarity (modeled as overlap in the two activated EC populations). An example

of two such input patterns and their corresponding output is schematically illustrated in

Figure 1B. The network model is deemed capable of separating similar input patterns if the

value of a distance metric f is substantially larger in the DG output (GC activity) compared to

its input (EC cell activity). Pattern separation is primarily estimated by looking at the

differences in the populations of neurons that encode each input ('population distance', f_1).

To account for the possibility of rate-based pattern separation (Deng et al., 2010), we also

simulate the network under conditions where the two input pairs are identical in terms of EC

'population distance' (i.e., $f_{1(input)} = 0$) but differ in their mean firing rates (graphically

depicted in Figure 1C). Pattern separation in this case is measured by looking at differences

in the average firing rate of the neurons encoding both inputs ('rate distance', f_2). The 'rate

distance' is estimated both at the input (EC cells) and the output (GCs) levels (see Materials

and Methods). Pattern separation is successfully performed if the 'rate distance' of the

output ($f_{2(output)}$) is larger than that of the input ($f_{2(input)}$). The network's ability to perform

pattern separation across all (200) input pairs measured with both metrics is shown in Table

2. These results demonstrate that two inputs that are quite similar in their topology (i.e.,

originate largely (f_1) or entirely from the same neurons, perhaps with different firing rates

(f_2)), induce substantially different activation patterns of the GCs that they impinge on

(outputs) in the DG network.

Understanding the role of Inhibition in Pattern separation

After establishing the network's ability to perform pattern separation, we tested its validity

against experimental data with respect to the role of inhibition in this phenomenon. The

network model reproduced the recent findings of Engin et al., whereby inhibition exerted by

BCs was critical for the sparse firing of GCs (Engin et al., 2015). Specifically, removal of all

BCs resulted in an over-excitation of the GC model population (more than 50% of GCs

responded strongly to any input). This over-excitation in turn impaired pattern separation, as

the GC populations responding to the two inputs became nearly identical and fired at very

similar, high frequencies. These findings are also in line with experimental evidence

reporting increased memory interference under conditions of reduced BC activity (Engin et

al., 2015).

Since MCs have also been suggested to control the excitability of DG granule cells (Jinde et

al., 2013), we simulated a complete MC-loss lesion (Figure 3A) as per (Ratzliff et al., 2004).

This manipulation led to an increase in the proportion of active GCs for all input patterns

tested (Figure 3B, 3D), and a decrease in pattern separation efficiency, measured either with

the population (f_1, Figure 3C) (Wilcoxon test, $W_1=2095.5$, $p_1≈10^{-9}$, 95% CI_1 [0.05, 0.09],

$W_2=2242.0$, $p_2≈10^{-11}$, 95% CI_2 [0.06, 0.10], $W_3=2363.0$, $p_3≈10^{-11}$, 95% CI_3 [0.07, 0.11],

$W_4=2444.0$, $p_4≈10^{-16}$, 95% CI_4 [0.08, 0.12]) or rate metrics (f_2, Figure 3E) (Wilcoxon test,

$W=930.0$, $p=0.0276$, 95% CI [0.01, 0.07]). The subscripts (form one to four) in the Wilcoxon

test statistic (W), p-value, confidence interval (CI) correspond to the four experiments with

different input overlaps (see Materials and Methods). The observed decrease in sparsity

under the MC-loss condition was accompanied by small increases in the excitability levels of

GC models. Specifically, for the population coding experiment, the mean GC firing frequency

increased from 3.5 Hz to 4.82 Hz, while for the rate-based coding experiment from 3.75 Hz

to 4.94 Hz for 40 Hz inputs and from 5.24 Hz to 8.07 Hz for 50 Hz inputs. These findings are

in line with the experimental data of (Ratzliff et al., 2004) where MC-loss did not lead to an

over-excitation of GCs.

Taken together, the proposed DG network model a) exhibits single-neuron response

properties that are in good agreement with experimental data, b) implements a connectivity

profile that relies on experimental observations, c) exhibits robust pattern separation and d)

replicates experimental data about the role of BCs and MC cells in the aforementioned task.

These features support the biological plausibility of the model which is next used to
investigate how dendrites contribute to pattern separation.

Dendrites and Pattern separation

Dendritic pruning

The main question of interest in this work is whether and how dendrites may contribute to
pattern separation. To answer this question we started by examining whether the number of
dendrites correlates with pattern separation performance as we prune the GC (sister)
branches from 12 (control) to 6 and 3 (Figure 4A). To assess the effect of dendritic number,
we kept all other parameters (maximum dendritic length, dendritic diameter in the IML, MML
and OML layers, membrane capacitance, "leak" conductance, number of activated synapses
and input firing rates) of the three GC models identical to those of the control. Dendritic
integration properties for the pruned GC models are shown in Figure 2-figure supplement 3
(B). The control DG network (12-dendrite model) was calibrated to have a mean population
sparsity level of 2-5% as per (Treves et al., 2008; Danielson et al., 2016), meaning that only
a small fraction of GCs were active for any given stimulus.

Interestingly, pattern separation of pairs of inputs with increasing similarity (measured either
by population distances, Figure 4C, or by rate distances, Figure 4E) was successfully
performed in the control as well as both pruned models for all pairs of inputs tested. The
efficiency of pattern separation however correlated with the number of dendrites in GCs
(Figure 4C, 4E), with the 12-dendrite GC model achieving the best performance for both
population (12-dendrite vs. 6-dendrite: Wilcoxon test, $W_1=1849.0$, $p_1 \approx 10^{-5}$, 95% CI_1 [0.02,
0.06], $W_2=1818.5$, $p_2 \approx 10^{-4}$, 95% CI_2 [0.02, 0.06], $W_3=2060.5$, $p_3 \approx 10^{-8}$, 95% CI_3 [0.04, 0.07],
$W_4=1948.0$, $p_4 \approx 10^{-6}$ 95% CI_4 [0.03, 0.06] and 12-dendrite vs. 3-dendrite: Wilcoxon test,
$W_1=2471.0$, $p_1 \approx 10^{-16}$, 95% CI_1 [0.11, 0.15], $W_2=2462.0$, $p_2 \approx 10^{-16}$, 95% CI_2 [0.11, 0.14],
$W_3=2473.0$, $p_3 \approx 10^{-16}$, 95% CI_3 [0.10, 0.13], $W_4=2451.0$, $p_4 \approx 10^{-16}$, 95% CI_4 [0.08, 0.11]) and
rate based-metrics (12-dendrite vs. 6-dendrite: Wilcoxon test, $W=1764.0$, $p=0.0004$, 95% CI

[0.03, 0.09] and 12-dendrite vs. 3-dendrite: Wilcoxon test, *W=2072.0, p≈10^{-8}, 95% CI [0.06,*

0.12]) (i.e., the highest $f_{1(output)}$ and $f_{2(output)}$, respectively).

These findings are better understood by looking at the sparsity levels exhibited by the three

GC network models. As shown in Figure 4B, the percentage of active GCs for the

population-based experiment increased substantially when the number of dendrites was

reduced. It rose from ~5%, to ~10% and ~20%, for GC model cells with 12, 6 and 3

dendrites, respectively. These differences in activity distributions were statistically significant

(12-dendrite vs. 6-dendrite: Kolmogorov-Smirnov test, *D=1.000, p≈10^{-16}* and 12-dendrite vs.

3-dendrite: Kolmogorov-Smirnov test, *D=1.000. p≈10^{-16}*). The activity levels of the three

models followed a similar pattern in the case of rate-based coding (Figure 4D); Input

patterns with high firing rates induced lower sparsity levels in the 12-dendrite GC model,

followed by the 6-dendrite model and then the 3-dendrite model (12-dendrite vs. 6-dendrite:

Kolmogorov-Smirnov test, *D=1.000, p≈10^{-16}* and 12-dendrite vs. 3-dendrite: Kolmogorov-

Smirnov test, *D=1.000. p≈10^{-16}*). These data reveal that high levels of sparsity (namely low

levels of GC activity) and pattern separation efficiency in the network model are a direct

consequence of having multiple dendrites and suggest that dendrites may contribute to

pattern separation through their effects on sparsity.

Dendritic growth

To further test whether the presence of dendrites helps pattern separation by increasing

sparsity, we also simulated the opposite process, namely the growth of dendrites. We built

GC models with 3, 6 and 12 dendrites with shapes that roughly mimic the stages of dendritic

growth (Figure 5A): starting with a GC model consisting of 3 thick dendritic compartments

and adding a branch point with two thinner sister branches at each terminal dendrite we end

up with a 12-dendrite model which is identical with our control one (see Materials and

Methods). Dendritic integration properties for the growth GC models are shown in Figure 2-

figure supplement 3 (C). Note that in this simulation, the number, length and mean diameter

of dendrites differ between the three models.

In line with the findings of the pruning experiment, the percentage of active GCs declined as

the number of dendrites increased (Figure 5B,D). The average GC activity for the

population-coding experiment started at ~70% for the network with 3 GC dendrites, dropped

to ~28% for the network with 6 GC dendrites and to ~5% for the 12-dendrite (control) model

(all differences were statistically significant, (12-dendrite vs. 6-dendrite: Kolmogorov-Smirnov

test, $D=1.000$, $p{\approx}10^{-16}$ and 12-dendrite vs. 3-dendrite: Kolmogorov-Smirnov test, $D=1.000$.

$p{\approx}10^{-16}$). Similar differences in GC sparsity were seen in the rate-based experiment (Figure

5D). In both cases, the 3-dendrite model exhibited much higher activity levels than the ones

seen in the pruning experiment (Figure 4B,D), primarily because of additional alterations in

dendritic length and diameter. Moreover, pattern separation measured by the population

metric was completely impaired in the 3-dendrite model (Figure 5C, blue line falls below the

diagonal) while performance based on rate coding remained above baseline (Figure 5E, blue

bar). Pattern separation was successfully performed in the 6-dendrite model albeit with lower

efficiency compared to the 12-dendrite model, as evaluated both with the population (12-

dendrite vs. 6-dendrite: Wilcoxon test, $W_1=2456.0$, $p_1{\approx}10^{-16}$, 95% CI_1 [0.12, 0.15],

$W_2=2421.0$, $p_2{\approx}10^{-15}$, 95% CI_2 [0.09, 0.13], $W_3=2466.0$, $p_3{\approx}10^{-16}$, 95% CI_3 [0.09, 0.13],

$W_4=2368.0$, $p_4{\approx}10^{-14}$, 95% CI_4 [0.06, 0.09] and 12-dendrite vs. 3-dendrite: Wilcoxon test, W_{1-}

$_4=2500.0$, $p_{1-4}{\approx}10^{-16}$, 95% CI_1 [0.44, 0.48], 95% CI_2 [0.40, 0.43], 95% CI_3 [0.36, 0.39], 95%

CI_4 [0.29, 0.32],) and rate distance metrics (12-dendrite vs. 6-dendrite: Wilcoxon test,

$W=1638.0$, $p=0.0076$, 95% CI [0.01, 0.07] and 12-dendrite vs. 3-dendrite: Wilcoxon test,

$W=2305.0$, $p{\approx}10^{-13}$, 95% CI [0.17, 0.25]) (Figure 5C,E).

Taken together, the "pruning" and "growth" simulations suggest a strong link between pattern

separation efficiency and GC population sparsity (Aimone et al., 2011, Deng et al., 2010)

and predict that dendrites may serve as a mechanism for increasing the sparsity of the GC

population which in turn enhances pattern separation.

Controlling sparsity with non-dendritic mechanisms

The above simulations predict that dendrites can be sufficient for mediating sparsity, which in turn enhances pattern separation. We next ask whether they are necessary for this task, as it is possible that a GC neuron can counteract the decrease in sparsity induced by having fewer/shorter dendrites through alternative mechanisms.

Input resistance does not explain differences in sparsity or pattern separation efficiency

First, we assessed the effect of input resistance differences among the various GC models on sparsity and pattern separation efficiency. A possible explanation of the above findings is that a GC model with a small dendritic tree has increased input resistance (as can be seen in Supplementary file 1A) which in turn leads to higher excitability and decreased sparsity. We thus corrected the input resistance in the 6- and 3-dendrite models to match the one in the 12-dendrite model (at the soma) by modifying a) the "leak" channel conductance (g_{leak}) or b) the size of the somatic compartments (see Supplementary file 1B for the corrected values and Figure 2-figure supplement 4 for the corrected dendritic integration profiles). Specifically, for the pruning models, g_{leak} increased by *1.230* and *1.635* for the 6- and 3-dendrite models while for the growth models, these numbers were *1.596* and *2.438*, respectively. Figure 6 shows the outcome of this correction for both pruning (A-B) and growth (C-D) cases. As evident from the figure, correcting the input resistance by increasing the "leak" conductance in the pruning case, reduced but did not eliminate differences in sparsity (Figure 6A) (12-dendrite vs. 6-dendrite: Kolmogorov-Smirnov test, *D=0.868, p≈10^{-16}* and 12-dendrite vs. 3-dendrite: Kolmogorov-Smirnov test, *D=1.000. p≈10^{-16}*) or pattern separation efficiency (Figure 6B) (12-dendrite vs. 6-dendrite: Wilcoxon test, *W_1=1636.5, p_1=0.0078, 95% CI_1 [0.01, 0.07], W_2=1606.5, p_2=0.0141, 95% CI_1 [0.01, 0.04], W_3=1732.5, p_3=0.0009, 95% CI_1 [0.02, 0.05], W_4=1668.5, p_4=0.0040 95% CI_4 [0.01, 0.04]* and 12-dendrite vs. 3-dendrite: Wilcoxon test, *W_1=2164.0, p_1≈10^{-10}, 95% CI_1 [0.06, 0.09],*

W_2=2177.5, $p_2 \approx 10^{-10}$, 95% CI_2 [0.05, 0.09], W_3=2284.0, $p_3 \approx 10^{-12}$, 95% CI_3 [0.06, 0.09],

W_4=2151.0, $p_4 \approx 10^{-10}$ 95% CI_4 [0.04, 0.07]). Similar findings were seen in the growth case,

both for the sparsity (Figure 6C) (12-dendrite vs. 6-dendrite: Kolmogorov-Smirnov test,

D=0.956, $p \approx 10^{-16}$ and 12-dendrite vs. 3-dendrite: Kolmogorov-Smirnov test, *D=1.000, $p \approx 10^{-16}$*

^{16}) and pattern separation efficiency (Figure 6D) (12-dendrite vs. 6-dendrite: Wilcoxon test,

W_1=2252.5, $p_1 \approx 10^{-12}$, 95% CI_1 [0.06, 0.10], W_2=2387.0, $p_2 \approx 10^{-15}$, 95% CI_2 [0.09, 0.12],

W_3=2408.5, $p_3 \approx 10^{-15}$, 95% CI_3 [0.07, 0.11], W_4=2258.5, $p_4 \approx 10^{-12}$, 95% CI_4 [0.05, 0.08] and

12-dendrite vs. 3-dendrite: Wilcoxon test, *W_{1-4}=2500.0, $p_{1-4} \approx 10^{-16}$, 95% CI_1 [0.24, 0.28], 95%*

CI_2 [0.22, 0.26], 95% CI_3 [0.20, 0.24], 95% CI_4 [0.16, 0.19]). Both sparsity and pattern

separation efficiency were highest in the 12-dendrite, followed by the 6-dendrite and the 3-

dendrite models. The same was seen when using the rate-based distance metric to evaluate

sparsity and pattern separation (Figure 6-figure supplement 1). In all cases, differences were

more pronounced in the growth compared to the pruning case, in line with the findings of

Figures 4-5.

Similar results were obtained when correcting the input resistance by increasing the

dimensions of the somatic compartment (Supplementary file 1B). For the pruning models,

the soma increased by a factor of *1.278* and *1.527* in the 6- and 3-dendrite models,

respectively while for the growth models, these factors were *1.513* and *1.746*, respectively.

As shown in Figure 7 (A-B), for the pruning case, differences in sparsity (12-dendrite vs. 6-

dendrite: Kolmogorov-Smirnov test, *D=0.820, $p \approx 10^{-16}$* and 12-dendrite vs. 3-dendrite,

Kolmogorov-Smirnov test, *D=1.000, $p \approx 10^{-16}$*) and pattern separation efficiency (12-dendrite

vs. 6-dendrite: Wilcoxon test, *W_1=1420.5, p_1=0.2412, 95% CI_1 [-0.01, 0.03], W_2=1588.0,*

p_2=0.0020, 95% CI_2 [0.01, 0.04], W_3=1713.5, p_3=0.0014, 95% CI_3 [0.02, 0.05], W_4=1600.5,

p_4=0.0158, 95% CI_1 [0.01, 0.04] and 12-dendrite vs. 3-dendrite: Wilcoxon test, *W_1=1914.0,*

$p_1 \approx 10^{-6}$, 95% CI_1 [0.03, 0.07], W_2=1878.0, $p_2 \approx 10^{-5}$, 95% CI_2 [0.02, 0.06], W_3=1979.5, $p_3 \approx 10^{-7}$,

95% CI_3 [0.03, 0.07], W_4=1891.5, $p_4 \approx 10^{-5}$, 95% CI_4 [0.02, 0.05]) among corrected models

decreased significantly but were not eliminated. Similar findings were seen in the growth

case, both for the sparsity (12-dendrite vs. 6-dendrite: Kolmogorov-Smirnov test, *D=1.000,*

p≈10^{-16} and 12-dendrite vs. 3-dendrite: Kolmogorov-Smirnov test, *D=1.000, p≈10^{-16}*) and

pattern separation efficiency (12-dendrite vs. 6-dendrite: Wilcoxon test, *W_1=1965.0, p_1≈10^{-13},*

95% CI_1 [0.04, 0.08], W_2=1961.5, p_2≈10^{-6}, 95% CI_2 [0.03, 0.07], W_3=2112.5, p_3≈10^{-9}, 95% CI_3

[0.04, 0.08], W_4=1919.5, p_4≈10^{-6}, 95% CI_4 [0.02, 0.06] and 12-dendrite vs. 3-dendrite:

Wilcoxon test, *W_1=2465.0, p_1≈10^{-16} 95% CI_1 [0.12, 0.15], W_2=2463.0, p_2≈10^{-16}, 95% CI_2*

[0.10, 0.14], W_3=2470.0, p_3≈10^{-16}, 95% CI_3 [0.09, 0.12], W_4=2388.0, p_4≈10^{-15}95% CI_4 [0.07,

0.10]) .Both sparsity and pattern separation efficiency remained highest in the 12-dendrite,

followed by the 6-dendrite and the 3-dendrite models. Similar findings were obtained when

using the rate-based distance metric to assess pattern separation (Figure 7-figure

supplement 1). Note that correcting the input resistance via increasing the somatic

compartment is more effective than increasing the leak conductance, as both C_m and g_{leak}

increase proportionally, keeping the membrane time constant identical across model cells.

Overall, these simulations suggest that while the input resistance is a key determinant of GC

neuron activity, it does not fully explain the differences in sparsity and pattern separation

efficiency among models with 3, 6 or 12 GC dendrites.

Sparsity is the key determinant of pattern separation efficiency

The question that arises naturally from the above findings is whether further manipulations of

the leak conductance and/or somatic dimensions could match sparsity across all models?

Moreover, would matching sparsity result in identical pattern separation efficiency, thus

making sparsity the key determinant of pattern separation (O'Reilly and McClelland, 1994;

Johnston et al., 2015)? To answer these questions we explored the effects of manipulating

intrinsic (g_{leak} and somatic dimensions) as well as extrinsic (synaptic weight) mechanisms in

the 3-, 6- and 12-dendrite GC models. Due to the consistent nature of the previous results,

the remaining simulations were performed using the pruning experiment configuration to

generate the 3- and 6-dendrite GC models and the population-based metric to assess

pattern separation. Their corresponding properties are shown in Supplementary file 1C and

the corrected dendritic integration profiles in Figure 2-figure supplement 5.

As shown in Figure 8Ai, increasing g_{leak} by a factor of *1.58* and *2.48*, in the 6- and 3-dendrite

GC models respectively, eliminated the differences in sparsity distributions compared to the

control (12-dendrite vs. 6-dendrite: Kolmogorov-Smirnov test, *D=0.096, p=0.1995* and 12-

dendrite vs. 3-dendrite: Kolmogorov-Smirnov test, *D=0.100, p=0.1641*). The same was

observed with respect to pattern separation efficiency (Figure 8Aii). All three models

exhibited identical performance across all difficulty levels (12-dendrite vs. 6-dendrite:

Wilcoxon test, *W_1=1148.5, p_1=0.4863, 95% CI_1 [-0.03, 0.01], W_2=1263.5, p_2=0.9286, 95%*

CI_2 [-0.02, 0.02], W_3=1337.5, p_3=0.5486, 95% CI_3 [-0.01, 0.02], W_4=1192.0, p_4=0.6918, 95%

CI_4 [-0.02, 0.02], 12-dendrite vs. 3-dendrite: Wilcoxon test, *W_1=1195.5, p_1=0.7097, 95% CI_1*

[-0.03, 0.01], W_2=1067.5, p_2=0.2096, 95% CI_2 [-0.04, 0.01], W_3=1326.5, p_3=0.6003, 95% CI_3

[-0.01, 0.02], W_4=1246.5, p_4=0.9835, 95% CI_4 [-0.02, 0.02]). Similarly, increasing the

diameter and length of the somatic compartment in the 6- and 3-dendrite models by *1.480*

and *1.870*, respectively (Figure 8Bi) resulted in similar sparsity (12-dendrite vs. 6-dendrite:

Kolmogorov-Smirnov test, *D=0.108, p=0.1083* and 12-dendrite vs. 3-dendrite: Kolmogorov-

Smirnov test, *D=0.040, p=0.9883*). Similarly, pattern separation efficiency was identical

across all models and difficulty levels (Figure 8Bii), (12-dendrite vs. 6-dendrite: Wilcoxon

test, *W_1=1092.0, p_1=0.2776, 95% CI_1 [-0.04, 0.01], W_2=1106.5, p_2=0.3242, 95% CI_2 [-0.03,*

0.01], W_3=1350.5, p_3=0.4906, 95% CI_3 [-0.01, 0.03], W_4=1378.0, p_4=0.3794, 95% CI_4 [-0.01,

0.03] and 12-dendrite vs. 3-dendrite: Wilcoxon test, *W_1=1144.0, p_1=0.4670, 95% CI_1 [-0.03,*

0.01], W_2=937.5, p_2=0.0315, 95% CI_2 [-0.04, 0.00], W_3=1281.0, p_3=0.8335, 95% CI_3 [-0.02,

0.02], W_4=1117.5, p_4=0.3628, 95% CI_4 [-0.03, 0.01]). It should be noted that the

abovementioned sizes are not realistic for the somata of GC neurons. Nevertheless, these

findings highlight the key role of sparsity in controlling pattern separation efficiency,

irrespectively of the number of GC dendrites. Moreover, these simulations predict that

intrinsic mechanisms of GC neurons could potentially be used to correct for morphological

alterations in order to control sparsity levels.

We next investigated whether sparsity levels can also be controlled via extrinsic (network)

rather than intrinsic mechanisms such as changes in the synaptic weights between EC and

GC neurons, namely by decreasing the strength of the input to the DG network. Specifically,

the synaptic weight of the EC → GC synapses in the 6- and 3-dendrite models was set to

0.75 and *0.56*, respectively (the control value in the 12-dendrite model was *1.00*). Again, as

shown in Figure 8Ci, differences in sparsity were eliminated among the three dendrite

models (12-dendrite vs. 6-dendrite: Kolmogorov-Smirnov test, *D=0.092, p = 0.2406* and 12-

dendrite vs. 3-dendrite: Kolmogorov-Smirnov test, *D=0.092, p=0.2406*). Finally, pattern

separation efficiency was nearly identical cross all models and difficulty levels (Figure 8Cii),

with a tiny difference in the 3-dendrite model for patterns overlapping by 90% (12-dendrite

vs. 6-dendrite: Wilcoxon test, $W_1=1085.5$, $p_1=0.2582$, 95% CI_1 [-0.04, 0.01], $W_2=1072.5$,

$p_2=0.2224$, 95% CI_2 [-0.03, 0.01], $W_3=1316.5$, $p_3=0.6491$, 95% CI_3 [-0.02, 0.02], $W_4=1122.0$,

$p_4=0.379495\%$ CI_4 [-0.03, 0.01] and 12-dendrite vs. 6-dendrite: Wilcoxon test, $W_1=1117.5$,

$p_1=0.3628$, 95% CI_1 [-0.04,0.01], $W_2=1052.5$, $p_2=0.1744$, 95% CI_2 [-0.04, 0.01], $W_3=966.0$,

$p_3=0.0507$, 95% CI_3 [-0.01, 0.01], $W_4=773.0$, $p_4=0.0010$, 95% CI_4 [-0.05, -0.01)]).

In sum, these simulations predict that sparsity is the key determinant of pattern separation

efficiency. Sparsity of the GC neuronal population can be controlled via multiple

mechanisms, including the growth of dendrites. Pruning of dendrites can be compensated by

the growth of very large somata, a large increase in the leak channel conductance or a

significant decrease in the strength of the EC input via synaptic weight changes.

What have we learnt from the model?

The above simulations suggest that dendrites contribute to pattern separation via enhancing

sparsity, yet they are not essential for this task. Different mechanisms, both intrinsic and

extrinsic, can achieve the same effects. What is not entirely clear is whether the relationship

between sparsity and pattern separation efficiency is identical for all three GC models across different sparsity levels. To test this we challenged the three models (deduced from the pruning experiment) with the task of separating pairs of inputs that overlap by 80%, while varying the synaptic weight of the EC → GC connections so as to induce different levels of sparsity. Only corrected models (namely exhibiting the same level of sparsity) were compared. As shown in Figure 9A, for any given sparsity level, pattern separation efficiency was almost identical for all three dendrite models (Wilcoxon test, *p>0.05*). These simulations suggest that indeed, when it comes to pattern separation, the key determinant is the level of GC population sparsity. Having multiple dendrites simply helps achieving high levels of sparsity because the probability of generating somatic spikes when inputs are distributed across many (rather than few) dendrites is significantly smaller.

Figure 9B summarizes the predictions made by our DG model, whereby sparsity plays a central role in pattern separation. We already knew from previous studies that inhibition is one of the key mechanisms mediating sparsity and in turn pattern separation (Myers and Scharfman, 2009, 2011; Jinde et al., 2012). What we propose here is that the presence of dendrites comes with a bonus in the DG network: it provides another mechanism for increasing sparsity and as such has a key role in pattern separation. While dendrites appear to be sufficient, they are not necessary for achieving high sparsity levels. Both intrinsic and network mechanisms can be used to achieve the same effects. In this study we identified some of these mechanisms but it is likely that there are many others and/or their combinations can also achieve similar results.

Discussion

Summary of the results

The model

The goal of this study was to reveal whether and how dendrites of principal (GC) cells can mediate pattern separation in the DG via the use of a computational approach. Towards this goal, we introduced a novel network model of the DG that includes the four major neuronal cell types found in this area, namely Granule Cells, Mossy Cells, Basket Cells, and HIPP cells. GCs were modeled as two stage integrators via the addition of dendritic branches whose properties are loosely constrained by electrophysiological and anatomical data. The rest of the neuronal types were simulated as exponential leaky integrate-and-fire neurons with adaptation. The proposed hybrid model serves as a bridge between simplified point neuron network models (Myers and Scharfman, 2009) and more detailed biophysical models (Santhakumar et al., 2005) of the DG. As such, it provides a biologically relevant and computationally efficient tool for the in depth exploration of different factors that may contribute to pattern separation, going beyond the scope of this particular study. The selective use of dendritic compartments only in GCs keeps the model complexity low while at the same time allowing the dissection of some basic GC dendritic mechanisms in pattern separation. To our knowledge this is the first DG network model of its kind.

Model predictions

Inhibition is known to control neuronal activity by increasing sparsity, as the number of active neurons is smaller for a given stimulus (Jung and McNaughton, 1993) therefore, enhancing pattern separation (Aimone et al., 2011). In the presented model, inhibition is provided to the network both directly via BCs (perisomatic inhibition) and HIPP cells (dendritic inhibition) and indirectly through the MC circuitry. We find that MC loss, while increasing GC cell activity in

line with the experimental data of (Ratzliff et al., 2002), does not lead to hyper-excitability yet

we do predict a measurable deficit on pattern separation.

To our knowledge, our model is the first to predict a role of GC dendrites in pattern

separation. This role is an indirect one and results from the inherent increase in sparsity of

the GC cells that is endowed by the presence of dendrites. Specifically, we show that the

number of GC dendrites correlates positively with pattern separation efficiency due to the

higher sparsity levels provided by having multiple dendrites. In our control model, higher

sparsity arises from the requirement of having at least two dendrites simultaneously active in

order to fire a GC model neuron. This emerged from the calibration of GC properties against

experimental data, thus is considered biologically relevant. As a result, GCs with large

numbers of dendrites have a lower probability of activation given a fixed number of afferents,

therefore increased network sparsity. We also predict that under conditions of dendritic

pruning and/or early in the growth stages of GCs, high sparsity can be achieved with

alternative mechanisms, both intrinsic (e.g., leak conductance, somatic dimensions) and

extrinsic (e.g., synaptic weights) making dendrites a sufficient but not necessary condition for

high pattern separation efficiency. These results support the hypothesis that sparsity in GC

activity improves pattern separation (O'Reilly and McClelland, 1994) and provide a list of

alternative mechanisms for controlling sparsity in the DG.

Implications for pathology

As part of the hippocampus, the DG is long hypothesized to play a key role in associative

memories, and especially when those are related with events (Morris, 2006). Furthermore,

the hippocampal DG has been implicated as the subregion most sensitive to the effects of

advancing age (Small et al., 2004). While the CA1 subregion is directly associated with

Alzheimer's Disease (AD) due to cell loss, as demonstrated in humans (West et al., 2006),

DG alterations have also been reported in patients with the aforementioned disease (Scheff

and Price, 2003), including changes in granule cell dendrites (Einstein et al., 1994).

Specifically, the dendrites of GCs in AD patients appear shorter, with fewer branches and fewer spines than those of matched controls. Moreover, in AD patients the dendrites of granular cells were reported to lose approximately 50% of their spines (Einstein et al., 1994). Our simulations show that the most important of the above-listed observations with respect to pattern separation performed by the DG would be the shortening of dendritic branches via the loss of branch points rather than the loss of side branches while mentioning the same dendritic length. Total dendritic length of GCs was previously linked to AD, which in turn is aligned with the evidence that patients with AD, who have extensive hippocampal and parahippocampal damage, lost their ability to encode information in distinct, orthogonal representations (Ally et al., 2013).

The DG is also associated with epileptogenesis in temporal lobe epilepsy (TLE) and hence, many computational models are used to investigate the effect of GC alterations in epilepsy (Tejada and Roque, 2014; Faghihi and Moustafa, 2015). Moreover, hilar cell loss has been reported in animal models after concussive head injury and also under TLE (Mathern et al., 1995). It remains unclear however, which hilar neurons are lost in animal models of TLE. As a result, there are currently three theories for TLE: a) the "dormant basket cell" hypothesis according to which the hyper-excitability in GC population is due to the loss of MCs which normally excite BCs which in turn provide inhibition to GCs. b) The "irritable mossy cell" hypothesis according to which surviving MCs hyper-excite GCs by sending uncontrolled excitation, and c) the MC loss-induced sprouting hypothesis (mossy fiber sprouting) (Ratzliff et al., 2002). We show that under conditions of MC loss, GCs exhibit increased activity (but not hyper-excitability) which should lead to pattern separation deficits. While our results are in agreement with the findings of (Ratzliff et al., 2002), more experiments need to be performed to revolve this debatable issue.

Simplifications of the model, and future directions

Several simplifications were made in modeling the individual cells and in implementing the DG network. First, we used simple point neurons in order to simulate the neuronal cells of DG. Although these models could capture the average spiking properties of a given neuron, it remains unclear how the geometrical characteristics of those neurons could affect their behavior. Another simplification concerns the effects of synaptic failure rates and receptor desensitization (Harney and Jones, 2002) in the DG, which were not included in the model.

An important aspect of DG function is the long-term synaptic plasticity, by which the connections from PP to GCs are modified. Previous DG models used a form of Hebbian learning that incorporates features of long-term potentiation and depression (Rolls, 2007). However, such a function is most likely to be relevant when stimuli are presented repetitively. On the contrary, the current model is used to distinguish patterns presented in single instances and accordingly, plasticity is not considered. Future work may address such issues along with including other interneuronal populations such as the Molecular layer Perforant Path-associated (MOPP) and the Hilar Commissural-Associational pathway related (HICAP) cells, especially when more data on their intrinsic and connectivity properties become known.

Moreover, GCs are among few cells that undergo neurogenesis in adulthood (Eriksson, 2003; Aimone et al., 2010). In a recent study by Nakashiba et al. (2012), the role of young GCs in pattern separation was investigated and it was concluded that new neurons are required for the discrimination of similar inputs. Since the presented model is used to examine specific alterations of GCs and their effect on pattern separation, neurogenesis is not incorporated but would be considered in the future. Overall, the abovementioned simplifications are unlikely to have a major effect on the basic conclusions about the contribution of morphological alterations of GC dendrites to pattern separation.

Materials and Methods

The model was developed mainly based on the structure and connectivity features described by Myers and Scharfman (2009), and incorporates the four major dentate cell types. These are the GCs, MCs, BCs and HIPP cells. As the Hilar Commissural-Associational Pathway (HICAP) cells are relatively rare and poorly understood (Sik et al., 1997), they are not included in the model. All simulations were performed using the BRIAN (BRIAN v1.4) network simulator (Goodman and Brette, 2009; Brette and Goodman, 2011) running on a High-Performance Computing Cluster (HPCC) with 312 cores under 64-bit CentOS Linux.

Model neurons

The four types of DG neurons were modeled as simplified phenomenological neurons of the integrate-and-fire (I&F) type (Izhikevich, 2003; Burkitt, 2006), with no internal geometry ("point neurons"). The GCs incorporated dendrites in order to study their role in pattern separation; however the MCs, BCs and HIPP cells were simulated as simple somatic compartments.

Modeling BC, MC and HIPP cells

Specifically, an adaptive exponential I&F model (aEIF) (Brette and Gerstner, 2005) was used to model MCs, BCs and HIPP cells. The model is mathematically described by the following differential equations (Equation 1, 2):

$$C_m \frac{dV_m}{dt} = -g_l(E_l - V_m) + g_l \Delta_T \exp\left(\frac{V_m - V_T}{\Delta_T}\right) + \sum I_{syn} - w \tag{1}$$

$$\tau_w \frac{dw}{dt} = \alpha(V_m - E_l) - w \tag{2}$$

where C_m is the membrane capacitance, V_m the membrane voltage, g_l the "leak" conductance, E_l the "leak" reversal potential (i.e., the resting potential), I_{syn} the synaptic

current flow onto the neuron, w the adaptation variable, Δ_T the slope factor, V_T the effective

threshold potential, α the adaptive coupling parameter, and τ_w is the adaptation time

constant.

The exponential nonlinearity describes the spike action potential and its upswing. In the

mathematical interpretation of the model a spike occurs at time t_{spike} when the membrane

voltage reaches a finite limit value, and thereafter the downswing of the action potential is

described by a reset fixed value V_{reset}, as follows:

$\text{at } t = t_{spike} \, (V_m \geq V_{threshold})$

$$\text{reset } V \leftarrow V_{reset}, w \leftarrow w + b \tag{3}$$

where $V_{threshold}$ is the voltage threshold above which the neuron fires a spike, and b is the

spike triggered adaptation parameter. For all neuron types the effective threshold is equal to

the voltage threshold (see Table 4 for model parameters).

Modeling principal neurons, GC

In order to investigate the role of GC dendrites in pattern separation, an extended point

neuron was implemented. The GC model consisted of a leaky Integrate-and-Fire somatic

compartment connected to a variable number of dendritic compartments whose morphology

relies on anatomical data (see Table 5 for structure characteristics). Furthermore, an

adaptation parameter w was used, only in the somatic compartment, to reproduce spike

frequency adaptation reported in these neurons. The equation that describes the membrane,

somatic and dendritic, potential of GC model cells is as follows:

$$C_m \frac{dV_m}{dt} = -g_l(E_l - V_m) + \sum I_{syn} - w \tag{4}$$

The adaptation parameter (w) was set to zero for the dendritic compartments. There is no

evidence for dendritic spikes in GCs (Krueppel et al., 2011), thus the spike mechanism was

only applied in the somatic equation.

The DG is divided into three distinct layers (Figure 1A), the molecular, granular, and

polymorphic (hilus) (Amaral et al., 2007). The GC dendrites extend in the molecular layer

(Amaral et al., 2007), which is further divided into the inner, middle, and outer molecular

layers, and therefore dendritic compartments are discretized accordingly. Table 5 lists the

morphological characteristics of the GC model. According to anatomical data, (Claiborne et

al., 1990) GCs have 10-15 dendrites; thus, the control GC model includes 12 dendrites and

its physiological responses are validated against experimental data (see Table 4 for GC

model parameters).

In order to investigate whether the number of GC dendrites affects pattern separation, we

used two different approaches; dendritic pruning and growth. First, two more GC models

were implemented which differ only in their number of dendrites (6 and 3), but the path

length remained the same across these models. Secondly, two GC models were

implemented which differ both in their dendritic number and their path length. The

morphological differences among the three models are shown in Table 5. The membrane

capacitance of the dendritic compartments was increased compared to the somatic one in

order to account for spines reported in GC dendrites (Aradi and Holmes, 1999).

The intrinsic model properties that were validated against experimental data are the input

resistance (R_{in}), the sag ratio, defined as the ratio between the exponentially extrapolated

voltage to the steady-state voltage, and the membrane time constant (τ_m). In line with

experimental procedures (Lübke et al., 1998), we used 1-second somatic current injection to

calculate the intrinsic properties. The input resistance is calculated by the equation $R_{in} =$

$\Delta V_m / I_{injected}$, where ΔV_m is the membrane response to current stimulation. Finally, the

membrane time-constant is approximated by the formula $\tau_m = R_{in} C_m$, which is a valid

approximation for passive compartments. As experimental data were obtained in the

presence of synaptic activity blockers, a somatic current injection at the model cell was used

to replicate those conditions.

Modeling Synapses

Since the DG network consists of both glutamatergic cells (GCs and MCs) and GABAergic

interneurons, AMPA, NMDA and GABA synapses were included in the network model.

Therefore, the total synaptic current (Equations 1 and 4) consisted of two components; the

excitatory current through AMPA receptors (I_{AMPA}) and NMDA receptors (I_{NMDA}), and the

inhibitory current via $GABA_{\alpha}$ receptors (I_{GABA}). The majority of ligand-gated ion channels

mediating synaptic transmission, such as AMPA and GABA receptors, display an

approximately linear current-voltage relationship when they open. We modeled these

channels as an ohmic conductance (g_{syn}) multiplied by the driving force:

$$I_{syn} = g_{syn}(t)\left[V_m(t) - E_{syn}\right] \tag{5}$$

563   where E_{syn} is the AMPA and GABA reversal potential, respectively.

564   The NMDA receptor-mediated conductance depends on the postsynaptic voltage due to the

565   gate blockage by a positively charged magnesium ion (Mg^{2+}). The fraction of NMDA

566   channels that are not blocked by Mg^{2+} can be fitted by a sigmoidal function (Jahr and

567   Stevens, 1990):

$$s(V) = \frac{1}{1 + \eta[Mg^{2+}]_o \exp(-\gamma V_m)} \tag{6}$$

where η is the sensitivity of Mg unblock, γ the steepness of Mg unblock, and $[Mg^{2+}]_o$ is the

outer magnesium (Mg) concentration. For NMDA receptors in MCs, BCs and HIPP cells we

used $\eta = 0.28\,\text{mM}^{-1}$, $[Mg^{2+}] = 1\text{mM}$, and $\gamma = 0.072\,\text{mV}^{-1}$. Instead, for GCs we tuned

these parameters in to match the latest experimental data found in literature (Krueppel et al.,

2011) with the corresponding values equal to $\eta = 0.2\,mM^{-1}$, $[Mg^{2+}] = 2mM$, and $\gamma =$

$0.04\,mV^{-1}$. Consequently, the NMDA synaptic current is calculated by the following

equation:

$$I_{syn} = g_{syn}(t)s(V)\big[V_m(t) - E_{syn}\big] \tag{7}$$

575   The ohmic conductance is simulated as a sum of two exponentials (Bartos et al., 2001), one

576   term based on rising and the other on the decay phase of the postsynaptic potential. This

577   function allows time constants to be set independently. We simulated such a function as a

578   system of linear differential equations (Roth and Van Rossum, 2009):

$$g_{syn}(t) = g_{max}u(t) \tag{8}$$

$$\frac{du}{dt} = -\frac{u}{\tau_{decay}} + h_0 v(1 - u), \frac{dv}{dt} = -\frac{v}{\tau_{rise}} \tag{9}$$

where τ_{rise} and τ_{decay} are the rise and decay constants respectively, h_0 a scaling factor and

$u(t)$ is the function of two exponentials $u(t) = \exp(-t/\tau_{decay}) - \exp(-t/\tau_{rise})$, which is

divided by its maximum amplitude. The scaling factor is set to $1\,ms^{-1}$ for all AMPA and

GABA receptors and all neuronal types. The NMDA scaling factor is set to $0.5\,ms^{-1}$ apart

from the synapses form on GCs where it is set to $2\,ms^{-1}$. Because axons of neurons are not

included in the model, a delay is used between pre- and postsynaptic transmission. The role

of the delay is to account for both the synaptic transmission and the axonal conduction

delay, and its value depends on the presynaptic and postsynaptic neuronal types. The peak

conductance (g_{max}), rise and decay time constants, and the delay of various network

connections were estimated from experimental data (Kneisler and Dingledine, 1995; Geiger

et al., 1997; Bartos et al., 2001; Schmidt-Hieber et al., 2007; Larimer and Strowbridge, 2008;

Schmidt-Hieber and Bischofberger, 2010; Krueppel et al., 2011; Chiang et al., 2012) and are

given in Table 6. Specifically, the GC peak conductance both for AMPA and NMDA, was

validated against experimental data (Krueppel et al., 2011), where it is evidenced that a

single synapse provokes a $0.6\,\mathrm{mV}$ somatic EPSP, and also the NMDA and AMPA peak

current ratio is equal to 1.08. These values were reproduced in the GC model cells. The

models also incorporate background activity, in order to simulate the experimental findings of

spontaneous activity in DG. Accordingly, we used Poisson independent spike trains in order

to reproduce the experimental data for MCs (2-4 Hz spontaneous activity) (Henze and

Buzsáki, 2007) and for BCs (1-2 Hz spontaneous activity) (Kneisler and Dingledine, 1995).

GCs infrequently generate spontaneously activity, even if inhibition is blocked (Lynch et al.,

2000). Thus, we implemented noisy inputs in order to only evoke spontaneous EPSPs (0.05

Hz spontaneous activity).

The DG network model

The DG network model consists of 2000 simulated GCs, a scale that represents 1/500 of the

one million GCs found in rat brains (West et al., 1991). The chosen number of GCs provides

enough power to explore pattern separation, while maintaining computational efficiency. The

population of GCs is organized in non-overlapping clusters, with each cluster containing 20

GCs, respectively. This kind of organization roughly corresponds to the lamellar organization

along the septotemporal extent of DG (Sloviter and Lømo, 2012).

Apart from the principal excitatory dentate cells (GCs), the model comprises two kind of

inhibitory interneurons, the perisomatic (BCs), which form synapses at the soma of the GCs,

and dendritic (HIPP) inhibitory cells, which contact the GCs at their distal dendritic

compartments. There is one BC per cluster of GCs, which in turn corresponds to 100

simulated BCs in the model. This is a form of "winner-take-all" competition (Coultrip et al.,

1992) in which all, but the most strongly activated GCs in a cluster, are silenced. Given 100

clusters in the model, and with one winner within each cluster, approximately 5% of GCs are

active for a given stimulus; this is in agreement with the theoretically and experimentally

estimation of 2-5% granular activity in the substrate (Treves et al., 2008; Danielson et al.,

2016). Moreover, the model includes simulated hilar MCs and HIPP cells. Estimated

numbers for these neuronal types vary from 30,000 to 50,000 MCs in rats (West et al., 1991;

Buckmaster and Jongen-Rêlo, 1999), which in turn corresponds to 3-5 MCs per 100 GCs.

Accordingly, the model includes 80 MCs per 2,000 GCs. Experimental counts for HIPP cells

vary significantly, but the latest estimates suggest about 12,000 HIPP cells in rats

(Buckmaster and Jongen-Rêlo, 1999) meaning less than 2 HIPP cells per 100 GCs. To

reflect this empirical data, we simulated 40 HIPP cells in the network model (Figure 1A).

External input to the network model is provided by 400 afferents representing the major input

that DG receives from Entorhinal Cortex (EC) Layer II cells, via the Perforant Path (PP). The

ratio of GCs to PP afferents is aligned with estimations of about 200,000 EC Layer II cells in

the rat (Amaral et al., 1990), suggesting a ratio of 20 EC cells per 100 GCs. Therefore, the

model incorporates synaptic input that corresponds to 400 EC Layer II cells. For simplicity,

the input cells are simulated as independent Poisson spike trains, with frequency of 40 Hz,

which is in line with experimental data (Hafting et al., 2005). Previous experimental studies

have shown that dentate GCs receive input from 10% of the 4,000 afferents that contact a

given GC in the rat during a task (McNaughton et al., 1991), which in turn suggests that an

approximate 10% of EC Layer II cells are active. The simulations reported here assume that

10% is the active PP afferents representing a given stimulus. According to McNaughton et

al., 1991, 10% of the total entorhinal input is necessary to discharge one GC. However, the

EC-GC connection is sparse, with each GC receiving input from about 2% of EC Layer II

neurons. Assuming only 400 input cells; one GC could have only 8 afferents from EC, which

in turn would make it impossible for the GC to become active. As a compromise, we used a

randomly determined 20% of EC Layer II cells as input to each GC and additionally, 20%

randomly determined EC Layer II cells as input to each HIPP cell; GCs contact each MC

with 20% probability; GCs and HIPP cells each feedback to contact a randomly determined

20% of GCs and finally, each MC connects with every BC in the network. Connections are

initialized randomly (uniform random distribution) before the start of the simulations and

remain fixed across all simulations (no rewiring). The connectivity matrix was the same for all

experiments and across all using GC models, apart from the PP→GC, and HIPP→GC

synapses due to the difference in GC number of dendrites.

Pattern Separation and Data Analysis

Generally, a network performs pattern separation whenever the similarity between two

distinct input patterns is higher than the similarity between the corresponding output patterns

(Figure 1B). In this work, the input patterns are presented as the activity along the 400 PP

afferents. Each input pattern has 40 active PP afferents (10% input density), an amount of

which are common between two patterns; hence the two patterns have a degree of

similarity. Active cell is considered every GC that fires at least one spike during stimulus

presentation (Myers and Scharfman, 2009), thus the output patterns correspond to the active

GCs. In order to quantify the pattern separation efficiency we used a metric denoted by f_1

('population distance'):

$$f_{1(i,o)} = \frac{\mathrm{HD}_{i,o}}{2(1 - s_{i,o})\mathrm{N}_{i,o}} \tag{10}$$

where the i and o subscripts denote input and output, respectively, s denotes the sparsity

(i.e., the ratio of silent neurons to all neurons), N the number of neurons, and HD is the

hamming distance between two binary patterns (Hamming, 1950), defined as the number of

positions at which the corresponding values are different. The factor of 2 in the denominator

is used to limit our distance measure at zero. Our network is said to perform pattern

separation if the input distance is smaller than the output distance, i.e., when $f_{1(i)} < f_{1(o)}$.

Thus far, the distance between two binary patterns is calculated using only the HD metric.

Although we construct the input patterns to have the same sparsity (i.e., 10% are active), the

corresponding output patterns do not necessarily have the same activity level. As we want to

examine the differences among the active neurons of each pattern we disengage the

dependence on sparsity by dividing the HD with the number of neurons that are active in a

pattern. In our case, the output patterns are vectors, each with length equal to 2000 and 2-

5% active neurons in 12 dendrites case, which in turn correspond to 40-100 neurons. The

total number of active neurons lies in the range of 40-200. For example, if the HD between

two patterns is 20, the old metric gives a distance equal to 0.01 whereas the f_1 metric

ranges from 0.10-0.25, depending on the percentage of active GC neurons. Thus, using the

proposed f_1 metric, differences only between active neurons are taken into account, making

the metric more robust across different levels of sparsity.

We constructed four groups of input pattern pairs, with different degrees of similarity and

calculated the input and the corresponding output population distances for each group

independently. Firstly, we constructed a variety of input patterns with input density 10% (i.e.,

40 active neurons) and consequently, four additional input patterns were built with 40 active

neurons, 8, 16, 24 and 32 of which are common between patterns, respectively, which, in

turn, corresponds to $f_{1(input)} = 0.4, 0.3, 0.2$ and 0.1. The reasoning behind this approach is to

examine highly overlapping patterns $(f_{1(input)} = 0.1, 0.2)$, as well as less similar ones

$(f_{1(input)} = 0.3, 0.4)$. Thus, each trial was composed of two simulations using two input

patterns within each group.

Whereas the f_1 metric quantifies the distance between two binary vectors containing active

and non-active neurons ('population distance'), we used an additional metric, denoted by f_2,

which quantifies the distance in the firing rates of common neurons that encode two patterns

by using their firing rates ('rate distance') (Figure 1C). The f_2 metric is calculated by dividing,

for each neuron that is active in both patterns, its mean firing rate given one stimulus (40 Hz)

by its mean firing rate given a stronger stimulus (50 Hz), and averaging these ratios across

the population of input and output neurons, respectively (Leutgeb et al., 2004). We subtract

this ratio from one in order to convert the 'rate similarity' into a 'rate distance':

$$f_2 = 1 - \frac{1}{N_c} \sum_{j=1}^{N_c} \frac{\left(r_{B,j} - \min r_B \right)}{\left(r_{A,j} - \min r_A \right)} \qquad (11)$$

where N_c denotes the number of common neurons that are active for bot inputs, r the firing

rate of *j-th* neuron using input A and B, respectively. Here, B represents the low firing

frequency input, while A the high frequency input. We subtract the global minimum firing rate

of GCs found in all trials in order to normalize the dynamic range of firing rates.

For this experiment, the population of active EC neurons in each pair of inputs was identical.

The network performs pattern separation if the input 'rate distance' is smaller than the

corresponding distance in the output, i.e., $f_{2(input)} < f_{2(output)}$. In each trial, the two stimuli

were presented to the network and their 'rate distance', both in input and output, was

estimated. The results were then averaged across all trials (50 trials for each model).

For each trial, the network was simulated for *850* milliseconds (ms). The first *300* ms were

simulated in order for the network to reach its stable state, so they were excluded from the

analysis. The input onset was at *300* ms and the stimulus was applied for *500* ms. The last

50 ms were simulated in order for the network to reach again its steady state and they were

excluded from the analysis as well. The time step for all simulations was set to *0.1* ms.

The data analysis and the figures describing the results were made using custom made

programs in python2.7.10™ (www.python.org) while the statistical analysis was made using

the R3.3.1 programming language (https://www.r-project.org). We used the two-sided, two-

sample Wilcoxon signed-rank test for the pattern separation efficiency comparison and the

two-sided, two-sample Kolmogorov-Smirnov test (K-S test) to compare the GC activity

probability density functions (Neuhäuser, 2011). The model will be available for download at

http://dendrites.gr/en/publications-8/software-23.

Acknowledgements

We would like to thank Alessandro Treves, Idan Segev and Nelson Spruston for their

valuable feedback on the project. In addition, we would like to thank Athanasia Papoutsi and

Panagiotis Bozelos for their helpful comments on the manuscript, as well as George

Kastellakis for his assistance in the software implementation and Pavlos Pavlidis for his

guidance on the statistical analysis and finally, all members of the Poirazi lab for their

comments during the various phases of the project.

Competing interests

723   The authors declare that no competing interests exist.

References

Aimone JB, Deng W, Gage FH. (2010). Adult neurogenesis: Integrating theories and separating functions. *Trends Cogn Sci* **14**:325–337. doi: http://dx.doi.org/10.1016/j.tics.2010.04.003

Aimone JB, Deng W, Gage FH. (2011). Resolving New Memories: A Critical Look at the Dentate Gyrus, Adult Neurogenesis, and Pattern Separation. *Neuron* **70**:589–596. doi: http://dx.doi.org/10.1016/j.neuron.2011.05.010

Ally BA, Hussey EP, Ko PC, Molitor RJ. (2013). Pattern separation and pattern completion in Alzheimer's disease: Evidence of rapid forgetting in amnestic mild cognitive impairment. *Hippocampus* **23**:1246–1258. doi: http://doi.wiley.com/10.1002/hipo.22162

Amaral DG, Ishizuka N, Claiborne B. (1990). Neurons, numbers and the hippocampal network. *Prog Brain Res* **83**:1–11. doi: http://www.ncbi.nlm.nih.gov/pubmed/2203093

Amaral DG, Scharfman HE, Lavenex P. (2007). The dentate gyrus: fundamental neuroanatomical organization (dentate gyrus for dummies). In: The Dentate Gyrus: A comprehensive guide to structure, function, and clinical implications (Scharfman HE, ed), pp 3–22. Amsterdam: Elsevier B.V.

Aradi I, Holmes WR. (1999). Role of multiple calcium and calcium-dependent conductances in regulation of hippocampal dentate granule cell excitability. *J Comput Neurosci* **6**:215–235.

Bakker A, Kirwan CB, Miller M, Stark CEL. (2008). Pattern separation in the human hippocampal CA3 and dentate gyrus. *Science* **319**:1640–1642.

Bartos M, Vida I, Frotscher M, Geiger JR, Jonas P. (2001). Rapid signaling at inhibitory synapses in a dentate gyrus interneuron network. *J Neurosci* **21**:2687–2698. doi: http://www.ncbi.nlm.nih.gov/pubmed/11306622

Bolz L, Heigele S, Bischofberger J. (2015). Running Improves Pattern Separation during

749   Novel Object Recognition. **1**:129–141.

750   Brette R, Gerstner W. (2005). Adaptive Exponential Integrate-and-Fire Model as an Effective

751   Description of Neuronal Activity. *J Neurophysiol* **94**:3637–3642. doi:

752   http://jn.physiology.org/cgi/doi/10.1152/jn.00686.2005

753   Brette R, Goodman DFM. (2011). Vectorized algorithms for spiking neural network

754   simulation. *Neural Comput* **23**:1503–1535. doi:

755   http://www.mitpressjournals.org/doi/abs/10.1162/NECO_a_00123

756   Buckmaster PS, Jongen-Rêlo AL. (1999). Highly specific neuron loss preserves lateral

757   inhibitory circuits in the dentate gyrus of kainate-induced epileptic rats. *J Neurosci*

758   **19**:9519–9529.

759   Burkitt AN. (2006). A Review of the Integrate-and-fire Neuron Model: I. Homogeneous

760   Synaptic Input. *Biol Cybern* **95**:1–19. doi: http://link.springer.com/10.1007/s00422-006-

761   0068-6

762   Chiang PH, Wu PY, Kuo TW, Liu YC, Chan CF, Chien TC, Cheng JK, Huang YY, Chiu C Di,

763   Lien CC. (2012). GABA Is Depolarizing in Hippocampal Dentate Granule Cells of the

764   Adolescent and Adult Rats. *J Neurosci* **32**:62–67.

765   Claiborne BJ, Amaral DG, Cowan WM. (1990). Quantitative, three-dimensional analysis of

766   granule cell dendrites in the rat dentate gyrus. *J Comp Neurol* **302**:206–219.

767   Coultrip R, Granger R, Lynch G. (1992). A cortical model of winner-take-all competition via

768   lateral inhibition. *Neural Networks* **5**:47–54.

769   Danielson NB, Kaifosh P, Zaremba JD, Lovett-Barron M, Tsai J, Denny CA, Balough EM,

770   Goldberg AR, Drew LJ, Hen R, Losonczy A, Kheirbek MA. (2016). Distinct Contribution of

771   Adult-Born Hippocampal Granule Cells to Context Encoding. *Neuron* **90**:101–112.

772   Das T, Ivleva EI, Wagner AD, Stark CEL, Tamminga CA. (2014). Loss of pattern separation

773   performance in schizophrenia suggests dentate gyrus dysfunction. *Schizophr Res*

774   **159**:193–197. doi: http://linkinghub.elsevier.com/retrieve/pii/S0920996414002357

Deng W, Aimone JB, Gage FH. (2010). New neurons and new memories: how does adult

hippocampal neurogenesis affect learning and memory? *Nat Rev Neurosci* **11**:339–350.

doi: http://dx.doi.org/10.1038/nrn2822

Einstein G, Buranosky R, Crain BJ. (1994). Dendritic pathology of granule cells in

Alzheimer's disease is unrelated to neuritic plaques. *J Neurosci* **14**:5077–5088.

Engin E, Zarnowska ED, Benke D, Tsvetkov E, Sigal M, Keist R, Bolshakov VY, Pearce RA,

Rudolph U. (2015). Tonic Inhibitory Control of Dentate Gyrus Granule Cells by 5-

Containing GABAA Receptors Reduces Memory Interference. *J Neurosci* **35**:13698–

13712. doi: http://www.jneurosci.org/cgi/doi/10.1523/JNEUROSCI.1370-15.2015

Erevitt BS, Skrondal A. (2010). The Cambridge dictionary of Statistics, 4th ed. Cambridge:

Cambridge University Press.

Eriksson PS. (2003). Neurogenesis and its implications for regeneration in the adult brain. *J*

Rehabil Med **35**:17–19.

Faghihi F, Moustafa AA. (2015). A computational model of pattern separation efficiency in

the dentate gyrus with implications in schizophrenia. *Front Syst Neurosci* **09**:1–12. doi:

http://journal.frontiersin.org/article/10.3389/fnsys.2015.00042

Geiger JRP, Lübke J, Roth A, Frotscher M, Jonas P. (1997). Submillisecond AMPA receptor-

mediated signaling at a principal neuron- interneuron synapse. *Neuron* **18**:1009–1023.

Goodman DFM, Brette R. (2009). The Brian simulator. *Front Neurosci* **3**:192–197. doi:

http://journal.frontiersin.org/article/10.3389/neuro.01.026.2009/abstract

Hafting T, Fyhn M, Molden S, Moser MB, Moser EI. (2005). Microstructure of a spatial map

in the entorhinal cortex. *Nature* **436**:801–806.

Hamming RW. (1950). Error Detecting and Error Correcting Codes. *Bell Syst Tech J*

29:147–160.

Harney SC, Jones M V. (2002). Pre- and postsynaptic properties of somatic and dendritic

inhibition in dentate gyrus. *Neuropharmacology* **43**:584–594.

Henze DA, Buzsáki G. (2007). Hilar mossy cells: functional identification and activity in vivo.

In: The Dentate Gyrus: A comprehensive guide to structure, function, and clinical

implications (Scharfman HE, ed), pp 199–216. Amsterdam: Elsevier B.V. doi:

http://linkinghub.elsevier.com/retrieve/pii/S007961230763012X

Ikrar T, Guo N, He K, Besnard A, Levinson S, Hill A, Lee HK, Hen R, Xu X, Sahay A. (2013).

Adult neurogenesis modifies excitability of the dentate gyrus. *Front Neural Circuits* **7**:204.

doi:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3872742&tool=pmcentrez&ren

dertype=abstract

Izhikevich EM. (2003). Simple model of spiking neurons. *IEEE Trans Neural Netw* **14**:1569–

1572. doi: http://www.ncbi.nlm.nih.gov/pubmed/18244602

Jahr CE, Stevens CF. (1990). Voltage dependence of NMDA-activated macroscopic

conductances predicted by single-channel kinetics. *J Neurosci* **10**:3178–3182.

Jain S, Yoon SY, Zhu L, Brodbeck J, Dai J, Walker D, Huang Y. (2012). Arf4 Determines

Dentate Gyrus-Mediated Pattern Separation by Regulating Dendritic Spine Development.

PLoS One **7**:1–15.

Jinde S, Zsiros V, Jiang Z, Nakao K, Pickel J, Kohno K, Belforte JE, Nakazawa K. (2012).

Hilar Mossy Cell Degeneration Causes Transient Dentate Granule Cell Hyperexcitability

and Impaired Pattern Separation. *Neuron* **76**:1189–1200. doi:

http://dx.doi.org/10.1016/j.neuron.2012.10.036

Jinde S, Zsiros V, Nakazawa K. (2013). Hilar mossy cell circuitry controlling dentate granule

cell excitability. *Front Neural Circuits* **7**:14. doi:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3569840&tool=pmcentrez&ren

dertype=abstract

Johnston ST, Shtrahman M, Parylak S, Gonçalves JT, Gage FH. (2015). Paradox of Pattern

Separation and Adult Neurogenesis: A Dual Role for New Neurons Balancing Memory

Resolution and Robustness. *Neurobiol Learn Mem* doi:

http://linkinghub.elsevier.com/retrieve/pii/S1074742715001938

Jung MW, McNaughton BL. (1993). Spatial selectivity of unit activity in the hippocampal

granular layer. *Hippocampus* **3**:165–182.

Kirwan CB, Stark CEL. (2007). Overcoming interference: an fMRI investigation of pattern

separation in the medial temporal lobe. *Learn Mem* **14**:625–633.

Kneisler TB, Dingledine R. (1995). Spontaneous and synaptic input from granule cells and

the perforant path to dentate basket cells in the rat hippocampus. *Hippocampus* **5**:151–

164.

Krueppel R, Remy S, Beck H. (2011). Dendritic integration in hippocampal dentate granule

cells. *Neuron* **71**:512–528. doi: http://dx.doi.org/10.1016/j.neuron.2011.05.043

Lacy JW, Yassa MA, Stark SM, Muftuler LT, Stark CEL. (2011). Distinct pattern separation

related transfer functions in human CA3/dentate and CA1 revealed using high-resolution

fMRI and variable mnemonic similarity. *Learn Mem* **18**:15–18.

Larimer P, Strowbridge BW. (2008). Nonrandom local circuits in the dentate gyrus. *J

Neurosci* **28**:12212–12223.

Leutgeb JK, Leutgeb S, Moser MB, Moser EI. (2007). Pattern separation in the dentate

gyrus and CA3 of the hippocampus. *Science* **315**:961–966.

Leutgeb JK, Leutgeb S, Treves A, Meyer R, Barnes CA, McNaughton BL, Moser MB, Moser

EI. (2005). Progressive transformation of hippocampal neuronal representations in

"morphed" environments. *Neuron* **48**:345–348.

Leutgeb S, Leutgeb JK, Treves A, Moser MB, Moser EI. (2004). Distinct Ensemble Codes in

Hippocampal Areas CA3 and CA1. *Science (80-)* **305**:1295–1298. doi:

http://www.sciencemag.org/cgi/doi/10.1126/science.1100265

Lübke J, Frotscher M, Spruston N. (1998). Specialized electrophysiological properties of

anatomically identified neurons in the hilar region of the rat fascia dentata. *J Neurophysiol*

79:1518–1534.

Lynch M, Sayin U, Golarai G, Sutula T. (2000). NMDA receptor-dependent plasticity of

granule cell spiking in the dentate gyrus of normal and epileptic rats. *J Neurophysiol*

84:2868–2879.

Marr D. (1971). Simple Memory : A Theory for Archicortex. *Philos Trans R Soc Lond B Biol*

Sci **262**:23–81.

Marrone DF, Adams AA, Satvat E. (2011). Increased pattern separation in the aged fascia

dentata. *Neurobiol Aging* **32**:2317.e23–e32. doi:

http://dx.doi.org/10.1016/j.neurobiolaging.2010.03.021

Mathern GW, Babb TL, Vickrey BG, Melendez M, Pretorius JK. (1995). The clinical-

pathogenic mechanisms of hippocampal neuron loss and surgical outcomes in temporal

lobe epilepsy. *Brain* **118**:105–118. doi:

http://www.brain.oxfordjournals.org/lookup/doi/10.1093/brain/118.1.105

McNaughton BL, Barnes CA, Mizumori SJY, Green EJ, Sharp PE. (1991). Contribution of

granule cells to spatial representations in hippocampal circuits: a puzzle. In: Kindling and

Synaptic Plasticity: The Legacy of Graham Goddar (Morrell F, ed), pp 110–123. Boston:

Springer-Verlag.

Morris RGM. (2006). Elements of a neurobiological theory of hippocampal function: the role

of synaptic plasticity, synaptic tagging and schemas. *Eur J Neurosci* **23**:2829–2846. doi:

http://doi.wiley.com/10.1111/j.1460-9568.2006.04888.x

Motley SE, Kirwan CB. (2012). A Parametric Investigation of Pattern Separation Processes

in the Medial Temporal Lobe. *J Neurosci* **32**:13076–13084.

Myers CE, Scharfman HE. (2009). A Role for hilar cells in pattern separation in the dentate

gyrus: A computational approach. *Hippocampus* **19**:321–337.

Myers CE, Scharfman HE. (2011). Pattern separation in the dentate gyrus: A role for the

CA3 backprojection. *Hippocampus* **21**:1190–1215.

Nakashiba T, Cushman JD, Pelkey KA, Renaudineau S, Buhl DL, McHugh TJ, Barrera VR,

Chittajallu R, Iwamoto KS, McBain CJ, Fanselow MS, Tonegawa S. (2012). Young

dentate granule cells mediate pattern separation, whereas old granule cells facilitate

pattern completion. *Cell* **149**:188–201. doi: http://dx.doi.org/10.1016/j.cell.2012.01.046

Neuhäuser M. (2011). Nonparametric Statistical Tests: A Computational Approach. New

York: CRC Press. doi: papers3://publication/uuid/1A2A31AE-B983-4060-BD27-

C5B73222FF1C

Nitz D, McNaughton B. (2004). Differential modulation of CA1 and dentate gyrus

interneurons during exploration of novel environments. *J Neurophysiol* **91**:863–872.

O'Reilly RC, McClelland JL. (1994). Hippocampal Conjunctive Encoding, Storage, and

Recall: Avoiding a Trade-Off. *Hippocampus* **4**:661–682. doi:

http://www.ncbi.nlm.nih.gov/pubmed/7704110

Olshausen BA, Field DJ. (2004). Sparse coding of sensory inputs. *Curr Opin Neurobiol*

14:481–487.

Petrantonakis PC, Poirazi P. (2014). A compressed sensing perspective of hippocampal

function. *Front Syst Neurosci* **8**:1–13. doi:

http://journal.frontiersin.org/Journal/10.3389/fnsys.2014.00141/abstract

Petrantonakis PC, Poirazi P. (2015). Dentate Gyrus Circuitry Features Improve Performance

of Sparse Approximation Algorithms Dehghani F, ed. *PLoS One* **10**:e0117023. doi:

http://dx.plos.org/10.1371/journal.pone.0117023

Ratzliff ADH, Howard AL, Santhakumar V, Osapay I, Soltesz I. (2004). Rapid deletion of

mossy cells does not result in a hyperexcitable dentate gyrus: implications for

epileptogenesis. *J Neurosci* **24**:2259–2269.

Ratzliff AH, Santhakumar V, Howard A, Soltesz I. (2002). Mossy cells in epilepsy: Rigor

mortis or vigor mortis? *Trends Neurosci* **25**:140–144.

Rolls ET. (2007). An attractor network in the hippocampus: theory and neurophysiology.

Learn Mem **14**:714–731.

Roth A, Van Rossum MCW. (2009). Modeling synapses. In: Computational modeling

methods for neuroscientists (De Shutter E, ed), pp 139–160. Cambridge, MA: MIT press.

Santhakumar V, Aradi I, Soltesz I. (2005). Role of mossy fiber sprouting and mossy cell loss

in hyperexcitability: a network model of the dentate gyrus incorporating cell types and

axonal topography. *J Neurophysiol* **93**:437–453.

Santoro A. (2013). Reassessing pattern separation in the dentate gyrus. *Front Behav*

Neurosci **7**:1–4. doi:

http://journal.frontiersin.org/article/10.3389/fnbeh.2013.00096/abstract

Satvat E, Schmidt B, Argraves M, Marrone DF, Markus EJ. (2011). Changes in task

demands alter the pattern of zif268 expression in the dentate gyrus. *J Neurosci* **31**:7163–

7167.

Scheff SW, Price DA. (2003). Synaptic pathology in Alzheimer's disease: A review of

ultrastructural studies. *Neurobiol Aging* **24**:1029–1046.

Schmidt-Hieber C, Bischofberger J. (2010). Fast sodium channel gating supports localized

and efficient axonal action potential initiation. *J Neurosci* **30**:10233–10242.

Schmidt-Hieber C, Jonas P, Bischofberger J. (2007). Subthreshold dendritic signal

processing and coincidence detection in dentate gyrus granule cells. *J Neurosci* **27**:8430–

8441.

Sik A, Penttonen M, Buzsaki G. (1997). Interneurons in the Hippocampal Dentate Gyrus : an

In Vivo Intracellular Study. *Eur J Neurosci* **9**:573–588.

Sloviter RS, Lømo T. (2012). Updating the lamellar hypothesis of hippocampal organization.

Front Neural Circuits **6**:102. doi:

http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3517983&tool=pmcentrez&ren

dertype=abstract

Small SA, Chawla MK, Buonocore M, Rapp PR, Barnes CA. (2004). Imaging correlates of

brain function in monkeys and rats isolates a hippocampal subregion differentially

vulnerable to aging. *Proc Natl Acad Sci U S A* **101**:7181–7186.

Squire LR, Stark CEL, Clark RE. (2004). The medial temporal lobe. *Annu Rev Neurosci*

27:279–306.

Tejada J, Roque AC. (2014). Computational models of dentate gyrus with epilepsy-induced

morphological alterations in granule cells. *Epilepsy Behav* **38**:63–70. doi:

http://dx.doi.org/10.1016/j.yebeh.2014.02.007

Treves A, Rolls ET. (1994). Computational Analysis of the Role of the Hippocampus in

Memory. *Hippocampus* **4**:374–391.

Treves A, Tashiro A, Witter MP, Moser EI. (2008). What is the mammalian dentate gyrus

good for? *Neuroscience* **154**:1155–1172. doi:

http://linkinghub.elsevier.com/retrieve/pii/S0306452208006660

West MJ, Kawas CH, Martin LJ, Troncoso JC. (2006). The CA1 Region of the Human

Hippocampus Is a Hot Spot in Alzheimer's Disease. *Ann N Y Acad Sci* **908**:255–259. doi:

http://doi.wiley.com/10.1111/j.1749-6632.2000.tb06652.x

West MJ, Slomianka L, Gundersen HJ. (1991). Unbiased stereological estimation of the total

number of neurons in thesubdivisions of the rat hippocampus using the optical

fractionator. *Anat Rec* **231**:482–497.

Yassa MA, Stark CEL. (2011). Pattern separation in the hippocampus. *Trends Neurosci*

34:515–525. doi: http://dx.doi.org/10.1016/j.tins.2011.06.006

Yim MY, Hanuschkin A, Wolfart J. (2015). Intrinsic rescaling of granule cells restores pattern

separation ability of a dentate gyrus network model during epileptic hyperexcitability.

Hippocampus **25**:297–308. doi: http://doi.wiley.com/10.1002/hipo.22373

Figures

Figure 1. A. Schematic representation of the implemented DG network model. Different shades of green

illustrate the layer division. PP: Perforant Path, GC: Granule Cells, BC: Basket Cells, MC: Mossy Cells, HIPP:

Hilar Perforant Path-associated cells. Perforant Path afferents curry the input to the network, and project on both

the GCs and the HIPP cells. MCs and GCs are connected in a recurrent manner. MCs also excite the BCs.

Inhibition of GCs is provided through BCs and HIPP cells directly and indirectly via MCs. Note that HIPP cells

contact the distal dendrites rather than the soma of GCs. GCs provide the output of the DG network. **B.**

Schematic representation of pattern separation using population-based coding. When two highly overlapping EC

inputs (input 1 & 2, with identical mean firing rates) arrive in DG, the corresponding outputs are highly dissimilar.

Note that the output pattern is sparse because of the low number of GCs that encode any given pattern. **C.**

Schematic representation of pattern separation using rate-based coding. When two highly overlapping EC inputs

(input 1 & 2, with different mean firing rates but identical input populations) arrive in DG, the corresponding

outputs are highly dissimilar in their firing rates but also likely to differ in the populations they activate. Rate

distances are estimated over the set of common neurons (red box).

Figure 2. Firing traces of the four model cells in response to current injection (1 second). Note that APs are not explicitly modeled in I&F neurons. **A:** The somatic membrane voltage of the granule cell (GC) model in response to +90 pA (top), and -50 pA (bottom) somatic current injections. **B.** Same as in **A,** for the mossy cell (MC) with +300 pA (top) and -300pA (bottom) current injections. **C.** Same as in **A,** for the Basket cell (BC), with +200 pA (top) and -50 pA (bottom) current injections. **D.** Same as in **A,** for the HIPP cell, with +300 pA (top) and -50 pA (bottom) current injections. Current – Voltage (I-V) curves are given in Figure 2-figure supplement 1, while Current – frequency plots are shown in Figure 2-figure supplement 2. Moreover, Figure 2-figure supplement 3 shows the somatic EPSP against Arithmetic sum for GCs under control, pruning and growth conditions. In Figure 2-figure supplement 4 and 5, the corresponding EPSP vs. Arithmetic sum curves for GCs after matching R_{in} and sparsity are given, respectively.

Figure 3. Complete mossy cell loss reduces pattern separation efficiency in the DG network. **A.** Schematic

representation of the 12-dendrite DG network with MC loss. **B.** The corresponding probability density functions of

GC activity in response to 40 Hz input for the control (dark green) and MC loss conditions (light green). The mean

activity is 5% and 9% for the control and MC loss networks, respectively. The respective mean firing rates of GCs

are shown in Table 3. The histograms were calculated with 20 bins each. **C.** Input / Output population distances

(f_1) for the control (dark green) and MC loss (light green) networks estimated using input patterns with increasing

similarity. The dashed line denotes the limit above which the model performs pattern separation. MC loss

reduces pattern separation efficiency for all input patterns tested. Error bars represent the standard error of the

mean (Erevitt and Skrondal, 2010). **D.** Probability density functions of GC activity using control and MC loss

models presented with two input patterns that differ only in their firing rates. Shades of green represent the high

frequency input (50 Hz), while shades of brown represent the low frequency input (40Hz). Dark and light shades

represent the control and MC loss condition, respectively. The respective mean firing rates of GCs are shown in

Table 3. **E.** Input / Output rate distances (f_2) for the control (dark green) and MC loss (light green) networks

estimated using two input patterns with different firing frequencies, 40 and 50 Hz respectively. MC loss slightly

reduces the efficiency of pattern separation.

Figure 4. Effect of GC dendrite pruning on pattern separation. **A**. A schematic drawing of the three GC models

with 12, 6 and 3 dendrites. **B.** Corresponding probability density functions of GC activity for the three GC models

in response to a single input pattern at 40 Hz. The mean activity is inversely analogous to the number of

dendrites: as the number of dendrites decreases, the GC population becomes more active, i.e., less sparse. The

respective mean firing rates of GCs are shown in Table 3. **C.** Input / Output population distances (f_1), for the 12-

dendrite (green), 6-dendrite (red) and 3-dendrite (blue) GC models in response to the presentation of two

overlapping input patterns at 40 Hz. Input pairs were generated so as to have decreasing amounts of overlap, as

depicted in Figure 1B. The dashed line denotes the limit above which the model performs pattern separation.

Performance declines as the dendritic tree has fewer dendrites. Error bars represent the standard error of the

mean across trials. **D.** Probability density functions of GC activity for the three GC models in response to

presentation of two input patterns with different firing rates (low rate = 40 Hz, high rate = 50 Hz), as depicted in

Figure 1C. GC activity increases with the number of dendrites and the input firing rate. The respective mean firing

rates of GCs are shown in Table 3. **E.** Input / Output rate distances (i.e., f_2) for the three models. All three models

perform pattern separation, as the rate distances in the input are significantly higher than the corresponding ones

in the output.

1012

Figure 5. Effect of GC dendrite growth on pattern separation. **A**. A schematic drawing of the three GC models

with 3, 6 and 12 dendrites. **B.** Corresponding probability density functions of GC activity for the three GC models

in response to a single input pattern at 40 Hz. The mean activity is inversely analogous to the number/length of

dendrites: as dendrites grow, the GC population becomes sparser, i.e., fewer GCs are active. The respective

mean firing rates of GCs are shown in Table 3. **C.** Input / Output population distances (f_1), for the 3-dendrite

(blue), 6-dendrite (red) and 12-dendrite (green) GC models in response to the presentation of two overlapping

input patterns at 40 Hz, with different degrees of overlap, as depicted in Figure 1B. The dashed line denotes the

limit above which the model performs pattern separation. Performance improves with the growth of dendrites. **D.**

Probability density functions of GC activity for the three GC models in response to presentation of two input

patters with different firing rates (low rate = 40 Hz, high rate = 50 Hz), as depicted in Figure 1C. GC activity

decreases with the number of dendrites and increases with the input firing rate. The respective mean firing rates

of GCs are shown in Table 3. **E.** Input / Output rate distances (i.e., f_2) for the three models. The number of GC

dendrites is again analogous to the pattern separation performance.

Dendritic pruning

Dendritic growth

1026

Figure 6. Effect of GC dendritic pruning (top panel) and growth (bottom panel) on pattern separation when the input resistance (R_{in}) is the same across models. To match R_{in}, the leak conductance (g_{leak}) was increased by a factor of 1.695 and 1.230, in the 3- and 6-dendrite models respectively. **A.** Corresponding probability density functions of GC activity for the three GC models in response to a single input pattern at 40 Hz. GC activity distributions become more similar but remain inversely analogous to the number of dendrites: as the number of dendrites increase, the GC population becomes sparser. The respective mean firing rates of GCs are shown in Table 3. **B.** Input / Output population distances (f_1), for the 3-dendrite (blue), 6-dendrite (red) and 12-dendrite (green) GC models in response to the presentation of two overlapping input patterns at 40 Hz as depicted in Figure 1B. The dashed line denotes the limit above which the model performs pattern separation. Performance improves with the number of dendrites. **C.** The same as in **A.** for the growth experiment. While distributions move closer to one another, the inverse relationship between dendritic number and mean sparsity is preserved. **D.** Same as in **B.** for the growth experiment. Pattern separation efficiency still correlates with the number of dendrites. Figure 6-figure supplement 1 depicts the corresponding pattern separation efficiency based on 'rate distance'.

Dendritic pruning

A.

B.

Dendritic growth

C.

D.

1041

Figure 7. Effect of GC dendritic pruning (top panel) and growth (bottom panel) on pattern separation when matching the input resistance (R_{in}) via increasing the somatic size. The neuronal soma of the 3- and 6-dendrite models was increased by a factor of 1.527 and 1.278, respectively. **A.** Corresponding probability density functions in response to a single input pattern at 40 Hz. While distributions move closer, the mean activity remains inversely analogous to the number of dendrites. The respective mean firing rates of GCs are shown in Table 3. **B.** Input / Output population distances (f_1), for the 3-dendrite (blue), 6-dendrite (red) and 12-dendrite (green) GC models in response to the presentation of two overlapping input patterns at 40 Hz as depicted in Figure 1B. The dashed line denotes the limit above which the model performs pattern separation. Performance becomes similar yet statistically different among the three corrected models and remains analogous to the number of dendrites. **C.** Same as in **A.** for the growth experiment. The inverse relationship between dendritic number and mean sparsity is preserved. **D.** Same as in **B.** for the growth experiment. Pattern separation efficiency remains different and analogous to the number of dendrites. Figure 7-figure supplement 1 depicts the corresponding pattern separation efficiency based on 'rate distance'.

A. Leak conductance (g_{leak})

i.

ii.

B. Somatic size

i.

ii.

C. Synaptic weight

i.

ii.

1055

Figure 8. Effect of matching sparsity on pattern separation efficiency. **A.** The "leak" conductance, g_{leak}, of the 6- and 3-dendrite GC models increases by a factor of 1.58 and 2.48, respectively. **A.i** Resulting GC activity distributions are not statistically different. **A.ii** Networks with corrected GC models have identical performance. Differences in pattern separation using the population metric are not statistically significant. **B.** GC activity distributions are matched by increasing the soma diameter and length of the 6- and 3-dendrite GC models by 1.48 and 1.87, respectively. **B.ii** The network with corrected 3-dendrite GC models has a slightly smaller pattern separation efficiency compared to the other two probably because its activity distribution is wider. Generally the three models have very similar pattern separation performance. **C.** GC activity distributions are matched by decreasing the EC → GC synaptic weight from 1.00, to 0.75 and 0.56 in the 6-, and 3- dendrite models, respectively. Pattern separation efficiency is statistically the same across all corrected GC models.

1066

Figure 9. A. The ratio of $f_{1(output)}$ to $f_{1(input)}$ for all three models as function of GC activity. Each dot represents

an average of 20 trials for a specific synaptic weight value. We used values in the range of 0.0 to 25.0. For any

given sparsity level, pattern separation efficiency is identical across all corrected GC models. **B.** Schematic

representation of our proposed working hypothesis based on model predictions: sparsity is the key determinant of

pattern separation. Sparsity is in turn controlled by a number of mechanisms, including inhibition, the presence of

dendrites as well as various intrinsic and extrinsic mechanisms.

Tables

Table 1. Physiological properties of real and model neurons

Cell type	Input resistance R_{in} (MΩ)		Membrane time constant, τ_m (ms)		Sag ratio		Maximum firing rate (Hz)	
	Model	Biological	Model	Biological	Model	Biological	Model	Biological
Granule cells	360	292 ± 34	41.2	31 ± 2	0.91	0.96 ± .1	60	70 ± 10
Mossy cells	105	199 ± 19	33.7	35 ± 5	0.98	0.81 ± .3	45	50 ± 6
Basket cells	55	56 ± 9	9.67	10 ± 1	0.99	0.97 ± .02	247	230 ± 15
HIPP cells	363	371 ± 47	21.4	15 ± 0	0.84	0.82 ± .04	113	101 ± 24
Biological value	Lubke et al., 1998, Krueppel et al., 2009		Lubke et al., 1997, Ratzliff et al., 2004		Lubke et al., 1998, Bartos et al., 2001		Lubke et al., 1997	

1075

1076   **Table 2. DG network (12-dendrite model) performance on a simple pattern separation task.**

Input Similarity ($f_{1(input)}$)	Output Similarity ($f_{1(output)}$) mean ± sem
0.10	0.43 ± 0.007
0.20	0.52 ± 0.007
0.30	0.59 ± 0.008
0.40	0.64 ± 0.008
Input Similarity ($f_{2(input)}$) mean ± sem	Output Similarity ($f_{2(output)}$) mean ± sem
0.16 ± 0.05	0.66 ± 0.01

1077

1078 **Table 3. Mean firing frequencies of GCs in response to the various input patterns used in**

1079 **Figures 3-8**

1080 *Population-distance Experiments (Pruning, Growth, MC loss)*

	Mean firing rate ± sem (Hz)		
GC models	Pruning	Growth	MC loss
12-dendrite	3.50±0.002	3.50±0.002	4.82±0.002
6-dendrite	3.65±0.009	3.85±0.007	-
3-dendrite	3.90±0.006	6.35±0.005	-

1081

1082 *Population-distance Experiments (Match R_{in})*

	Mean firing rate ± sem (Hz)			
	Pruning		Growth	
GC models	g_{leak}	Soma dimensions	g_{leak}	Soma dimensions
6-dendrite	3.61±0.009	3.58±0.009	3.66±0.008	3.60±0.008
3-dendrite	3.75±0.007	3.58±0.007	4.59±0.006	3.78±0.006

1083

1084 *Population-distance Experiments (Match sparsity)*

	Mean firing rate ± sem (Hz)		
GC models	g_{leak}	Soma dimensions	Synaptic weight
6-dendrite	3.47±0.010	3.44±0.010	3.31±0.010
3-dendrite	3.47±0.010	3.31±0.010	3.12±0.008

1085

1086 *Rate-distance Experiments (Control, Growth, MC loss)*

	Mean firing rate ± sem (Hz)					
	Control		Growth		MC loss	
GC models	Low input rate (40Hz)	High input rate (50Hz)	Low input rate (40Hz)	High input rate (50Hz)	Low input rate (40Hz)	High input rate (50Hz)
12-dendrite	3.75±0.036	5.24±0.048	3.75±0.036	5.24±0.048	4.94±0.057	8.07±0.110
6-dendrite	3.98±0.028	5.20±0.035	3.83±0.023	5.03±0.030	-	-
3-dendrite	4.13±0.019	5.48±0.023	6.43±0.015	9.07±0.017	-	-

1087

 Table 4. Model parameters for all neuronal types

Model Parameter	Granule cells		Mossy cells	Basket cells	HIPP cells
	Soma	dendrites			
E_l (mV) Resting potential	-87	-82	-64	-52	-59
g_l (nS) "Leak" conductance	0.00003[a]	0.00001[a]	4.53	18.054	1.930
C_m (nF) Membrane capacitance	1.0[b]	2.5[b]	0.621	0.1793	0.0584
V_{reset} (mV) Reset voltage	-74		-49	-45	-56
$V_T = V_{thr}$ (mV) Threshold voltage	-56		-42	-39	-50
Δ_T (mV) Slope factor	-		2	2	2
α (nS) Adaptation coupling parameter	2.0		2	0.1	0.82
τ_w (ms) Adaptation time constant	45		180	100	93
b (nS) Spike triggered adaptation	0.0450		0.0829	0.0205	0.015

1089   [a]For the GC model the "leak" conductance is given in Siemens/cm^2

1090   [b]For the GC model the membrane capacitance is given in uF/cm^2

1091 **Table 5. Morphological structure of GC models**

Structure of GC models	Control	Pruning		Growth	
	12 dendrites	6 dendrites	3 dendrites	6 dendrites	3 dendrites
# of compartments					
total	21	15	9	9	3
proximal	3	3	3	3	3
medial	6	6	3	6	-
distal	12	6	3	-	-
length per compartment (um)	83	83	83	83	83
total dendritic length (um)	1743	1245	747	747	249
diameter per compartment (um)					
proximal	1.0	1.0	1.0	1.0	1.0
medial	0.9	0.9	0.9	0.9	-
distal	0.8	0.8	0.8	-	-

1092

 Table 6. Synaptic parameters of the dentate network

From (column) / To (row)	Granule cells	Mossy cells	Basket cells	HIPP cells
AMPA				
Perforant Path				
g_{max} (nS)	0.8066			0.240
τ_{rise} (ms)	0.1			2.0
τ_{decay} (ms)	2.5			11.0
delay (ms)	3.0			3.0
Granule cells				
g_{max} (nS)		0.500	0.210	
τ_{rise} (ms)		0.5	2.5	
τ_{decay} (ms)		6.2	3.5	
delay (ms)		1.5	0.8	
Mossy cells				
g_{max} (nS)	0.1066		0.350	
τ_{rise} (ms)	0.1		2.5	
τ_{decay} (ms)	2.5		3.5	
delay (ms)	3.0		3.0	
NMDA				
Perforant Path				
g_{max} (nS)	0.8711			0.276
τ_{rise} (ms)	0.33			4.8
τ_{decay} (ms)	50.0			110.0
delay (ms)	3.0			3.0
Granule Cells				
g_{max} (nS)		0.525	0.231	
τ_{rise} (ms)		4.0	10.0	
τ_{decay} (ms)		100.0	130.0	
delay (ms)		1.5	0.8	
Mossy Cells				
g_{max} (nS)	0.1151		0.385	
τ_{rise} (ms)	0.33		10.0	
τ_{decay} (ms)	50.0		130.0	
delay (ms)	3.0		3.0	
GABA$_A$				
Basket cells				
g_{max} (nS)	14.0			
τ_{rise} (ms)	0.9			
τ_{decay} (ms)	6.8			
delay (ms)	0.85			
HIPP cells				
g_{max} (nS)	0.12			
τ_{rise} (ms)	0.9			
τ_{decay} (ms)	6.8			
delay (ms)	1.6			

1094