1 **Genomic insights into adaptive divergence and speciation between**
2 **malaria vectors of the *Anopheles nili* group**
3

Colince Kamdem[1]*, Caroline Fouet[1], Stephanie Gamez[1], Bradley J. White[1,2]*

[1]Department of Entomology, University of California, Riverside, CA 92521

[2]Center for Disease Vector Research, Institute for Integrative Genome Biology, University of

California, Riverside, CA 92521

*Corresponding authors: colincek@ucr.edu; bwhite@ucr.edu

Abstract

Ongoing speciation in most African malaria vectors gives rise to cryptic species that differ

remarkably in their behaviour, ecology and capacity to vector malaria parasites. Any vector

control measure can be undermined if cryptic populations exist and are mistargeted within

vector species. To examine population structure and the potential impacts of recent large-

scale control programs, we have investigated genomic patterns of differentiation in

mosquitoes belonging to a large taxonomic group that diverged ~3-Myr ago. Using 4343

single nucleotide polymorphisms (SNPs), we detected strong population structure

characterized by high F_{ST} values between multiple divergent populations. Delineating

cryptic species within the *Anopheles nili* group is challenging due to incongruence between

morphology, ribosomal DNA and SNP markers consistent with incomplete lineage sorting

and/or gene flow. Throughout the genome, a very high proportion of loci are fixed (F_{ST} = 1)

within putative species, which suggests that ecological and/or reproductive barriers are

maintained by strong selection on a substantial number of genes.

Key words: *Anopheles nili*, divergent selection, high- FST regions, speciation.

1. Introduction

One of the principal goals of population genetics is to summarize genetic similarities and differences between populations (Wright 1984). This task can be relatively straightforward for some taxa, but the genetic relationship among populations can also be difficult to summarize, especially for species whose evolutionary history is complex and reticulate. Most *Anopheles* mosquitoes have very complex range-wide population structure due to cryptic speciation and ongoing gene flow, which complicate species and population delimitations (Krzywinski & Besansky 2003; Harbach 2013). For example, almost all major African malaria vectors belong to large taxonomic groups encompassing multiple incipient species relatively isolated reproductively and geographically from one another (Reviewed by Sinka *et al.* 2010; Antonio-nkondjio & Simard 2013; Dia *et al.* 2013; Lanzaro & Lee 2013). These characteristics make them promising model systems to study speciation and the processes which contribute to reproductive barriers (e.g. Turner *et al.* 2005; Lawniczak *et al.* 2010; Neafsey *et al.* 2010; Fontaine *et al.* 2015), but can also have far-reaching practical consequences.

When the cryptic genetic structure underlying variations in behavior, distribution and vectorial capacity within mosquito species is not taken into account, vector populations can be mistargeted and major efforts at malaria control undermined (e.g. Molineaux & Gramiccia 1980; Van Bortel *et al.* 2001). The genetic structure of *Anopheles* species is most often based on macrogeographic or regional subdivisions (e.g. Touré *et al.* 1998; Wondji *et al.* 2005; Antonio-Nkondjio *et al.* 2007; Slotman *et al.* 2007; Ayala *et al.* 2011; Ndo *et al.* 2013; Pinto *et al.* 2013; Caputo *et al.* 2014; Kamdem *et al.* 2017), but can also involve more subtle differentiation between larvae and adults, or between adult populations found in or

around human dwellings (e.g. Riehle *et al.* 2011). This type of microgeographic genetic

structure can have potentially significant impacts on malaria epidemiology and insecticide

resistance because the main vector control measures target exclusively indoor mosquito

populations (WHO 2016).

The recent scaling up of insecticide-treated nets usage and indoor insecticide

spraying to a lesser extent have led to a dramatic reduction of malaria morbidity across the

continent, but have also inevitably increased insecticide resistance among vectors

(Hemingway *et al.* 2016; Ranson & Lissenden 2016). Although a direct causal link is not

established, populations from urban areas and regions where pesticides are intensely used

are strongly differentiated genetically implying a certain correlation between insecticide

exposure and local adaptation (Kamdem *et al.* 2017). It has also been shown that this

increased usage of xenobiotics can drive selective sweeps across genomes of the two major

vectors *An. gambiae* and *An. funestus* (Clarkson *et al.* 2014; Barnes *et al.* 2017; Kamdem *et*

al. 2017). The success of current malaria control measures relies among others on accurate

knowledge of population structure and the environmental drivers that contribute or have

contributed to genetic differentiation between mosquito populations. Extensive sampling

combined with recent genomic tools and analytical approaches have helped resolve fine-

scale population structure and genomic targets of selection in *An. gambiae* and *An. funestus*

(White *et al.* 2011; Reidenbach *et al.* 2012; O'Loughlin *et al.* 2014; Barnes *et al.* 2017; Fouet

et al. 2017; Kamdem *et al.* 2017), but important taxa remain understudied.

The present work focused on a set of African malaria vectors representing a large

taxonomic unit named "*An. nili* group". Species that compose this group are characterized

by reticulate evolution and complex phylogenies that have been challenging to resolve

(Kengne *et al.* 2003; Awono-Ambene *et al.* 2004, 2006, Ndo *et al.* 2010, 2013; Peery *et al.*

2011; Sharakhova *et al.* 2013). To date, four species distinguishable by slight morphological

variations are known within the group and occur in forested areas of Central and West

Africa: *An. nili sensu stricto* (hereafter *An. nili*), *An. ovengensis*, *An. carnevalei*, and *An.*

somalicus. Populations of *An. nili* and *An. ovengensis* are very anthropophilic and efficient

vectors of *Plasmodium* in rural areas where malaria prevalence is particularly high

(Antonio-Nkondjio *et al.* 2006).

To delineate genomic patterns of differentiation, we sampled mosquito populations

throughout the range of species of the *An. nili* group in Cameroon and used reduced

representation sequencing to develop genome-wide SNP markers that we genotyped in 145

individuals. We discovered new strongly differentiated subpopulations within *An.*

ovengensis and *An. nili*. We analyzed the genetic differentiation at 4343 SNPs and revealed

the presence of a very high number of outlier loci that are targets of selection. Further

studies using a more complete reference genome will help answer important questions

concerning the functional and phenotypic characteristics of differentiated loci as well as the

contribution of recent selective events.

2. Materials and methods

(a) Mosquito species

We surveyed 28 locations within the geographic ranges of species of the *An. nili*

group previously described in Cameroon (Figure 1) (Awono-Ambene *et al.* 2004, 2006;

Antonio-Nkondjio *et al.* 2009; Ndo *et al.* 2010, 2013). We collected larvae and adult

mosquitoes in and around human dwellings using several sampling techniques (Service

1993) (Table S1). The four currently known members of the *An. nili.* group were identified

using morphological keys and a diagnostic PCR, which discriminates species on the basis of

point mutations of the ribosomal DNA (Gillies & De Meillon 1968; Gillies & Coetzee 1987;

Kengne *et al.* 2003; Awono-Ambene *et al.* 2004).

(b) Library preparation, sequencing and SNP discovery

We created double-digest RAD libraries using a modified version of the protocol

described by Peterson *et al.* 2012. Genomic DNA of mosquitoes was extracted using the

DNeasy Blood and Tissue kit (Qiagen) and the Zymo Research MinPrep kit for larvae and

adult samples respectively. Approximately 50 ng (10μl) of DNA of each mosquito was

digested simultaneously with *MluC1* and *NlaIII* restriction enzymes. Digested products were

ligated to adapter and barcode sequences enabling identification of individuals. Samples

were pooled, purified, and 400-bp fragments selected. The resulting libraries were

amplified via PCR and purified, and fragment size distribution was checked on BioAnalyzer

(Agilent Technologies, Inc., USA). PCR products were quantified, diluted and sequenced on

Illumina HiSeq2000 (Illumina Inc., USA).

(b) SNP discovery and genotyping

The *process_radtags* program of the Stacks v 1.35 pipeline (Catchen *et al.* 2011, 2013) was used to demultiplex and clean raw reads. Reads that passed quality filters were aligned to the *An. nili* Dinderesso draft genome assembly made up of 51048 short contigs (~200-30512bp long) using Gsnap (Wu & Nacu 2010). The *ref_map.pl* program of Stacks was used to call SNPs within consensus RAD loci. We set the minimum number of reads required to form a stack to three and allowed two mismatches during catalogue creation. We generated SNP files in different formats for further downstream analyses using the *populations* program of Stacks and Plink v1.09 (Purcell *et al.* 2007).

(d) Population genomics analyses

We analyzed the genetic structure of *An. nili* sensu lato (s.l.) populations using Principal Component Analysis (PCA) and Neighbor-Joining trees (NJ). We also examined ancestry proportions and admixtures between populations in Admixture v1.23 (Alexander *et al.* 2009) and Structure v2.3.4 (Pritchard *et al.* 2000). We used the R package *adegenet* (Jombart 2008) to implement the PCA. Neighbor-Joining trees were generated from SNP allele frequencies via a matrix of Euclidian distance using the R package *ape* (Paradis *et al.* 2004). We ran Admixture with 10-fold cross-validation for values of k from 1 through 8. We analyzed patterns of ancestry from k ancestral populations in Structure, testing five replicates of k = 1-8. We used 200000 iterations and discarded the first 50000 iterations as burn-in for each Structure run. Clumpp v1.1.2 (Jakobsson & Rosenberg 2007) was used to summarize assignment results across independent runs. To identify the optimal number of genetic clusters in our sample, we applied simultaneously the lowest cross-validation error

in Admixture, the ad-hoc statistic deltaK (Evanno *et al.* 2005; Earl & VonHoldt 2012) and

the Discriminant Analysis of Principal Component (DAPC) method implemented in

adegenet. To examine the extent of genome divergence, we estimated locus-specific genetic

differentiation (F_{ST}) using the *populations* program of the Stacks pipeline. Mean F_{ST} values

were also used to assess pairwise divergence among populations. To infer the demographic

history of populations, we used the diffusion approximation method implemented in the

∂a∂i package (Gutenkunst *et al.* 2009). Single-population models were fitted to allele

frequency spectra and the best model was selected with the lowest likelihood and Akaike

Information Criterion as well as visual inspections of residuals.

3. Results

(a) SNP genotyping

We collected mosquitoes from four locations out of 28 sampling sites (Figure 1,

Table S1) and sequenced 145 individuals belonging, according to morphological

identifications and diagnostic PCRs, to two species (*An. nili* (n = 24) and *An. ovengensis* (n =

121)). We assembled 197724 RAD loci that mapped to unique positions throughout the

reference genome. We applied stringent filtering rules and retained 408 loci present in all

populations and in at least 50% of individuals in each population. Within these loci, we

identified 4343 high-quality biallelic markers that were used to analyze population

structure and genetic differentiation.

(b) Morphologically-defined species do not correspond to genetic clusters

PCA and NJ trees show that the genetic variation across 4343 SNPs is best explained

by more than two clusters, implying subdivisions within *An. nili* and *An. ovengensis* (Figure

2). Three subgroups are apparent within *An. nili* while two distinct clusters segregate in *An.*

ovengensis. These five subpopulations are associated with the different sampling sites

suggesting local adaptation of divergent populations. Importantly, Structure and Admixture

analyses reveals that, at k = 2, one population identified by morphology and diagnostic

PCRs as *An. nili* has almost the same ancestry pattern as the largest *An. ovengensis* cluster

(Figure 3). Such discrepancies between morphology-based and molecular taxonomies can

be due to a variety of processes including phenotypic plasticity, introgressive hybrydization

or incomplete lineage sorting (i.e., when independent loci have different genealogies by

chance) (Arnold 1997; Combosch & Vollmer 2015; Fontaine *et al.* 2015; Weng *et al.* 2016).

At k = 2 and k = 3, some populations also depict half ancestry from each morphological

species suggestive of gene flow. We found a conflicting number of genetic clusters in our

samples reflecting the complex history of subdivisions and admixtures among populations

(Figure 4). The Evanno et al. method, which highlights early divergence between *An. nili* and

An. ovengensis indicates two probable ancestors. DAPC and the Admixture cross-validation

error, which are more sensitive to recent hierarchical population subdivisions, show five or

more distinct clusters as revealed by PCA and NJ trees (Figure 4).

As suggested by the long internal branches connecting subpopulations on NJ trees,

there is strong differentiation between and within morphological species characterized by

globally high F_{ST} values (Table 1). Relatively lower F_{ST} observed between certain clusters

can be due to greater inter-population migration and intermixing or more recent

divergence. Overall, both genetic structure and differentiation reveal a set of populations

whose ancestries and species status are likely confounded by hybridization and/or

incomplete lineage sorting. The current taxonomy based on morphology and ribosomal

DNA does not effectively describe the optimal reproductive units among populations of this

group of mosquitoes.

(c) Genomic signatures of divergent selection and demographic history

Estimates of genetic differentiation varied substantially across 4343 SNP loci (Figure

5). The distribution of locus-specific F_{ST} between the five subpopulations revealed a bimodal

shape characterized by two peaks around 0 and 1. The great majority of SNPs have low to

moderate divergence, but a substantial number of variants are extremely differentiated

between populations. The maximum F_{ST} among SNPs is 1 and the number of loci with $F_{ST} = 1$

varies from 406 between the populations we termed *An. nili* goup 1 and *An. nili* group 2 to

838 between the subgroups called *An. nili* group 3 and *An. ovengensis* group 2. This pattern

of genome-wide divergence suggests that a very high number of sites with abrupt

differentiation, which likely contain genes that contribute to divergent selection and/or

reproductive isolation, coexist with regions of weak divergence that can be freely

exchanged between species.

Models of population demography indicate that all subgroups have experienced an

increase in effective size in more or less recent past (Table 2). Nevertheless, confidence

intervals of population parameters are high in some populations, and our results should be

interpreted with the necessary precautions. The population growth is less significant in *An.*

nili group 1.

4. Discussion

(a) Genetic differentiation

Advances in sequencing and analytical approaches have opened new avenues for the study of genomes of disease vectors. We have focused on malaria mosquitoes of the *An. nili* group, whose taxonomy and population structure have been challenging to resolve with low-resolution markers. We analysed genetic structure using genome-wide SNPs and found strong differentiation and local adaption among populations belonging to the two morphological species *An. nili* and *An. ovengensis*. The exact number of subpopulations remains contentious, with the suggested number of divergent clusters varying from two to five within both species. Significant population structure at eight microsatellite loci has been described among *An. nili* populations from Cameroon, with F_{ST} values as high as 0.48 between samples from the forest area (Ndo *et al.* 2013). By contrast, *An. ovengensis* was discovered recently and the population structure of this vector remains understudied. This species was initially considered as a sibling of *An. nili* (Kengne *et al.* 2003; Awono-Ambene *et al.* 2004, 2006), but more recent studies have started to challenge the assumed relatedness between the two species due to the high divergence of their polytene chromosomes (Sharakhova *et al.* 2013). Our findings call for careful review of the current taxonomy within this group of species, which is a necessary first step for accurate delineation of the role played by the different subpopulations in malaria transmission.

Our samples were collected from locations characterized by more or less degraded forest in the equatorial region of Cameroon. In those habitats, larvae of *An. nili* s.l. exploit relatively similar breeding sites consisting of slow-moving rivers (Antonio-Nkondjio *et al.* 2009). Given the similarity of habitats among populations, it is hard to pinpoint the

environmental variables that may be correlated with adaptive divergence. One of the most

expected outcomes of current large-scale malaria control measures that are underway in

Sub-Saharan African countries concerns the effects of increased insecticide exposure on

genetic diversity and population structure of vectors. The inferred demographic history of

the different subpopulations of the *An. nili* group suggests that there are currently no

bottlenecks that can be potentially linked with increased usage of insecticides and

insecticide-treated nets (e.g. Athrey *et al.* 2012). Nevertheless, further study is needed to

understand whether recent selection associated with human intervention is among the

drivers of local adaptation as observed in *An. gambiae* (Kamdem *et al.* 2017).

(b) Genomic architecture of reproductive isolation

The genomic architecture of reproductive isolation can reveal crucial information on

the mode of speciation and has been the focus of much research over the last few decades

(e.g. Turner *et al.* 2005; Harr 2006; Nadeau *et al.* 2012; Ellegren *et al.* 2012; Carneiro *et al.*

2014; Burri *et al.* 2015). One influential concept of speciation coined the "genic view of

species" proposes that boundaries between species are properties of individual genes or

genome regions and not of whole organisms or lineages (Key 1968; Barton & Hewitt 1985;

Harrison 1990; Rieseberg *et al.* 1999; Wu 2001; Nosil & Feder 2012; Harrison & Larson

2014). We have found that a high number of SNPs are strongly differentiated between

populations and often fixed within subgroups. These variants can presumably be

assimilated with regions of the genome bearing signals of reproductive isolation. However,

recent studies analyzing interactions between pairs of related taxa at their hybrid zone

indicate a complex relationship between genetic differentiation and gene flow at the

genome level (e.g. Gompert *et al.* 2012; Hamilton *et al.* 2013a; b, Larson *et al.* 2013, 2014;

Parchman *et al.* 2013; Taylor *et al.* 2014). Highly divergent genomic regions do not

necessarily coincide with regions of reduced gene flow. Several alternative interpretations

exist for the numerous high-F_{ST} regions we detected in all pairwise comparisons (Noor &

Bennett 2009; Nachman & Payseur 2012; Cruickshank & Hahn 2014; Delmore *et al.* 2015),

but careful examination of these outlier of differentiation will reveal significant insights into

the markers of ecological divergence and/or reproductive isolation. More complete genome

assemblies will be necessary to better delineate specific regions of the genome under

natural selection, and thereby clarify the genomic basis of phenotypic fitness differences

between divergent populations.

Signals consistent with gene flow between *An. nili* and *An. ovengensis* are apparent

in our data despite significant time since divergence (~3M-yr) (Ndo *et al.* 2013). Some

individuals depict almost half ancestry from each morphological species. The disagreement

observed between morphology/PCR and molecular taxonomies also suggests that

incongruent genealogies may be widespread along chromosomes due to hybridization

and/or incomplete lineage sorting. However, hybridization can be difficult to detect

because other factors such as incomplete lineage sorting or technical artefacts can leave

signatures that are similar to those of interspecific gene flow (Patterson *et al.* 2012; Liu *et*

al. 2014). A complete reference genome is also necessary to analyze the detailed

distribution of genealogies across genomic windows and to disentangle the relative

contribution of processes that generate the putative admixtures and species confusion

observed among divergent populations (Fontaine *et al.* 2015; Weng *et al.* 2016).

5. Conclusions and implications

Delineating the fine-scale population structure of mosquito populations is crucial to understand their epidemiological significance and their potential response to vector control measures. Moreover, recent malaria control efforts are affecting interspecific gene flow, genetic differentiation, population demography and natural selection in mosquitoes (Athrey *et al.* 2012; Clarkson *et al.* 2014; Norris *et al.* 2015; Barnes *et al.* 2017; Kamdem *et al.* 2017). Deciphering the signatures of all these processes across mosquito genome is important to minimize their negative impacts on vector control. Our findings shed some light on the complex evolutionary history of members of the *An. nili* group and provide a framework for future investigations into the genetic basis of ecological and reproductive barriers.

Acknowledgements

Funding for this project was provided by the University of California Riverside and NIH grants 1R01AI113248 and 1R21AI115271 to BJW. We thank populations and administrative authorities of the sampling sites included in this study for their collaboration.

Data Archiving Statement

Data for this study are available at: to be completed after manuscript is accepted for publication

References

Alexander DH, Novembre J, Lange K (2009) Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, **19**, 1655–1664.

Antonio-Nkondjio C, Kerah CH, Simard F *et al.* (2006) Complexity of the malaria vectorial system in Cameroon: contribution of secondary vectors to malaria transmission. *Journal of medical entomology*, **43**, 1215–1221.

Antonio-Nkondjio C, Ndo C, Awono-Ambene P *et al.* (2007) Population genetic structure of the malaria vector Anopheles moucheti in south Cameroon forest region. *Acta Tropica*, **101**, 61–68.

Antonio-Nkondjio C, Ndo C, Costantini C *et al.* (2009) Distribution and larval habitat characterization of Anopheles moucheti, Anopheles nili, and other malaria vectors in

river networks of southern Cameroon. *Acta Tropica*, **112**, 270–276.

Antonio-nkondjio C, Simard F (2013) Highlights on Anopheles nili and Anopheles moucheti,
Malaria Vectors in Africa. In: *Anopheles mosquitoes - New insights into malaria vectors*
(ed Manguin S)

Arnold ML (1997) *Natural Hybridization and Evolution*.

Athrey G, Hodges TK, Reddy MR *et al.* (2012) The effective population size of malaria
mosquitoes: large impact of vector control. *PLoS genetics*, **8**, e1003097.

Awono-Ambene HP, Kengne P, Simard F, Antonio-Nkondjio C, Fontenille D (2004)
Description and bionomics of Anopheles (Cellia) ovengensis (Diptera: Culicidae), a new
malaria vector species of the Anopheles nili group from south Cameroon. *Journal of*
medical entomology, **41**, 561–568.

Awono-Ambene HP, Simard F, Antonio-Nkondjio C *et al.* (2006) Multilocus enzyme
electrophoresis supports speciation within the Anopheles nili group of malaria vectors
in Cameroon. *The American journal of tropical medicine and hygiene*, **75**, 656–658.

Ayala D, Fontaine MC, Cohuet A *et al.* (2011) Chromosomal inversions, natural selection and
adaptation in the malaria vector Anopheles funestus. *Molecular biology and evolution*,
28, 745–758.

Barnes KG, Weedall GD, Ndula M *et al.* (2017) Genomic Footprints of Selective Sweeps from
Metabolic Resistance to Pyrethroids in African Malaria Vectors Are Driven by Scale up
of Insecticide-Based Vector Control. *PLoS Genetics*, **13(2): e10**, 1–22.

Barton N, Hewitt G. (1985) Analysis of hybrid zones. *Annual Review of Ecology and*
Systematics, **16**, 113–148.

Van Bortel W, Harbach RE, Trung HD *et al.* (2001) Confirmation of Anopheles varuna in
vietnam, previously misidentified and mistargeted as the malaria vector Anopheles
minimus. , **65**, 729–732.

Burri R, Nater A, Kawakami T *et al.* (2015) Linked selection and recombination rate
variation drive the evolution of the genomic landscape of differentiation across the
speciation continuum of Ficedula flycatchers. *Genome Research*, **25**, 1656–1665.

Caputo B, Nwakanma D, Caputo FP *et al.* (2014) Prominent intraspecific genetic divergence
within Anopheles gambiae sibling species triggered by habitat discontinuities across a
riverine landscape. *Molecular ecology*, **23**, 4574–89.

Carneiro M, Albert FW, Afonso S *et al.* (2014) The genomic architecture of population
divergence between subspecies of the European Rabbit. *PLoS Genetics*, **10**.

Catchen JM, Amores A, Hohenlohe P, Cresko W, Postlethwait JH (2011) Stacks: building and
genotyping Loci de novo from short-read sequences. *G3 (Bethesda, Md.)*, **1**, 171–82.

Catchen J, Hohenlohe P a., Bassham S, Amores A, Cresko W. (2013) Stacks: An analysis tool
set for population genomics. *Molecular Ecology*, **22**, 3124–3140.

Clarkson CS, Weetman D, Essandoh J *et al.* (2014) Adaptive introgression between
Anopheles sibling species eliminates a major genomic island but not reproductive
isolation. *Nature communications*, **5**, 4248.

Combosch DJ, Vollmer S V. (2015) Trans-Pacific RAD-Seq population genomics confirms
introgressive hybridization in Eastern Pacific Pocillopora corals. *Molecular*
Phylogenetics and Evolution, **88**, 154–162.

Cruickshank TE, Hahn MW (2014) Reanalysis suggests that genomic islands of speciation
are due to reduced diversity, not reduced gene flow. *Molecular ecology*, **23**, 3133–57.

Delmore KE, Hübner S, Kane NC *et al.* (2015) Genomic analysis of a migratory divide reveals

candidate genes for migration and implicates selective sweeps in generating islands of
differentiation. *Molecular Ecology*, 1873–1888.
Dia I, Guelbeogo M, Ayala D (2013) Advances and Perspectives in the Study of the Malaria
Mosquito Anopheles funestus. In: *Anopheles mosquitoes - New insights into malaria*
vectors (ed Manguin S)
Earl DA, VonHoldt BM (2012) STRUCTURE HARVESTER: a website and program for
visualizing STRUCTURE output and implementing the Evanno method. *Conservation*
Genetics Resources, **4**.
Ellegren H, Smeds L, Burri R *et al.* (2012) The genomic landscape of species divergence in
Ficedula flycatchers. *Nature*, **491**, 756–60.
Evanno G, Goudet J, Regnaut S (2005) Detecting the number of clusters of individuals using
the software structure: a simulation study. *Molecular Ecology*, **14**, 2611–2620.
Fontaine MC, Pease JB, Steele a. *et al.* (2015) Extensive introgression in a malaria vector
species complex revealed by phylogenomics. *Science*, **347**, 1258524–1258524.
Fouet C, Kamdem C, Gamez S, White BJ (2017) Extensive genetic diversity among
populations of the malaria mosquito Anopheles moucheti revealed by population
genomics. *Infection, Genetics and Evolution*, **48**, 27–33.
Gillies MT, Coetzee M (1987) *A supplement to the Anophelinae of Africa south of the Sahara*.
The South African Institute for Medical Research, Johannesburg.
Gillies MT, De Meillon B (1968) *The Anophelinae of Africa South of the Sahara*. Publications
of the South African Institute for Medical Research, Johannesburg.
Gompert Z, Lucas LK, Nice CC *et al.* (2012) Genomic regions with a history of divergent
selection affect fitness of hybrids between two butterfly species. *Evolution*, **66**, 2167–
2181.
Gutenkunst RN, Hernandez RD, Williamson SH, Bustamante CD (2009) Inferring the joint
demographic history of multiple populations from multidimensional SNP frequency
data. *PLoS genetics*, **5**, e1000695.
Hamilton JA, Lexer C, Aitken SN (2013a) Genomic and phenotypic architecture of a spruce
hybrid zone (Picea sitchensis × P. glauca). *Molecular Ecology*, **22**, 827–841.
Hamilton J a., Lexer C, Aitken SN (2013b) Differential introgression reveals candidate genes
for selection across a spruce (Picea sitchensis × P. glauca) hybrid zone. *New Phytologist*,
197, 927–938.
Harbach RE (2013) The Phylogeny and Classification of Anopheles. In: *Anopheles*
mosquitoes - New insights into malaria vectors (ed Manguin S)
Harr B (2006) Genomic islands of differentiation between house mouse subspecies. , 730–
737.
Harrison RG (1990) Hybrid zones: windows on evolutionary process. In: (eds Futuyma D, J.
Antonovics), pp. 69–128. New York.
Harrison RG, Larson EL (2014) Hybridization, introgression, and the nature of species
boundaries. *Journal of Heredity*, **105**, 795–809.
Hemingway J, Ranson H, Magill A *et al.* (2016) Averting a malaria disaster: Will insecticide
resistance derail malaria control? *The Lancet*, **387**, 1785–1788.
Jakobsson M, Rosenberg N (2007) CLUMPP: a cluster matching and permutation program
for dealing with multimodality in analysis of population structure. *Bioinformatics*, **23**,
1801– 1806.
Jombart T (2008) adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics, **24**, 1403–1405.

Kamdem C, Fouet C, Gamez S, White BJ (2017) Pollutants and insecticides drive local
adaptation in African malaria mosquitoes. *Molecular Biology and Evolution*, **Accepted**.

Kengne P, Awono-Ambene, H. P. Antonio-Nkondjio C, Simard F, Fontenille D (2003)
Molecular identification of the Anopheles nili group African malaria vectors. *Med Vet*
Entomol, **17**, 67–74.

Key KHL (1968) The concept of stasipatric speciation. *Syst. Zool.*, **17**, 14–22.

Krzywinski J, Besansky NJ (2003) Molecular systematics of Anopheles : from subgenera to
subpopulations. *Annual Review of Entomology*, **48**, 111–139.

Lanzaro GC, Lee Y (2013) Speciation in Anopheles gambiae — The Distribution of Genetic
Polymorphism and Patterns of Reproductive Isolation Among Natural Populations. In:
Anopheles mosquitoes - New insights into malaria vectors (ed Manguin S)

Larson EL, Andrés JA, Bogdanowicz SM, Harrison RG (2013) Differential introgression in a
mosaic hybrid zone reveals candidate barrier genes. *Evolution*, **67**, 3653–3661.

Larson EL, White TA, Ross CL, Harrison RG (2014) Gene flow and the maintenance of
species boundaries. *Molecular Ecology*, **23**, 1668–1678.

Lawniczak MKN, Emrich SJ, Holloway a K *et al.* (2010) Widespread divergence between
incipient Anopheles gambiae species revealed by whole genome sequences. *Science*
(New York, N.Y.), **330**, 512–4.

Liu KJ, Dai J, Truong K *et al.* (2014) An HMM-Based Comparative Genomic Framework for
Detecting Introgression in Eukaryotes. *PLoS Computational Biology*, **10**.

Molineaux L, Gramiccia G (1980) *The Garki Project. Research on the epidemiology and*
control of malaria in the sudan savanna of West Africa. World Health Organization.

Nachman MW, Payseur BA (2012) Recombination rate variation and speciation : theoretical
predictions and empirical results from rabbits and mice. , 409–421.

Nadeau NJ, Whibley A, Jones RT *et al.* (2012) Genomic islands of divergence in hybridizing
Heliconius butterflies identified by large-scale targeted sequencing. , 343–353.

Ndo C, Antonio-Nkondjio C, Cohuet A *et al.* (2010) Population genetic structure of the
malaria vector Anopheles nili in sub-Saharan Africa. *Malaria journal*, **9**, 161.

Ndo C, Simard F, Kengne P *et al.* (2013) Cryptic Genetic Diversity within the Anopheles nili
group of Malaria Vectors in the Equatorial Forest Area of Cameroon (Central Africa).
PLoS ONE, **8**, 1–12.

Neafsey DE, Lawniczak MKN, Park DJ *et al.* (2010) SNP genotyping defines complex gene-
flow boundaries among African malaria vector mosquitoes. *Science (New York, N.Y.)*,
330, 514–7.

Noor M a F, Bennett SM (2009) Islands of speciation or mirages in the desert? Examining
the role of restricted recombination in maintaining species. *Heredity*, **103**, 439–444.

Norris LC, Main BJ, Lee Y *et al.* (2015) Adaptive introgression in an African malaria
mosquito coincident with the increased usage of insecticide-treated bed nets.
Proceedings of the National Academy of Sciences, 201418892.

Nosil P, Feder JL (2012) Widespread yet heterogeneous genomic divergence. *Molecular*
Ecology, **21**, 2829–2832.

O'Loughlin SM, Magesa S, Mbogo C *et al.* (2014) Genomic Analyses of Three Malaria Vectors
Reveals Extensive Shared Polymorphism but Contrasting Population Histories.
Molecular biology and evolution, 1–14.

Paradis E, Claude J, Strimmer K (2004) Analyses of Phylogenetics and Evolution in R

language. *Bioinformatics*, **20**, 289–290.

Parchman TL, Gompert Z, Braun MJ *et al.* (2013) The genomic consequences of adaptive
divergence and reproductive isolation between species of manakins. *Molecular ecology*,
22, 3304–17.

Patterson N, Moorjani P, Luo Y *et al.* (2012) Ancient admixture in human history. *Genetics*,
192, 1065–1093.

Peery A, Sharakhova M V, Antonio-Nkondjio C *et al.* (2011) Improving the population
genetics toolbox for the study of the African malaria vector Anopheles nili:
microsatellite mapping to chromosomes. *Parasites & Vectors*, **4**, 202.

Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012) Double Digest RADseq: An
Inexpensive Method for De Novo SNP Discovery and Genotyping in Model and Non-
Model Species. *PLoS ONE*, **7**, e37135.

Pinto J, Egyir-Yawson A, Vicente J *et al.* (2013) Geographic population structure of the
African malaria vector Anopheles gambiae suggests a role for the forest-savannah
biome transition as a barrier to gene flow. *Evolutionary applications*, **6**, 910–24.

Pritchard JK, Stephens M, Donnelly P (2000) Inference of population structure using
multilocus genotype data. *Genetics*, **155**, 945–959.

Purcell S, Neale B, Todd-Brown K *et al.* (2007) PLINK: a toolset for whole-genome
association and population-based linkage analysis. *American Journal of Human*
Genetics, **81**.

Ranson H, Lissenden N (2016) Insecticide Resistance in African Anopheles Mosquitoes: A
Worsening Situation that Needs Urgent Action to Maintain Malaria Control. *Trends in*
Parasitology, **32**, 187–196.

Reidenbach KR, Neafsey DE, Costantini C *et al.* (2012) Patterns of genomic differentiation
between ecologically differentiated M and S forms of Anopheles gambiae in West and
Central Africa. *Genome biology and evolution*, **4**, 1202–12.

Riehle MM, Guelbeogo WM, Gneme A *et al.* (2011) A cryptic subgroup of Anopheles gambiae
is highly susceptible to human malaria parasites. *Science (New York, N.Y.)*, **331**, 596–8.

Rieseberg LH, Whitton J, Gardner K (1999) Hybrid Zones and the Genetic Architecture of a
Barrier to Gene Flow Between Two Sunflower Species.

Service MW (1993) *Mosquito ecology: field sampling methods* (UK Elsevier Applied Science,
London, Ed,).

Sharakhova M V., Peery A, Antonio-Nkondjio C *et al.* (2013) Cytogenetic analysis of
Anopheles ovengensis revealed high structural divergence of chromosomes in the
Anopheles nili group. *Infection, Genetics and Evolution*, **16**, 341–348.

Sinka ME, Bangs MJ, Manguin S *et al.* (2010) The dominant Anopheles vectors of human
malaria in Africa, Europe and the Middle East: occurrence data, distribution maps and
bionomic précis. *Parasites & vectors*, **3**, 117.

Slotman M a, Tripet F, Cornel a J *et al.* (2007) Evidence for subdivision within the M
molecular form of Anopheles gambiae. *Molecular ecology*, **16**, 639–49.

Taylor SA, Curry RL, White TA, Ferretti V, Lovette I (2014) Spatiotemporally consistent
genomic signatures of reproductive isolation in a moving hybrid zone. *Evolution*, **68**,
3066–3081.

Touré Y, Petrarca V, Traoré S *et al.* (1998) The distribution and inversion polymorphism of
chromosomally recognized taxa of the Anopheles gambiae complex in Mali, West
Africa. *Parassitologia*, **40**, 477–511.

Turner TL, Hahn MW, Nuzhdin S V. (2005) Genomic islands of speciation in Anopheles
gambiae. *PLoS Biology*, **3**, 1572–1578.
Weng D, Yu Y, Hahn MW, Nakhleh L (2016) Reticulate evolutionary history and extensive
introgression in mosquito species revealed by phylogenetic network analysis.
Molecular Ecology, **Special is**.
White B, Lawniczak M, Cheng C *et al.* (2011) Adaptive divergence between incipient species
of Anopheles gambiae increases resistance to Plasmodium. *Proceedings of the National*
Academy of Sciences, **108**, 244–249.
WHO (2016) *World malaria report 2016*.
Wondji C, Simard F, Petrarca V (2005) Species and populations of the Anopheles gambiae
complex in Cameroon with special emphasis on chromosomal and molecular forms of
Anopheles gambiae ss. *Journal of medical entomology*.
Wright S (1984) *Evolution and the Genetics of Populations: Genetics and Biometric*
Foundations v. 1 (Genetic & Biometric Foundations); New Edition. University of Chicago
press.
Wu C (2001) The genic view of the process of speciation. *Journal of Evolutionary Biology*, **14**,
851–865.
Wu TD, Nacu S (2010) Fast and SNP-tolerant detection of complex variants and splicing in
short reads. *Bioinformatics*, **26**, 873–881.
508

509 **Author contributions**
510 Conceived and designed the experiments: CK CF BJW. Performed the experiments: CK CF SG
511 BJW. Analyzed the data: CK CF. Wrote the paper: CK CF BJW.
512
513
514

Tables

Table 1: Pairwise F_{ST} between divergent subpopulations.

F_{ST}	An. nili group 1	An. nili group 2	An. nili group 3	An. ovengensis group 1	An. ovengensis group 2
An. nili group 1	-				
An. nili group 2	0.374	-			
An. nili group 3	0.506	0.552	-		
An. ovengensis group 1	0.135	0.275	0.364	-	
An. ovengensis group 2	0.432	0.458	0.492	0.349	-

Table 2: Demographic models of different subgroups.

Population	Best Model	Log Likelihood	Final Population Size[a] (95% CI)	Time[b] (95% CI)
An. nili group 1	*Growth*	-18.42	6.41 (5.326 - 20.71)	3.70 (1.11 - 13.31)
An. nili group 2	*Two-epoch*	-19.97	17.87 (9.33 - 35.50)	11.27 (4.93 - 19.64)
An. ovengensis group 1	*Growth*	-112.18	13.04 (12.15 - 17.26)	0.70 (0.58 - 1.08)
An. Ovengensis group 2	*Growth*	-22.98	19.95 (14.45 - 45.70)	5.11 (2.33 - 15.13)

[a] Relative to ancestral population size.

[b] Expressed in units 2Ne generations from start of growth to present.

 Figures
521 **Figure 1**: Map showing the sampling locations and the relative frequencies of *An. nili* and
522 *An. ovengensis*. Small and large black dots indicate respectively the 28 locations surveyed
523 and the four sampling sites where mosquitoes were collected.
524

525
526

527 **Figure 2**: Population genetic structure inferred from 4343 SNPs using PCA (A) and a
528 neighbor-joining tree (B). The percentage of variance explained is indicated on each PCA
529 axis.
530

531
532
533

534 **Figure 3:** Ancestry proportions inferred in ADMIXTURE with k = 2 – 8.
535

536
537
538

539 **Figure 4:** Identification of the optimal number of genetic clusters using the delta k method
540 of Evanno et al (A), DAPC (B) and 10-fold cross-validation in ADMIXTURE (C). The lowest
541 BIC and CV error and the highest delta k indicate the most probable number of clusters.
542
543

544
545
546

Figure 5: Distribution of F_{ST} values across 4343 SNPs between *An. nili* group 1 and *An. nili*
group 2 (A); *An. nili* group 1 and *An. nili* group 3 (B); *An. nili* group 1 and *An. ovengensis*
group 1 (C); *An. nili* group 1 and *An. ovengensis* group 2 (D); *An. nili* group 2 and *An. nili*
group 3 (E); *An. nili* group 2 and *An. ovengensis* group 1 (F); *An. nili* group 2 and *An.*
ovengensis group 2 (G); *An. nili* group 3 and *An. ovengensis* group 1 (H); *An. nili* group 3 and
An. ovengensis group 2 (I); *An. ovengensis* group 1 and *An. ovengensis* group 2 (J). The
number of SNPs with F_{ST} = 1 is indicated in each pairwise comparison.
554

 Supplemental information

 Table S1: Information on mosquito samples included in this study.

Sampling locations	Geographic coordinates	Sampling methods			Total
		HLC-OUT	**HLC-IN**	**LC**	
Ebebda	4°20'00"N, 11°17'00"E			6	6
Nkoteng	4°31'00"N, 12°02'00"E			8	8
Nyabessan	2°24'00"N, 10°24'00"E	63	44		107
Mbébé	4°10'00"N, 11°04'00"E	13	3	8	24
Total		76	47	22	145

HLC-OUT, human landing catches performed outdoor; HLC-IN, human landing catches performed indoor; LC, larval collection